Taxonomic distribution of metabolic functions underpins nutrient cycling in *Trichodesmium* consortia

Coco Koedooder[1,2,3*], **Futing Zhang**[1,2], **Siyuan Wang**[1,2], **Subhajit Basu**[1,2,4], **Sheean T. Haley**[5], **Nikola Tolic**[6], **Carrie D. Nicora**[6], **Sonya T. Dyhrman**[5,8], **Martha Gledhill**[9], **Rene M. Boiteau**[10,11], **Maxim Rubin-Blum**[3], **Yeala Shaked**[1,2]

[1]*The Fredy and Nadine Herrmann Institute of Earth Sciences, Hebrew University of Jerusalem, Jerusalem, Israel*
[2]*The Interuniversity Institute for Marine Sciences in Eilat, Eilat, Israel*
[3]*Israel Oceanographic and Limnological Research, Haifa, Israel*
[4]*Microsensor Research Group, Max Planck Institute for Marine Microbiology, Bremen, Germany*
[5]*Lamont-Doherty Earth Observatory, Columbia University, New York, NY, United States*
[6]*Biological Sciences Laboratory Pacific Northwest National Laboratory, Richland, WA, United States*
[7]*Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, MI, United States*
[8]*Department of Earth and Environmental Sciences, Columbia University, New York, NY, United States*
[9]*GEOMAR, Helmholtz Center for Ocean Research, Kiel, Germany*
[10]*Environmental Molecular Sciences Laboratory Pacific Northwest National Laboratory, Richland, WA, United States*
[11]*College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, Corvallis, OR, United States*

* Correspondence: Coco Koedooder (coco.koedooder@mail.huji.ac.il)

Running Title: Associated Bacteria of Red Sea *Trichodesmium* populations

Keywords: *Trichodesmium*, metagenomes, associated bacteria, siderophore biosynthesis, vitamin B, denitrification, dissimilatory nitrate fixation to ammonia, phosphonate, nitrogen fixation

Abstract

The photosynthetic and diazotrophic cyanobacterium *Trichodesmium* is a key contributor to marine biogeochemical cycles in the subtropical-oligotrophic oceans. *Trichodesmium* forms colonies that harbor a distinct microbial community, which expands their functional potential and is predicted to influence the cycling of carbon, nitrogen, phosphorus and iron (C, N, P, and Fe). To link key traits to taxa and elucidate how community structure influences nutrient cycling, we assessed Red Sea *Trichodesmium* colonies using metagenomics and metaproteomics. This diverse consortium comprises bacteria that typically associate with algae and particles, such as the ubiquitous *Alteromonas macleodii,* but also lineages specific to *Trichodesmium*, such as members from the order Balneolales. These bacteria carry functional traits that would influence resource cycling in the consortium, including siderophore biosynthesis, reduced phosphorus metabolism, vitamins, denitrification, and dissimilatory-nitrate-reduction-to-ammonium (DNRA) pathways. Denitrification and DNRA appeared to be modular as bacteria collectively completed the steps for these pathways. The vast majority of associated bacteria were auxotrophic for vitamins, indicating the interdependency of consortium members. *Trichodesmium* in turn may rely on associated bacteria to meet its high Fe demand as several lineages can synthesize the photolabile siderophores vibrioferrin, rhizoferrin, and petrobactin, enhancing the bioavailability of particulate-Fe to the entire consortium. Our results highlight that *Trichodesmium* is a hotspot for C, N, P, Fe, and vitamin exchange. The functional redundancy of nutrient cycling in the consortium likely underpins its resilience within an ever-changing global environment.

Importance (150 words)

Colonies of the cyanobacteria *Trichodesmium* act as a biological hotspot for the usage and recycling of key resources such as C, N, P and Fe within an otherwise oligotrophic environment. While *Trichodesmium* colonies are known to interact with a unique community of algae and particle-associated microbes, our understanding of the taxa that populate these colonies and the gene functions they encode is still limited. Characterizing the taxa and adaptive strategies that influence consortium physiology and its concomitant biogeochemistry is critical in a future ocean predicted to have increasing particulate fluxes and resource-depleted regions.

Introduction

Trichodesmium spp. are a globally relevant group of cyanobacteria that can form surface blooms visible from space (1). Owing to their high abundance and capability to fix both carbon and nitrogen, they are considered key players in the biogeochemical cycling of C and N within the oligotrophic tropical and subtropical oceans (2–5). A key trait underpinning the functionality of Trichodesmium is that its filaments can cluster together to form large colonies of 1-2 mm in size. The formation of colonies allows Trichodesmium to capture dust that is deposited on the ocean surface and Trichodesmium is reported to actively select for Fe and P-rich particles and shuffle them to their colony core (6–8). Colony formation hereby adds an intriguing spatial component where Trichodesmium spp. is predicted to act as a biological hotspot not only for C and N, but also for the uptake and recycling of key limiting nutrients such as Fe and P.

Trichodesmium colonies harbor a unique microbial consortium of epibiotic bacteria. Amplicon sequencing has shown these associated bacteria are distinct from populations in the surrounding oligotrophic surface waters (9–11). At a genetic level, the associated microbial community was found to enrich the total functional potential of the collective colony beyond the traits of N_2 and C fixation (12–14). By enriching the genetic repertoire of metabolic functions, or traits, related to the nutrients C, N, Fe, and P, consortia members were predicted to influence the internal cycling of these nutrients (14–17). To understand the biogeochemical contributions of Trichodesmium within the global ocean it is therefore critical to look beyond the metabolic functions present in Trichodesmium and towards the processes taking place within the larger microbial community. While metagenomic studies of Trichodesmium colonies in the Atlantic (13) and the Pacific Oceans (12, 17, 18) highlighted key functional traits within the consortium, only one study linked pathways to the associated bacteria from 8

metagenome-assembled bins (13). Expanding these studies, and characterizing how functional

traits are distributed across different taxa is critical to understanding the role of *Trichodesmium*

associated bacteria in consortium physiology and its concomitant influence over the cycling of

key resources such as C and N.

Recent evidence from the Red Sea highlights the unique ability of *Trichodesmium* consortia

members to acquire particulate Fe from dust through the production of Fe-chelating

siderophore molecules (7, 8, 16, 19, 20). Siderophores dissolve particulate Fe from dust and

are predicted to increase its bioavailability for both *Trichodesmium* and its associated bacteria

(16, 21, 22). In this manner, *Trichodesmium* spp. interacts with their associated bacteria in

order to meet the high Fe-requirements for the processes of photosynthesis and N_2 fixation (23).

This finding is particularly relevant in light of climate model simulations, which predict an

increase in regional dust emissions in areas such as the Red Sea where *Trichodesmium* is known

to occur (24, 25). While *Trichodesmium* blooms are a re-occurring phenomenon in this area

(26), it represents an under sampled region where the taxonomic composition of its consortium

has not been explored genetically. Furthermore, the presence and taxonomic distribution of

siderophore biosynthesis pathways within *Trichodesmium* colonies has not yet been

interrogated in depth. Understanding which consortium members positively enhance the

bioavailability of Fe within the *Trichodesmium* consortium will help elucidate key interactions

and community dynamics taking place within the colony that influence the ability of

Trichodesmium to fix C and N. This information is critical to evaluating *Trichodesmium*

contributions to C and N cycling in the future ocean.

In this study, we conducted an in-depth metagenomic and proteomic study of *Trichodesmium*

colonies collected from the Red Sea. To elucidate how the community structure influences the

functional dynamics of the *Trichodesmium* consortium we identified the different taxa making

up the *Trichodesmium* consortium and examined the presence of functional traits involved in

nutrient cycling. We probed our dataset for genes, proteins and pathways involved in the uptake

and metabolism of N, P, Fe, and vitamins. Finally, we contextualize these findings by

comparing them to previous metagenomic studies of *Trichodesmium* colonies obtained from

the Atlantic and the Pacific oceans.

Results and Discussion

A total of 52 metagenomic-assembled genomes (MAGs) were obtained from *Trichodesmium*

colonies, of which 42 MAGs were of high quality (90% complete and <5% redundancy

(Supplementary Table 1) (27). The relative abundance of the different MAGs was consistent

across the three samples with *T. thiebautii* MAG 52 making up 71% ± 6 of the total relative

abundance (Supplementary Table 1). In parallel to the metagenomic analysis, a proteomic

dataset consisting of 14161 peptide sequences matching to protein sequences of 52 different

MAGs further probed the activity within the Red Sea *Trichodesmium* consortia

(Supplementary Table 2; Supplementary Table 3).

Figure 1. Phylogenetic tree of the 52 MAGs assembled from *Trichodesmium* colonies collected in the Red Sea.
The tree is rooted by the outgroup *Fuseobacterium nucleatum*. Known bacterial orders are marked in different
colours and several MAGs could taxonomically be identified at the genus level. The relative abundance of each
MAG within the *Trichodesmium* consortium, excluding the abundance of Trichodesmium MAG 52 (±70% of
samples), is presented as a bubble chart. The 11 MAGs that matched to MAGs assembled from a previous
metagenomic dataset of Red Sea *Trichodesmium* colonies (28) are highlighted in blue and tentatively represent
the core-consortium (see Supplementary Figure 2; Supplementary Table 4). MAGs that did not meet the
requirements of a high-quality genome (>90% completeness; <5% redundancy) are indicated in grey. Note that
re-occurring MAGs are not necessarily the most abundant MAGs present within the Trichodesmium colony. (a
full taxonomic description of each MAG in this dataset can be found in Supplementary Table 1).

Trichodesmium colonies host diverse and flexible associated bacteria

Trichodesmium colonies consisted of a single *Trichodesmium* species (*T. thiebautii* MAG 52)

and a diverse consortium of 51 associated bacteria spanning at least 10 different taxonomic

orders (Figure 1). Unlike a previous study (28), our samples did not include any non-

diazotrophic *Trichodesmium* populations and the 51 MAGs hereby represent the associated bacteria of *T. thiebautii* MAG 52. The majority of the associated bacterial MAGs belonged to taxonomic orders known to be present in *Trichodesmium* colonies and included the orders Rhodobacterales, Pseudomonadales, Balneolales, Rhodospirillales, Flavobacterales, Enterobacterales, Rhizobiales, and Sphingomonadales (9, 11–13).

An intercomparison between the MAGs observed here and a previous metagenomic dataset from the Red Sea (28) , identified a total of 11 MAGs present in both datasets (Supplementary Figure 2; Supplementary Table 4). These 11 MAGs included the particle- and algae-associated lineages (29–31) of *Marinobacter* sp. (MAG23) and *Alcanivorax* sp. (MAG 47) from the order Pseudomonadales; *Rhodobacter* sp. (MAG 28) and *Sulfitobacter* sp. (MAG29) from the order Rhodobacterales, and *Alteromonas macleodii* (MAG 10), a ubiquitous bacterium often found in association with phototrophs (32) and previously isolated from *T. erythraeum* IMS101 cultures (33). If these observations are consistent across a larger time-period of observations, these MAGs may represent a core component of the *Trichodesmium* consortium. Further comparison of our metagenomic dataset with those obtained from the Atlantic, North or South Pacific oceans, and the Red Sea (12, 13, 17, 18, 28), showed taxonomic similarities at the order level rather than at the species level. While most taxonomic orders re-occurred in all datasets, only one Balneolales genome (MAG 33) was present amongst all *Trichodesmium* metagenomic datasets. As none of the re-occurring MAGs represented highly abundant genomes within our samples (Figure 1), this may partially be the result of a higher sequencing depth (500 million reads) compared to past datasets. Nonetheless, the wide diversity coupled with the recurrence of only a few MAGs indicates the presence of a flexible assemblage of associated bacteria, which may have important implications for the community structure and functioning of the *Trichodesmium* consortium.

Trichodesmium associated bacteria synthesize diverse photolabile siderophores, which may increase iron availability from dust

We explored the different Fe-uptake mechanisms present within the *Trichodesmium*

consortium, as N_2 and C fixation in *Trichodesmium* is often constrained by Fe availability (23,

34). Despite the high Fe-requirements of *Trichodesmium* (23)*, T. thiebautii* (MAG 52) only

contained genes for the uptake of free Fe^{+2} and Fe^{+3}, but not organic Fe, suggesting it is limited

to a narrow range of forms it can directly take up (Figure 2). In contrast, MAGs of the

associated bacteria contained a larger diversity of Fe-uptake genes which included the ability

to take up organically bound Fe, such as heme, citrate, and siderophores (Figure 2). The

proteome of the *Trichodesmium* consortium indicated that, at the time of sampling,

Trichodesmium was actively taking up Fe through the presence of a ferric Fe-uptake

transporters (FbpC, FbpA) and Fe-storage proteins (Dps, Bfr) (Supplementary Table 2).

The genomes of 10 *Trichodesmium* associated bacteria spread across 6 different taxonomic

orders encoded different siderophore biosynthesis gene clusters (BGCs) (Figure 2). The

presence of a siderophore biosynthesis pathway was strain-specific and not present in an entire

clade or lineage. Similar to previous studies, *T. thiebautii* MAG 52 did not contain any known

siderophore biosynthesis pathways (12, 35). Siderophore BGCs could further be separated into

4 NIS-type and 6 NRPS-type siderophores (36, 37). Based on the presence of one or more

iucA/iucC domains, the 4 NIS-types BGCs matched closely to petrobactin (*Alteromonas*

macleodii MAG 10), vibrioferrin (*Marinobacter salarius* MAG 23; *Limnobacter sp.* MAG 11),

and rhizoferrin (Rhodospirillales MAG 08) biosynthesis pathways (Figure 2). From the 6

different NRPS-type siderophores, only 2 matched a known BGC, namely that of the

membrane-bound amphibactin (100%) (*Alcanivorax venustensis* MAG 47) and one resembling

the BGC of the siderophores agrobactin (93%) or enterobactin (±83%) (*Roseibium* sp MAG 18) (Figure 2). Our results additionally indicated the presence of several novel NRPS-type siderophore biosynthesis pathways (Supplementary Figure 3). NRPS-type-siderophore BGCs are not as well characterized as NIS-type-siderophores BGCs and can be difficult to distinguish from other secondary metabolites such as toxins and antibiotics (38). While the 4 NRPS-type-siderophore BGCs could not be linked to known siderophore BGCs, several unknown metallophores have previously been identified and associated with *Trichodesmium* colonies and we speculate that some may be linked to these uncharacterized NRPS BGCs (19). The presence of multiple NRPS pathways containing a putative siderophore TonB-dependent receptor (SMCOG1082) likely involved in Fe uptake, highlights the potential for novel siderophore pathways within the *Trichodesmium* consortium and indicates the importance of additional culture-based research linking such pathways to a siderophore molecule.

Our results indicate that the *Trichodesmium* consortium in the Red Sea may interact with photolabile siderophore-producing bacteria to mine Fe from dust. The re-occurring consortium members *Marinobacter salarius* (MAG 23), *Alcanivorax venustensis* (MAG 47), and *Alteromonas macleodii* (MAG 10) are all able to synthesize the NIS-type siderophores petrobactin, vibrioferrin, and rhizoferrin (Figure 2) whose citrate-moiety results in their photolability (39). In the natural illumination of surface waters, where *Trichodesmium* resides, sunlight can reductively disassociate Fe from these siderophore complexes and provide a flux of highly available dissolved inorganic Fe (40, 41). While *Trichodesmium* does not contain a TonB-receptor, the presence of photolabile siderophores implies that Fe is also bioavailable to consortium members that lack a specific TonB-receptor for that particular siderophore. These photolabile siderophores act as a "common good" within the *Trichodesmium* consortia, where only a few members carrying the trait are needed to benefit the entire consortium (42). This

contrasts with the uptake mode of photostable siderophores, such as desferrioxamine-B

(DFOB), which require a specialized TonB-siderophore transport system (43). Similarly,

amphibactin is a membrane-bound siderophore (44), which prevents its diffusion in the

environment and subsequently its ability to act as a public good to the *Trichodesmium*

consortium. A recent gene-knockout study in *Alteromonas macleodii* verified the ability of

petrobactin to enhance the bioavailability of particulate Fe (22). This further supports the idea

that *Trichodesmium* can interact with consortium members that produce photolabile

siderophores to acquire Fe from dust.

Most of the siderophore BGCs detected here were identified in ubiquitous and copiotrophic

bacteria typically found in association with particles and algae (11, 22) , including

Marinobacter salarius (MAG 23), *Alcanivorax venustensis* (MAG 47), and *Alteromonas*

macleodii (MAG 10) (Figure 2). Siderophore-production within the *Trichodesmium*

consortium appears to be linked to a lifestyle associated with a rapidly changing environment

rich in N and C, where regulating Fe-bioavailability can provide a competitive advantage. In

line with particle and algae associations, siderophore production has also been associated with

other ecological functions, including heavy metal detoxification and intra-specific

communication (45) (Supplementary Figure 4; Supplementary Table 5). Whether siderophore

production can benefit the *Trichodesmium* consortium in functions other than Fe-dissolution

from dust requires further investigation. Nonetheless, the presence of a few, yet reoccurring,

siderophore producing MAGs augments previous measurements of siderophore production

within the consortium (16, 19, 20) and reveals additional layers of *Trichodesmium* colonies'

intriguing ability to sequester Fe from dust.

Figure 2 (below). MAGs were probed for Fe-uptake and siderophore biosynthesis genes which are noted in
Supplementary Table 5. **(a)** *Trichodesmium* MAG 52 does not contain a wide diversity of Fe-uptake genes, and
the majority of MAGs of associated bacteria contain an uptake receptor for organic Fe (heme, citrate,
siderophores). Several MAGs of associated bacteria contained an NRPS or NIS-type siderophore biosynthesis
pathways (BGCs). **(b)** Siderophore biosynthetic gene clusters (BGCs) of known NIS-type siderophore BGCs can

be clustered according to the presence of one or multiple *iuc*A/*iuc*C domains which enables one to target the closest siderophore BGC match (52). Using this method, we were able to link these pathways to the synthesis of the photolabile **(i)** petrobactin, **(ii)** vibrioferrin and **(iii)** rhizoferrin. **(c)** Siderophore BGCs of known NRPS siderophores. Using antiSMASH 7.0, which aligns BGCs to known siderophores within the MIBiG database, we were able to link 2 of the 6 putative siderophore NRPS-type pathways to **(i)** the membrane-bound amphibactin and **(ii)** agrobactin/enterobactin. (See Supplementary Figure 3 for more details on the siderophore biosynthesis pathways).

The *Trichodesmium* consortium is heterogeneous in genes related to the uptake and use of inorganic and organic phosphorous.

We predicted that the majority of bacteria associated with *Trichodesmium* have the ability to metabolize phosphite or phosphonate. This can offer a key competitive advantage amongst consortium members in a P-limited environment (18, 46) including the Gulf of Aqaba in the Red Sea (47). Within a low-P environment, dissolved inorganic phosphorus is scarce, and bacteria must scavenge phosphate from dissolved organic phosphorus to meet their P-requirements (48). *Trichodesmium* is highly efficient at scavenging and utilizing organic-P through the production of alkaline phosphatase (AP), and the uptake of phosphonate and phosphite (49). Multiple studies have shown that *Trichodesmium* colonies in the Atlantic and Pacific Ocean are hotspots for the reduced P-cycling of phosphite or phosphonate (46, 50) suggesting that the majority of consortium members can process reduced-P.

Figure 3. MAGs were probed for several different P-metabolism genes which are noted in Supplementary Table 5. *Trichodesmium* is known to be highly competitive for P, as exemplified by the multitude of **(a)** P-uptake systems present in its genome for phosphate (*pst*S, *pst*B), phosphonate (*phn*CDE), and phosphonate/phosphite (*phn*CDE/*ptx*ABC). **(b)** Within the *Trichodesmium* consortium, the ability to metabolize a broad range of phosphonate using a broad C-P lyase (*phn*GHIJKLM) is more prominent than the substrate specific hydrolysis of

242 2-aminoethylphosphonate (2-AEP; *phnWXYZA*) or phosphonopyruvate (*pal*A) and the phosphite metabolism
243 (*ptx*D).
244
Confirming past studies (18, 51), *T. thiebautii* MAG 52 encodes a set of functions to transport

and metabolize diverse forms of P (Figure 3). This includes the presence of a high-affinity

phosphate transporter (*pst*SCAB), and a phosphonate/phosphite transporter

*(phn*CDE/*pxt*ABC). In addition to these transport systems, genes related to utilization and

metabolism of organic-P included alkaline phosphatase (AP) (*pho*A, *pho*X), and a carbon-

phosphorus (C-P) lyase (*phn*GHIJKLM) which hydrolyses a broad range of phosphonate

compounds. *T. thiebautii* MAG 52 also encodes phosphite dehydrogenase (*ptxD)* which

catalyzes the oxidation of phosphite to phosphate. The presence of PstB, PhoX, and PhnD

within the proteome (Supplementary Table 2), suggests *Trichodesmium* was taking up

phosphate, phosphonate, and hydrolyzing phosphoester compounds to meet its P demand.

Several of these proteins are regulated by P (51), and their detection in the proteome is

consistent with intense competition for P in the low P Red Sea.

As KEGG annotation cannot accurately distinguish between a phosphonate (*phn*CDE) and a

phosphite (*pxt*ABC) transporter, the presence of adjacent P-metabolism genes can further

indicate the P-source linked to it. Specifically, the presence of phosphite dehydrogenase (*ptxD)*

points to phosphite metabolism while a broad-specificity C-P lyase (*phnGHIJKLM*) or the

substrate-specific hydrolysis of phosphonopyruvate (*pal*A) or 2-aminoethylphosphonate (2-

AEP) to acetaldehyde (*phn*WX), acetate (*phn*WYA), or glycine (*phn*Y'Z), can point to the

metabolism of phosphonate (52, 53). Our results showed that the presence of a

*phn*CDE/*ptx*ABC uptake system was specific to certain taxonomic orders including

Pseudomonadales, Rhodospirillales, Rhodobacterales, Caulobacterales, and Rhizobiales

(Figure 3). Within the *Trichodesmium* consortium, these taxonomic orders may have a

competitive advantage over members that lack them, particularly in low P conditions. Of these

25 MAGs, 16 could be linked to phosphonate metabolism through the presence of a broad-

specificity C-P lyase. In contrast, the presence of phosphite dehydrogenase was species-

specific where only 4 MAGs appeared to be able to process phosphite spread across different

taxonomic orders (Figure 3). Within the *Trichodesmium* consortium, phosphonate metabolism

appears to be the more common reduced P-currency of the two. 7 MAGs contained genes

associated with the substrate-specific hydrolysis of 2-AEP but pathways were not complete,

making it difficult to draw conclusions (Figure 3). Nonetheless, the presence of a broad-

specificity C-P lyase appeared to be the most prevalent strategy for phosphonate utilization

within the *Trichodesmium* consortium. This finding is in contrast to a meta-analysis of marine

metagenomes where the most common phosphonate degradation strategy within the water

column was the substrate-specific hydrolysis of 2-AEP (54). 2-AEP is the most abundant

phosphonate source in the marine environment (55), and the differences in enzyme distribution

within the Trichodesmium consortium may reflect heterogeneity in the phosphonate

compounds present in the colony, relative to the water column. Differences in enzyme

distribution may also be a function of the taxonomic composition of *Trichodesmium* associated

bacteria relative to those in in the water column. Regardless, these observations suggest that a

diversity of P compounds are being cycled by members of the consortia in the low P Red Sea

and offer a new reference dataset for future comparisons with *Trichodesmium* consortia from

other regions.

Denitrification and dissimilatory nitrate reduction to ammonia (DNRA) pathways are

modular in the *Trichodesmium* consortium.

We explored the presence of multiple N-metabolic pathways as they are predicted to influence

the biogeochemical contribution of *Trichodesmium* regarding N_2 fixation. Our results showed

that, in the *Trichodesmium* colonies sampled from the Red Sea, N_2-fixation appears to be

uniquely attributed to *T. thiebautii* MAG 52 (Figure 4). This contrasts with a previous study

that has observed *nifH* genes in both *Trichodesmium* and their associated bacteria (12). In our

study, however, only *T. thiebautii* (MAG 52) contained the full N_2 fixation pathway

(*nifDKHEB*: M00175) and therefore the ability to fix N_2 within the colony. Similar to other

studies (12, 17, 56), genes of the two additional N-transformation pathways of dissimilatory

nitrate reduction to ammonium (DNRA) and denitrification were present within the

Trichodesmium consortium (Figure 4). Respectively, denitrification and DNRA are N-loss and

N-recycling processes and are predicted to enable the residing community to conserve and fully

utilize the N_2 fixed by *Trichodesmium* (12) but also alter the net contribution of *Trichodesmium*

colonies to local N-cycling dynamics (15, 57). Our results indicated a relatively high number

of 24 MAGs encoding genes involved in DNRA and 10 MAGs encoding genes involved in

denitrification (Figure 4). Intriguingly, none of the MAGs analyzed contained the complete

pathways for denitrification and DNRA, with the exception of Rhizobiales for the latter (Figure

4). Collectively, MAGs were able to complete all steps within either pathway, indicating that

these pathways appear to be modular within the *Trichodesmium* consortium, where a bacteria

can perform one step within the pathway but not necessarily all steps. While Rhizobiales (MAG

16, 17 and 44) contained the full DNRA pathway, which includes the reduction of nitrate to

nitrite (*napAB, narGH*), and nitrite to ammonium (*nrfAH, nirBD*), the remaining MAGs

contained only one of two steps (Figure 4). Similar to DNRA, several MAGs from the phyla

Alphaproteobacteria contained a part of the denitrification steps, including the reduction of

nitrite to nitric-oxide (*nir*KS) and nitric-oxide to nitrous oxide (*nor*BC), while members of the

phyla Bacteroidetes could perform the final step within the denitrification pathway (*nosZ)*

which converts nitrous oxide to nitrogen (Figure 4).

Figure 4. MAGs were probed for N-metabolic pathways and the genes for each step within the pathway are noted in Supplementary Table 5. *Trichodesmium* MAG 52 is the only MAG with a N$_2$ fixation pathway. Several MAGs contained one or two steps within the pathways of denitrification or dissimilatory nitrate to reduction to ammonia (DNRA) indicating their modularity. Collectively, MAGs completed the denitrification and DNRA pathways, but none contained the genes for the entire pathway.

While the presence of modularity in denitrification and DNRA pathways itself is not

uncommon (58–60) finding it present within the *Trichodesmium* consortium highlights the

spatial role *Trichodesmium* plays in providing a platform where each step can be performed by

a distinct subset of the microbial consortium. While the re-occurring consortium members

Alteromonas macleodii (MAG 10)*, Marinobacter salarius* MAG 23)*,* and *Alcanivorax sp*

(MAG 47, MAG 48) encoded genes (*nir*BD) within the DNRA pathway, it did not appear to

be a key characteristic within the *Trichodesmium* associated bacteria of the Red Sea. The

presence of DNRA and denitrification pathways is rare in the oxic surface ocean, as these

processes require anoxic and suboxic conditions (61). In the surface waters, suboxic as well as

anoxic micro-environments were detected in marine aggregates, such as marine snow and fecal

pellets, where enhanced microbial respiration results in the consumption of O$_2$ (62, 63).

Trichodesmium colonies may represent an additional environment where such processes may

be taking place within the surface waters of the ocean. While it is not entirely clear when and

where these N-cycling pathways are actively taking place within the colony studies have

measured DNRA and denitrification processes within natural *Trichodesmium* colonies (15, 57). While the obligatory suboxic conditions for denitrification and DNRA may occur in the presence of dust and the rapid O_2 consumption by bacteria (64), the formation of these microenvironments have thus far not been observed in *Trichodesmium* colonies (57). Altogether, these findings pinpoint the intricate community structure needed for such modular processes to take place within the consortium. The exact tradeoffs between *Trichodesmium* and denitrifying or DNRA bacteria, particularly in light of a changing ocean with higher dust deposition requires further investigation.

Trichodesmium consortium members are interdependent for vitamins B1, B7 and B12.

We explored the presence of vitamin auxotrophy within the consortium as an indicator for community interdependence. Previous studies have shown that *Trichodesmium* can synthesize and secrete the vitamin cobalamin (B12) to the environment (17). Our results show that while *Trichodesmium* MAG 52 encodes the biosynthesis pathways for B12, biotin (B7) and thiamin (B1), the majority of associated bacteria (49 of the 52 MAGs) lack a biosynthesis pathway for at least one of these vitamins (Figure 5). Bacteroidetes, Sphingomonadales, and Enterobacterales species, including *Alteromonas macleodii* (MAG 10), were all auxotrophic for vitamin B12. MAGs from the order Rhodospirillales lacked the ability to synthesize vitamin B7, whereas Rhizobiales and Rhodobacterales appeared to be auxotrophic for both vitamin B1 and B7. Only *Pseudoalteromonas* encoded all three biosynthesis pathways, including the re-occurring taxa *Alcanivorax* sp. (MAG 47) and *Marinobacter salarius* (MAG 23). These consortium members may also contribute to the synthesis and provision of vitamins B1, B7, and B12 to their auxotrophic counterparts (Figure 5) but, based on overall relative abundance, *Trichodesmium* represents the most likely source of B vitamins. In sum, the majority of associated bacteria (49 of 52 MAGs) were predicted to be auxotrophic for B12 and likely

363 dependent on *Trichodesmium* to meet their intracellular requirements. Our results indicate that

364 the production and secretion of different vitamins by *Trichodesmium* is a key determinant of

365 community composition, stimulating vitamin-based interactions among auxotrophic

366 consortium members.

Figure 5. MAGs were probed for vitamin B1, B7, and B12 biosynthesis and uptake pathways. The genes used to
indicate the biosynthesis or uptake of these vitamins are noted in Supplementary Table 5. *Trichodesmium* contains
the biosynthesis pathways for all three vitamins, but the vast majority of associated bacteria were auxotrophic for
one or more vitamins. These results suggest that the associated bacteria rely on *Trichodesmium* to obtain their
vitamin requirements.

Interactions between copiotrophic microbes underpin nutrient cycling in *Trichodesmium* consortia

A literature study of the bacterial lineages present within the *Trichodesmium* consortium

describes these taxonomic orders as typical primary colonizers of marine surfaces such as algae

or sinking particles (29–31). Strains of *Alteromonas*, *Marinobacter,* and *Pseudomonas* often

co-occur with phototrophs such as cyanobacteria as well as oil spills, likely as a result of their

ability to degrade hydrocarbons and aromatic compounds (32, 65). While the prevalence of

these copiotrophic lifestyles within the *Trichodesmium* consortium has been observed in

previous studies (11, 13) it is still unclear whether their association with *Trichodesmium* is

specific or reflective of a more general interaction between algae or particle associations.

Copiotrophy is a widely encompassing term, but it generally concerns opportunistic bacteria

with a versatile metabolic capacity to use a wide range of substrates, acting as important remineralizers by rapidly responding towards areas enriched in organic matter (66, 67). It is therefore possible to consider the majority of the consortium as active remineralizers of the C, N, P, Fe, and vitamins *Trichodesmium* is providing. Viewing *Trichodesmium* as a hotspot for nutrients is supported by our proteomic samples where the majority of protein hits attributed to *T. thiebautii* MAG 52 were related to photosynthetic (K05376; K05377; K02284) and diazotrophic machinery (K03839, K02588, K02591) indicating that the processes of N and C fixation were actively taking place at the time of collection (Supplementary Table 2). In turn, the majority of protein hits associated to consortia members were involved in the uptake of amino-acids (K01999; K09969), sugars (K02055; K02027; K10552), and peptides (K02035; K02032) which is reflective of typical copiotrophic activity (Supplementary Table 3).

Copiotrophic bacteria are also characterised by their strong ability to interact within their associations (68). This is supported by the prominence of several bacterial-interactive and particle-associated traits in the MAGs studied here (68, 69) (Supplementary Figure 4; Supplementary Table 5). The presence of bacterial-associated interactions taking place between *Trichodesmium,* and their consortium members could further be implied through the presence of several proteins involved in pilus formation (K02658, K02662, K02669) and secretion systems (K11003; K03072; K03070, K03110, K03073; K03217) within *T. thiebautii* MAG 52 (Supplementary Table 2). Collectively, these findings are indicative of an intricate network of interaction between *Trichodesmium* and its associated bacteria. For example, *T. thiebautii* MAG 52 and several MAGs, including re-occurring members from the order Rhodobacterales, contained a homoserine lactone biosynthesis pathway (Supplementary Figure 4a). Previous studies have predicted that consortium members can directly interact with *Trichodesmium* through the production of homoserine lactones (HLs), where the presence of

HLs was shown to modulate N_2 fixation rates (14) and increase alkaline phosphatase (AP)

activity in *Trichodesmium* (70). The functional association of poorly studied bacterial clades,

in particular Balneolales (Supplementary Table 1), however, remained more elusive and

indicated that we still lack a clear understanding of these bacteria and their role within the

Trichodesmium consortium. Further investigations will be required to untangle the different

dynamics of both general and specific interactions taking place between members of the

Trichodesmium consortium.

Conclusions and Future Perspectives

The complex *Trichodesmium* consortium represents an ideal system to study the intricate

interactions taking place within a particle-rich system (Figure 6), asking how these processes

may impact the C, N, P and Fe-cycles in a rapidly changing ocean where both microbial

community composition and changes in microbial abundance are expected to occur (25, 71).

The *Trichodesmium* consortium is redundant for multiple traits related to nutrient cycling

including siderophore production, denitrification, DNRA, and reduced phosphate uptake. The

processes of denitrification and DNRA were modular, indicating an interactive and spatial

coupling between different consortium members. The associated bacteria were largely

auxotrophic for one or more vitamins that *Trichodesmium* can synthesize, and this is likely a

major trait influencing the community structure (Figure 6). The vast majority of the associated

bacteria could further be characterized according to a copiotrophic lifestyle where

Trichodesmium represents a hotspot for C, N, P, Fe, and vitamins to the residing community.

On the other hand, Fe-uptake through the production of siderophores appeared to be a key

beneficial function *Trichodesmium* can obtain from its residing community. *Trichodesmium*

harbored several bacteria that can secrete photolabile siderophores, supporting the idea that

Trichodesmium can meet their high Fe-demand by interacting with siderophore-producing

bacteria to mine Fe from dust particles collected within the colony. Other, as of yet

uncharacterized, traits will likely further elucidate the factors that drive the community

structure of *Trichodesmium* colonies. Our results highlight the presence of a dynamic and

flexible consortium where functional traits are well conserved within the *Trichodesmium*

consortium, underpinning the resilience of the colony within an ever-changing ocean.

440
Figure 6. **Schematic summary of the potential interactions taking place within the *Trichodesmium* consortia**
as discussed in this study. Each blue bubble indicates a different function, and each blue circle pools together
the different MAGs containing a particular trait for that function. Each MAG is represented by a coloured dot
reflective of the taxonomic order it is associated with (grey dots represent MAGs that were not attributed to a
specific taxonomic order). *Trichodesmium* is highlighted in cobalt-blue, and the important functions
Trichodesmium is known to display are highlighted with a dotted circle. (See Supplementary Table 5 for the full
list of functional traits and their presence/absence). **(a)** Several siderophore biosynthesis genes are present in a
diverse set of MAGs. Siderophores are predicted to mine Fe from dust that has been collected and centered within
Trichodesmium colonies. 6 of the 4 were identified as known NIS-type siderophores of which petrobactin,
vibrioferrin and rhizoferrin are photolabile. In the presence of sunlight, the Fe complexed to these siderophores
disassociates and acts as a public good to the larger consortium. The genes and pathways for **(b-i)** denitrification,
(b-ii) N_2 fixation, and **(b-iii)** dissimilatory- nitrate- reduction- to- nitrate (DNRA) are present in several MAGs of
the consortium. These processes appear to be modular within the *Trichodesmium* consortium as different
taxonomic groups complete the steps for these pathways. **(c)** Several MAGs contain a homoserine lactone

synthesis pathway (HL), including *Trichodesmium*. HL acts as an intercommunicating quorum sensing molecule
and has been shown to influence the production of alkaline phosphatase (AP) and N_2 fixation in *Trichodesmium*.
(d) *Trichodesmium* is capable of producing the vitamins cobalamin (B12), thiamin (B1) and biotin (B7). Different
subsets of MAGs lack the biosynthesis pathways for one or more of these vitamins with very few capable of
synthesizing all three (associated bacteria can also produce vitamins but is not depicted here - see Figure 2).
Consortia members and *Trichodesmium* likely exhibit an interdependent relationship for these vitamins. **(e)**
Trichodesmium colonies have been shown to take up and utilize phosphonate and phosphite in a low-P
environment and MAGs containing these pathways may have a competitive advantage over members that lack
them.

Material and Methods

Trichodesmium metagenomic sampling and extraction

Trichodesmium puff colonies were hand-picked during the spring bloom (6th May, 2019) and

separated into three samples for metagenomics (~100-200 colonies each) using a 100 μm

phytoplankton net at 20 m depth in the Gulf of Aqaba (Eilat, Israel) (29.56ºN, 34.95ºE).

Colonies were washed 3 times, by gently picking colonies with a pipette and placing them in

new petri dishes with fresh 0.2 μm filtered seawater, before being filtered on a 0.2 μm, 45 mm

polycarbonate (PC) filter using a vacuum pump. Filters were flash frozen in liquid N_2 and kept

at -80ºC. DNA was extracted from the three *Trichodesmium* samples using the DNeasy Plant

Pro Kit (Qiagen), with a minor modification to the lysis procedure. The kit-provided tissue

disruption tubes were not used. Rather, ~250 μL zirconia/silica beads (0.5 mm) were added

to each sample tube before the addition of Solution CD1, and then samples were vortexed for

5 min. The resulting lysate was processed as per the remainder of the manufacturer instructions.

DNA was quantified fluorometrically using the dsDNA HS kit (ThermoFisher Scientific) on a

Qubit. From the three samples, metagenomic libraries were prepared and sequenced with 0.7

billion reads 2×151 bp reads on Illumina NovaSeq S4 at the DOE Joint Genome Institute

(California, USA). This study can be found under JGI Gold Study ID number Gs0149370.

Metagenomic raw read assembly and binning

Sequences were analysed from this study and compared to previously published metagenomic

studies from the Red Sea (PRJNA804487) (28) and metagenomes of *Trichodesmium* colonies

isolated from the Pacific (PRJNA435427) (17) (PRJNA358796) (12) and the Atlantic

(PRJNA330990) (13) Oceans. All raw sequences were analyzed using the same protocol

described below.

Metagenomes were assembled, binned, and quantified using the ATLAS (v2) pipeline (72).

Briefly, raw sequences underwent quality control through the BBTools suite (73, 74) and were

assembled using metaSPAdes (75) (k-mer lengths: 21, 33, 55, 99, 121 bp). MAGs were binned

from each sample using MetaBAT 2 (76), MaxBin 2.0 (77) and VAMB (78). The completeness

and redundancy of each bin were subsequently assessed using CheckM2 v0.1.2 (79). MAGs

were considered of high-quality if they reached 90% completeness and 10% redundancy and

were highlighted throughout our results and figures (27). A non-redundant set of bins was

produced using DAS Tool (80) and dRep (81) based on an average nucleotide identity (ANI)

cutoff of 97.5%. Genes were predicted using Prodigal (82). The coverage of each MAG was

quantified across each sample by mapping reads back to each MAG. We refer to the MAGs

from this study as MAG XX, bacterial MAGs from a previous Red Sea study as MAG R-XX

(28) and bacterial MAGs from other metagenomic studies from the Pacific and Atlantic Oceans

(12, 13, 17) as MAG T-XX.

Phylogenetic Diversity

Metagenomically assembled genomes (MAGs) were taxonomically characterized using the

genome taxonomy database tool kit GTDB-tk v2.1.1 (83). MAGs were classified according to

GTDB taxonomy and NCBI taxonomy (Supplementary Table 1; Supplementary Table 4).

GTDB taxonomy is used throughout the analysis, but to simplify comparisons with the

literature, we used the larger NCBI taxonomic order Rhodospirillales which encompasses the

smaller GTDB taxonomic groups of Kiloniellales and Thalassobaculaceae.

Phylogeny was inferred using GTo-Tree (v.1.16.12; default settings) (84) from a concatenation

of 74 conserved single-copy HMM markers for bacteria using the best-fit model Q.pfam+R7

model in IQ-Tree (v2.1.3), using the Bayesian information criterion (BIC) (85, 86).

Shimodaira–Hasegawa approximate likelihood-ratio test (SH-aLRT) and ultrafast bootstrap

approximation (UFBoot) branch support values were estimated from 1000 bootstraps. The tree

was visualized using FigTree (v1.4.4) and rooted using the genome of *Fuseobacterium*

nucleatum (PRJNA1419) as an outgroup according to (87). MAG were considered to re-occur

in another metagenomic dataset if its genome sequence identity was >97.5% similar

(Supplementary Figure 1).

Functional Diversity

The amino-acid sequence for each MAG was annotated using hmmscan and GhostKoala under

default parameters (50% identity cutoff), utilizing the PFAM and KEGG databases,

respectively (88, 89). Secondary metabolites (e.g. biosynthesis of siderophores and homoserine

lactones) were screened for each MAG using AntiSMASH 7.0. In certain cases, the Rapid

Annotation using Subsystem Technology (RAST) server was used to annotate amino-acid

sequences using the collection of protein families, FIGfams (90, 91). In addition, each MAG

was assessed using METABOLIC (v4.0) which lists the presence of interactive KEGG

modules for each MAG using a default completeness cutoff of 0.75 (92). Based on these

different annotation formats, we inspected each MAG for the presence of functional traits. For

each functional pathway or gene, the annotation method and coinciding KEGG and PFAM ID

number can be found in Supplementary Table 5. The presence and absence of different traits

for each MAG was displayed using the Interactive Tree Of Life (iTOL) visualization tool (93).

 Iron metabolism

 To explore the relationship between *Trichodesmium* and its associated bacteria for the

 bioavailability of Fe from dust, we searched MAGs for the KEGG IDs of multiple Fe-uptake

 pathways. Fe-uptake receptors were separated according to abiotic Fe^{+2} (*feoB*:K04759), Fe^{+3}

 (*afuA*:K02012), and organic Fe uptake which is further separated into Fe-heme

 (*hemR*:K16087), Fe-citrate (*fecA*:K16091) and Fe-siderophores (*fevS*:K02016). The presence

 of an Fe:siderophore receptor K02016 should be taken cautiously as its KEGG annotation does

 not clearly distinguish between siderophores or cobalamin as a substrate.

 Siderophore biosynthesis pathways within MAGs were identified using a conservative

 approach that required a combination of both KEGG and HMM annotation for each MAG and

 the use of AntiSMASH 7.0 and FeGenie tools (94) (Supplementary Figure 4). Pathways were

 separately analyzed for both Non-Ribosomal Peptide Synthetase (NRPS) Type and NRPS

 Independent Synthetase (NIS) Type biosynthesis pathways. NIS-type siderophore biosynthesis

 pathways are more easily identifiable of the two, and we searched for the presence of the Fe-

 uptake chelate (*iuc)A/iuc*C domain (PF04183). NIS genes were clustered into different types

 using known NIS biosynthesis genes as defined and described by Carroll and Moore (36).

 Briefly, genes were aligned using the online Multiple Alignment Fast Fourier Transformation

 (MAFFT v7.490; L-INS-i) software (95). A phylogenetic tree was constructed from this

 alignment using IQtree2 (v2.1.3). The resulting best-fit substitution model Q.pfam+F+R4 was

 selected using Bayesian information criteria (BIC). Branches were assigned using Shimodira–

 Hasegawa approximate likelihood-ratio test (SH-aLRT) and ultrafast bootstrap approximation

 (UFBoot) branch support values were estimated from 1,000 bootstraps. The resulting

 consensus tree was visualized using FigTree (v1.4.4).

The identification of NRPS-type siderophore biosynthesis pathways required FeGenie, which

utilizes biosynthetic pHMMs, and was used to highlight putative NRPS-type siderophores that

were further inspected using AntiSMASH 7.0. MAGs containing an NRPS pathway coupled

with a siderophore transport-receptor (SMCOG1082) within the gene-cluster were marked as

putative NRPS-siderophore producers. These pathways were further compared to known

NRPS siderophore pathways using the MiBIG database (96) and the KEGG pathway map for

NRPS-type siderophores biosynthesis gene cluster (BGC) map01053.

Vitamin B1, B7, and B12 biosynthesis and uptake

We assume that a bacterial strain is auxotrophic for thiamin (B1), biotin (B7) or cobalamin

(B12) if their coinciding genome lacks a cobalamin biosynthesis pathway and contains a

transporter for its uptake. We searched MAGs for the KEGG IDs and determined the presence

of a vitamin biosynthesis pathway using a KEGG map for each vitamin. The presence of a

thiamin biosynthesis pathway (M00127) was determined by the presence of *thi*DCEGL

(K00946; K00788; K03149; K00941; K03147). A thiamin uptake system was present if a

coinciding transporter system (M00192) was observed. The presence of a biotin biosynthesis

pathway (M00581) was determined by the presence of *bio*ABDFH (K01012; K01935;

K00833; K00652; K02170). A biotin uptake system was present if a coinciding transporter

system (M00581) was observed. The presence of a cobalamin biosynthesis pathway was

determined by the presence of *cob*ACDUQS (K19221; K02226; K02227; K02231:K02233).

A putative vitamin B12 uptake system was considered present in a MAG if the specific

transporter *btu*B (K16092) was observed.

N-metabolism pathways om MAGs were compiled from a selected set of KEGG ID numbers

using METABOLIC-C (v4.0) and the KEGG pathway map ko00910. MAGs were queried for

the presence of genes related to N_2 fixation, dissimilatory nitrate reduction to ammonium

(DNRA), and denitrification. The ability to fix N_2 was determined through the presence of

*nif*HKD (K02588, K02591, K02586). DNRA can be separated into two steps and includes the

reduction of nitrate to nitrite (*nap*AB: K02567, K02568*; nar*GH: K00370-K00374) and nitrite

to ammonia (*nrf*AH: K15876, K03385*, nir*BD: K00362, K00363). Denitrification can further

be separated into the reduction of nitrite to nitric oxide (*nir*S:K15864*; nir*K: K00368), nitric

oxide to nitrous oxide (*nor*BC: K04561, K02305) and nitrous oxide to N_2 (*nos*Z: K00376).

Phosphorus metabolism

The presence of multiple P-metabolic pathways within the *Trichodesmium* consortium were

explored. We explored several transporters for high-affinity phosphate (*pst*SCAB: K02036-

K02038, K02040), low-affinity phosphate (*pitA*:K03306), and a

phosphite/phosphonate transporter (*phn*CDE: K02041; K02042; K02044). We searched

MAGs for KEGG ID numbers for the presence of phosphite dehydrogenase (*ptx*D:K18916),

which oxidizes phosphite to phosphate, and the broad specificity C-P lyase (*phn*GHIJKLM:

K06162-6, K05780-1), which hydrolyses phosphonate bonds in organic P. Additionally,

MAGs were queried for the presence of the substrate-specific phosphonate metabolism

enzymes including *phn*A (K19670), *phn*X (K05306), and *phn*W (K03430) and collectively

these genes were used to indicate the functional potential for reduced P-metabolism. Lastly,

MAGs were examined for the presence of alkaline phosphatases (AP) *pho*A (K01077), *pho*X

(K07093), and *pho*D (K01113). AP hydrolyses phosphoesters enabling the metabolism of
organic-P.

Bacterial and Particle Interaction traits

Known traits that are indicative of putative bacterial interactions were previously summarized
and identified by Zoccarato *et al* for marine bacterial genomes (68). In addition to siderophore
biosynthesis and vitamin B1, B7 and B12 biosynthesis (see previously), MAGs were probed
for the presence of homoserine lactone (HLs) biosynthesis genes using AntiSMASH 7.0. The
presence of auxin efflux genes was determined according to RAST annotation. MAGs were
screened for KEGG ID numbers for the presence of the quorum sensing regulation gene *lux*R
(K07782) and *bja*R1 (K18098), secretion systems (KEGG map03070), and motility and
adhesion genes (KEGG map02020) (Supplementary Table 5). Traits involved in particle-
interaction were previously described and summarized by Debeljak *et al* (69). In addition to
Fe-uptake and siderophore biosynthesis pathways, MAGs were screened for KEGG ID
numbers involved in metal-response, metal-efflux and metal-storage pathways (Supplementary
Table 5).

Trichodesmium proteomic sampling and extraction

In parallel to the metagenomic sampling, 20 puff colonies were picked, in triplicate, for
proteomic analysis at the Environmental Molecular Sciences Laboratory (EMSL) at Pacific
Northwest National Laboratory according to standardized protocols. Colonies were washed 3
times with filtered seawater, as described above. Samples were centrifuged, and the resulting
pellet was diluted in 200 µl 8 M urea and transferred to 2 mL pre-filled Micro-Organism Lysing
Mix glass bead tubes and bead beat in a Bead Ruptor Elite bead mill homogenizer (OMNI
International, Kennesaw Georgia) at speed 5.5 for 45 sec. After bead beating, the lysate was

transferred to a new 4 mL tube and immediately placed in an ice block and spun at 1,000 x g

for 10 mins at 4°C. 200 µl of the lysed samples were transferred into 2 mL centrifuge tubes. A

bicinchoninic acid (BCA) assay (Thermo Scientific, MA USA) was performed to determine

protein concentration. Following the assay, 10 mM dithiothreitol (DTT) was added to the

samples and incubated at 60°C for 30 mins with constant shaking at 800 rpm. Samples were

diluted 8-fold for preparation for digestion with 100 mM NH_4HCO_3, 1 mM $CaCl_2$ and

sequencing grade trypsin (Promega, WI) was added to all protein samples at a 1:50 (w/w)

trypsin-to-protein ratio for 3 hrs at 37˚C, 450 rpm. Digested samples were desalted using a 4-

probe positive pressure Gilson GX-274 ASPEC™ system (Gilson Inc., WI) with Discovery

C18 50 mg/1 mL solid phase extraction tubes (Supelco, MO), using the following protocol: 3

mL of methanol was added for conditioning followed by 3 mL of 0.1% trifluoroacetic acid

(TFA) in H_2O. The samples were then loaded onto each column followed by 4 mL of 95:5:

H_2O:ACN, 0.1% TFA. Samples were eluted with 1mL 80:20 ACN:H_2O, 0.1% TFA. The

samples were concentrated using a speed vac and a final BCA was performed to determine the

peptide concentration and samples were fractionated into 12 fractions for LC-MS/MS analysis.

647
MS analysis was performed using a Q-Exactive Plus mass spectrometer (Thermo Scientific)

outfitted with a homemade nano-electrospray ionization interface. Electrospray emitters used

150 µm o.d. × 20 µm i.d. chemically etched fused silica (126). The ion transfer tube

temperature and spray voltage were 300°C and 2.2 kV, respectively. Data was collected for

120 min following a 10 min delay after completion of sample trapping and start of gradient.

FT-MS spectra were acquired from 300 to 1800 m/z at a resolution of 70 k (AGC target 3e6)

and the top 12 FT-HCD-MS/MS spectra were acquired in data-dependent mode with an

isolation window of 1.5 m/z at a resolution of 17.5 k (AGC target 1e5) using a normalized

collision energy of 30, dynamic exclusion time of 30 s, and detected charge state of an ion 2 or

higher. The resulting proteomic samples from this study was deposited in MassIVE

(MSV000091416).

Proteomic Analysis

Peptide matching was performed using MSGF+ software (97) against the amino-acid

sequences obtained from the matching metagenome dataset through PROKKA (98) using the

ATLAS pipeline (v2) (72) resulting in 201374 protein sequences. MSGF+ was used in

target/decoy mode with 20 ppm parent ion tolerance, partial tryptic rule, and M-oxidation

(+15.9949) as a dynamic modification. Best matches from the MSGF+ searches were filtered

at 1% FDR based on the target/decoy model; this set of peptides was used in consequent

quantitation analysis. Spectral counts were performed without consideration of peptide

specificity for proteins and MAGs. 92% of peptides mapped to unique proteins while 93% of

peptides were found in only one MAG. Spectral count normalization was performed using the

normalized spectral abundance factor (NSAF) approach (99). Briefly, counts were divided by

the matching peptide length and their relative abundance was calculated from the total number

of spectral hits per sample.

Acknowledgements

This research was performed on a project award (EMSL 50403) under the FICUS program and

used resources at the DOE Joint Genome Institute and the Environmental Molecular Sciences

Laboratory, which are DOE Office of Science User Facilities. Both facilities are sponsored by

the Biological and Environmental Research program and operated under Contract Nos. DE-

AC02-05CH11231 (JGI) and DE-AC05-76RL01830 (EMSL). This research was supported by

Grant no. 2020041 awarded to YS and RB from the United States-Israel Binational Science

Foundation (BSF). This research was supported by Grant no. 2020041 awarded to YS and RB

from the United States-Israel Binational Science Foundation (BSF) and the Israel Ministry of

Science and Technology (MOST) Grant no. 001126 to MR-B. Additional support was

provided by a grant from the Simons Foundation award ID 721225 to STD.

Competing Interests

The authors declare no conflicting interests.

Data Availability Statement

Raw sequences are available under the Gold Study ID number Gs0149370. The 52
metagenomic assembled genomes analysed in this study are available at NCBI under the
BioProject number PRJNA944101. Proteomic samples are available at MassIVE under the
project accession number MSV000091416.

Author Contributions

YS, RB, MG, SB, conceived of this study. SB, FZ, SW collected the DNA and protein samples.
SB performed the experiments together with FZ, SW and YS. SD, SH, TG extracted and
sequenced the DNA. CDN extracted and processed protein samples that were analyzed by NT
and RB. MRB and CK performed the metagenomic analysis. CK wrote the manuscript under
guidance of YS, MRB, RB, MG, SD, with further input from SW, SB, FZ, NT.
698
699

1.   Bergman B, Sandh G, Lin S, Larsson J, Carpenter EJ. 2013. *Trichodesmium*– a
widespread marine Cyanobacterium with unusual nitrogen fixation properties. FEMS
Microbiol Rev 37:286–302.
2.   Zehr JP. 2011. Nitrogen fixation by marine cyanobacteria. Trends Microbiol 19:162–
173.
3.   Capone DG, Burns JA, Montoya JP, Subramaniam A, Mahaffey C, Gunderson T,
Michaels AF, Carpenter EJ, Capone C, Burns JA, Montoya JP, Subramaniam A,
Mahaffey C, Gunderson T, Michaels AF. 2005. Nitrogen fixation by *Trichodesmium*
spp: An important source of new nitrogen to the tropical and subtropical North
Atlantic Ocean. Global Biogeochem Cycles 19:1–17.
4.   Hutchins DA, Fu FX, Zhang Y, Warner ME, Feng Y, Portune K, Bernhardt PW,
Mulholland MR. 2007. CO_2 control of *Trichodesmium* N_2 fixation, photosynthesis,
growth rates, and elemental ratios: Implications for past, present, and future ocean
biogeochemistry. Limnol Oceanogr 52:1293–1304.
5.   Thompson AW, Zehr JP. 2013. Cellular interactions: lessons from the nitrogen-fixing
cyanobacteria. J Phycol 49:1024–1035.
6.   Wang S, Koedooder C, Zhang F, Kessler N, Eichner M, Shi D, Shaked Y. 2022.
Colonies of the marine cyanobacterium *Trichodesmium* optimize dust utilization by
selective collection and retention of nutrient-rich particles. iScience 25:103587.

7. Kessler N, Armoza-Zvuloni R, Wang S, Basu S, Weber PK, Stuart RK, Shaked Y.
2019. Selective collection of iron-rich dust particles by natural *Trichodesmium*
colonies. The ISME Journal 2019 14:1 14:91–103.

8. Rubin M, Berman-Frank I, Shaked Y. 2011. Dust- and mineral-iron utilization by the
marine dinitrogen-fixer *Trichodesmium*. Nature Geoscience 2011 4:8 4:529–534.

9. Hmelo L, Van Mooy B, Mincer T. 2012. Characterization of bacterial epibionts on the
cyanobacterium Trichodesmium. Aquatic Microbial Ecology 67:1–14.

10. Rouco M, Haley ST, Dyhrman ST. 2016. Microbial diversity within the
Trichodesmium holobiont. Environ Microbiol 18:5151–5160.

11. Lee MD, Walworth NG, McParland EL, Fu FX, Mincer TJ, Levine NM, Hutchins DA,
Webb EA. 2017. The *Trichodesmium* consortium: Conserved heterotrophic co-
occurrence and genomic signatures of potential interactions. ISME Journal 11:1813–
1824.

12. Gradoville MR, Crump BC, Letelier RM, Church MJ, White AE. 2017. Microbiome of
Trichodesmium colonies from the North Pacific Subtropical Gyre. Front Microbiol 8.

13. Frischkorn KR, Rouco M, Van Mooy BAS, Dyhrman ST. 2017. Epibionts dominate
metabolic functional potential of *Trichodesmium* colonies from the oligotrophic ocean.
The ISME Journal 2017 11:9 11:2090–2101.

14. Frischkorn KR, Rouco M, Van Mooy BAS, Dyhrman ST. 2018. The *Trichodesmium*
microbiome can modulate host N_2 fixation. Limnol Oceanogr Lett 3:401–408.

15. Klawonn I, Eichner MJ, Wilson ST, Moradi N, Thamdrup B, Kümmel S, Gehre M,
Khalili A, Grossart HP, Karl DM, Ploug H. 2019. Distinct nitrogen cycling and steep
chemical gradients in *Trichodesmium* colonies. The ISME Journal 2019 14:2 14:399–
412.

16. Basu S, Gledhill M, de Beer D, Prabhu Matondkar SG, Shaked Y. 2019. Colonies of
marine cyanobacteria *Trichodesmium* interact with associated bacteria to acquire iron
from dust. Communications Biology 2019 2:1 2:1–8.

17. Frischkorn KR, Haley ST, Dyhrman ST. 2018. Coordinated gene expression between
Trichodesmium and its microbiome over day-night cycles in the North Pacific
Subtropical Gyre. ISME Journal 12:997–1007.

18. Frischkorn KR, Krupke A, Guieu C, Louis J, Rouco M, Salazar Estrada AE, Van
Mooy BAS, Dyhrman ST. 2018. *Trichodesmium* physiological ecology and phosphate
reduction in the western tropical South Pacific. Biogeosciences 15:5761–5778.

19. Gledhill M, Basu S, Shaked Y. 2019. Metallophores associated with *Trichodesmium*
erythraeum colonies from the Gulf of Aqaba. Metallomics 11:1547–1557.

20. Basu S, Shaked Y. 2018. Mineral iron utilization by natural and cultured
Trichodesmium and associated bacteria. Limnol Oceanogr 63:2307–2320.

21. Polyviou D, Baylay AJ, Hitchcock A, Robidart J, Moore CM, Bibby TS. 2018. Desert
dust as a source of iron to the globally important diazotroph *Trichodesmium*. Front
Microbiol 8:2683.

22. Manck LE, Park J, Tully BJ, Poire AM, Bundy RM, Dupont CL, Barbeau KA. 2021.
Petrobactin, a siderophore produced by *Alteromonas*, mediates community iron
acquisition in the global ocean. The ISME Journal 2021 16:2 16:358–369.

23. Berman-Frank I, Quigg A, Finkel Z V., Irwin AJ, Haramaty L. 2007. Nitrogen-fixation
strategies and Fe requirements in cyanobacteria. Limnol Oceanogr 52:2260–2269.

24. Jickells TD, An ZS, Andersen KK, Baker AR, Bergametti C, Brooks N, Cao JJ, Boyd
PW, Duce RA, Hunter KA, Kawahata H, Kubilay N, LaRoche J, Liss PS, Mahowald
N, Prospero JM, Ridgwell AJ, Tegen I, Torres R. 2005. Global iron connections
between desert dust, ocean biogeochemistry, and climate. Science (1979) 308:67–71.

25.　Kok JF, Ward DS, Mahowald NM, Evan AT. 2018. Global and regional importance of
the direct dust-climate feedback. Nature Communications 9:1 9:1–11.
26.　Post AF, Dedej Z, Gottlieb R, Li H, Thomas DN, El-Absawi M, El-Naggar A, El-
Gharabawi M, Sommer U. 2002. Spatial and temporal distribution of *Trichodesmium*
spp. in the stratified Gulf of Aqaba, Red Sea. Mar Ecol Prog Ser 239:241–250.
27.　Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy TBK,
Schulz F, Jarett J, Rivers AR, Eloe-Fadrosh EA, Tringe SG, Ivanova NN, Copeland A,
Clum A, Becraft ED, Malmstrom RR, Birren B, Podar M, Bork P, Weinstock GM,
Garrity GM, Dodsworth JA, Yooseph S, Sutton G, Glöckner FO, Gilbert JA, Nelson
WC, Hallam SJ, Jungbluth SP, Ettema TJG, Tighe S, Konstantinidis KT, Liu WT,
Baker BJ, Rattei T, Eisen JA, Hedlund B, McMahon KD, Fierer N, Knight R, Finn R,
Cochrane G, Karsch-Mizrachi I, Tyson GW, Rinke C, Lapidus A, Meyer F, Yilmaz P,
Parks DH, Eren AM, Schriml L, Banfield JF, Hugenholtz P, Woyke T. 2017.
Minimum information about a single amplified genome (MISAG) and a metagenome-
assembled genome (MIMAG) of bacteria and archaea. Nature Biotechnology 35:8
35:725–731.
28.　Koedooder C, Landou E, Zhang F, Wang S, Basu S, Berman-Frank I, Shaked Y,
Rubin-Blum M. 2022. Metagenomes of Red Sea subpopulations challenge the use of
marker genes and morphology to assess *Trichodesmium* diversity. Front Microbiol
13:1712.
29.　Dang H, Lovell CR. 2000. Bacterial primary colonization and early succession on
surfaces in marine waters as determined by amplified rRNA gene restriction analysis
and sequence analysis of 16S rRNA Genes. Appl Environ Microbiol 66:467.
30.　Dang H, Li T, Chen M, Huang G. 2008. Cross-ocean distribution of Rhodobacterales
bacteria as primary surface colonizers in temperate coastal marine waters. Appl
Environ Microbiol 74:52–60.
31.　Duret MT, Lampitt RS, Lam P. 2019. Prokaryotic niche partitioning between
suspended and sinking marine particles. Environ Microbiol Rep 11:386–400.
32.　Koch H, Germscheid N, Freese HM, Noriega-Ortega B, Lücking D, Berger M, Qiu G,
Marzinelli EM, Campbell AH, Steinberg PD, Overmann J, Dittmar T, Simon M, Wietz
M. 2020. Genomic, metabolic, and phenotypic variability shapes ecological
differentiation and intraspecies interactions of *Alteromonas macleodii*. Scientific
Reports 2020 10:1 10:1–14.
33.　Hou S, López-Pérez M, Pfreundt U, Belkin N, Stüber K, Huettel B, Reinhardt R,
Berman-Frank I, Rodriguez-Valera F, Hess WR. 2018. Benefit from decline: The
primary transcriptome of *Alteromonas macleodii* str. Te101 during *Trichodesmium*
demise. ISME Journal 12:981–996.
34.　Held NA, Webb EA, McIlvin MM, Hutchins DA, Cohen NR, Moran DM, Kunde K,
Lohan MC, Mahaffey C, Malcolm E, Saito MA. 2020. Co-occurrence of Fe and P
stress in natural populations of the marine diazotroph *Trichodesmium*. Biogeosciences
17:2537–2551.
35.　Chappell PD, Webb EA. 2010. A molecular assessment of the iron stress response in
the two phylogenetic clades of *Trichodesmium*. Environ Microbiol 12:13–27.
36.　Carroll CS, Moore MM. 2018. Ironing out siderophore biosynthesis: a review of non-
ribosomal peptide synthetase (NRPS)-independent siderophore synthetases. Crit Rev
Biochem Mol Biol. Taylor and Francis Ltd
https://doi.org/10.1080/10409238.2018.1476449.
37.　Gulick AM. 2017. Nonribosomal peptide synthetase biosynthetic clusters of ESKAPE
pathogens. Nat Prod Rep 34:981–1009.

38. Reitz ZL, Medema MH. 2022. Genome mining strategies for metallophore discovery.
Curr Opin Biotechnol 77:102757.

39. Butler A, Harder T, Ostrowski AD, Carrano CJ. 2021. Photoactive siderophores:
structure, function and biology. J Inorg Biochem 221:111457.

40. Amin SA, Green DH, Küpper FC, Carrano CJ. 2009. Vibrioferrin, an unusual marine
siderophore: iron binding, photochemistry, and biological implications. Inorg Chem
48:11451–11458.

41. Amin SA, Green DH, Hart MC, Küpper FC, Sunda WG, Carrano CJ. 2009. Photolysis
of iron-siderophore chelates promotes bacterial-algal mutualism. Proc Natl Acad Sci U
S A 106:17071–17076.

42. Cordero OX, Ventouras LA, DeLong EF, Polz MF. 2012. Public good dynamics drive
evolution of iron acquisition strategies in natural bacterioplankton populations. Proc
Natl Acad Sci U S A 109:20059–20064.

43. Barbeau K, Rue EL, Trick CG, Bruland KW, Butler A. 2003. Photochemical reactivity
of siderophores produced by marine heterotrophic bacteria and cyanobacteria based on
characteristic Fe(III) binding groups. Limnol Oceanogr 48:1069–1078.

44. Martinez JS, Carter-Franklin JN, Mann EL, Martin JD, Haygood MG, Butler A. 2003.
Structure and membrane affinity of a suite of amphiphilic siderophores produced by a
marine bacterium. Proc Natl Acad Sci U S A 100:3754–3759.

45. Kramer J, Özkaya Ö, Kümmerli R. 2020. Bacterial siderophores in community and
host interactions. Nat Rev Microbiol. Nature Research https://doi.org/10.1038/s41579-
019-0284-4.

46. Van Mooy BAS, Krupke A, Dyhrman ST, Fredricks HF, Frischkorn KR, Ossolinski
JE, Repeta DJ, Rouco M, Seewald JD, Sylva SP. 2015. Phosphorus cycling: a major
role of planktonic phosphate reduction in the marine phosphorus redox cycle. Science
348:783–785.

47. Mackey KRM, Labiosa RG, Calhoun M, Street JH, Post AF, Paytan A. 2007.
Phosphorus availability, phytoplankton community dynamics, and taxon-specific
phosphorus status in the Gulf of Aqaba, Red Sea. Limnol Oceanogr 52:873–885.

48. Karl DM, Björkman KM. 2015. Dynamics of dissolved organic phosphorus.
Biogeochemistry of marine dissolved organic matter: Second Edition 233–334.

49. Orchard ED, Webb EA, Dyhrman ST. 2009. Molecular analysis of the phosphorus
starvation response in *Trichodesmium* spp. Environ Microbiol 11:2400–2411.

50. Frischkorn KR, Krupke A, Guieu C, Louis J, Rouco M, Salazar Estrada AE, Van
Mooy BAS, Dyhrman ST. 2018. *Trichodesmium* physiological ecology and phosphate
reduction in the western tropical South Pacific. Biogeosciences 15:5761–5778.

51. Frischkorn KR, Haley ST, Dyhrman ST. 2019. Transcriptional and proteomic
choreography under phosphorus deficiency and re-supply in the N2 fixing
cyanobacterium *Trichodesmium erythraeum*. Front Microbiol 10:330.

52. Stosiek N, Talma M, Klimek-Ochab M. 2020. Carbon-Phosphorus Lyase—the State of
the Art. Appl Biochem Biotechnol 190:1525–1552.

53. Kim AD, Baker AS, Dunaway-Mariano D, Metcalf WW, Wanner BL, Martin BM.
2002. The 2-aminoethylphosphonate-specific transaminase of the 2-
aminoethylphosphonate degradation pathway. J Bacteriol 184:4134–4140.

54. Lockwood S, Greening C, Baltar F, Morales SE. 2022. Global and seasonal variation
of marine phosphonate metabolism. The ISME Journal 2022 16:9 16:2198–2212.

55. Hilderbrand RL. 2018. The effects of synthetic phosphonates on living systems: the
role of phosphonates in living systems 139–169.

56. Coates CJ, Wyman M. 2017. A denitrifying community associated with a major,
marine nitrogen fixer. Environ Microbiol 19:4978–4992.

57. Eichner M, Basu S, Wang S, de Beer D, Shaked Y. 2020. Mineral iron dissolution in
Trichodesmium colonies: The role of O2 and pH microenvironments. Limnol
Oceanogr 65:1149–1160.
58. Pajares S, Ramos R. 2019. Processes and microorganisms involved in the marine
nitrogen cycle: knowledge and gaps. Front Mar Sci 6:739.
59. Graf DRH, Jones CM, Hallin S. 2014. Intergenomic comparisons highlight modularity
of the denitrification pathway and underpin the importance of community structure for
N$_2$O Emissions. PLoS One 9:e114118.
60. Kreimer A, Borenstein E, Gophna U, Ruppin E. 2008. The evolution of modularity in
bacterial metabolic networks. Proc Natl Acad Sci U S A 105:6976–6981.
61. Hutchins DA, Capone DG. 2022. The marine nitrogen cycle: new developments and
global change. Nature Reviews Microbiology 2022 20:7 20:401–414.
62. Smriga S, Ciccarese D, Babbin AR. 2021. Denitrifying bacteria respond to and shape
microscale gradients within particulate matrices. Communications Biology 2021 4:1
4:1–9.
63. Stief P, Kamp A, Thamdrup B, Glud RN. 2016. Anaerobic nitrogen turnover by
sinking diatom aggregates at varying ambient oxygen levels. Front Microbiol 7:98.
64. Paerly HW, Bebout BM, Prufert LE. 1989. Bacteria associations with marine
Oscillatoria sp. (*Trichodesmium* sp.) populations: ecophysiological implications. J
Phycol 25:773–784.
65. McGenity TJ. 2019. Taxonomy, genomics and ecophysiology of hydrocarbon-
degrading microbes. Springer International Publishing. Retrieved 18 January 2023.
66. Ho A, Di Lonardo DP, Bodelier PLE. 2017. Revisiting life strategy concepts in
environmental microbial ecology. FEMS Microbiol Ecol 93:6.
67. Lauro FM, McDougald D, Thomas T, Williams TJ, Egan S, Rice S, DeMaere MZ,
Ting L, Ertan H, Johnson J, Ferriera S, Lapidus A, Anderson I, Kyrpides N, Munk AC,
Detter C, Han CS, Brown M V., Robb FT, Kjelleberg S, Cavicchioli R. 2009. The
genomic basis of trophic strategy in marine bacteria. Proc Natl Acad Sci U S A
106:15527–33.
68. Zoccarato L, Sher D, Miki T, Segrè D, Grossart HP. 2022. A comparative whole-
genome approach identifies bacterial traits for marine microbial interactions.
Communications Biology 2022 5:1 5:1–13.
69. Debeljak P, Blain S, Bowie A, van der Merwe P, Bayer B, Obernosterer I. 2021.
Homeostasis drives intense microbial trace metal processing on marine particles.
Limnol Oceanogr 66:3842–3855.
70. Van Mooy BAS, Hmelo LR, Sofen LE, Campagna SR, May AL, Dyhrman ST,
Heithoff A, Webb EA, Momper L, Mincer TJ. 2011. Quorum sensing control of
phosphorus acquisition in *Trichodesmium* consortia. The ISME Journal 2012 6:2
6:422–429.
71. Azam. F., Fenchel T, Field JG, Azam F, Field JG, Graf JS, Meyer-Rei LA, Thingstad
F. 1983. The Ecological Role of Water-Column Microbes in the Sea. Ecol Prog Ser,
Mar 10:257–263.
72. Kieser S, Brown J, Zdobnov EM, Trajkovski M, McCue LA. 2020. ATLAS: a
Snakemake workflow for assembly, annotation, and genomic binning of metagenome
sequence data. BMC Bioinformatics 21.
73. B. Bushnell. BBTools software package. sourceforge.net/projects/bbmap/. Retrieved
16 December 2021.
74. Bushnell B, Rood J, Singer E. 2017. BBMerge – Accurate paired shotgun read
merging via overlap. PLoS One 12:e0185056.

75.   Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a new
917   versatile metagenomic assembler. Genome Res 27:824–834.
918   76.   Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, Wang Z. 2019. MetaBAT 2: An
919   adaptive binning algorithm for robust and efficient genome reconstruction from
920   metagenome assemblies. PeerJ 2019:e7359.
921   77.   Wu YW, Simmons BA, Singer SW. 2016. MaxBin 2.0: an automated binning
922   algorithm to recover genomes from multiple metagenomic datasets. Bioinformatics
923   32:605–607.
924   78.   Nissen JN, Johansen J, Allesøe RL, Sønderby CK, Armenteros JJA, Grønbech CH,
925   Jensen LJ, Nielsen HB, Petersen TN, Winther O, Rasmussen S. 2021. Improved
926   metagenome binning and assembly using deep variational autoencoders. Nat
927   Biotechnol 39:555–560.
928   79.   Chklovski A, Parks DH, Woodcroft BJ, Tyson GW. 2022. CheckM2: a rapid, scalable
929   and accurate tool for assessing microbial genome quality using machine learning.
930   bioRxiv 2022.07.11.499243.
931   80.   Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, Banfield JF.
932   2018. Recovery of genomes from metagenomes via a dereplication, aggregation and
933   scoring strategy. Nature Microbiology 2018 3:7 3:836–843.
934   81.   Olm MR, Brown CT, Brooks B, Banfield JF. 2017. dRep: a tool for fast and accurate
935   genomic comparisons that enables improved genome recovery from metagenomes
936   through de-replication. The ISME Journal 2017 11:12 11:2864–2868.
937   82.   Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal:
938   Prokaryotic gene recognition and translation initiation site identification. BMC
939   Bioinformatics 11:1–11.
940   83.   Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. 2020. GTDB-Tk: a toolkit to
941   classify genomes with the Genome Taxonomy Database. Bioinformatics 36:1925–
942   1927.
943   84.   Lee MD. 2019. GToTree: a user-friendly workflow for phylogenomics. Bioinformatics
944   35:4162–4164.
945   85.   Kalyaanamoorthy S, Minh B, Wong T, von Haeseler A, Jermiin L. ModelFinder: fast
946   model selection for accurate phylogenetic estimates. Nat Methods 14:587–589.
947   86.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2014. IQ-TREE: A fast and
948   effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol
949   Biol Evol 32:268–274.
950   87.   Coleman GA, Davín AA, Mahendrarajah TA, Szánthó LL, Spang A, Hugenholtz P,
951   Szöllsi GJ, Williams TA. 2021. A rooted phylogeny resolves early bacterial evolution.
952   Science (1979) 372.
953   88.   Kanehisa M, Sato Y, Morishima K. 2016. BlastKOALA and GhostKOALA: KEGG
954   Tools for Functional Characterization of Genome and Metagenome Sequences. J Mol
955   Biol 428:726–731.
956   89.   Bateman A, Coin L, Durbin R, Finn RD, Hollich V, Griffiths-Jones S, Khanna A,
957   Marshall M, Moxon S, Sonnhammer ELL, Studholme DJ, Yeats C, Eddy SR. 2004.
958   The Pfam protein families database. Nucleic Acids Res 32:D138–D141.
959   90.   Aziz RK, Bartels D, Best A, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S,
960   Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA,
961   McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R,
962   Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST Server: Rapid
963   annotations using subsystems technology. BMC Genomics 9:1–15.
964   91.   Meyer F, Overbeek R, Rodriguez A. 2009. FIGfams: yet another set of protein
965   families. Nucleic Acids Res 37:6643–6654.

92. Zhou Z, Tran PQ, Breister AM, Liu Y, Kieft K, Cowley ES, Karaoz U, Anantharaman K. 2022. METABOLIC: high-throughput profiling of microbial genomes for functional traits, metabolism, biogeochemistry, and community-scale functional networks. Microbiome 10:33.

93. Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. Nucleic Acids Res 49: W293–W296.

94. Garber AI, Nealson KH, Okamoto A, McAllister SM, Chan CS, Barco RA, Merino N. 2020. FeGenie: A comprehensive tool for the identification of iron genes and iron gene neighborhoods in genome and metagenome assemblies. Front Microbiol 11:37.

95. Katoh K, Kuma KI, Toh H, Miyata T. 2005. MAFFT version 5: improvement in accuracy of multiple sequence alignment. Nucleic Acids Res 33:511–518.

96. Kautsar SA, Blin K, Shaw S, Navarro-Muñoz JC, Terlouw BR, van der Hooft JJJ, van Santen JA, Tracanna V, Suarez Duran HG, Pascal Andreu V, Selem-Mojica N, Alanjary M, Robinson SL, Lund G, Epstein SC, Sisto AC, Charkoudian LK, Collemare J, Linington RG, Weber T, Medema MH. 2019. MIBiG 2.0: a repository for biosynthetic gene clusters of known function. Nucleic Acids Res 48: D454–D458.

97. Kim S, Gupta N, Pevzner PA. 2008. Spectral probabilities and generating functions of tandem mass spectra: A strike against decoy databases. J Proteome Res 7:3354–3363.

98. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–2069.

99. Florens L, Carozza MJ, Swanson SK, Fournier M, Coleman MK, Workman JL, Washburn MP. 2006. Analyzing chromatin remodeling complexes using shotgun proteomics and normalized spectral abundance factors. Methods 40:303–311.

FIGURES

Figure 1. Phylogenetic tree of the 52 MAGs assembled from *Trichodesmium* colonies collected in the Red Sea. The tree is rooted by the outgroup *Fuseobacterium nucleatum*. Known bacterial orders are marked in different colors and several MAGs could taxonomically be identified at the genus level. The relative abundance of each MAG within the *Trichodesmium* consortium, excluding the abundance of *Trichodesmium* MAG 52 (±70% of samples), is presented as a bubble chart. The 11 MAGs that matched to MAGs assembled from a previous metagenomic dataset of Red Sea *Trichodesmium* colonies (28) are highlighted in blue and tentatively represent the core-consortium (see Supplementary Figure 2; Supplementary Table 4). MAGs that did not meet the requirements of a high-quality genome (>90% completeness; <5% redundancy) are indicated in grey. Note that re-occurring MAGs are not necessarily the most abundant MAGs present within the *Trichodesmium* colony. (a full taxonomic description of each MAG in this dataset can be found in Supplementary Table 1).

Figure 2. MAGs were probed for Fe-uptake and siderophore biosynthesis genes which are noted in Supplementary Table 5. **(a)** *T. thiebautii* MAG 52 does not contain a wide diversity of Fe-uptake genes, and the majority of MAGs of associated bacteria contain an uptake receptor for organic Fe (heme, citrate, siderophores). Several MAGs of associated bacteria contained an NRPS or NIS-type siderophore biosynthesis pathways (BGCs). **(b)** Siderophore biosynthetic gene clusters (BGCs) of known NIS-type siderophore BGCs can be clustered according to the presence of one or multiple iucA/iucC domains which enables one to target the closest siderophore BGC match (36). Using this method, we were able to link these pathways to the synthesis of the photolabile **(i)** petrobactin, **(ii)** vibrioferrin and **(iii)** rhizoferrin. (c) Siderophore BGCs of known NRPS siderophores. Using antiSMASH 7.0, which aligns BGCs to known siderophores within the MIBiG database, we were able to link 2 of the 6

putative siderophore NRPS-type pathways to **(i)** the membrane-bound amphibactin and **(ii)** agrobactin/enterobactin. (See Supplementary Figure 3 for more details on the siderophore biosynthesis pathways).

Figure 3. MAGs were probed for several different P-metabolism genes which are noted in Supplementary Table 5. *Trichodesmium* is known to be highly competitive for P, as exemplified by the multitude of **(a)** P-uptake systems for phosphate (*pst*S, *pst*B), phosphonate (*phn*CDE), and phosphonate/phosphite (*phn*CDE/*ptxABC*). **(b)** Within the *Trichodesmium* consortium, the ability to metabolize a range of phosphonate compounds using a broad specificity C-P lyase (*phn*GHIJKLM), is more prominent than the substrate-specific hydrolysis of 2-aminoethylphosphonate (2-AEP; *phn*WXYZA) or phosphonopyruvate (*pal*A) and phosphite metabolism (*ptx*D).

Figure 4. MAGs were probed for N-metabolic pathways and the genes for each step within the pathway are noted in Supplementary Table 5. *T. thiebautii* MAG 52 is the only MAG with a N_2 fixation pathway. Several MAGs contained one or two steps within the pathways of denitrification or dissimilatory nitrate to reduction to ammonia (DNRA) indicating their modularity. Collectively, MAGs completed the denitrification and DNRA pathways, but none contained the genes for the entire pathway.

Figure 5. MAGs were probed for vitamin B1, B7, and B12 biosynthesis and uptake pathways. The genes used to indicate the biosynthesis or uptake of these vitamins are noted in Supplementary Table 5. *Trichodesmium* contains the biosynthesis pathways for all three vitamins but the vast majority of associated bacteria were auxotrophic for one or more vitamins. These results suggest that the associated bacteria rely on *Trichodesmium* to obtain their vitamin requirements.

Figure 6. Schematic summary of the potential interactions taking place within the Trichodesmium consortia as discussed in this study. Genes for each trait are noted in Supplementary Table 5. Each blue bubble indicates a different function, and each blue circle pools together the different MAGs containing a particular trait for that function. Each MAG is represented by a colored dot reflective of the taxonomic order it is associated with (grey dots represent MAGs that were not attributed to a specific taxonomic order). *Trichodesmium* is highlighted in cobalt-blue, and the important functions *Trichodesmium* is known to display are highlighted with a dotted circle. (See Supplementary Table 5 for the full list of functional traits and their presence/absence). **(a)** Several siderophore biosynthesis genes are present in a diverse set of MAGs. Siderophores are predicted to mine Fe from dust that has been collected and centered within *Trichodesmium* colonies. 6 of the 4 were identified as known NIS-type siderophores of which petrobactin, vibrioferrin and rhizoferrin are photolabile. In the presence of sunlight, the Fe complexed to these siderophores disassociates and acts as a public good to the larger consortium. The genes and pathways for **(b-i)** denitrification, **(b-ii)** N_2 fixation, and **(b-iii)** dissimilatory- nitrate- reduction- to- nitrate (DNRA) are present in several MAGs of the consortium. These processes appear to be modular within the *Trichodesmium* consortium as different taxonomic groups complete the steps for these pathways. **(c)** Several MAGs contain a homoserine lactone synthesis pathway (HL), including *Trichodesmium*. HL acts as an intercommunicating quorum sensing molecule and has been shown to influence the production of alkaline phosphatase (AP) and N_2 fixation in *Trichodesmium*. (associated bacteria can also produce AP but is not depicted here - see Figure 3). **(d)** *Trichodesmium* is capable of producing the vitamins cobalamin (B12), thiamin (B1) and biotin (B7). Different subsets of MAGs lack the biosynthesis pathways for one or more of these vitamins with very few capable of

synthesizing all three (associated bacteria can also produce vitamins but is not depicted here - see Figure 5). Consortia members and *Trichodesmium* likely exhibit an interdependent relationship for these vitamins. **(e)** *Trichodesmium* colonies have been shown to take up and utilize phosphonate and phosphite in a low-P environment and MAGs containing these pathways may have a competitive advantage over members that lack them.

SUPPLEMENTARY FIGURES

Supplementary Figure 1. Relative abundance of bacterial MAGs from sampled *Trichodesmium* colonies indicate the associated bacterial distribution is consistent. The 51 MAGs are stacked, and colors reflect the different bacterial orders each MAG belongs to and 'other' in grey. The exact relative abundance value (%) for each MAG and its standard deviation can be viewed in Supplementary Table 1.

Supplementary Figure 2. Phylogenetic tree of the 52 MAGs together with MAGs assembled from 3 other Trichodesmium metagenomic datasets from colonies collected in the Red Sea (28), the Pacific (12, 17) and the Atlantic (13) Oceans. MAGs from the 3 other datasets are highlighted in grey. MAGs deriving from samples of the Atlantic or Pacific Ocean are listed as MAG T-XX and those from the Red Sea are listed as MAG R-XX. The 11 MAGs that matched to those assembled from previous metagenomic datasets, using a 97.5 % identity cutoff, are highlighted in blue. Known bacterial orders are marked using different colored squares. The tree is rooted by the outgroup *Fuseobacterium nucleatum*. The full taxonomic description of each MAG in this dataset and others can be found in Supplementary Table 4.

Supplementary Figure 3. (a) NIS and NRPS-type siderophore biosynthesis pathways identified in MAGs of associated bacteria. (b) The siderophore BGC analysis tools used to probe and identify the different siderophore BGCs. Most tools are limited in their ability to discover siderophore BGCs while antiSMASH 7.0 was the most conclusive but also time-consuming (53). **(c)** Putative NRPS-type siderophore BGCs that did not match to a known siderophore within the MIBiG database but did containing a TonB-siderophore receptor (SMCOG 1208). This made us speculate that these are putative siderophores whose identity remains to be resolved.

Supplementary Figure 4. (a) MAGs were probed for bacterial-interactive traits (68)**.** MAGs from the phyla Proteobacter, contain the presence several interactive traits, including quorum sensing, gliding motility, adhesion, cell-to-cell signaling, and antibiotic resistance marker genes. Intriguingly, both the Trichodesmium MAG and several Proteobacter MAGs (e.g. Rhodobacter) contain genes for homoserine lactones (HLs), which are putative quorum sensing molecules. **(b) MAGs were probed for particle-interactive traits** (69)**.** In comparison to *T. thiebautii* MAG 52, many MAGs from associated bacteria contained a wide variety of heavy metal efflux systems and metal-transcription factors. While *Trichodesmium*, like most Cyanobacteria, is known to be sensitive to heavy metals, our results indicate that the consortium is attuned towards an environment rich in metals, including particulate dust. Genes for each trait are noted in Supplementary Table 5

SUPPLEMENTARY TABLES

Supplementary Table 1. General information about the MAGs making up the Red Sea *Trichodesmium* consortium. The relative abundance of associated bacteria (%) and standard

deviation of the triplicates (SD) excludes the abundance of *T. thiebautii* (MAG 52). The relative
abundance of MAGs was consistent across our samples. MAGs that did not meet the high-
quality standard of 90% completeness and <5% redundancy are marked in green (10 of 52) and
the functional analysis from these MAGs needs to be taken with more caution. The % of the
genomes annotated by RAST is also indicated, showcasing how MAGs from the phyla
Bacteroidetes are marked with a low annotation percentage. The closest taxonomic identity is
given using both NCBI and GTDB-taxonomy. Throughout this paper we use GTDB-taxonomy
except for the order Rhodospirillales.

Supplementary Table 2: Top proteomic hits matching to *T. thiebautii* MAG52. Spectrum
counts that mapped to KEGG ID numbers mentioned in Supplementary Table 5 are highlighted
in blue. Most proteins are geared towards the processes of photosynthesis, nitrogen fixation,
carbon metabolism, amino acid uptake, protein synthesis, and phosphate uptake. Total spectral
counts (T-total) and their associated KEGG or PFAM ID number and their description are
presented together with the average relative contribution (T0%) of each protein hit to all
Trichodesmium-related proteins and for all three samples (%T1-3). Collectively, the top 50
proteins represented >50% of the total spectral counts.

Supplementary Table 3: Top proteomic hits matching to MAGs of *Trichodesmium* associated
bacteria. Many proteins are geared towards a typical heterotrophic and copiotrophic lifestyle
including proteins related to amino-acid uptake and carbon metabolism are highlighted in bold.
Total spectral counts (%) and their associated KEGG or PFAM ID numbers and their
description are presented together with the average relative contribution (Avg%) of each
protein hit to all Trichodesmium-related proteins and for all three samples (%T1-3).

Supplementary Table 4. Taxonomic identity of MAGs used in this study are compared to
MAGs assembled from other metagenomic studies of *Trichodesmium* colonies (Red Sea
samples: MAG R-XX (28), the Pacific (12, 17) and Atlantic samples (13): MAG T-XX. Their
closest taxonomic identity is given using both NCBI and GTDB-taxonomy. Throughout this
paper we use GTDB-taxonomy except for the order Rhodospirillales. Highlighted in blue are
the 11 MAGs from this dataset that matched to those assembled from these other datasets (in
bold), using a 97.5 % identity cutoff. The phylogenetic tree of all MAGs from all datasets
combined can be seen in Supplementary Figure 2.

Supplementary Table 5. The presence and absence of functional traits separated into larger
groups including Fe-uptake, P-metabolism, N-metabolism, Vitamin B synthesis and uptake,
metal regulators, metal efflux, metal storage, signaling, secretion systems. motility and
adhesion genes. The associated analysis tool, KEGG id number or module is given for each
trait. The total number of MAGs containing a specific trait and (MAGs %) is calculated and
show below. Traits that were detected within our proteomic samples are also highlighted in
blue.

1155
1156 **Supplementary Figure 1. Relative abundance of bacterial MAGs from sampled *Trichodesmium* colonies**
1157 **indicate the associated bacterial distribution is consistent.** The 51 MAGs are stacked, and colors reflect the
1158 different bacterial orders each MAG belongs to and 'other' in grey. The exact relative abundance value (%) for
1159 each MAG and its standard deviation can be viewed in Supplementary Table 1.
1160

1161
Phylogenetic tree of the 52 MAGs together with MAGs assembled from 3 other *Trichodesmium*
metagenomic datasets from colonies collected in the Red Sea (28)**, the Pacific and the Atlantic Ocean.** MAGs
from the 3 other datasets are highlighted in grey. MAGs deriving from samples of the Atlantic or Pacific Ocean
are listed as MAG T-XX and those from the Red Sea are listed as MAG R-XX. The 11 MAGs that matched to
those assembled from previous metagenomic datasets, using a 97.5 % identity cutoff, are highlighted in blue.
Known bacterial orders are marked using different colored squares. The tree is rooted by the outgroup
Fuseobacterium nucleatum. The full taxonomic description of each MAG in this dataset and others can be found
in Supplementary Table 4.
1170

Supplementary Figure 3. (a) NIS and NRPS-type siderophore biosynthesis pathways identified in MAGs of associated bacteria. **(b)** The siderophore BGC analysis tools used to probe and identify the different siderophore BGCs. Most tools are limited in their ability to discover siderophore BGCs while antiSMASH 7.0 was the most conclusive but also time-consuming. **(c)** Putative NRPS-type siderophore BGCs that did not match to a known siderophore within the MIBiG database but did containing a TonB-siderophore receptor (SMCOG 1208). This made us speculate that these are putative siderophores whose identity remains to be resolved.

Supplementary Figure 4. (a) MAGs were probed for bacterial-interactive traits (68). MAGs from the phyla Proteobacter, contain the presence several interactive traits, including quorum sensing, gliding motility, adhesion, cell-to-cell signaling, and antibiotic resistance marker genes. Intriguingly, both the Trichodesmium MAG and several Proteobacter MAGs (e.g. Rhodobacter) contain genes for homoserine lactones (HLs), which are putative quorum sensing molecules. **(b)** MAGs were probed for particle-interactive traits (69). In comparison to T. thiebautii MAG 52, many MAGs from associated bacteria contained a wide variety of heavy metal efflux systems and metal-transcription factors. While *Trichodesmium*, like most Cyanobacteria, is known to be sensitive to heavy metals, our results indicate that the consortium is attuned towards an environment rich in metals, including particulate dust. Genes for each trait are noted in Supplementary Table 5.