An ancestral SARS-CoV-2 vaccine induces anti-Omicron variants antibodies by hypermutation

Running Title: Hypermutation and antibody specificity

Seoryeong Park[1,2*], Jaewon Choi[3,4*], Yonghee Lee[5], Jinsung Noh[5,6], Namphil Kim[5], JinAh Lee[7], Geummi Cho[1,8], Sujeong Kim[1,8], Duck Kyun Yoo[1,8], Chang Kyung Kang[9], Pyoeng Gyun Choe[9], Nam Joong Kim[9], Wan Beom Park[9], Seungtaek Kim[7], Myoung-don Oh[9], Sunghoon Kwon[3,5,6], Junho Chung[1,2,8]

[1]Department of Biochemistry and Molecular Biology, Seoul National University College of Medicine, Seoul, Republic of Korea

[2]Interdisciplinary Program in Cancer Biology Major, Seoul National University College of Medicine, Seoul, Republic of Korea

[3]Interdisciplinary Program in Bioengineering, Seoul National University, Seoul, Republic of Korea

[4]Integrated Major in Innovative Medical Science, Seoul National University, Seoul, Republic of Korea

[5]Department of Electrical and Computer Engineering, Seoul National University, Seoul, Republic of Korea

[6]Bio-MAX Institute, Seoul National University, Seoul, Republic of Korea

[7]Zoonotic Virus Laboratory, Institut Pasteur Korea, Seongnam, Republic of Korea

[8]Department of Biomedical Science, Seoul National University College of Medicine, Seoul,

Republic of Korea

[9]Department of Internal Medicine, Seoul National University College of Medicine, Seoul,

Republic of Korea

*These authors contributed equally to this work.

Corresponding author

S. Kim (seungtaek.kim@ip-korea.org), M. Oh (mdohmd@snu.ac.kr), S. Kwon

(skwon@snu.ac.kr) or J. Chung (jjhchung@snu.ac.kr).

The immune escape of Omicron variants significantly subsides by the third dose of an

mRNA vaccine. However, it is unclear how Omicron variant-neutralizing antibodies

develop under repeated vaccination. We analyzed blood samples from 41 BNT162b2

vaccinees following the course of three injections and analyzed their B-cell receptor (BCR)

repertoires at six time points in total. The concomitant reactivity to both ancestral and

Omicron receptor-binding domain (RBD) was achieved by a limited number of BCR

clonotypes depending on the accumulation of somatic hypermutation (SHM) after the third

dose. Our findings suggest that SHM accumulation in the BCR space to broaden its

specificity for unseen antigens is a counterprotective mechanism against virus variant

immune escape.

Main

Since the emergence of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), over 14

million sequences of variants have been collected and shared via the Global Initiative on Sharing

All Influenza Data (GISAID) [1]. While most mutations have little effect or are detrimental to the

virus, a small subset of mutations may provide a selective advantage leading to a higher

reproductive rate[2]. The spike protein, a viral coat protein that mediates viral attachment to host

cells and fusion between the virus and the cell membrane, is the primary target of neutralizing

antibodies[3]. Serological analysis has shown that the receptor-binding domain (RBD) of the spike

protein is the target of 90% of neutralizing activity in the immune sera[4,5]. In this context, the

RBD has become the essential component of most mRNA-, adenovirus-, and recombinant

protein-based vaccines[6]. However, Omicron variant BA.1 has accumulated 15 mutations in

RBD[7], resulting in a 22-fold reduction in neutralization by plasma from vaccinees receiving two

doses of the BNT162b2 vaccine[8]. This has led to reduced efficacy of most commercially

available monoclonal antibodies and antibodies under development against BA.1[7,9]. Although

bivalent vaccines have been developed to overcome the immune evasion of Omicron variant[10-13],

the majority of the population has received only monovalent vaccines to date. Fortunately, it has

been proven that a third dose of the BNT162b2 monovalent vaccine can neutralize BA.1[14-18].

However, the mechanism by which Omicron variant-neutralizing antibodies are generated

through repeated exposure to the ancestral spike protein remains unclear. In this study, we

analyzed the chronological B-cell receptor (BCR) repertoires of BNT162b2 vaccinees and traced

the development of Omicron variant-neutralizing antibodies.

Vaccination and plasma antibody levels

We followed 41 healthcare workers from Seoul National University Hospital who received two doses of the BNT162b2 mRNA vaccine with a three-week interval and a third dose at approximately nine months after the first dose (Fig. 1a). Peripheral blood samples were collected six times: once before the first dose, three times after each dose, and two times between the second and third doses. The vaccinees did not report a history of COVID-19 infection, and the absence of plasma IgG against the nucleocapsid (N) protein of SARS-CoV-2 was confirmed in all vaccinees (Extended Data Fig. 1). In an enzyme-linked immunosorbent assay (ELISA), plasma levels of IgA and IgG against the ancestral RBD were significantly increased after the second dose (Fig. 1b and Extended Data Fig. 2). However, when vaccination ELISA was performed against BA.1 RBD, a third dose was essential to achieve elevated levels of IgA and IgG, which also reacted to the RBD of Omicron subvariant BQ.1.1 (Fig. 1b, Extended Data Figs. 2 and 3). This observation was in accord with prior reports that neutralizing activity of plasma against the Omicron variant was detectable only after the third dose[19,20], while the second dose could induce Omicron variant-neutralizing memory B cells[21].

Selection and characterization of BA.1 RBD-binding antibody clones

We selected six vaccinees and generated a single-chain variable fragment (scFv) phage display library using cDNA prepared from the sixth blood sample. For the selection of these vaccinees, we reconstituted the *in silico* chronological BCR repertoires of vaccinees by next-generation sequencing (Supplementary Table 1) and compared the frequency of BCR heavy chain (HC) clonotypes listed in the CoV-Ab Dab[22] database binding the SARS-CoV-2 spike protein. Vaccinee 32, 35, 39 and 43 and vaccine 22 and 27 showed a higher frequency of clonotypes

binding to the BA.1 and ancestral spike proteins, respectively, and were selected. In this study,

the BCR HC clonotype was defined as a set of immunoglobulin heavy chain sequences that

exhibit the same immunoglobulin heavy variable (IGHV) and immunoglobulin heavy joining

(IGHJ) genes and homologous heavy chain complementarity-determining region 3 (HCDR3)

amino acid sequences with a minimum sequence identity of 80% to a reference sequence.

From six libraries, nine BA.1 RBD-binding scFv clones were selected based on biopanning and

phage ELISA. A cluster of scFv clones composed of 27-60 clones and 51 other clones sharing

IGHV3-53/3-66, IGHJ6, and the same HCDR sequences at the amino acid level was selected

from five libraries (Table 1 and Supplementary Table 2). We have previously reported that an

IGHV3-53/3-66 and IGHJ6 BCR HC clonotype is present not only in the majority of

convalescent patients from ancestral strains but also in normal human populations not exposed to

SARS-CoV-2 and can neutralize the virus[23]. The frequent use of the IGHV3-53/3-66 and IGHJ6

gene pair in SARS-CoV-2 neutralizing antibodies was further confirmed by a subsequent

report[24-26]. Six scFv clones carrying the IGHV1-69 gene were selected from two libraries, which

frequently encoded RBD-reactive antibodies found in convalescent patients from both the

ancestral strain and BA.1 variant[27]. Two additional scFv clones were selected from the two

libraries.

In the recombinant scFv-human Fc-hemagglutinin (HA) fusion protein format, seven scFv clones

showed reactivity against the RBD of the ancestral virus, Alpha, Beta, Gamma, and Delta

variants and Omicron-sublineage variants (BA.1, BA.2, BA.4/5 and BQ.1.1) (Extended Data Fig.

4a), with half-maximal effective concentrations (EC50) below 350 pM. The 27-60 clone and 35-

15 clone showed dramatically reduced reactivity to BA.4/5 and BQ.1.1 RBDs and BQ.1.1 RBD,

respectively. In a microneutralization assay, all clones neutralized the ancestral SARS-CoV-2

strain with a half-maximal inhibition concentration (IC50) range of 0.44 to 10.03 μg/ml

(Extended Data Fig. 4b). Four clones (35-8, 35-46, 35-68, and 35-78) showed slightly decreased

neutralizing activity against the BA.1 variant with an IC50 below 15 μg/ml, while the activity of

others was increased to 25.79 to 74.12 μg/ml. Three clones (35-8, 35-68, and 35-78) could also

neutralize Alpha, Beta, Gamma and Delta variants with less IC50 values under 12.12 ug/ml.

Omicron RBD-reactive clonotypes in BCR HC repertoires

BCR HC clonotypes of Omicron RBD-binding clones (Omicron RBD-reactive BCR HC

clonotype) were mapped to 293 sequences in the repertoire of 23 vaccinees (Table 1 and

Supplementary Table 3). The most frequently identified BCR HC clonotype was 27-60, found in

19 vaccinees (46%), followed clonotypes by 43-09 and 35-46, found in five and two vaccinees

(12% and 5%), respectively (Supplementary Table 4). Six BCR HC clonotypes were found only

in one vaccinee. Five BCR HC clonotypes, including 27-60, were not found in vaccinees in

whom the antibody clones were found. This type of discrepancy is expected to inevitably

originate from the incomplete *in silico* reconstitution of the BCR HC repertoire due to the

limitation of the throughput of next-generation sequencing[28]. In this regard, the BCR HC

clonotypes are expected to be present in a larger population than we observed herein.

Among the 293 BCR HC sequences, 57 and 216 BCR HC sequences were present in the third

and sixth repertoires, respectively. Only 20 BCR HC sequences were found in other

chronological BCR repertoires. This skewed distribution originated from the expansion and

proliferation of B cells encoding the Omicron RBD-reactive BCR HC clonotype immediately

after the second and third doses of the vaccine. Thus, we limited our statistical analysis to BCR

HC sequences present in the third and sixth repertoires. Rapid class switch recombination (CSR)

following vaccination was evident, as the main immunoglobulin subtype of Omicron RBD-

reactive BCR HC sequences was IgG_1 in the third repertoire, followed by IgG_2, which was

maintained in the sixth repertoire (Supplementary Table 4 and Extended Data Fig. 5a). This rapid

CSR resulting in the dominance of the IgG_1 subtype among RBD-reactive BCR HC sequences

was also observed in convalescent patients infected with the ancestral virus[23].

Additionally, somatic hypermutation (SHM) of Omicron RBD-reactive BCR HC clonotypes

occurred mainly after the third dose. The average number of SHMs in the IGHV gene of

Omicron RBD-reactive BCR HC clonotypes in the sixth repertoire was dramatically increased

compared to that in the third repertoire with statistical significance (Extended Data Fig. 5b). The

diversity of the HCDR3 sequence also increased from the third repertoire to the sixth repertoire

(Extended Data Fig. 5c). Five Omicron RBD-reactive BCR HC clonotypes were mapped to eight

vaccinees in both the third and sixth repertoires (Extended Data Fig. 5d). The increase in the

average number of SHMs was evident, and statistical significance was confirmed in three

vaccinees with the 27-60 clonotype and one vaccinee with the 43-34 clonotype.

Development of concomitant reactivity to ancestral and BA.1 RBDs via SHM

Clone 27-60 was encoded by IGHV3-53/IGHV3-66 (Fig. 2a), in which CDR1 and CDR2 are

known to have two key motifs providing reactivity to ancestral RBD[23,29]. The BCR HC of clone

27-60 had two SHMs in its IGHV gene sequence, and when we back-mutated the V27 residue in

CDR1 or the F58 residue adjacent to CDR2 to the germline sequence, its affinity for the ancestral

and BA.1 RBDs was decreased 8.7-fold and greatly diminished, respectively (Fig. 2b). This

observation showed that the reactivity of clone 27-60 toward the BA.1 RBD depended on the

SHM to a greater degree than its reactivity to the ancestral RBD.

Thereafter, we selected the most frequent BCR HC sequences of the 27-60 clonotype in the third

and sixth BCR repertoires of six vaccinees and analyzed their sequences (Fig. 2a). All the BCR

HC sequences found in the sixth repertoires had more SHMs than their pairs in the third

repertoires. All six BCR HC sequences in the sixth repertoire gained the Y58F mutation,

increasing the affinity of IGHV3-53/IGHV3-66 antibodies for the ancestral RBD[30,31]. There was

also diversity in their HCDR3 sequences limited to a motif of three amino acid residues. As the

HCDR3 sequence can diversify either at the stage of germline VDJ gene recombination by P-

and N-nucleotide addition/deletion or later by SHM, it was not possible to determine the

germline HCDR3 sequence of clones 27-60. To analyze the influence of the HCDR3 sequence on

reactivity to ancestral and BA.1 RBDs, we generated a scFv phage display with the randomized

HCDR3 sequence of ARDLXX(A/V)GGMDV, arbitrarily selected 672 phage clones and

checked their reactivity to both RBDs. For the ancestral RBD, 200 phage clones (30%) showed

positive reactivity (Fig. 2c, Extended Data Fig. 6 and Supplementary Table 5). However, for the

BA.1 RBD, only 18 phage clones (3%) were reactive. All 18 clones were dual-reactive to both

RBDs. Six mono-reactive and six dual-reactive phage clones were arbitrarily selected for

expression as recombinant scFv-hFc-HA fusion proteins and tested for their affinity for both

RBDs. The EC50 values of mono-reactive scFv proteins to the ancestral and BA.1 RBDs were in

the ranges of 0.84–3.80 nM and 3.83–17.25 nM, respectively. In addition, dual-reactive scFv

proteins showed EC50 values of 0.16–1.29 nM and 0.35–1.26 nM for the ancestral and BA.1

RBDs, respectively (Fig. 2d). These observations proved that it is relatively rare for ancestral

RBD-reactive BCR HC clonotypes encoded by the IGHV3-53/3-66 and IGHJ6 genes with the

HCDR3 sequence of ARDLXX(A/V)GGMDV (BCR HC XXA/V clonotypes) to develop

increased compatible reactivity to the BA.1 RBD. This dual reactivity is likely to be achieved

among the large population of antigen-stimulated and proliferating B cells through SHM, rather

than in a small number of B cells in the stage of germline VDJ gene rearrangement. Indeed, only

one BCR HC XXA/V sequence was found in only one vaccinee in the pre-immune repertoire

(Extended Data Fig. 7 and Supplementary Table 6). After the first, second, and third injections, 3,

58, and 138 BCR HC XXA/V sequences were found in two, ten and sixteen vaccinees,

respectively. The average frequency of BCR HC XXA/V sequences also increased from 6.06×10^{-6}

under pre-immune status to 2.85×10^{-5}, 1.91×10^{-5}, and 2.41×10^{-5} after the first, second, and third

doses, respectively.

To check the effect of SHM accumulation on the affinity for RBDs in 27-60 BCR HC

clonotypes, six pairs of BCR HC sequences were coupled with the light chain gene of clone 27-

60 (Supplementary Table 7) and expressed as recombinant scFv-hFc-HA fusion proteins. Then,

their affinity for the ancestral RBD and BA.1 RBD was determined by ELISA. For the ancestral

RBD, all six scFvs of the third repertoire showed notable affinities, with EC50 values less than

7.5 nM. However, only one scFv from vaccinee 1 showed comparable affinity for BA.1 RBD

(Fig. 2e and Supplementary Table 8). In the sixth repertoire, four additional scFvs showed

affinity for the BA.1 RBD, with EC50 values less than 4.5 nM.

We also analyzed the BCR HC sequences of four additional clonotypes (35-15, 35-46, 43-09 and

43-34) found in both the third and sixth repertoires of identical vaccinees (Extended Data Fig.

8a). All BCR HC sequences in the sixth repertoires showed more SHMs than their paired

sequences in the third repertoires. Then, we prepared eight recombinant scFv-hFc-HA fusion

proteins in pairs with the light chains of four scFv phage clones. In ELISA, all scFvs from the

third repertoire showed a remarkable level of affinity for the ancestral RBD, with EC50 values

less than 200 pM, which were well maintained within a 2-fold difference in their paired

sequences in the sixth repertoire (Extended Data Fig. 8b). It was exceptional that the BCR HC

sequences of the 35-46 clonotype in the third library were encoded by IGHV1-69/IGHJ3 genes

without any SHM and showed a significantly high affinity for BA.1 RBD. In the 35-15, 43-09

and 43-34 BCR HC clonotype pairs, the affinity for BA.1 RBD was significantly increased from

0.59, 1.12 and 7.91 nM in the third repertoire to 0.13, 0.09 and 0.14 nM in the sixth repertoire,

respectively.

Expansion of the BCR HC repertoire to RBDs of Omicron subvariants by SHM

We studied how the diversification of the BCR HC repertoire by SHM expands its reactivity to

RBDs of Omicron subvariants. We first searched the BCR HC sequences of the 27-60, 35-15,

35-46, 43-09 and 43-34 clonotypes in the third repertoire and found them in nine, one, one, one,

and one of the vaccinees, respectively (Supplementary Table 4). Then, we selected the BCR HC

sequences with the highest frequency in each vaccinee and used them to construct a new set of

BCR HC clonotypes (designated clonotypes-3rd) confined to each vaccinee. The greatest

numbers of BCR HC sequences in the 27-60, 35-15, 35-46, 43-09 and 43-34 clonotypes-3rd were

found in vaccinees 35, 43, 35, 23, and 43, consisting of 86, 8, 24, 2, and 18 sequences,

respectively. Based on their HCDR3 sequences, these sequences in the clonotypes-3rd were

subdivided into 11, 4, 11, 2, and 9 clusters, respectively. From each cluster, the BCR HC

sequences with the highest frequency that were present in the third and sixth BCR HC repertoires

were selected, and their genes were synthesized (Extended Data Fig. 9, 10 and Supplementary

Table 7, 9). As expected, all the scFv clones in the sixth BCR HC repertoires showed more SHM

than those in the third repertoires (Fig. 3), and some of them showed dramatically enhanced

binding toward specific types of Omicron variant RBDs. Two scFv clones (35-15/6th-43 and 35-

46/6th-35-2) showed potent reactivity confined to either BA.1 or XBB.1.5 RBD with an EC50

below 0.100 nM. The other two scFv clones (35-46/6th-35-1 or 43-09/6th-23) showed the

equivalent EC50 values toward RBDs of Omicron subvariants (either BA.1 or BQ.1.1 in addition

to XBB.1.5, individually). Another two scFv clones (43-34/6th-43-1 and 43-34/6th-43-3) reacted

with the RBDs of all Omicron subvariants tested with same potency. These findings suggest that

the accumulation of SHM after the third dose of ancestral SARS-CoV-2 vaccine generated clones

with divergent mosaic pattern of specificity toward three major Omicron subvariants. More

interestingly, clones reactive to BQ.1.1 and XBB.1.5 achieved their specificity much earlier than

their emergence.

Our study showed that the third dose of the mRNA vaccine encoding the ancestral viral spike

protein induced the accumulation of SHM in ancestral RBD-reactive antibody clones and

generated a panel of BCRs with additional divergent reactivity to RBDs of Omicron subvariants,

which may contribute to the dramatically increased plasma level of Omicron RBD–reactive

antibodies and may have the potential to counteract novel SARS-CoV-2 variants yet to emerge.

We believe that this expansion in the specificity of the BCR repertoire to unseen antigens via

SHM is a protective immune mechanism that evolved in response to the challenge of viral

immune escape.

References

1   Meredith, L. W. *et al.* Rapid implementation of SARS-CoV-2 sequencing to invest

igate cases of health-care associated COVID-19: a prospective genomic surveillanc

e study. *The Lancet Infectious Diseases* **20**, 1263-1271 (2020). https://doi.org:10.10

16/s1473-3099(20)30562-4

2   Harvey, W. T. *et al.* SARS-CoV-2 variants, spike mutations and immune escape.

Nature Reviews Microbiology **19**, 409-424 (2021). https://doi.org:10.1038/s41579-02

1-00573-0

3   Letko, M., Marzi, A. & Munster, V. Functional assessment of cell entry and recep

tor usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nature Microbio

logy* **5**, 562-569 (2020). https://doi.org:10.1038/s41564-020-0688-y

4   Liu, L. *et al.* Potent neutralizing antibodies against multiple epitopes on SARS-Co

V-2 spike. *Nature* **584**, 450-456 (2020). https://doi.org:10.1038/s41586-020-2571-7

5   Piccoli, L. *et al.* Mapping Neutralizing and Immunodominant Sites on the SARS-

CoV-2 Spike Receptor-Binding Domain by Structure-Guided High-Resolution Serolo

gy. *Cell* **183**, 1024-1042.e1021 (2020). https://doi.org:10.1016/j.cell.2020.09.037

6   Jackson, L. A. *et al.* An mRNA Vaccine against SARS-CoV-2 - Preliminary Repo

rt. *N Engl J Med* **383**, 1920-1931 (2020). https://doi.org:10.1056/NEJMoa2022483

7   Cao, Y. *et al.* Omicron escapes the majority of existing SARS-CoV-2 neutralizing

antibodies. *Nature* **602**, 657-663 (2022). https://doi.org:10.1038/s41586-021-04385-3

8   Cele, S. *et al.* Omicron extensively but incompletely escapes Pfizer BNT162b2 ne

utralization. *Nature* **602**, 654-656 (2022). https://doi.org:10.1038/s41586-021-04387-1

9   Dejnirattisai, W. *et al.* SARS-CoV-2 Omicron-B.1.1.529 leads to widespread escape

from neutralizing antibody responses. *Cell* **185**, 467-484.e415 (2022). https://doi.or

g:10.1016/j.cell.2021.12.046

10 Chalkias, S. *et al.* A Bivalent Omicron-Containing Booster Vaccine against Covid-

19. *New England Journal of Medicine* **387**, 1279-1291 (2022). https://doi.org:10.10

56/nejmoa2208343

11 Tseng, H. F. *et al.* Effectiveness of mRNA-1273 vaccination against SARS-CoV-2

omicron subvariants BA.1, BA.2, BA.2.12.1, BA.4, and BA.5. *Nature Communicati*

ons **14** (2023). https://doi.org:10.1038/s41467-023-35815-7

12 Zou, J. *et al.* Neutralization of BA.4-BA.5, BA.4.6, BA.2.75.2, BQ.1.1, and XBB.

1 with Bivalent Vaccine. *N Engl J Med* **388**, 854-857 (2023). https://doi.org:10.10

56/NEJMc2214916

13 Scheaffer, S. M. *et al.* Bivalent SARS-CoV-2 mRNA vaccines increase breadth of

neutralization and protect against the BA.5 Omicron variant in mice. *Nat Med* **29**,

247-257 (2023). https://doi.org:10.1038/s41591-022-02092-8

14 Evans, J. P. *et al.* Neutralization of SARS-CoV-2 Omicron sub-lineages BA.1, BA.

1.1, and BA.2. *Cell Host Microbe* **30**, 1093-+ (2022). https://doi.org:10.1016/j.chom

.2022.04.014

15 Garcia-Beltran, W. F. *et al.* mRNA-based COVID-19 vaccine boosters induce neutr

alizing immunity against SARS-CoV-2 Omicron variant. *Cell* **185**, 457-466.e454 (2

022). https://doi.org:10.1016/j.cell.2021.12.033

16 Nemet, I. *et al.* Third BNT162b2 Vaccination Neutralization of SARS-CoV-2 Omic

ron Infection. *New England Journal of Medicine* **386**, 492-494 (2022). https://doi.o

rg:10.1056/nejmc2119358

17   Muik, A. *et al.* Neutralization of SARS-CoV-2 Omicron by BNT162b2 mRNA vac

cine-elicited human sera. *Science* **375**, 678-+ (2022). https://doi.org:10.1126/science.

abn7591

18   Carreño, J. M. *et al.* Activity of convalescent and vaccine serum against SARS-C

oV-2 Omicron. *Nature* **602**, 682-688 (2022). https://doi.org:10.1038/s41586-022-0439

9-5

19   Yu, J. *et al.* Neutralization of the SARS-CoV-2 Omicron BA.1 and BA.2 Variants.

New England Journal of Medicine **386**, 1579-1580 (2022). https://doi.org:10.1056/

nejmc2201849

20   Goel, R. R. *et al.* Efficient recall of Omicron-reactive B cell memory after a third

dose of SARS-CoV-2 mRNA vaccine. *Cell* **185**, 1875-1887.e1878 (2022). https://d

oi.org:10.1016/j.cell.2022.04.009

21   Kotaki, R. *et al.* SARS-CoV-2 Omicron-neutralizing memory B cells are elicited b

y two doses of BNT162b2 mRNA vaccine. *Science immunology* **7**, eabn8590 (202

2).

22   Raybould, M. I. J., Kovaltsuk, A., Marks, C. & Deane, C. M. CoV-AbDab: the c

oronavirus antibody database. *Bioinformatics* **37**, 734-735 (2021). https://doi.org:10.

1093/bioinformatics/btaa739

23   Kim, S. I. *et al.* Stereotypic neutralizing V-H antibodies against SARS-CoV-2 spik

e protein receptor binding domain in patients with COVID-19 and healthy individu

als. *Sci Transl Med* **13** (2021). https://doi.org:ARTN eabd6990 10.1126/scitranslmed

.abd6990

24   Barnes, C. O. *et al.* SARS-CoV-2 neutralizing antibody structures inform therapeut

ic strategies. *Nature* **588**, 682-687 (2020). https://doi.org:10.1038/s41586-020-2852-1

25 Cao, Y. *et al.* Potent Neutralizing Antibodies against SARS-CoV-2 Identified by H

igh-Throughput Single-Cell Sequencing of Convalescent Patients' B Cells. *Cell* **182**

, 73-84.e16 (2020). https://doi.org:10.1016/j.cell.2020.05.025

26 Zhang, Q. *et al.* Potent and protective IGHV3-53/3-66 public antibodies and their

shared escape mutant on the spike of SARS-CoV-2. *Nature Communications* **12** (2

021). https://doi.org:10.1038/s41467-021-24514-w

27 Yan, Q. *et al.* Shared IGHV1-69-encoded neutralizing antibodies contribute to the

emergence of L452R substitution in SARS-CoV-2 variants. *Emerging Microbes &a*

mp; Infections **11**, 2749-2761 (2022). https://doi.org:10.1080/22221751.2022.2140611

28 Lee, H. *et al.* Optimization of peripheral blood volume for in silico reconstitution

of the human B-cell receptor repertoire. *FEBS Open Bio* **12**, 1634-1643 (2022).

https://doi.org:10.1002/2211-5463.13467

29 Yuan, M. *et al.* Structural basis of a shared antibody response to SARS-CoV-2. *Sc*

ience **369**, 1119-1123 (2020). https://doi.org:10.1126/science.abd2321

30 Tan, T. J. C. *et al.* Sequence signatures of two public antibody clonotypes that bi

nd SARS-CoV-2 receptor binding domain. *Nature Communications* **12** (2021). https

://doi.org:10.1038/s41467-021-24123-7

31 Tian, X. *et al.* The prominent role of a CDR1 somatic hypermutation for converg

ent IGHV3-53/3-66 antibodies in binding to SARS-CoV-2. *Emerging Microbes &a*

mp; Infections **11**, 1186-1190 (2022). https://doi.org:10.1080/22221751.2022.2063074

334
335

336 **Tables**

337 **Table 1.** Omicron RBD-reactive scFv clones and their BCR-HC clonotypes

scFv clone	scFv library (vaccinees)	HCDR3 sequences	IGHV	IGHJ	No. of vaccinees with the corresponding BCR HC clonotype
27-60	22, 27, 32, 39, 43	ARDLMEAGGMDV	*IGHV3-53/3-66*	*IGHJ6*	19
35-08	35	AREIGYSGSGSAKYFDP	*IGHV1-69*	*IGHJ5*	1
35-15	35	ARAEDHGTYYSDSSGYHFDY	*IGHV1-69*	*IGHJ4*	1
35-46	35	ARVHGYSGYGANDAFDI	*IGHV1-69*	*IGHJ3*	2
35-68	35	AREVGYSGFGASPKFDP	*IGHV1-69*	*IGHJ5*	1
35-78	35	AREPGILGYCSSTSCYID	*IGHV1-69*	*IGHJ4*	1
43-09	43	ARTGSGWTDAFDI	*IGHV3-30*	*IGHJ3*	5
43-27	43	ATTYHYDTDGPYGEFYY	*IGHV5-51*	*IGHJ4*	1

| 43-34 | 43 | ARVRGYSGYGASGYFDN | *IGHV1-69* | *IGHJ4* | 1 |

Figures

Fig. 1. Vaccination, blood sampling and plasma antibody levels. a, Vaccinees received the

first two doses at a three-week interval and the third dose between the 31st and 39th weeks after

the first dose. Blood was collected six times: pre-vaccination (1st); 1 week after the first dose

(2nd); 1, 6, and 30 weeks after the second dose (3rd, 4th, and 5th); and one to four weeks after the

third dose (6th). **b,** Plasma levels of antibodies against SARS-CoV-2 ancestral and BA.1 RBDs

were measured after 2,500-fold dilution. All experiments were performed in duplicate, and the

average value for each vaccinee was plotted. The P value was calculated using one-way ANOVA.

ns, not significant, $p > 0.05$; *, $p < 0.05$; ***, $p < 0.001$; ****, $p < 0.0001$. The increases and

decreases in antibody levels are marked with red and green asterisks, respectively.

Fig. 2. Characterization of the 27-60 BCR HC clonotype. a, BCR HC sequences found in six

vaccinees in their third and sixth BCR repertoires with the highest frequency. **b,** Reactivity of the

recombinant 27-60 scFv-hFc-HA protein and its clones back mutated to the germline sequence.

c, Reactivity of scFv-displaying phage clones from the HCDR3-randomized library to the

ancestral and BA.1 RBDs. The clones labeled in red and green were selected for expression as

recombinant scFv-hFc-HA protein. **d,** Reactivity of recombinant scFv fusion proteins of

HCDR3-randomized (ARDLXX(A/V)GGMDV) clones to the ancestral and BA.1 RBDs. **e,**

Reactivity of recombinant scFv-hFc-HA proteins encoded by individual BCR HC sequences and

the light chain gene of clone 27-60 to the ancestral and BA.1 RBDs.

Clonotype		Ancestral	BA.1	BQ.1.1	XBB.1.5
27-60	3rd-35	2.778	15.640	18.160	20.040
	3rd-35-1	81.320	54.730	73.220	60.480
	6th-35	0.098	0.179	19.460	25.400
	6th-35-1	0.067	3.997	29.340	38.380
	6th-35-2	0.071	2.067	8.850	16.820
	6th-35-3	0.074	0.630	12.870	15.270
	6th-35-4	0.107	4.522	83.480	77.860
	6th-35-5	1.005	55.400	90.360	81.240
	6th-35-6	0.854	11.330	65.310	67.820
	6th-35-7	0.220	0.890	25.760	32.590
	6th-35-8	0.156	2.250	37.720	34.140
35-15	3rd-43	0.105	3.408	134.100	81.640
	6th-43	0.050	0.091	17.660	11.370
	6th-43-1	0.188	3.208	6.809	7.444
	6th-43-2	0.134	2.459	10.330	11.470
35-46	3rd-35	0.149	0.564	4.309	0.316
	6th-35	0.081	0.154	0.263	0.101
	6th-35-1	0.075	0.099	0.130	0.066
	6th-35-2	0.044	0.104	0.330	0.065
	6th-35-3	0.103	0.704	42.720	1.277
	6th-35-4	0.123	0.552	3.022	0.226
	6th-35-5	0.290	0.875	2.303	0.495
	6th-35-6	0.130	0.329	3.505	0.168
	6th-35-7	0.229	0.344	0.815	0.213
	6th-35-8	0.074	0.168	1.833	0.108
	6th-35-9	0.089	0.177	1.024	0.240
43-09	3rd-23	0.150	16.650	0.210	0.254
	6th-23	0.079	0.487	0.080	0.073
43-34	3rd-43	0.401	2.412	8.159	3.557
	6th-43	0.118	0.138	0.133	0.079
	6th-43-1	0.100	0.094	0.075	0.080
	6th-43-2	0.089	0.100	0.119	0.090
	6th-43-3	0.108	0.097	0.075	0.069
	6th-43-4	0.072	0.212	0.375	0.476
	6th-43-5	0.125	0.236	0.347	0.822
	6th-43-6	0.075	0.462	0.218	0.200
	6th-43-7	0.167	0.269	0.381	0.493

EC50 (nM)

Legend: 0 – 0.05 | 0.05 – 0.10 | 0.10 – 0.50 | 0.50 – 1.00 | 1.00 – 10.00 | 10.00 – 100.00 | 100.00 – 200.00

Clonotype	HCDR3 sequence	No. of SHM
27-60	ARDLAAAGGMDV	5
	ARGLAAAGGMDV	5
	ARDLQEAGGMDV	21
	ARDLEAAGGMDV	7
	ARDLLEAGGMDV	8
	ARDLAEAGGMDV	14
	ARDLEAAGAMDV	17
	ARDLAEAGAMDV	10
	ARDLAETGGMDV	10
	ARDLEMAGGMDV	16
	ARDLEVAGGMDV	12
35-15	ARAEDHGTYYSDSSGYHFDF	4
	ARAEDHGTYYSDSSGYHFDY	17
	ARPEDHGTYYSDSSGYHFDY	18
	ARPEDHGTYYSDSSGYHFDH	18
35-46	ARVRGYSGYGANDAFDI	0
	ARVHGYSGYGANDAFDL	14
	ARVHGYSGYGANDAFDI	10
	ARVRGYSDYGANDFFDM	20
	ARVRGYSDYGANDAFDI	13
	ARVRGYSDYGANDAFDM	14
	TRVRGYSAYGANDAFDM	20
	ARVRGYTGYGANDAFDI	27
	ARVRGYSGYGANDAFDM	21
	ARVRGYSDYGANDAFDL	10
	ARVRGYSGYGANEAFDI	16
43-09	ARTNSGWSDAFDI	3
	ARTNSGWTDALDI	16
43-34	ARARGYSGYDASGYFDY	1
	ARVRGYSGYGASGYFDN	10
	ARARGYSGYGASAYLDY	16
	ARGRGYSGYGASGYFDN	22
	ARARGYSGYGASAYFDY	17
	ARGRGYSGYGASGYFDY	17
	ARGRGYSAYGASGYFDY	17
	ARARGYSDYGASGYFDS	9
	ARARGYSDYGASGYFDY	11

362 **Fig. 3. The reactivity of the scFv clones of HCDR3-based clusters in the 27-60, 35-15, 35-46,**

363 **43-09, and 43-34 BCR HC clonotypes-3rd.** The EC50 values to Ancestral, BA.1, BQ.1.1, and

364 XBB.1.5 RBDs of the scFv clones with the highest frequencies of HCDR3-based clusters in each

365 clonotype-3rd are displayed with their HCDR3 sequences and the quantities of SHM.

Methods

Study participants

Peripheral blood sampling was approved by the Institutional Ethics Review Board of Seoul

National University Hospital (IRB approval number, 2102-032-1193). The vaccinees had a

median age of 30 years (range 23-62) and showed a nearly equal distribution of males and

females (46% and 54%, respectively). Peripheral blood mononuclear cells (PBMCs) and plasma

were separated using Lymphoprep (STEMCELL) Ficoll (Cytiva) following the manufacturer's

instructions. Total RNA was isolated using TRIzol Reagent (Invitrogen) according to the

manufacturer's instructions.

ELISA

Plasma and phage ELISAs were performed in 96-well microtiter plates (Corning, 3690) coated

with 100 ng of recombinant SARS-CoV-2 proteins (Sino Biological, Ancestral RBD, 40592-

V08H; Alpha RBD, 40592-V08H82; Beta RBD, 40592-V08H85; Gamma RBD, 40592-V08H86;

Delta RBD, 40592-V08H90; Omicron BA.1 RBD, 40592-V08H121; Omicron BA.2 RBD,

40592-V08H123, Omicron BA.4/5 RBD, 40592-V08H130; Omicron BQ.1.1 RBD, 40592-

V08H143; Ancestral N, 40588-V08B) in coating buffer (0.1 M sodium bicarbonate, pH 8.6) as

described previously [23]. Briefly, the plates were coated with the antigen by incubation at 4°C

overnight and blocked with 3% bovine serum albumin (BSA) in PBS for 1 hour at 37°C. Then,

serially diluted serum (100-, 500-, 2,500-fold) or phage supernatant (twofold) in blocking buffer

was added to the wells of microtiter plates, followed by incubation for 1 hour at 37°C. Then, the

plates were washed three times with 0.05% PBST. Horseradish peroxidase (HRP)-conjugated

goat anti-human IgM and IgA (Invitrogen, A18835 and A18781, 1:5,000), rabbit anti-human IgG

antibody (Invitrogen, 31423, 1:20,000) and HRP-conjugated anti-M13 antibody (Sino

Biological, 11973-MM05T-H, 1:4,000) were used to determine the amount of bound antibody or

M13 bacteriophage. A 3,3′,5,5′-tetramethylbenzidine liquid substrate solution (Thermo Fisher

Scientific Inc.) was used as an HRP substrate, and the absorbance was measured at 450 or 650

nm using a microplate spectrophotometer (Thermo Fisher Scientific Inc., Multiskan GO)

depending on the use of 2 M sulfuric acid as the stop solution. All assays were performed in

duplicate.

In ELISA for the recombinant scFv-hFc-HA fusion proteins, the wells of microtiter plates

(Corning) were first coated with 100 ng of mouse anti-His antibody (Invitrogen, MA1-21315)

and blocked. Then, the recombinant SARS-CoV-2 RBD protein with a polyhistidine tag (100

nM) was added to the wells. After brief washing, the scFv-hFC-HA fusion proteins were serially

diluted fourfold from 1 μM to 0.24 pM in blocking buffer and added to wells of a microtiter

plate. HRP-conjugated rat anti-HA antibody (Roche, 12013819001, 1:1,000) was used to

determine the amount of bound antibody. All assays were performed in duplicate.

Next-generation sequencing

Genes encoding the variable domain of the heavy chain (V_H) and part of the first constant

domain of the heavy chain (CH1) domain (V_H-CH1) were amplified using specific primers, as

described previously[23]. All primers used are listed in Supplementary Table 10. Briefly, total RNA

was used as a template to synthesize complementary DNA (cDNA) using the SuperScript IV

First-Strand Synthesis System (Invitrogen) with specific primers (primer No. 1–8) targeting the

CH1 domain of each isotype (IgM, IgD, IgG, IgA, and IgE) according to the manufacturer's

protocol. After cDNA synthesis, 1.8 volumes of SPRI beads (Beckman Coulter, AMPure XP)

were used to purify cDNA, which was eluted in 35 μl of water. The purified cDNA (15 μl) was

subjected to second-strand synthesis in a 25 μl reaction volume using IGHV gene–specific

primers (primer No.9–14) and a KAPA Biosystems kit (Roche, KAPA HiFi HotStart). The PCR

conditions were as follows: 95°C for 3 min, 98°C for 30 s, 60°C for 45 s, and 72°C for 6 min.

After second-strand synthesis, dsDNA was purified using 1 volume of SPRI beads, as described

above. V_H-CH1 genes were amplified using purified dsDNA (15 µl) in a 25 µl reaction volume

using primers containing indexing sequences (primers 15 and 16) and a KAPA Biosystems kit.

The PCR conditions were as follows: 95°C for 3 min; 25 cycles of 98°C for 30 s, 60°C for 30 s,

and 72°C for 1 min; and 72°C for 5 min. PCR products were subjected to electrophoresis on a

1.5% agarose gel and purified using a QIAquick gel extraction kit (QIAGEN Inc.) according to

the manufacturer's instructions. The gel-purified PCR products were purified again using 1

volume of SPRI beads and eluted in 20 μl water. The SPRI-purified sequencing libraries were

quantified with a 4200 TapeStation System (Agilent Technologies) using a D1000 ScreenTape

assay and subjected to next-generation sequencing on the Illumina NovaSeq platform.

NGS data processing

Raw sequencing reads obtained from sequencing the V_H-CH1 region of B cells in peripheral

blood were processed using a custom pipeline. The pipeline included adapter trimming and

quality filtering; unique molecular identifier (UMI) processing; V(D)J gene annotation;

clustering; quality control; and diversity analysis.

The forward reads (R1) and reverse reads (R2) of the raw NGS data were merged using paired-

end read merger (PEAR) v0.9.10 with the default settings[32]. The merged reads were q-filtered

under the q20p95 condition, resulting in 95% of base pairs in the reads having a Phred score

greater than 20. Primer positions were identified in the quality-filtered reads, and primer regions

were trimmed to remove the effects of primer synthesis errors while allowing one substitution or

deletion.

Based on the primer recognition results, UMI sequences were extracted, and reads were clustered

according to the UMI sequences. To eliminate index misassignment, we subclustered the

clustered reads based on the similarity of the reads (allowing 5 mismatches in each subcluster)

and matched the majority subcluster to the UMI. The subclustered reads were aligned using the

multiple sequence alignment tool Clustal Omega v1.2.4 with the default settings[33,34]. Consensus

calling was performed by selecting major frequency bases at every position of the aligned

sequences. The number of reads in the consensus sequence was redefined as the number of UMI

subclusters belonging to the consensus sequence.

Sequence annotation consisted of isotype annotation and V(D)J annotation. The consensus

sequence was divided into a V(D)J region and a constant region. The isotype of the consensus

sequence was annotated by aligning the extracted constant region with the constant gene of the

International Immunogenetics Information System (IMGT)[35]. Then, the V(D)J region of the

consensus sequence was annotated using an updated version of IgBLAST (v1.17.1)[36]. Among

the annotation results, the IGHV genes, IGHJ genes, HCDR3 sequences, and number of SHMs

were extracted for further analysis. The number of SHMs was counted only in the aligned IGHV

genes. The nonfunctional consensus reads were defined and filtered using the following criteria:

(i) sequence length shorter than 250 base pairs, (ii) presence of a stop codon or frameshift in the

entire amino acid sequence, (iii) failure to annotate one or more HCDR1, HCDR2 and HCDR3

regions, and (iv) failure to annotate the isotype.

Construction of scFv phage display library

Six human scFv phage display libraries were constructed using the total RNA prepared from the

sixth blood sample for vaccine Nos. 22, 27, 32, 35, 39 and 43 as described previously[23]. Briefly,

for the V_H and V_κ/V_λ genes, total RNA was employed to synthesize cDNA using the SuperScript

IV First-Strand Synthesis System (Invitrogen) with gene-specific primers targeting the J_H and

$C_{\kappa/\lambda}$ genes (primer No.17–22), respectively. After cDNA synthesis, 1.8 volumes of SPRI beads

(Beckman Coulter) were used to purify cDNA, which was eluted in 20 μl of water. The purified

cDNA (11.25 μl) of the V_H gene was subjected to second-strand synthesis in a 25 μl reaction

volume using IGHV gene-specific primers (primer No. 23–32, 7.5 μl) and a KAPA Biosystems

kit (Roche). The reaction conditions were as follows: 95°C for 3 min, 98°C for 1 min, 60°C for 1

min, and 72°C for 5 min. In the case of the V_κ/V_λ gene, the eluted cDNA (17.25 μl) was used for

the first round of PCR synthesis in a 25 μl reaction volume using V_κ/V_λ and J_κ/J_λ gene-specific

primers (primer No. 33–69, 0.75 μl). The PCR conditions were as follows: 95°C for 3 min; 4

cycles of 98°C for 1 min, 60°C for 1 min, and 72°C for 1 min; and 72°C for 10 min. After

second-strand synthesis for the V_H gene or PCR amplification of the V_κ/V_λ gene, double-stranded

DNA (dsDNA) was purified with 1 volume of SPRI beads and eluted in 40 μl of water. V_H and

V_κ/V_λ genes were amplified using 10 μl of purified dsDNA, 2.5 pmol of the primers (primer No.

70–73), and KAPA Biosystems kit components in a 50 μl total reaction volume with the

following thermal cycling program: 95°C for 3 min; 30 cycles of 98°C for 30 s, 60°C for 30 s,

and 72°C for 1 min; and 72°C for 10 min. Then, the amplified V_H and V_κ/V_λ genes were

subjected to electrophoresis on a 1.5% agarose gel and purified using a QIAquick gel extraction

kit (QIAGEN Inc.) according to the manufacturer's instructions. The purified V_H and V_κ/V_λ gene

fragments (100 ng) were mixed and subjected to overlap extension PCR to generate scFv genes

using 2.5 pmol of the overlap extension primers (primer No. 74 and 75) using the KAPA

Biosystems kit. The PCR conditions were as follows: 95°C for 3 min; 25 cycles of 98°C for 20 s,

65°C for 15 s, and 72°C for 1 min; and 72°C for 10 min. The amplified scFv gene was purified

and cloned into a phagemid vector as described previously[37].

Selection of BA.1RBD-binding clones

Human scFv phage display libraries with 7.1×10^8, 6.1×10^8, 8.1×10^8, 8.3×10^8,

7.9×10^8, and 7.4×10^8 colony-forming units were generated using cDNA prepared from

vaccinee Nos. 22, 27, 32, 35, 39 and 43, respectively. The libraries were subjected to five rounds

of biopanning against the recombinant SARS-CoV-2 BA.1 RBD protein (Sino Biological Inc.) as

described previously[38]. Immune tubes (SPL, 43015) coated with 17 µg of the BA.1 RBD were

used for the first round, and 5×10^6 magnetic beads (Invitrogen, Dynabeads M-270 epoxy)

conjugated with 1.4 µg of the BA.1 RBD protein were used for the other rounds. After each

round of biopanning, the bound phages were eluted and amplified for the next round of

biopanning. For the selection of BA.1 RBD-reactive scFv phage clones, individual phage clones

were amplified from the titration plate of the last round and subjected to phage ELISA as

described previously[39]. The genes encoding BA.1 RBD-reactive scFv clones were identified

using phagemid DNA prepared from phage clones and Sanger nucleotide sequencing as

described previously[38]. A recombinant scFv protein fused with human IgG1 FC and the HA

peptide (scFv-hFc-HA) was expressed using a mammalian expression system and purified as

described previously[39].

HCDR3-randomized scFv phage display library

The gene fragment encoding the V_H region with a randomized HCDR3 sequence and another

gene fragment encoding the rest of the scFv were amplified using phagemid DNA of clone 27-

60, primers for randomization (primer No.76-79) and the KAPA Biosystems kit. The PCR

conditions were as follows: 95°C for 3 min; 25 cycles of 98°C for 20 s, 65°C for 15 s, and 72°C

for 1 min; and 72°C for 10 min. Then, the amplified genes were subjected to electrophoresis on a

1.5% agarose gel and purified using a QIAquick gel extraction kit (QIAGEN Inc.) according to

the manufacturer's instructions. The purified gene fragments (200 ng) were mixed and subjected

to overlap extension PCR to generate scFv genes using 2.5 pmol of the overlap extension primers

(primer No.76 and 79) and the KAPA Biosystems kit. The PCR conditions were as follows: 95°C

for 3 min; 25 cycles of 98°C for 20 s, 65°C for 15 s, and 72°C for 1 min; and 72°C for 5 min.

The amplified scFv genes were purified and cloned into a phagemid vector as described

previously[37]. For phage ELISA, individual phage clones were amplified from the titration plate

and subjected to phage ELISA as described previously[40].

Microneutralization assay

The ancestral SARS-CoV-2 (βCoV/Korea/KCDC03/2020 NCCP43326), Alpha B.1.1.7 (hCoV-

19/Korea/KDCA51463/2021 (NCCP 43381), Beta B.1.351 (hCoV-19/Korea/KDCA55905/2021

(NCCP 43382), Gamma P.1 (hCoV-19/Korea/KDCA95637/2021 (NCCP 43388), Delta B.1.617.2

(hCoV-19/Korea/KDCA119861/2021 (NCCP 43390), and Omicron B.1.1.529 (hCoV-

19/Korea/KDCA447321/2021 NCCP43408) viruses were obtained from the Korea Disease

Control and Prevention Agency. The viruses were propagated in Vero cells (ATCC, CCL-81) in

Dulbecco's Modified Eagle's Medium (DMEM, Welgene) in the presence of 2% fetal bovine

serum (Gibco, Thermo Fisher Scientific Inc.), as described previously[23,41].

Neutralization assays were performed as described previously[23,42]. Briefly, Vero cells were

seeded in 96-well plates (1.5×10^4 or 0.5×10^4 cells/well) in Opti-PRO SFM (Thermo Fisher

Scientific Inc.) supplemented with 4 mM L-glutamine and $1\times$ Antibiotics-Antimycotic (Thermo

Fisher Scientific Inc/) and grown for 24 h at 37°C in a 5% CO_2 environment. Recombinant scFv-

hFc proteins were diluted from 100 to 0.1953 µg/ml (twofold) in phosphate-buffered saline

(PBS, Welgene) and mixed with 100 or 500 $TCID_{50}$ of SARS-CoV-2. Then, the mixture was

incubated for 30 min at 37°C and added to the cells in tetrads, followed by incubation for 4 or 6

days at 37°C in a 5% CO_2 environment. The cytopathic effect (CPE) in each well was visualized

following crystal violet staining 4 or 6 days post infection. The IC_{50} values were calculated using

the dose□response inhibition equation of GraphPad Prism 6 (GraphPad Software).

Construction of BCR HC clonotype-3^{rd}

The most frequent BCR HC sequences of the 27-60, 35-15, 35-46, 43-09, and 43-34 clonotypes

in the third BCR repertoires were mapped to the chronological BCR repertoires following the

same definition of BCR HC clonotypes. All the mapped nucleotide sequences were aligned using

the multiple sequence alignment tool Clustal Omega v1.2.4 with the default settings[33,34]. The

aligned mapped nucleotide sequences were interpreted using a phylogenetic tree generated by

IgPhyML v1.1.3 052020 using the HLP model option[43]. The phylogenetic trees were plotted

using Interactive Tree of Life (iTOL)[44].

Data availability

All sequencing data are available from the National Center for Biotechnology Information

(www.ncbi.nlm.nih.gov/) under accession number PRJNA945512 (SRA).

Code availability

NGS data preprocessing was performed as described in a previous study (Kim, S. I. *et al*. Sci

Transl Med 13 (2021)). The code use in this study is available upon request for academic

research purposes.

Methods references

23 Kim, S. I. *et al.* Stereotypic neutralizing V-H antibodies against SARS-CoV-2 spik

e protein receptor binding domain in patients with COVID-19 and healthy individu

als. *Sci Transl Med* **13** (2021). https://doi.org:ARTN eabd6990 10.1126/scitranslmed

.abd6990

32 Zhang, J. J., Kobert, K., Flouri, T. & Stamatakis, A. PEAR: a fast and accurate I

llumina Paired-End reAd mergeR. *Bioinformatics* **30**, 614-620 (2014). https://doi.or

g:10.1093/bioinformatics/btt593

33 Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple sequenc

e alignments using Clustal Omega. *Mol Syst Biol* **7** (2011). https://doi.org:ARTN 5

39 10.1038/msb.2011.75

34 Sievers, F. & Higgins, D. G. Clustal Omega for making accurate alignments of m

any protein sequences. *Protein Sci* **27**, 135-145 (2018). https://doi.org:10.1002/pro.3

290

35 Lefranc, M. P. IMGT® databases, web resources and tools for immunoglobulin an

d T cell receptor sequence analysis. *Leukemia* **17**, 260–266 (2003).

36 Ye, J., Ma, N., Madden, T. L. & Ostell, J. M. IgBLAST: an immunoglobulin vari

able domain sequence analysis tool. *Nucleic Acids Res* **41**, W34-W40 (2013). https

://doi.org:10.1093/nar/gkt382

37 Andris-Widhopf, J., Steinberger, P., Fuller, R., Rader, C. & Barbas, C. F. Generati

on of Human scFv Antibody Libraries: PCR Amplification and Assembly of Light

- and Heavy-Chain Coding Sequences. *Cold Spring Harbor Protocols* **2011**, pdb.pr

ot065573 (2011). https://doi.org:10.1101/pdb.prot065573

Lee, Y., Kim, H. & Chung, J. An antibody reactive to the Gly63–Lys68 epitope of NT-proBNP exhibits O-glycosylation-independent binding. *Experimental & Molecular Medicine* **46**, e114-e114 (2014). https://doi.org:10.1038/emm.2014.57

C. F. Barbas, D. R. B., J. K. Scott, G. J. Silverman. *Phage Display: A Laboratory Manual*. Vol. 76 (2001).

Park, S., Lee, D. H., Park, J. G., Lee, Y. T. & Chung, J. A sensitive enzyme immunoassay for measuring cotinine in passive smokers. *Clin Chim Acta* **411**, 1238-1242 (2010). https://doi.org:10.1016/j.cca.2010.04.027

REED, L. J. & MUENCH, H. A SIMPLE METHOD OF ESTIMATING FIFTY PER CENT ENDPOINTS12. *American Journal of Epidemiology* **27**, 493-497 (1938). https://doi.org:10.1093/oxfordjournals.aje.a118408

Jiang, L. W. *et al.* Potent Neutralization of MERS-CoV by Human Neutralizing Monoclonal Antibodies to the Viral Spike Glycoprotein. *Sci Transl Med* **6** (2014). https://doi.org:ARTN 234ra59 10.1126/scitranslmed.3008140

Hoehn, K. B. *et al.* Repertoire-wide phylogenetic models of B cell molecular evolution reveal evolutionary signatures of aging and vaccination. *Proceedings of the National Academy of Sciences* **116**, 22664-22672 (2019). https://doi.org:10.1073/pnas.1906020116

Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nucleic Acids Res* **49**, W293-W296 (2021). https://doi.org:10.1093/nar/gkab301

Acknowledgements

This research was supported by the Korea Dementia Research Center (KDRC, HU20C0339), Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI, HI23C0521), the BK21 FOUR program, the National Research Foundation of Korea (NRF-2020R1A3B3079653 NRF-2020M3H1A1073304, NRF-2017M3A9G6068245, NRF-2022M3A9J1081343 and NRF-2023M3A9G6057281), the SNUH Research Fund (03-2021-0080), and the Inter-university Semiconductor Research Center. The pathogen resources (NCCP43326, NCCP43381, NCCP43382, NCCP43388, NCCP43390, NCCP43408) for this study were provided by the National Culture Collection for Pathogens. J. Choi is grateful for financial support from the Hyundai Motor Chung Mong-Koo Foundation.

Author contributions

S.P. designed and conducted all experiments, performed analyses, interpreted experimental results, and wrote and revised the paper. J. Choi performed the bioinformatic analyses, visualized and interpreted the results, and wrote and revised the paper. Y.L., J.N. and N.P.K. performed the bioinformatic analysis. J.L. conducted experiments related to the microneutralization assay. D.Y. conducted experiments related to the construction of repertoire libraries for NGS. G.C. and S.J.K. contributed to experiments related to the expression of recombinant proteins. K.K., P.C., N.J.K., and W.P. enrolled the study participants and collected the study samples. M.O. conceived and designed the study, and interpreted all results. S.T.K. supervised all the experiments with infectious viruses and microneutralization assay. S. Kwon conceived the study and designed and supervised the bioinformatics analysis. J. Chung conceived the study, designed and supervised all experiments, interpreted all results, and wrote the paper.

Competing interests

S.P., J. Choi, Y.L., J.N., N.P.K., J.L., G.C., W.P., S.T.K., M.O., S. Kwon and J. Chung are

inventors on a patent application for the antibodies described in this article. The remaining

authors declare no conflict of interest.

Additional information

Extended data

Extended Data Fig. 1 Plasma levels of antibodies against the SARS-CoV-2 N protein.

Reactivity of plasma from 41 vaccinees to the SARS-CoV-2 N protein (Sino Biological) was

tested by ELISA. The wells of microtiter plates were coated with the SARS-CoV-2 N protein and

blocked. After washing, plasma samples diluted in blocking buffer (1:2,500-fold) were added to

wells. The amount of bound antibody was determined using a horseradish peroxidase (HRP)-

conjugated rabbit anti-human IgG antibody (Invitrogen).

Extended Data Fig. 2 Plasma levels of antibodies against ancestral and BA.1 RBD. Plasma levels of antibodies to SARS-CoV-2 ancestral and BA.1 RBDs were measured in 41 vaccinees after (**a**) 100-fold or (**b**) 500-fold dilution. All experiments were performed in duplicate, and the average value for each vaccinee was plotted. The P value was calculated using one-way ANOVA. ns, not significant, $p > 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$. The increases and decreases in antibody levels are marked with red and green asterisks, respectively

Extended Data Fig. 3 Plasma levels of antibodies against the RBD of Omicron subvariant

BQ.1.1. Plasma levels of antibodies to the Omicron subvariant BQ.1.1. RBD (Sino Biological)

were determined after (**a**) 100-, (**b**) 500- or (**c**) 2,500-fold dilution. All experiments were

performed in duplicate, and the average value for each vaccinee was plotted. The P value was

calculated using one-way ANOVA. ns, not significant, $p > 0.05$; ***, $p < 0.001$; ****, $p <$

0.0001.

Extended Data Fig. 4 Characteristics of BA.1 RBD-binding scFv clones. **a,** Reactivity of

BA.1 RBD-binding scFv clones from the phage display library was tested against SARS-CoV-2

ancestral, Alpha, Beta, Gamma, Delta, and Omicron-sublineage (BA.1, BA.2, BA.4/5, BQ.1.1

and XBB.1.5) RBD proteins in the form of a recombinant scFv-hFc-HA protein. **b,** The

neutralization efficacy of BA.1 RBD-reactive scFv clones was tested in a microneutralization

assay using ancestral SARS-CoV-2 (βCoV/Korea/KCDC03/2020 NCCP43326), Alpha B.1.1.7

(hCoV-19/Korea/KDCA51463/2021 (NCCP 43381), Beta B.1.351 (hCoV-

19/Korea/KDCA55905/2021 (NCCP 43382), Gamma P.1 (hCoV-19/Korea/KDCA95637/2021

(NCCP 43388), Delta B.1.617.2 (hCoV-19/Korea/KDCA119861/2021 (NCCP 43390), and

Omicron B.1.1.529 (hCoV-19/Korea/KDCA447321/2021 NCCP43408) strains.

Extended Data Fig. 5 The characteristics of the BCR HC sequences of nine Omicron RBD-

reactive clonotypes mapped to the third and sixth repertoires. a, The isotype compositions of

the BCR HC sequences in the third and sixth repertoires (n = 57 and 216, respectively). **b,**

Numbers of SHMs in BCR HC sequences at the third and sixth repertoires (n = 57 and 216,

respectively). The P value was calculated using a two-tailed independent t test. ****, $p < 0.0001$.

c, Diversification of BCR HC sequences occurred from the third to the sixth repertoire through

SHM and HCDR3 sequence variation. The numbers of unique BCR HC sequences and HCDR3

sequences are counted at the nucleotide level. **d,** Numbers of SHMs in the BCR HC sequences of

clonotypes found in the third and sixth repertoires of the same vaccinee. The P value was

calculated using the Mann☐Whitney U test. In the case of repertoires for which statistical

significance is not indicated, statistical significance was not calculated, as the number of clones

included in the third or sixth repertoire was only one or two. ns, not significant, $p > 0.05$; *, $p <$

0.05; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

Extended Data Fig. 6 Determination of the threshold for RBD-reactive clones in the

HCDR3-randomized scFv phage display library. After phage ELISA using both ancestral and

BA.1 RBDs, the absorbance of each well was determined. The absorbance value was

approximated with a Gaussian distribution by the binned kernel estimation method under the

Gaussian kernel setting (black line). The chosen threshold absorbance value for a positive clone

was 0.81 (red vertical line), which corresponded to the valley point in the distribution plot.

Extended Data Fig. 7 The frequency distribution and sequence diversity of BCR-HC

XXA/V clonotypes after the first, second, and third injections. a, The frequency distribution

of BCR HC XXA/V clonotypes. The BCR HC XXA/V clonotypes with the same HCDR3 amino

acid sequence are plotted in the same color. **b,** Diversity of BCR HC XXA/V sequences in

individual repertoires. The numbers of unique BCR HC sequences and HCDR3 sequences are

counted at the nucleotide level.

Extended Data Fig. 8 a, BCR HC sequences of four clonotypes found in vaccinees in their third

and sixth BCR repertoires with the highest frequency. **b,** Reactivity of recombinant scFv-hFc-HA

proteins encoded by individual BCR HC sequences and the light chain gene of each scFv clone

to the ancestral and BA.1 RBDs.

Extended Data Fig. 9 The BCR HC sequences of the 27-60, 35-15, 35-46, 43-09, and 43-34

clonotypes-3[rd] found in vaccinees 35, 43, 35, 23, and 43.

a

27-60 No.35

Tree scale : 0.1

b

35-15 No.43

c

35-46 No.35

720 **Extended Data Fig.10 Phylogenetic trees of the 27-60, 35-15, 35-46, 43-09, and 43-34**

721 **clonotypes-3rd found in vaccinees 35, 43, 35, 23, and 43.** The BCR sequences that were

722 synthesized to confirm the reactivity to ancestral RBD and those of Omicron subvariants are

723 shown in bold.