Human antigen R differentially modulates the translation of SARS-CoV-2 genomic and

sub-genomic RNAs

Harsha Raheja[1], Risabh Sahu[1], Trinath Ghosh[2], Priya Rani[1], Biju George[1], Shashank

Tripathi[1,3] and Saumitra Das[1,2*]

1. Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore-

560012

2. National Institute of Biomedical Genomics, Kalyani, West Bengal

3. Centre for Infectious Disease Research, Indian Institute of Science, Bangalore-560012

* To whom correspondence should be addressed.

Prof. Saumitra Das

Department of Microbiology and Cell Biology

Indian Institute of Science, Bangalore-560012, India

Phone: +91 80 2293 2886, Fax: +91 80 2360 2697

Email: sdas@iisc.ac.in

Running title: Role of HuR in SARS-CoV-2 translation

Abstract

Severe acute respiratory syndrome-coronavirus-2 (SARS-CoV-2) viral RNA associates with different RNA-binding host proteins at each stage of its life cycle, although the potential role of human antigen R (HuR) is unclear. Here, we found that HuR bound the 5′-untranslated region (5′-UTR) of SARS-CoV-2, and knockdown and knockout studies revealed the importance of such binding in viral translation. We identified 5′-UTR mutations in SARS-CoV-2 variants of concern that altered the HuR-binding affinity. Interestingly, HuR promoted non-structural protein translation through the genomic 5′-UTR and suppressed the structural protein translation from the sub-genomic 5′-UTR, which required polypyrimidine tract-binding protein binding to the 5′-UTR. HuR knockout increased the sensitivity to remdesivir treatment by decreasing its half-maximal inhibitory concentration by 10-fold. An antisense oligonucleotide (whose binding site overlapped the HuR-binding site) reduced viral RNA production and viral titers in wild-type cells but not HuR-knockout cells, further suggesting that HuR binds the SARS-CoV-2 5′-UTR and promotes replication. Our results indicate that HuR supports SARS-CoV-2 life cycle by promoting differential translational reprogramming of genomic and sub-genomic RNAs, implying that HuR can potentially be targeted for therapeutic interventions.

Keywords 5′-untranslated region / genomic RNA / human antigen R / severe acute respiratory syndrome-coronavirus-2 / sub-genomic RNA / translation

Introduction

Severe acute respiratory syndrome-coronavirus-2 (SARS-CoV-2) is a positive-strand RNA virus with a 30-kilobase genome. The viral life cycle begins with attachment of the virus to the angiotensin-converting enzyme-2 (Ace-2) receptor. That interaction is followed by receptor-mediated endocytosis, where the virus particles are internalized into endosomes, which release viral RNA into the host cell cytoplasm. Viral RNA undergoes translation to generate a polyprotein that codes for nonstructural proteins. These non-structural proteins initiate replication on the viral 3′-untranslated region (3′-UTR), generating negative-strand genomic RNA. Multiple small RNA fragments coding for structural viral proteins are synthesized from negative-strand RNA and positive-strand genomic RNA. These are the sub-genomic RNAs and the structural proteins thus formed coat the viral positive-strand RNA to form infectious virions, which are released from cells[1].

SARS-CoV-2 RNA and proteins interact with several host proteins to modulate viral pathogenesis. The viral RNA possesses binding sites for cellular RNA-binding proteins (RBPs), which assist the above-mentioned processes at different levels[2,3]. The viral coding region is flanked by UTRs at both the 5′ and 3′ ends, which provide scaffolds for viral and cellular RBPs to bind and regulate translation, replication, or packaging. A dynamic association exists between RBPs and viral RNA during different stages of viral infection. High-throughput screens have identified many RBPs, such as IMP-1, DDX-1, and hnRNPs, that bind to SARS-CoV-2 genomic RNA[4,5]. Here, we studied RBPs predicted to bind to the viral 5′-UTR. Of these, we selected human antigen R (HuR; also known as ELAVL1) for further study.

HuR is an RBP that binds to AU-rich elements in the 3′-UTRs of target mRNAs and stabilizes mRNAs[6,7]. HuR shuttles between the nucleus and cytoplasm and participates in the cellular transport of various RNAs. Different post-translation modifications of HuR have

been linked to different functional roles and subcellular-localization sites in cells [8]. HuR can

regulate the replication of hepatitis C virus[9], coxsackievirus B3[10], and adenovirus[11]; stabilize

Sindbis virus RNA[12]; and modulate human immunodeficiency virus reverse transcription[13]. It

was also shown to bind the SARS-CoV-2 genome in genome-wide screens [4,14]. We confirmed

that HuR can bind the viral 5′-UTR through multiple independent assays and studied its

importance in the SARS-CoV-2 life cycle. We examined the potential roles of HuR in

regulating the translation of both structural and non-structural proteins, which are translated

from separate sub-genomic and genomic RNAs, respectively. Further, we blocked HuR–

SARS-CoV-2 genome interactions with specific antisense oligonucleotides (ASOs),

suggesting that they can be explored as therapeutics for treating COVID19 disease.

Results

HuR regulated the SARS-CoV-2 life cycle

The 5′-UTR of SARS-CoV-2 is 265 nucleotides (nt) long. Enzymatic probing showed that

the initial 300 nt of the SARS-CoV-2 genome at the 5′ end forms five stem-loop secondary

structures that are conserved across all emerging variants [15]. The RNA-binding protein

database (RBPDB) was used to identify trans-acting host factors that interact with the 5′-UTR

of SARS-CoV-2. Several proteins showed 100% relative binding scores, including A2BP1,

ZRANB2, sap-49, FUS, PUM2, SRSF9, MBNL1, KHSRP, RBMX, SFRS1, and HuR (Fig

S1A, B). Bioinformatics-based predictions of RBPs with the RBPDB were based on

matching linear sequence motifs.

To study the role of HuR in the viral life cycle, Ace-2- expressing HEK-293T cells

were transfected with HuR short-interfering RNA (siRNA), which partially knocked down

HuR expression by 70% (si HuR) or control siRNA (Nsp si) , and infected with SARS-CoV-2

at a multiplicity of infection (MOI) of 0.1. The cells were harvested 48 h post-infection, and

the viral titers in the supernatants of siHuR-treated cells and non-specific siRNA-treated cells

were compared. We observed that siHuR treatment decreased the viral titer by more than 1

log (Fig 1A, B), emphasizing its important role in the viral life cycle. Parallel experiments

were performed to verify the effect of HuR on the life cycle of different viral variants.

Silencing of HuR decreased the viral titers of all variants, but the effect was delayed with the

alpha variant, with no difference was observed after 24 h and a clear difference observed after

48 h (Fig 1C-E). To confirm the role of HuR, we generated HuR-knock out (KO) HEK-293T-

Ace-2 cells using a guide RNA targeting the HuR coding region. HuR KO was confirmed by

western blotting using an anti-HuR antibody (Fig 1F) and by sequencing (Fig 1G). We

infected wild-type (WT) HuR-positive cells and HuR-KO cells with SARS-CoV-2 at an MOI

of 0.1 and examined differences in the supernatant viral titers and cellular SARS-CoV-2

RNA levels. Knocking out HuR reduced the cellular viral RNA levels by >80% (Fig 1H) and

decreased the viral titer by >2 logs (Fig 1I). The abundances of SARS-CoV-2 sub-genomic

RNAs (sgRNAs) encoding the Spike and Nucleocapsid proteins were decreased similarly to

that of genomic RNA in the HuR KO cell line (Fig 1J).

HuR bound the 5′-UTR of SARS-CoV-2

To verify that HuR can bind the SARS-CoV-2 5′-UTR, we performed ultraviolet (UV)-

crosslinking assays with a SARS-CoV-2 5′-UTR probe (transcribed *in vitro* with P^{32}) and

recombinant HuR protein. Direct UV crosslinking showed that HuR bound the labeled

SARS-CoV-2 5′-UTR probe in a dose-dependent manner (Fig 2A). The interaction was

further confirmed by performing competitive UV-crosslinking experiments, where 100-fold

and 150-fold excesses of unlabeled self-RNA showed increasing competition, which was not

observed with an unlabeled non-specific RNA (Fig 2B). These results confirmed the *in vitro*

binding of HuR to the SARS-CoV-2 5′-UTR.

To verify the binding in cell lysates, we performed UV-crosslinking

immunoprecipitation (IP) experiments wherein HuR was pulled down from the cytoplasmic

S10 lysates following UV crosslinking with the SARS-CoV-2 5′-UTR, using an anti-HuR

antibody. Specific HuR binding to the SARS-CoV-2 5′-UTR was demonstrated by comparing

the electrophoretic bands between experimental (after crosslinking and IP) and input samples.

The band at approximately 36 kDa, which was absent from the flow-through beads, depicted

the HuR band in the cell lysate (Fig 2C). *Ex vivo* binding of HuR to the SARS-CoV-2 5′-

UTR in infected cells was demonstrated by performing IP-reverse transcription (RT)

experiments where HuR was pulled down using an anti-HuR antibody and the associated

SARS-CoV-2 5′-UTR was amplified using specific primers. Amplification of the SARS-

CoV-2 5′-UTR specifically in the pull-down fractions confirmed HuR binding in SARS-

CoV-2 5′-UTR in infected cells (Fig 2D).

HuR bound around the C241 site in the SARS-CoV-2 5′-UTR

To identify the HuR-binding site in the SARS-CoV-2 5′-UTR, we used a catRAPID fragment

software that considers the secondary structure when determining the binding affinity of a

protein. HuR showed very high affinity for the region spanning nts 150-250, which

corresponds to stem-loop 5 of the SARS-CoV-2 5′-UTR (Fig S2). Two HuR-binding motifs

(199-GUUU-202 and 237-GUUU-240) were found within the region spanning nts 150-250

nts, which overlapped with the loop regions of SL5A and SL5B, respectively. However, the

catRAPID fragment showed lower affinity for the third 8-GUUU-11 motif, which base paired

with the SL1 hairpin.

To assess the impact of HuR binding, we evaluated the presence of mutations in the 5′-

UTRs of different SARS-CoV-2 variants. We aligned the 5′-UTRs of the alpha, beta, gamma,

and delta variants and found two mutations between them. The alpha variant acquired a

C241U mutation, which was retained in the other variants. The beta and delta variants

acquired another mutation, namely G210U (Fig 3A).

We performed Surface Plasmon Resonance (SPR) studies to analyze the effect of these

mutations on the HuR-binding affinities (Fig 3B-D). Incorporation of the C241U mutation

decreased the binding affinity of HuR from 32.5 +/- 13 nM (WT strain) to 338.6 +/- 63 nM,

whereas the incorporation of G210U along with C241U increased back the binding affinity to

72.9 +/- 7.2 nM (Fig 3E). These results could indicate that the second mutation (G210U)

generated an additional binding site for HuR. These findings corroborate the variant data

shown in Figure 1, and the lower binding affinity of HuR to the alpha variant might explain

its delayed effect. Furthermore, the 5′-UTRs of the alpha and gamma variants contained the

same mutations, but the gamma variant did not show lesser HuR dependency at 24h post

infection.. The results could reflect other direct or indirect effects of HuR silencing. To

confirm the binding of HuR at these sites, we performed toe-printing assays with the WT 5′-

UTR, which revealed multiple binding sites, with C241 being the major binding site (Fig 3F).

These data provide insights into the differential roles of HuR in SARS-CoV-2 variants due to

the altered binding affinity of HuR.

158   HuR promoted translation from the SARS-CoV-2 5′-UTR

SARS-CoV-2 replication can be impacted through the regulation of many different steps in

the viral life cycle. Because HuR bound to the SARS-CoV-2 5′-UTR (where translation of

the genome begins), we investigated whether HuR regulates this step. To this end, we

generated a luciferase reporter construct wherein the viral 5′-UTR was followed by the

luciferase gene. Luciferase activities under different conditions revealed various degrees of

SARS-CoV-2 5′-UTR translation.

A549 cells were transfected with a 5′-UTR-luc construct in the context of either partial HuR knockdown (Fig 4A) or HuR overexpression (Fig 4B), and the luciferase activities were measured. To normalize the luciferase activities, the cells in each well were co-transfected with 2 ng of pRLTK (encoding Renilla luciferase), and dual-luciferase readings were taken after 8 or 12 h. Partial HuR knockdown reduced translation from the SARS-CoV-2 5′-UTR (Fig 4A), and HuR overexpression increased translation (Fig 4B). These results suggest that HuR promoted viral RNA translation by binding to the 5′-UTR. This effect was also observed in HuR-KO cells. The inclusion of SARS-CoV-2 5′-UTR in luciferase reporter construct showed increased luciferase activity in WT cells (lanes 1 and 2), suggesting that the presence of the SARS-CoV-2 5′-UTR might have recruited additional factors needed for enhancing cap-dependent viral translation. Furthermore, this increase was completely abrogated in HuR KO cells (lanes 3 and 4), implying the importance of HuR as the factor behind increased translation from the viral 5′-UTR (Fig 4C). Notably, translation from the parental luciferase construct was reduced in HuR KO cells, which could be represent an independent effect of HuR knockdown.

The role of HuR in SARS-CoV-2 translation was further studied by analyzing viral RNA loaded into the polysomes in WT cells containing WT and HuR-KO cells. The reduction in polysomal peaks suggested that SARS-CoV-2 infection led to a global reduction in translation (Fig 4D). This reduction was compromised in HuR-KO cells (Fig 4E). RNA was isolated from pooled monosome and polysome fractions, and the abundances of the SARS-CoV-2 5′-UTR were quantified by real-time polymerase chain reaction (PCR) analysis using specific primers. HuR knockout led to a 70% decrease in SARS-CoV-2 RNA loading into polysomes (Fig 4F, G). This finding confirms the important role of HuR in promoting translation of the SARS-CoV-2 5′-UTR.

HuR regulated the translation of SARS-CoV-2 sub-genomic RNAs

We observed that HuR KO led to decreased viral genomic and sgRNA production (Fig 1J).

Since they are independent RNA species with differences in the 5′-UTR, we examined the

effect of HuR KO on translation on the sg 5′-UTR. To this, end we infected WT and HuR KO

cells with SARS-CoV-2 at an MOI of 1.0 and harvested the cells and culture supernatants

after 24 h. We observed a >10-fold reduction in viral titers in HuR-KO cells (Fig 5A).

Furthermore, the genomic and sub-genomic RNA levels decreased to the same extent in

HuR-KO cells (Fig 5B). We quantified the expression levels of non-structural and structural

proteins (produced by the translation of genomic and sub-genomic 5′-UTRs, respectively) in

WT and HuR-KO cells by western blot analysis. The Nucleocapsid protein represents sub-

genomic translation, and the Nsp5 protein represents genomic translation. We observed that

the Nucleocapsid: Nsp5 protein ratio was higher in HuR-KO cells than in WT cells (Fig 5C),

even though no difference was evident in their RNA levels. These findings suggested that

both RNA species were differentially regulated at the level of translation. To study sub-

genomic RNA translation, we performed capped *in vitro* transcription of sub-genomic SARS-

CoV-2 5′-UTR (nts 1-70) followed by luciferase RNA and used *in vitro*-transcribed

luciferase RNA alone as a control. These RNAs were transfected into WT and HuR-KO

HEK-293T-Ace-2 cells, and luciferase activities were measured at 8 h post-transfection. As

observed previously, transfecting the parental luciferase reporter RNA alone exhibited lower

readings in the HuR-KO cell line. This decrease was rescued for sg 5′-UTR-luc, suggesting

the increase in translation from the sg 5′-UTR in the HuR-KO cell line (Fig 5D). These data

explain the increased ratio of structural to non-structural proteins in the HuR-KO cell line and

provide the first evidence of differential translational regulation of SARS-CoV-2 genomic

RNA and sgRNA by a host protein.

HuR regulated SARS-CoV-2 translation by assisting binding of the polypyrimidine
215

tract-binding (PTB) protein
216

To further investigate the mechanism of action of HuR, we analyzed the binding of different

cellular proteins to the 5′-UTR of SARS-CoV-2. We purified cellular S10 extracts from three

different cell lines (HEK-293T-Ace-2, A549, and Huh7.5) and analyzed their binding to the

SARS-CoV-2 5′-UTR by performing UV-crosslinking experiments. We observed that

multiple proteins bound to the 5′-UTR in a dose-dependent manner (Fig 6A). The band

marked with an asterisk has a mass that corresponded to the size of HuR, as verified in UV-IP

experiments (Fig 2C).

We checked whether the addition of exogenous HuR to the S10 lysate would alter the

binding to any other protein and observed a band, approximately 63kDa in size, whose

intensity increased with increasing concentrations of HuR (Fig 6B). We previously

characterized PTB as an important protein with a band in this size range[9]. Therefore, we

purified recombinant PTB and HuR proteins and performed competitive-crosslinking

experiments with increasing HuR concentrations, while keeping the PTB concentration

constant. HuR binding promoted PTB binding to the SARS-CoV-2 5′-UTR, corroborating

our previous data (Fig 6C). We analyzed whether this competition occurs *ex vivo* in cells by

performing IP-RT experiments, where PTB was immunoprecipitated from cells expressing

WT HuR and HuR-KO cells (Fig 6D). Thereafter, the associated 5′-UTR was amplified by

PCR. The absence of HuR in the knockout cell line prevented the PTB binding to 5′-UTR

(Fig 6D, upper panel), confirming that HuR functions by promoting the binding the PTB.

Because HuR is primarily a nuclear protein that relocalizes to the cytoplasm upon

infection with different viruses, we checked the localization of HuR at different time points

following infection by conducting immunofluorescence and nuclear-cytoplasmic

fractionation experiments (Fig S3). In the immunofluorescence assays, HuR localized to the

nucleus in mock-infected cells, and we did not observe any changes in HuR localization

when HEK-293T-Ace-2 cells were infected with SARS-CoV-2 at different timepoints (Fig

S3A). Nuclear-cytoplasmic fractionation experiments revealed that a fraction of HuR

localized to the cytoplasm and that its cytoplasmic abundance did not increase upon SARS-

CoV-2 infection. These findings suggest that cytoplasmic HuR is sufficient for promoting

PTB binding and influencing viral translation.

We then examined whether PTB overexpression could rescue the decrease in viral titer

in HuR-KO cells. PTB overexpression increased the viral titer by more than 1 log in both the

WT HuR and HuR-KO cell lines (Fig 6E). PTB overexpression demonstrated a similar

increase in viral RNA levels in both cell lines (Fig 6F, G). Luciferase constructs were used to

examine whether these effects were mediated through translational regulation. PTB

overexpression increased luciferase activity to a greater extent in the HuR-KO cell line (Fig

6H). These results suggest that the presence and binding of HuR to the SARS-CoV-2 5′-UTR

could guide PTB binding and thereby promote viral translation. The same effect was

observed by increasing PTB expression in cells, which would increase its binding propensity

and affinity.

HuR can potentially be used as a target for developing better therapeutics

Because HuR played an important role in guiding SARS-CoV-2 viral translation by binding

to viral RNA, we explored whether it could alter the activity of remdesivir in viral inhibition.

We used a concentration gradient of remdesivir and examined decreases in viral RNA in WT

and HuR-KO cells. We found that 200 nM remdesivir completely inhibited viral RNA

production in both cell lines; however, at lower remdesivir concentrations, the HuR-KO cell

line showed substantially lower viral RNA levels (Fig 7A). The half-maximal inhibitory

concentration (IC_{50}) of remdesivir was calculated based on the resulting data, and we found

that the IC_{50} was 10.46 nM in WT cells but only 1.82 nM in HuR-KO cells (Fig 7B). These

data suggest the potential of targeting HuR as an additional therapy in combination with

remdesivir.

Recently, ASOs directed against different regions of SARS-CoV-2 RNA have been

demonstrated to inhibit SARS-CoV-2 replication in cell culture and in animal models [16]. Of

the ASOs used, ASO5 targeted the HuR-binding site (encompassing 241 sites) and reduced

SARS-CoV-2 RNA levels. Therefore, we investigated the activity of ASO5 using WT and

HuR-KO cell lines. As observed previously, ASO5 transfection reduced viral RNA levels in a

dose-dependent manner in WT cells. However, such a decrease was not observed in the HuR-

KO cell line (Fig 7C), strengthening the evidence supporting the location of the HuR-binding

site, its role in the viral life cycle, and its use as a potential therapeutic target.

Discussion

Host proteins are important factors that regulate the virus life cycle. RNA-binding proteins

bind to viral RNAs at different stages of their life cycles, thereby modulating their cellular

interactions. RBP screening has identified common proteins involved in the life cycle of

multiple viruses. HuR is one such protein that has been shown to regulate the translation and

replication of viruses, including hepatitis C Virus, coxsackievirus, and Sindbis virus. We

studied the role of this important protein in regulating the SARS-CoV-2 life cycle. We found

that HuR bound to the viral 5′-UTR and regulated its life cycle by promoting translation. It

did so by promoting the binding of another RBP, PTB, which can enhance ribosome

recruitment to the viral 5′-UTR. HuR binding occurred near 241 site in the 5′-UTR, which is

mutated in some SARS-CoV-2 variants of concern. This mutation alters the binding affinity

for HuR and the dependency of the alpha variant on HuR, which could be one of the

mechanisms behind the longer generation time and hence lesser transmission of the alpha

variant as compared to the delta variant[17]. Knocking out HuR increased the sensitivity of

infected cells towards remdesivir treatment for viral clearance. The HuR-binding site lies in

the target region of anti-sense oligo ASO5, which inhibits the viral life cycle and therefore is

an attractive candidate for add-on therapy. We further addressed the HuR-mediated

translational regulation of SARS-CoV-2 genomic and sgRNAs. We showed that while HuR

promoted translation from the genomic 5′-UTR, it inhibits sg 5′-UTR translation. Previous

reports have demonstrated the translational regulation by viral proteins such as Nsp1[18,19] and

ORF6[20], and we propose that HuR could represent the cellular arm of such regulation. The

genomic and sub-genomic 5′-UTRs vary in the length. The sub-genomic UTR harbors one of

the 4 predicted HuR binding sites (8-10) and that site resides in the stem region of the

secondary structure while the other 3 sites, which are present only in the FL UTR reside in

the loop region in the secondary structure [15]. This could be responsible for the differential

effect of HuR binding. Furthermore, it could also affect the PTB recruitment capability of

HuR, altering the translation regulation.

HuR does not relocalize to the cytoplasm upon SARS-CoV-2 infection in HEK-293T-

Ace2 cells, but the small amount of HuR present in the cytoplasm due to nuclear-cytoplasmic

shuttling might be sufficient to promote PTB binding, which appears to relocalize to the

cytoplasm upon infection. This possibility was supported by our competition-crosslinking

assays, where a small quantity of HuR greatly enhanced PTB binding. Furthermore, HuR is

known to alter the stability of cellular transcripts. The cytokine storm and disease severity

observed upon SARS-CoV-2 infection could be regulated by HuR by virtue of its binding to

mRNAs encoding pro-inflammatory cytokines such as interleukin (IL)-6 and IL-8[21,22]. We

also observed that HuR levels increased following SARS-CoV-2 infection. The regulatory

mechanism could be interesting, because miRNAs, especially miR-125b-5p, have been

shown to target HuR translation[23].

Taken together, our observations provide important insights and open multiple avenues

for targeting the SARS-CoV-2 life cycle by modulating an important host factor, HuR.

Targeting HuR through ASOs could prevent its binding to the genomic 5′-UTR while

increasing its ability to bind to the sub-genomic SARS-CoV-2 5′-UTR, which would decrease

structural protein translation and thereby help in clearing SARS-CoV-2 infection.

Materials and Methods

Plasmids and constructs: To make the template for *in vitro* transcription, the SARS-CoV-2

5'UTR was amplified with 5'AATT<u>GCTAGC</u>ATTAAAGGTTTATACCTTCCCAGGTA3'

and 5'AATT<u>AAGCTT</u>CTCGTTGA AACCAGGGACAAG3' from pCBB 5'UTR (Kind gift

from Dr. Milan Surjit, THSTI) and subcloned at the downstream of T7 promoter in NheI and

HindIII cloning site of pcDNA3.1+. For luciferase assay, SARS-CoV-2 5'UTR was

amplified   with   5'AATT<u>AAGCTT</u>ATTAAAGGTTTATACCTTCCCAGGTAAC3'   and

5'AATT<u>CCATGG</u>TTCTCGTTGAAACCAGGGACAAG 3' to subclone in pGL3 control

vector.

pcDNA3.1-myc-HuR and pET28a-HuR were generated as described previously.

pcDNA3.1-myc-PTB was a kind gift from Dr. Douglas Black's laboratory.

Cell lines and transfection: HEK-293T cells expressing Ace-2 (obtained from BEI

resources), A549(ATCC) lung epithelial cells and Vero E6 cells were grown in DMEM

media (Gibco) supplemented with 10% FBS (Gibco). For luciferase assays in A549 cells, day

before transfection 1.3×10^5 cells were distributed per well of 24 well tissue culture plate.

Before transfection DMEM was replaced with Opti-MEM I reduced serum media (Gibco).

Lipofectamine 2000 (Invitrogen) was used for all transfection experiment as per

manufacturer's protocol. To check the effect of HuR overexpression, we co-transfected

250ng of pGL3-SARS-CoV-2 5'UTR with either 250ng of pcDNA3.1 or pCDNA3.1-HuR.

pGL3 control vector was used for control set. 2ng pRLTK was also co-transfected in each

well to normalize the firefly luciferase activity. For HuR knockdown experiment we

transfected 150nM of either DsiHuR: 5'AUUUCUGAAUCUGUGACGCAAGAAT 3'(IDT)

or Nonspecific (Eurogentec) siRNA 24 hrs prior to luciferase construct transfection.

Virus stock preparation and infection.

For virus experiments, HEK-293T-Ace2 cells were seeded in 24-well plates. The cells were transfected with 150 nM of Nsp si or si HuR using Lipofectamine 2000 as per the manufacturer's protocol. 16h after transfection, cells were infected with virus stock at an MOI of 0.1 or 1 as described. Virus stocks were obtained from BEI resources and maintained by Viral BSL3 repository at IISc.

Dual luciferase assay: Luciferase assay was done by dual luciferase assay kit (Promega) according to manufacturer's protocol. For HuR overexpression and siHuR experiment, assays were done 12 and 8 hrs post-transfection of luciferase constructs, respectively. Luciferase reading was taken with Glomax Luminometer (Promega).

Protein purification: *E. coli BL21* cells were transformed with pET28a containing His tagged HuR. Recombinant protein was purified with Ni-NTA agarose. The detail of purification was described earlier. Briefly, the expression of HuR was induced at an optical density of 0.6 at 600 nm with 0.5 mM isopropyl-1-thio-β-D-galactopyranoside (IPTG) for 4h. The cells were pelleted and resuspended in lysis buffer (50 mM Tris, pH 7.5, 300 mM NaCl, 0.1 mM phenylmethylsulfonyl fluoride [PMSF]) containing 1X Bacterial protease Inhibitor (Sigma) and lysed by sonication on ice. The supernatant obtained thereafter was incubated with Ni-NTA slurry for 3 h and bound protein was eluted with 500 mM Imidazole after washing the beads with lysis buffer containing 40 mM imidazole. The eluted protein was dialysed in 10 times volume dialysis buffer (25 mM Tris, pH 7.4, 100 mM KCl, 7 mM β-mercaptoethanol [β-ME], 10% glycerol), aliquoted and stored in -70℃.

In vitro **transcription:** The pcDNA3.1-SARS-CoV-2 5'UTR was linearized with HindIII digestion and purified with phenol-chloroform method. The linearised DNA was used as a template for synthesis of RNA using T7 RNA Polymerase (ThermoScientific) and ^{32}P uridine triphosphate. The transcription reactions were carried out under standard conditions at 37 ºC for 1.5 h. After alcohol precipitation, RNA was resuspended in nuclease free water. For radiolabelled RNA, 1 μl the prepared RNA was spotted onto DE81 filter paper, washed with phosphate buffer and dried and incorporated radioactivity was measured using scintillation counter.

UV induced crosslinking of RNA and protein: UV-induced cross-linking was carried out as described previously. Briefly, - ^{32}P-labeled RNA probes were allowed to form complexes with recombinant HuR in RNA binding buffer (5 mM HEPES, pH 7.6, 25 mM KCl, 2 mM

MgCl2, 3.8% glycerol, 2 mM dithiothreitol [DTT], and 0.1 mM EDTA) and then UV

irradiated for 20 min. The mixture was treated with 50 µg of RNase A (Sigma), separated on

an SDS–10% polyacrylamide gel (SDS-PAGE), and band analyzed by phosphorimaging.

Surface Plasmon Resonance: Surface plasmon resonance spectroscopy was performed

using a BIAcore3000 optical biosensor (GE Healthcare Lifescience) to study the binding

kinetics of HuR with SARS CoV2 5'UTR RNA. Biotin-labeled SARS CoV2 5'UTR RNA

was immobilized on streptavidin-coated sensor chips (GE Healthcare Lifescience) to a final

concentration of 300 Resonance Units (RU)/flow cell. RNA–protein interactions were carried

out in a continuous flow of Tris buffer (25 mM Tris (pH 7.5), 100 mM KCl, 7 mM β-

mercaptoethanol, and 10% glycerol) at 25 °C at a flow rate of 10 µl/min. Increasing

concentrations of HuR protein loaded on the biosensor chip for 100 s (characterized as the

association phase), followed by a dissociation phase of 300 s with buffer alone. For

normalising background non-specific interaction, a blank surface without any RNA was used

for simultaneous injections of the sample during the experiment. BIAevaluation software

(version 3.0) was used to determine the on rate, $k$on (M^{-1} s^{-1}), and off rate, $k$off (s^{-1}), using a

1:1 Langmuir binding model. The binding affinity, Kd was determined using the following

equation: $Kd = k$off/$k$on.

Plaque forming assay: Plaque assay was performed in Vero E6 cells in 12- well plates.

$0.5*10^6$ cells were seeded to give 90% confluency. After 12-16 h, cells were incubated with

100 µl of increasing dilution of supernatant obtained from Nsp si or si HuR treated Ace2-

HEK-293T cell infected with SARS-CoV2 or supernatant from mock infected cells. After 1 h

of incubation with the cells at 37°C, viral inoculum was removed and cells overlayed with

DMEM containing 0.8% Agarose and kept at 37°C. After 48 h of incubation, cells were fixed

with 4% paraformaldehyde for 1 h and stained with Crystal violet for 20 mins. Overlay was

removed and the number of plaques counted. Following calculation was used to determine

PFU/ml: PFU/ml= No. of plaques ÷ (Dilution factor × Vol of virus inoculum in ml)

Immunofluorescence: For immunofluorescence staining, HEK-293T-Ace2 cells were

seeded in a 24-well plate on coverslips coated with poly-L-lysine for 14-16 h followed by

infection in SARS-CoV2 virus at an MOI of 0.1. After desired time of infection, cells were

washed twice with 1X PBS and fixed using 4 % formaldehyde at room temperature for 20

min. After permeabilization by 0.1 % Triton X-100 for 5 min at room temperature, cells were

incubated with 3 % BSA at 37 °C for 1 h followed by incubation with the indicated antibody

for 2 h at 4 °C and then detected by Alexa-633-conjugated anti-mouse or Alexa-488

conjugated anti-rabbit secondary antibody for 30 min (Invitrogen). Images were taken using

Zeiss microscope and image analysis was done using the Zeiss LSM or ZEN software tools.

HuR knockout cell line preparation: The guide RNA targeting HuR sequence were

designed using Zhang lab software. These gRNAs were cloned in the pSpCas9(BB)-2A-GFP

(PX458) vector as per the described protocol. The clones were transfected in HEK-293T-

Ace-2 cells and 48h post transfection, the cells expressing GFP were sorted as one cell per

well in a 96-well plate. These cells were grown and analysed for HuR knockout using

western blotting and confirmed though sequencing. For control, untransfected cells were also

sorted as one cell per well in parallel and one of the clones, expressing Ace-2 receptor was

used.

gRNA1: ACCACATGGCCGAAGACTGC

gRNA2: CTTATTCGGGATAAAGTAGC

IP-RT: HEK-293T-Ace-2 cells were infected with the SARS-CoV-2 virus at an MOI of 0.1

and after completion of the experiment, 1% formaldehyde was added to the media and

incubated at 4 °C for 10 mins. Thereafter, the reaction was quenched with 0.1 M Glycine

incubated at 4 °C for 10 mins. The cells were then washed with PBS and lysed in polysome

lysis buffer (100 mM KCl, 10 mM HEPES pH 7.0, 5 mM $MgCl_2$, 0.5 % NP-40, 1 mM DTT,

100 U/ml RNasin). The supernatant obtained was precleared with Protein G Sepharose beads.

In parallel, the Protein G Sepharose beads were incubated with 1 ug of HuR/ PTB antibody

overnight at 4 °C and added to the precleared lysates followed by overnight incubation with

continuous mixing on a rotator device at 4°C. The beads were then washed three times with

polysome lysis buffer. SDS sample buffer was added to the 20% beads and boiled to release

the immunoprecipitated protein, and the supernatant was electrophoresed on SDS-12 %

PAGE. The RNA was isolated from remaining 80% beads. The beads were incubated with

0.1% SDS and 30 µg proteinase K at 50 °C for 30 mins and RNA isolated after addition of 3

times volume of Tri reagent.

RNA isolation followed by Real time PCR: The total RNA from Ace2-HEK-293T cells

was isolated using TRI Reagent (Sigma) as per manufacturer's protocol. cDNA was

synthesised using following primers: CoV2 Fwd: 5' TGTCGTTGACAGGACACGAG 3',

CoV2   Rev: 5'   TTACCTTTCGGTCACACCCG   3',   GAPDH   Fwd: 5'

CAGCCTCAAGATCATCAGCAAT 3', GAPDH Rev: 5' GGTCATGAGTCCTTCCACGA

3'. For quantification of viral positive strand, CoV2 Rev and GAPDH Rev primers were used

for cDNA synthesis. For quantification of viral negative strand, CoV-2 Fwd and GAPDH

Rev primers were used for cDNA synthesis. Moloney murine leukemia virus(M-MLV)

reverse transcriptase was used for cDNA synthesis at 42°C for 1 h using 600 ng of total RNA.

Quantitative RT-PCR (qRT-PCR) was done using DyNAmo HS SYBR green qPCR kit

(ThermoScientific). qRT-PCR was done using 2 µl of cDNA in a 10-µl reaction mixture

according to the manufacturer's instructions for 40 cycles. The comparative threshold cycle

(CT) method was used to calculate fold change in SARS CoV2 RNA levels (2^ΔΔCT) and

normalized to GAPDH.

Western blotting: Protein concentrations of the extracts were assayed by Bradford reagent

(Bio-Rad) and equal amounts of cell extracts were separated by SDS-10 % PAGE and

transferred onto a nitrocellulose membrane (Sigma). Samples were then analyzed by western

blot using the desired antibodies, anti-SARS-CoV-2 N protein (40143-MM05, Sino

Biological), anti-HuR antibody (3A2, Santa Cruz), anti-Myc tag antibody (ab9106) or anti-

Ace-2 antibody (ab15348) followed by the respective secondary antibodies (horseradish

peroxidase-conjugated anti-mouse or anti-rabbit IgG; Sigma). Mouse-monoclonal anti-β-

actin- peroxidase antibody (A3854, Sigma) was used as a control for equal loading of total

cell extracts. Antibody complexes were detected using the ImmobilonTM Western systems

(Millipore).

Sucrose gradient polysome fractionation: After completion of the experiment, cells were

treated with 100 µg/ml cycloheximide for 10min at 37°C. Thereafter, media was removed

and cells washed first with ice cold PBS containing 100µg/ml cycloheximide and then with

1X hypotonic buffer (5 mM TRIS-HCl pH-7.5, 5 mM MgCl$_2$ and 1.5 mM KCl) containing

100 µg/ml cycloheximide. Cells were scraped in 350µL ice cold lysis buffer (5 mM TRIS-

HCl pH-7.5, 5 mM MgCl$_2$, 1.5 mM KCl, 100 µg/ml cycloheximide, 1mM DTT, 200 U/ml

RNase in from Promega, 0.5% Sodiumdeoxycholate, 0.5% Triton X -100, 200µg t- RNA and

1X protease inhibitor cocktail) and incubated for 15 min on ice. The KCl concentration in the

lysate was adjusted to 150 mM. The lysate was spun for 8min at 3000 g at 4°C and

supernatant collected. 500 µg of the lysate was loaded on the top of 15–50% sucrose gradient

containing 100µg/ml cycloheximide and the gradients were centrifuged at 36,000 rpm for 2 h

at 4°C in SW41 rotor (Beckman). Density Gradient Fractionation System (ISCO) was used to

fractionate the gradients at a flow rate of 0.3 mm/sec with the UV detector sensitivity set at

1.0. Monosome and Polysomes fractions were pooled to isolate RNA using Trizol (Sigma).

SARS-CoV-2 RNA level was quantified using real time PCR.

Toeprinting: The Unlabeled SARS-CoV2 5′-UTR RNA (10 pmoles) was incubated in

presence of BSA or increasing concentration of purified recombinant HuR protein and then

annealed to an end labeled primer (100 fmoles) with sequence complimentary to the 3′ end of

the RNA. AMV-RT (Promega) was used to reverse-transcribe the RNA-protein complex

reaction at 30 °C for 1 h. This was followed by denaturation of the enzyme at 95 °C for

5 min. 0.3 M sodium acetate and 2.5 volumes of absolute alcohol was used to pellet the

cDNA, which was resuspended in DEPC treated water and resolved onto a 7 M urea 6%

polyacrylamide sequencing gel. In parallel, sequencing reaction (fmol kit, Promega) using the

same end labeled primer was run to identify the exact position of RT stops.

Primers for PCR:

Genomic 5'UTR Forward: TGTCGTTGACAGGACACGAG

Genomic 5'UTR Rev: TTACCTTTCGGTCACACCCG

N sgmRNA LS Forward: CGATCTCTTGTAGATCTGTTC

N sgmRNA Rev: AGCGGTGAACCAAGACGCA

S sgmRNA LS Forward: CCAACTTTCGATCTCTTGTAG

S sgmRNA Rev: AGAACAAGTCCTGAGTTGAATG

Primers for HuR gene PCR:

Forward primer: TACTTGCTCTTTTTCTCTTGGC

Reverse primer: CACAGCCATCGTTTCAAGGC

Anti-sense oligos:

Ctrl ASO: CGTTAGATTACCGCG

ASO5: CAACACGGACGAAACC

Softwares: All emerging lineage of SARS-CoV-2 were downloaded from Global Initiative

on Sharing All Influenza Data (GISAID, https://www.gisaid.org/). For Multiple sequence

alignment Clastal Omega online tool (https://www.ebi.ac.uk/Tools/msa/clustalo/) was used.

To predict the secondary structure RNAstructure online tool was used (https://rna.urmc.

rochester.edu/RNAstructureWeb/Servers/). RNA binding protein was predicted with RNA

Binding Protein Database (RBPDB, http://rbpdb.ccbr.utoronto.ca/). Binding region on the

RNA sequence was predicted with Cat-Rapid fragment

(http://service.tartaglialab.com/page/catrapid_group).

Data was plotted using GraphPad Prism.

Acknowledgements

SD acknowledges the J.C. Bose Fellowship from Department of Science and Technology

(DST), India, for research support. This study was also supported by the DBT-IISc

partnership program, DST Fund for Improvement of Science and Technology Infrastructure

(DST- FIST) level II infrastructure, and the University Grants Commission Centre of

Advanced Studies. HR is supported by the research fellowship from the Council of Scientific

and Industrial Research (CSIR-SPM). We recognize the SPR facility at the Biological

Science division at IISc. We acknowledge Prof. Adolfo Garcia-Sastre, Icahn School of

Medicine, for his insightful inputs. We acknowledge viral BSL3 repository at IISc for

maintaining SARS-CoV-2 virus stocks. We would like to thank Editage (www.editage.com)

for English language editing.

Disclosure and competing interests statement

The authors declare no competing interests.

References

1   V'kovski, P., Kratzel, A., Steiner, S., Stalder, H. & Thiel, V. Coronavirus biology and replication:
526   implications for SARS-CoV-2. *Nature Reviews Microbiology* **19**, 155-170, doi:10.1038/s41579-
527   020-00468-6 (2021).
528   2   Embarc-Buh, A., Francisco-Velilla, R. & Martinez-Salas, E. RNA-Binding Proteins at the Host-
529   Pathogen Interface Targeting Viral Regulatory Elements. *Viruses* **13**, doi:10.3390/v13060952
530   (2021).

3 Lisy, S., Rothamel, K. & Ascano, M. RNA Binding Proteins as Pioneer Determinants of
Infection: Protective, Proviral, or Both? *Viruses* **13**, 2172, doi:10.3390/v13112172 (2021).
4 Kamel, W. *et al.* Global analysis of protein-RNA interactions in SARS-CoV-2-infected cells
reveals key regulators of infection. *Molecular cell* **81**, 2851-2867.e2857,
doi:10.1016/j.molcel.2021.05.023 (2021).
5 Gordon, D. E. *et al.* A SARS-CoV-2 protein interaction map reveals targets for drug
repurposing. *Nature* **583**, 459-468, doi:10.1038/s41586-020-2286-9 (2020).
6 Mukherjee, N. *et al.* Integrative Regulatory Mapping Indicates that the RNA-Binding Protein
HuR Couples Pre-mRNA Processing and mRNA Stability. *Molecular cell* **43**, 327-339,
doi:https://doi.org/10.1016/j.molcel.2011.06.007 (2011).
7 Lebedeva, S. *et al.* Transcriptome-wide Analysis of Regulatory Interactions of the RNA-
Binding Protein HuR. *Molecular cell* **43**, 340-352,
doi:https://doi.org/10.1016/j.molcel.2011.06.008 (2011).
8 Grammatikakis, I., Abdelmohsen, K. & Gorospe, M. Posttranslational control of HuR function.
Wiley Interdiscip Rev RNA **8**, 10.1002/wrna.1372, doi:10.1002/wrna.1372 (2017).
9 Shwetha, S. *et al.* HuR Displaces Polypyrimidine Tract Binding Protein To Facilitate La Binding
to the 3′ Untranslated Region and Enhances Hepatitis C Virus Replication. **89**,
11356-11371, doi:doi:10.1128/JVI.01714-15 (2015).
10 George, B., Dave, P., Rani, P., Behera, P. & Das, S. Cellular Protein HuR Regulates the
Switching of Genomic RNA Templates for Differential Functions during the Coxsackievirus B3
Life Cycle. *J Virol* **95**, e0091521, doi:10.1128/jvi.00915-21 (2021).
11 Jehung, J. P. *et al.* Adenovirus infection induces HuR relocalization to facilitate virus
replication. *Biochemical and biophysical research communications* **495**, 1795-1800,
doi:10.1016/j.bbrc.2017.12.036 (2018).
12 Sokoloski, K. J. *et al.* Sindbis virus usurps the cellular HuR protein to stabilize its transcripts
and promote productive infections in mammalian and mosquito cells. *Cell host & microbe* **8**,
196-207, doi:10.1016/j.chom.2010.07.003 (2010).
13 Lemay, J. *et al.* HuR interacts with human immunodeficiency virus type 1 reverse
transcriptase, and modulates reverse transcription in infected cells. *Retrovirology* **5**, 47,
doi:10.1186/1742-4690-5-47 (2008).
14 Eskandarzade, N. *et al.* Network for network concept offers new insights into host- SARS-
CoV-2 protein interactions and potential novel targets for developing antiviral drugs. *Comput*
Biol Med **146**, 105575-105575, doi:10.1016/j.compbiomed.2022.105575 (2022).
15 Miao, Z., Tidu, A., Eriani, G. & Martin, F. Secondary structure of the SARS-CoV-2 5'-UTR. *RNA*
biology **18**, 447-456, doi:10.1080/15476286.2020.1814556 (2021).
16 Zhu, C. *et al.* An intranasal ASO therapeutic targeting SARS-CoV-2. *Nature Communications*
13, 4503, doi:10.1038/s41467-022-32216-0 (2022).
17 Hart, W. S. *et al.* Generation time of the alpha and delta SARS-CoV-2 variants: an
epidemiological analysis. *The Lancet Infectious Diseases* **22**, 603-610,
doi:https://doi.org/10.1016/S1473-3099(22)00001-9 (2022).
18 Tidu, A. *et al.* The viral protein NSP1 acts as a ribosome gatekeeper for shutting down host
translation and fostering SARS-CoV-2 translation. *Rna* **27**, 253-264,
doi:10.1261/rna.078121.120 (2020).
19 Schubert, K. *et al.* SARS-CoV-2 Nsp1 binds the ribosomal mRNA channel to inhibit
translation. *Nature Structural & Molecular Biology* **27**, 959-966, doi:10.1038/s41594-020-
0511-8 (2020).
20 Kehrer, T. *et al.* Impact of SARS-CoV-2 ORF6 and its variant polymorphisms on host
responses and viral pathogenesis. *bioRxiv*, 2022.2010.2018.512708,
doi:10.1101/2022.10.18.512708 (2022).
21 Srikantan, S. & Gorospe, M. HuR function in disease. *Front Biosci (Landmark Ed)* **17**, 189-205,
doi:10.2741/3921 (2012).

22   Banerjee, A. *et al.* Experimental and natural evidence of SARS-CoV-2-infection-induced
583   activation of type I interferon responses. *iScience* **24**, 102477,
584   doi:https://doi.org/10.1016/j.isci.2021.102477 (2021).
585   23   Shwetha, S. *et al.* Interaction of miR-125b-5p with Human antigen R mRNA: Mechanism of
586   controlling HCV replication. *Virus Res* **258**, 1-8, doi:10.1016/j.virusres.2018.09.006 (2018).

Figure legends

Fig 1. Role of HuR in SARS-CoV-2 lifecycle. (A) HEK-293T-Ace2 cells were transfected with either non-specific (Nsp si) or siRNA targeting HuR. 16 hrs post transfection, cells were infected with 0.1 MOI of SARS-CoV-2 virus. Viral titre in supernatant was determined using plaque assay. (B) Western blotting was done to confirm HuR silencing using anti-HuR antibody. (C) Effect of HuR silencing on different SARS-CoV-2 VoCs. HEK-293T-Ace2 cells were transfected with either non-specific siRNA (Nsp si) or siRNA targeting HuR (si HuR). 16 hrs post transfection cells were infected with different SARS-CoV-2 VoCs at 0.1 MOI. 48 hrs post infection cell supernatant was collected to determine the viral titres by plaque assay. (D) Representative images for plaque assay with VOCs. (E) Western blotting using anti-HuR antibody was performed for confirmation of HuR silencing. (F) HuR KO was confirmed by western blotting using anti-HuR antibody. (G) Sequence alignment of HuR gene being targeted by the CRISPR gRNA. The blanks in KO sequence represent the deletion of the region between the two gRNAs. (H) Positive strand RNA levels were determined using primers specific for SARS-CoV-2 5'UTR from WT and HuR KO cells harvested after 48 hrs of infection. (I) WT HuR and HuR KO HEK-293T-Ace-2 cells were infected with SARS-CoV-2 virus at 0.1 MOI and 48 hrs post infection, viral titre in supernatant was determined using plaque assay. (J) RNA levels of genomic 5'UTR and that of sub-genomic RNAs for Spike and Nucleocapsid were quantified using specific primers.

Fig 2. HuR binds to SARS-CoV-2 5'UTR. (A, B) *in vitro* interaction of recombinant HuR with radiolabelled SARS-CoV-2 5'UTR using UV-crosslinking assay. (C) UV-crosslinking of S10 extract with radiolabelled SARS-CoV-2 5'UTR, followed by immune pull down of HuR. (D) *ex vivo* interaction of HuR with CoV-2 5'UTR in SARS-CoV-2 virus infected cells by immune-pulldown using anti-HuR antibody. The upper panel denoted PCR amplification of CoV-2 5'UTR and bottom panel represents the western blot for HuR pull-down.

Fig 3. HuR binding site on SARS-CoV-2 5'UTR. (A) Multiple sequence alignment of 5'UTR of alpha, beta, gamma and delta variants of SARS-CoV-2. (B, C, D) Sensorgrams for

indicated biotinylated SARS-CoV-2 5'UTR binding to recombinant HuR using SPR. Colors

denote increasing HuR concentration (E) Binding affinity of mutants obtained by SPR. (F)

Toe-printing assay was performed with end labelled primer complementary to 3' end of the

SARS-CoV-2 5'UTR and increasing concentration of HuR. BSA was used as a control for

nonspecific binding. Sequencing reaction was conducted in the presence of ddNTPs and the

bases labelled on the top represent the base complementary to the ddNTP used in a particular

reaction. The asterisks denote toeprints around C241 in 5'UTR.

Fig 4. HuR promotes SARS-CoV-2 translation by binding to its 5'UTR. (A) A549 cells

were transfected with either non-specific (Nsp si) or siRNA targeting HuR (siHuR). 16 hrs

post transfection, cells were transfected with either pGL3-SV40-Luc (Control) or pGL3-

SV40-SARS-CoV-2 5'UTR-Luc (SARS-CoV-2 Luc) and harvested after 12 hrs of

transfection and luciferase readings measured. pRLTK was co-transfected for transfection

control and used for normalization of luciferase values. Effect of si HuR was calculated with

respect to the luciferase in Nsp si condition. (B) HuR overexpression was performed in place

of silencing in the previous panel. Effect of overexpression was calculated with respect to

luciferase in no overexpression condition. (C) WT HuR and HuR KO HEK-293T-Ace2 cells

were transfected with either pGL3-SV40-Luc (Luc alone) or pGL3-SV40-SARS-CoV-2

5'UTR-Luc (CoV-2 Luc) and luciferase values were measured 24 hrs post-transfection. The

graph represents absolute values obtained upon normalization of luciferase readings with

protein concentration. (D) WT HEK-293T-Ace2 cells were infected with SARS-CoV-2 virus

at 0.1 MOI and 48 hrs post infection cells were harvested for sucrose gradient polysome

fractionation. The lines represent the absorbance at 254 nm v/s the distance from the top of

gradient. (E) HuR KO HEK-293T-Ace2 cells were infected with SARS-CoV-2 virus at 0.1

MOI and 48 hrs post infection cells were harvested for sucrose gradient polysome

fractionation. The lines represent the absorbance at 254 nm v/s the distance from the top of

gradient. (F) Western blotting was performed for the lysates used for polysome fractionation

using anti-HuR antibody. (G) RNA was isolated from monosome and polysomal fractions of

infected lysate from panel (D) and (E) and the abundance of SARS-CoV-2 5'UTR quantified

by specific primers in real time PCR.

Fig 5. HuR inhibits translation of SARS-CoV-2 sub-genomic RNAs. (A) WT HuR and

HuR KO HEK-293T-Ace-2 cells were infected with SARS-CoV-2 virus at MOI of 1.0 and

24 h post infection, viral titre in supernatant was determined using plaque assay. The below

panel shows representative plaque assay images for the same. (B) RNA levels of genomic

5'UTR and that of sub-genomic RNAs for Spike and Nucleocapsid were quantified in the

same experiment using specific primers. (C) Protein levels of Nucleocapsid and Nsp5 were

determined in the above experiment by western blotting using specific antibodies. Ponceau

staining of the bot represents equal loading. The values at the bottom represent densitometry

values after quantifying the intensities of Nucleocapsid and Nsp5 bands. (D) WT HuR and

HuR KO HEK-293T-Ace2 cells were transfected with either *in vitro* transcribed capped

Luciferase RNA (Luc alone) or *in* vitro transcribed capped sub-genomic 5'UTR followed by

Luciferase RNA (sg 5'UTR luc) and luciferase values were measured 8 h post-transfection.

The graph represents Fold change in luciferase values obtained upon normalization of

luciferase readings with protein concentration.

Fig 6. HuR assists PTB binding to 5'UTR to promote translation. (A) UV-crosslinking

profiles with increasing amount of S10 extracts of different cell lines as indicated. Asterisk

indicates the band at the size of HuR. (B) UV-crosslinking was performed with S10 extracts

of different cell lines and increasing amount of HuR. Asterisk indicates the band at the size of

PTB. (C) Competition UV-crosslinking was performed with constant amount of PTB and

increasing amounts of recombinant HuR. (D) HEK-293T-Ace-2 with WT and HuR KO were

infected with 0.1 MOI of SARS-CoV-2. 48 hrs post infection, cells were harvested and PTB

immunoprecipitated. The associated RNA was amplified by PCR using SARS-CoV-2 5'UTR

specific primers (Upper panel). The immune pull down was confirmed by western blot using

anti-PTB antibody (Lower panel). (E) WT and HuR KO HEK-293T-Ace2 cells were

transfected with myc-tagged HuR and PTB overexpression constructs. 16 hrs post-

transfection, cells were infected with SARS-CoV-2 at 0.1 MOI. Cell supernatant was

collected after 48 hrs of infection and Viral titres determined using plaque assay. (F) Viral

RNA levels were quantified in cells from the previous experiment. (G) Western blotting was

performed using anti-myc antibody confirmation of overexpression. (H) WT and HuR KO

HEK-293T-Ace2 cells were transfected with myc-tagged HuR and PTB overexpression

constructs. 16 hrs post-transfection, cells were transfected with pGL3-SV40-Luc (Luc alone)

or pGL3-SV40-SARS-CoV-2 5'UTR-Luc (CoV-2 Luc) and harvested after 24 hrs of

transfection and luciferase readings measured.

Fig 7. Therapeutic significance of the role of HuR in SARS-CoV-2 life cycle. (A) Effect

of HuR KO on remdesivir sensitivity of viral infection. WT and HuR KO HEK-293T-Ace-2

cells were infected with SARS-CoV-2 virus at an MOI of 0.1 and the mentioned

concentrations of Remdesivir added after 1 hr of virus inoculation. Cells were harvested after

48 hrs of treatment and RNA isolated to detected SARS-CoV-2 5'UTR. (B) Table for IC50

of remdesivir for CoV-2 infection in WT and HuR KO cell line. (C) Effect of anti-sense

oligos (ASOs) targeting HuR binding site on viral titres. WT and HuR KO HEK-293T-Ace-2

cells were transfected with the mentioned concentrations of ASO and 12 hrs post transfection,

cells were infected with SARS-CoV-2 virus at 0.1 MOI. 48 hrs post infection, cells were

harvested, and SARS-CoV-2 RNA quantified using real time PCR.

Fig S1. Prediction of HuR binding to SARS-CoV-2 5'UTR. (A) Bioinformatics prediction

for binding of RNA binding proteins to SARS-CoV-2 5'UTR using RBPDB. (B) Secondary

structure of CoV-2 5'UTR. Red boxes denote the binding sites as found in (A).

Fig S2. Predicted HuR binding sites on SARS-CoV-2 5'UTR. (A, B) Binding of HuR to

SARS-CoV-2 5'UTR using Catrapid software.

Fig S3. HuR does not relocalise of cytoplasm upon SARS-CoV-2 infection. (A)HEK-

293T-Ace-2 cells were infected SARS-CoV-2 virus at an MOI of 0.1. Cells were harvested at

indicated timepoints and immunofluorescence staining performed using Alexa Fluor

conjugated secondary antibodies against HuR (Red) and SARS-CoV-2 Spike protein (Green).

The nucleus was counterstained with DAPI. Scale bar represents 10µm. Nuclear-cytoplasmic

fractionation was performed at (B) 24 hrs, (C) 48 hrs post infection. Western blotting was

performed using anti-HuR antibody to detect the presence of HuR. GAPDH served as

cytoplasmic marker and Lamin as nuclear marker.

Figure 1.

Figure 2.

A.

Direct UV-crosslinking:

Probe: SARS-CoV2 5'UTR
Protein: rHuR

B.

Competition UV-crosslinking:

Probe: SARS-CoV2 5'UTR
Protein: rHuR

C.

D.

Semi-quantitative PCR

Western blot

Figure 3.

Figure 4.

Figure 5.

A.

B.

C.

D.

Figure 6.

Figure 7.

A.

B.

	Control	HuR k/o
IC50 at MOI 0.1	10.46 nM	1.82 nM