RESEARCH REPORT

NAC1 directs CEP1-CEP3 peptidase expression and decreases cell wall extensins linked to root hair growth in Arabidopsis

Diana R. Rodríguez-García[1], Eliana Marzol[1], Yossmayer del Carmen Rondón Guerrero[1], Lucía Ferrero[2], Andrés Hugo Rossi[1], Esteban A. Miglietta[1], Ariel. A. Aptekmann[3,4], Javier Martínez Pacheco[1], Mariana Carignani[1], Victoria Berdión Gabarain[1], Leonel E. Lopez[1], Gabriela Diaz Dominguez[1], José Juan Sánchez-Serrano[6], Lin Xu[5], Alejandro D. Nadra[3,4], Federico Ariel[2], Enrique Rojo[6], José M. Estevez[1,7,8,9,†]

[1]Fundación Instituto Leloir and IIBBA-CONICET. Av. Patricias Argentinas 435, Buenos Aires C1405BWE, Argentina.

[2]Instituto de Agrobiotecnología del Litoral, CONICET, Universidad Nacional del Litoral, Colectora Ruta Nacional 168 km 0, 3000, Santa Fe, Argentina.

[3]Departamento de Fisiología, Biología Molecular y Celular, Instituto de Biociencias, Biotecnología y Biología Traslacional (iB3). Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Ciudad Universitaria, Buenos Aires C1428EGA, Argentina.

[4]Departamento de Química Biológica, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires (IQUIBICEN-CONICET), Ciudad Universitaria, Buenos Aires C1428EGA, Argentina.

[5]National Laboratory of Plant Molecular Genetics, CAS Center for Excellence in Molecular Plant Sciences, Institute of Plant Physiology and Ecology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 200032, China

[6]Centro Nacional de Biotecnología, Consejo Superior de Investigaciones Científicas, Cantoblanco, E-28049 Madrid, Spain

[7]Centro de Biotecnología Vegetal, Facultad de Ciencias de la Vida, Universidad Andrés Bello and Millennium Institute for Integrative Biology (iBio), Santiago, Chile

[8]ANID – Millennium Institute for Integrative Biology (iBio), 7500000 Santiago, Chile

[9]ANID –Millennium Nucleus for the Development of Super Adaptable Plants (MN-SAP), 8331150 Santiago, Chile

[†] Correspondence should be addressed. Email: <u>jestevez@leloir.org.ar</u> (J.M.E).

Word count: x,xxx

Running Head: AtCEP1-AtCEP3 negatively modulate root hair growth.

Author for Correspondence:

José M. Estevez

Fundación Instituto Leloir, Av. Patricias Argentinas 435, Buenos Aires C1405BWE, Argentina.
TE: 54-115238-7500 EXT. 3206

Centro de Biotecnología Vegetal, Facultad de Ciencias de la Vida, Universidad Andrés Bello
and Millennium Institute for Integrative Biology (iBio), Santiago CP 8370146, Chile.

Email: jestevez@leloir.org.ar / jose.estevez@unab.cl

Special Issue: Plant cell walls

Abstract Word count: 265

Text Word count: 2,739

Figures 4

Figure S1-S7

Tables S1-S3

References: 48

52
53

Abstract

Plant genomes encode a unique group of papain-type Cysteine EndoPeptidases (CysEPs) containing a KDEL endoplasmic reticulum (ER) retention signal (KDEL-CysEPs or CEPs). CEPs process the cell-wall scaffolding EXTENSIN proteins (EXTs), which regulate *de novo* cell wall formation and cell expansion. Since CEPs are able to cleave EXTs and EXT-related proteins, acting as cell wall-weakening agents, they may play a role in cell elongation. *Arabidopsis thaliana* genome encodes three CEPs (AtCPE1-AtCEP3). Here we report that the three Arabidopsis CEPs, *AtCEP1-AtCEP,* are highly expressed in root-hair cell files. Single mutants have no evident root-hair phenotype, but *atcep1-3 atcep3-2* and *atcep1-3 atcep2-2* double mutants have longer root hairs (RHs) than wild type (Wt) plants, suggesting that expression of *AtCEPs* in root trichoblasts restrains polar elongation of the RH. We provide evidence that the transcription factor *NAC1* activates AtCEPs expression in roots to limit RH length. Chromatin immunoprecipitation indicates that NAC1 binds the promoter of *AtCEP2* and may directly regulate its expression. Indeed, Inducible *NAC1* overexpression increases *AtCEP1 and AtCEP2* transcript levels in roots and leads to reduced RH growth while the loss of function *nac1-2* mutation reduces *AtCEP1-AtCEP3* gene expression and enhances RH growth. Likewise, expression of a dominant chimeric NAC1-SRDX repressor construct leads to increased RH length. Finally, we show that RH cell walls in the *atcep1-1 atcep3-2* double mutant have reduced levels of EXT secretion/insolubilization, suggesting that the defects in RH elongation are linked to alterations in EXT processing and accumulation. Taken together, our results support the involvement of the NAC1-AtCEP regulatory module in controlling RH polar growth through EXT processing and insolubilization at the cell wall.

.

Introduction

In plants there is a unique group of papain-type Cysteine EndoPeptidases (CysEPs) containing a KDEL endoplasmic reticulum (ER) retention signal (KDEL-CysEPs or CEPs) for which no homologous genes have been identified in mammals or yeast (Gietl et al. 2000). CEPs are synthesized as pre-pro-enzyme and co-translationally translocated into the ER, where the pre-sequence is removed and the pro-enzyme is finally released from the ricinosomes (ER-associated structures) upon vacuolar collapse and acidification of the cytosol, triggering the maturation of the enzyme (Schmid et al. 1999, 2001; Beers et al. 2000; Zhang et al. 2014). AtCEP1 (At5g50260), AtCEP2 (At3g48340), and AtCEP3 (At3g48350) are three *Arabidopsis thaliana* AtCEPs with broad expression patterns in vegetative and reproductive tissues along the plant, including root tissues (Helm et al. 2008; Hierl et al. 2014; Zhou et al. 2016). Several CEPs have been identified in cells or tissues associated with programmed cell death (PCD), where they play crucial roles in intracellular protein degradation (Tanaka 1991; Becker et al. 1997; He and Kermode 2003; Zhang et al. 2014; Olvera-Carrillo Y et al 2015; Chen et al. 2016). AtCEP1 was found to be a central mediator of tapetal PCD, allowing tapetal cell degeneration and functional pollen formation (Zhang et al. 2014). In addition, AtCEP1 and AtCEP2 in Arabidopsis were found to be functional in the *de novo* emergence of adventitious root tips associated with EXT-degradation and regulated by the NAC1 transcription factor (Chen et al. 2016). More recently, it was reported that *AtCEP1* and *AtCEP2* are expressed in root epidermal cells that separate to allow LR emergence, and that loss of function of *AtCEP1* or *AtCEP2* causes delayed emergence of LR primordia, suggesting that these KDEL-CysEPs might be involved in cell-wall remodelling for cell separation during development (Howing et al. 2018).

In addition to the papain-type preference for neutral amino acids with large aliphatic and non-polar (Leu, Val, Met) or aromatic (Phe, Tyr, Trp) side chains in the P2 position, Ricinus CEP (RcCysEP) exhibits an unusually broad substrate specificity. This broad substrate specificity is a result of the active site cleft of KDEL-CysEPs, which accepts a wide variety of amino acids, including proline and the glycosylated hydroxyproline of hydroxyproline-rich glycoproteins (HRGP) of the cell wall (Than et al. 2004). The amino acid residues that are essential for this generally more open structure of the active site cleft, as well as those that define the catalytic pocket, are highly conserved among known KDEL-CysEPs (Hierl et al. 2014). It can recognize the Ser-(Hyp)$_{3-5}$ repeats, *O*-glycosylated Hyp, and prolines at one-two amino acids relative to the cleavage site (Than et al., 2004; Helm et al. 2008; Hierl et al. 2014). These Ser-(Hyp)$_{3-5}$ repeats with *O*-glycosylated modifications are frequently observed in structural *O*-glycoproteins Extensins (EXTs) and possibly in a large number of uncharacterized apoplastic EXT-related proteins (e.g. PERK, Formins, AGP-EXTs hybrids) (Borassi et al. 2016). 59 encoded EXTs in *Arabidopsis thaliana* contain a Tyr-crosslinking motif close to an *O*-glycosylated Ser-(Hyp)$_{3-5}$ (Showalter et al., 2010; Marzol et al. 2018). EXT Tyr-mediated crosslinking is catalyzed by apoplastic peroxidases (Schnabelrauch et al., 1996; Jackson et al., 2001) and allows them to form glycoprotein networks in the cell wall, which influences de novo plant cell wall

formation during embryo development (Cannon et al., 2008) and in polar cell expansion
processes (Velasquez et al. 2011; 2015). Since CEPs can cleave *O*-glycosylated EXTs (Hierl et
al., 2014), thus acting as cell wall-weakening agents, this lends credence to the idea that CEPs
may play a pivotal role in cell elongation. Prior to this study, we determined that at least six
EXTs (EXT6–7, 12–14, 18) co-regulated at the transcriptional level play a crucial role in the
polar-cell expansion process, specifically in RHs (Velasquez et al. 2011; Velasquez et al 2015;
Marzol et al. 2018) In this process, Leucine Rich Extensins 1 and 2 (LRX1 and LRX2) were also
found to be essential (Baumberger 2001, 2003; Ringli 2010). Pollen EXTs (PEXs) and LRXs were
also recently linked to polar growth regulation in pollen tubes (Sede et al. 2018; Ndinyanka
Fabrice et al., 2017; Wang et al., 2018), highlighting a conserved role of these EXT and EXT-
related proteins in polar growth.

On the basis of these previous findings, we hypothesized that Arabidopsis AtCEPs may play a
role in polar-growth regulation linked to the processing of *O*-glycosylated EXT and EXT-related
proteins processing and possibly other substrates during their maturation along the secretory
pathway. Here, we provide evidence that, indeed, AtCEPs negatively regulate levels of EXT
secretion/insolubilization at the cell-wall of root hairs (RHs) and retrain their growth.

Results and Discussion

Expression of Arabidopsis AtCEPs has been previously analysed using functional reporter
constructs that include the coding sequences of AtCEP1 and AtCEP2 under the control of their
respective regulatory regions of 2.0 Kb fragment upstream of the start codon, with HA and
fluorescent protein tags, either GFP or mCherry (AtCEP1pro:PRE-PRO-3xHA-EGFP-AtCEP1-
KDEL and AtCEP2pro:PRE-PRO-3xHA-mCherry-AtCEP2-KDEL) inserted between the pro-
peptide and the mature enzyme sequence (Howing et al. 2014 and Hierl et al. 2014).
Intriguingly, it was reported that AtCEP2pro:PRE-PRO-3xHA-mCherry-AtCEP2-KDEL is
expressed specifically in non-protruding cell files of the hypocotyl, the cell files where stomata
are formed, but not in protruding cell files (Hierl et al., 2014). Interestingly, the alternating
pattern of non-protruding and protruding cell files in the hypocotyl is controlled by common
mechanisms to the alternating pattern of root-hair cell (root trichoblast) and non-root-hair
cell (root atrichoblast) files in the root. Hypocotyl non-protruding cells and root trichoblasts
are always positioned over an anticlinal cell wall of the underlying cortex and lack expression
of cell fate regulators of GL2 and WER and the enhancer-trap marker J2301, whereas
hypocotyl protruding cells and root atrichoblast are positioned over a periclinal cortex cell
wall and express the J2301 marker and GL2 and WER, which block development of stomata or
RHs in these cell files (Berger et al.,1998, Grierson et al., 2014). Importantly, we found that
AtCEP2pro:PRE-PRO-3xHA-mCherry-AtCEP2-KDEL is expressed in root epidermis exclusively in
the trichoblasts (**Figure 1A-B; Figure S1-S2**), further supporting that root-hair cell files share
specification mechanisms and characteristics with non-protruding cell files of the hypocotyl

(Berger et al.,1998, Grierson et al., 2014). AtCEP1pro:PRE-PRO-3xHA-EGFP-AtCEP1-KDEL was
also specifically expressed in root trichoblasts (**Figure 1A-B; Figure S1**). In these cells, the
mCherry-AtCEP2 fusion protein was found in mobile punctate structures, consistent with it
being localized in the endomembrane system. To determine the nature of those organelles,
we analysed for co-localization with established markers for different compartments (**Figure
2A**). The AtCEP2 labelled puncta did not co-localize with markers for the Golgi apparatus
NAG1-GFP (0.06+-0.13) and N-ST-YFP (0.29+-0.16) (**Figure 2A**). However, we found higher co-
localization levels in round or spindled shaped compartments with the ER lumenal marker SP-
GFP-KDEL (0.63+-0.16), indicating that AtCEP2 might reside in an ER-derived compartment in
root trichoblasts (**Figure 2B**). Indeed, AtCEP2 was observed in ER-derived compartments that
resembled ricinosomes and ER-Bodies in leaf, hypocotyl, and root cap cells (Hierl et al. 2014;
Howing et al. 2014). In addition, CEP2 mCherry was not detected in the apoplast space when
plasmolysis was performed in the root hairs (**Figure 2C**). On the other hand, apoplast
targeting of AtCEPs has not been reported previously, possibly indicating that AtCEPs process
their substrates within the secretory pathway. This confirms that AtCEP2 is primarily targeted
to the ER compartment in root trichoblasts. We cannot rule out the possibility of a low level
of an apoplast AtCEP2 expression. To analyse the expression of *AtCEP3*, we used a promoter
fragment of 2-0Kb to drive GFP expression. *AtCEP3pro::GFP* fluorescence was also high in root
trichoblasts, although not specific to this root epidermal cell-type (**Figure 1A-B; Figure S1-S2**).
However, the specific expression of AtCEP1 and AtCEP2 in root trichoblast cells and the high
expression of AtCEP3 indicates that these genes may play a role in growth of these specialized
cells. To address if they are involved in RH polar growth we isolated *cep* T-DNA mutants
(**Table S2** and **Figure 1C**). We characterized at least two T-DNA alleles for each *AtCEP* gene.
Single mutants for *AtCEP1*, *AtCEP2* and *AtCEP3* showed similar phenotype to Wt Col-0 (**Figure
S3**) while the double mutants *atcep1-3 atcep3-2* and *atcep1-3 atcep2-2* showed increased RH
growth (up to 20% longer) when compared to Wt Col-0 (**Figure 1D**). These suggested that all
three AtCEPs, AtCEP1-AtCEP3, redundantly restrict RH growth. We reasoned then that
increased levels of these proteins should shorten RH length. Together, these results indicate
that Arabidopsis AtCEP1-AtCEP3 proteins are expressed in RHs where they negatively regulate
RH cell growth.

NAC1 was previously identified as a regulator of lateral root development and *de novo* root
organogenesis (Xie et al., 2000; Chen et al., 2016). Since NAC1 was shown to control *AtCEP1*
and *AtCEP2* expression during *de novo* root organogenesis (Chen et al. 2016), we tested if a
similar regulation could be taking place in RHs. Indeed, *in silico* analysis (eFP browser and
Root Cell Atlas) and characterization of a *NAC1pro:GFP* reporter line supports that *NAC1* is
highly expressed in root trichoblasts, specifically in the later stages of their development
(**Figure 3A-B**; **Figure S4**). Moreover, a *nac1-2* null mutant displays significantly longer RHs than
Wt plants (**Figure 3B**). In addition, when we expressed *NAC1* fused to the repression domain
SRDX (Hiratsu et al., 2003) to specifically suppress the expression of NAC1 target gene
(35Spro::NAC1-SRDX1-1 and 35Spro::NAC1-SRDX1-2 lines) we observed an elongated RH

phenotype (**Figure 3B**). Then, to analyse if *NAC1* overexpression would trigger a direct effects
on RH growth possibly by upregulation of AtCEP1/AtCEP2, we used the pER8:3pro::×FLAG-
NAC1 β-estradiol-inducible line (Zuo et al., 2000). In the presence of estradiol, an 75-fold
induction of *NAC1* levels was observed (**Figure S5**). We then tested if the transcription factor
NAC1 activates *AtCEP1-AtCEP3* expression in developing roots, as was shown for *de novo*
adventitious root formation assay (Chen et al. 2016). The estradiol-inducible NAC1-FLAG line
increased *CEP1* expression by 1.6 folds and *CEP2* by 4-folds while no changes were detected
for *CEP3*. (**Figure S5**). The *nac1-2* mutant showed significantly lower levels of transcripts of all
three AtCEPs, AtCEP1-AtCEP3 (**Figure 3C**). Together, these results indicate that NAC1 activates
the expression of *AtCEP1-AtCEP3*, and in consequence, controls RH growth, possibly through
their EXT-processing activity. To determine if NAC1 controls the expression of *AtCEP* genes by
direct binding to their promoter regions, we searched for putative NAC-binding sites in open
chromatin regions, according to publicly available ATAC-Seq datasets (Maher et al., 2018).
According to ChIP-qPCR using pER8:3pro::×FLAG-NAC1 plants (60x induction) and anti-FLAG
antibodies, *AtCEP2* and *AtCEP3* appeared as direct targets (**Figure 3D**), as revealed in
comparison to the previously identified direct target *E2Fa* (Xie et al. 2023).

In previous studies, AtCEPs were shown to be involved in the processing of EXT proteins
(Greenwood et al., 2005; Helm et al., 2008; Hierl et al., 2012). We thus tested if the reduction
of AtCEP activity in the mutants had an effect on the EXTs secreted and insolubilized in the RH
cell walls. To this end, we used an EXT-reporter carrying a tdTomato tag (SS-TOM-Long-EXT)
that is resistant to the acidic pH, characteristic of cell wall apoplast (**Figure 4A**). Importantly,
expression of the reporter did not affected the polar growth of RHs (Martinez Pacheco et al.
2022), making it an ideal probe for monitoring *in situ* alterations in the arrangement of cell
wall EXTs. We measured the cell wall fluorescence signal from the SS-TOM-Long-EXT
construct in the apical zones of RHs under plasmolysis. Plasmolysis allowed us to retract the
plasma membrane and detect the EXT-signal coming specifically from the cell walls.
Interestingly, cell wall stabilization/insolubility of SS-TOM-Long-EXT in the RH tip was
drastically reduced in *cep1 cep3* double mutant when compared to Wt Col-0 plants. Thus, the
SS-TOM-Long-EXT reporter tested in the apical zone of the RHs is modified by deficient AtCEPs
activities possibly during the EXT processing in the secretory pathway. Since EXTs
insolubilization in the cell walls of growing RHs is regulated by several factors including the
Reactive Oxygen Species (ROS) homeostasis (Martinez Pacheco et al. 2022; Marzol et al.
2022), we tested if the effect of these AtCEPs might affect global ROS levels. We measured
total cytoplasmic ROS ($_{cyt}H_2O_2$) with the hydrogen peroxide-selective dye Peroxy-Orange 1
(PO1), as a permeable boronate-based dye that is non-fluorescent in its reduced form, but
becomes fluorescent when irreversibly oxidized by H_2O_2 (Dickinson et al., 2010). Apoplastic
ROS ($_{apo}H_2O_2$) levels were determined in the RH tips with the cell-impermeable Amplex™
UltraRed Reagent (Mangano et al. 2016; Matinez-Pacheco et al. 2022; Marzol et al. 2022)
(**Figure 4B**). The *atcep1-3 atcep3-2* showed higher levels of $_{apo}H_2O_2$ in RH tips compared to
Col-0, whereas the $_{cyt}H_2O_2$ were similar in both genotypes. This indicates that ROS

homeostasis is also changed in the mutant background, which can also affect the EXTs
insolubilization in the cell walls.

Finally, to test if AtCEP1-AtCEP3 might be able to interact with single-chain EXTs, we
performed homology modeling with one cysteine endopeptidase of RcCysEP and a cysteine
protease from *Ambrosia artemisiifolia* (pdbs 1S4V and 5EF4 respectively) known to be able to
cleave and process EXT-like substrates under *in vitro* conditions. AtCEP1-AtCEP3 proteins
share a sequence identity of 66-74% with the most well-characterized *Ricinus communis*
RcCysEP (**Figure S6**). By *docking* analysis, we obtained interaction energies (Kcal/mol) for all
three AtCEPs and they were compared to RcCysEP 1SV4. We analyzed docking with four
different short EXT peptides from non-hydroxylated to fully *O*-glycosylated peptide (Marzol et
al. 2016) (**Figure S7**). It was previously shown that mutants carrying under-*O*-glycosylated
EXTs and related EXT-proteins have severe defects in RH growth (Velasquez et al. 2011;
Velasquez et al. 2015). Our docking results for these three AtCEPs showed consistent
interaction energy differences that depend on the EXT glycosylation state, being higher for
non-*O*-glycosylated species. In general, we observed higher interaction energies (higher
negative values) for (non)-hydroxylated EXT species than for *O*-glycosylated EXT variants.
When we compared interaction energies among different AtCEPs interacting with EXT
substrates, we observed that AtCEP1 and AtCEP3 displayed the highest interaction activity
with the (non)-hydroxylated EXT species (**Figure S7**). Overall, this likely indicates that AtCEP1
and AtCEP3 might interact with EXT substrates and possibly catalyse proteolysis in open
regions of the EXT backbones with little or no *O*-glycosylation while CEP2 might prefer *O*-
Glycosylated regions. This is in agreement with previous studies suggesting that high levels of
O-glycosylation in certain proteins physically restrict its degradation.

Conclusions

It was previously considered that up-regulation of AtCEPs was related to the processing of EXT
since EXT promotes wound healing, and this might be a barrier for the emergence of
regenerated root tips. Therefore, it was hypothesized that the NAC1-AtCEPs pathway
antagonizes EXT-mediated wound healing, and this allows the emergence of regenerated root
tips (Chen et al. 2016). However the overall results in our study suggest that NAC1 directly
activates AtCEP1 and AtCEP2 expression, which in turn represses polar-cell expansion in
growing RHs. By a colocalization analysis with several subcellular markers, we detected
AtCEP2 in ER-derived compartments but not in the apoplast. Based on our results, it is
plausible that AtCEP1-AtCEP3 could act together as components on the EXT and EXT-related
protein quality control program and proper EXTs protein processing. So far there is no
information regarding the existence of a quality control pathway for *O*-glycosylated proteins
in the same manner to the well-known and conserved CNX (calnexin)/calreticulin lectins
linked to ERAD (ER-associated degradation)-program (e.g. with OS9, HRD3/SEL1L components)

as it happens for misfolded *N*-glycosylated proteins (Su et al. 2011; Su et al. 2012; Strasser
2016). It is also unknown how the *O*-glycosylation machinery required for a proper processing
of EXTs and EXT-related proteins in the secretory pathway (Velásquez et al. 2012; Marzol et al.
2018) are coordinated and controlled. Our findings pave the way for the discovery of novel
functions for AtCEPs in cells that are not undergoing PCD but are characterized by extensive
cell wall expansion or remodelling, possibly as a result of EXTs cleavage. The involvement of a
protease in cell wall remodelling indicates the significance of the protein component of the
cell wall in this process. Insights into the underlying mechanisms, such as the membrane
fusion of ricinosomes with the plasma membrane during root development, similar to other
ER-derived vesicles, will be intriguing hypothesis to investigate. Recently, a vacuolar
processing enzyme (βVPE) was found to control the maturation of AtCEP1 transforming this
pro-protease into mature protease while in *vpe* mutants, the maturation of AtCEP1 and other
proteases was severely inhibited (Cheng et al. 2020). It is still unclear if VPE may be also acting
in root trichoblasts to control AtCEPs processing. Root trichoblasts provide an excellent model
system to dissect the molecular components of the NAC1-AtCEPs-EXT *O*-glycosylation
pathway required for polar-growth.

Experimental Procedures

Plant and Growth Conditions. *Arabidopsis thaliana* Columbia-0 (Col-0) was used as the Wt
genotype in all experiments. All mutants and transgenic lines tested are in this ecotype.
Seedlings were germinated on agar plates in a Percival incubator at 22°C in a growth room
with 16h light/8h dark cycles for 7-10 days at 140 μmol m^{-2}s^{-1} light intensity. Plants were
transferred to soil for growth under the same conditions as previously described at 22°C. For
identification of T-DNA knockout lines, genomic DNA was extracted from rosette leaves.
Confirmation by PCR of a single and multiple T-DNA insertions in the target AtCEP and NAC
genes were performed using an insertion-specific LBb1 or LBb1.3 (for SALK lines) or Lb3 (for
SAIL lines) primer in addition to one gene-specific primer. To ensure gene disruptions, PCR
was also run using two gene-specific primers, expecting bands corresponding to fragments
larger than in Wt. In this way, we isolated homozygous lines (for all the genes mentioned
above). Mutant list is detailed in **Table S1**. Lines expressing AtCEP2pro:pre-pro-3xHA-
mCherry-AtCEP2-KDEL were crossed with ER and Golgi marker lines to obtain the double
transgenic AtCEP2 + organelle marker

Root Hair Phenotype. Seeds were surface sterilized and stratified in darkness for 3 d at 4°C
and they were grown *in vitro* on a specific condition and medium in a plant growth chamber
in continuous light (120 mol s-1 m-2) at 22°C. The quantitative analyses of RH phenotypes of
Col-0 and transgenic lines were made the last day of the growth conditions described in the
two previous sections. Images were captured using an Olympus SZX7 Zoom Stereo
Microscope (Olympus, Tokyo, Japan) equipped with a Q-Colors digital camera and Q CAPTURE

PRO 7 software (Olympus). Results were expressed as the mean SD using the GRAPHPAD
PRISM 8.0.1 (GraphPad Software, Boston, MA, USA) statistical analysis software. Results are
representative of three independent experiments, each involving 15–20 roots.

AtCEP imaging. Root hairs were ratio imaged with the Zeiss LSM 510 laser scanning confocal
microscope (Carl Zeiss) using a 40X oil-immersion, 1.2 numerical aperture objective. EGFP
(473–505nm) and mCherry (526–536 nm) emissions were collected using a 458-nm primary
dichroic mirror and the Meta-detector of the microscope. Bright-field images were acquired
simultaneously using the transmission detector of the microscope. For time-lapse analysis,
images were collected every 3 or 5 s. Image sequences were analyzed using the Template
Matching and Slice Alignment plug-in for ImageJ. Fluorescence intensity was measured in 7
μm ROI at the RH apex.

Transcriptional reporter generation. Vectors based on the Gateway cloning technology
(Invitrogen) were used for all manipulations. Promoter regions (2Kb) were PCR-amplified with
AttB recombination sites. PCR products were first recombined in pDONOR207 and transferred
into pMDC111. Transgenic lines used in this study are described in **Table S2**.

AtCEPs expression. To quantify AtCEP1 and AtCEP2 expression levels, lines AtCEP1pro:3xHA-
EGFP-KDEL AtCEP2pro: 3xHA-mCherry-KDEL were used. The fluorescence intensity of GFP and
mCherry was measured respectively. All RHs growth stages were measured. Ten days old
seedlings were removed from the medium and placed in a slide containing a drop of liquid MS
0.5X in the dark, and images were obtained using a Zeiss Imager A2 epifluorescence
microscope. The objective used was 10X, 0.3 numerical aperture, exposure time 2 seconds.
The lasers used were suitable for each fluorophore: GFP (GFP λ max ex. = 488 nm, λ max em. =
507 nm) and mCherry (λ max ex. = 536 nm, λ max em. = 632 nm) The Wt Col-0 line was used
to rule out autofluorescence noise. The images were analyzed using ImageJ 1.50b software.
To measure fluorescence intensity levels (represented in arbitrary units: UA), both a circular
region of interest within the RH apex (ROI) and the total area of RH were selected. All RHs
from 6 seedlings were analyzed. The reported values are the mean ± standard error (mean ±
SEM).

Subcellular localization of AtCEP2. Lines expressing CEP2pro:pre-pro-3xHA-mCherry-CEP2-
KDEL were crossed with ER and Golgi marker lines to obtain the double transgenic CEP2 +
organelle marker. Co-localization analysis was performed using the BIOP implementation
(Battistella et al. 2019) of the JACoP-plugin (Bolte and Cordelières 2006) in ImageJ (1.53t) on
individual root hairs for which ROIs (Region Of Interest) were drawn manually. Background
intensity was subtracted from each channel as the mean intensity of the autofluorescence
control + 2SD. Different marker expression levels excluded the usage of fixed cutoff values for
all images, so the histogram-derived Otsu method was chosen based on visual inspection of
the thresholded images for both channels. To compare the relative localization of
CEP2pro:pre-pro-3xHA-mCherry-CEP2-KDEL (Ch2) in each of the different subcellular

compartments (Ch1), we used Mander's M2 overlap coefficient (Manders et al. 1993) to
measure the proportion of positive pixels in Ch2 that co-occur with positive pixels in Ch1.
Confocal microscopy images were acquired on a Zeiss LSM710 microscope with an EC Plan-
Neofluar 40x/1.30 oil objective. Images were acquired sequentially on different tracks in
order to avoid excitation and emission bleed-through using the following emission ranges for
the individual channels (8-bit pixel depth): Ch1 519–589 nm, Ch2 594-690 nm. Pixel size was
set at 100 nm following Nyquist sampling criterion and pinhole was adjusted to obtain a 3.6
µm thick optical slice. For each experimental replicate, 15-25 total individual root hairs from
five different plants per subcellular compartment marker were imaged.

Chromatin immunoprecipitation Assay. Chromatin immunoprecipitation (ChIP) assays were
performed on pER8:3xFLAG-NAC1 plants (Chen et.al 2016, DOI: 10.1104/pp.15.01733) mainly
as described in Ariel et al. (2020). Plants were grown for 10 days in plates containing MS 0,5X
medium (pH 5,7; 0.8% agar) placed vertically in a culture chamber at 22°C and continuous
light (140µmol/m2.sec). After 10 days, the plates were placed horizontally and treated with β-
estradiol 10 µM solution for 3h. The expression of NAC1 was checked by qPCR
(supplementary table Sx). Chromatin was cross-linked with formaldehyde 1% for 10 min at
room temperature. Cross-linking was stopped by adding glycine (125 mM final concentration)
and incubating for 10 min at room temperature. Crosslinked chromatin was extracted by cell
resuspension, centrifugation, cell membrane lysis, and sucrose gradient as previously
described (Ariel et al., 2020). Nuclei were resuspended in Nuclei Lysis Buffer and chromatin
was sonicated using a water bath Bioruptor Pico (Diagenode; 30s on / 30s off pulses, at high
intensity for 10 cycles). Chromatin samples were incubated for 12h at 4°C with Protein G
Dynabeads (Invitrogen) precoated with the antibodies anti-FLAG (Abcam ab18230) or anti-IgG
(Abcam ab6702) as a negative control. Immunoprecipitated DNA was recovered using
Phenol:Chloroform:Isoamilic Acid (25:24:1; Sigma) and analyzed by qPCR using the primers
listed in Supplementary Table Sx. A region upstream E2Fa gene was used as a positive control
(Xie et al. 2023). ACTIN was used as a negative control. Untreated sonicated chromatin was
processed in parallel and considered the input sample. The GraphPad Prism 6 software was
used to analyze the data and produce the graphs.

Apoplastic and cytoplasmic ROS measurements. To measure ROS levels in root trichoblasts, 8
days-old Arabidopsis seedlings grown in continuous light were used. For cytoplasmic ROS, the
seedlings were incubated in darkness for 10 min with H_2O_2 and were visualized with Peroxy-
Orange 1 (PO1). PO1 was dissolved in DMSO to make a 500 µM stock and was further diluted
in water to make a 50 µM working solution. Seedlings were incubated in PO1 for 15 min in the
dark and were then rinsed with water and mounted in water for imaging and observed with
Zeiss Imager A2 Epifluorescence Microscope (Zeiss, Germany) (Plain Apochromat 40X/1.2 WI
objective, exposure time 25 ms). Images were analyzed using ImageJ software. To measure
ROS levels, a circular region of interest was chosen in the zone of the RH tip cytoplasm. To
measure apoplastic ROS, the seedlings were incubated with 50 µM Amplex™ UltraRed
Reagent (AUR) (Molecular Probes, Invitrogen) for 15 min in darkness and rinsed with liquid

0.5X MS media (Duchefa, Netherlands). Root hairs were imaged with a Zeiss LSM5 Pascal
(Zeiss, Germany)) laser scanning confocal microscope (Excitation 543 nm argon laser;
Emission: 560–610 nm, Plain Apochromat 40X/1.2 WI objective). Quantification of the AUR
probing fluorescence signal was restricted to apoplastic spaces at the RH tip and quantified
using the ImageJ software. Fluorescence AU were expressed as the mean ± SD using the
GraphPad Prism 8.0.1 (USA) statistical analysis software. Results of both ROS measurements
are representative of two independent experiments, each involving 10–15 roots and
approximately, between 10 and 20 RHs per root were observed.

Treatments with *β*-Estradiol. *β*- Estradiol (Sigma-Aldrich,) was added to 0.5X MS media at a
final concentration of 10 µM. The lines ER8pro:AtCEP2 and ER8pro:3×FLAG-NAC1 were grown
at 22°C for 4 days + 3 days of *β*- Estradiol induction. Root hairs phenotype was then
quantified.

Quantitative reverse transcriptase PCR (qRT-PCR). Total RNA was isolated from 10-d-old
seedling roots (40 for each line) using the RNeasy Plant Mini Kit (Qiagen, Germany). cDNA was
synthesized using TOPscript^TM RT DryMIX (dT18, Enzynomics, Korea). qRT-PCR analyses were
performed using TOPrealTM qPCR 2x PreMIX (SYBR Green, Enzynomics, Korea) and
Chromo4™ Four-Color Real-Time Detector (Bio-Rad, USA). Gene-specific signals were
normalized relatively to *PP2A* (At1G69960) signals. Each qRT-PCR reaction was performed in
triplicate, and each experiment was repeated three times using independent preparations of
RNA. Primers used are as listed in **Table S3**.

Modelling and molecular docking between AtCEP1-AtCEP3 and EXTs. Modeling and
molecular docking: cDNA sequences of AtCEPs were retrieved from TAIR (AtCEP1:
AT1G05240, AtCEP2: AT3G50990, AtCEP3: AT4G26010) and NCBI Nucleotide DB. Homology
modeling was performed using an AtCEP from *Ricinus communis* and from *Ambrosia*
artemisiifolia using modeller 9.14 (Sali et al. 1993), using the crystal structures 1S4V and 5EF4
as templates, available at the protein data bank. 100 structures were generated for each
protein and the best scoring one (according to DOPE score) was picked. The *receptor* for the
docking runs was generated by the prepare_receptor4 script from the autodock suite, adding
hydrogens   and   constructing   bonds.   Peptides   based   on   the   sequence
PYYSPSPKVYYPPPSSYVYPPPPS were used, replacing proline by hydroxyproline, and/or adding
O-Hyp glycosylation with up to four arabinoses per hydroxyproline in the fully glycosylated
peptide and a galactose on the serine, as it is usual in plant *O*-Hyp (Strasser 2016). Ligand
starting structure was generated as the most stable structure by molecular dynamics
(Velasquez et al. 2015a). All ligand bonds were set to be able to rotate. Docking was
performed in two steps, using Autodock vina (Trott et al. 2010). First, an exploratory search
over the whole protein surface (exhaustiveness 4) was done, followed by a more exhaustive
one (exhaustiveness 8), reducing the search space to a 75x75x75 box centered over the most
frequent binding site found in the former run.

429
430

Acknowledgements

We thank ABRC (Ohio State University) for providing T-DNA lines seed lines and Gietl for providing some of the materials used in this study. J.M.E. is investigator of the National Research Council (CONICET) from Argentina. This work was supported by grants from ANPCyT (PICT 2016-0132 and PICT2019-00015 to J.M.E and PICT2018-00577 to E.M.) and from the Spanish Ministry of Science and Innovation MCIN/AEI/10.13039/501100011033 and FEDER "una manera de hacer Europa" (PID2021-128078NB-I00 to E.R.). In addition, this research is also funded by ANID – Programa Iniciativa Científica Milenio ICN17_022, NCN2021_010 and Fondo Nacional de Desarrollo Científico y Tecnológico [1200010] to J.M.E. The authors gratefully acknowledge the Microscopy and Imaging Facility at the Leloir Institute Foundation (FIL) for their support and assistance in the present work.

Author Contribution

D.R.G performed all the experiments and analyzed the data. E.M performed AtCEP expression and analyzed the data. Y.d.C.R. performed ROS experiments. A.H.R and E.A.M. carried out co-localization studies. A.A.A. and A.D.N. carried out the molecular modeling of AtCEPs-EXTs. L.F and F.A. carried out the ChIP experiment and analyzed the data. J.M. P., M.C., V.B.G., L.E.L and G.D.D. analyzed the data. L.X., E.R and J.J.S.S. provided materials, analyzed the co-localization data and wrote the paper. E.R. wrote the paper. J.M.E. designed research, analyzed the data, supervised the project, and wrote the paper. All authors commented on the results and the manuscript. This manuscript has not been published and is not under consideration for publication elsewhere. All the authors have read the manuscript and have approved this submission.

Competing financial interest

The authors declare no competing financial interests. Correspondence and requests for materials should be addressed to J.M.E. (Email: jestevez@leloir.org.ar).

460
461
Figure 1. AtCEP1, AtCEP2 and AtCEP3 are expressed in root hairs and negatively regulate
root hair growth. (**A**) AtCEP1 and AtCEP2 are expressed in root trichoblasts. Translational
reporters for AtCEP1 (AtCEP1pro::PRE-PRO-GFP-KDEL) and AtCEP2 (AtCEP2pro::PRE-PRO-
mCherry-AtCEP2-KDEL). AtCEP3 are expressed in both atricoblasts and tricoblasts.
Transcriptional reporters of AtCEP3 (AtCEP3proGFP) in the root differentiation zone and
specifically in RHs. Scale bar = 200 µm. (*) Indicates lack of expression in atrichoblasts. (**B**)
Scheme of AtCEP1, AtCEP2 and AtCEP3 genes showing introns (thin lines), exons (rectangles)
and positions of T-DNA insertions. (**C**) Quantitative analysis of RH length (mean ± s.e.m., n=
200) in Wt Col-0 and *ceps* mutants. NS= not significant difference. Data are shown as the
mean ± SEM, (n=20). Asterisks indicate significant differences from the Wt according to an
ANOVA test with p<0.05. (On the right) Selected images of RHs in Wt and in single and double
cep mutants. Scale bar = 200 µm.
474

475
476
Figure 2. AtCEP2 colocalizes mostly with an ER-membrane marker and, to a lower extent, to Golgi markers in root hair cells. (**A**) Co-localization of AtCEP2-mCherry with markers for different subcellular compartments. Seven-day-old AtCEP2mCherry seedlings were grown (n= 10 roots with 1–5 root hair cells each). HDEL-GFP is a Lumen Endoplasmic reticulum (ER) marker and KKXX-GFP is an ER membrane marker. NAG-GFP is a cis-Golgi marker and ST-YFP is a trans-Golgi network marker. Scale bar 5 µm. (**B**) Quantification of colocalization using Manders correlation coefficient. (**C**) Plasmolyzed root hair to show the lack of signal in the apoplastic regions of AtCEP2-mCherry. Scale bar 20 µm. (CW) Cell Wall; (PM) Plasma membrane; (Cyt) Cytoplasm.
486
487

489
490

491
Figure 3. NAC1 is expressed and regulates root hair growth linked to AtCEP1-AtCEP3
493 **expression.** (**A**) NAC1pro:GFPexpression in roots and RHs. (**B**) RH phenotype in *nac1-2* mutant
494 and two constitutive negative NAC1 lines. Quantitative analysis of RH length (mean ± s.e.m.,

n= 200). Data are shown as the mean ± SEM, (n=20). Asterisks indicate significant differences from the Wt Col-0 according to an ANOVA test with p<0.05. (**C**) qPCR analysis of AtCEP1, AtCEP2, and AtCEP3 in Wt Col-0 and *nac1-2* mutants. RNA was extracted from root of seedlings. *PP2A* was used as a control and amplification was performed for 30 cycles. Black arrowheads represent the region amplified by the primers used for the RT-PCR. (**D**) ChIP-qPCR analysis of NAC1 binding to CEP2 and CEP3 promoter regions. Schemes of the loci showing the location of the fragments analyzed by ChIP-qPCR are shown in the upper part. Primers were designed analyzing ATAC-seq experiments in regions where the chromatin is accessible (Maher et al., 2018). The enrichment was measured relative to the negative control ACTIN.

Figure 4. **EXT secretion/insolubilization and apoplastic ROS are affected by AtCEP1-AtCEP3 in root hairs**. (**A**) Signal of SS-EXT LONG-TOM in the apical zone of RHs in the *cep1-3 cep3-2* double mutant. Cells were plasmolyzed with a mannitol 8% solution. In the images: (*) indicates cell surface including the plant cell walls, (**) indicates the retraction of the plasma membrane, (ap) apoplastic space delimited between the plant cell wall and the retracted plasma membrane. Each point in the graph is the signal derived from a single RH tip. Fluorescence AU data are the mean ± SD SD (N = 3), two-way ANOVA followed by a Tukey–Kramer test; (***) p<0.01. Results are representative of two independent experiments. NS = non-significant differences. Scale bars = 5 μm. (**B**) Intensity profiles across a dotted line in the RH tips. (**C**) Cytoplasmic ROS ($_{cyt}$ROS) levels were measured using the hydrogen peroxide-selective dye Peroxy-Orange 1 (PO1) in apical areas of RHs in wild-type (Columbia Col-0) and in the double mutant *cep1-3 cep3-2*. Each point is the signal derived from a single RH tip. Fluorescence AU data are the mean ± SD (N= 20 RHs), two-way ANOVA followed by a Tukey–

Kramer test; (**) p < 0.01. Results are representative of two independent experiments.
Asterisks indicate significant differences. NS = non-significant differences. Scale bars = 2 µm.
(**D**) Apoplastic ROS ($_{apo}$ROS) levels were measured with Amplex™ UltraRed in apical areas of
RHs in wild-type (Columbia Col-0) and in the double mutant *cep1-3 cep3-2*. Each point in (**D**) is
the signal derived from a single RH tip. Fluorescence AU data are the mean ± SD (N =3), two-
way ANOVA followed by a Tukey–Kramer test. Results are representative of two independent
experiments. Scale bars = 2 µm.

530
531

532
533 **Figure S1.** The *in silico* analysis of AtCEPs gene expression using Tissue Specific Root eFP
534 (http://bar.utoronto.ca/eplant/). EXP7 was included as a root hair cell marker.

535
536 **Figure S2**. The *in silico* analysis of AtCEP1-AtCEP3 expression using Root Cell Atlas
537 (https://rootcellatlas.org/). EXP7 was included as a root hair cell marker.

538
539
540 **Figure S3**. **Single *cep* mutants do not perturb root hair growth.** Quantitative analysis of RH
541 length (mean ± s.e.m., n= 200) in Wt Col-0 and in single *cep* mutants. NS= not significant
542 difference. Data are shown as the mean ± SEM, (n=20).
543

544
545
546   **Figure S4**. The *in silico* analysis of NAC1 expression using Tissue Specific Root eFP
547   (http://bar.utoronto.ca/eplant/) (**A**) and using Root Cell Atlas (https://rootcellatlas.org/) (**B**).
548   EXP7 was included as a root hair cell marker.
549

550
551
552 **Figure S5**. Induction of CEP1 and CEP2 by NAC1-FLAG inducible line (PER8pro:3×FLAG-NAC1).
553 Gene-specific signals were normalized relatively to ACT2.
554

CEP3.1ΔKDEL 275YSEVIFGSLSLRQKTSYEKITYQC299

Figure S6. Protein alignment and domains of AtCEP1-AtCEP3 sequences from *Arabidopsis thaliana* with Ricinus CEP 1SV4 (RcCysEP). (**A**) Phylogenetic tree of Arabidopsis AtCEPs and Ricinus AtCEP. The phylogenetic analysis was carried out with MEGA6 (Tamura et al., 2007) using the Maximum Similarity method (Maximum Likelihood) (Saitou and Nei, 1987). The numbers in the nodes indicate the bootstrap values obtained for 1000 iterations. Scale represents the evolutionary distance, expressed as the number of substitutions per amino acid. (**B**) AtCEP protein domains. AtCEPs are synthesized as pre-pro-enzyme, which is then co-translationally synthesized into the ER, where the pre-sequence is removed, and pro-enzyme is finally released from the ricinosomes. KDEL is an ER retention signal peptide. (**C**) AtCEP1-AtCEP3 Protein alignment. PRE, PRO and KDEL domains are indicated.

Figure S7. Interaction by an *in silico* docking approach of AtCEP1, AtCEP2 and AtCEP3 with EXT peptides. (**A,B**) Ten docking results for each EXT *O*-glycosylation state are shown superimposed on the AtCEP1 protein surface to evaluate the consistency of docking sites. (**A**) Model of AtCEP1 (protein surface shown in grey) complexed to a non-*O*-glycosylated EXT substrate (SPPPYVY)$_3$ (in magenta). (**B**) Model of AtCEP1 (protein surface shown in gray) complexed to an *O*-glycosylated-EXT substrate (protein and *O*-glycans shown in light red, both depicted as sticks). Arabino-galactosylated EXT peptide = [(SOOOYVY)$_3$-AG]. (**C**) Comparison of the binding energy of different AtCEP1-AtCEP3 and 1S4V (Ricinus AtCEP included as a control) to EXT substrates with different degrees of *O*-glycosylation. (from left to right) A non-hydroxylated EXT peptide (SPPPYVY)$_3$, a hydroxylated but not *O*-glycosylated EXT peptide [(SOOOYVY)$_3$; O=hydroxyproline], only arabinosylated EXT-peptide [(SOOOYVY)$_3$-A], and arabino-galactosylated EXT peptide [(SOOOYVY)$_3$-AG] were analyzed.

582   **Table S1.** Mutant lines used in this study.

NAME	ATG code		Mutant line	Reference
NAC1	*At1g56010*	*nac1-2*	CS1013025	This work
AtCEP1	*At5g50260*	*cep1-2*	SALK_137016	This work
		cep1-3	SAIL_158_B06	(Höwing et al. 2014)
AtCEP2	*At3g48340*	*cep2-1*	SALK_079519	This work
		cep2-2	CSHET_6591	(Hierl et al. 2014)
AtCEP3	*At3g48350*	*cep3-1*	SALK_024692	This work
		cep3-2	SALK_016791	This work

585 **Table S2.** Transgenic lines used in this study.

NAME	CONSTRUCTION	BACKGROUND	REFERENCE
PER8pro:AtCEP2	promoterPER8::AtCEP2	Wt Col-0	(Chen et al. 2016)
PER8pro:AtCEP1	promoterPER8::AtCEP1	Wt Col-0	(Chen et al. 2016)
PER8pro:3×FLAG-NAC1	promoterPER8::3×FLAG-NAC1	Wt Col-0	(Chen et al. 2016)
NAC1pro:NAC1-GUS	promoterNAC1::NAC1-GUS	Wt Col-0	(Chen et al. 2016)
AtCEP1pro:3xHA-EGFP-AtCEP1-KDEL	promoterAtCEP1:: PRE-PRO-3xHA-EGFP-AtCEP1-KDEL	*cep1-3*	(Höwing et al. 2014)
AtCEP1pro:3xHA-EGFP-KDEL	promoterAtCEP1:: PRE-PRO-3xHA-EGFP-KDEL	*cep1-3*	(Höwing et al. 2014)
AtCEP2pro: 3xHA-mCherry-KDEL	promoterAtCEP2::PRE-PRO-3xHA-mCherry-KDEL	*cep2-2*	(Hierl et al. 2014)
AtCEP2pro:3xHA-mCherry -AtCEP2-KDEL	promoterAtCEP2::PRE-PRO-3xHA-mCherry-AtCEP2KDEL	Wt Col-0	(Hierl et al. 2014)
AtCEP3pro:GFP	promoterAtCEP3::GFP	Wt Col-0	This work
NAC1pro:GFP	promoterNAC1::GFP	Wt Col-0	This work

PP2A RT-F	GTCGACCAAGCGGTTGTGGAGA
PP2A RT-R	ACGCCCAACGAACAAATCACAGA
AtCEP1 RT-F	AAACCAAGGCCAATGCGGGAGTTG
AtCEP1 RT-R	TTCCGCATCTCCCGGTAAACACTC
AtCEP2 RT-F	CCGGTTCCAACATCAAGCATCAC
AtCEP2 RT-R	AGATCCGTAAACACTCCCTCTG
NAC1 RT-F	TGAGCTCTCCAAAGGAAGACTGG
NAC1 RT-R	ACAGAATGAGTCGAGGCCTGTG
CEP2 CHIP A F	ATGGGCTGCGCCAAATGTTC
CEP2 CHIP A R	ACGTCTTAAACCATGCACCGA
CEP2 CHIP B F	TCGGACTTTTTGATTGGAATCCTCT
CEP2 CHIP B R	TCGTGGACCGATCCAAACAG
CEP3 CHIP A F	GAGGACCCAACCCATCTTCATC
CEP3 CHIP A R	ACACGTACGTCCATTCCCATTT
E2Fa CHIP B- F	TTGCAGAGAATTGTGATT
E2Fa CHIP B- R	GAGAATCCGATCATAGAC
NAC1_RT-qPCR FW	CTCCTCGAGGCCGTAAAACC
NAC1_RT-qPCR RV	AGACCCAGTCTTCCTTTGGAG

Actin F	GGTAACATTGTGCTCAGTGGTGG
Actin R	AACGACCTTAATCTTCATGCTGC

Table S3. List of primer used.

REFERENCES

Battistella C, Guiet R, Burri O, Seitz A, Escrig S, Knott GW, Meibom A, Klok HA. (2019) Cellular
Uptake and Intracellular Trafficking of Poly(N-(2-Hydroxypropyl) Methacrylamide).
Biomacromolecules. 20(1):231-242. doi: 10.1021/acs.biomac.8b01372.

Baumberger N, Ringli C, Keller B. (2001). The chimeric leucinerich repeat/extensin cell wall
protein LRX1 is required for root hair morphogenesis in Arabidopsis thaliana. *Genes &*
Development 15, 1128–1139
Baumberger N, Steiner M, Ryser U, Keller B, Ringli C. (2003). Synergistic interaction of the two
paralogous Arabidopsis genes LRX1 and LRX2 in cell wall formation during root hair
development. *Plant Journal* 35, 71–81
Becker, C., Senyuk, V.I., Shutov, A.D., Nong, V.H., Fischer, J., Horstmann, C., and Müntz, K.
(1997). Proteinase A, a storage globulin-degrading endopeptidase of vetch (Vicia sativa
L.) seeds, is not involved in early steps of storage-protein mobilization. Eur. J. Biochem.
248: 304–312
Beers, E.P., Woffenden, B.J., Zhao C. (2000). Plant proteolytic enzymes: possible role during
programmed cell death. Plant Mol. Biol. 44, 399–415.doi: 10.1023/A:1026556928624
Berger F, Linstead P, Dolan L, Haseloff J. (1998) Stomata patterning on the hypocotyl of
Arabidopsis thaliana is controlled by genes involved in the control of root epidermis
patterning. *Dev Biol.* 194(2):226-34. doi: 10.1006/dbio.1997.8836.
Bolte S, Cordelières FP. (2006) A guided tour into subcellular colocalization analysis in light
microscopy. J Microsc. 224(Pt 3):213-32. doi: 10.1111/j.1365-2818.2006.01706.x.
Borassi C, Sede AR, Mecchia MA, Salgado Salter JD, Marzol E, Muschietti JP, Estevez JM.
(2016) An update on cell surface proteins containing extensin-motifs. J Exp Bot.
67(2):477-87. doi: 10.1093/jxb/erv455
Cannon MC et al (2008) Self-assembly of the plant cell wall requires an extensin scaffold. Proc
Natl Acad Sci USA 105:2226–2231.doi:10.1073/pnas.0711980105
Chen X., Cheng J., Chen L., Zhang G., Huang, H., Zhang Y., Xu L. (2016) Auxin-independent *NAC*
Pathway Acts in Response to Explant-Specific Wounding and Promotes Root Tip
Emergence during *De Novo* Root Organogenesis in *Arabidopsis*. Plant Physiol.
DOI:10.1104/pp.15.01733
Cheng Z, Guo X, Zhang J, Liu Y, Wang B, Li H, Lu H. (2020) βVPE is involved in tapetal
degradation and pollen development by activating proprotease maturation in
Arabidopsis thaliana. J Exp Bot. 25;71(6):1943-1955. doi: 10.1093/jxb/erz560
Fabrice T, Vogler H, Draeger C, Munglani G, Gupta S, Herger A, Knox P, Grossniklaus U,
Ringli C. (2018). LRX Proteins play a crucial role in pollen grain and pollen tube cell wall
development. *Plant Physiology* 176: 1981–1992
Grebe M, Xu J, Möbius W, Ueda T, Nakano A, Geuze HJ, Rook MB, Scheres B. (2003)
Arabidopsis sterol endocytosis involves actin-mediated trafficking via ARA6-positive
early endosomes. Curr Biol. 13(16):1378-87. doi: 10.1016/s0960-9822(03)00538-4.
Greenwood JS, Helm M, Gietl C (2005) Ricinosomes and endosperm transfer cell structure in
programmed cell death of the nucellus during Ricinus seed development. Proc Natl Acad
Sci USA 102:2238–2243. doi:10.1073/pnas.0409429102
Grierson C, Nielsen E, Ketelaarc T, Schiefelbein J. Root hairs. (2014) Arabidopsis Book.
12:e0172. doi: 10.1199/tab.0172.

Hall Q, Cannon MC. (2002) The cell wall hydroxyproline-rich glycoprotein RSH is essential for
normal embryo development in Arabidopsis. Plant Cell. 14(5):1161-72. doi:
10.1105/tpc.010477

He, X., and Kermode, A.R. (2003). Proteases associated with programmed cell death of
megagametophyte cells after germination of white spruce (Picea glauca) seeds. Plant
Mol. Biol. 52: 729–744

Helm M et al (2008) KDEL-tailed cysteine endopeptidases involved in programmed cell death,
intercalation of new cells, and dismantling of extensin scaffolds. Am. J. Bot. 95:1049–
1062. doi:10.3732/ajb.2007404

Hierl G, Vothknecht U, Gietl C (2012) Programmed cell death in *Ricinus* and *Arabidopsis*: the
function of KDEL cysteine peptidases in development. Physiol Plant 145:103–113.
doi:10.1111/j.1399-3054.2012.01580.x

Hierl G, Howing T, Isono E, Lottspeich F, Gietl C (2014) Ex vivo processing for maturation of
Arabidopsis KDEL-tailed cysteine endopeptidase 2 (AtCEP2) pro-enzyme and its storage
in endoplasmic reticulum derived organelles. Plant Mol Biol 84:605–620.
doi:10.1007/s11103-013-0157-6

Howing T, Huesmann C, Hoefle C, Nagel MK, Isono E, Huckelhoven R, Gietl C (2014)
Endoplasmic reticulum KDEL-tailed cysteine endopeptidase 1 of Arabidopsis (AtCEP1) is
involved in pathogen defense. Front Plant Sci 5:58. doi:10.3389/fpls.2014. 00058

Howing T, Dann M, Muller B, Helm M, Scholz S, Schneitz K, et al. (2018) The role of KDEL-
tailed cysteine endopeptidases of Arabidopsis (AtCEP2 and AtCEP1) in root
development. PLoS ONE 13(12): e0209407.
https://doi.org/10.1371/journal.pone.0209407

Jackson, P.A., Galinha, C.I., Pereira, C.S., Fortunato, A., Soares, N.C., Amâncio, S.B., and Pinto
Ricardo, C.P. (2001). Rapid deposition of extensin during the elicitation of grapevine
callus cultures is specifically catalyzed by a 40-kilodalton peroxidase. *Plant Physiol.* 127:
1065–1076

Maher KA, Bajic M, Kajala K, Reynoso M et al. (2018) Profiling of Accessible Chromatin Regions
across Multiple Plant Species and Cell Types Reveals Common Gene Regulatory
Principles and New Control Modules. *Plant Cell* 30(1):15-36.

Manders EMM, Verbeek FJ, Aten JA. (1993) Measurement of co-localization of objects in dual-
colour confocal images. *J Microsc*. 169(3):375-382. doi: 10.1111/j.1365-
2818.1993.tb03313.x.

Martinez-Pacheco J, Ranocha P, Kasulin L, Fusari CM, et al. (2022) Apoplastic class III
peroxidases PRX62 and PRX69 promote Arabidopsis root hair growth at low
temperature. Nat Commun. 14;13(1):1310. doi: 10.1038/s41467-022-28833-4

Marzol E, Borassi C, Bringas M, Sede A, Rodríguez Garcia DR, Capece L, Estevez JM. (2018).
Filling the Gaps to Solve the Extensin Puzzle. *Mol Plant*. 11(5):645-658

Marzol E, Borassi C, Carignani Sardoy M, Ranocha P et al. (2022) Class III Peroxidases PRX01,
PRX44, and PRX73 Control Root Hair Growth in *Arabidopsis thaliana*. Int J Mol Sci.
11;23(10):5375. doi: 10.3390/ijms23105375

Merkouropoulos G, Shirsat AH. (2003) The unusual Arabidopsis extensin gene atExt1 is
expressed throughout plant development and is induced by a variety of biotic and
abiotic stresses. Planta. 217(3):356-66. doi: 10.1007/s00425-003-1002-y

Ndinyanka Fabrice T, Kaech A, Barmettler G, Eichenberger C, Knox JP, Grossniklaus U, Ringli C
(2017) Efficient preparation of Arabidopsis pollen tubes for ultrastructural analysis using
chemical and cryo-fixation. BMC Plant Biol 17: 176

Olvera-Carrillo Y et al (2015) A conserved core of programmed cell death indicator genes
discriminates developmentally and environmentally induced programmed cell death in
plants. Plant Physiol 169:2684–2699. doi:10.1104/pp.15.00769

Ringli C. (2010). The hydroxyproline-rich glycoprotein domain of the Arabidopsis LRX1
requires Tyr for function but not for insolubilization in the cell wall. *Plant Journal* 63,
662–669

Sede AR, Borassi C, Wengier DL, Mecchia MA, Estevez JM, Muschietti JP (2018) Arabidopsis
pollen extensins LRX are required for cell wall integrity during pollen tube growth. FEBS
Lett 592:233-243

Schmid M, Simpson D, Gietl C (1999) Programmed cell death in castor bean endosperm is
associated with the accumulation and release of a cysteine endopeptidase from
ricinosomes. Proc Natl Acad Sci USA 96:14159–14164. doi:10.1073/pnas.96.24.14159

Schmid M, Simpson DJ, Sarioglu H, Lottspeich F, Gietl C (2001) The ricinosomes of senescing
plant tissue bud from the endoplasmic reticulum. Proc Natl Acad Sci USA 98:5353–5358.
doi:10.1073 /pnas.061038298

Schnabelrauch LS, Kieliszewski M, Upham BL, Alizedeh H, Lamport DT. (1996) Isolation of pI
4.6 extensin peroxidase from tomato cell suspension cultures and identification of Val-
Tyr-Lys as putative intermolecular cross-link site. Plant J. 9(4):477-89. doi:
10.1046/j.1365-313x.1996.09040477.x.

Showalter AM, Keppler B, Lichtenberg J, Gu D, Welch LR. (2010). A bioinformatics approach to
the identification, classification, and analysis of hydroxyproline-rich glycoproteins. Plant
physiology 153, 485–513

Strasser, R. (2016). Plant protein glycosylation. Glycobioogy 1014. doi:
10.1093/glycob/cww023

Su W, Liu Y, Xia Y, Hong Z, Li J. (2012). The Arabidopsis homolog of the mammalian OS-9
protein plays a key role in the endoplasmic reticulum associated degradation of
misfolded receptor-like kinases. Mol Plant. 5:929–940

Su W, Liu Y, Xia Y, Hong Z, Li J. (2011). Conserved endoplasmic reticulum-associated
degradation system to eliminate mutated receptor-like kinases in Arabidopsis. Proc Natl
Acad Sci USA. 108:870–875

Tanaka, T., Yamauchi, D., and Minamikawa, T. (1991). Nucleotide sequence of cDNA for an
endopeptidase (EP-C1) from pods of maturing Phaseolus vulgaris fruits. Plant Mol. Biol.
16: 1083–1084

Than ME, Helm M, Simpson DJ, Lottspeich F, Huber R, Gietl C (2004) The 2.0 A crystal
structure and substrate specificity of the KDEL-tailed cysteine endopeptidase
functioning in programmed cell death of Ricinus communis endosperm. J Mol Biol
336:1103–1116. doi:10.1016/j.jmb.2003.12.075

Velasquez SM, Ricardi MM, Dorosz JG, *et al*. (2011). O-Glycosylated Cell Wall Proteins Are
Essential in Root Hair Growth. Science 332, 1401–1403.

Velasquez SM, Marzol E, Borassi C, Pol-Fachin L, Ricardi MM, et al. (2015). Low sugar is not
always good: Impact of specific O-glycan defects on tip growth in Arabidopsis. Plant
physiology 168, 808–813

Velasquez M, Salter JS, Dorosz JG, Petersen BL, Estevez JM. (2012). Recent advances on the
posttranslational modifications of EXTs and their roles in plant cell walls. Frontiers in
plant science 3, 93

Wang, X., Wang, K., Yin, G., Liu, X., Liu, M., Cao, N., Duan, Y., Gao, H., Wang, W., Ge, W., et al.
(2018). Pollen-expressed leucine-rich repeat extensins are essential for pollen
germination and growth. *Plant Physiol*. 176, 1993–2006

Xie Q, Frugis G, Colgan D, Chua NH (2000) Arabidopsis NAC1 transduces auxin signal
downstream of TIR1 to promote lateral root development. Genes Dev 14: 3024–3036

Xie C, Li C, Wang F, Zhang F, Liu J, Wang J, Zhang X, Kong X, Ding Z. (2023) NAC1 regulates root
ground tissue maturation through coordinating with SCR/SHR-CYCD6;1 module in
Arabidopsis. Mol Plant. S1674-2052(23)00039-4. doi: 10.1016/j.molp.2023.02.006.

Zhang D et al (2014) The cysteine protease AtCEP1, a key executor involved in tapetal
programmed cell death, regulates pollen development in Arabidopsis. Plant Cell
26:2939–2961. doi:10.1105/tpc.114.127282

Zhou L-Z, Howing, T., Muller B., Hammes U.Z, Gietl C., Dresselhaus T. (2016) Expression
analysis of KDEL-CysEPs programmed cell death markers during reproduction in
Arabidopsis. Plant Reprod DOI 10.1007/s00497-016-0288-4

Zuo J, Niu QW, Chua NH (2000) Technical advance: An estrogen receptor-based transactivator
XVE mediates highly inducible gene expression in transgenic plants. Plant J. 24: 265-273

Parsed Citations

Battistella C, Guiet R, Burri O, Seitz A, Escrig S, Knott GW, Meibom A, Klok HA. (2019) Cellular Uptake and Intracellular Trafficking of Poly(N-(2-Hydroxypropyl) Methacrylamide). Biomacromolecules. 20(1):231-242. doi: 10.1021/acs.biomac.8b01372.
  Google Scholar: Author Only Title Only Author and Title

Baumberger N, Ringli C, Keller B. (2001). The chimeric leucinerich repeat/extensin cell wall protein LRX1 is required for root hair morphogenesis in Arabidopsis thaliana. Genes & Development 15, 1128–1139
  Google Scholar: Author Only Title Only Author and Title

Baumberger N, Steiner M, Ryser U, Keller B, Ringli C. (2003). Synergistic interaction of the two paralogous Arabidopsis genes LRX1 and LRX2 in cell wall formation during root hair development. Plant Journal 35, 71–81
  Google Scholar: Author Only Title Only Author and Title

Becker, C., Senyuk, V.I., Shutov, A.D., Nong, V.H., Fischer, J., Horstmann, C., and Müntz, K. (1997). Proteinase A, a storage globulin-degrading endopeptidase of vetch (Vicia sativa L.) seeds, is not involved in early steps of storage-protein mobilization. Eur. J. Biochem. 248: 304–312
  Google Scholar: Author Only Title Only Author and Title

Beers, E.P., Woffenden, B.J., Zhao C. (2000). Plant proteolytic enzymes: possible role during programmed cell death. Plant Mol. Biol. 44, 399–415.doi: 10.1023/A:1026556928624
  Google Scholar: Author Only Title Only Author and Title

Berger F, Linstead P, Dolan L, Haseloff J. (1998) Stomata patterning on the hypocotyl of Arabidopsis thaliana is controlled by genes involved in the control of root epidermis patterning. Dev Biol. 194(2):226-34. doi: 10.1006/dbio.1997.8836.
  Google Scholar: Author Only Title Only Author and Title

Bolte S, Cordelières FP. (2006) A guided tour into subcellular colocalization analysis in light microscopy. J Microsc. 224(Pt 3):213-32. doi: 10.1111/j.1365-2818.2006.01706.x.
  Google Scholar: Author Only Title Only Author and Title

Borassi C, Sede AR, Mecchia MA, Salgado Salter JD, Marzol E, Muschietti JP, Estevez JM. (2016) An update on cell surface proteins containing extensin-motifs. J Exp Bot. 67(2):477-87. doi: 10.1093/jxb/erv455
  Google Scholar: Author Only Title Only Author and Title

Cannon MC et al (2008) Self-assembly of the plant cell wall requires an extensin scaffold. Proc Natl Acad Sci USA 105:2226–2231.doi:10.1073/pnas.0711980105
  Google Scholar: Author Only Title Only Author and Title

Chen X., Cheng J., Chen L., Zhang G., Huang, H., Zhang Y., Xu L. (2016) Auxin-independent NAC Pathway Acts in Response to Explant-Specific Wounding and Promotes Root Tip Emergence during De Novo Root Organogenesis in Arabidopsis. Plant Physiol. DOI:10.1104/pp.15.01733
  Google Scholar: Author Only Title Only Author and Title

Cheng Z, Guo X, Zhang J, Liu Y, Wang B, Li H, Lu H. (2020) βVPE is involved in tapetal degradation and pollen development by activating proprotease maturation in Arabidopsis thaliana. J Exp Bot. 25;71(6):1943-1955. doi: 10.1093/jxb/erz560
  Google Scholar: Author Only Title Only Author and Title

Fabrice T, Vogler H, Draeger C, Munglani G, Gupta S, Herger A, Knox P, Grossniklaus U, Ringli C. (2018). LRX Proteins play a crucial role in pollen grain and pollen tube cell wall development. Plant Physiology 176: 1981–1992
  Google Scholar: Author Only Title Only Author and Title

Grebe M, Xu J, Möbius W, Ueda T, Nakano A, Geuze HJ, Rook MB, Scheres B. (2003) Arabidopsis sterol endocytosis involves actin-mediated trafficking via ARA6-positive early endosomes. Curr Biol. 13(16):1378-87. doi: 10.1016/s0960-9822(03)00538-4.
  Google Scholar: Author Only Title Only Author and Title

Greenwood JS, Helm M, Gietl C (2005) Ricinosomes and endosperm transfer cell structure in programmed cell death of the nucellus during Ricinus seed development. Proc Natl Acad Sci USA 102:2238–2243. doi:10.1073/pnas.0409429102
  Google Scholar: Author Only Title Only Author and Title

Grierson C, Nielsen E, Ketelaarc T, Schiefelbein J. Root hairs. (2014) Arabidopsis Book. 12:e0172. doi: 10.1199/tab.0172.
  Google Scholar: Author Only Title Only Author and Title

Hall Q, Cannon MC. (2002) The cell wall hydroxyproline-rich glycoprotein RSH is essential for normal embryo development in Arabidopsis. Plant Cell. 14(5):1161-72. doi: 10.1105/tpc.010477
  Google Scholar: Author Only Title Only Author and Title

He, X., and Kermode, A.R. (2003). Proteases associated with programmed cell death of megagametophyte cells after germination of white spruce (Picea glauca) seeds. Plant Mol. Biol. 52: 729–744
  Google Scholar: Author Only Title Only Author and Title

Helm M et al (2008) KDEL-tailed cysteine endopeptidases involved in programmed cell death, intercalation of new cells, and dismantling of extensin scaffolds. Am. J. Bot. 95:1049–1062. doi:10.3732/ajb.2007404
Google Scholar: Author Only Title Only Author and Title

Hierl G, Vothknecht U, Gietl C (2012) Programmed cell death in Ricinus and Arabidopsis: the function of KDEL cysteine peptidases in development. Physiol Plant 145:103–113. doi:10.1111/j.1399-3054.2012.01580.x
Google Scholar: Author Only Title Only Author and Title

Hierl G, Howing T, Isono E, Lottspeich F, Gietl C (2014) Ex vivo processing for maturation of Arabidopsis KDEL-tailed cysteine endopeptidase 2 (AtCEP2) pro-enzyme and its storage in endoplasmic reticulum derived organelles. Plant Mol Biol 84:605–620. doi:10.1007/s11103-013-0157-6
Google Scholar: Author Only Title Only Author and Title

Howing T, Huesmann C, Hoefle C, Nagel MK, Isono E, Huckelhoven R, Gietl C (2014) Endoplasmic reticulum KDEL-tailed cysteine endopeptidase 1 of Arabidopsis (AtCEP1) is involved in pathogen defense. Front Plant Sci 5:58. doi:10.3389/fpls.2014. 00058
Google Scholar: Author Only Title Only Author and Title

Howing T, Dann M, Muller B, Helm M, Scholz S, Schneitz K, et al. (2018) The role of KDEL-tailed cysteine endopeptidases of Arabidopsis (AtCEP2 and AtCEP1) in root development. PLoS ONE 13(12): e0209407. https://doi.org/10.1371/journal.pone.0209407
Google Scholar: Author Only Title Only Author and Title

Jackson, P.A., Galinha, C.I., Pereira, C.S., Fortunato, A, Soares, N.C., Amâncio, S.B., and Pinto Ricardo, C.P. (2001). Rapid deposition of extensin during the elicitation of grapevine callus cultures is specifically catalyzed by a 40-kilodalton peroxidase. Plant Physiol. 127: 1065–1076
Google Scholar: Author Only Title Only Author and Title

Maher KA, Bajic M, Kajala K, Reynoso M et al. (2018) Profiling of Accessible Chromatin Regions across Multiple Plant Species and Cell Types Reveals Common Gene Regulatory Principles and New Control Modules. Plant Cell 30(1):15-36.
Google Scholar: Author Only Title Only Author and Title

Manders EMM, Verbeek FJ, Aten JA. (1993) Measurement of co-localization of objects in dual-colour confocal images. J Microsc. 169(3):375-382. doi: 10.1111/j.1365-2818.1993.tb03313.x.
Google Scholar: Author Only Title Only Author and Title

Martinez-Pacheco J, Ranocha P, Kasulin L, Fusari CM, et al. (2022) Apoplastic class III peroxidases PRX62 and PRX69 promote Arabidopsis root hair growth at low temperature. Nat Commun. 14;13(1):1310. doi: 10.1038/s41467-022-28833-4
Google Scholar: Author Only Title Only Author and Title

Marzol E, Borassi C, Bringas M, Sede A, Rodríguez Garcia DR, Capece L, Estevez JM. (2018). Filling the Gaps to Solve the Extensin Puzzle. Mol Plant. 11(5):645-658
Google Scholar: Author Only Title Only Author and Title

Marzol E, Borassi C, Carignani Sardoy M, Ranocha P et al. (2022) Class III Peroxidases PRX01, PRX44, and PRX73 Control Root Hair Growth in Arabidopsis thaliana. Int J Mol Sci. 11;23(10):5375. doi: 10.3390/ijms23105375
Google Scholar: Author Only Title Only Author and Title

Merkouropoulos G, Shirsat AH. (2003) The unusual Arabidopsis extensin gene atExt1 is expressed throughout plant development and is induced by a variety of biotic and abiotic stresses. Planta. 217(3):356-66. doi: 10.1007/s00425-003-1002-y
Google Scholar: Author Only Title Only Author and Title

Ndinyanka Fabrice T, Kaech A, Barmettler G, Eichenberger C, Knox JP, Grossniklaus U, Ringli C (2017) Efficient preparation of Arabidopsis pollen tubes for ultrastructural analysis using chemical and cryo-fixation. BMC Plant Biol 17: 176
Google Scholar: Author Only Title Only Author and Title

Olvera-Carrillo Y et al (2015) A conserved core of programmed cell death indicator genes discriminates developmentally and environmentally induced programmed cell death in plants. Plant Physiol 169:2684–2699. doi:10.1104/pp.15.00769
Google Scholar: Author Only Title Only Author and Title

Ringli C. (2010). The hydroxyproline-rich glycoprotein domain of the Arabidopsis LRX1 requires Tyr for function but not for insolubilization in the cell wall. Plant Journal 63, 662–669
Google Scholar: Author Only Title Only Author and Title

Sede AR, Borassi C, Wengier DL, Mecchia MA, Estevez JM, Muschietti JP (2018) Arabidopsis pollen extensins LRX are required for cell wall integrity during pollen tube growth. FEBS Lett 592:233-243
Google Scholar: Author Only Title Only Author and Title

Schmid M, Simpson D, Gietl C (1999) Programmed cell death in castor bean endosperm is associated with the accumulation and release of a cysteine endopeptidase from ricinosomes. Proc Natl Acad Sci USA 96:14159–14164. doi:10.1073/pnas.96.24.14159
Google Scholar: Author Only Title Only Author and Title

Schmid M, Simpson DJ, Sarioglu H, Lottspeich F, Gietl C (2001) The ricinosomes of senescing plant tissue bud from the endoplasmic reticulum. Proc Natl Acad Sci USA 98:5353–5358. doi:10.1073 /pnas.061038298

Google Scholar: Author Only Title Only Author and Title

Schnabelrauch LS, Kieliszewski M, Upham BL, Alizedeh H, Lamport DT. (1996) Isolation of pI 4.6 extensin peroxidase from tomato cell suspension cultures and identification of Val-Tyr-Lys as putative intermolecular cross-link site. Plant J. 9(4):477-89. doi: 10.1046/j.1365-313x.1996.09040477.x.

Google Scholar: Author Only Title Only Author and Title

Showalter AM, Keppler B, Lichtenberg J, Gu D, Welch LR. (2010). A bioinformatics approach to the identification, classification, and analysis of hydroxyproline-rich glycoproteins. Plant physiology 153, 485–513

Google Scholar: Author Only Title Only Author and Title

Strasser, R. (2016). Plant protein glycosylation. Glycobiogy 1014. doi: 10.1093/glycob/cww023

Google Scholar: Author Only Title Only Author and Title

Su W, Liu Y, Xia Y, Hong Z, Li J. (2012). The Arabidopsis homolog of the mammalian OS-9 protein plays a key role in the endoplasmic reticulum associated degradation of misfolded receptor-like kinases. Mol Plant. 5:929–940

Google Scholar: Author Only Title Only Author and Title

Su W, Liu Y, Xia Y, Hong Z, Li J. (2011). Conserved endoplasmic reticulum-associated degradation system to eliminate mutated receptor-like kinases in Arabidopsis. Proc Natl Acad Sci USA. 108:870–875

Google Scholar: Author Only Title Only Author and Title

Tanaka, T., Yamauchi, D., and Minamikawa, T. (1991). Nucleotide sequence of cDNA for an endopeptidase (EP-C1) from pods of maturing Phaseolus vulgaris fruits. Plant Mol. Biol. 16: 1083–1084

Google Scholar: Author Only Title Only Author and Title

Than ME, Helm M, Simpson DJ, Lottspeich F, Huber R, Gietl C (2004) The 2.0 A crystal structure and substrate specificity of the KDEL-tailed cysteine endopeptidase functioning in programmed cell death of Ricinus communis endosperm. J Mol Biol 336:1103–1116. doi:10.1016/j.jmb.2003.12.075

Google Scholar: Author Only Title Only Author and Title

Velasquez SM, Ricardi MM, Dorosz JG, et al. (2011). O-Glycosylated Cell Wall Proteins Are Essential in Root Hair Growth. Science 332, 1401–1403.

Google Scholar: Author Only Title Only Author and Title

Velasquez SM, Marzol E, Borassi C, Pol-Fachin L, Ricardi MM, et al. (2015). Low sugar is not always good: Impact of specific O-glycan defects on tip growth in Arabidopsis. Plant physiology 168, 808–813

Google Scholar: Author Only Title Only Author and Title

Velasquez M, Salter JS, Dorosz JG, Petersen BL, Estevez JM. (2012). Recent advances on the posttranslational modifications of EXTs and their roles in plant cell walls. Frontiers in plant science 3, 93

Google Scholar: Author Only Title Only Author and Title

Wang, X., Wang, K., Yin, G., Liu, X., Liu, M., Cao, N., Duan, Y., Gao, H., Wang, W., Ge, W., et al. (2018). Pollen-expressed leucine-rich repeat extensins are essential for pollen germination and growth. Plant Physiol. 176, 1993–2006

Google Scholar: Author Only Title Only Author and Title

Xie Q, Frugis G, Colgan D, Chua NH (2000) Arabidopsis NAC1 transduces auxin signal downstream of TIR1 to promote lateral root development. Genes Dev 14: 3024–3036

Google Scholar: Author Only Title Only Author and Title

Xie C, Li C, Wang F, Zhang F, Liu J, Wang J, Zhang X, Kong X, Ding Z. (2023) NAC1 regulates root ground tissue maturation through coordinating with SCR/SHR-CYCD6;1 module in Arabidopsis. Mol Plant. S1674-2052(23)00039-4. doi: 10.1016/j.molp.2023.02.006.

Google Scholar: Author Only Title Only Author and Title

Zhang D et al (2014) The cysteine protease AtCEP1, a key executor involved in tapetal programmed cell death, regulates pollen development in Arabidopsis. Plant Cell 26:2939–2961. doi:10.1105/tpc.114.127282

Google Scholar: Author Only Title Only Author and Title

Zhou L-Z, Howing, T., Muller B., Hammes U.Z, Gietl C., Dresselhaus T. (2016) Expression analysis of KDEL-CysEPs programmed cell death markers during reproduction in Arabidopsis. Plant Reprod DOI 10.1007/s00497-016-0288-4

Google Scholar: Author Only Title Only Author and Title

Zuo J, Niu QW, Chua NH (2000) Technical advance: An estrogen receptor-based transactivator XVE mediates highly inducible gene expression in transgenic plants. Plant J. 24: 265-273

Google Scholar: Author Only Title Only Author and Title