CanFlyet: Habitat Zone and Diet Trait Dataset for Diptera Species of Canada and Greenland

Majoros, S.E.[1] & Adamowicz, S.J.[1]

[1]University of Guelph

Abstract

Background

True flies (Diptera) are an ecologically important group that play a role in agriculture, public health and ecosystem functioning. As researchers continue to investigate this order, it is beneficial to link the growing occurrence data to biological traits. However, large-scale ecological trait data are not readily available for fly species. While some databases and datasets include fly data, many ecologically relevant traits for taxa of interest are not included. In this study we create a dataset containing ecological traits (habitat and diet) for fly species of Canada and Greenland having occurrence records on the Barcode of Life Data Systems (BOLD). We present a dataset containing trait information for 983 Diptera species.

New Information

Diptera were chosen for the dataset based on the occurrence records available for Diptera species from Canada and Greenland on the Barcode of Life Data Systems (BOLD). Trait data were then compiled based on literature searches conducted from April 2021 - January 2023 and assigned at the lowest taxonomic level possible. Three biological traits were included: habitat, larval diet, and adult diet. The dataset contains traits for 983 species across 380 genera, 25 subfamilies, and 61 families. This dataset allows for assignment of traits to occurrence data for Diptera species and can be used for further research into the ecology, evolution, and conservation of this order.

Keywords

Diptera, Canada, Greenland, dataset, habitat, diet, ecological traits

Introduction

True flies (Diptera) are a diverse, widespread taxonomic group, occurring across a wide range of ecological niches and geographic regions (Adler & Courtney, 2019, Marshall, 2012). This important group occupies various ecological roles, many of which have an impact on the environment and agricultural practices. Fly species can act as pests to crops, such as *Liriomyza bryoniae* on cabbages, lettuce, and tomatoes (Bragard et al., 2020); *Delia radicum* on broccoli and cauliflower (Mesmin et al., 2019); *Dasineura mali* on apples (Wearing et al., 2013); and *Bradysia ocellaris* and *Lycoriella ingenua* on mushrooms (Shamshad 2010). Diptera also includes pests of livestock such as the blood-feeding *Stomoxys calcitrans* (Olafson et al., 2021). Flies are also a group of interest with regard to public health, as some, like *Calliphora vomitoria*, are vectors of pathogenic microorganisms (Bedini et al., 2017). Despite these detrimental examples, flies also provide many important ecological benefits. Flies are an important pollinator group, especially in Arctic and Northern environments (Tiusanen et al., 2016). The

family Muscidae includes species that are pollinators of key plant species in this region and play an important role in the ecosystem (Tiusanen et al., 2016). Flies also fill various other ecological roles, such as parasitoids of other insects that act as vectors for pathogenic bacteria (Molinatto et al., 2020). Many species act as ecosystem engineers; changing environments through suspension-feeding, grazing, burrowing, and predation, as well as serving as an important food source for other organisms (Adler & Courtney, 2019). The saprophagous diet of many flies is also important for the breakdown of decaying organic material (Adler & Courtney, 2019). As biocontrol agents and bioindicators of water quality, flies act as a valuable resource for food production and research (Adler & Courtney, 2019).

This ecologically important group has been the focus of large DNA barcoding efforts, such as the Global Malaise Trap Program, which seeks to document Diptera diversity (which is likely under described) and open doors for ecological research and monitoring (deWaard et al., 2018, Geiger et al., 2016, Global Malaise Program, n.d.). DNA barcoding involves using a standardized gene region to identify and delineate species (Hebert et al., 2003), and there is a large amount of DNA barcoding data available on databases such as the Barcode of Life Data System (BOLD; Ratnasingham and Hebert, 2007), which contained 3 235 922 Diptera records representing 30 259 species as of Jan 18th, 2023. There is also a large amount of species occurrence data on databases such as the Global Biodiversity Information Facility (GBIF; GBIF.org, 2023), which contained 16 372 311 occurrences as of Feb 13th, 2023. As Diptera research continues, it is valuable to link such occurrence records with biological data.

However, finding biological trait data for fly species can be a challenge. There are various databases or datasets currently available for Diptera and other insect groups, covering a varying number of traits and taxa (Table 1). There are few databases that focus solely on Diptera, and while flies are included in some databases, not all species and traits of interest are included. A larger focus also appears to be on morphological traits, such as body size, with information on ecological traits, such as habitat and diet, being less readily available. Finally, few databases focus on Diptera from Canada or Greenland. Overall, traits for many Diptera species are very difficult to find and, when known, are typically present in taxon-specific studies rather than compiled datasets. The goal of this study was to create a dataset containing ecological traits (habitat and diet) for selected Diptera species found in Canada and Greenland. Species were selected that have multiple occurrence records and high-quality DNA barcode sequences on BOLD (following criteria outlined in Majoros et al. (2023)), which likely includes many of the most common species in the study region. This dataset can be used in future research to further our knowledge and understanding of this important taxonomic group.

General Description

This study provides a dataset containing the habitat and adult and larval diet for Diptera species of Canada and Greenland having occurrence records on BOLD. The traits were assigned to taxa using the currently available literature. This dataset will aid in further research that involves the ecological traits of Diptera species.

Project Description

Title: CanFlyet: Habitat Zone and Diet Trait Dataset for Diptera Species of Canada and Greenland

Personnel: Samantha E. Majoros & Sarah J. Adamowicz

Design Description: The dataset provides habitat, larval diet, and adult diet for Diptera species found in Canada and Greenland. Traits were determined based on a literature search.

Funding: Funding support for this project comes from the Natural Sciences and Engineering Research Council of Canada; the Government of Canada through Genome Canada and Ontario Genomics; the Ontario Ministry of Economic Development, Job Creation and Trade; and Food from Thought: Agricultural Systems for a Healthy Planet Initiative program funded by the Government of Canada through the Canada First Research Excellence Fund.

Sampling Methods

Description: The fly species were chosen for inclusion in this dataset by first downloading data for Diptera from Canada and Greenland from BOLD directly into R using BOLD's application programming interface (API) on June 24th, 2021. The records were filtered based on the requirements outlined in Majoros et al. (2023), and the remaining species were chosen for analysis and inclusion in this dataset. In Majoros et al. (2023), species were represented by Barcode Index Numbers (BINS), which are operational taxonomic units (OTUs) used by BOLD that are clusters of barcode sequences similar to species (Ratnasingham & Hebert, 2013). For other portions of the analysis, records were clustered by a 4% clustering threshold, and this was used in the place of BINs. Records needed to possess high quality DNA sequence data to be included in the analysis. BINs and clusters were included in the analysis if they possessed at least 20 records and were found in at least two geographic regions meeting the requirements of Majoros et al. (2023). They also needed to be represented by at least 10 records in each region they were found in. The species represented by these BINs and clusters were chosen and trait data was obtained. Additional species from Greenland were chosen based on occurrence records from GBIF (GBIF.org (June 24th, 2021)) GBIF Occurrence Download (https://doi.org/10.15468/dl.mk52hp) and included in the dataset. The traits were assigned to taxa using a series of literature searches conducted between April 2021 and January 2023.

Sampling Description: The biological traits for each species were determined and assigned through literature searches conducted from April 2021 - January 2023. Through the Omni Academic search tool available through the University of Guelph and Google Scholar, traits were found using the following search terms: trait AND "Taxonomic name", habitat AND "Taxonomic name", diet AND "Taxonomic name", and "Feeding mode" AND "Taxonomic name". The title and DOI of each reference are included in the dataset, and the full references are available in Appendix A. Traits were assigned to the lowest taxonomic level possible; however, not all traits could be assigned at the species level. For these species, traits were assigned using data from the next lowest level available, whether genus, subfamily or family. For example, if the authors of a given study mentioned that a particular genus has a specific trait, then species belonging to that genus were assigned that trait. The taxonomic level of the trait data is included in the dataset.

For habitat, species were classified as terrestrial (defined as taxa that live primarily in land habitats), aquatic (defined as taxa that live primarily in water bodies or associated habitats), or semi-aquatic (defined as species that require both aquatic and terrestrial habitats). These assignments were

made based on the larval habitat requirements. For adult and larval diet, species could be classified as predaceous (defined as taxa who prey on insects or other animals), phytophagous (defined as taxa who feed on plant material), fungivores (defined as those who feed on fungi), saprophagous (defined as those who feed on decaying organic matter), omnivores (defined as those who consume material of both plant and animal origin), parasites (defined as those who live and feed in or on an organism of another species), parasitoid (defined as organism whose young develop within a host, eventually resulting in host death), or nectivore (defined as taxa that primarily feed on nectar). Taxa can also be assigned a combination of the diet categories.

Geographic Coverage

Description: This dataset includes Diptera species found in Canada and Greenland that also have multiple occurrence records and DNA barcode sequences publicly available on BOLD.

Taxonomic Coverage

Description: The dataset contains traits for 983 species across 380 genera, and 61 families. Traits are also provided for 25 subfamilies. The families are represented by varying numbers of species within the dataset (Fig. 1). For several families, the dataset only contains information for one species. The most species-rich family in this dataset is Chironomidae, which is represented by 182 species.

Trait Coverage

Description: The dataset contains information for three biological traits: habitat, adult diet, and larval diet. For habitat, 656 species were terrestrial, 235 were aquatic, and 92 were semi-aquatic (Fig 2a). For adult diet, 101 were saprophagous, 42 were phytophagous, 247 were nectivores, 138 were predaceous, 4 were fungivores, 1 was an omnivore, 36 were both nectivores and predaceous, 1 was both phytophagous and a fungivore, 4 were both fungivores and nectivores, 24 were both nectivores and saprophagous, and 385 were non-feeding (Fig. 2b). For larval diet, 296 were saprophagous, 164 were phytophagous, 209 were predaceous, 62 were fungivores, 2 were parasitic, 21 were omnivores, 36 were parasitoids, 30 were both phytophagous and fungivores, and 163 were both phytophagous and saprophagous (Fig. 2c).

Temporal Coverage

Notes: The literature searches were conducted from April 2021 to January 2023. The papers referenced were published from 1955 – 2022.

Usage licence

Usage licence: Creative Commons Public Domain Wavier (CC-Zero)

Data Resources

Data package title: CanFlyet: Habitat Zone and Diet Trait Dataset for Diptera Species of Canada and Greenland

Number of data sets: 1

Data set name: CanFlyet: Habitat Zone and Diet Trait Dataset for Diptera Species of Canada and Greenland

Data format: csv, xlsx, tsv

Data format version: xlsx is version 2302. Csv and tsv are UTF-8.

Description: The dataset presents habitat, adult diet, and larval diet for 983 species across 380 genera, 25 subfamilies, and 61 families. A table showing the column labels and descriptions is provided in Table 2. The Diptera species included are from Canada and Greenland and have multiple occurrence records and DNA sequence data on BOLD. The traits were assigned to taxa using a series of literature searches conducted between April 2021 and January 2023. This dataset can be used by researchers to determine biological traits for Diptera species and conduct research that involves ecological trait data.

Additional Information

For this study, we created a dataset containing habitat and adult and larval diet for 983 Diptera species across 61 families found in Canada and Greenland. This dataset allows for assignment of traits to a large variety of species. This is a valuable resource for bioinformatics work, as well as for ecological studies. This dataset was originally used to determine the relationship between biological traits and population genetic structure (Majoros et al., 2023) and can be applied to a wide range of other studies and research. These traits can also be linked to other types of data, such as species occurrence or community composition data.

The species included in this dataset possessed a wide variety of different habitat and dietary requirements. Most species were terrestrial, non-feeding at the adult life stage or saprophagous at the larval stage. Other diets were far less represented, such as those with a parasitic larval stage and those with a fungivore adult diet.

Trait data has been used to investigate various ecological and evolutionary patterns and topics, such as community assembly (Kraft & Ackerly, 2010, Zhang et al., 2020), molecular evolution (May et al., 2020), phylogenetic assemblage structure (Barnagaud et al., 2014), and how populations change over time or in response to ecological change (Coulthard et al., 2019, De Palma et al., 2015). Similarly, this dataset can be used to answer a variety of questions and is a useful tool for future research. It can be used for various studies that relate to ecology, evolution, biogeography, and conservation of fly species and include the habitat and diet of flies. The dataset can be used to investigate the relationship between traits and population genetic structure or phylogenetic community structure. The dataset could also be used to find relationships between traits, such as in studies like Freire et al. (2021). A visualization is provided in Figure 3 of the phylogenetic relationships between species possessing different traits. This could be built upon to look at the relationship between traits and evolutionary history and to investigate

the phylogenetic relationships, such as in studies like Rainford & Mayhew (2015). The trait dataset can also be used to investigate the diversity of different fly traits in an area of interest, to determine how these traits impact occurrence patterns, how traits may relate to responses to environmental change, or to determine which species may be most likely to colonize these areas in the future. Trait data could also prove useful for biomonitoring of insect diversity in agricultural areas, such as research into which farming practices are associated with a balance of ecological functions in associated insect communities.

In this paper we presented a trait dataset containing habitat and diet information for a large number of Diptera taxa from Canada and Greenland that also have occurrence records and DNA barcoding sequences available on BOLD. This dataset will be made publicly available on Data Dryad but is currently available at https://github.com/S-Majoros/Diptera_Dataset_Phylogenetic_Tree. The file is available in csv, xlsx, and tsv format in order to make it as accessible as possible for future users. This dataset is publicly available so that other researchers can use it to conduct further studies, answer more questions, and improve our knowledge of this ecologically important group.

Acknowledgements

The authors acknowledge financial support from the Natural Sciences and Engineering Research Council of Canada; the Government of Canada through Genome Canada and Ontario Genomics; the Ontario Ministry of Economic Development, Job Creation and Trade; and Food from Thought: Agricultural Systems for a Healthy Planet Initiative program funded by the Government of Canada through the Canada First Research Excellence Fund. The authors would like to thank Tyler Elliott for reviewing the manuscript and dataset and providing helpful insight. The authors would also like to thank the researchers from whose work these trait data were gathered, as well as all of the researchers who contribute publicly available data to databases such as BOLD.

Conflict of Interest

The Authors declare that there are no conflicts of interest.

Author Contributions

Samantha E. Majoros – project design, dataset construction, manuscript writing and editing.

Sarah J. Adamowicz – project design, manuscript editing.

Table 1: Overview of some of the databases and datasets that include insects currently publicly available. Number of records are as of Feb 2[nd], 2023.

Database Name	Taxonomic Focus	Geographic Focus	Types of Traits	Number of Traits	Number of Records	Data Availability	References
DISPERSE[1]	Annelida, Mollusca, Platyhelminthes & Arthropoda	Europe	Dispersal	9	480	Download as xlsx file	Sarremejane et al. (2020)
Freshwater Biological Traits[2]	Freshwater macroinvertebrate taxa	North America	Morphology, life history, mobility, environmental tolerance, resource acquisition/ preference	26	11912	Select taxa, region and traits on website. Download as an xlsx file	United States Environmental Protection Agency (2012)
Animal Traits[3]	Arthropoda, Chordata, Mollusca and Annelida	Global	Body mass, metabolic rate, brain size	3	3580	Download xlsx or csv file	Herberstein et al. (2022)
CONUS[4]	Arthropoda	United States of America	Life history, dispersal, morphology, ecology	11	2.05 million	Download csv file	Twardochleb et al. (2021)
Lotic Invertebrate Traits for	Invertebrates	North America	Ecology, morphology, behaviour, physiology	62	14127	Download as a txt file	Vieira et al. (2006)

[1]https://figshare.com/articles/dataset/DISPERSE_a_trait_database_to_assess_the_dispersal_potential_of_European_aquatic_macroinvertebrates/12417251/1

[2] https://cfpub.epa.gov/ncea/global/traits/search.cfm

[3] https://animaltraits.org/

[4] https://portal.edirepository.org/nis/mapbrowse?packageid=edi.481.5

Database Name	Taxonomic Focus	Geographic Focus	Types of Traits	Number of Traits	Number of Records	Data Availability	References
North America[5]							
European & Maghreb Butterfly Trait Database[6]	Lepidoptera	Europe and North Africa	Life history, morphology, resource-based, behaviour	25	542	Download as xlsx file	Middleton-Welling et al. (2020)
GlobalAnts[7]	Formicidae	Global	Morphology, ecology, life history	23	82910	Need to make an account to access. Can search for desired traits and download as a csv file	Parr et al. (2016)
The Odonate Phenotypic Database[8]	Odonata	Global	Morphology, life history, behaviour	35	3978	Download as a xlsx file	Waller et al. (2019)
Morphological trait database of saproxylic beetles[9]	Saproxylic Coleoptera	Europe	Morphology	13	1376	Download as a csv file	Hagge et al. (2021)
Data from: Sensitivity of functional diversity metrics to	Carabidae	The Netherlands	Morphology	7	73	Download as txt file	van der Plas et al. (2017)

[5] https://pubs.er.usgs.gov/publication/ds187
[6] https://butterflytraits.github.io/European-Butterfly-Traits/index.html
[7] http://globalants.org/
[8] http://www.odonatephenotypicdatabase.org/shiny/odonates/?_inputs_&choose_species=%22%22
[9] https://datadryad.org/stash/dataset/doi:10.5061/dryad.2fqz612p3

Database Name	Taxonomic Focus	Geographic Focus	Types of Traits	Number of Traits	Number of Records	Data Availability	References
sampling intensity[10]							
Data from: A summary of eight traits of Coleoptera, Hemiptera, Orthoptera and Araneae, occurring in Grasslands in Germany[11]	Arthropoda	Germany	Morphology, ecology	8	1230	Download as txt file	Gossner et al. (2015)

[10] https://datadryad.org/stash/dataset/doi:10.5061/dryad.1fn46
[11] https://datadryad.org/stash/dataset/doi:10.5061/dryad.53ds2

Table 2: The column labels and descriptions included within the dataset.

Column Label	Description
order	The order level taxonomic classification
family	The family level taxonomic classification
subfamily	The subfamily level taxonomic classification
genus	The genus level taxonomic classification
specificEpithet	The species level taxonomic classification
scientificName	The scientific name of the species
habitat	The habitat in which the taxon is found. Taxa can be assigned terrestrial (taxa that live primarily in land habitats), aquatic (taxa that live primarily in water bodies or associated habitats) or semi-aquatic (species that require both aquatic and terrestrial habitats).
traitName	The name of the biological trait. The traits included are adultDiet and larvalDiet.
traitValue	The trait category assigned to the species. Taxa can be assigned predaceous (taxa who prey on insects or other animals), phytophagous (taxa who feed on plant material), fungivores (those who feed on fungi), saprophagous (those who feed on decaying organic matter), omnivores (those who consume material of both plant and animal origin), nectivore (taxa that primarily feed on nectar), parasites (those who live and feed in or on an organism of another species) or parasitoid (organism whose young develop within a host, eventually resulting in host death). The taxa can also be assigned a combination of these categories. In this case, the categories are separated by "_and_". For example, "nectivore_and_predaceous".
traitAssignmentLevel	The taxonomic level mentioned in the publication from which the traits were obtained and assigned.
basisOfRecord	Where the trait data was obtained from.
referenceNumber	The number of sources referenced when obtaining the trait information for each taxon.
referenceTitle	The title of the reference used for the trait assignment. Full references are included in the manuscript. The number after referenceTitle refers to the number of each reference. For example, if there was a 3 in the referenceNumber column, and this was the 3rd reference, the column would be called referenceTitle3.
referenceDOI	The DOI of the reference used for the trait assignment. Full references are included in the manuscript. The number after referenceDOI refers to the number of each reference. For example, if

Column Label	Description
	there was a 3 in the referenceNumber column, and this was the 3rd reference, the column would be called referenceDOI3

Figure 1: The number of Diptera species from Canada and Greenland of each family that is present in the dataset. The selected species included here also have multiple occurrence records and DNA barcode sequences publicly available on BOLD. For several small families, only one species is included. Chironomidae is the largest family within the dataset, with 182 species present.

Figure 2: The number of Diptera species from Canada and Greenland in this dataset having each trait category for: a) habitat, b) adult diet, and c) larval diet. The majority of species included in the dataset are terrestrial. Most species are non-feeding at the adult life stage, and few are omnivores or phytophagous and fungivores as adults. The majority of species included in the dataset are saprophagous as larva, and few are parasitic.

a) Habitat

Habitat
- Terrestrial
- Aquatic
- Semi-Aquatic

b) Adult Diet

Adult Diet
- Saprophagous
- Nectivore
- Fungivore
- Predaceous
- Phytophagous
- Non-Feeding
- Nectivore and Predaceous
- Nectivore and Saprophagous
- Fungivore and Nectivores
- Phytophagous and Fungivore

c) Larval Diet

Larval Diet
- Phytophagous
- Predaceous
- Saprophagous
- Fungivore
- Omnivores
- Parasitoid
- Parasitic
- Phytophagous and Saprophagous
- Phytophagous and Fungivore

Figure 3: Phylogenetic trees of Diptera species from Canada and Greenland included in this dataset; these species also have multiple occurrence records and DNA barcode sequences publicly available on BOLD. The different colours of the tips represent species with different: a) habitat categories, b) adult diet categories, and c) larval diet categories. The traits provided in the dataset could be used to investigate the relationship between traits and evolutionary history. These maximum likelihood trees were created using cytochrome c oxidase subunit 1 (CO1) sequences from BOLD and functions from the package phangorn version 2.11.1(Schliep, 2011) in the R programming language (R Core Team, 2023). The tips were coloured using the package ggtree version 3.6.2 (Yu et al., 2017). The code for how to create this tree is provided at https://github.com/S-Majoros/Diptera_Dataset_Phylogenetic_Tree

References

Adler, P.H. & Courtney, G.W. (2019). Ecological and societal services of aquatic Diptera. *Insects, 10*(3), 70. doi: 10.3390/insects10030070

Barnagaud, J., Kissling, W.D., Sandel, B., Eiserhardt, W.L., Şekercioğlu, C.H., Enquist, B.J., Tsirogiannis, C. & Svenning, J. (2014). Ecological traits influence the phylogenetic structure of bird species co-occurrences worldwide. *Ecology Letters, 17*(7), 811-820. doi: 10.1111/ele.12285

Bedini, S., Flamini, G., Cosci, F., Ascrizzi, R., Echeverria, M.C., Guidi, L., Landi, M., Lucchi, A. & Conti, B. (2017). *Artemisia* spp. essential oils against the disease-carrying blowfly *Calliphora vomitoria. Parasites & Vectors, 10*(1), 80. doi: 10.1186/s13071-017-2006-y

Bragard, C., Dehnen-Schmutz, K., Di Serio, F., Gonthier, P., Jacques, M., Jaques Miret, J..A., Justesen, A.F., Magnusson, C.S., Milonas, P., Navas-Cortes, J.A., Parnell, S., Potting, R., Reignault, P.L., Thulke, H., Van der Werf, W., Vicent Civera, A., Yuen, J., Zappalà, L., Czwienczek, E., Streissl, F. & MacLeod, A. (2020). Pest categorisation of *Liriomyza bryoniae. EFSA Journal, 18* (3), p.e06038-n/a. doi: 10.2903/j.efsa.2020.6038

Coulthard, E., Norrey, J., Shortall, C. & Harris, W.E. (2019). Ecological traits predict population changes in moths. *Biological Conservation, 233,* 213-219. doi: 10.1016/j.biocon.2019.02.023

De Palma, A., Kuhlmann, M., Roberts, S.P.M., Potts, S.G., Börger, L., Hudson, L.N., Lysenko, I., Newbold, T. & Purvis, A. Ecological traits affect the sensitivity of bees to land-use pressures in European agricultural landscapes. *Journal of Applied Ecology, 52*(6), 1567-1577. doi: 10.1111/1365-2664.12524

deWaard, J.R., Levesque-Beaudin, V., deWaard, S.L., Ivanova, N.V., McKeown, J.T.A., Miskie, R., Naik, S., Perez, K.H.J., Ratnasingham, S., Sobel, C.N., Sones, J.E., Steinke, C., Telfer, A.C., Young, A.D., Young, M.R., Zakharov, E.V. & Hebert, P.D.N. (2018). Expedited assessment of terrestrial arthropod diversity by coupling Malaise traps with DNA barcoding. *Genome, 62*(3), 85-95. doi: 10.1139/gen-2018-0093

Freire, Jr., G.B., Silva, T., Oliveira, H., Collier, C., Rodrigues, H.P., Dias, J.P., Santos, J.P., Marini-Filho, O.J., Freitas, A.V.L., Smilanich, A.M., Dyer, L.A. & Diniz, I.R. (2021). Good Things Come in Larger Packages: Size Matters for Adult Fruit-Feeding Butterfly Dispersal and Larval Diet Breadth. *Diversity, 13*(12), 664. doi: 10.3390/d13120664

GBIF.org (2020). *GBIF Home Page.* Avaliable from https://www.gbif.org. [13 February 2023]

Geiger, M.F., Moriniere, J., Hausmann, A., Haszprunar, G., Wägele, W., Hebert, P.D.N. & Rulik, B. (2016). Testing the Global Malaise Trap Pogram-How well does the current barcode reference library identify flying insects in Germany? *Biodiversity Data Journal, 4*(4), e10671. doi: 10.3897/BDJ.4.e10671

Global Malaise Trap Program (n.d.). *Global Malaise Trap Program.* Retrieved from http://globalmalaise.org/homecopy/

Gossner, M.M., Simons, N.K., Achtziger, R., Blick, T., Dorow, W.H.O., Dziock, F., Köhler, F., Rabitsch, W. & Weisser, W.W. (2015). A summary of eight traits of Coleoptera, Hemiptera, Orthoptera and

Araneae, occurring in grasslands in Germany. *Scientific Data, 2*(1), 150013. doi: 10.1038/sdata.2015.13

Hagge, J., Müller, J., Birkemoe, T., Buse, J., Christensen, R.H.B., Gossner, M.M., Gruppe, A., Heibl, C., Jarzabek-Müller, A., Seibold, S., Siitonen, J. Soutinho, J.G., Sverdrup-Thygeson, A., Thorn, S. & Drag, L. (2021). What does a threatened saproxylic beetle look like? Modelling extinction risk using a new morphological trait database. *Journal of Animal Ecology, 90*(8), 1934-1947. doi: 10.1111/1365-2656.13512

Hebert, P.D.N., Cywinska, A., Ball, S.L. & DeWaard, J.R. (2003). Biological identifications through DNA barcodes. *Proceedings of the Royal Society B: Biological Sciences, 270*(1512), 313– 21. doi: 10.1098/rspb.2002.2218

Herberstein, M.E., McLean, D.J., Lowe, E., Wolff, J.O., Khan, M.K., Smith, K., Allen, A.P., Bulbert, M., Buzatto, B.A., Eldridge, M.D.B., Falster, D., Winzer, L.F., Griffith, S.C., Madin, J.S., Narendra, A., Westoby, M., Whiting, M.J., Wright, I.J. & Carthey, A.J.R. (2022). AnimalTraits – a curated animal trait database for body mass, metabolic rate and brain size. *Scientific Data, 9*(1), 265. doi: 10.1038/s41597-022-01364-9

Kraft, N.J.B. & Ackerly, D.D. (2010). Functional trait and phylogenetic tests of community assembly across spatial scales in an Amazonian forest. *Ecological Monographs, 80*(3), 401-422. doi: 10.1890/09-1672.1

Majoros, S.E., Adamowicz, S.J. & Cottenie, K. (2023). Novel pipeline for large-scale comparative population genetics. bioRxiv. doi: 10.1101/2023.01.23.524574

Marshall, S.A. (2012). *Flies. The natural history and diversity of Diptera.* Firefly Books Ltd.

May, J.A., Feng, Z., Orton, M.G. & Adamowicz, S.J. (2020). The effects of ecological traits on the rate of molecular evolution in ray-finned fishes: a multivariable approach. *Journal of Molecular Evolution, 88*(8-9), 689-702. doi: 10.1007/s00239-020-09967-9

Mesmin, X., Vincent, M., Tricault, Y., Estorgues, V., Daniel, L., Cortesero, A., Faloya, V. & Le Ralec, A. (2019). Assessing the relationship between pest density and plant damage: a case study with the belowground herbivore *Delia radicum* (Diptera: Anthomyiidae) on broccoli. *Applied Entomology and Zoology, 54*(2), 155-165. doi: 10.1007/s13355-019-00607-3

Middleton-Welling, J., Dapporto, L., García-Barros, E., Wiemers, M., Nowicki, P., Plazio, E., Bonelli, S., Zaccagno, M., Šašić, M., Liparova, J., Schweiger, O., Harpke, A., Musche, M., Settele, J., Schmucki, R. & Shreeve, T. (2020). A new comprehensive trait database of European and Maghreb butterflies, Papilionoidea. *Scientific Data*, *7*(1), 351. doi: 10.1038/s41597-020-00697-7

Molinatto, G., Demichelis, S., Bodino, N., Giorgini, M., Mori, N. & Bosco, D. (2020). Biology and prevalence in Northern Italy of *Verrallia aucta* (Diptera, Pipunculidae), a parasitoid of *Philaenus spumarius* (Hemiptera, Aphrophoridae), the main vector of *Xylella fastidiosa* in Europe. *Insects, 11*(9), 607. doi: 10.3390/insects11090607

Olafson, PU., Aksoy, S., Attardo, G.M., Buckmeier, G., Chen, X., Coates, C.J., Davis, M., Dykema, J., Emrich, S.J., Friedrich, M., Holmes, C.J., Ioannidis, P., Jansen, E.N., Jennings, E.C., Lawson, D.,

Martinson, E.O., Maslen, G.L., Meisel, R.P., Murphy, T.D., Nayduch, D., Nelson, D.R., Oyen, K.J., Raszick, T.J., Ribeiro, J.M.C., Robertson, H.M., Rosendale, A.J., Sackton, T.B., Saelao, P., Swiger, S.L., Sze, S., Tarone, A.M., Taylor, D.B., Warren, W.C., Waterhouse, R.M., Weirauch, M.T., Werren, J.H., Wilson, R.K., Zdobnov, E.M. & Benoit, J.B. (2021). The genome of the stable fly, *Stomoxys calcitrans*, reveals potential mechanisms underlying reproduction, host interactions, and novel targets for pest control. *BMC Biology, 19*(1), 41. doi: 10.1186/s12915-021-00975-9

Parr, C.L., Dunn, R.R., Sanders, N.J., Weiser, M.D., Photakis, M., Bishop, T.R., Fitzpatrick, M.C., Arnan, X., Baccaro, F., Brandão, C.R.F., Chick, L., Donoso, D.A., Fayle, T.M., Gómez, C., Grossman, B., Munyai, T.C., Pacheco, R., Retana, J., Robinson, A., Sagata, K., Silva, R.R., Tista, M., Vasconcelos, H., Yates, M. & Gibb, H. (2016). GlobalAnts: a new database on the geography of ant traits (Hymenoptera: Formicidae). *Insect Conservation and Diversity, 10*(1), 5-20. doi: 10.1111/icad.12211

R Core Team. (2022). *R: A language and environment for statistical computing.* R Foundation for Statistical Computing, Vienna, Austria. URL http://www.R-project.org/

Rainford, J.L. & Mayhew, P.J. (2015). Diet evolution and clade richness in Hexapoda: a phylogenetic study of higher taxa. *The American Naturalist, 186*(6), 771-791. doi: 10.1086/683461

Ratnasingham, S. & Hebert, P.D.N. (2007). Bold: the barcode of life data system. *Molecular Ecology Notes, 7*(3), 355-364. doi:10.1111/j.1471-8286.2007.01678.

Ratnasingham, S. & Hebert, P.D.N. (2013). A DNA-based registry for all animal species: The Barcode Index Number (BIN) system. *PLoS ONE, 8*(7), e66213.

Sarremejane, R., Cid, N., Stubbington, R., Datry, T., Alp, M., Cañedo-Argüelles, M., Cordero-Rivera, A. Csabai, Z., Gutiérrez-Cánovas, C., Heino, J., Forcellini, M., Millán, A., Paillex, A., Pařil, P., Polášek, M., Tierno de Figueroa, J.M., Usseglio-Polatera, P., Zamora-Muñoz, C. & Bonada, N. (2020) DISPERSE, a trait database to assess the dispersal potential of European aquatic macroinvertebrates. *Scientific Data, 7*(1), 386. doi: 10.1038/s41597-020-00732-7

Schliep K.P. (2011). phangorn: phylogenetic analysis in R. *Bioinformatics, 27*(4), 592-593. doi: 10.1093/bioinformatics/btq706

Shamshad, A. (2010). The development of integrated pest management for the control of mushroom sciarid flies, *Lycoriella ingenua* (Dufour) and *Bradysia ocellaris* (Comstock), in cultivated mushrooms. *Pest Management Science, 66*(10), 1063-1074. doi: 10.1002/ps.1987

Tiusanen, M., Hebert, P.D.N., Schmidt, N.M. & Roslin, T. (2016). One fly to rule them all – muscid flies are the key pollinators in the Arctic, *Proceedings of the Royal Society: Biological Sciences, 283*(1839), 20161271. doi: 10.1098/rspb.2016.1271

Twardochleb, L., Hiltner, E., Pyne, M. & Zarnetske, P. (2021). Freshwater insects CONUS: A database of freshwater insect occurrences and traits for the contiguous United States. *Global Ecology and Biogeography, 30*(4), 826-841. doi: 10.1111/geb.13257

United States Environmental Protection Agency (2012). Freshwater Biological Trait Database. (Report No. EPA/600/R-11/038F). Retrieved from https://cfpub.epa.gov/ncea/global/recordisplay.cfm?deid=241813

van der Plas, F., van Klink, R., Manning, P., Olff, H. & Fischer, M. (2017). Sensitivity of functional diversity metrics to sampling intensity. *Methods in Ecology and Evolution, 8*(9), 1072-1080. doi: 10.1111/2041-210X.12728

Vieira, N.K.M., Poff, N.L., Carlisle, D.M., Moulton II, S.R., Koski, M.L. & Kondratieff, B.C. (2006). A database of lotic invertebrate traits for North America. *United States Geographical Survey.* https://pubs.usgs.gov/ds/ds187/pdf/ds187.pdf

Waller, J.T., Willink, B., Tschol, M. & Svensson, E.I. (2019). The odonate phenotypic database, a new open data resource for comparative studies of an old insect order. *Scientific Data*, *6*(1), 316. doi: 10.1038/s41597-019-0318-9

Wearing, C. H., Marshall, R. R., Attfield, B. & Colhoun, C. (2013). Phenology and distribution of the apple leafcurling midge (*Dasineura mali* (Kieffer)) (Diptera: Cecidomyiidae) and its natural enemies on apples under biological and integrated pest management in Central Otago, New Zealand. *New Zealand Entomologist, 36*(2), 87-106. doi: 10.1080/00779962.2012.712887

Yu G., Smith, D., Zhu, H., Guan, Y. & Lam, T.T. (2017). ggtree: an R package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods in Ecology and Evolution, 8*(1), 28-36. doi: 10.1111/2041-210X.12628

Zhang, J., Swenson, N.G., Liu, J., Liu, M., Qiao, X. & Jiang, M. (2020). A phylogenetic and trait-based analysis of community assembly in a subtropical forest in central China. *Ecology and Evolution, 10*(15), 8091-8104. doi: 10.1002/ece3.6465

APPENDICES

Appendix A: References for Diptera habitat and diet traits

Acartophthalmidae

Roháček, J. (2012). The fauna of the opomyzoid families Clusiidae, Acartophthalmidae, Anthomyzidae, Opomyzidae, Stenomicridae, Periscelididae, Asteiidae (Diptera) in the Gemer area (Central Slovakia). *Časopis Slezského Zemského Muzea, 61*(2), 97-111. doi: 10.2478/v10210-012-0011-5

Roháček, J. (2013). The fauna of the Acalyptrate families Micropezidae, Psilidae, Clusiidae, Acartophthalmidae, Anthomyzidae, Aulacigastridae, Periscelididae and Asteiidae (Diptera) in the Gemer area (Central Slovakia): supplement 1. C*asopis Slezskeho Zemskeho Muzea, 62*(2), 125-136. doi: 10.2478/cszma-2013-0014

Agromyzidae

Ayabe, Y., Ueno, T. & Harvey, J.A. (2012). Complex feeding tracks of the sessile herbivorous insect *Ophiomyia maura* as a function of the defense against insect parasitoids. *PLoS one, 7*(2), p.e32594-e32594. doi: 10.1371/journal.pone.0032594

Bañuelos, M. & Kollmann, J. (2011). Effects of host-plant population size and plant sex on a specialist leaf-miner. *Acta Oecologica, 37*(2), 58-64. doi: 10.1016/j.actao.2010.11.007

Bragard, C., Dehnen-Schmutz, K., Di Serio, F., Gonthier, P., Jacques, M., Jaques Miret, J..A., Justesen, A.F., Magnusson, C.S., Milonas, P., Navas-Cortes, J.A., Parnell, S., Potting, R., Reignault, P.L., Thulke, H., Van der Werf, W., Vicent Civera, A., Yuen, J., Zappalà, L., Czwienczek, E., Streissl, F. & MacLeod, A. (2020). Pest categorisation of *Liriomyza bryoniae. EFSA Journal, 18* (3), p.e06038-n/a. doi: 10.2903/j.efsa.2020.6038

Carapelli, A., Soltani, A., Leo, C., Vitale, M., Amri, M. & Mediouni-Ben Jemâa, J. (2018). Cryptic Diversity Hidden within the Leafminer Genus *Liriomyza* (Diptera: Agromyzidae). *Genes, 9*(11), 554. doi: 10.3390/genes9110554

Eber, S. (2004). Bottom-up density regulation in the holly leaf-miner *Phytomyza ilicis. The Journal of Animal Ecology, 73*(5), 948-958. doi: 10.1111/j.0021-8790.2004.00867.x

Gratton, C. & Welter, S.C. (1998). Oviposition preference and larval performance of *Liriomyza helianthi* (Diptera: Agromyzidae) on normal and novel host plants. *Environmental Entomology, 27*(4), 926-935. doi: 10.1093/ee/27.4.926

Hébert, J.B., Scheffer, S.J. & Hawthorne, D.J. (2016). Evidence for ecological speciation via a host shift in the holly leaf miner, *Phytomyza glabricola* (Diptera: Agromyzidae). *Ecology and Evolution, 6*(18), 6565-6577. doi: 10.1002/ece3.2358

Lambkin, C.L., Fayed, S.A., Manchester, C., La Salle, J., Scheffer, S.J. & Yeates, D.K. (2008). Plant hosts and parasitoid associations of leaf mining flies (Diptera: Agromyzidae) in the Canberra region of Australia. *Australian Journal of Entomology, 47*(1), 13-19. doi: 10.1111/j.1440-6055.2007.00622.x

Lonsdale, O. (2015). A new afrotropical species of *Phytomyza* in the subgenus *Ptochomyza* (Diptera: Agromyzidae). *African Invertebrates, 56*(3), 637-643. doi: 10.5733/afin.056.0309

Lonsdale, O. (2021). Manual of North American Agromyzidae (Diptera, Schizophora), with revision of the fauna of the "Delmarva" states. *ZooKeys, 1051*, 1-481. doi: 10.3897/zookeys.1051.64603

Marino, P.C. & Cornell, H.V. (1992). Adult movement of the native holly leafminer, *Phytomyza ilicicola* Loew (Diptera: Agromyzidae): Consequences for host choice within and between habitats. *Oecologia, 92*(1), 76-82. doi: 10.1007/BF00317265

Marino, P.C., Cornell, H.V. & Kahn, D.H. (1993). Environmental and clonal influences on host choice and larval survival in a leafmining insect. *Journal of Animal Ecology, 62*(3), 503-510. doi: 10.2307/5199

Masetti, A., Lanzoni, A. & Burgio, G. (2010). Effects of flowering plants on parasitism of lettuce leafminers (Diptera: Agromyzidae). *Biological Control, 54*(3), 263-269. doi: 10.1016/j.biocontrol.2010.05.016

Nzama, S., Olckers, T. & Zachariades, C. (2014). Seasonal activity, habitat preferences and larval mortality of the leaf-mining fly *Calycomyza eupatorivora* (Agromyzidae), a biological control agent established on *Chromolaena odorata* (Asteraceae) in South Africa. *Biocontrol Science and Technology, 24*(11), 1297-1307. Doi: 10.1080/09583157.2014.935293

Okoth, C.A., Deng, A.L., Tabu, I.M., Akutse, K.S. & Fiaboe, K.K.M. (2014). Effect of host plant on feeding, biological and morphological parameters of *Liriomyza huidobrensis* Blanchard (Diptera: Agromyzidae). *African Entomology, 22*(3), 577-588. doi: 10.4001/003.022.0315

Piwowarczyk, R., Mielczarek, L. & Guzikowski, S. (2018). First report of *Phytomyza orobanchia* (Diptera: Agromyzidae) from Poland and *Chymomyza amoena* (Diptera: Drosophilidae) on *Phelipanche ramose* (Orobanchaceae). *Florida Entomologist, 101*(3), 540-542. doi: 10.1653/024.101.0333

Roíguez-Castañeda, G., MacVean, C., Cardona, C. & Hof, A.R. (2017). What limits the distribution of *Liriomyza huidobrensis* and its congener *Liriomyza sativae* in their native niche: when temperature and competition affect species' distribution range in Guatemala. *Journal of Insect Science, 17*(4). doi: 10.1093/jisesa/iex059

Scheffer, S.J. & Hawthorne, D.J. (2007). Molecular evidence of host-associated genetic divergence in the holly leafminer *Phytomyza glabricola* (Diptera: Agromyzidae): apparent discordance among marker systems. *Molecular Ecology, 16*(13), 2627-2637. doi: 10.1111/j.1365-294X.2007.03303.x

Scheffer, S.J., Lewis, M.L., Hébert, J.B. & Jacobsen, F. (2020). Diversity and host plant-use North American *Phytomyza* holly leafminers (Diptera: Agromyzidae): colonization, divergence, and specificity

in a host-associated radiation. *Annals of the Entomological Society of America, 114*(1), 56-69. doi: 10.1093/aesa/saaa03

Scheffer, S.J. & Lonsdale, O. (2011). *Phytomyza omlandi* spec. nov. – The first species of Agromyzidae (Diptera: Schizophora) reared from the family Gelsemiaceae (Asteridae). *Proceedings of the Entomological Society of Washington, 113*(1), 42-49. doi: 10.4289/0013-8797.113.1.42

Scheffer, S.J. & Wiegmann, B.M. (2000). Molecular phylogenetics of the holly leafminers (Diptera: Agromyzidae: *Phytomyza*): species limits, speciation, and dietary specialization. *Molecular Phylogenetics and Evolution, 17*(2), 244-255. doi: 10.1006/mpev.2000.0830

Soltani, A., Ben Abda, M., Amri, M., Carapelli, A. & Mediouni Ben Jemâa, J. (2020). Seasonal incidence of the leaf miner *Liriomyza cicerina* Rond (Diptera: Agromyzidae) in chickpea fields and effects of climatic parameters, chickpea variety, and planting date on the leaf miner infestation rate. *Euro-Mediterranean journal for environmental integration, 5*(3). doi: 10.1007/s41207-020-00198-4

Williams, E. C. & Macdonald, O. C. (1995). Critical factors required by the nematode *Steinernema feltiae* for the control of the leafminers *Liriomyza huidobrensis*, *Liriomyza bryoniae* and *Chromatomyia syngenesiae*. *Annals of Applied Biology, 127*(2), 329-341. doi: 10.1111/j.1744-7348.1995.tb06677.x

Anisopodidae

Kolcsár, L., Dvořák, L. & Beuk, P.L. (2016). New records of *Sylvicola* (Diptera: Anisopodidae) from Romania. *Biodiversity Data Journal, 4*(4), p.e7861-e7861. doi: 10.3897/BDJ.4.e7861.

Ševčík, J. (2011). Anisopodidae and Dixidae (Diptera) of the Gemer region, with a new record for Slovakia. *Časopis Slezského Zemského Muzea, 60*(2), 181-184. doi: 10.2478/v10210-011-0019-2

Soli, G. & Rindal, E. (2014). The genus Sylvicola Harris, 1780 (Diptera, Anisopodidae) in Norway – with a key to the North European species. *Norwegian Journal of Entomology, 61*(2), 190-200.

Anthomyiidae

Bažok, R., Ceranić-Sertić, M., Barčić, J.I., Borošić, J., Kozina, A., Kos, T., Lemić, D. & Čačija, M. (2012). Seasonal flight, optimal timing and efficacy of selected insecticides for cabbage maggot (*Delia radicum* L., Diptera: Anthomyiidae) control. *Insects, 3*(4), 1001-1027. doi: 10.3390/insects3041001

Deasy, W., Shepherd, T., Alexander, C.J., Birch, A.N.E. & Evans, K.A. (2016). Field-based evaluation of a novel SPME-GC-MS method for investigation of below-ground interaction between brassica roots and larvae of cabbage root fly, *Delia radicum* L. *Phytochemical Analysis, 27*(6), 343-353. doi: 10.1002/pca.2634

Dreves, A.J., Dalthorp, D., Stone, A.G. & Fisher, G. (2006). Spring emergence and seasonal flight of *Delia radicum* L. (Diptera: Anthomyiidae) in western Oregon. *Environmental Entomology, 35*(2), 465-477. doi: 10.1603/0046-225X-35.2.465

Gouinguene, S.P & Stadler, E. (2006). Oviposition in *Delia platura* (Diptera, Anthomyiidae): The role of volatile and contact cues of bean. *Journal of Chemical Ecology, 32*(7), 1399-1413. doi: 10.1007/s10886-006-9058-3

Guerra, P.C., Keil, C.B., Stevenson, P.C., Mina, D., Samaniego, S., Peralta, E., Mazon, N. & Chancellor, T.C.B. (2016). Larval performance and adult attraction of *Delia platura* (Diptera: Anthomyiidae) in a native and an introduced crop. *Journal of Economic Entomology, 110*(1), 186-191. doi: 10.1093/jee/tow237

Kim, H.S., Cho, J.R., Kim, J.J., Lee, M. & Byun, M.W. (2001). Optimal radiation dose of Cobalt60 to improve the sterile insect technique for *Delia antiqua* and *Delia platura. Journal of Asia-Pacific Entomology, 4*(1), 11-16. doi: 10.1016/S1226-8615(08)60095-3

Lamb, R.J. & Boivin, G. (2018). Population variability of three Delia species (Diptera: Anthomyiidae) from the same agricultural habitat in Quebec, Canada. *Canadian Entomologist, 150*(1), 80-96. doi: 10.4039/tce.2017.52

Leuchtmann, A. (2007). Botanophila flies on Epichloë host species in Europe and North America: no evidence for co-evolution. *Entomologia Experimentalis et Applicata, 123*(1), 13-23. doi: 10.1111/j.1570-7458.2006.00518.x
Lukwinski, A.T., Hill, J.E., Khachatourians, G.G., Hemmingsen, S.M. & Hegedus, D.D. (2006). Biochemical and taxonomic characterization of bacteria associated with the crucifer root maggot (*Delia radicum*). *Canadian Journal of Microbiology, 52*(3), 197-208. doi: 10.1139/w05-123

Mesmin, X., Vincent, M., Tricault, Y., Estorgues, V., Daniel, L., Cortesero, A., Faloya, V. & Le Ralec, A. (2019). Assessing the relationship between pest density and plant damage: a case study with the belowground herbivore *Delia radicum* (Diptera: Anthomyiidae) on broccoli. *Applied Entomology and Zoology, 54*(2), 155-165. doi: 10.1007/s13355-019-00607-3

Mlynarek, J.J., MacDonald, M., Sim, K., Hiltz, K., McDonald, M.R. & Blatt, S. (2020). Oviposition, feeding preferences and distribution of *Delia* species (Diptera: Anthomyiidae) in Eastern Canadian onions. *Insects, 11*(11), 780. doi: 10.3390/insects11110780

Nieto, D.J., Letourneau, D.K., Toyama, L., Slay, C. & Kula, R.R. (2021). Surveying populations of *Delia radicum* (Diptera: Anthomyiidae) and its associated natural enemies in organic cauliflower in California. *Biocontrol Science and Technology, 31*(1), 112-118. doi: 10.1080/09583157.2020.1845609

Nilsson, U., Rännbäck, L.M., Anderson, P. & Rämert, B. (1986). Herbivore response to habitat manipulation with floral resources: a study of the cabbage root fly. *Journal of Applied* Entomology, 136(7), 481-489. doi: 10.1111/j.1439-0418.2011.01685.x

Ohta, S., Yousuke S., Koichiro T., Yukio I. & Takashi M. (2015). Comprehensive identification of odorant-binding protein genes in the seed fly, *Delia platura* (Diptera: Anthomyiidae). *Applied Entomology and Zoology, 50*(4), 457-463. doi: 10.1007/s13355-015-0353-8

Rao, S. & Baumann, D. (2004). The interaction of a Botanophila fly species with an exotic Epichloë fungus in a cultivated grass: fungivore or mutualist? *Entomologia Experimentalis et Applicata, 112*(2), 99-105. doi: 10.1111/j.0013-8703.2004.00189.x

Rao, S., Alderman, S.C., Takeyasu, J. & Matson, B. (2005). Botanophila-Epichloe association in cultivated Festuca in Oregon: evidence of simple fungivory. *Entomologia Experimentalis et Applicata, 115*(3), 427-433. doi: 10.1111/j.1570-7458.2005.00309.x

Saumure, R.A., Walde, A.D. & Wheeler, T.A. (2006). Nonpredatory fly larvae (*Delia platura*: Anthomyiidae) in a nest of a Northern map turtle (*Graptemys geographica*). *Chelonian Conservation and Biology, 5*(2), 274-275. doi: 10.2744/10718443(2006)5[274:NFLDPA]2.0.CO2

Silver, N., Hillier, K. & Blatt, S. (2018). Management of Delia (Diptera: Anthomyiidae) through selectively timed planting of *Phaseolus vulgaris* (Fabaceae) in Atlantic Canada. *Canadian Entomologist, 150*(5), 663-674. doi: 10.4039/tce.2018.36

Sontowski, R., Poeschl, Y., Okamura, Y., Vogel, H., Guyomar, C., Cortesero, A. & Dam, N.M. (2022). A high-quality functional genome assembly *of Delia radicum* L. (Diptera: Anthomyiidae) annotated from egg to adult. *Molecular Ecology Resources, 22*(5), 1954-1971. doi: 10.1111/1755-0998.13594

Van der Heyden, H., Fortier, A. & Savage, J. (2020). A HRM assay for rapid identification of members of the seedcorn maggot complex (*Delia florilega* and *D. platura*) (Diptera: Anthomyiidae) and evidence of variation in temporal patterns of larval occurrence. *Journal of Economic Entomology, 113*(6), 2920-2930. doi: 10.1093/jee/toaa230

Anthomyzidae

RohaAek, J., Petrakova, L. & Tothova, A. (2019). Molecular phylogeny and timing of evolution of *Anthomyza* and related genera (Diptera: Anthomyzidae). *Zoologica Scripta, 48*(6), 745-760. doi: 10.1111/zsc.12373

Roháček, J. (2012). The fauna of the opomyzoid families Clusiidae, Acartophthalmidae, Anthomyzidae, Opomyzidae, Stenomicridae, Periscelididae, Asteiidae (Diptera) in the Gemer area (Central Slovakia). *Časopis Slezského Zemského Muzea, 61*(2), 97-111. doi: 10.2478/v10210-012-0011-5

Asilidae

Dennis, D.S., Lavigne, R.J. & Dennis, J.G. (2012). Spiders (Araneae) as prey of robber flies (Diptera: Asilidae). *Journal of Entomological Research Society, 14*(1), 65-76.

Pollock, D.A. (2021). Robber flies (Diptera: Asilidae) as predators of harvester ant workers (Hymenoptera: Formicidae: Pogonomyrmex) in Eastern New Mexico and West Texas. *The Southwestern Naturalist, 65*(1), 19-27. doi: 10.1894/0038-4909-65.1.3

Asteiidae

Roháček, J. (2012). The fauna of the opomyzoid families Clusiidae, Acartophthalmidae, Anthomyzidae, Opomyzidae, Stenomicridae, Periscelididae, Asteiidae (Diptera) in the Gemer area (Central Slovakia). *Časopis Slezského Zemského Muzea, 61*(2), 97-111. doi: 10.2478/v10210-012-0011-5

Aulacigastridae

Roháček, J. (2013). The fauna of the Acalyptrate families Micropezidae, Psilidae, Clusiidae, Acartophthalmidae, Anthomyzidae, Aulacigastridae, Periscelididae and Asteiidae (Diptera) in the Gemer area (Central Slovakia): supplement 1. C*asopis Slezskeho Zemskeho Muzea, 62*(2), 125-136. doi: 10.2478/cszma-2013-0014

Bibionidae

Frouz, J., Lin, Q., Li, X., Abakumov, E., Brune, A. & Šustr, V. (2020). Utilization of dietary protein in the litter-dwelling larva of *Bibio marci* (Diptera: Bibionidae). *Eurasian Soil Science, 52*(12), 1583-1587. doi: 10.1134/S1064229319120032

Šustr, V., Stingl, U. & Brune, A. (2014). Microprofiles of oxygen, redox potential, and pH, and microbial fermentation products in the highly alkaline gut of the saprophagous larva of *Penthetria holosericea* (Diptera: Bibionidae). *Journal of Insect Physiology, 67*, 64-69. doi: 10.1016/j.jinsphys.2014.06.007

Calliphoridae

Andere, A.A., Platt, R.N., Ray, D.A. & Picard, C.J. (2016). Genome sequence of *Phormia regina* Meigen (Diptera: Calliphoridae): implications for medical, veterinary and forensic research. *BMC Genomics, 17*(1), 842. doi: 10.1186/s12864-016-3187-z

Bedini, S., Flamini, G., Cosci, F., Ascrizzi, R., Echeverria, M.C., Guidi, L., Landi, M., Lucchi, A. & Conti, B. (2017). Artemisia spp. essential oils against the disease-carrying blowfly *Calliphora vomitoria. Parasites & Vectors, 10*(1), 80. doi: 10.1186/s13071-017-2006-y

Bennett, G.F. & Whitworth, T.L. (1991). Studies on the life history of some species of *Protocalliphora* (Diptera: Calliphoridae). *Canadian Journal of Zoology, 69*(8), 2048-2058. doi: 10.1139/z91-286

Berrigan, D. & Lighton, J.R.B. (1994). Energetics of pedestrian locomotion in adult male blowflies, *Protophormia terraenovae* (Diptera: Calliphoridae). *Physiological Zoology, 67*(5), 1140-1153. doi: 10.1086/physzool.67.5.30163886

Brodie, B.S., Wong, W.H.L., VanLaerhoven, S. & Gries, G. (2015). Is aggregated oviposition by the blow flies *Lucilia sericata* and *Phormia regina* (Diptera: Calliphoridae) really pheromone-mediated? *Insect Science, 22*(5), 651-660. doi: 10.1111/1744-7917.12160

Burgess, I.F. (1987). Opportunistic myiasis of nestling common blackbirds by *Lucilia silvarum. Parasitology Research, 121*(2), 763-764. doi: 10.1007/s00436-021-07402-6

Dufek, M.I., Oscherov, E.B., Damborsky, M.P. & Mulieri, P.R. (2019). Calliphoridae (Diptera) in human-transformed and wild habitats: diversity and seasonal fluctuations in the humid Chaco Ecoregion of South America. *Journal of Medical Entomology, 56*(3), 725-736. doi: 10.1093/jme/tjy234

Dykstra, C.R., Hays, J.L., Simon, M.M. & Wegman, A.R. (2012). *Protocalliphora* (Diptera: Calliphoridae) infestations of nestling red-shouldered hawks in Southern Ohio. *The Wilson Journal of Ornithology, 124*(4), 783-787. doi: 10.1676/1559-4491-124.4.783

Garrido-Bautista, J., Moreno-Rueda, G., Baz, A., Canal, D., Camacho, C., Cifrián, B., Nieves-Aldrey, J.L., Carles-Tolrá, M. & Potti, J. (1987). Variation in parasitoidism of *Protocalliphora azurea* (Diptera: Calliphoridae) by *Nasonia vitripennis* (Hymenoptera: Pteromalidae) in Spain. *Parasitology Research, 119*(2), 559-566. doi: 10.1007/s00436-019-06553-x

Gentes, M., Whitworth, T.L., Waldner, C., Fenton, H. & Smits, J.E. (2007). Tree swallows (*Tachycineta bicolor*) nesting on wetlands impacted by oil sands mining are highly parasitized by the bird blow fly *Protocalliphora* sp. *Journal of Wildlife Diseases, 43*(2), 167-178. doi: 10.7589/0090-3558-43.2.167

Gunn, A. & Bird, J. (2010). The ability of the blowflies *Calliphora vomitoria* (Linnaeus), *Calliphora vicina* (Rob-Desvoidy) and *Lucilia sericata* (Meigen) (Diptera: Calliphoridae) and the muscid flies *Muscina stabulans* (Fallén) and *Muscina prolapsa* (Harris) (Diptera: Muscidae) to colonise buried remains. *Forensic Science International, 207*(1), 198-204. doi: 10.1016/j.forsciint.2010.10.008

Johnson, L.S., Eastman, M.D. & Kermott, L.H. (1991). Effect of ectoparasitism by larvae of the blow fly *Protocalliphora parorum* (Diptera: Calliphoridae) on nestling house wrens, *Troglodytes aedon. Canadian Journal of Zoology, 69*(6), 1441-1446. doi: 10.1139/z91-204

Kelly, M.A., Usher, M.J., Ujvari, B., Madsen, T., Wallman, J.F., Buttemer, W.A. & Hulbert, A.J. (2014). Diet fatty acid profile, membrane composition and lifespan: An experimental study using the blowfly (*Calliphora stygia*). *Mechanisms of Ageing and Development, 138,* 15-25. doi: 10.1016/j.mad.2014.02.001

Maeda, T., Tamotsu, M., Yamaoka, R. & Ozaki, M. (2015). Effects of floral scents and their dietary experiences on the feeding preference in the blowfly, *Phormia regina. Frontiers in Integrative Neuroscience, 9*, 59. doi: 10.3389/fnint.2015.00059

Magni, P.A., Pacini, T., Pazzi, M., Vincenti, M. & Dadour, I.R. (2014). Development of a GC–MS method for methamphetamine detection in *Calliphora vomitoria* L. (Diptera: Calliphoridae). *Forensic Science International, 241*, 96-101. doi: 10.1016/j.forsciint.2014.05.004

Parker, F.D., Welch, J.B. & Matlock, R.B. Jr. (1993). Influence of habitat, season and attractant on adult behavior of the screwworm (Diptera: Calliphoridae) in a tropical dry zone in Costa Rica. *Journal of Economic Entomology, 86*(5), 1359-1375. doi: 10.1093/jee/86.5.1359

Pérez, C., Segura, N.A., Patarroyo, M.A. & Bello, F.J. (2016). Evaluating the biological cycle and reproductive and population parameters of *Calliphora vicina* (Diptera: Calliphoridae) reared on three different diets. *Journal of Medical Entomology, 53*(6), 1268-1275. doi: 10.1093/jme/tjw114

Pölkki, M., Kangassalo, K. & Rantala, M.J. (2012). Transgenerational effects of heavy metal pollution on immune defense of the blow fly *Protophormia terraenovae*. *PLoS ONE, 7*(6), p.e38832. doi: 10.1371/journal.pone.0038832

Wesolowski, T. (1987). Host–parasite interactions in natural holes: marsh tits (*Parus palustris*) and blow flies (*Protocalliphora falcozi*). *Journal of Zoology, 255*(4), 495-503. doi: 10.1017/S0952836901001571

Whitworth, T.L., Dawson, R.D., Magalon, H. & Baudry, E. (2007). DNA barcoding cannot reliably identify species of the blowfly genus *Protocalliphora* (Diptera: Calliphoridae). *Proceedings of the Royal Society. B, Biological sciences, 274*(1619), 1731-1739. doi: 10.1098/rspb.2007.0062

Cecidomyiidae

Aggarwal, R., Subramanyam, S., Zhao, C., Chen, M., Harris, M.O., Stuart, J.J. & Castagnone-Sereno, P. (2014). Avirulence effector discovery in a plant galling and plant parasitic arthropod, the Hessian fly (*Mayetiola destructor*). *PLoS One, 9*(6), p.e100958. doi: 10.1371/journal.pone.0100958

Chen, M., Liu, S., Wang, H., Cheng, X., El Bouhssini, M., Whitworth, R. J. & Escriva, H. (2016). Massive shift in gene expression during transitions between developmental stages of the gall midge, *Mayetiola destructor. PLoS One, 11*(5), p.e0155616. doi: 10.1371/journal.pone.0155616

Dorchin, N., Harris, K.M. & Stireman, J.O. (2019). Phylogeny of the gall midges (Diptera, Cecidomyiidae, Cecidomyiinae): Systematics, evolution of feeding modes and diversification rates. *Molecular Phylogenetics and Evolution, 140,* 106602. doi: 10.1016/j.ympev.2019.106602

Heath, J.J. & Stireman, J.O. III (2010). Dissecting the association between a gall midge, *Asteromyia carbonifera*, and its symbiotic fungus, *Botryosphaeria dothidea, 137*(1), 36-49. doi: 10.1111/j.1570-7458.2010.01040.x

Johnson, A. J., Shukle, R. H., Chen, M.S., Srivastava, S., Subramanyam, S., Schemerhorn, B. J., Weintraub, P. G., Abdel Moniem, H. E. M., Flanders, K. L., Buntin, G. D. & Williams, C. E. (2015). Differential expression of candidate salivary effector proteins in field collections of hessian fly, *Mayetiola destructor. Insect Molecular Biology, 24*(2), 191-202. doi: 10.1111/imb.12148

Sá, C.E.M., Silveira, F.A.O., Santos, J.C., Isaias, R.M.S. & Fernandes, G.W. (2009). Anatomical and developmental aspects of leaf galls induced by *Schizomyia macrocapillata* Maia (Diptera: Cecidomyiidae) on *Bauhinia brevipes* Vogel (Fabaceae). *Revista Brasileira de Botânica, 32*(2), 319-327. doi: 10.1590/S0100-84042009000200011

Sadeghi, R., Odubiyi, S., Nikoukar, A., Schroeder, K.L. & Rashed, A. (2021). *Mayetiola destructor* (Diptera: Cecidmyiidae) host preference and survival on small grains with respect to leaf reflectance and phytohormone concentrations. *Scientific Reports, 11*(1), 4761. doi: 10.1038/s41598-021-84212-x

Stireman, J.O., Janson, E.M., Carr, T.G., Devlin, H. & Abbot, P. (2008). Evolutionary radiation of *Asteromyia carbonifera* (Diptera: Cecidomyiidae) gall morphotypes on the goldenrod *Solidago altissima* (Asteraceae). *Biological Journal of the Linnean Society, 95*(4), 840-858. doi: 10.1111/j.1095-8312.2008.01101.x

Subramanyam, S., Shreve, J.T., Nemacheck, J.A., Johnson, A.J., Schemerhorn, B., Shukle, R.H. & Williams, C.E. (2018). Modulation of nonessential amino acid biosynthetic pathways in virulent hessian fly larvae (*Mayetiola destructor),* feeding on susceptible host wheat (*Triticum aestivum*). *Journal of Insect Physiology, 105*, 54-63. doi: 10.1016/j.jinsphys.2018.01.001

Wearing, C. H., Marshall, R. R., Attfield, B. & Colhoun, C. (2013). Phenology and distribution of the apple leafcurling midge (*Dasineura mali* (Kieffer)) (Diptera: Cecidomyiidae) and its natural enemies on apples under biological and integrated pest management in Central Otago, New Zealand. *New Zealand Entomologist, 36*(2), 87-106. doi: 10.1080/00779962.2012.712887

Ceratopogonidae

Bakhoum, M.T., Fall, A.G., Fall, M., Bassene, C.K., Baldet, T., Seck, M.T., Bouyer, J. Garros, C. & Gimonneau, G. (2016). Insight on the larval habitat of Afrotropical *Culicoides Latreille* (Diptera: Ceratopogonidae) in the Niayes area of Senegal, West Africa. *Parasites & Vectors, 9*(1), 462. doi: 10.1186/s13071-016-1749-1

Barceló, C. & Miranda, M.A. (2021). Development and lifespan of *Culicoides obsoletus* s.s. (Meigen) and other livestock-associated species reared at different temperatures under laboratory conditions. *Medical and Veterinary Entomology, 35*(2), 187-201. doi: 10.1111/mve.12487

Chen, H., Chou, J., Lin, C., Wen, Y. & Wang, W. (2016). Seasonal yeast compositions in *Forcipomyia taiwana* (Diptera: Ceratopogonidae). *Journal of Asia-Pacific Entomology, 19*(2), 509-514. doi: 10.1016/j.aspen.2016.04.020

Díaz, F., Mangudo C., Gleiser, R.M. & Ronderos, M.M. (2019). Redescription of immatures of *Dasyhelea flavifrons* Guérin-Méneville (Culicomorpha: Ceratopogonidae) and new contribution to the knowledge of its larval habitats. *Anais da Academia Brasileira de Ciências, 91*(1). doi: 10.1590/0001-3765201920180047

Erram, D., Blosser, E.M. & Burkett-Cadena, N. (2019). Habitat associations of *Culicoides* species (Diptera: Ceratopogonidae) abundant on a commercial cervid farm in Florida, USA. *Parasites & Vectors, 12*(1), 367. doi: 10.1186/s13071-019-3626-1

Erram, D. & Burkett-Cadena, N. (2021). Oviposition of *Culicoides insignis* (Diptera: Ceratopogonidae) under laboratory conditions with notes on the developmental life history traits of its immature stages. *Parasites & Vectors, 14*(1), 522. doi: 10.1186/s13071-021-05025-5

Erram, D., Burkett-Cadena, N. & Foil, L. (2019). Laboratory rearing of *Culicoides stellifer* (Diptera: Ceratopogonidae), a suspected vector of orbiviruses in the United States. *Journal of Medical Entomology, 57*(1), 25-32. doi: 10.1093/jme/tjz154

Erram, D. & Zurek, L. (2017). Larval development of *Culicoides sonorensis* (Diptera: Ceratopogonidae) in mud supplemented with manure of various farm animals. *Journal of Medical Entomology, 55*(1), 43-50. doi: 10.1093/jme/tjx197

Kameke, D., Kampen, H., Wacker, A. & Werner, D. (2021). Field studies on breeding sites of *Culicoides* LATREILLE (Diptera: Ceratopogonidae) in agriculturally used and natural habitats. *Scientific Reports, 11*(1), 10007. doi: 10.1038/s41598-021-86163-9

Liu, Y., Tao, H., Yu, Y., Yue, L., Xia, W., Zheng, W., Ma, H., Liu, X. & Chen, H. (2018). Molecular differentiation and species composition of genus *Culicoides* biting midges (Diptera: Ceratopogonidae) in different habitats in southern China. *Veterinary Parasitology, 254*, 49-57. doi: 10.1016/j.vetpar.2018.02.035

Riddin, M. A., Venter, G. J., Labuschagne, K. & Villet, M. H. (2019). Bloodmeal analysis in *Culicoides* midges collected near horses, donkeys and zebras in the Eastern Cape, South Africa. *Medical and Veterinary Entomology, 33*(4), 467-475. doi: 10.1111/mve.12381

Steinke, S., Lühken, R., Balczun, C. & Kiel, E. (2016). Emergence of *Culicoides obsoletus* group species from farm-associated habitats in Germany: Emergence of *Culicoides obsoletus* group species. *Medical and Veterinary Entomology, 30*(2), 174-184. doi: 10.1111/mve.12159

Urbanek, A. & Rost-Roszkowska, M.M. (2015). Ultrastructural studies on the midgut of biting midge *Forcipomyia nigra* (Winnertz) (Diptera: Ceratopogonidae). *Micron, 69,* 25-34. doi: 10.1016/j.micron.2014.11.003

Urbanek, A., Szadziewski, R., Stepnowski, P., Boros-Majewska, J., Gabriel, I., Dawgul, M., Kamysz, W., Sosnowska, D. & Gołębiowski, M. (2012). Composition and antimicrobial activity of fatty acids detected in the hygroscopic secretion collected from the secretory setae of larvae of the biting midge *Forcipomyia nigra* (Diptera: Ceratopogonidae). *Journal of Insect Physiology*, *58*(9), 1265-1276. doi: 10.1016/j.jinsphys.2012.06.014

Van den Eynde, C., Sohier, C., Matthijs, S. & De Regge, N. (2021). Temperature and food sources influence subadult development and blood-feeding response *of Culicoides obsoletus* (sensu lato) under laboratory conditions. *Parasites & Vectors, 14*(1), 1-300. doi: 10.1186/s13071-021-04781-8

Veiga, J., Martínez-de la Puente, J., Václav, R., Figuerola, J. & Valera, F. (2018*). Culicoides paolae* and *C. circumscriptus* as potential vectors of avian haemosporidians in an arid ecosystem. *Parasites & Vectors, 11*(1), 524. doi: 10.1186/s13071-018-3098-8

Werner, D., Groschupp, S., Bauer, C. & Kampen, H. (2020). Breeding habitat preferences of major Culicoides species (Diptera: Ceratopogonidae) in Germany. *International Journal of Environmental Research and Public Health, 17*(14), 5000. doi: 10.3390/ijerph17145000.

Yeh, W.-B., Lee, H.-M, Tu, W.-C, Tang, L.-C & Lee, P.-Y. (2009). Molecular differentiation and diversity of *Forcipomyia taiwana* (Diptera: Ceratopogonidae) based on the mitochondrial cytochrome oxidase II sequence. *Journal of Medical Entomology, 46*(2), 249-256. doi: 10.1603/033.046.0209

Chaoboridae

Barth, L.E., Sprules, W.G., Wells, M., Coman,M. & Prairie, Y. (2014). Seasonal changes in the diel vertical migration of *Chaoborus punctipennis* larval instars. *Canadian Journal of Fisheries and Aquatic Sciences, 71*(5), 665-674. doi: 10.1139/cjfas-2013-0440

Berendonk, T.U. & Spitze, K. (2006). Linking lakes? The population genetic structure of *Chaoborus flavicans. Journal of Plankton Research, 28*(11), 1015-1025. doi: 10.1093/plankt/fbl033

Härkönen, L., Pekcan-Hekim, Z., Hellén, N. & Horppila, J. (2013). Feeding efficiency of *Chaoborus flavicans* (Insecta, Diptera) under turbulent conditions. *Hydrobiologia, 722*(1), 9-17. doi: 10.1007/s10750-013-1670-y

Jäger, I.S., Hölker, F., Flöder, S. & Walz, N. (2011). Impact of *Chaoborus flavicans* -predation on the zooplankton in a mesotrophic lake - a three year study. *International Review of Hydrobiology, 96*(1), 191-208. doi: 10.1002/iroh.201011253

Moore, M.V. (1988). Differential use of food resources by the instars of *Chaoborus punctipennis. Freshwater Biology, 19*(2), 249-268. doi: 10.1111/j.1365-2427.1988.tb00346.x

Schroder, A. & Dam, H.G. (2013). Density- and size-dependent winter mortality and growth of late *Chaoborus flavicans* larvae. *PLoS One, 8*(10), p.e75839. doi: 10.1371/journal.pone.0075839

Chironomidae

Anderson, A.M., Stur, E. & Ekrem, T. (2013). Molecular and morphological methods reveal cryptic diversity and three new species of Nearctic *Micropsectra* (Diptera:Chironomidae). *Freshwater Science, 32*(3), 892-921. doi: 10.1899/12-026.1

Baniszewski, J., Miller, N., Kariuki, E.M., Cuda, J.P. & Weeks, E.N.I. (2020). *Cricotopus lebetis* intraspecific competition and damage to *Hydrilla. The Florida Entomologist, 103*(1), 32-37. doi: 10.1653/024.103.0405

Baniszewski, J., Weeks, E.N.I. & Cuda, J.P. (2016). *Bacillus thuringiensis* subspecies Kurstaki reduces competition by *Parapoynx diminutalis* (Lepidoptera: Crambidae) in colonies of the *Hydrilla* biological control agent *Cricotopus lebetis* (Diptera: Chironomidae). *The Florida Entomologist, 99*(4), 644-647. doi: 10.1653/024.099.0410

Belle, S., Millet, L., Gillet, F., Verneaux, V. & Magny, M. (2015). Assemblages and paleo-diet variability of subfossil Chironomidae (Diptera) from a deep lake (Lake Grand Maclu, France). *Hydrobiologia, 755*(1), 145-160. doi: 10.1007/s10750-015-2222-4

Boggero, A., Füreder, L., Lencioni, V., Simcic, T., Thaler, B., Ferrarese, U., Lotter, A. F. & Ettinger, R. (2006). Littoral chironomid communities of Alpine lakes in relation to environmental factors. *Hydrobiologia, 562*(1), 145-165. doi: 10.1007/s10750-005-1809-6

Boonsoong, B. (2016). Phoretic associations between *Nanocladius asiaticus* (Diptera, Chironomidae) and its hosts *Gestroiella* (Heteroptera, Naucoridae) and *Euphaea masoni* (Odonata, Euphaeidae) in streams in Western Thailand. *Annales de Limnologie, 52*, 163-169. doi: 10.1051/limn/2015025

Brown, N.E., Mitchell, S.C. & Garbary, D.J. (2013). Host selection by larvae of a marine insect *Halocladius variabilis*: nutritional dependency or escape from predation? *Journal of the Marine Biological Association of the United Kingdom, 93*(5), 1373-1379. doi: 10.1017/S0025315412001634

Cranston, P.S. & Krosch, M.N. (2015). DNA sequences and austral taxa indicate generic synonymy of *Paratrichocladius* Santos-Abreu with *Cricotopus* Wulp (Diptera: Chironomidae). *Systematic Entomology, 40*(4), 719-732. doi: 10.1111/syen.12130

Delettre, Y.R. (2005). Short-range spatial patterning of terrestrial Chironomidae (Insecta: Diptera) and farmland heterogeneity. *Pedobiologia, 49*(1), 15-27. Doi: 10.1016/j.pedobi.2004.06.010

Diserud, O.H., Stur, E. & Aagaard, K. (2012). How reliable are malaise traps for biomonitoring? -A bivariate species abundance model evaluation using alpine Chironomidae (Diptera). *Insect Conservation and Diversity, 6*(5), 561-571. doi: 10.1111/icad.12012

Dosdall, L.M. & Parker, D.W. (1998). First report of a symphoretic association between *Nanocladius branchicolus* Saether (Diptera: Chironomidae) and *Argia moesta* (Hagen) (Odonata: Coenagrionidae). *The American Midland Naturalist, 139*(1), 181-185. Doi: 10.1674/0003-0031(1998)139[0181:FROASA]2.0.CO2

Durnova, N. A., Demin, A. G., Polukonova, N. V. & Mugue, N. S. (2014). The time of origin of the capacity to accrete and mine submerged substrates in the midge subfamilies Chironominae Macquart, 1838 and Orthocladiinae Lenz, 1921 (Diptera, Chironomidae): Analysis of mitochondrial genes COI and COII. *Entomological Review, 94*(7), 949-958. doi: 10.1134/S0013873814070045

Earle, W., Mangan, R., O'Brien, M. & Baars, J. (2013). Biology of *Polypedilum* n. sp. (Diptera: Chironomidae), a promising candidate agent for the biological control of the aquatic weed *Lagarosiphon* major (Hydrocharitaceae) in Ireland. *Biocontrol Science and Technology, 23*(11), 1267-1283. doi: 10.1080/09583157.2013.826344

Egan, A.T. & Ferrington, L.C. (2015). Chironomidae (Diptera) in freshwater coastal rock pools at Isle Royale, Michigan. *Transactions of the American Entomological Society, 141*(1), 1-25. doi: 10.3157/061.141.0102

Egan, A.T. & Ferrington, L.C. (2019). Chironomidae of the upper Saint Croix river, Wisconsin. *Transactions of the American Entomological Society, 145*(3), 353-384. doi: 10.3157/061/145/0307

Enushchenko, I.V. (2018). Most interesting findings from remnants of Chironomid (Insecta: Diptera: Chironomidae) larvae in layers of a 200-year-old stratum of lake Oron bottom sediments. *Russian Journal of Ecology, 49*(1), 69-74. doi: 10.1134/S1067413618010058

Ghonaim, M., Ali, A. & Salem, M. (2004). *Tanytarsus* (Diptera: Chironomidae) from Egypt with description of a new species. *The Florida Entomologist, 87*(4), 571-575. doi: 10.1653/0015-4040(2004)087[0571:TDCEWD]2.0.CO2

Goffová, K., Čiamporová-Zat'ovičová, Z. & Čiampor, F. (2015). Twenty-one new microsatellite markers for the ecologically important midge *Heterotrissocladius marcidus*, and their use in studies of alpine lakes. *Biochemical Systematics and Ecology, 61*, 329-335. doi: 10.1016/j.bse.2015.06.037

Hänel, C. & Chown, S.L. (1997). The impact of a small, alien invertebrate on a sub-Antarctic terrestrial ecosystem: *Limnophyes minimus* (Diptera, Chironomidae) at Marion Island. *Polar Biology, 20*(2), 99-106. doi: 10.1007/s003000050282

Hazra, N., Saha, G.K. & Chaudhuri, P.K. (2002). Records of Orthoclad species from the Darjeeling-Sikkim Himalayas of India (Diptera: Chironomidae), with notes on their ecology. *Hydrobiologia, 474*(1), 41-55. doi: 10.1023/A:1016511702944

Hayford, B. (2012). Parochlus Kiefferi (Garrett, 1925) in Nebraska (Diptera: Chrionomidae). *Great Plains Research, 22*(1), 27-33.

Hershey, A.E. (1986). Selective predation by *Procladius* in an Arctic Alaskan lake. *Canadian Journal of Fisheries and Aquatic Sciences, 43*(12), 2523-2528. doi: 10.1139/f86-312

Ingvason, H.R., Ólafsson, J.S. & Gardarsson, A. (2004). Food selection of *Tanytarsus gracilentus* larvae (Diptera: Chironomidae): An analysis of instars and cohorts. *Aquatic Ecology, 38*(2), 231-237. doi: 10.1023/B:AECO.0000032053.67992.03

Johnson, R.K. (1987). Seasonal variation in diet of *Chironomus plumosus* (L.) and *C. anthracinus* Zett. (Diptera: Chironomidae) in mesotrophic Lake Erken. *Freshwater Biology, 17*(3), 525-532. doi: 10.1111/j.1365-2427.1987.tb01073.x

Lencioni, V. & Bernabò, P. (2015). Thermal survival limits of young and mature larvae of a cold stenothermal chironomid from the Alps (Diamesinae: Pseudodiamesa branickii [Nowicki, 1873]. *Insect Science, 24*(2), 314-324. doi: 10.1111/1744-7917.12278

Li, X. & Wang, X. (2014). New species and records of *Metriocnemus* van der Wulp s. str. from China (Diptera, Chironomidae). *ZooKeys, 387*(387), 73-87. doi: 10.3897/zookeys.387.6408

Lilley, T.M., Ruokolainen, L., Pikkarainen, A., Laine, V.N., Kilpimaa, J., Rantala, M.J., Nikinmaa, M. (2012). Impact of Tributyltin on immune response and life history traits of *Chironomus riparius*: single and multigeneration effects and recovery from pollution. *Environmental Science & Technology, 46*(13), 7382-7389. doi: 10.1021/es300536t

Malchow, D.E., Knight, A.W. & Maier, K.J. (1995). Bioaccumulation and toxicity of selenium in *Chironomus decorus* larvae fed a diet of seleniferous *Selenastrum capricornutum. Archives of Environmental Contamination and Toxicology, 29*(1), 104-109. doi: 10.1007/BF00213094

McLachlan, A.J. (1986). Sexual dimorphism in midges: strategies for flight in the rain-pool dweller *Chironomus imicola* (Diptera: Chironomidae). *The Journal of Animal Ecology, 55*(1), 261-267. doi: 10.2307/4706

Medeiros, A.S., Milošević, D., Francis, D.R., Maddison, E., Woodroffe, S., Long, A., Walker, I.R., Hamerlík, L., Quinlan, R., Langdon, P., Brodersen, K.P. & Axford, Y. (2020). Arctic chironomids of the northwest North Atlantic reflect environmental and biogeographic gradients. *Journal of Biogeography, 48*(3), 511-525. doi: 10.1111/jbi.14015

Medeiros, A.S. & Quinlan, R. (2011). The distribution of the Chironomidae (Insecta: Diptera) along multiple environmental gradients in lakes and ponds of the eastern Canadian Arctic. *Canadian Journal of Fisheries and Aquatic Sciences, 68*(9), 1511-1527. doi: 10.1139/F2011-076

Moubayed-Breil, J. & Bitušík, P. (2019). Taxonomic notes on the genus *Chaetocladius* (laminatus-group) (Diptera: Chironomidae, Orthocladiinae) II. Descriptions of *C. bitusiki* sp. n. and *C. mantetensis* sp. n., two relict species inhabiting cold alpine springs and streams. *Biologia, 74*(11), 1489-1500. doi: 10.2478/s11756-019-00253-8

Namayandeh, A. & Beresford, D.V. (2019). Chironomidae of Fosheim Peninsula, Ellesmere Island: first female description of *Chaetocladius* (Chaetocladius) *glacialis* (Lundström, 1915) (Diptera: Chironomidae: Orthocladiinae) and new faunistic records from the Nunavut (Canada) and the Nearctic Region. *Aquatic Insects, 40*(1), 1-18. doi: 10.1080/01650424.2018.1523435

Niedrist, G.H. & Füreder, L. (2018). When the going gets tough, the tough get going: The enigma of survival strategies in harsh glacial stream environments. *Freshwater Biology, 63*(10), 1260-1272. doi: 10.1111/fwb.13131

Nieto, E., Corada-Fernández, C., Hampel, M., Lara-Martín, P.A., Sánchez-Argüello, P. & Blasco, J. (2017). Effects of exposure to pharmaceuticals (diclofenac and carbamazepine) spiked sediments in the midge, *Chironomus riparius* (Diptera, Chironomidae). *The Science of the Total Environment, 609*, 715-723. doi: 10.1016/j.scitotenv.2017.07.171

Nolte, U. & Hoffmann, T. (1992). Fast life in cold water*: Diamesa incallida* (Chironomidae). *Ecography, 15*(1), 25-30. doi: 10.1111/j.1600-0587.1992.tb00004.x

Paasivirta, L. (2014). Checklist of the family Chironomidae (Diptera) of Finland. *ZooKeys, 441*(441), 63-90. doi: 10.3897/zookeys.441.7461

Prat, N. & García-Roger, E.M. (2018). Are there so many congeneric species of chironomid larvae in a small stream? *Journal of Limnology, 77*(s1), 65-80. doi: 10.4081/jlimnol.2018.1757

Ricciardi, A. (1994). Occurrence of chironomid larvae (*Paratanytarsus* sp.) as commensals of dreissenid mussels (*Dreissena polymorpha* and *D. bugensis*). *Canadian Journal of Zoology, 72*(6), 1159-1162. doi: 10.1139/z94-155

Silva, C.J.M., Silva, A.L.P., Gravato, C. & Pestana, J.L.T. (2019). Ingestion of small-sized and irregularly shaped polyethylene microplastics affect *Chironomus riparius* life-history traits. *The Science of the Total Environment, 672*, 862-868. doi: 10.1016/j.scitotenv.2019.04.017

Som, D.K., Das, N. & Hazra, N. (2013). Systematics and biology of Metriocnemus clarivirgulus sp. n. (Diptera: Chrionomidae) from Darjeeling, India with revised keys to male and female adults of Metriocnemus van der Wulp. *Deutsche Entomologische Zeitschrift*, *60*(1), 111-121. doi: 10.1002/mmnd.201300014

Stoaks, R. (2009). Rare and unusual arthropod ecology, behavior and occurrence at hummingbird feeders in northern California. *The Pan-Pacific Entomologist, 85*(4), 194-198. doi: 10.3956/2009-18.1

Stur, E. & Ekrem, T. (2006). A revision of West Palaearctic species of the *Micropsectra atrofasciata* species group (Diptera: Chironomidae). *Zoological Journal of the Linnean Society, 146*(2), 165-225. doi: 10.1111/j.1096-3642.2006.00198.x

Stur, E. & Ekrem, T. (2015). A review of Norwegian *Gymnometriocnemus* (Diptera, Chironomidae) including the description of two new species and a new name for *Gymnometriocnemus volitans* (Goetghebuer) sensu Brundin. *ZooKeys, 508*(508), 127-142. doi: 10.3897/zookeys.508.9874

Syrjämäki, J. (1968). Diel patterns of swarming and other activities of two Arctic Dipterans (Chironomidae and Trichoceridae) on Spitsbergen. *Oikos, 19*(2), 250-258. doi: 10.2307.3565012

Syrovátka, V. (2018). The predatory behaviour of *Monopelopia tenuicalcar* (Kieffer, 1918) larvae in a laboratory experiment. *Journal of Limnology, 77.* doi: 10.4081/jlimnol.2018.1792

Tarakhovskaya, E.R. & Garbary, D.J. (2009). *Halocladius variabilis* (Diptera: Chironomidae): a marine insect symbiotic with seaweeds from the White Sea, Russia. *Journal of the Marine Biological Association of the United Kingdom, 89*(7), 1381-1385. doi: 10.1017/S0025315409000071

Tarkowska-Kukuryk, M. (2014). Spatial distribution of epiphytic chironomid larvae in a shallow macrophyte-dominated lake: effect of macrophyte species and food resources. *Limnology, 15*(2), 141-153. doi: 10.1007/s10201-014-0425-4

Torrisi, G.J., Hoback, W.W., Foster, J.E., Heinrichs, E.A. & Higley, L.G. (2015). Predation of introduced mosquito larvae by the midge *Metriocnemus knabi* in the Phytotelma of the pitcher plant Sarracenia purpurea and colonization following dry conditions. *Northeastern Naturalist, 22*(3), 513-520. doi: 10.1656/045.022.0307

Tucker, T.R., Hudson, P.L. & Riley, S.C. (2016). Observations of cocooned *Hydrobaenus* (Diptera: Chironomidae) larvae in Lake Michigan. *Journal of Great Lakes Research*, *42*(5), 1129-1135. doi: 10.1016/j.jglr.2016.07.013

Vinogradova, E.M., Riss, H.W. & Spies, M. (2009). New species of *Tanytarsus* van der Wulp, 1874 (Diptera: Chironomidae) from Central America. *Aquatic Insects, 31*(1), 11-17. doi: 10.1080/01650420802502477

Wrubleski, D.A. & Roback, S.S. (1987). Two species of *Procladius* (Diptera:Chironomidae) from a Northern prairie marsh: descriptions, phenologies and mating behaviour. *Journal of the North American Benthological Society, 6*(3), 198-212. doi: 10.2307/1467511

Younc, R.M. (1969). Field observations on a midwinter breeding flight of Diamesa arctica (Diptera: Chironomidae). *Annals of the Entomological Society of America, 62*(5), 1204. doi: 10.1093/aesa/62.5.1204

Chloropidae

Allen, W.A. & Pienkowski, R.L. (1974). The biology and seasonal abundance of the fruit fly, *Oscinella* in reed canarygrass in Virginia. *Annals of the Entomological Society of America, 67*(4), 539-544. doi: 10.1093/aesa/67.4.539

Jones, M.G. (1969). Effect of weather on frit fly (*Oscinella* frit L.) and its predators. *The Journal of Applied Ecology, 6*(3), 425-441. doi: 10.2307/2401509

Nartshuk, E.P. (2014). Grass-fly larvae (Diptera, Chloropidae): Diversity, habitats, and feeding specializations. *Entomological Review, 94*(4), 514-525. doi: 10.1134/S001387381404006X

Nartshuk, E.P. & Khruleva, O.A. (2011). Plant-feeding dipterans (Diptera, Chloropidae, Agromyzidae) from Wrangel Island (the Chukchi Sea). *Entomological Review, 91*(7), 849-854. doi: 10.1134/S0013873811070062

Riccardi, P.R. & Amorim, D.S. (2020). Phylogenetic relationships and classification of the Chloropinae of the world (Diptera: Chloropidae). *Zoological Journal of the Linnean Society, 190*(3), 889-941. doi: 10.1093/zoolinnean/zlaa007

Triseleva, T.A., Petrosyan, V.G., Yatsuk, A.A. & Safonkin, A.F. (2021). The role of plants in the formation of species-specific features in grass flies (Diptera, Chloropidae, Meromyza). *Biodiversity Data Journal, 9*(20), p.e78017. doi: 10.3897/BDJ.9.e78017

Vandevyvere, I. & Bruyn, L. de. (1998). Morphological and histochemical analysis of galls of *Lipara lucens* (Diptera, Chloropidae) on *Phragmites australis* (Poaceae). *Canadian Journal of Botany, 76*(8), 1374-1384. doi: 10.1139/b98-143

Clusiidae

Roháček, J. (2012). The fauna of the opomyzoid families Clusiidae, Acartophthalmidae, Anthomyzidae, Opomyzidae, Stenomicridae, Periscelididae, Asteiidae (Diptera) in the Gemer area (Central Slovakia). Časopis Slezského Zemského Muzea, 61(2), 97-111. doi: 10.2478/v10210-012-0011-5

Coelopidae

Edward, D.A., Blyth, J.E., Mckee, R. & Gilburn, A.S. (2007). Change in the distribution of a member of the strand line community: the seaweed fly (Diptera: Coelopidae). *Ecological Entomology, 32*(6), 741-746. doi: 10.1111/j.1365-2311.2007.00919.x

Culicidae

Andrew, E.S., Xu, G. & Rich, S.M. (2014). Microbial communities within field-collected *Culiseta melanura* and *Coquillettidia perturbans. Medical and Veterinary Entomology, 28*(2), 125-132. doi: 10.1111/mve.12017

Anderson, I.H. (1992). The effect of sugar meals and body size on fecundity and longevity of female *Aedes communis* (Diptera: Culicidae). *Physiological Entomology, 17*(3), 203-207. Doi: 10.1111/j.1365-3032.1992.tb01011.x

Bartlett-Healy, K., Crans, W. & Gaugler, R. (2009). Vertebrate hosts and phylogenetic relationships of amphibian trypanosomes from a potential invertebrate vector, *Culex territans* Walker (Diptera: Culicidae). *The Journal of Parasitology, 95*(2), 381-387. doi: 10.1645/GE-1793.1

Benedict, M.Q., Hunt, C.M., Vella, M.G., Gonzalez, K.M., Dotson, E.M., Collins, C.M. & Hansen, I.A. (2020). Pragmatic selection of larval mosquito diets for insectary rearing of *Anopheles gambiae* and *Aedes aegypti. PLoS One, 15*(3), p.e0221838. doi: 10.1371/journal.pone.0221838

Friend, W.G., Smith, J.J.B., Schmidt, J.M. & Tanner, R.J. (1989). Ingestion and diet destination in *Culiseta inornata*: responses to water, sucrose and cellobiose. *Physiological Entomology, 14*(2), 137-146. doi: 10.1111/j.1365-3032.1989.tb00945.x

Gadawski, R.M. & Smith, S.M. (1992). Nectar sources and age structure in a population of *Aedes provocans* (Ditpera: Culicidae). *Journal of Medical Entomology, 29*(5), 879-886. doi: 10.1093/jmedent/29.5.879

Greenberg, J.A., Lujan, D.A., DiMenna, M.A., Wearing, H.J. & Hofkin, B.V. (2013). Identification of blood meal sources in *Aedes vexans* and *Culex quinquefasciatus* in Bernalillo County, New Mexico. *Journal of Insect Science, 13*(75), 1-12. Doi: 10.1673/031.013.7501

Ribeiro, J.M.C., Martin-Martin, I., Arcà, B., Calvo, E. & Hansen, I.A. (2016). A deep insight into the sialome of male and female *Aedes aegypti* mosquitoes. *PLoS One, 11*(3), p.e0151400. doi: 10.1371/journal.pone.0151400

Saab, S.A., zu Dohna, H., Nilsson, L.K.J., Onorati, P., Nakhleh, J., Terenius, O. & Osta, M.A. (2020). The environment and species affect gut bacteria composition in laboratory co-cultured *Anopheles gambiae* and *Aedes albopictus* mosquitoes. *Scientific Reports, 10*(1), 3352. doi: 10.1038/s41598-020-60075-6

Schaefer, M.L. & Lundstroem, J.O. (2006). Different responses of two floodwater mosquito species, *Aedes vexans* and *Ochlerotatus sticticus* (Diptera: Culicidae), to larval habitat drying. *Journal of Vector Ecology, 31*(1), 123-128.

Sherwood, J. A., Stehman, S., Howard, J. J. & Oliver, J. (2020). Cases of Eastern equine encephalitis in humans associated with *Aedes canadensis*, *Coquillettidia perturbans* and *Culiseta melanura* mosquitoes

with the virus in New York State from 1971 to 2012 by analysis of aggregated published data. *Epidemiology and Infection, 148*, p.e72. doi: 10.1017/S0950268820000308

Smith, S.M. & Gadawski, R.M. (1994). Nectar feeding by the early-spring mosquito *Aedes provocans. Medical and Veterinary Entomology, 8*(3), 201-213. doi: 10.1111/j.1365-2915.1994.tb00499.x

Yee, W.L., Foster, W.A., Howe, M.J. & Hancock, R.G. (1992). Simultaneous field comparison of evening temporal distributions of nectar and blood feeding by *Aedes vexans and Ae. trivittatus* (Diptera: Culicidae) in Ohio. *Journal of Medical Entomology, 29*(2), 356-360. doi: 10.1093/jmedent/29.2.356

Dolichopodidae

Bernasconi, M.V., Pollet, M., Varini-Ooijen, M., Irvine Ward, P. (2007). Phylogeny of European *Dolichopus* and *Gymnopternus* (Diptera: Dolichopodidae) and the significance of morphological characters inferred from molecular data. *European Journal of Entomology, 104*(3), 601-617. doi: 10.14411/eje.2007.075

Chursina, M.A. & Negrobov, O.P. (2020). Morphometric variations in leg segments' length in the family Dolichopodidae (Diptera). *Moscow University Biological Sciences Bulletin, 75*(3), 110-116. doi: 10.3103/S0096392520030049

Gelbic, I. & Olejnicek, J. (2011). Ecology of Dolichopodidae (Diptera) in a wetland habitat and their potential role as bioindicators. *Central European Journal of Biology, 6*(1), 118-129. doi: 10.2478/s11535-010-0098-x

Hulcr, J., Pollet, M., Ubik, K. & Vrkoč, J. (2005). Exploitation of kairomones and synomones by *Medetera* spp. (Diptera: Dolichopodidae), predators of spruce bark beetles. *European Journal of Entomology, 102*(4), 655-662. Doi: 10.14411/eje.2005.093

Kautz, A.R. & Gardiner, M.M. (2019). Agricultural intensification may create an attractive sink for Dolichopodidae, a ubiquitous but understudied predatory fly family. *Journal of Insect Conservation, 23*(3), 453-465. doi: 10.1007/s10841-018-0116-2

Martin, O.Y., Puniamoorthy, N., Gubler, A., Wimmer, C. & Bernasconi, M.V. (2013). Infections with Wolbachia, Spiroplasma, and Rickettsia in the Dolichopodidae and other Empidoidea. *Infection, Genetics and Evolution, 13*, 317-330. doi: 10.1016/j.meegid.2012.11.005

Mcghehey, J.H. & Nagel, W.P. (1966). A technique for rearing larvae of *Medetera aldrichii* (Diptera: Dolichopodidae). *Annals of the Entomological Society of America, 59*(2), 290-292. doi: 10.1093/aesa/59.2.290

Drosophilidae

Band, H.T., Bachli, G. & Band, R.N. (2005). Behavioral constancy for interspecies dependency enables Nearctic *Chymomyza amoena* (Loew) (Diptera: Drosophilidae) to spread in orchards and forests in Central and Southern Europe. *Biological Invasions, 7*(3), 509-530. doi: 10.1007/s10530-004-6352-2

Church, S.H., Extavour, C.G. & Rosenberg, M. (2022). Phylotranscriptomics reveals discordance in the phylogeny of Hawaiian *Drosophila* and *Scaptomyza* (Diptera: Drosophilidae). *Molecular Biology and Evolution, 39*(3). doi: 10.1093/molbev/msac012

Dyer, K.A. (2012). Local selection underlies the geographic distribution of sex-ratio drive in *Drosophila neotestacea*. *Evolution, 66*(4), 973-984. doi: 10.1111/j.1558-5646.2011.01497.x

Dyer, KA., White, B.E., Sztepanacz, J.L., Bewick, E.R. & Rundle, H.D. (2014). Reproductive character displacement of epicuticular compounds and their contribution to mate choice in *Drosophila subquinaria* and *Drosophila recens*. *Evolution, 68*(4), 1163-1175. doi: 10.1111/evo.12335

Jaenike, J. (1985). Genetic and environmental determinants of food preference in *Drosophila tripunctata. Evolution, 39*(2), 362-369. doi: 10.1111/j.1558-5646.1985.tb05673.x

James, A.C., Jakubczak, J., Riley, M.P. & Jaenike, J. (1988). On the causes of monophagy in *Drosophila quinaria. Evolution, 42*(3), 626-630. doi: 10.1111/j.1558-5646.1988.tb04166.x

Jaramillo, S.L., Mehlferber, E. & Moore, P.J. (2015). Life-history trade-offs under different larval diets in *Drosophila suzukii* (Diptera: Drosophilidae). *Physiological Entomology, 40*(1), 2-9. doi: 10.1111/phen.12082

Koštál, V., Mollaei, M. & Schöttner, K. (2016). Diapause induction as an interplay between seasonal token stimuli, and modifying and directly limiting factors: hibernation in *Chymomyza costata*: Environmental factors inducing diapause. *Physiological Entomology, 41*(4), 344-357. doi: 10.1111/phen.12159

Stehlík, J., Závodská, R., Shimada, K., Šauman, I. & Koštál, V. (2008). Photoperiodic induction of diapause requires regulated transcription of timeless in the larval brain of *Chymomyza costata. Journal of Biological Rhythms, 23*(2), 129-139. doi: 10.1177/0748730407313364

Trajković, J., Vujić, V., Miličić, D., Gojgić-Cvijović, G., Pavković-Lučić, S. & Savić, T. (2017). Fitness traits of *Drosophila melanogaster* (Diptera: Drosophilidae) after long-term laboratory rearing on different diets. *European Journal of Entomology, 144*(1), 222-229. doi: 10.14411/eje.2017.027

Worthen, W.B., Jones, M.T. & Jetton, R.M. (1998). Community structure and environmental stress: desiccation promotes nestedness in mycophagous fly communities [*Drosophila tripunctata, Drosophila putrida*, nested-subset pattern, depauperate assemblage]. *Oikos, 81*(1), 45-54. doi: 10.2307/3546466

Dryomyzidae

Gibson, J.F. & Choong, H.H.C. (2021). New range records and life history observations of insects (Diptera: Dryomyzidae, Chironomidae; Coleoptera: Staphylinidae) associated with barnacles (Balanomorpha:

Balanidae, Chthamalidae) on the Pacific coasts of North America and Japan. *Canadian Entomologist, 153*(2), 196-210. doi: 10.4039/tce.2020.69

Harley, C.D.G. & Lopez, J.P. (2003). The natural history, thermal physiology, and ecological impacts of intertidal mesopredators, *Oedoparena* spp. (Diptera: Dryomyzidae). *Invertebrate Biology, 122*(1), 61-73. doi: 10.1111/j.1744-7410.2003.tb00073.x

Empididae

Alcock, J. (2016). The mating behavior of an undescribed species of *Rhamphomyia* (Diptera: Empididae). *Journal of Insect Behavior, 29*(2), 153-161. doi: 10.1007/s10905-016-9545-5

Bussière, L.F., Gwynne, D.T. & Brooks, R. (2008). Contrasting sexual selection on males and females in a role-reversed swarming dance fly, *Rhamphomyia longicauda* Loew (Diptera: Empididae). *Journal of Evolutionary Biology, 21*(6), 1683-1691. doi: 10.1111/j.1420-9101.2008.01580.x

Evans, H.E. (1988). Observations on swarms of *Rhamphomyia sociabilis* (Williston) (Diptera: Empididae). *Journal of the New York Entomological Society, 96*(3), 316-322.

Gwynne, D.T. & Bussière, L.F. (2002). Female mating swarms increase predation risk in a 'role-reversed' dance fly (Diptera: Empididae: *Rhamphomyia longicauda* Loew). *Behaviour, 139*(11-12), 1425-1430. doi: 10.1163/15685390260514690

Ivkovic, M., Perovic, M., Grootaert, P. & Pollet, M. (2021). High endemicity in aquatic dance flies of Corsica, France (Diptera, Empididae, Clinocerinae and Hemerodromiinae), with the description of a new species of *Chelipoda. ZooKeys, 1039*(1039), 177-197. doi: 10.3897/zookeys.1039.66493

Khruleva, O. A., Shamshev, I. V. & Sinclair, B. J. (2021). The Empidoid Flies (Diptera: Brachystomatidae, Empididae, Hybotidae) of Wrangel Island (Chukotka Autonomous Okrug): composition and distribution of the fauna. *Entomological Review, 101*(6), 792-819. doi: 10.1134/S0013873821060063

Poinar, G.O. & Vega, F.E. (2021). A new genus of Empididae (Diptera) with enlarged postpedicels in mid-Cretaceous Burmese amber. *Historical Biology, 33*(9), 1830-1835. doi: 10.1080/08912963.2020.1743700

Shamshev, I.V. & Selitskaya, O.G. (2016). Methyl salicylate as an attractant for the dance fly *Rhamphomyia gibba* (Fallen) (Diptera, Empididae). *Entomological Review, 96*(8), 1003-1007. doi: 10.1134/S0013873816080054

Shamshev, I.V. & Sinclair, B.J. (2009). Revision of the *Iteaphila setosa* group (Diptera: Empididae). *European Journal of Entomology, 106*(3), 441-450. doi: 10.14411/eje.2009.055

Ephydridae

Bownes, A. (2018). Effects of *Hydrilla verticillata* (L.f.) Royle (Hydrocharitaceae) growing conditions and nutrient content on the performance of a leaf-mining fly, *Hydrellia purcelli* Deeming (Diptera: Ephydridae). *Biocontrol Science and Technology, 28*(3), 278-292. doi: 10.1080/09583157.2018.1441370

Cabrera Walsh, G., Magalí Dalto, Y., Mattioli, Federico M., Carruthers, Raymond I. & Anderson, L.W. (2013). Biology and ecology of Brazilian elodea *(Egeria densa)* and its specific herbivore*, Hydrellia* sp., in Argentina. *BioControl, 58*(1), 133-147. doi: 10.1007/s10526-012-9475-x

Prado e Castro, C., Santos, V. & Ameixa, O.M.C.C. (2022). Shore fly (Diptera: Ephydridae) communities from Ria de Aveiro, with new records for Portugal. *Fragmenta Entomologica, 54*(1). doi: 10.13133/2284-4880/715

Purcell, M.F., Brown, B.T., Harms, N.E., Sun-Hee, H. & McCulloch, G.A. (2021). Biology and preliminary host range of a Korean leaf-mining *Hydrellia* sp. (Diptera: Ephydridae) rejected as a potential biological control agent for monoecious *Hydrilla verticillata* in the United States. *Biocontrol Science and Technology, 31*(4), 343-356. doi: 10.1080/09583157.2020.1853051

Kubatova-Hirsova, H. (2005). Temporal patterns in shore fly (Diptera, Ephydridae) community structure in a salt marsh habitat. *Ecological Entomology, 30*(2), 234-240. doi: 10.1111/j.0307-6946.2005.00602.x

Mangan, R., Dirilgen, T. & Baars, J. (2015). Responses of adult *Hydrellia lagarosiphon* to a revised diet: implications for life cycle studies and laboratory culturing techniques. *Entomologia Experimentalis et Applicata, 157*(2), 164-169. doi: 10.1111/eea.12350

Morton, A. & García del Pino, F. (2007). Susceptibility of shore fly *Scatella stagnalis* to five entomopathogenic nematode strains in bioassays. *BioControl, 52*(4), 533-545. doi: 10.1007/s10526-006-9047-z

Scheiring, J.F. (1974). Diversity of shore flies (Diptera: Ephydridae) in inland freshwater habitat. *Journal of the Kansas Entomological Society, 47*(4), 485-491.

Scheiring, J.F. (1975). A multivariate analysis of similarity in shore fly habitats (Diptera: Ephydridae). *Journal of the Kansas Entomological Society, 48*(2), 232-243.

Scheiring, J.F. (1977). Geographic variation in *Scatella stagnalis* (Diptera: Ephydridae) [North America, insects]. *Annals of the Entomological Society of America, 70*(4), 511-523. doi: 10.1093/aesa/70.4.511

Smith, R., Mangan, R. & Coetzee, J.A. (2019). Risk assessment to interpret the physiological host range of *Hydrellia egeriae*, a biocontrol agent for *Egeria densa. BioControl, 64*(4), 447-456. Doi: 10.1007/s10526-019-09942-4

Vanninen, I. (2001). Biology of the shore fly *Scatella stagnalis* in rockwool under greenhouse conditions. *Entomologia Experimentalis et Applicata, 98*(3), 317-328. doi: 10.1046/j.1570-7458.2001.00788.x

Fanniidae

Velásquez, Y., Martínez-Sánchez, A. & Rojo, S. (2013). First record of *Fannia leucosticta* (Meigen) (Diptera: Fanniidae) breeding in human corpses. *Forensic Science International, 229*(1), p.e13-e15. doi: 10.1016/j.forsciint.2013.03.022

Wang, M., Li, W., Zhu, W. & Zhang, D. (2016). Review of the *Fannia postica*-group Chillcott, 1961 of the genus *Fannia* Robineau-Desvoidy, 1830, with description of two new species from the Palearctic and Oriental regions (Diptera, Fanniidae). *ZooKeys, 598*(598), 113-128. doi: 10.3897/zookeys.598.7983

Heleomyzidae

Rotheray, G.E. (2012). Morphology of the puparium and breeding sites of eight species of Heleomyzidae (Diptera). *Journal of Natural History, 46*(33-34), 2075-2102. doi: 10.1080/00222933.2012.707241

Soszynska-Maj, A. & Woznica, A.J. (2016). A case study of Heleomyzidae (Diptera) recorded on snow in Poland with a review of their winter activity in Europe. *European Journal of Entomology, 113*(1), 279-294. doi: 10.14411/eje.2016.035

Tuno, N., Sagara, N. & Okadome, T. (2003). Territorial behaviour of *Suillia* males on basidiocarps of *Hebeloma radicosum* in central Japan. *Mycologist, 17*(3), 122-125. doi: 10.1017/S0269915X03002143

Hybotidae

Bartak, M. & Kubik, S. (2015). Three new species of European *Platypalpus* (Diptera, Hybotidae). *ZooKeys, 470*(470), 145-155. doi: 10.3897/zookeys.470.8967

Jaume-Schinkel, S., Machado Barros, L., Breno Graça, M. & Mota Soares, M.M. (2022). In natura sit-and-wait behaviour and predation success of a Neotropical dance fly (Diptera: Hybotidae). *Journal of Natural History, 56*(1-4), 1-14. doi: 10.1080/00222933.2022.2032443

Khruleva, O. A., Shamshev, I. V. & Sinclair, B. J. (2021). The Empidoid Flies (Diptera: Brachystomatidae, Empididae, Hybotidae) of Wrangel Island (Chukotka Autonomous Okrug): composition and distribution of the fauna. *Entomological Review, 101*(6), 792-819. doi: 10.1134/S0013873821060063

Keroplatidae

Sivinski, J. (1982). Prey attraction by luminous larvae of the fungus gnat *Orfelia fultoni. Ecological Entomology, 7*(4), 443-446. doi: 10.1111/j.1365-2311.1982.tb00686.x

Viviani, V.R., Silva, J.R., Amaral, D.T., Bevilaqua, V.R., Abdalla, F.C., Branchini, B.R. & Johnson, C.H. (2020). A new brilliantly blue-emitting luciferin-luciferase system from *Orfelia fultoni* and Keroplatinae (Diptera). *Scientific Reports, 10*(1), 9608. doi: 10.1038/s41598-020-66286-1

Lauxaniidae

Davies, G.B.P. & Miller, R.M. (2009). Revision of Afrotropical *Pachycerina* Macquart (Diptera: Lauxaniidae). *African Invertebrates, 50*(2), 295-320. doi: 10.10520/EJC84650

Hesler, L.S. & Fernandes, G.W. (2016). Capture of nontarget flies (Diptera: Lauxaniidae, Chloropidae, and Anthomyiidae) on traps baited with volatile chemicals in field-crop habitats. *Psyche, 2016,* 1-8. doi: 10.1155/2016/6938368

Limoniidae

Bilalli, A., Ibrahimi, H., Musliu, M., Grapci-Kotori, L., Geci, D., Slavevska-Stamenkovic, V., Hinic, J., Mitic-Kopanja, D. & Keresztes, L. (2021). New records of the craneflies (Diptera: Limoniidae, Tipulidae) from the Western Balkans. *Journal of Entomological Research Society, 23*(2), 141-152. Doi: 10.51963/jers.v23i2.1929

Ferreira, S., Oosterbroek, P., Stary, J., Sousa, P., Mata, V.A., da Silva, L.P., Pauperio, J. & Beja, P. (2021). The InBIO Barcoding Initiative Database: DNA barcodes of Portuguese Diptera 02-Limoniidae, Pediciidae and Tipulidae. *Biodiversity Data Journal, 9*(4), p.e69841. doi: 10.3897/BDJ.9.e69841

Kolcsár, L., Ivković, M. & Ternjej, I. (2015). New records of Limoniidae and Pediciidae (Diptera) from Croatia. *ZooKeys, 513*(513), 23-37. Doi: 10.3897/zookeys.513.10066

Ozgul, O. & Koc, H. (2014). Four new species of Limoniidae (Diptera, Nematocera) from the Inner-West Anatolian Subregion of Turkey. *The Florida Entomologist, 97*(2), 620-625. doi: 10.1653/024.097.0238

Petersen, M.J. (2008). A systematic investigation of the crane fly subfamily Limoniinae (Tipuloidea: Limoniidae). Iowa State University. doi: 10.31274/rtd-180813-6702

Podenas, S. & Gelhaus, J. (2011). Three new species of Chioneinae crane flies (Diptera: Limoniidae) from North-Central Mongolia. *Proceedings of the Academy of Natural Sciences of Philadelphia, 161*(1), 73-86. doi: 10.1635/053.161.0105

Podeniene, V., Podenas, S. & Gelhaus, J.K. (2002). The first description of the larva of the crane fly genus *Gonempeda* Alexander, 1924 (Limoniidae: Chioneinae), with new information for understanding the phylogenetic relationships of the genus. *Proceedings of the Academy of Natural Sciences of Philadelphia, 152*(1), 67-73. doi: 10.1635/0097-3157(2002)152[0067:TFDOTL]2.0.CO2

Podeniene, V., Podenas, S. & Gelhaus, J.K. (2004). First record of a crane fly larva (Diptera, Limoniidae: Chioneinae) from Baltic amber. *Annals of the Entomological Society of America, 97*(6), 1126-1128. doi: 10.1603/0013-8746(2004)097[1126:FROACF]2.0.CO2

Suetsugu, K., Tetsu, S. & Sueyoshi, M. (2019). Fruit- and seed-feeding habit of the crane fly *Libnotes puella* (Diptera: Limoniidae) in Ryukyu Islands, Japan. *Entomological Science, 22*(4), 413-417. doi: 10.1111/ens.12387

Lonchopteridae

Stalker, H.D. (1956). On the evolution of parthenogenesis in *Lonchoptera* (Diptera). *Evolution, 10*(4), 345-359. doi: 10.1111/j.1558-5646.1956.tb02862.x

Micropezidae

Barnes, J.K. (2015). Biology and immature stages of *Compsobata univitta* (Walker, 1849) (Diptera: Micropezidae: Calobatinae). *Proceedings of the Entomological Society of Washington, 117*(4), 421-434. doi: 10.4289/0013-8797.117.4.421

Milichiidae

Giordani, G., Tuccia, F., Zoppis, S., Vecchiotti, C. & Vanin, S. (2019). Record of *Leptometopa latipes* (Diptera: Milichiidae) from a human cadaver in the Mediterranean area. *Forensic Sciences Research, 4*(4), 341-347. doi: 10.1080/20961790.2018.1490473

Xi, Y., Yang, D. & Yin, X. (2018). The genus *Phyllomyza* Fallén from China, with descriptions of three new species (Diptera, Milichiidae). *ZooKeys, 760*(760), 143-157. doi: 10.3897/zookeys.760.22595

Yusah, K.M. & Fayle, T.M. (2014). The first record of a fly of the family Milichiidae (Diptera) interacting with an ant of the genus *Polyrhachis* Smith, 1857 (Hymenoptera: Formicidae). *Biodiversity Data Journal, 2*(2), p.e4168. doi: 10.3897/BDJ.2.e4168

Muscidae

Achint, R. & Singh, D. (2021). Application of COI gene for identification of some economically and forensically important muscid flies of India (Diptera: Muscidae). *International Journal of Tropical Insect Science, 41*(4), 3023-3029. doi: 10.1007/s42690-021-00494-8

Baleba, S.B.S., Torto, B., Masiga, D., Weldon, C.W. & Getahun, M.N. (2019). Egg-laying decisions based on olfactory cues enhance offspring fitness in *Stomoxys calcitrans* L. (Diptera: Muscidae). *Scientific Reports, 9*(1), 3850. doi: 10.1038/s41598-019-40479-9

Bautista-Martínez, N., Illescas-Riquelme, C.P. & García-Ávila, C., de J. (2017). First report of "hunter-fly" *Coenosia attenuata* (Diptera: Muscidae) in Mexico. *The Florida Entomologist, 100*(1), 174-175. doi: 10.1653/024.100.0126

Butterworth, N.J. & Wallman, J.F. (2022). Flies getting filthy: The precopulatory mating behaviours of three mud-dwelling species of Australian *Lispe* (Diptera: Muscidae). *Ethology, 128*(4), 369-377. doi: 10.1111/eth.13236

Fowler, F.E. & Mullens, B.A. (2016). Dividing the pie: differential dung pat size utilization by sympatric *Haematobia irritans* and *Musca autumnalis*: Dung pat size use by horn flies and face flies. *Medical and Veterinary Entomology, 30*(2), 185-192. doi: 10.1111/mve.12166

Getahun, M.N., Ahuya, P., Ngiela, J., Orone, A., Masiga, D. & Torto, B. (2020). Shared volatile organic compounds between camel metabolic products elicits strong *Stomoxys calcitrans* attraction. *Scientific Reports, 10*(1), 21454. doi: 10.1038/s41598-020-78495-9

Grzywacz, A., Hall, M.J.R. & Pape, T. (2015). Morphology successfully separates third instar larvae of *Muscina. Medical and Veterinary Entomology, 29*(3), 314-329. doi: 10.1111/mve.12117

Gunn, A. (2016). The colonisation of remains by the muscid flies *Muscina stabulans* (Fallén) and *Muscina prolapsa* (Harris) (Diptera: Muscidae). *Forensic Science International, 266*, 349-356. doi: 10.1016/j.forsciint.2016.06.013

Hieu, T.T., Jung, J., Kim, S.., Ahn, Y. & Kwon, H.W. (2014). Behavioural and electroantennogram responses of the stable fly (*Stomoxys calcitrans* L.) to plant essential oils and their mixtures with attractants. *Pest Management Science, 70*(1), 163-172. doi: 10.1002/ps.3547

Ivkovic, M. & Pont, A.C. (2016). Long-time emergence patterns of *Limnophora* species (Diptera, Muscidae) in specific karst habitats: tufa barriers. *Limnologica, 61*, 29-35. doi: 10.1016/j.limno.2016.09.003

Krafsur, E.S. & Moon, R.D. (1997). Bionomics of the face fly, *Musca autumnalis*. *Annual Review of Entomology, 42*(1), 503-523. doi: 10.1146/annurev.ento.42.1.503

Krivosheina, N.P. (2013). On the ecology of *Phaonia* larvae (Diptera, Muscidae). *Entomological Review, 93*(3), 324-333. doi: 10.1134/S0013873813030068

Krivosheina, N.P. (2014). Morpho-ecological characteristic of immature stages of the genus *Mydaea* (Diptera, Muscidae). *Entomological Review, 94*(5), 675-686. doi: 10.1134/S0013873814050042

Krivosheina, N.P. (2019). Morpho-ecological characteristics of immature stages of Xytobiont species of the genus *Phaonia* Robineau-Desvoidy, 1830 (Muscidae, Diptera) inhabiting tree holes and sap accumulations. Entomological Review, 99(2), 250-261. doi: 10.1134/S0013873819020118

Kutty, S.N., Pont, A.C., Meier, R. & Pape, T. (2014). Complete tribal sampling reveals basal split in Muscidae (Diptera), confirms saprophagy as ancestral feeding mode, and reveals an evolutionary correlation between instar numbers and carnivory. *Molecular Phylogenetics and Evolution, 78*, 349-364. doi: 10.1016/j.ympev.2014.05.027

Liénard, E., Salem, A., Jacquiet, P., Grisez, C., Prévot, F., Blanchard, B., Bouhsira, E. & Franc, M. (1987). Development of a protocol testing the ability of *Stomoxys calcitrans* (Linnaeus, 1758) (Diptera:

Muscidae) to transmit *Besnoitia besnoiti* (Henry, 1913) (Apicomplexa: Sarcocystidae). *Parasitology Research, 112*(2), 479-486. doi: 10.1007/s00436-012-3157-6

Liu, G., Wang, Q., Liu, X., Li, X., Pang, X. & Zhang, D. (2021). Antennal and palpal sensilla of three predatory *Lispe* species (Diptera: Muscidae): an ultrastructural investigation. *Scientific Reports, 11*(1), 18357. doi: 10.1038/s41598-021-97677-7

Olafson, PU., Aksoy, S., Attardo, G.M., Buckmeier, G., Chen, X., Coates, C.J., Davis, M., Dykema, J., Emrich, S.J., Friedrich, M., Holmes, C.J., Ioannidis, P., Jansen, E.N., Jennings, E.C., Lawson, D., Martinson, E.O., Maslen, G.L., Meisel, R.P., Murphy, T.D., Nayduch, D., Nelson, D.R., Oyen, K.J., Raszick, T.J., Ribeiro, J.M.C., Robertson, H.M., Rosendale, A.J., Sackton, T.B., Saelao, P., Swiger, S.L., Sze, S., Tarone, A.M., Taylor, D.B., Warren, W.C., Waterhouse, R.M., Weirauch, M.T., Werren, J.H., Wilson, R.K., Zdobnov, E.M. & Benoit, J.B. (2021). The genome of the stable fly, *Stomoxys calcitrans*, reveals potential mechanisms underlying reproduction, host interactions, and novel targets for pest control. *BMC Biology, 19*(1), 41. doi: 10.1186/s12915-021-00975-9

Olesen, A.S. Lohse, L., Hansen, M.F., Boklund, A., Halasa, T., Belsham, G.J., Rasmussen, T.B., Bøtner, A. & Bødker, R. (2018). Infection of pigs with African swine fever virus via ingestion of stable flies (*Stomoxys calcitrans*). *Transboundary and Emerging Diseases, 65*(5), 1152-1157. doi: 10.1111/tbed.12918

Phillips, H.N., Moon, R.D., Sorge, U.S. & Heins, B.J. (2020). Efficacy of broilers as a method of face fly (*Musca autumnalis* De Geer) larva control for organic dairy production. *Animals, 10*(12), 2429. doi: 10.3390/ani10122429

Pont, A.C. & Ivković, M. (2013). The Hunter-flies of Croatia (Diptera: Muscidae: genus *Limnophora* Robineau-Desvoidy). *Journal of Natural History, 47*(15-16), 1069-1082. doi: 10.1080/00222933.2012.750775

Rotheray, G.E. & Wilkinson, G. (2015). Trophic structure and function in the larvae of predatory muscid flies (Diptera, Muscidae). Zoomorphology, 134(4), 553-563. doi: 10.1007/s00435-015-0284-5

Savage, J. & Sorokina, V.S. (2021). Review of the North American fauna of *Drymeia* Meigen (Diptera, Muscidae) and evaluation of DNA barcodes for species-level identification in the genus. *ZooKeys, 1024*(5), 31-89. doi: 10.3897/zookeys.1024.60393

Savage, J., Wheeler, T.A. & Wiegmann, B.M. (2004). Phylogenetic analysis of the genus *Thricops rondani* (Diptera: Muscidae) based on molecular and morphological characters. *Systematic Entomology, 29*(3), 395-414. doi: 10.1111/j.0307-6970.2004.00252.x

Seabra, S.G., Martins, J., Bras, P., Tavares, A.M., Freitas, I., Barata, A., Rebelo, M.T., Mateus, C., Paulo, O.S. & Figueiredo, E. (2021). PCR-based detection of prey DNA in the gut contents of the tiger-fly, *Coenosia attenuata* (Diptera: Muscidae), a biological control agent in Mediterranean greenhouses. *European Journal of Entomology, 118*(1), 335-343. doi: 10.14411/eje.2021.035

Sommer, C., Jensen, K.-M.V. & Jespersen, J.B. (2001). Topical treatment of calves with synthetic pyrethroids: effects on the non-target dung fly *Neomyia cornicina* (Diptera: Muscidae). *Bulletin of Entomological Research, 91*(2), 131-137. doi: 10.1079/BER200079

Sorokina, V.S. & Ovtshinnikova, O.G. (2020). The position of the Azeliinae in the Muscidae (Diptera) based on musculature of the male terminalia. *ZooKeys, 975*(975), 87-110. doi: 10.3897/zookeys.975.55502

Sowig, P., Himmelsbach, R. & Himmelsbach, W. (1997). Predator-prey relationship between insect larvae: growth of *Sphaeridium* larvae (Coleoptera: Hydrophilidae) under time constraints through predation on *Musca autumnalis* maggots (Diptera: Muscidae). *Canadian Journal of Zoology, 75*(12), 2069-2076. doi: 10.1139/z97-841

Xue, W., Zhang, L. & Wang, M. (2008). A key to the species of *Spilogona* Schnabl from China, with descriptions of four new species from the Qinghai-Xizang (Tibetan) Plateau (Diptera: Muscidae). *Entomological Science, 11*(1), 87-95. doi: 10.1111/j.1479-8298.2007.00257.x

Vikhrev, N.E. (2011). Review of the Palaearctic members of the *Lispe tentaculata* species-group (Diptera, Muscidae): revised key, synonymy and notes on ecology. *ZooKeys, 84*(84), 59-70. doi: 10.3897/zookeys.84.819

Vikhrev, N.E. (2012). Revision of the *Lispe longicollis*-group (Diptera, Muscidae). *ZooKeys, 235*(235), 23-39. doi: 10.3897/zookeys.235.3306

Wang, Y., Zhang, Y., Wang, M., Hu, G., Fu, Y., Zhi, R., Wang, J. & Lysyk, T. (2020). Development of *Hydrotaea spinigera* (Diptera: Muscidae) at constant temperatures and its significance for estimating postmortem interval. *Journal of Medical Entomology, 58*(1), 56-63. doi: 10.1093/jme/tjaa162

Zou, D., Coudron, T.A., Xu, W., Xu, J. & Wu, H. (2020). Performance of the tiger-fly *Coenosia attenuata* Stein reared on the alternative prey, *Chironomus plumosus* (L.) larvae in coir substrate. *Phytoparasitica, 49*(1), 83-92. doi: 10.1007/s12600-020-00866-9

Zou, D., Coudron, T.A., Zhang, L., Xu, W., Xu, J., Wang, M., Xiao, X. & Wu, H. (2021). Effect of prey species and prey densities on the performance of adult *Coenosia attenuata. Insects, 12*(8), 669. doi: 10.3390/insects12080669

Mycetophilidae

Bauernfeind, R., Schneeberg, K. & Beutel, R.G. (2015). The larval head of *Exechia* (Mycetophilidae) and *Bibio* (Bibionidae) (Diptera). *Arthropod Structure and Development, 44*(4), 326-345. doi: 10.1016/j.asd.2015.04.005

Bouju, V., Rosse-Guillevic, S., Griffon, M., Bojarski, B., Szwedo, J. & Perrichot, V. (2021). The genus *Allodia* (Diptera: Mycetophilidae) in Miocene Ethiopian amber. *Mitteilungen aus dem Museum für Naturkunde in Berlin, 24*(2), 339-346. doi: 10.5194/fr-24-339-2021

Jürgenstein, S., Kurina, O. & Põldmaa, K. (2015). The *Mycetophila ruficollis* Meigen (Diptera, Mycetophilidae) group in Europe: elucidating species delimitation with COI and ITS2 sequence data. *ZooKeys, 508*(508), 15-51. doi: 10.3897/zookeys.508.9814

Kaspřák, D., Kerr, P., Sýkora, V., Tóthová, A. & Ševčík, J. (2019). Molecular phylogeny of the fungus gnat subfamilies Gnoristinae and Mycomyinae, and their position within Mycetophilidae (Diptera): Molecular phylogeny of Gnoristinae and Mycomyinae. *Systematic Entomology, 44*(1), 128-138. doi: 10.1111/syen.12312

Kurina, O. (2021). A contribution towards checklist of fungus gnats (Diptera, Diadocidiidae, Ditomyiidae, Bolitophilidae, Keroplatidae, Mycetophilidae) in Georgia, Transcaucasia. *ZooKeys, 1026*(1026), 69-142. doi: 10.3897/zookeys.1026.63749

Martinsson, S., Kjærandsen, J. & Sundberg, P. (2011). Towards a molecular phylogeny of the fungus gnat genus *Boletina* (Diptera: Mycetophilidae). *Zoologica Scripta, 40*(3), 272-281. doi: 10.1111/j.1463-6409.2011.00474.x

Opomyzidae

Roháček, J. (2012). The fauna of the opomyzoid families Clusiidae, Acartophthalmidae, Anthomyzidae, Opomyzidae, Stenomicridae, Periscelididae, Asteiidae (Diptera) in the Gemer area (Central Slovakia). *Časopis Slezského Zemského Muzea, 61*(2), 97-111. doi: 10.2478/v10210-012-0011-5

Oestridae

Martín-Vega, D., Clark, B., Ferrer, L. M., López-Tamayo, S., Panadero, R., Cepeda-Palacios, R., Colwell, D. D. & Hall, M. J. R. (2021). Major differences in the larval anatomy of the digestive and excretory systems of three Oestridae species revealed by micro-CT. *Medical and Veterinary Entomology, 35*(1), 106-120. doi: 10.1111/mve.12476

Pediciidae

Ferreira, S., Oosterbroek, P., Stary, J., Sousa, P., Mata, V.A., da Silva, L.P., Pauperio, J. & Beja, P. (2021). The InBIO Barcoding Initiative Database: DNA barcodes of Portuguese Diptera 02-Limoniidae, Pediciidae and Tipulidae. *Biodiversity Data Journal*, *9*(4), p.e69841. doi: 10.3897/BDJ.9.e69841

Krivosheina, N.P. (2011). New data on the larval morphology of limoniid flies of the genus *Ula* (Diptera, Pediciidae). *Entomological Review, 91*(4), 432-443. doi: 10.1134/S001387381104004X

Phoridae

Castillo-Alanis, L.A., González-Hernández, A.E., Quijano-Mateos, A., Pedraza-Lara, C.S., Villavicencio-Queijeiro, A. & Bravo-Gómez, M.E. (2020). Standardization of a culture medium for *Megaselia scalaris* (Diptera: Phoridae) for entomotoxicological studies. *Journal of Medical Entomology, 57*(5), 1421-1431. doi: 10.1093/jme/tjaa088

Disney, R.H.L. (2007). Natural history of the scuttle fly, *Megaselia scalaris*. *Annual Review of Entomology, 53*(1), 39-60. doi: 10.1146/annurev.ento.53.103106.093415

Feng, D. & Liu, G. (2012). A study on morphology of immature stages of *Diplonevra peregrina* (Wiedemann) (Diptera: Phoridae). *Microscopy Research and Technique, 75*(10), 1425-1431. doi: 10.1002/jemt.22085

Giangaspero, A., Barlaam, A., Pane, S., Marchili, M.R., Muda, A.O., Putignani, L. & Hall, M.J.R. (2020). Accidental nasal Myiasis caused by *Megaselia rufipes* (Diptera: Phoridae) in a child. *Journal of Medical Entomology, 58*(1), 121-124. doi: 10.1093/jme/tjaa184

Pennington, M.J., Rothman, J.A., Jones, M.B., McFrederick, Q.S., Gan, J. & Trumble, J.T. (2017). Effects of contaminants of emerging concern on *Megaselia scalaris* (Lowe, Diptera: Phoridae) and its microbial community. *Scientific Reports, 7*(1), 1-12. doi: 10.1038/s41598-017-08683-7

Sorensen, M.A., Chase-Dunn, C.M. & Trumble, J.T. (2009). Chronic exposure to elevated levels of manganese and nickel is not harmful to a cosmopolitan detritivore, Megaselia scalaris (Diptera: Phoridae). *Insect Science, 16*(1), 73-79. doi: 10.1111/j.1744-7917.2009.00256.x

Stoaks, R. (2009). Rare and unusual arthropod ecology, behavior and occurrence at hummingbird feeders in northern California. *The Pan-Pacific Entomologist, 85*(4), 194-198. doi: 10.3956/2009-18.1

Piophilidae

Martín-Vega, D. & Baz, A. (2011). Could the 'vulture restaurants' be a lifeboat for the recently rediscovered bone-skippers (Diptera: Piophilidae)? *Journal of Insect Conservation, 15*(5), 747-753. doi: 10.1007/s10841-011-9429-0

Oudin, M.J., Bonduriansky, R. & Rundle, H.D. (2015). Experimental evidence of condition-dependent sexual dimorphism in the weakly dimorphic antler fly *Protopiophila litigata* (Diptera: Piophilidae). *Biological Journal of the Linnean Society, 116*(1), 211-220. doi: 10.1111/bij.12549

Taleb, M., Tail, G. & Açıkgöz, H.N. (2019). DNA barcoding of *Stearibia nigriceps* (Meigen) and *Piophila casei* (Linnaeus) (Diptera: Piophilidae) from Algeria and the first African report of *Stearibia nigriceps. International Journal of Legal Medicine, 134*(3), 895-902. doi: 10.1007/s00414-019-02223-w

Pipunculidae

Jervis, M.A. (1992). A taxonomic revision of the pipunculid fly genus *Chalarus* Walker, with particular reference to the European fauna. *Zoological Journal of the Linnean Society, 105*(3), 243-352. doi: 10.1111/j.1096-3642.1992.tb01232.x

Kehlmaier, C. (2014). Checklist of the family Pipunculidae (Diptera) of Finland. *ZooKeys, 441*(441), 225-232. doi: 10.3897/zookeys.441.7278

Molinatto, G., Demichelis, S., Bodino, N., Giorgini, M., Mori, N. & Bosco, D. (2020). Biology and prevalence in Northern Italy of *Verrallia aucta* (Diptera, Pipunculidae), a parasitoid of *Philaenus*

spumarius (Hemiptera, Aphrophoridae), the main vector of *Xylella fastidiosa* in Europe. *Insects, 11*(9), 607. doi: 10.3390/insects11090607

Platystomatidae

McMichael, B.A., Foote, B.A. & Bowker, B.D. (1990). Biology of *Rivellia melliginis* (Diptera: Platystomatidae), a consumer of the nitrogen-fixing root nodules of black locust (Leguminosae). *Annals of the Entomological Society of America, 83*(5), 967-974. doi: 10.1093/aesa/83.5.967

Polleniidae

Taleb, M., Tail, G. & Açıkgöz, H.N. (2022). Molecular identification of the potentially forensically relevant cluster flies *Pollenia rudis* (Fabricius) and *Pollenia vagabunda* (Meigen) (Diptera: Polleniidae) - non-recorded species in Algeria. *Forensic Sciences Research, 7*(1), 69-77. doi: 10.1080/20961790.2020.1857937

Vezsenyi, K.A., Langer, S.V., Samkari, B.A. & Beresford, D.V. (2022). The history and current state of cluster flies (Diptera: Polleniidae: *Pollenia*) in North America, with new Canadian provincial records. *Canadian Entomologist, 154*(1). doi: 10.4039/tce.2022.11

Psilidae

Degen, T., Städler, E. & Ellis, P.R. (1999). Host-plant susceptibility to the carrot fly, *Psila rosae*. 2. Suitability of various host species for larval development. *Annals of Applied Biology, 134*(1), 13-26. doi: 10.1111/j.1744-7348.1999.tb05231.x

Städler, E. (2012). An improved mass-rearing method of the carrot rust fly, *Psila rosae* (Diptera: Psilidae). *The Canadian Entomologist, 103*(7), 1033-1038. doi: 10.4039/Ent1031033-7

Psychodidae

Ali, A. & Kok-Yokomi, M.L. (1991). Preliminary population assessment of *Psychoda alternata* (Diptera: Psychodidae) in soil irrigated with wastewater for turf cultivation. *The Florida Entomologist, 74*(4), 591-596. doi: 10.2307/3495415

Azmiera, N., Low, V.L. & Heo, C.C. (2021). Colonization of rabbit carcasses by drain fly larvae, *Psychoda* sp. (Diptera: Psychodidae): The first report. *Acta Parasitologica, 66*(2), 706-709. doi: 10.1007/s11686-020-00313-z

Ptychopteridae

Wolf, B., Zwick, P. & Marxsen, J. (1997). Feeding ecology of the freshwater detritivore *Ptychoptera paludosa* (Diptera, Nematocera). *Freshwater Biology, 38*(2), 375-386. doi: 10.1046/j.1365-2427.1997.00250.x

Rhagionidae

Imada, Y., Kato, M. & Brady, S. (2016). Bryophyte-Feeders in a basal Brachyceran lineage (Diptera: Rhagionidae: Spaniinae): Adult oviposition behavior and changes in the larval mouthpart morphology accompanied with the diet shifts. *PLoS One, 11*(11), p.e0165808. doi: 10.1371/journal.pone.0165808

Roberts, M.J. (1987). Structure of the mouthparts of the larvae of the flies *Rhagio* and *Sargus* in relation to feeding habits. *Journal of Zoology, 159*(3), 381-398. doi: 10.1111/j.1469-7998.1969.tb08453.x

Sarcophagidae

Blanckenhorn, W.U., Pemberton, A.J., Bussière, L.F., Roembke, J. & Floate, K.D. (2010). A review of the natural history and laboratory culture methods for the yellow dung fly, *Scathophaga stercoraria. Journal of Insect Science, 10*(11), 1-17. doi: 10.1673/031.010.1101

Buenaventura, E. (2021). Museomics and phylogenomics with protein-encoding ultraconserved elements illuminate the evolution of life history and phallic morphology of flesh flies (Diptera: Sarcophagidae). *BMC Ecology and Evolution, 21*(1), 70. doi: 10.1186/s12862-021-01797-7

Buenaventura, E., Szpila, K., Cassel, B.k., Wiegmann, B.M. & Pape, T. (2020). Anchored hybrid enrichment challenges the traditional classification of flesh flies (Diptera: Sarcophagidae). *Systematic Entomology, 45*(2), 281-301. doi: 10.1111/syen.12395

Buenaventura, E., Whitmore, D. & Pape, T. (2017). Molecular phylogeny of the hyperdiverse genus *Sarcophaga* (Diptera: Sarcophagidae), and comparison between algorithms for identification of rogue taxa. *Cladistics, 33*(2), 109-133. doi: 10.1111/cla.12161

Danyk, T., Mackauer, M. & Johnson, D.L. (2005). The influence of host suitability on the range of grasshopper species utilized by *Blaesoxipha atlanis* (Diptera: Sarcophagidae) in the field. *Bulletin of Entomological Research, 95*(6), 571-578. doi: 10.1079/BER2005388

Dufek, M.I., Mello-Patiu, C.A. & Mulieri, P.R. (2020). Inventory of Sarcophaginae (Diptera: Sarcophagidae) for the Humid Chaco, a poorly surveyed ecoregion of South America. *Journal of Natural History, 54*(5-6), 367-403. doi: 10.1080/00222933.2020.1764646

Miura, K. & Ohsaki, N. (2013). Mortality effects of the parasitoid flesh fly *Blaesoxipha japonensis* (Diptera: Sarcophagidae) in relation to body size of the adult grasshopper *Parapodisma tanbaensis* (Orthoptera: Catantopidae). *Applied Entomology and Zoology, 49*(1), 171-176. doi: 10.1007/s13355-013-0236-9

Miura, K. & Ohsaki, N. (2014). The cost of autotomy caused by the parasitoid fly *Blaesoxipha japonensis* (Diptera: Sarcophagidae): an interspecific comparison between two sympatric grasshopper host species. *Ecological Research, 30*(1), 33-39. doi: 10.1007/s11284-014-1202-x

Otronen, M. (1995). Energy reserves and mating success in males of the yellow dung fly, *Scathophaga stercoraria. Functional Ecology, 9*(4), 683-688. doi: 10.2307/2390161

Rees, N.E. (1970). Suitability of selected North American grasshopper species as hosts for the Eurasian parasites *Acridomyia sacharovi1* and *Blaesoxipha lineata2* (Diptera). *Annals of the Entomological Society of America, 63*(3), 901-903. doi: 10.1093/aesa/63.3.901

Ren, L., Shang, Y., Yang, L., Wang, S., Wang, X., Chen, S., Bao, Z., An, D., Meng, F., Cai, J. & Guo, Y. (2021). Chromosome-level de novo genome assembly of *Sarcophaga peregrina* provides insights into the evolutionary adaptation of flesh flies. *Molecular Ecology Resources, 21*(1), 251-262. doi: 10.1111/1755-0998.13246

Souza, C.M., Madeira-Ott, T., Masiero, F.S., Bunde, P.R.S., Ribeiro, G.A., Thyssen, P.J. & Geden, C. (2020). Synanthropy of Sarcophaginae (Diptera: Sarcophagidae) from Southern Brazil and its sanitary implications. *Journal of Medical Entomology, 58*(2), 913-920. doi: 10.1093/jme/tjaa243

Stamper, T., Dahlem, G.A., Cookman, C. & Debry, R.W. (2013). Phylogenetic relationships of flesh flies in the subfamily Sarcophaginae based on three mtDNA fragments (Diptera: Sarcophagidae). *Systematic Entomology, 38*(1), 35-44. doi: 10.1111/j.1365-3113.2012.00646.x

Valverde Castro, C., Buenaventura, E., Sánchez-Rodríguez, J.D. & Wolff, M. (2017). Flesh flies (Diptera: Sarcophagidae: Sarcophaginae) from the Colombian Guajira biogeographic province, an approach to their ecology and distribution. *Zoologia, 34,* 1-11. doi: 10.3897/zoologia.34.e12277

Scathophagidae

Blanckenhorn, W.U., Pemberton, A.J., Bussière, L.F., Roembke, J. & Floate, K.D. (2010). A review of the natural history and laboratory culture methods for the yellow dung fly, *Scathophaga stercoraria. Journal of Insect Science, 10*(11), 1-17. doi: 10.1673/031.010.1101

Kutty, S.N., Bernasconi, M.V., Šifner, F. & Meier, R. (2007). Sensitivity analysis, molecular systematics and natural history evolution of Scathophagidae (Diptera: Cyclorrhapha: Calyptratae). *Cladistics, 23*(1), 64-83. doi: 10.1111/j.1096-0031.2006.00131.x

Ovchinnikov, A.N. & Ovtshinnikova, O.G. (2013). The muscular system of the ovipositor of *Spaziphora hydromyzina* (Fallen) (Diptera, Scathophagidae). *Entomological Review*, *93*(8), 987-990. doi: 10.1134/S001387381308006X

Ozerov, A.L. (2017). A new species of the genus *Scathophaga* Meigen, 1803 (Diptera, Scathophagidae) from Yakutia (Russia). *Entomological Review, 97*(1), 132-135. doi: 10.1134/S0013873817010134

Rohner, P.T., Blanckenhorn, W.U. & Schäfer, M.A. (2017). Critical weight mediates sex-specific body size plasticity and sexual dimorphism in the yellow dung fly *Scathophaga stercoraria* (Diptera: Scathophagidae). *Evolution & Development, 19*(3), 147-156. doi: 10.1111/ede.12223

Webb, L., Beaumont, D.J., Nager, R.G. & McCracken, D.I. (2007). Effects of avermectin residues in cattle dung on yellow dung fly *Scathophaga stercoraria* (Diptera: Scathophagidae) populations in grazed pastures. *Bulletin of Entomological Research, 97*(2), 129-138. doi: 10.1017/S0007485307004683

Sciaridae

Badshah, K., Ullah, F., Ahmad, B., Ahmad, S., Alam, S., Ullah, M., Jamil, M. & Sardar, S. (2021). Management of *Lycoriella ingenua* (Diptera: Sciaridae) on oyster mushroom (*Pleurotus ostreatus*) through different botanicals. *International Journal of Tropical Insect Science, 41*(2), 1435-1440. doi: 10.1007/s42690-020-00341-2

Cloonan, K.R., Andreadis, S.S. & Baker, T.C. (2016). Attraction of female fungus gnats, *Lycoriella ingenua*, to mushroom-growing substrates and the green mold *Trichoderma aggressivum. Entomologia Experimentalis et Applicata, 159*(3), 298-304. doi: 10.1111/eea.12439

Cloonan, K.R., Andreadis, S.S. & Baker, T.C. (2019). Little effect of delayed mating on fecundity or fertility of female fungus gnats *Lycoriella ingenua*. *Physiological Entomology, 44*(1), 60-64. doi: 10.1111/phen.12276

Gou, Y., Guo, S., Wang, G. & Liu, C. (1986). Effects of short-term heat stress on the growth and development of *Bradysia cellarum* and *Bradysia impatiens. Journal of Applied Entomology, 144*(4), 315-321. doi: 10.1111/jen.12733

Gou, Y., Quandahor, P., Zhang, Y., Coulter, J.A., Liu, C. & Gao, Y. (2020). Host plant nutrient contents influence nutrient contents in *Bradysia cellarum* and *Bradysia impatiens. PLoS One, 15*(4), p.e0226471. doi: 10.1371/journal.pone.0226471

Heiri, O. & Lotter, A.F. (2007). Sciaridae in lake sediments: indicators of catchment and stream contribution to fossil insect assemblages. *Journal of Paleolimnology, 38*(2), 183-189. doi: 10.1007/s10933-006-9068-8

Kecskeméti, S., Fail, J. & Geösel, A. (2018). Development of *Lycoriella ingenua* and *Bradysia impatiens* on different phases of agaricus composts. *Review on Agriculture and Rural Development, 7*(1-2), 55.

Kecskeméti, S., Szelényi, M.O., Erdei, A.L., Geösel, A., Fail, J. & Molnár, B.P. (2020). Fungal volatiles as olfactory cues for female fungus gnat, *Lycoriella ingenua* in the avoidance of Mycelia colonized compost. *Journal of Chemical Ecology, 46*(10), 917-926. doi: 10.1007/s10886-020-01210-5

Sawangproh, W. & Cronberg, N. (2016). Life history traits of the liverwort herbivore *Scatopsciara cunicularius* (Diptera: Sciaridae). *Annals of the Entomological Society of America, 109*(3), 343-349. doi: 10.1093/aesa/saw004

Sawangproh, W., Ekroos, J. & Cronberg, N. (2016). The effect of ambient temperature on larvae of *Scatopsciara cunicularius* (Diptera: Sciaridae) feeding on the thallose liverwort Marchantia polymorpha. *European Journal of Entomology, 113*(1), 259-264. doi: 10.14411/eje.2016.030

Shamshad, A. (2010). The development of integrated pest management for the control of mushroom sciarid flies, *Lycoriella ingenua* (Dufour) and *Bradysia ocellaris* (Comstock), in cultivated mushrooms. *Pest Management Science, 66*(10), 1063-1074. doi: 10.1002/ps.1987

Shin, S., Jung, S., Heller, K., Menzel, F., Hong, T. K., Shin, J. S., Lee, S. H., Lee, H. & Lee, S. (1986). DNA barcoding of *Bradysia* (Diptera: Sciaridae) for detection of the immature stages on agricultural crops. *Journal of Applied Entomology, 139*(8), 638-645. doi: 10.1111/jen.12198

Shin, S., Jung, S., Menzel, F., Heller, K., Lee, H. & Lee, S. (2013). Molecular phylogeny of black fungus gnats (Diptera: Sciaroidea: Sciaridae) and the evolution of larval habitats. *Molecular Phylogenetics and Evolution, 66*(3), 833-846. doi: 10.1016/j.ympev.2012.11.008

Vilkamaa, P., Hippa, H. & Taylor, S.J. (2011). The genus *Camptochaeta* in Nearctic caves, with the description of *C. prolixa* sp. n. (Diptera, Sciaridae). *ZooKeys, 135*(135), 69-75. doi: 10.3897/zookeys.135.1624

Xi, L., Zhou, X., Liu, Y., Li, Y., Shi, H., Wu, H. & Huang, J. (1986). Behavioural response of the fungus gnat, *Bradysia impatiens* (Diptera: Sciaridae) towards certain edible mushrooms and saprophytic fungi. *Journal of Applied Entomology, 145*(5), 458-466. doi: 10.1111/jen.12857

Yan, H., Zhang, B., Wang, E., Xu, X. & Wei, G. (2021). Combining predatory mites and film mulching to control *Bradysia cellarum* (Diptera: Sciaridae) on Chinese chives, *Allium tuberosum. Experimental & Applied Acarology, 86*(1), 117-127. doi: 10.1007/s10493-021-00681-9

Zhu, G., Luo, Y., Xue, M., Zhao, H., Sun, X. & Wang, X. (2017). Effects of feeding on different host plants and diets on *Bradysia odoriphaga* population parameters and tolerance to heat and insecticides. *Journal of Economic Entomology, 110*(6), 2371-2380. doi: 10.1093/jee/tox242

Sciomyzidae

Bistline-East, A., Carey, J.G.J., Colton, A., Day, M.F. & Gormally, M.J. (2018). Catching flies with honey(dew): adult marsh flies (Diptera: Sciomyzidae) utilize sugary secretions for high-carbohydrate diets. *Environmental Entomology, 47*(6), 1632-1641. doi: 10.1093/ee/nvy155

Chapman, E.G., Foote, B.A., Malukiewicz, J. & Hoeh, W.R. (2006). Parallel evolution of larval morphology and habitat in the snail-killing fly genus *Tetanocera. Journal of Evolutionary Biology, 19*(5), 1459-1474. doi: 10.1111/j.1420-9101.2006.01132.x

Chapman, E.G., Przhiboro, A.A., Harwood, J.D., Foote, B.A. & Hoeh, W.R. (2012). Widespread and persistent invasions of terrestrial habitats coincident with larval feeding behavior transitions during snail-killing fly evolution (Diptera: Sciomyzidae). *BMC Evolutionary Biology, 12*(1), 175. doi: 10.1186/1471-2148-12-175

Murphy, W.L., Knutson, L.V., Chapman, E.G. Mc Donnell, R.J., Williams, C.D., Foote, B.A. & Vala, J. (2012). Key aspects of the biology of snail-killing Sciomyzidae flies. *Annual Review of Entomology, 57*(1), 425-447. doi: 10.1146/annurev-ento-120710-100702

Sepsidae

Rohner, P.T., Haenni, J., Giesen, A., Busso, J.P., Schäfer, M.A., Püchel-Wieling, F. & Blanckenhorn, W.U. (2019). Temporal niche partitioning of Swiss black scavenger flies in relation to season and substrate age (Diptera, Sepsidae). *Alpine Entomology, 3,* 1-10. doi: 10.3897/alpento.3.28366

Roy, J., Blanckenhorn, W.U. & Rohner, P.T. (2018). Largely flat latitudinal life history clines in the dung fly *Sepsis fulgens* across Europe (Diptera: Sepsidae). *Oecologia, 187*(3), 851-862. doi: 10.1007/s00442-018-4166-7

Zeender, V., Roy, J., Wegmann, A., Schäfer, M.A., Gourgoulianni, N., Blanckenhorn, W.U. & Rohner, P.T. (2019). Comparative reproductive dormancy differentiation in European black scavenger flies (Diptera: Sepsidae). *Oecologia, 189*(4), 905-917. doi: 10.1007/s00442-019-04378-0

Simuliidae

Adler, P.H., Cheke, R.A., Post, R.J. (2010). Evolution, epidemiology, and population genetics of black flies (Diptera: Simuliidae). *Infection, genetics and evolution, 10*(7), 846-865. doi: 10.1016/j.meegid.2010.07.003

Adler, P.H., Huang, Y., Reeves, W.K., Kim, S.K., Otsuka, Y., Takaoka, H. & Laudet, V. (2013). Macrogenomic evidence for the origin of the black fly *Simulium suzukii* (Diptera: Simuliidae) on Okinawa Island, Japan. *PLoS One, 8*(8), p.e70765. doi: 10.1371/journal.pone.0070765

Al-Shaer, L., Pierce, A.K., Larson, D. & Hancock, R. (2015). Notes on facultative predation in *Prosimulium* larvae (Diptera: Simuliidae) in alpine and subalpine streams in Colorado. *Journal of the American Mosquito Control Association, 31*(1), 113-116. doi: 10.2987/14-6460.1

Brabrand, Å., Bremnes, T., Koestler, A. G., Marthinsen, G., Pavels, H., Rindal, E., Raastad, J. E., Saltveit, S. J. & Johnsen, A. (2014). Mass occurrence of bloodsucking blackflies in a regulated river reach: localization of oviposition habitat of *Simulium truncatum* using DNA barcoding: mass occurrence bloodsucking blackflies. *River Research and Applications, 30*(5), 602-608. doi: 10.1002/rra.2669

Finn, D.S. & Alder, P.H. (2006). Population genetic structure of a rare high-elevation black fly, *Metacnephia coloradensis*, occupying Colorado lake outlet streams. *Freshwater Biology, 51*(12), 2240-2251. doi: 10.1111/j.1365-2427.2006.01647.x

Gaudreau, C., Larue, B. & Charpentier, G. (2010). Molecular comparison of Quebec and Newfoundland populations of the blackfly, *Simulium vittatum*, species complex. *Medical and Veterinary Entomology, 24*(2), 214-217. doi: 10.1111/j.1365-2915.2009.00844.

Hall, R.J., Bailey, R. C. & Findeis, J. (1988). Factors affecting survival and cation concentration in the blackflies *Prosimulium fuscum/mixtum* and the mayfly *Leptophlebia cupida* during spring snowmelt. *Canadian Journal of Fisheries and Aquatic Sciences, 45*(12), 2123. doi: 10.1139/f88-247

Hunter, F.F., Sutcliffe, J.F. & Downe, A.E.R. (1993). Blood-feeding host preferences of the isomorphic species *Simulium venustum* and *S. truncatum. Medical and Veterinary Entomology, 7*(2), 105-110.

Iburg, J.P., Gray, E.W. & Noblet, R. (2016). Feeding behavior of *Simulium vittatum* larvae in response to various suspended materials: implications for control using biological larvicides. *Entomologia Experimentalis et Applicata, 159*(1), 70-76. doi: 10.1111/eea.12414

Karjalainen, A.K., Salmelin, J., Dimock, B. & Hintelmann, H. (2020). Blackfly larvae (*Simulium* spp.) can intensify methylmercury biomagnification in boreal food webs. *Water, Air, and Soil Pollution, 231*(1). doi: 10.1007/s11270-020-04717-5

Malmqvist, B. (1999). Life history of *Metacnephia lyra*, a blackfly highly characteristic of large Swedish river rapids at the time of maximum discharge (Diptera: Simuliidae). *Aquatic Insects, 21*(2), 89-98. doi: 10.1076/aqin.21.2.89.4534

Martin, P.J.S. & Edman, J.D. (1993). Assimilation rates of different particulate foods for *Simulium verecundum* (Diptera: Simuliidae). *Journal of Medical Entomology, 30*(4), 805-809. doi: 10.1093/jmedent/30.4.805

McCreadie, J.W. & Colbo, M.H. (1991). Spatial distribution patterns of larval cytotypes of the *Simulium venustum/verecundum* complex (Diptera: Simuliidae) on the Avalon Peninsula, Newfoundland: factors associated with occurrence. *Canadian Journal of Zoology, 69*(10), 2651-2659. doi: 10.1139/z91-373

McCreadie, J.W. & Colbo, M.H. (1993). Larval and pupal microhabitat selection by *Simulium truncatum* Lundström, *S. rostratum* Lundström, and *S. verecundum* AA (Diptera: Simuliidae). *Canadian Journal of Zoology, 71*(2), 358-367. doi: 10.1139/z93-050

McCreadie, J.W. & Colbo, M.H. (1993). Seasonal succession and spatial-temporal distribution patterns of six larval cytospecies of the *Simulium venustum/verecundum* complex (Diptera: Simuliidae). *Canadian Journal of Zoology, 71*(1), 116-124. doi: 10.1139/z93-017

McCreadie, J.W., Williams, R.H., Stutsman, S., Finn, D.S. & Adler, P.H. (2018). The influence of habitat heterogeneity and latitude on gamma diversity of the Nearctic Simuliidae, a ubiquitous group of stream-dwelling insects. *Insect Science, 25*(4), 712-720. doi: 10.1111/1744-7917.12442

Pangjanda, S. & Pramual, P. (2016). Trait-based and phylogenetic community ecology of black flies (Diptera: Simuliidae) in tropical streams of Thailand. *Hydrobiologia, 763*(1), 345-356. doi: 10.1007/s10750-015-2389-8

Sitarz, M., Buczek, A.M., Buczek, W., Buczek, A. & Bartosik, K. (2022). Risk of attacks by blackflies (Diptera: Simuliidae) and occurrence of severe skin symptoms in bitten patients along the Eastern border of the European Union. *International Journal of Environmental Research and Public Health, 19*(13), 7610. doi: 10.3390/ijerph19137610

Ya'cob, Z., Low, V.L., Tan, T.K., Noor-Izwan, A., Lourdes, E.Y., Ramli, R., Takaoka, H. & Adler, P.H. (1987). Sexually anomalous individuals of the black fly *Simulium trangense* (Diptera: Simuliidae) infected with mermithid parasites (Nematoda: Mermithidae). *Parasitology Research, 120*(5), 1555-1561. doi: 10.1007/s00436-021-07087-x

Zhang, Y. (2006). Balancing food availability and hydrodynamic constraint: phenotypic plasticity and growth in *Simulium noelleri* blackfly larvae. *Oecologia, 147*(1), 39-46. doi: 10.1007/s00442-005-0243-9

Zettler, J.A., Adler, P.H. & McCreadie, J.W. (1998). Factors influencing larval color in the *Simulium vittatum* Complex (Diptera: Simuliidae). *Invertebrate Biology, 117*(3), 245-252. doi: 10.2307/3226989

Sphaeroceridae

Marshall, S.A. (1982). A revision of the Nearctic *Leptocera (Thoracochaeta Duda)* (Diptera: Sphaeroceridae). *Canadian Entomologist, 114*(1), 63-78. doi: 10.4039/Ent11463-1

Pitkin, B.R. (1986). Bait, habitat preferences and the phenology of some lesser dung flies (Diptera: Sphaeroceridae) in Britain. *Journal of Natural History, 20*(6), 1283-1295. doi: 10.1080/00222938600770851

Stratiomyidae

Fontenelle, J.C.R., Viana-Silva, F.E.C., Martins, R.P. & Del-Claro, K. (2012). Use of plant resources by *Merosargus* (Diptera, Stratiomyidae, Sarginae) larvae. *Psyche, 2012*, 1-10. doi: 10.1155/2012/690203

Gligorescu, A., Toft, S., Hauggaard-Nielsen, H., Axelsen, J.A., & Nielsen, S. A. (2018). Development, metabolism and nutrient composition of black soldier fly larvae (*Hermetia illucens*; Diptera: Stratiomyidae) in relation to temperature and diet. *Journal of Insects as Food and Feed, 4*(2), 123-133. doi: 10.3920/JIFF2017.0080

Syrphidae

Branquart, E. & Hemptinne, J. (2008). Selectivity in the exploitation of floral resources by hoverflies (Diptera: Syrphinae). *Ecography, 23*(6), 732-742. doi: 10.1111/j.1600-0587.2000.tb00316.x

Budenberg, W.J. & Powell, W. (1992). The role of honeydew as an ovipositional stimulant for two species of syrphids. *Entomologia Experimentalis et Applicata, 64*(1), 57-61. doi: 10.1111/j.1570-7458.1992.tb01594.x

Campbell, D.R., Bischoff, M., Lord, J.M. & Robertson, A.W. (2010). Flower color influences insect visitation in alpine New Zealand. *Ecology, 91*(9), 2638-2649. doi: 10.1890/09-0941.1

Campoy, A., Aracil, A., Pérez-Bañón, C. & Rojo, S. (2020). An in-depth study of the larval head skeleton and the external feeding structures related with the ingestion of food particles by the eristaline flower flies *Eristalis tenax* and *Eristalinus aeneus. Entomologia Experimentalis et Applicata, 168*(10), 783-798. doi: 10.1111/eea.12974

Dziock, F. (2005). Evolution of prey specialization in aphidophagous syrphids of the genera *Melanostoma* and *Platycheirus* (Diptera: Syrphidae) 1. Body size, development and prey traits. European Journal of Entomology, 102(3), 413-421. doi: 10.14411/eje.2005.059

Francuski, L., Matić, I., Ludoški, J. & Milankov, V. (2011). Temporal patterns of genetic and phenotypic variation in the epidemiologically important drone fly, *Eristalis tenax. Medical and Veterinary Entomology, 25*(2), 135-147. doi: 10.1111/j.1365-2915.2011.00956.x

Gojkovic, N., Francuski, L., Ludoski, J. & Milankov, V. (2020). DNA barcode assessment and population structure of aphidophagous hoverfly *Sphaerophoria scripta*: Implications for conservation biological control. *Ecology and Evolution, 10*(7), 9428-9443. doi: 10.1002/ece3.6631

Jamali, R.A., Memom, N., Shah, M.A., Khan, K. & Ansari, A. (2018). Prevalence of aphidophagous hoverflies (Syrphidae: Syrphinae) in relation to their prey, green aphids on Brassica in Dadu. *Journal of Animal and Plant Sciences, 28*(5), 1447.

Jiang, S., Li, H., He, L. & Wu, K. (2022). Population fitness of *Eupeodes corollae* Fabricius (Diptera: Syrphidae) feeding on different species of aphids. *Insects, 13*(6), 494. doi: 10.3390/insects13060494

Jovičić, S., Burgio, G., Diti, I., Krašić, D., Markov, Z., Radenković, S. & Vujić, A.(2017). Influence of landscape structure and land use on M*erodon* and *Cheilosia* (Diptera: Syrphidae): contrasting responses of two genera. *Journal of Insect Conservation, 21*(1), 53-64. doi: 10.1007/s10841-016-9951-1

Khaganinia, S. & Kazerani, F. (2014). A review of the genus *Cheilosia* (Diptera, Syrphidae) from Iran. *Vestnik Zoologii, 48*(5), 401-410. doi: 10.2478/vzoo-2014-0048

Koval, A.G., Guseva, O.G. & Shpanev, A.M. (2018). Hoverflies (Diptera, Syrphidae) in Agrolandscapes of St. Petersburg and Leningrad Province. *Entomological Review, 98*(6), 702-708. doi: 10.1134/S0013873818060064

Krivosheina, N.P. (2021). Ecological relations of the hoverfly larvae (Diptera, Syrphidae, Eristalinae) bark inhabitants with xylobiont insects. *Biology Bulletin of the Russian Academy of Sciences, 47*(6), 605-616. doi: 10.1134/S1062359020060096

Lillo, I., Perez-Bañón, C. & Rojo, S. (2021). Life cycle, population parameters, and predation rate of the hover fly *Eupeodes corollae* fed on the aphid *Myzus persicae. Entomologia Experimentalis et Applicata, 169*(11), 1027-1038. doi: 10.1111/eea.13090

Mengual, X., Stahls, G. & Rojo, S. (2008). First phylogeny of predatory flower flies (Diptera, Syrphidae, Syrphinae) using mitochondrial COI and nuclear 28S rRNA genes: conflict and congruence with the current tribal classification. *Cladistics, 24*(4), 543-562. doi: 10.1111/j.1096-0031.2008.00200.x

Mielczarek, L.E. (2013). *Melangyna soszynskii* sp. n. (Diptera: Syrphidae) from the Sayan Mountains. *Polskie Pismo Entomologiczne, 82*(4), 339-342. doi: 10.2478/v10200-012-0047-2

Milic, D., Radenkovic, S., Radisic, D., Andric, A., Nikolic, T., Vujic, A. & Silva, D.P. (2019). Stability and changes in the distribution of *Pipiza* hoverflies (Diptera, Syrphidae) in Europe under projected future climate conditions. *PLoS One, 14*(9), p.e0221934. doi: 10.1371/journal.pone.0221934

Nedeljković, Z., Ačanski, J., Đan, M., Obreht-Vidaković, D., Ricarte, A., Vujić, A. & Biesmeijer, J.C. (2015). An integrated approach to delimiting species borders in the genus *Chrysotoxum* Meigen, 1803 (Diptera: Syrphidae), with description of two new species. *Contributions to Zoology, 84*(4), 285-304. doi: 10.1163/18759866-08404002

Nedeljković, Z., Ačanski, J., Vujić, A., Obreht, D., Đan, M., Ståhls, G. & Radenković, S. (2013). Taxonomy of *Chrysotoxum festivum* Linnaeus, 1758 (Diptera: Syrphidae) - an integrative approach. *Zoological Journal of the Linnean Society, 169*(1), 84-102. doi: 10.1111/zoj.12052

Reemer, M. & Rotheray, G.E. (2009). Pollen feeding larvae in the presumed predatory syrphine genus *Toxomerus* Macquart (Diptera, Syrphidae). *Journal of Natural History, 43*(15-16), 939-949. doi: 10.1080/00222930802610576

Rizza, A., Campobasso, G., Dunn, P.H. & Stazi, M. (1988). *Cheilosia corydon* (Diptera: Syrphidae), a candidate for the biological control of musk thistle in North America. *Annals of the Entomological Society of America, 81*(2), 225-232. doi: 10.1093/aesa/81.2.225

Rotheray, E. L., Lepais, O., Nater, A., Krützen, M., Greminger, M., Goulson, D. & Bussière, L. F. (2012). Genetic variation and population decline of an endangered hoverfly *Blera fallax* (Diptera: Syrphidae). *Conservation Genetics, 13*(5), 1283-1291. doi: 10.1007/s10592-012-0371-9

Tijana, N., D.R., Dubravka, M., V.M., Sonja, T., Snežana, J., Smiljka, S. & A.V. (2013). Models of the potential distribution and habitat preferences of the genus *Pipiza* (Syrphidae: Diptera) on the Balkan peninsula. *Archives of Biological Sciences, 65*(3), 1037-1052. doi: 10.2298/ABS1303037N

Vanhaelen, N., Francis, F. & Haubruge, E. (2004). Purification and characterization of glutathione S-transferases from two syrphid flies (*Syrphus ribesii* and *Myathropa florae*). *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology, 137*(1), 95-100. doi: 10.1016/j.cbpc.2003.10.006

Yan, Y., Hu, L. & National Library of Medicine. (2021). The complete mitochondrial genome of *Platycheirus albimanus* (Diptera: Syrphidae: Syrphinae) and phylogenetic analysis of the Syrphidae. *Mitochondrial DNA, 6*(2), 528-530. doi: 10.1080/23802359.2021.1872455.

Tabanidae

Kelly-Hope, L., Paulo, R., Thomas, B., Brito, M., Unnasch, T.R. & Molyneux, D. (2017). Loa loa vectors *Chrysops* spp.: perspectives on research, distribution, bionomics, and implications for elimination of lymphatic filariasis and onchocerciasis. *Parasites & Vectors, 10*(1), 172. doi: 10.1186/s13071-017-2103-y

Lall, S.B. (1969). Effect of starvation on the sensitivity of tarsal and labellar contact chemoreceptors in *Hybomitra lasiophthalma* (Diptera: Tabanidae). *Annals of the Entomological Society of America, 62*(5), 1217. doi: 10.1093/aesa/62.5.1217

Ossowski, A. & Hunter, F.F. (2012). Distribution patterns, body size, and sugar-feeding habits of two species of *Chrysops* (Diptera: Tabanidae). *The Canadian Entomologist, 132*(2), 213-221. doi: 10.4039/Ent132213-2

Teskey, H.J., Shemanchuk, J.A. & Weintraub, J. (1987). *Hybomitra agora*, a new species of Tabanidae (Diptera) from Western North America. *Canadian Entomologist, 119*(12), 1117-1122. doi: 10.4039/Ent1191117-12

Tachinidae

Al-Dobai, S., Reitz, S. & Sivinski, J. (2012). Tachinidae (Diptera) associated with flowering plants: estimating floral attractiveness. Biological Control, 61(3), 230-239. doi: 10.1016/j.biocontrol.2012.02.008

Bora, D. & Deka, B. (2014). Role of visual cues in host searching behaviour of *Exorista sorbillans* Widemann, a parasitoid of Muga Silk Worm, *Antheraea assama* Westwood. *Journal of Insect Behavior, 27*(1), 92-104. doi: 10.1007/s10905-013-9409-1

Cerretti, P., O'Hara, J.E., Wood, D.M., Shima, H., Inclan, D.J. & Stireman III, J.O. (2014). Signal through the noise? Phylogeny of the Tachinidae (Diptera) as inferred from morphological evidence. Systematic Entomology, 39(2), 335-353. doi: 10.1111/syen.12062

Dindo, M.L., Rezaei, M. & De Clercq, P. (2019). Improvements in the rearing of the Tachinid parasitoid *Exorista larvarum* (Diptera: Tachinidae): Influence of adult food on female longevity and reproduction capacity. *Journal of Insect Science, 19*(2). doi: 10.1093/jisesa/iey122

Foerster, L.A. & Doetzer, A.K. (2002). Host instar preference of *Peleteria robusta* (Wiedman) (Diptera: Tachinidae) and development in relation to temperature. *Neotropical Entomology, 31*(3), 405-409. doi: 10.1590/S1519-566X2002000300009

Kara, K., Tschosnig, H. & Atay, T. (2020). Checklist of Turkish Tachinidae (Insecta, Diptera) with new records. *Journal of the Entomological Research Society*, 22(2), 163-190.

Martel, V., Thireau, J. & Régnière, J. (2022). Manual inoculation of host larvae with first instar maggots as a rearing technique for the larval parasitoid *Actia interrupta* (Diptera: Tachinidae). *Biocontrol Science and Technology, 32*(1), 110-113. doi: 10.1080/09583157.2021.1967292

Regniere, J., Thireau, J., Saint-Amant, R. & Martel, V. (2021). Modeling climatic influences on three parasitoids of low-density spruce budworm populations. Part 3: *Actia interrupta* (Diptera: Tachinidae). *Forests, 12*(11), 1471. doi: 10.3390/f12111471

Sahebari, F.S., Khaghaninia, S. & Talebi, A.A. (2018). New data of the subfamily Tachininae (Diptera: Tachinidae) from north-western Iran. *Zoology and Ecology*, 28(3), 252-258. doi: 10.1080/21658005.2018.1490107

Scaccini, D. (2018). Remarks on the biology of *Dinera ferina* (Diptera Tachinidae) as parasitoid of the two Italian *Platycerus* species (Coleoptera Lucanidae). *Bulletin of Insectology, 71*(1), 39.

Tachi, T. & Shima, H. (2010). Molecular phylogeny of the subfamily Exoristinae (Diptera, Tachinidae), with discussions on the evolutionary history of female oviposition strategy. *Systematic Entomology, 35*(1), 148-163. doi: 10.1111/j.1365-3113.2009.00497.x

Tipulidae

Bearup, D., Petrovskii, S. Blackshaw, R. & Hastings, A. (2013). Synchronized dynamics of *Tipula paludosa* metapopulation in a Southwestern Scotland agroecosystem: Linking pattern to process. *The American Naturalist, 182*(3), 393-409. doi: 10.1086/671162

Blackshaw, R.P. & Moore, J.P. (2011). Within-generation dynamics of leatherjackets (*Tipula paludosa* Meig.) *Journal of Applied Entomology, 136*(8), 605-613. doi: 10.1111/j.1439-0418.2011.01696.x

Cheever, B.M. & Webster, J.R. (2014). Effects of consumers and nitrogen availability on heterotrophic microbial activity during leaf decomposition in headwater streams. *Freshwater Biology, 59*(8), 1768-1780. doi: 10.1111/fwb.12383

Cook, D.M., Henriksen, E.D., Upchurch, R. & Peterson, J.B.D. (2020). Isolation of polymer-degrading bacteria and characterization of the hindgut bacterial community from the detritus-feeding larvae of *Tipula abdominalis* (Diptera: Tipulidae). *Applied and Environmental Microbiology, 73*(17), 5683-5686. doi: 10.1128/AEM.00213-07

Jo, J. (2017). Description of larval and pupal stages of *Tipula (Nippotipula) sinica* (Diptera, Tipulidae) from South Korea with ecological notes. *Animal Systematics, Evolution and Diversity, 33*(1), 56-59. doi: 10.5635/ASED.2017.33.1.036

Kania, I. & Nel, A. (2013). New fossil Tipulidae from the volcano-sedimentary latest Oligocene of Bes-Konak (Turkey). *Polish Journal of Entomology, 82*(4), 327-338. doi: 10.2478/v10200-012-0046-3

Meats, A. (1975). The development dynamics of *Tipula paludosa* and the relation of climate to its growth pattern, flight season and geographical distribution. *Oecologia, 19*(2), 177-128. doi: 10.1007/BF00369096

Peck, D.C. & Olmstead, D. (2009). Invasive *Tipula* (Diptera: Tipulidae) in turfgrass of the Northeast United States: Geographic distribution and local incidence three years after detection. *Journal of Economic Entomology, 102*(2), 652-658. doi: 10.1603/029.102.0225

Peck, D.C., Olmstead, D. & Morales, A. (2008). Application timing and efficacy of alternatives for the insecticidal control of *Tipula paludosa* Meigen (Diptera: Tipulidae), a new invasive pest of turf in the northeastern United States. *Pest Management Science, 64*(10), 989-1000. doi: 10.1002/ps.1592

Petersen, M.J. (2013). Evidence of a climatic niche shift following North American introductions of two crane flies (Diptera; genus *Tipula*). *Biological Invasions, 15*(4), 885-897. doi: 10.1007/s10530-012-0337-3

Podeniene, V., Gelhaus, J.K. & Yadamsuren, O. (2006). The last instar larvae and pupae of *Tipula (Arctotipula)* (Diptera, Tipulidae) from Mongolia. *Proceedings of the Academy of Natural Sciences of Philadelphia, 155*(1), 79-105. doi: 10.1635/i0097-3157-155-1-79.1

Pritchard, G. (1983). Biology of Tipulidae. *Annual Review of Entomology, 28*(1), 1-22. doi: 10.1146/annurev.en.28.010183.000245

Rao, S., Liston, A., Crampton, L. & Takeyasu, J. (2006). Identification of larvae of exotic *Tipula paludosa* (Diptera: Tipulidae) and *T. oleracea* in North America using mitochondrial cytB sequences. *Annals of the Entomological Society of America, 99*(1), 33-40. doi: 10.1603/0013-8746(2006)099[0033:IOLOET]2.0.CO2

Rogers, T.E. & Doran-Peterson, J. (2010). Analysis of cellulolytic and hemicellulolytic enzyme activity within the *Tipula abdominalis* (Diptera: Tipulidae) larval gut and characterization of *Crocebacterium ilecola* gen. nov., sp. nov., isolated from the *Tipula abdominalis* larval hindgut. *Insect Science, 17*(3), 291-302. doi: 10.1111/j.1744-7917.2010.01346.x

Salmela, J. & Petrašiūnas, A. (2014). Checklist of the infraorder Tipulomorpha (Trichoceridae, Tipuloidea) (Diptera) of Finland. *ZooKeys, 441*(441), 21-36. doi: 10.3897/zookeys.441.7533

Smith, R.M., Young, M.R. & Marquiss, M. (2008). Bryophyte use by an insect herbivore: does the crane-fly *Tipula montana* select food to maximise growth? *Ecological Entomology, 26*(1), 83-90. doi: 10.1046/j.1365-2311.2001.00297.x

Smith-Herron, A.J., Cook, J.L. & Cook, T.J. (2014). *Brevispina tasii* (Brevispinidae) n. fam., n. gen., n. sp (Apicomplexa: Eugregarinidae: Stenophoroidea) parasitizing the crane fly *Tipula umbrosa* (Diptera: Tipulidae: Tipulinae) from East Texas leaf litter. *Comparative Parasitology, 81*(1), 1-9. doi: 10.1654/4656.1

Trichoceridae

Laurence, B.R. (1955). On the life history of *Trichocera saltator* (Harris) (Diptera, Trichoceridae). *Proceedings of the Zoological Society of London, 126*(2), 235-244. doi: 10.1111/j.1096-3642.1956.tb00434.x

Ulidiidae

Goyal, G., Nuessly, G.S., Seal, D.R., Steck, G.J., Capinera, J.L. & Boote, K.J. (2017). Developmental studies of maize-infesting picture-winged flies (Diptera: Ulidiidae). *Environmental Entomology, 46*(4), 946-953. doi: 10.1093/ee/nvx082

Kaplin, V. G., Nartshuk, E. P., Volodina, I. A. & Kuryanovich, A. A. (2022). A new alfalfa nectarophage from the genus *Timia* Wiedemann (Diptera, Ulidiidae) in the forest-steppe of the Middle Volga Region. *Entomological Review, 101*(9), 1304-1309. doi: 10.1134/S0013873821090086

Owens, D., Larsen, N. & Nuessly, G.S. (2017). Post-harvest crop destruction effects on picture-winged fly (Diptera: Ulidiidae) emergence. *The Florida Entomologist, 100*(2), 422-425. doi: 10.1653/024.100.0203

Xylophagidae

Lee, J., Kim, Y. & Suh, S.J. (2022). First record of the awl fly genus *Xylophagus* (Diptera: Xylophagidae) from Korea. *Animal Systematics, Evolution and Diversity, 38*(1), 34-38.

Nagatomi, A. & Liu, N. (1995). Spermatheca and female terminalia of Pantophthalmidae and xylophagidae s. lat. (Diptera). *Annals of the Entomological Society of America, 88*(5), 603-626. doi: 10.1093/aesa/88.5.603