A Novel Silver-Containing Antimicrobial potentiates aminoglycoside
activity against *Pseudomonas aeruginosa.*

Gracious Yoofi Donkor[a], Greg M. Anderson[a], Michael Stadler[a], Patrick Ofori Tawiah[a], Carl

D. Orellano[a], Kevin A. Edwards[b] and Jan-Ulrik Dahl[a#]

[a]School of Biological Sciences, Illinois State University, Microbiology, Normal, IL, USA

[b]School of Biological Sciences, Illinois State University, Cell Biology, Normal, IL, USA

RUNNING TITLE: Reactive oxygen species in antimicrobial therapy

[#]Address correspondence to Jan-Ulrik Dahl, email: jdahl1@ilstu.edu, phone: +1-(309)-

438-7694

Word count:

Abstract: 244

Importance: 130

Manuscript: 7158

ABSTRACT

The rapid dissemination of antibiotic resistance combined with the decline in the discovery of novel antibiotics represents a major challenge for infectious disease control that can only be mitigated by investments into novel treatment strategies. Alternative antimicrobials, including silver, have regained interest due to their diverse mechanisms of inhibiting microbial growth. One such example is AGXX®, a broad-spectrum silver containing antimicrobial that produces highly cytotoxic reactive oxygen species (ROS) to inflict extensive macromolecular damage. Due to connections identified between ROS production and antibiotic lethality, we hypothesized that AGXX® could potentially increase the activity of conventional antibiotics. Using the gram-negative pathogen *Pseudomonas aeruginosa*, we screened possible synergistic effects of AGXX® on several antibiotic classes. We found that the combination of AGXX® and aminoglycosides tested at sublethal concentrations led to a rapid exponential decrease in bacterial survival and restored sensitivity of a kanamycin-resistant strain. ROS production contributes significantly to the bactericidal effects of AGXX®/aminoglycoside treatments, which is dependent on oxygen availability and can be reduced by the addition of ROS scavengers. Additionally, *P. aeruginosa* strains deficient in ROS detoxifying/repair genes were more susceptible to AGXX®/aminoglycoside treatment. We further demonstrate that this synergistic interaction was associated with significant increase in outer and inner membrane permeability, resulting in increased antibiotic influx. Our study also revealed that AGXX®/aminoglycoside-mediated killing requires an active proton motive force across the bacterial membrane. Overall, our findings provide an understanding of cellular targets that could be inhibited to increase the activity of conventional antimicrobials.

IMPORTANCE

The emergence of drug-resistant bacteria coupled with the decline in antibiotic development highlights the need for novel alternatives. Thus, new strategies aimed at repurposing conventional antibiotics have gained significant interest. The necessity of these interventions is evident especially in gram-negative pathogens as they are particularly difficult to treat due to their outer membrane. This study highlights the effectiveness of the silver containing antimicrobial AGXX® in potentiating aminoglycoside

activities against *P. aeruginosa*. The combination of AGXX® and aminoglycosides not
only reduces bacterial survival rapidly but also significantly re-sensitizes aminoglycoside-
resistant *P. aeruginosa* strains. In combination with gentamicin, AGXX® induces
increased endogenous oxidative stress, membrane damage and iron sulfur cluster
disruption. These findings emphasize AGXX®'s potential as a route of antibiotic adjuvant
development and shed light into potential targets to enhance aminoglycoside activity.
67
68

INTRODUCTION
The spread of antibiotic resistance has now reached a large number of bacterial
pathogens and evolved into a pertinent global health challenge (*1*). Over the past decade,
resistance has been reported against all classical antibiotics, including last resort
treatments such as polymyxins (*2*). The resistance crisis is further exacerbated in gram-
negative pathogens due to the low permeability of their outer membrane and the
extensive arsenal of drug resistance mechanisms that these critters employ. One
example of a difficult to treat gram-negative bacterium is the opportunistic pathogen *P.*
aeruginosa, a common cause of acute (e.g., wounds, burns) and chronic infections (e.g.,
diabetic ulcers, cystic fibrosis) (*3*). *P. aeruginosa* is characterized by its high intrinsic and
acquired resistance mechanisms, allowing the pathogen to thrive in the presence of a
large number of antibiotics (*4*).
81

In light of these challenges, recent studies have focused on alternative treatment
strategies to limit bacterial infection and colonization (*5*, *6*). Transition metals, such as
silver and copper, have long been recognized for their antimicrobial activities and were
already used by the ancient Greeks for would healing (*7*). Despite its long-standing history
and high efficiency against bacteria, the antimicrobial mode of action of silver is poorly
understood. Pleiotropic effects have been described and include changes in DNA
condensation, membrane alteration and protein damage (*7*, *8*). Silver ions have a
particularly high affinity for cysteine thiols, disrupt exposed iron-sulfur clusters of
dehydratase enzymes, and replace metal-containing cofactors, thus affecting a wide
range of cytoplasmic and membrane proteins (*9*, *10*). More recently, silver derivates have

received increased attention in medical applications, e.g. as antimicrobial surface-
coatings on catheters and implants to protect against biofilm-forming bacteria and to
reduce the risk of nosocomial infections (*5, 11*). One such promising silver-based
antimicrobial with broad-spectrum activity is AGXX®. AGXX® consists of micro-galvanic
elements of silver (Ag) and ruthenium (Ru), which are surface conditioned with ascorbic
acid (*12, 13*). AGXX® has been used in the form of a powder and implemented as an
electroporated sheet on a variety of surfaces, including steel meshes, ceramics and water
pipes, to limit bacterial colonization. AGXX®'s antimicrobial is not entirely dependent on
the release of silver ions; instead it is proposed to generate reactive oxygen species
(ROS), such as hydroxyl radicals (•OH) and superoxide (O_2^-) through a series of redox
reactions where the oxidized Ag component is reduced by organic matter and donates
electrons to the valent Ru, which subsequently generates O_2^- and other ROS (*12–16*).
Previous studies revealed that, compared to classical silver and other metals, AGXX®
was significantly more bactericidal against gram-positive bacteria such as
Staphylococcus aureus and *Enterococcus faecalis* (*12*, *13*). Surprisingly, the antimicrobial
effects of AGXX® on gram-negative pathogens remain largely unexplored.

Compounds that generate ROS and/or stimulate endogenous oxidative stress in bacteria
have gained interest as potential antibiotic adjuvants (*17*). Adjuvants increase the efficacy
of antibiotics by targeting metabolic processes or cellular networks that ultimately lead to
a synergistic increase in antibiotic potency (*18*). A classic example for the synergy
between adjuvants and antibiotic is the combination of amoxicillin, a beta-lactam
antibiotic, and clavulanic acid. Due to clavulanic acid's high affinity for beta-lactamase
enzymes, its combination with the beta-lactam antibiotics, such as amoxicillin,
synergistically potentiate their activity against penicillin-resistant bacteria (*19*). The
hypothesis behind the synergistic effects of ROS-generating or -inducing compounds is
based on the multimodal action of these compounds, which could potentially disrupt
bacterial targets necessary for the defense of antibiotics (*20*). Recent studies on the
bactericidal mode of action of aminoglycoside, fluoroquinolone and beta-lactam
antibiotics have also proposed increased endogenous ROS stress as an additional
mechanism for bacterial killing (*5, 20, 21*). Notably, activation of the bacterial envelope

stress response, hyperactivation of the electron transport chain, and damage of iron sulfur
clusters have been shown to contribute to an increase in endogenous ROS level (5).
Although these findings are controversial, extensive evidence has been presented for the
integral role of ROS-mediated damage in antibiotic-induced cell death (6). More
importantly, recent studies employing ROS-generating compounds have reported
promising results on their potential in sensitizing a wide variety of multidrug-resistant
bacteria to both bactericidal and bacteriostatic antibiotics (5).
130
Given that AGXX®'s proposed mode of action involves ROS production and studies with
focus on evaluating potential synergistic effects of AGXX® on conventional antibiotics are
lacking, we started to investigate possible potentiating effects of AGXX® on members of
several antibiotic classes, using the *P. aeruginosa* strain PA14 as a model. Exploring
potential synergies between AGXX® and antibiotics could provide viable answers to the
antimicrobial discovery drought for instance by reducing the minimal inhibitory
concentrations of conventional antibiotics required for treatment or possibly overriding
and/or delaying antibiotic resistance development (22). By exposing PA14 to sublethal
concentrations of AGXX® and the antibiotics of interest alone and in combination, we
found that bacterial survival was exponentially reduced when the cells were exposed to
combinations of AGXX® and aminoglycoside antibiotics. Moreover, combined treatment
of both compounds re-sensitized a kanamycin resistant PA14 strain to sublethal
concentrations of the antibiotic. We further demonstrate that the combined treatment of
AGXX® and aminoglycosides resulted in increased endogenous oxidative stress and a
subsequent disruption of iron homeostasis, potentially providing an explanation for the
elevated ROS level. The synergy was associated with a significant increase in outer and
inner membrane permeability, which facilitates antibiotic influx. Moreover, our studies
revealed that the synergy between AGXX® and aminoglycosides relies on an active
proton motive force across the bacterial membrane.
150

RESULTS
AGXX® is more efficient in killing P. aeruginosa than silverdene and silver nitrate.
Studies with silver ions suggest that gram-negative bacteria are more susceptible than

gram-positive bacteria (*23*). In this study, we investigated the effect of AGXX® on *P.*
aeruginosa using PA14 as model strain. To compare the effective antimicrobial
concentrations of different AGXX® formulations against PA14, we first performed survival
analyses in the presence and absence of AGXX383®, AGXX394®, AGXX823®, and
AGXX720C®, respectively. While these AGXX® formulations all consist of the galvanized
silver/ruthenium complex, they differ in various aspects such as silver ratio, particle size,
and production procedure, which may affect their antimicrobial activity. Our time killing
assays revealed that AGXX383® and AGXX394® have comparable antimicrobial
activities against PA14 and are considerably more potent in inhibiting PA14 growth and
survival compared to AGXX823® and AGXX720C® (**Supplementary FIG S1**). Silver
derivates have received increased attention in medical applications, e.g. as antimicrobial
surface-coatings on catheters that protect from biofilm-forming bacteria and reduce the
risk of nosocomial infections (*24*). Moreover, silver is used in topicals to prevent and/or
treat infections in wounds (*25*). One such example is silver sulfadiazine (silverdene), the
gold standard for treating and preventing *P. aeruginosa* infections in burn wound patients
(*26*). However, silverdene is associated with complications such as allergic reactions to
the sulfadiazine moiety emphasizing the need for novel treatment therapies (*26*). To
compare the antimicrobial activities of AGXX394®, silver nitrate ($AgNO_3$) and silverdene,
we conducted survival studies of PA14 in the presence and absence of 25 μg/ml of each
compound. Survival of PA14 was highly compromised when cells were exposed to
AGXX394® (**FIG.1**). A direct comparison of the killing efficiencies between AGXX394 and
silverdene and silver nitrate after 3 hours of treatment revealed a ~1-log and 2-log higher
bactericidal activity for AGXX394®, respectively, potentially making AGXX® an attractive
treatment/prevention alternative (**FIG. 1).** AGXX treatment has been shown to elicit
oxidative stress responses shock responses and other general stress response regulons
(*13*). To confirm ROS production of AGXX394®-treated PA14, we quantified intracellular
ROS levels using the redox-sensitive molecular probe dichlorodihydrofluorescein
diacetate (H_2DCFDA). H_2DCFDA has been extensively demonstrated to detect various
ROS, including peroxides as well as peroxyl- and •OH radicals (*27*). When exposed to
ROS, the nonfluorescent fluorescent H_2DCFDA, is oxidized to highly green fluorescent
2',7'-dichlorofluorescein (DCF) (*27*). We found that exposure to both sublethal (20 µg/ml)

and bactericidal (40 µg/ml) levels of AGXX394® caused a 2.5 and 4-fold increase in
endogenous ROS levels **(Supplementary FIG S2A)**. Pretreating PA14 with thiourea, an
ROS scavenger, resulted in reduced ROS levels and recovered survival at bactericidal
AGXX394® concentrations (40 µg/ml) **(Supplementary FIG S2A&B)**.

AGXX® exponentially increases the activity of aminoglycosides against P.
aeruginosa. Our next goal was to determine how the presence of AGXX would affect
antibiotic activity against *P. aeruginosa.* Specifically, we were interested in determining
whether combining AGXX® with members of different antibiotic classes, such as
inhibitors of DNA replication (fluoroquinolones), cell wall biosynthesis (β-lactams), folate
biosynthesis, membrane integrity (polymyxins), and translation (aminoglycosides), show
increased bactericidal activity against *P. aeruginosa*. Here, we used AGXX720C®, a
formulation with significantly lower antimicrobial activity that allowed us to control dosage
of AGXX® more reliably (**Supplementary FIG S1**). We determined the minimal inhibitory
concentrations (MIC) of AGXX720C® and ten members of five different antibiotic classes
in PA14, which were cultivated under aerobic conditions in Mueller-Hinton broth
(**Supplementary Table S1**). Using time killing assays, we exposed PA14 to sublethal
concentrations of AGXX720C® (75 µg/ml), the indicated antibiotics, and their
combinations at the same sublethal concentrations that were used for the individual
treatments, respectively. We monitored colony forming units counts every 60 min over a
time course of three hours (**FIG 2; Supplementary FIG S3**) and calculated the percent
survival for each sample at the 3-hour time point relative to the untreated control (**FIG 2**).
None of the individual treatments with either 75 µg/ml AGXX720C® or any of the
antibiotics resulted in substantial killing of PA14 (**FIG 2; Supplementary FIG S3**). The
combination of 75 µg/ml AGXX720C® with 35 ng/ml ciprofloxacin and 100 ng/ml
norfloxacin, respectively, resulted in PA14 survival comparable to the individual
treatments suggesting that AGXX® does not potentiate the bactericidal activities of
fluoroquinolones against PA14 (**FIG 2A; Supplementary FIG S3A, B**). Likewise, the
combined treatments of AGXX720C® and 78 µg/ml carbenicillin, 0.156 µg/ml imipenem,
or 125 µg/ml trimethoprim did not significantly change PA14 survival compared to their
individual treatments (**FIG 2B, C; Supplementary FIG S3C**). On the other hand, we found

that the bactericidal activity of the membrane targeting antibiotic polymyxin B was
increased by the presence of AGXX720C® as evidenced by a 1-log reduction in PA14
survival as compared to AGXX720C® or polymyxin B alone, which each caused less than
5% killing (**FIG 2D; Supplementary FIG S3D**). However, we observed the most drastic
decrease in PA14 survival when AGXX720C® was combined with a member of
aminoglycoside antibiotics, even at concentrations far below the MIC. Co-treatment of
AGXX720C® with 0.4 µg/ml gentamicin (Gm) (0.2x MIC) reduced PA14 survival by as
much as 4 logs after three hours of treatment (**FIG 2E; Supplementary FIG S3E**), while
combinations of AGXX720C® with 2 µg/ml amikacin (0.27x MIC), 1 µg/ml tobramycin
(0.4x MIC), or 3 µg/ml streptomycin (0.15x MIC) caused up to 3 log reduction in survival
(**FIG 2E, Supplementary FIG S3F-H**).
It has been reported that $AgNO_3$'s synergizing effects on aminoglycosides was reduced
in minimal media (*28*). We therefore exposed PA14 sub-cultured in MOPS-glucose
(MOPSg) minimal medium to increasing sublethal AGXX concentrations alone or in
combination with a sublethal gentamicin (**Supplementary FIG S4A**). Increasing AGXX®
concentrations significantly reduced PA14 survival in a dose-response like manner
(**Supplementary FIG S4A**). Considering that casamino acid (CAS) supplementation was
also reported to partially restore $AgNO_3$'s effects in minimal media, we replicated our time
killing assays in MOPSg media in the presence of 0.2% CAS. Surprisingly, addition of
0.2% CAS decreased AGXX®'s effects on Gm in MOPSg medium (**Supplementary FIG
S4B**). Although, we do not probe the reasons behind the effects of casamino acids in
minimal media, it is evident that AGXX® synergizes with aminoglycosides in both rich and
minimal media.
239
AGXX® increases the sensitivity of P. aeruginosa strain PA14 to kanamycin.
Considering the significant increase in lethality upon concurrent exposure of PA14 to
AGXX720C® and aminoglycosides, we sought to determine whether the addition of
AGXX720C® could reintroduce sensitivity in aminoglycoside-resistant *P. aeruginosa*
strains. Like many other *P. aeruginosa* strains, PA14 is intrinsically resistant to many
different antibiotics, including the aminoglycoside kanamycin. Under the conditions
tested, our PA14 strain showed a MIC of 240 µg/ml for kanamycin. We then exposed

PA14 to 75 µg/ml AGXX720C®, 50 µg/ml kanamycin, or a combination of 75 µg/ml

AGXX720C® and 50 µg/ml kanamycin and determined their survival over the time course

of 3 hours as described before. As early as 60 minutes post treatment, the combination

of AGXX720C® and kanamycin reduced colony survival by ~1.5 log, which increased to

about 4 log (10,000-fold) difference in survival after 3 hours (**FIG 3**). In summary, we

conclude from our data that AGXX® potentiates the activity of a wide range of

aminoglycoside antibiotics, potentially making aminoglycoside-resistant *P. aeruginosa*

strains more sensitive again.

The combination of sublethal concentrations of AGXX® and aminoglycosides

increases ROS formation and causes upregulation of genes indicative of DNA

damage and protein aggregation. AGXX®'s main antimicrobial mode of action is

mediated by the generation of ROS (39). Increased endogenous oxidative stress has also

been linked to the lethality of aminoglycosides and other bactericidal antibiotics (*29, 30*).

It may therefore be possible that a disruption in the redox balance during antibiotic

exposure increases lethality by amplifying the adventitious generation of ROS by

antibiotics. We treated exponentially growing PA14 for 60 min with sublethal

concentrations of Gm (0.25 µg/ml), AGXX720C® (50 µg/ml), or their combination and

compared H2DCFDA fluorescence in each sample to the untreated control. Individual

treatments with either Gm or AGXX720C® did not result in a significant increase in

H_2DCFDA fluorescence, excluding the possibility that substantial amounts of ROS were

formed at these sublethal concentrations (**FIG 4A**). However, when applied in

combination, we observed an 8-fold increase in H_2DCFDA fluorescence indicative of

increased ROS production under these conditions. Pre-treatment of PA14 with the ROS

scavenger thiourea (*31*) or the hydrogen peroxide (H_2O_2) -detoxifying enzyme catalase

resulted in a significant decline in H_2DCFDA fluorescence (**FIG 4A**) as well as restored

PA14 survival by 2- to 4-log, respectively (**FIG 4B; Supplementary FIG S5A**). Thus, our

data suggest that the high antimicrobial activity of the combinational treatment can at least

in parts be explained by increased ROS formation. However, H_2DCFDA is rather

unspecific and detects numerous ROS compounds. We therefore followed up on our

observation with the use of additional, more specific ROS-detecting fluorescent dyes. We

used the boronate-based peroxy orange 1 (PO1) dye as well as hydroxyphenyl
fluorescein (HPF) to examine intracellular concentrations of H_2O_2 and •OH levels,
respectively (*32, 33*). After 1 hour of treatment, we detected a significant shift in PO1 and
HPF fluorescence, indicating that both H_2O_2 and •OH are produced in AGXX720C®/Gm-
treated PA14 (**FIG 4C, D; Supplementary FIG S5B, C**).

To deal with the negative consequences of ROS and eliminate ROS-mediated damage,
microorganisms have evolved intricate systems to maintain and restore a balanced redox
homeostasis and repair ROS-mediated damage [recently reviewed in (*34–39*)]. Proteins
and nucleic acids represent the most prevalent targets of ROS (*40, 41*). To test whether
the elevated ROS production causes macromolecular damage, we analyzed the
transcript level of *ibpA* and *sulA,* two genes that have previously been shown to be
upregulated when cells experience severe oxidative stress (*42, 43*), in PA14 cells treated
with Gm and AGXX alone and in combination. *ibpA* encodes a molecular chaperone,
which plays a significant role in protecting bacteria from proteotoxic stressors including
ROS (*44*). *sulA* encodes the cell division inhibitor SulA, which is part of the SOS response
and induced when the cell experiences DNA damage, a possible consequence of
oxidative stress (*45*). We exposed exponentially growing PA14 cells to sublethal
concentrations of AGXX720C®, Gm, or the combination thereof for 60 min and quantified
ibpA/sulA mRNA level. While individual treatments with sublethal concentrations of
AGXX720C® or Gm did not cause significant changes in *ibpA/sulA* expression, their
transcript levels were approximately 5-fold (*sulA*) and 100-fold (*ibpA*) increased when
PA14 was treated with the combination of AGXX720C® and gentamicin, indicating
significant macromolecular damage **(FIG 4E)**.

Anaerobic growth and antioxidant systems provide protection against ROS-
mediated damage caused by a combinational treatment of AGXX® and
aminoglycosides*. To provide additional evidence that ROS contribute to the synergy
between aminoglycosides and AGXX®, we determined whether their killing efficiency
depends on the presence of molecular oxygen, which is essential for ROS production.
PA14 was grown in MHB under either aerobic or anaerobic conditions and treated with

sublethal concentrations of gentamicin, AGXX720C® or the combination thereof. When

grown anaerobically, MHB was supplemented with 1% KNO_3 to stimulate the proton

motive force (PMF). Anaerobically respiring PA14 cells tolerated treatments with AGXX®

and 0.5 µg/ml Gm (0.25x MIC) fairly well and showed about 2.5 log higher survival relative

to aerobically grown cells exposed to the same treatment. However, when we increased

the Gm concentration to 1 µg/ml (0.5x MIC) in the in the presence of AGXX®, PA14

survival also declined albeit at a lower rate than aerobically growing cells under the same

conditions **(FIG. 5A)**.

Next, we tested the susceptibility of PA14 strains with transposon insertions in genes that

have previously been identified as major oxidative stress response and/or defense

systems. We exposed PA14 transposon mutant strains (*46*) with defects in *oxyR*

(encodes ROS-sensing transcriptional regulator OxyR), *katA* (encodes catalase KatA),

katB (encodes catalase KatB), and *dps* (encodes an iron acquisition protein in response

to oxidative stress) to sublethal concentrations of AGXX720C® (25 µg/ml), tobramycin

(0.25 µg/ml), or the combination thereof for 3 hours and compared their colony survival

to that of the corresponding wildtype strain PA14 (**FIG. 5B**). We found that strains deficient

in functional copies of *katA* and *dps* showed a little over 1 log reduced survival compared

to wildtype cells, while no difference in survival was observed for cells lacking the

hydrogen peroxide global regulator OxyR and catalase KatB, respectively (**FIG. 5B**).

Taken together, our findings point towards a relevant role for ROS generation in the

synergistic interaction between aminoglycosides and AGXX®.

The synergy between AGXX® and aminoglycosides on PA14 killing is in parts

mediated by a disruption in iron homeostasis. Fe/S cluster are essential cofactors in

various metabolic enzymes, including members of the electron transport chain and the

tricarboxylic acid cycle (TCA) cycle (*47*). Fe/S cluster are particularly vulnerable to ROS,

which negatively impacts the activity of metabolic enzymes and ultimately cellular

metabolism during oxidative stress. Likewise, silver has been found to disrupt Fe/S

clusters in proteins (*10*), resulting in a cellular increase in free iron, which in turn can

stimulate •OH formation via Fenton reaction and cause extensive macromolecular

damage (*7*). Moreover, Fe/S disruption and Fenton-mediated •OH production have been

proposed as a downstream consequences of aminoglycoside killing (*30*). Given the

elevated •OH level (**FIG 4D; Supplementary FIG S5C**) and increased susceptibility of a

dps-deficient strain upon PA14 exposure to the AGXX®/Gm combination (**FIG 5B**), we

wondered whether the increased ROS production impairs the activity of Fe/S cluster-

containing enzymes. We prepared cell lysates from PA14 exposed to individual or

combined AGXX® and Gm, and measured the activity of aconitase, an Fe/S-containing

enzyme of the TCA cycle, whose activity depends on the presence of the Fe/S-cluster.

While individual treatments with sublethal AGXX720C® or Gm concentrations had no

impact on aconitase activity, exposure to a combination of AGXX720C® and Gm resulted

in ~75% loss in activity (**FIG 6A**). The loss in aconitase activity could be explained by the

release of iron atoms, as it occurs during Fe/S cluster oxidation under O_2^- and H_2O_2 stress

(*47, 48*). To test the role of free iron for the synergistic effect between AGXX720C® and

aminoglycosides, we treated PA14 with the iron chelator 2',2' bipyridyl prior to their

exposure to sublethal concentrations of AGXX720C® and Gm. Indeed, we found that

PA14 survival was less impaired when cells had been treated with 2',2' bipyridyl prior to

the presence of the AGXX720C®/Gm cocktail **(FIG 6B)**. However, iron chelators, such as

2',2' bipyridyl, have been reported to reduce aminoglycoside killing in *E. coli,* a

phenomenon we confirmed in *P. aeruginosa* **(Supplementary FIG S5D)**. Overall, our

data suggest a potential role of free iron for the ROS-mediated toxicity of aminoglycoside

exposure when in synergy with AGXX®.

Combined AGXX® and aminoglycoside treatment induces significant membrane

damage. In the initial stages of aminoglycoside uptake, the polycationic aminoglycoside

electrostatically interacts with the bacterial outer membrane, displacing membrane

divalent cations and inevitably increasing membrane permeability (*49*). Thus, compounds

with membrane permeabilizing properties have been reported as potent aminoglycoside

adjuvants (*50, 51*). To probe the role of membrane permeability, we evaluated outer

membrane disruption using the hydrophobic fluorescent probe N-phenyl-1-napthylamine

(NPN), which has diminished fluorescence in the presence of an intact outer membrane

but significantly increases when bound to exposed phospholipid groups in a disrupted

lipopolysaccharide monolayer (*52*). We found that in contrast to individual treatments with

sublethal AGXX720C® and Gm concentrations, a combined treatment significantly
increased NPN uptake, resulting in ~7-fold higher NPN fluorescence (**FIG 7A**). Next, we
examined the inner membrane permeability of PA14 cells that were subjected to
AGXX720C® and Gm treatment alone and in combination, using the fluorescent probe
propidium iodide (PI). Due to its size and charge, PI can only cross compromised inner
membranes where it binds nonspecifically to nucleic acids enhancing its fluorescence
exponentially (*53*). Individual treatments with sublethal AGXX720C® and gentamicin
concentrations did not result in significantly increased PI fluorescence (**FIG 7B**)
suggesting that at these concentrations none of them causes significant plasma
membrane damage. A combined treatment, however, led to substantially increased PI
fluorescence. Our spectrophotometric findings were complemented by fluorescent
microscopy analyses of PA14 cells that were treated as described before, washed in PBS,
stained with Syto9/PI (live/dead stain), incubated in the dark for 15 minutes at room
temperature, mounted onto a glass slide with 1% agarose, and imaged at 63x
magnification via inverted confocal microscopy. While the individual treatments with 0.25
µg/ml Gm or 50 µg/ml AGXX720C® only caused increased PI fluorescence in very few
cells, the number of PI-stained cells was substantially higher in PA14 cells that were
treated with a combination of sublethal AGXX720C®/Gm concentrations (**FIG 7C**).

AGXX® increases aminoglycoside uptake and lethality through increased activity
of the proton motive force (PMF). Considering the significance of membrane
permeability for aminoglycoside uptake, we reasoned that the increased membrane
disruption PA14 experiences during a combined treatment of AGXX® and
aminoglycosides would facilitate intracellular accumulation of aminoglycosides. Using
flow cytometry, we evaluated the uptake of Texas-Red-labelled Gentamicin (TR-Gm) in
exponentially growing PA14 in the presence or absence of AGXX720C®. Treatment of
PA14 with a combination of AGXX720C® and TR-Gm resulted in a significant increase in
TR-Gm uptake similar to treatments with the membrane-targeting antibiotic polymyxin B
(**FIG 8A, Supplementary FIG S6A).** The increased uptake of TR-Gm in the presence of
AGXX® resulted in a 2.5 log reduction in bacterial survival compared to PA14 that were
only exposed to TR-Gm alone **(Supplementary FIG S6B**). Moreover, the secondary and

tertiary stage of aminoglycoside import appears to depend on an active membrane
potential and occurs therefore only in actively respiring cells (*49, 54*). As such, increasing
cellular respiration or stimulation of membrane potential has been found to increase
aminoglycoside lethality (*55*). Using the protonophore CCCP, a compound that efficiently
inhibits the proton motive force (PMF), we determined whether the synergy between
AGXX® and aminoglycosides is dependent on the membrane potential. Not surprisingly,
PA14 killing by a lethal dose of gentamicin (1 µg/ml) could be reduced by almost 2 logs
when the cells were pre-incubated with CCCP (**Supplementary FIG S6C).** Likewise,
pretreatment with CCCP restored PA14 survival to a similar degree in the presence of
0.25 µg/ml Gm and 50 µg/ml AGXX720C® for three hours **(Fig. 8B).** Surprisingly,
combining polymyxin B at 1x MIC and 2x MIC with sublethal Gm concentrations did not
result in increased killing relative to polymyxin B alone **(Fig. S2D)**. Overall, our findings
indicate that AGXX®'s synergistic effect on aminoglycoside lethality does not only involve
increased membrane permeability but also requires an active proton motive force across
the bacterial membrane.
417

DISCUSSION
In the present study, we demonstrate evidence that the novel silver containing
antimicrobial AGXX® may potentially be used an antibiotic adjuvant with activity on the
opportunistic pathogen *P. aeruginosa*. We show that AGXX® serves as an efficient tool
to potentiate the activity of aminoglycoside antibiotics by reducing bacterial survival up to
50,000-fold. Aside from a 1-log increase in killing found when the Ag/Ru complex was
combined with the membrane-targeting antibiotic polymyxin B, AGXX® did not influence
the activity of antibiotics targeting DNA replication, cell wall synthesis, or folate
metabolism **(FIG 2)** raising the possibility that AGXX®'s potentiating effects could be
linked to a malfunction in protein synthesis. AGXX®'s potentiating effect was not limited
to select aminoglycosides but applied to all members tested, including gentamicin,
streptomycin, amikacin, and tobramycin **(FIG 3)**. We observed the exponential killing of
the AGXX®/aminoglycoside combination in complex and minimal media independent of
the presence or absence of amino acids, indicating that the media composition has only
minor effects on the synergistic effect (**Supplementary FIG S3)**. Additional support for

AGXX®'s stimulating effect on aminoglycoside activity was provided by its ability to re-

sensitize a kanamycin resistant *P. aeruginosa* strain (**FIG 4**). Notably, both AGXX® and

aminoglycoside antibiotics were present at concentrations far below their MICs (0.1-0.3x

MIC) indicating that the combination between the compounds was quite powerful with

regards to increasing the bactericidal activity of aminoglycosides.

Based on our findings, we propose the following model for the synergy between AGXX®

and aminoglycoside antibiotics (**FIG 9**): The antimicrobial action of AGXX® is proposed

to involve O_2^- and H_2O_2, which are generated in a redox cycle between silver (Ag) and

ruthenium (Ru^{x+1}) (*15, 16*). While the location of AGXX®-mediated ROS production was

not the focus of this study, we propose that ROS is mainly generated in the extracellular

part before the compounds penetrate the cell. This assumption is based on our finding

that exogenous addition of catalase was highly effective in reducing ROS level and

increasing *P. aeruginosa* survival upon treatment with a combination of AGXX® and Gm

(**Supplementary FIG S5A**), given that catalase cannot readily cross the bacterial cell

envelope. The combination of AGXX® and aminoglycosides concertedly increased

endogenous ROS levels, including H_2O_2 and •OH. Potentially facilitated by released silver

ions, the increased ROS level may disrupt iron-sulfur clusters in metabolic enzymes such

as aconitase, resulting in the release of free iron, which ultimately triggers •OH formation

in a Fenton reaction (*56*). Increasing ROS levels can inflict macromolecule damage such

as DNA damage, protein aggregation, and membrane damage, as evidenced by *(i)* the

increased expression of members of the SOS and heat shock responses, and *(ii)* the

possibility that the observed damage on outer and inner membrane is mediated by ROS.

In either way, increased ROS production contributes to the enhanced killing of *P.*

aeruginosa upon treatment with a combination of AGXX® and aminoglycosides. The

synergistic effect between AGXX® and aminoglycosides is also mediated by an increased

uptake and the cellular accumulation of aminoglycosides, which can be attenuated by

disrupting the bacterial membrane potential with ionophores such as CCCP.

The antimicrobial effects of silver and silver containing agents have been extensively

documented (*57–59*). However, in spite of the increasing use of silver in antimicrobial

applications, its mechanism of action remains to be poorly understood (*24, 60, 61*). Silver

along with other ROS inducing agents have been proposed to induce an array of

multimodal cytotoxic events, including the mis-metalation of proteins, DNA damage, and

imbalanced redox homeostasis, which ultimately disrupt multiple bacterial networks

leading to cell death (*5, 17, 62*). Thus, we investigated whether AGXX®, through its

multimodal effects as an ROS inducing compound, could increase that activity of

conventional antibiotics against *P. aeruginosa*. AGXX® itself has been proposed to exert

its antimicrobial action through the generation of ROS rather than the release of silver

ions. AGXX®-mediated ROS production likely starts with the generation of O_2^-, which

subsequently can be dismutated by cellular superoxide dismutases into hydrogen

peroxide (*15, 63*). Likewise, aminoglycosides have also been posited to induce metabolic

alterations that induce endogenous ROS levels (*64*). Growth of *P. aeruginosa* under

anaerobic conditions provided a protective effect against the bactericidal effects of

AGXX®/aminoglycoside killing **(FIG 5A)**. Considering that the availability of nitrate as an

alternative electron acceptor would facilitate PMF activity during anaerobic growth, a

decrease in killing efficiency points towards a relevant role for ROS in the synergistic

combination of the two compounds. At higher gentamicin concentrations, however,

AGXX® sensitized *P. aeruginosa* to gentamicin **(FIG 6A),** which could be potentially

explained with an increase in aminoglycoside uptake due to the high gentamicin

availability.

We found a significant increase in hydrogen peroxide and •OH radical signals upon

exposure of PA14 to a combination of AGXX® and aminoglycoside as evidenced by

increased PO1 and HPF fluorescence **(FIG 4)** and decreased survival of transposon

insertion strains defective in KatA and Dps **(FIG 5B)**. Surprisingly, a transposon insertion

into the *oxyR* gene, encoding the major hydrogen response regulator OxyR, did not show

a significant difference in survival relative to the wildtype under combined

AGXX®/aminoglycoside exposure **(FIG 5B).** However, this is consistent with a previous

study on the aminoglycoside potentiating effects of silver nitrate, in which neither *oxyR-*

deficient strains nor strains that constitutively express OxyR had significant effects on

bacterial survival (*28*). On the other hand, in a different study overexpression of the H_2O_2

detoxifying gene, *ahpF,* attenuated the lethality and proteotoxic effects of
aminoglycosides, providing evidence for the relevance of oxidative stress in
aminoglycoside toxicity (*65*). Previous studies have demonstrated that pretreatment with
antioxidant molecules such as thiourea reduced intracellular •OH radical levels and
increased bacterial survival during aminoglycoside and silver stress (*17, 30, 66, 67*).
Consistent with these findings, we made similar observations as the addition of either
catalase or thiourea to AGXX®-aminoglycoside-treated PA14 resulted in a significant
reduction in endogenous ROS level as well as bacterial killing relative to the combined
treatment alone **(FIG 4A, B; Supplementary FIG S5A, B)**. Consistent with our previous
report that AGXX® causes protein aggregation (*68*), we found significant transcriptional
upregulation of *ibpA*, which encodes the molecular chaperone IbpA*,* during exposure to
AGXX® and Gm. Given that IbpA serves as the front line defense to limit the aggregation
of unfolded proteins (*44*), AGXX®/Gm-induced oxidative stress likely causes an
imbalance of proteostasis and increases the cellular demand for chaperone expression.
Interestingly, heat-shock mediated protein aggregation was found to exponentially
increase aminoglycoside lethality in several gram-negative bacteria (*69*). Like our study,
this novel mechanism was attenuated under ROS quenching conditions (*69*), providing
additional evidence for the relevance of ROS in aminoglycoside killing. Considering that
individual AGXX® or gentamicin treatments did not result in a significant increase in ROS
signals, it is plausible that applying these antimicrobials at concentrations far below the
MIC would not induce the physiological/metabolic alterations necessary for aggravating
endogenous ROS levels.
517
Considering that both O_2^- and silver ions have been reported to destabilize Fe/S cluster
proteins resulting in the release of solvent-exposed iron (*10, 70*), we investigated the
effects of individual and combinational treatments with AGXX® and/or aminoglycoside
antibiotics on the activity of aconitase, an Fe-S cluster containing enzyme. We found
aconitase to be extremely sensitive towards exposure to both antimicrobials when
administered in combination (**FIG 6A**). Disrupted Fe-S clusters could potentially lead to
an increase in intracellular iron levels and subsequent •OH generation through Fenton
reaction as it has been demonstrated under hydrogen peroxide stress in ROS-sensitive

E. coli strains (*71*). This would explain the significantly higher •OH levels detected in PA14

cells that were exposed to a combined AGXX® and gentamicin mixture **(FIG 4D)**, which

could cause significant DNA damage and induce the SOS response. In fact, we found

that the combination of sublethal AGXX® and gentamicin significantly induced the

expression of *sulA,* an SOS response marker of DNA damage **(FIG 4E)**. Likewise,

addition of 2',2' bipyridyl resulted in a 1-log increase in bacterial survival when cells were

exposed to the toxic cocktail of gentamicin and AGXX® suggesting that free iron

contributes to the bactericidal effects **(FIG 6B)**. However, a similar protective effect was

observed in the presence of lethal Gm concentrations, suggesting that the possibility that

the rescuing effect in the presence of BIP could be due to its limiting effect on

aminoglycoside uptake and not necessarily a quenching of hydroxyl radical generation

through Fenton reactions. The Ezraty lab has demonstrated that bipyridyl inhibits Fe-S

biosynthesis under iron limiting conditions, resulting in reduced respiratory complex I and

II activities and decreased PMF, which is necessary for aminoglycoside uptake (*28*). All

in all, our findings highlight the relevance of oxidative stress in the enhancement of

antibiotic lethality in *P. aeruginosa* exposed to sub-inhibitory concentrations of AGXX®

and aminoglycosides.

Compared to other gram-negative pathogens, *P. aeruginosa* possesses high intrinsic

resistance mechanisms towards a variety of antibiotics (*72*). This resistance is mediated

in part by its extensive arsenal of efflux pumps and significantly lower outer membrane

permeability (*72*). We found that combining AGXX® and gentamicin increased both inner

and outer membrane permeability in *P. aeruginosa* leading to a subsequent increase in

aminoglycoside uptake **(FIG 7, FIG 8)**. Aminoglycosides are polycationic in nature and

known to displace divalent cations that cross-bridge lipopolysaccharides, which makes

the outer membrane substantially more permeable (*73*, *74*). It is therefore possible that

AGXX®, much like aminoglycosides, disrupts the outer leaflet of the outer membrane,

with the result that lower amounts of aminoglycosides are required to permeabilize *P.*

aeruginosa. If this were true, we would suspect that the initial stages of aminoglycoside

uptake could likely be accelerated causing a more rapid antibiotic influx. Surprisingly, we

found that the addition of sublethal polymyxin B concentrations did not affect gentamicin

lethality indicating that the mechanism behind the potentiating effects of AGXX® extends
beyond increased membrane permeability **(Supplementary FIG S6D)**. We further found
that disrupting membrane potential using an ionophore increased bacterial survival under
combined AGXX® and gentamicin stress **(FIG 8)**, suggesting an important role for the
energy dependent phase of aminoglycoside uptake AGXX®'s potentiating effect, as has
been found with strategies that increase aminoglycoside lethality via TCA cycle mediated
PMF generation (*55*).

By definition, antibiotic adjuvants do not possess inherent antimicrobial activity (*75*). Even
though AGXX® can technically not be defined as an antibiotic adjuvant given its own
antimicrobial activity (*16*), our use of sublethal AGXX® concentrations to screen for
potentiating effects satisfied this definition. Notably, the antimicrobial properties of silver
have been accepted for a long time and taken advantage of in topical ointments such as
silverdene cremes to prevent and treat *P. aeruginosa* infections in burn wound patients
(*26*). A comparison to silver nitrate and silver sulfadiazine (silverdene) revealed an
increased potency of some of the AGXX® formulations resulting in a lower survival rate
of PA14 **(FIG 1)**. Moreover, given that antibacterial combination therapies typically involve
significantly lower concentrations of the antimicrobials compared to what is needed for
individual treatments, such approaches would potentially delay resistance development.
575

MATERIALS AND METHODS

Bacterial strains and growth conditions. Unless stated otherwise, overnight *P.*
aeruginosa PA14 strains **(Supp. Table 2)** were grown under aerobic conditions in Luria-
Bertani broth (LB, Millipore Sigma) at 37°C for 16-20 hours at 300 rpm. For subsequent
assays, overnight cultures were diluted into Mueller-Hinton broth (MHB) or 3-(N-
morpholino) propanesulfonic acid minimal media containing 0.2% glucose, 1.32 mM
K_2HPO_4 and 10 µM thiamine (MOPSg) and incubated at 37 °C at 150 rpm. Growth under
anaerobic conditions was performed as described previously (*76*). In brief, PA14 was
grown anaerobically in the presence of 1% KNO_3 in a 50 ml falcon tube that was
additionally sealed with parafilm. Growth was performed in DuoClick™ (Thomas
Scientific) screw cap culture tubes filled with minimal room. Overnight PA14 cultures were

diluted to an OD_{600}=0.002 in MHB containing 1% KNO_3 and grown until OD_{600} ~0.2 after

which the treatments were applied for four hours.

Minimum Inhibitory Concentration (MIC). MIC assays were performed in 96-well plates

in a total volume of 200 µl per well. Overnight PA14 cultures were diluted into MHB to an

OD_{600} of 0.002 and distributed into 96 well plates that contained increasing concentrations

of ciprofloxacin, nalidixic acid, carbenicillin, imipenem, trimethoprim, polymyxin B,

gentamicin, amikacin, tobramycin, streptomycin, and kanamycin, respectively. Plates

were subsequently incubated at 37 °C for 16 hours at 300 rpm. MIC assays were

performed in duplicates. The MIC is defined as the lowest antibiotic concentration that

inhibited growth.

Time Killing Assays. Overnight PA14 cultures were diluted ~25-fold into MHB by

normalizing cultures to an OD_{600} = 0.1 in a 6-well sterile cell culture plate. Sublethal

concentrations of AGXX720C® and antibiotics of interest were added individually and in

combination as indicated and plates incubated at 37°C for 3 hours at 150 rpm under

aerobic conditions. For thiourea, CCCP and BIP pretreatments, overnight PA14 was

grown in MOPSg to exponential phase. Thiourea, CCCP or BIP was then added 60

minutes prior to AGXX®/aminoglycoside treatments whenever indicated. At 1-hour

intervals, OD_{600} of cultures were recorded, sample volumes were normalized to the lowest

optical density measured, serially diluted in PBS (pH = 7.4) and plated on LB agar to

quantify colony forming units (CFU) after 16 hours incubation. Percent survival was

calculated as the ratio of surviving colonies from treated to untreated samples.

Intracellular ROS level. Intracellular ROS levels were quantified using the redox-

sensitive dye 2',7'-dichlorodihydrofluorescein diacetate (H_2DCFDA) (Thermofisher

Scientific). Exponentially growing PA14 cultures were left untreated or treated with 0.25

µg/ml Gm, 50 µg/ml AGXX720C® or a combination of both for 1 hour. Samples were

collected and normalized to an OD_{600}=1.0. Cells were washed twice, resuspended in

prewarmed PBS containing 10 µM H_2DCFDA, and incubated in the dark at 37 °C for 30

min before samples were washed twice again in PBS and DCF fluorescence measured

at excitation/emission wavelengths of 485/535 nm (Tecan 200 plate reader). To quench
cellular ROS, cells were pretreated with 50 mM thiourea prior to the stress treatments.
620

Quantification of hydrogen peroxide level. The monoborate fluorescent probe peroxy
orange 1 (PO1) was used to measure hydrogen peroxide level. Exponentially growing
PA14 cultures were treated with 10 µM PO1 prior to their exposure to 0.25 µg/ml Gm, 50
µg/ml AGXX720C®, or a combination of both for 1 hour. Cells were washed twice and
resuspended in prewarmed PBS. Samples were analyzed using the flow cytometer (BD
FACS Melody™) in the PE-CF594 channel. At least 10,000 events were recorded, and
figures generated using FCSalyzer (*77*).

Quantification of Hydroxyl radical level. The fluorescent probe hydroxyphenyl
fluorescein (HPF; Invitrogen) was used to determine the amount of cellular hydroxyl
radicals (•OH) produced (*33*). Exponentially growing PA14 cultures were either left
untreated or treated for 1 hour with 0.25 µg/ml Gm, 50 µg/ml AGXX720C®, or the
combination thereof. Samples were collected and normalized to an OD_{600nm} of 1.0 after
washing twice with PBS (pH 7.4). Cells were then stained with 10 µM HPF and incubated
in the dark for 30 minutes at 37 °C. Cells were washed twice and resuspended in
prewarmed PBS. Samples were analyzed using the flow cytometer (BD FACS Melody™)
in the FITC channel. At least 10,000 events were recorded, and figures generated using
FCSalyzer (*77*).

Gene expression analyses by qRT-PCR. Overnight PA14 cultures were diluted into
MOPSg media to an OD_{600}=0.1 and grown to mid-log phase (OD_{600nm}=0.3). Cultures were
left untreated or treated with 0.25 µg/ml Gm, 50 µg/ml AGXX720C®, or the combination
thereof for 60 minutes. Transcription was stopped by the addition of an equal volume of
ice-cold methanol. Total RNA was extracted from three biological replicates using a
commercially available RNA extraction kit (Macherey & Nagel). Remaining DNA was
removed using the TURBO DNA-free kit (Thermo Scientific). mRNA was reverse
transcribed into cDNA using the PrimeScript cDNA synthesis kit (Takara). The following
primer pairs were used for gene amplification:

ibpA - TTCCGTCATTCCGTAGG/ AGGTCTTCTTCCTGG ;

sulA - ACTGTTCCAGGAAGCGTTCT/ AGCGAAAGTTCGCTAAAGGC;

rrsD - TATCAGATGAGCCTAGGTCGGATTA/ TTTACAATCCGAAGACCTTCTTCAC.

qRT-PCRs were set up according to the manufacturer's instructions (Alkali Scientific).

Transcript levels of the indicated genes were normalized against transcript levels of the

16S rRNA-encoding *rrsD* gene and relative fold changes in gene expression were

calculated using the $2^{-\Delta\Delta CT}$ method(*78*).

Aconitase Assay. Exponentially growing PA14 were treated with sublethal

concentrations of AGXX720C® (100 µg/ml), Gm (0.6 µg/ml) or the combination thereof.

Aconitase activity was measured from cell lysates using the Aconitase Assay Kit (Abcam)

according to manufacturer's instructions. One unit of aconitase is defined as the amount

of enzyme that isomerizes 1 µmol of citrate to isocitrate per min at pH 7.4 and 25 °C.

Outer membrane permeability assay. The N-phenyl-1-naphthylamine (NPN) uptake

assay was used to detect outer membrane damage as described in (*79*). Exponential

phase PA14 were treated with 0.25 µg/ml Gm, 50 µg/ml AGXX720C®, a combination

thereof. Addition of 2 µg/ml polymyxin B was used as a positive control. Cells were

collected after 60 min of treatment, washed, and resuspended to an OD_{600}=0.5 in HEPES-

sodium buffer (pH 7.2). 10 µM NPN was added, and samples incubated in the dark for 15

min. NPN fluorescence was measured at excitation/emission wavelengths of 350/420 nm.

Increased NPN uptake (indicating outer membrane damage) was calculated using the

following equation (*80*):

$$NPN\ uptake = \frac{Sample_{+NPN} - Sample_{-NPN}}{Hepes\ Buffer_{+NPN} - Hepes\ Buffer_{-NPN}}$$

Inner membrane disruption assay. Inner membrane integrity was determined by the

cellular uptake of propidium iodide following antimicrobial treatments. Exponentially

growing PA14 cells were either left untreated or treated with 0.25 µg/ml Gm, 50 µg/ml

AGXX720C®, or a combination thereof. Cells were harvested after 1 hour, washed twice,

and resuspended in PBS (pH 7.4) at an OD_{600}=0.5. Propidium iodide (Thermo Fisher

Scientific) was added to a final concentration of 500 nM and samples incubated in the
dark for 30 min. Fluorescence intensities were measured at excitation/emission
wavelengths of 535/617 nm. Samples treated with polymyxin B at a sublethal
concentration (2 μg/ml) were included as a positive control.
683
LIVE/DEAD staining. Exponentially growing PA14 cells were either left untreated or
treated with 0.25 µg/ml Gm, 50 µg/ml AGXX720C®, or a combination thereof. Cells were
harvested after 1 hour, washed twice, and resuspended in PBS (pH 7.4) at an OD_{600}=0.2.
Samples were stained with SYTO9 (6µM) and PI (30 µM) and incubated in the dark for
15 min at room temperature. Cells were then transferred onto a glass slide and covered
with 1% agarose prior to visualization using a Leica SP8 confocal system equipped with
a DMi8 CS inverted microscope. Samples treated with polymyxin B at a sublethal
concentration (2 μg/ml) were included as a positive control.
692
Texas Red-Gentamicin Uptake Assay. Texas Red-succinimidyl ester (Invitrogen) was
dissolved to a final concentration of 20 mg/ml in high quality anhydrous N,N-
dimethylformamide at 4 °C (*81*). Gm was dissolved in 100 mM K_2CO_3 (pH 8.5) to a final
concentration of 10 mg/ml at 4 °C. 20 µl Texas Red was slowly added dropwise to 700 ul
Gm to allow for the conjugation reaction to obtain the Texas-Red labeled gentamicin (TR-
Gm) (*81*). The TR-Gm conjugate was diluted in water and stored at -20 °C protected from
light. Exponentially growing PA14 cells were first treated with 1 µg/ml TR-Gm followed by
exposure to either 50 µg/ml AGXX720C® or 2 µg/ml polymyxin B (positive control). After
1 hour incubation in the dark at 37 °C, cells were collected and analyzed in the flow
cytometer (BD FACS Melody™) using the PE-Cy5 channel. At least 10,000 events were
recorded, and figures generated using FCSalyzer (*77*).
704
705 **Statistical analyses:** All statistical analyses were performed in GraphPad prism version
706 8.0.
707

ACKNOWLEDGEMENTS

This work was supported by the NIAID grant R15AI164585 and the Illinois State University Pre-Tenure Faculty Initiative Grant (to J.-U. D.). G.Y.D. was supported by Weigel and Mockford-Thompson by the Phi-Sigma Biological Sciences Honors Society, and the SIGMA Xi and BIRDFeeder grants. G.M.A. and C.D.O. were supported by the Illinois State University Undergraduate Research Support Program. Dr. Uwe Landau and his team from Largentec GmbH are acknowledged for scientific discussion and providing the AGXX® formulations. We further thank the ISU Confocal Microscopy Facility which was funded by NSF grant DBI-1828136, for their support. We also thank Sadia Sultana for critically proof-reading the manuscript and providing invaluable feedback. Figure 9 was generated with Adobe Illustrator.

FIGURE LEGENDS

FIG 1 AGXX394® is more efficient in killing *P. aeruginosa* than silverdene and silver nitrate. Overnight PA14 cultures were diluted ~25-fold into MOPSg media (OD_{600} = 0.1) and treated for three hours with 25 µg/ml $AgNO_3$ (black square), silverdene (grey square), and AGXX394® (red square), respectively. Colony survival was evaluated every hour by serially diluting samples and plating them onto LB agar. Percent survival was calculated relative to the untreated control (*n=4, ± S.D).*

FIG 2 AGXX® exponentially increases the activity of aminoglycosides against *P. aeruginosa*. In our time killing assay, overnight PA14 cultures were diluted ~25-fold (OD600 = 0.1) into MHB and exposed to 75 µg/ml AGXX720C® (black bars), a sublethal concentration of the indicated antibiotic (grey bar), or the combined treatment of both (red bars) for 3 hours. Samples were taking every 60 min, serial diluted, plated on LB agar, and incubated for 20 hours for CFU counts. Percent survival was calculated relative to the untreated control for: (A) 35 ng/ml ciprofloxacin and 100 ng/ml nalidixic acid, respectively; (B) 78 µg/ml carbenicillin and 0.156 µg/ml imipenem, respectively; (C) 125 µg/ml trimethoprim; (D) 1.5 µg/ml polymyxin B; and (E) 0.4 µg/ml gentamicin, 2.0 µg/ml amikacin, 1.0 µg/ml tobramycin, and 3 µg/ml streptomycin, respectively. All experiments

were performed in at least three biological replicates and error bars represent mean (±

SD. * p < 0.05, ** p < 0.01, *** p < 0.001; student's t test, calculated relative to cultures

treated with antibiotics alone).

FIG 3 AGXX® increases the sensitivity of *P. aeruginosa* strain PA14 to kanamycin.

Overnight PA14 cultures were diluted ~25-fold into Mueller-Hinton media (OD_{600}=0.1) and

either left untreated or exposed to 75 µg/ml AGXX720C®, 50 µg/ml kanamycin, or the

combination thereof for 3 hours. Samples were taken every 60 minutes, serial diluted and

plated on LB agar for CFU counts, (*n=4, ± S.D*).

FIG 4 The combination of sublethal concentrations of AGXX® and gentamicin

increases ROS levels and causes DNA damage and protein aggregation. Mid-log

PA14 cells were treated with sublethal concentrations of Gm (0.25 µg/ml), AGXX720C®

(50 µg/ml), the combination thereof, or left untreated. **(A)** Intracellular ROS levels were

quantified by H_2DCFDA fluorescence. 50mM thiourea was used as a ROS quencher (*n=3,*

± S.D.). **(B)** Samples were serially diluted in PBS after 60 min of incubation, spot-titered

onto LB agar and incubated for 20 hours. One representative of three independent

experiments with similar outcomes. **(C, D)** Cells were strained with **(C)** 10 µM peroxy-

orange 1 (PO1) and **(D)** 10µM hydroxyphenyl fluorescein (HPF) for 60 min and

fluorescence was measured via flow cytometry (*n=3, ± S.D.)*. **(E)** The induction of *sulA*

(white bar) and *ibpA* (black bar) transcript levels was determined by qRT-PCR. Gene

expression was normalized to the housekeeping gene *rrsD* and calculated as fold

changes based on expression levels in the untreated control (*n=3, ± S.D.*; One-way

ANOVA, Dunnett's posttest; *ns=p>0.05, * p < 0.05, ** p < 0.01, *** p < 0.001, **** p <*

0.0001).

FIG 5 Anaerobic growth and antioxidant systems provide protection against ROS-

mediated damage caused by a combinational treatment of AGXX® and

aminoglycosides. (A) PA14 were diluted approximately 1000-fold (OD_{600}=0.002) in MHB

supplemented with 1% KNO_3 and grown under aerobic (black bars) and anaerobic (grey

bars) conditions, respectively. At OD_{600} ~0.2, cultures were treated with either 0.5 µg/ml

or 1.0 µg/ml Gm alone or in combination with 75 µg/ml AGXX® for four hours. Cells were subsequently plated onto LB agar for 20 hours at 37°C to enumerate surviving colonies (*n=3, ± S.D.*). **(B)** Overnight cultures of PA14 wildtype and mutant strains with MrT7 transposon insertions in *oxyR, katA, katB* and *dps* were diluted into MOPSg to an OD_{600}=0.01 and grown under aerobic conditions until OD_{600}=0.1. Cultures were either left untreated (grey bars) or treated with a combination of 0.25 µg/ml tobramycin and 25 µg/ml AGXX720C® (white bars). Bacterial survival was quantified after two hours by serially diluting cells in PBS and plating on LB agar for 20 hours at 37°C (*n=3, ± S.D*; Student t-test, * *p* < 0.05).

FIG 6 The synergy between AGXX® and aminoglycosides on PA14 killing is in parts mediated by a disruption in iron homeostasis. Overnight PA14 cultures were diluted into MOPSg and incubated under aerobic conditions until exponential phase was reached. **(A)** Cells were left untreated or treated with 100 µg/ml AGXX720C®, 0.6 µg/ml Gm, or the combination thereof for 1 hour. Aconitase activities were determined in crude extracts (*n=4, ± S.D*). One-way ANOVA, Dunnett's posttest; *ns p>0.05,* ** *p* < 0.01. **(B)** Cultures were either left untreated (white square) or treated with 0.25 µg/ml Gm (white diamond), 50µg/ml AGXX720C® (white circle), or the combination thereof (white triangle) for three hours. Survival was determined each hour by serially diluting samples in PBS and plating onto LB agar for overnight growth. The impact of free iron on the increased killing by AGXX®/Gm cotreatments was tested by the absence (white triangle) and presence (black diamonds) of 125 µM 2',2' bipyridyl (*n=3, ± S.D.*). Student's t test, * *p*<0.05.

 FIG 7 Combined AGXX® and aminoglycoside treatment induces significant membrane damage. PA14 cells grown to mid-log phase in MOPSg media were left untreated or treated with sublethal concentrations of Gm (0.25 µg/ml), AGXX720C® (50 µg/ml), and the combination thereof. Cells were harvested after 1 hour of treatment, washed in PBS, and stained with **(A)** 10 µM N-phenyl-1-naphthylamine (NPN) dye, and **(B)** 0.5 µM propidium iodide (PI). Fluorescence intensities were determined at excitation/emission wavelengths of **(A)** 350/420 nm and **(B)** 535/617 nm, respectively,

($n=4$, $\pm$ S.D.). One-way ANOVA, Dunnett's posttest; ns $p>0.05$, * $p < 0.05$, **** $p < 0.0001$.

(C) Samples were washed in PBS, stained with PI/Syto9, incubated in the dark for 15

minutes at room temperature, mounted onto a glass slide with 1% agarose, and imaged

at 63x magnification via inverted confocal microscopy. One representative of three

independent experiments with similar outcomes.

FIG 8 AGXX® increases aminoglycoside uptake and lethality through increased

activity of the proton motive force (PMF). (A) Mid-log PA14 cells were treated with 1.0

µg/ml TR-Gm, 1.0 µg/ml TR-Gm + 50 µg/ml AGXX720C®, and 1.0 µg/ml TR-Gm + 2.0

µg/ml polymyxin B (PMB) for 1 hour, respectively. TR-Gm uptake was measured via flow

cytometry. **(B)** Mid-log phase PA14 were left untreated (control) or exposed to 0.25 µg/ml

Gm, 50 µg/ml AGXX720C® or 0.25 µg/ml Gm + 50 µg/ml AGXX720C® for 3 hours.

Samples were serial diluted, plated on LB agar, and incubated for 20 hours for CFU

counts. To test the impact of the PMF on the killing of a combination of AGXX® and

aminoglycosides, PA14 were pretreated with or without 50 µM carbonyl cyanide m-

chlorophenyl hydrazone (CCCP) prior to AGXX®/Gm exposure ($n=3$, $\pm$ S.D.). Student's t

test, ** $p<0.01$.

FIG 9 Proposed model for the synergy between AGXX® and aminoglycoside

antibiotics. The antimicrobial action of AGXX® is mediated by superoxide (O_2^-) and

hydrogen peroxide (H_2O_2), which are generated in a redox cycle between silver (Ag) and

ruthenium (Ru^{x+1}). The combination of AGXX® and aminoglycosides concertedly

increases endogenous ROS levels. As such anaerobic growth suppressed the synergistic

interaction between the two antimicrobials at lower Gm concentrations. Potentially

facilitated by a release of silver ions, the increased ROS level may disrupt iron-sulfur

clusters in metabolic enzymes such as aconitase, resulting in the release of free iron,

which ultimately triggers hydroxyl radical (•OH) formation in a Fenton reaction. Increasing

ROS levels can inflict macromolecule damage such as DNA damage and protein

aggregation, contributing to increased killing as observed for $P.$ $aeruginosa$ that were

treated with a combination of AGXX® and aminoglycosides. The synergistic effect

between AGXX® and aminoglycosides is also mediated by an increased uptake and the

cellular accumulation of aminoglycosides, which can be attenuated by disrupting the

bacterial membrane potential with ionophores such as CCCP.

References

1. C. Llor, L. Bjerrum, Antimicrobial resistance: risk associated with antibiotic overuse and initiatives to reduce the problem. *Ther. Adv. Drug Saf.* **5**, 229–41 (2014).

2. B. Spellberg, The future of antibiotics. *Crit. Care.* **18**, 228 (2014).

3. K. H. Turner, J. Everett, U. Trivedi, K. P. Rumbaugh, M. Whiteley, Requirements for Pseudomonas aeruginosa Acute Burn and Chronic Surgical Wound Infection. *PLOS Genet.* **10**, e1004518 (2014).

4. Z. Pang, R. Raudonis, B. R. Glick, T. J. Lin, Z. Cheng, Antibiotic resistance in Pseudomonas aeruginosa: mechanisms and alternative therapeutic strategies. *Biotechnol. Adv.* **37**, 177–192 (2019).

5. I. Chopra, L. Hesse, A. J. O'Neill, Exploiting current understanding of antibiotic action for discovery of new drugs. *Symp. Ser. Soc. Appl. Microbiol.* **31**, 4S-15S (2002).

6. D. Shang, Y. Liu, F. Jiang, F. Ji, H. Wang, X. Han, Synergistic Antibacterial Activity of Designed Trp-Containing Antibacterial Peptides in Combination With Antibiotics Against Multidrug-Resistant Staphylococcus epidermidis. *Front. Microbiol.* **0**, 2719 (2019).

7. F. Barras, L. Aussel, B. Ezraty, Silver and antibiotic, new facts to an old story. *Antibiotics.* **7**, 1–10 (2018).

8. S. Georgios A, P. Sotiris E, Antibacterial activity of nanosilver ions and particles. *Environ. Sci. Technol.* **44**, 5649–5654 (2010).

9. S. Y. Liau, D. C. Read, W. J. Pugh, J. R. Furr, A. D. Russell, Interaction of silver nitrate with readily identifiable groups: Relationship to the antibacterial action of silver ions. *Lett. Appl. Microbiol.* **25**, 279–283 (1997).

10. F. F. Xu, J. A. Imlay, Silver(I), Mercury(II), Cadmium(II), and Zinc(II) Target Exposed Enzymic Iron-Sulfur Clusters when They Toxify Escherichia coli. *Appl. Environ. Microbiol.* **78**, 3614 (2012).

11. D. Church, S. Elsayed, O. Reid, B. Winston, R. Lindsay, Burn wound infections. *Clin. Microbiol. Rev.* **19**, 403–434 (2006).

12. E. Clauss-Lendzian, A. Vaishampayan, A. de Jong, U. Landau, C. Meyer, J. Kok, E. Grohmann, Stress response of a clinical Enterococcus faecalis isolate subjected to a novel antimicrobial surface coating. *Microbiol. Res.* **207**, 53–64 (2018).

13. V. V. Loi, T. Busche, T. Preuß, J. Kalinowski, J. Bernhardt, H. Antelmann, The AGXX® Antimicrobial Coating Causes a Thiol-Specific Oxidative Stress Response

and Protein S-bacillithiolation in Staphylococcus aureus. *Front. Microbiol.* **9**, 3037 (2018).

14. L.-Y. Sobisch, K. M. Rogowski, J. Fuchs, W. Schmieder, A. Vaishampayan, P. Oles, N. Novikova, E. Grohmann, Biofilm Forming Antibiotic Resistant Gram-Positive Pathogens Isolated From Surfaces on the International Space Station. *Front. Microbiol.* **10**, 543 (2019).

15. N. Linzner, H. Antelmann, The Antimicrobial Activity of the AGXX® Surface Coating Requires a Small Particle Size to Efficiently Kill Staphylococcus aureus. *Front. Microbiol.* **12** (2021), doi:10.3389/FMICB.2021.731564/FULL.

16. A. Guridi, A. K. Diederich, S. Aguila-Arcos, M. Garcia-Moreno, R. Blasi, M. Broszat, W. Schmieder, E. Clauss-Lendzian, T. Sakinc-Gueler, R. Andrade, I. Alkorta, C. Meyer, U. Landau, E. Grohmann, New antimicrobial contact catalyst killing antibiotic resistant clinical and waterborne pathogens. *Mater. Sci. Eng. C*. **50**, 1–11 (2015).

17. J. Ruben Morones-Ramirez, J. A. Winkler, C. S. Spina, J. J. Collins, Silver enhances antibiotic activity against gram-negative bacteria. *Sci. Transl. Med.* **5** (2013), doi:10.1126/scitranslmed.3006276.

18. L. Kalan, G. D. Wright, Antibiotic adjuvants: multicomponent anti-infective strategies. *Expert Rev. Mol. Med.* **13** (2011), doi:10.1017/S1462399410001766.

19. A. Huttner, J. Bielicki, M. N. Clements, N. Frimodt-Møller, A. E. Muller, J.-P. Paccaud, J. W. Mouton, Oral amoxicillin and amoxicillin–clavulanic acid: properties, indications and usage. *Clin. Microbiol. Infect.* **26**, 871–879 (2020).

20. M. P. Brynildsen, J. A. Winkler, C. S. Spina, I. C. MacDonald, J. J. Collins, Potentiating antibacterial activity by predictably enhancing endogenous microbial ROS production. *Nat. Biotechnol.* **31**, 160 (2013).

21. J. J. Foti, B. Devadoss, J. A. Winkler, J. J. Collins, G. C. Walker, Oxidation of the guanine nucleotide pool underlies cell death by bactericidal antibiotics. *Science*. **336**, 315–319 (2012).

22. G. Soothill, Y. Hu, A. Coates, Can We Prevent Antimicrobial Resistance by Using Antimicrobials Better? *Pathog. 2013 Vol 2 Pages 422-435*. **2**, 422–435 (2013).

23. Q. L. Feng, J. Wu, G. Q. Chen, F. Z. Cui, T. N. Kim, J. O. Kim, A mechanistic study of the antibacterial effect of silver ions on Escherichia coli and Staphylococcus aureus. *J. Biomed. Mater. Res.* **52**, 662–668 (2000).

24. S. Silver, L. T. Phung, G. Silver, Silver as biocides in burn and wound dressings and bacterial resistance to silver compounds. *J. Ind. Microbiol. Biotechnol.* **33**, 627–634 (2006).

25. S. Egger, R. P. Lehmann, M. J. Height, M. J. Loessner, M. Schuppler, Antimicrobial Properties of a Novel Silver-Silica Nanocomposite Material. *Appl. Environ. Microbiol.* **75**, 2973–2976 (2009).

26. F. W. Fuller, The side effects of silver sulfadiazine. *J. Burn Care Res. Off. Publ. Am. Burn Assoc.* **30**, 464–470 (2009).

27. H. Kim, X. Xue, Detection of Total Reactive Oxygen Species in Adherent Cells by 2',7'-Dichlorodihydrofluorescein Diacetate Staining. *J. Vis. Exp. JoVE.* **2020**, 1–5 (2020).

28. M. Herisse, Y. Duverger, I. Martin-Verstraete, F. Barras, B. Ezraty, Silver potentiates aminoglycoside toxicity by enhancing their uptake. *Mol. Microbiol.* **105**, 115–126 (2017).

29. M. A. Kohanski, A. Tharakan, A. P. Lane, M. Ramanathan, Bactericidal Antibiotics Promote Reactive Oxygen Species Formation and Inflammation in Human Sinonasal Epithelial Cells. *Int. Forum Allergy Rhinol.* **6**, 191 (2016).

30. M. A. Kohanski, D. J. Dwyer, B. Hayete, C. A. Lawrence, J. J. Collins, A Common Mechanism of Cellular Death Induced by Bactericidal Antibiotics. *Cell.* **130**, 797–810 (2007).

31. M. J. Kelner, R. Bagnell, K. J. Welch, Thioureas react with superoxide radicals to yield a sulfhydryl compound. Explanation for protective effect against paraquat. *J. Biol. Chem.* **265**, 1306–1311 (1990).

32. B. C. Dickinson, C. Huynh, C. J. Chang, A Palette of Fluorescent Probes with Varying Emission Colors for Imaging Hydrogen Peroxide Signaling in Living Cells. *J. Am. Chem. Soc.* **132**, 5906 (2010).

33. M. Price, J. J. Reiners, A. M. Santiago, D. Kessel, Monitoring Singlet Oxygen and Hydroxyl Radical Formation with Fluorescent Probes During Photodynamic Therapy. *Photochem. Photobiol.* **85**, 1177–1181 (2009).

34. S. Sultana, J.-U. Dahl, A Year at the Forefront of Bacterial Defense Systems Against Neutrophilic Oxidants. *Biol. Open.* **12**, bio059809 (2023).

35. S. Sultana, A. Foti, J.-U. Dahl, Bacterial Defense Systems against the Neutrophilic Oxidant Hypochlorous Acid. *Infect. Immun.* **88** (2020), doi:10.1128/IAI.00964-19.

36. M. Varatnitskaya, A. Degrossoli, L. I. Leichert, Redox regulation in host-pathogen interactions: thiol switches and beyond. *Biol. Chem.* **402**, 299–316 (2021).

37. A. Sen, J. A. Imlay, How Microbes Defend Themselves From Incoming Hydrogen Peroxide. *Front. Immunol.* **12**, 667343 (2021).

38. M. J. Gray, W.-Y. Wholey, U. Jakob, Bacterial Responses to Reactive Chlorine Species. *Annu. Rev. Microbiol.* **67**, 141–160 (2013).

39. J.-U. Dahl, M. J. Gray, U. Jakob, Protein Quality Control under Oxidative Stress Conditions. *J. Mol. Biol.* **427**, 1549–1563 (2015).

40. B. Ezraty, A. Gennaris, F. Barras, J. F. Collet, Oxidative stress, protein damage and repair in bacteria. *Nat. Rev. Microbiol. 2017 157.* **15**, 385–396 (2017).

41. M. S. Cooke, M. D. Evans, M. Dizdaroglu, J. Lunec, Oxidative DNA damage: mechanisms, mutation, and disease. *FASEB J.* **17**, 1195–1214 (2003).

42. B. Groitl, J.-U. Dahl, J. W. Schroeder, U. Jakob, Pseudomonas aeruginosa defense systems against microbicidal oxidants. *Mol. Microbiol.* **106**, 335–350 (2017).

43. S. Sultana, M. E. Crompton, K. Meurer, O. Jankiewicz, G. H. Morales, C. Johnson, E. Horbach, K. P. Hoffmann, P. Kr, R. Shah, G. M. Anderson, N. T. Mortimer, J. E. Schmitz, M. Hadjifrangiskou, A. Foti, J.-U. Dahl, Redox-Mediated Inactivation of the Transcriptional Repressor RcrR is Responsible for Uropathogenic Escherichia coli's Increased Resistance to Reactive Chlorine Species. *mBio.* **13**, e01926-22 (2022).

44. M. Kitagawa, Y. Matsumura, T. Tsuchido, Small heat shock proteins, IbpA and IbpB, are involved in resistances to heat and superoxide stresses in Escherichia coli. *FEMS Microbiol. Lett.* **184**, 165–171 (2000).

45. K. Lewis, Programmed Death in Bacteria. *Microbiol. Mol. Biol. Rev.* **64**, 503–514 (2000).

46. N. T. Liberati, J. M. Urbach, S. Miyata, D. G. Lee, E. Drenkard, G. Wu, J. Villanueva, T. Wei, F. M. Ausubel, An ordered, nonredundant library of Pseudomonas aeruginosa strain PA14 transposon insertion mutants. *Proc. Natl. Acad. Sci.* **103**, 2833–2838 (2006).

47. K. Keyer, J. A. Imlay, Superoxide accelerates DNA damage by elevating free-iron levels. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 13635–13640 (1996).

48. S. I. Liochev, I. Fridovich, The role of O2.- in the production of HO.: in vitro and in vivo. *Free Radic. Biol. Med.* **16**, 29–33 (1994).

49. H. W. Taber, J. P. Mueller, P. F. Miller, A. S. Arrow, Bacterial uptake of aminoglycoside antibiotics. *Microbiol. Rev.* **51**, 439–457 (1987).

50. W. Kim, W. Zhu, G. L. Hendricks, D. Van Tyne, A. D. Steele, C. E. Keohane, N. Fricke, A. L. Conery, S. Shen, W. Pan, K. Lee, R. Rajamuthiah, B. B. Fuchs, P. M. Vlahovska, W. M. Wuest, M. S. Gilmore, H. Gao, F. M. Ausubel, E. Mylonakis, A new class of synthetic retinoid antibiotics effective against bacterial persisters. *Nat. 2018 5567699.* **556**, 103–107 (2018).

51. D. Pletzer, S. C. Mansour, R. E. W. Hancock, Synergy between conventional antibiotics and anti-biofilm peptides in a murine, sub-cutaneous abscess model caused by recalcitrant ESKAPE pathogens. *PLOS Pathog.* **14**, e1007084 (2018).

52. C. R. MacNair, J. M. Stokes, L. A. Carfrae, A. A. Fiebig-Comyn, B. K. Coombes, M. R. Mulvey, E. D. Brown, Overcoming mcr-1 mediated colistin resistance with colistin in combination with other antibiotics. *Nat. Commun. 2018 91*. **9**, 1–8 (2018).

53. D. J. Arndt-Jovin, T. M. Jovin, Chapter 16 Fluorescence Labeling and Microscopy of DNA. *Methods Cell Biol.* **30**, 417–448 (1989).

54. L. E. Bryan, H. M. Van Den Elzen, Streptomycin accumulation in susceptible and resistant strains of Escherichia coli and Pseudomonas aeruginosa. *Antimicrob. Agents Chemother.* **9**, 928–938 (1976).

55. S. Meylan, C. B. M. Porter, J. H. Yang, P. Belenky, A. Gutierrez, M. A. Lobritz, J. Park, S. H. Kim, S. M. Moskowitz, J. J. Collins, S. M. Conceptualization, J. J. C. ; Methodology, J. J. C. Investigation, F. Acquisition, J. J. C. ; Resources, S. M. M. ; Supervision, Carbon Sources Tune Antibiotic Susceptibility in Pseudomonas aeruginosa via Tricarboxylic Acid Cycle Control HHS Public Access. *Cell Chem Biol*. **24**, 195–206 (2017).

56. S. I. Liochev, The mechanism of "Fenton-like" reactions and their importance for biological systems. A biologist's view. *undefined*. **36**, 1–39 (1999).

57. B. Nowack, H. F. Krug, M. Height, 120 years of nanosilver history: implications for policy makers. *Environ. Sci. Technol.* **45**, 1177–1183 (2011).

58. W. K. Jung, H. C. Koo, K. W. Kim, S. Shin, S. H. Kim, Y. H. Park, Antibacterial Activity and Mechanism of Action of the Silver Ion in Staphylococcus aureus and Escherichia coli. *Appl. Environ. Microbiol.* **74**, 2171–2178 (2008).

59. R. B.-K. Wakshlak, R. Pedahzur, D. Avnir, Antibacterial activity of silver-killed bacteria: the "zombies" effect. *Sci. Rep.* **5**, 9555 (2015).

60. A. Panáček, L. Kvítek, M. Smékalová, R. Večeřová, M. Kolář, M. Röderová, F. Dyčka, M. Šebela, R. Prucek, O. Tomanec, R. Zbořil, Bacterial resistance to silver nanoparticles and how to overcome it. *Nat. Nanotechnol. 2017 131*. **13**, 65–71 (2017).

61. H. Wang, M. Wang, X. Xu, P. Gao, Z. Xu, Q. Zhang, H. Li, A. Yan, R. Y. T. Kao, H. Sun, Multi-target mode of action of silver against Staphylococcus aureus endows it with capability to combat antibiotic resistance. *Nat. Commun. 2021 121*. **12**, 1–16 (2021).

62. R. Tambosi, S. Liotenberg, M.-L. Bourbon, A.-S. Steunou, M. Babot, A. Durand, N. Kebaili, S. Ouchane, Silver and Copper Acute Effects on Membrane Proteins and

Impact on Photosynthetic and Respiratory Complexes in Bacteria. *mBio*. **9**, e01535-18 (2018).

63. A. Vaishampayan, A. de Jong, D. J. Wight, J. Kok, E. Grohmann, A novel antimicrobial coating represses biofilm and virulence-related genes in methicillin-resistant Staphylococcus aureus. *Front. Microbiol.* **9**, 1–14 (2018).

64. D. J. Dwyer, P. A. Belenky, J. H. Yang, I. Cody MacDonald, J. D. Martell, N. Takahashi, C. T. Y. Chan, M. A. Lobritz, D. Braff, E. G. Schwarz, J. D. Ye, M. Pati, M. Vercruysse, P. S. Ralifo, K. R. Allison, A. S. Khalil, A. Y. Ting, G. C. Walker, J. J. Collins, Antibiotics induce redox-related physi. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E2100–E2109 (2014).

65. J. Ling, C. Cho, L.-T. Guo, H. Aerni, J. Rinehart, D. Söll, Protein aggregation caused by aminoglycoside action is prevented by a hydrogen peroxide scavenger. *Mol. Cell*. **48**, 713–722 (2012).

66. J. Ye, Y. Su, X. Lin, S. Lai, W. Li, F. Ali, J. Zheng, B. Peng, Alanine Enhances Aminoglycosides-Induced ROS Production as Revealed by Proteomic Analysis. *Front. Microbiol.* **9** (2018) (available at https://www.frontiersin.org/articles/10.3389/fmicb.2018.00029).

67. X. Ji, J. Zou, H. Peng, A.-S. Stolle, R. Xie, H. Zhang, B. Peng, J. J. Mekalanos, J. Zheng, Alarmone Ap4A is elevated by aminoglycoside antibiotics and enhances their bactericidal activity. *Proc. Natl. Acad. Sci.* **116**, 9578–9585 (2019).

68. S. Sultana, G. M. Anderson, K. P. Hoffmann, J.-U. Dahl, Extraction and Visualization of Protein Aggregates after Treatment of Escherichia coli with a Proteotoxic Stressor. *JoVE J. Vis. Exp.*, e62628 (2021).

69. B. Lv, X. Huang, C. Lijia, Y. Ma, M. Bian, Z. Li, J. Duan, F. Zhou, B. Yang, X. Qie, Y. Song, T. K. Wood, X. Fu, Heat shock potentiates aminoglycosides against gram-negative bacteria by enhancing antibiotic uptake, protein aggregation, and ROS. *Proc. Natl. Acad. Sci.* **120**, e2217254120 (2023).

70. J. A. Imlay, *How oxygen damages microbes: Oxygen tolerance and obligate anaerobiosis* (2002), vol. 46.

71. S. Park, X. You, J. A. Imlay, Substantial DNA damage from submicromolar intracellular hydrogen peroxide detected in Hpx- mutants of Escherichia coli. *Proc. Natl. Acad. Sci.* **102**, 9317–9322 (2005).

72. K. Poole, Pseudomonas aeruginosa: Resistance to the max. *Front. Microbiol.* **2**, 65 (2011).

73. R. E. W. Hancock, S. W. Farmer, Z. Li, K. Poole, Interaction of aminoglycosides with the outer membranes and purified lipopolysaccharide and OmpF porin of Escherichia coli. *Antimicrob. Agents Chemother.* **35**, 1309–1314 (1991).

74. R. E. Hancock, V. J. Raffle, T. I. Nicas, Involvement of the outer membrane in gentamicin and streptomycin uptake and killing in Pseudomonas aeruginosa. *Antimicrob. Agents Chemother.* **19**, 777–785 (1981).

75. G. D. Wright, Antibiotic Adjuvants: Rescuing Antibiotics from Resistance. *Trends Microbiol.* **24**, 862–871 (2016).

76. M. Wu, T. Guina, M. Brittnacher, H. Nguyen, J. Eng, S. I. Miller, The Pseudomonas aeruginosa Proteome during Anaerobic Growth. *J. Bacteriol.* **187**, 8185–8190 (2005).

77. Mostböck, Sven, FCSalyzer. *SourceForge* (2021), (available at https://sourceforge.net/projects/fcsalyzer/).

78. M. W. Pfaffl, A new mathematical model for relative quantification in real-time RT–PCR. *Nucleic Acids Res.* **29**, e45 (2001).

79. I. M. Helander, T. Mattila-Sandholm, Fluorometric assessment of Gram-negative bacterial permeabilization. *J. Appl. Microbiol.* **88**, 213–219 (2000).

80. A. Sabnis, K. L. H. Hagart, A. Klöckner, M. Becce, L. E. Evans, R. C. D. Furniss, D. A. I. Mavridou, R. Murphy, M. M. Stevens, J. C. Davies, G. J. Larrouy-Maumus, T. B. Clarke, A. M. Edwards, Colistin kills bacteria by targeting lipopolysaccharide in the cytoplasmic membrane. *eLife*. **10** (2021), doi:10.7554/ELIFE.65836.

81. H. Saito, K. Inui, R. Hori, Mechanisms of gentamicin transport in kidney epithelial cell line (LLC-PK1). *J. Pharmacol. Exp. Ther.* **238** (1986).

FIG 1 AGXX394® is more efficient in killing *P. aeruginosa* than silverdene and silver nitrate. Overnight PA14 cultures were diluted ~25-fold into MOPSg media (OD_{600} = 0.1) and treated for three hours with 25 µg/ml $AgNO_3$ (black square), silverdene (grey square), and AGXX394® (red square), respectively. Colony survival was evaluated every hour by serially diluting samples and plating them onto LB agar. Percent survival was calculated relative to the untreated control (*n=4, ± S.D).*

FIG 2 AGXX® exponentially increases the activity of aminoglycosides against *P. aeruginosa*. In our time killing assay, overnight PA14 cultures were diluted ~25-fold (OD600 = 0.1) into MHB and exposed to 75 µg/ml AGXX720C® (black bars), a sublethal concentration of the indicated antibiotic (grey bar), or the combined treatment of both (red bars) for 3 hours. Samples were taking every 60 min, serial diluted, plated on LB agar, and incubated for 20 hours for CFU counts. Percent survival was calculated relative to the untreated control for: (A) 35 ng/ml ciprofloxacin and 100 ng/ml nalidixic acid, respectively; (B) 78 µg/ml carbenicillin and 0.156 µg/ml imipenem, respectively; (C) 125 µg/ml trimethoprim; (D) 1.5 µg/ml polymyxin B; and (E) 0.4 µg/ml gentamicin, 2.0 µg/ml amikacin, 1.0 µg/ml tobramycin, and 3 µg/ml streptomycin, respectively. All experiments were performed in at least three biological replicates and error bars represent mean (± SD. * p < 0.05, ** p < 0.01, *** p < 0.001; student's t test, calculated relative to cultures treated with antibiotics alone).

FIG 3 AGXX® increases the sensitivity of *P. aeruginosa* strain PA14 to kanamycin.
Overnight PA14 cultures were diluted ~25-fold into Mueller-Hinton media (OD_{600}=0.1) and either left untreated or exposed to 75 µg/ml AGXX720C®, 50 µg/ml kanamycin, or the combination thereof for 3 hours. Samples were taken every 60 minutes, serial diluted and plated on LB agar for CFU counts, (*n=4, ± S.D)*.

FIG 4 The combination of sublethal concentrations of AGXX® and gentamicin increases ROS levels and causes DNA damage and protein aggregation. Mid-log PA14 cells were treated with sublethal concentrations of Gm (0.25 µg/ml), AGXX720C® (50 µg/ml), the combination thereof, or left untreated. **(A)** Intracellular ROS levels were quantified by H_2DCFDA fluorescence. 50mM thiourea was used as a ROS quencher (*n=3, ± S.D.)*. **(B)** Samples were serially diluted in PBS after 60 min of incubation, spot-titered onto LB agar and incubated for 20 hours. One representative of three independent experiments with similar outcomes. **(C, D)** Cells were strained with **(C)** 10 µM peroxy-orange 1 (PO1) and **(D)** 10µM hydroxyphenyl fluorescein (HPF) for 60 min and fluorescence was measured via flow cytometry (*n=3, ± S.D.)*. **(E)** The induction of *sulA* (white bar) and *ibpA* (black bar) transcript levels was determined by qRT-PCR. Gene expression was normalized to the housekeeping gene *rrsD* and calculated as fold changes based on expression levels in the untreated control (*n=3, ± S.D.;* One-way

ANOVA, Dunnett's posttest; *ns=p>0.05,* * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$).

FIG 5 Anaerobic growth and antioxidant systems provide protection against ROS-mediated damage caused by a combinational treatment of AGXX® and aminoglycosides. (A) PA14 were diluted approximately 1000-fold (OD$_{600}$=0.002) in MHB supplemented with 1% KNO$_3$ and grown under aerobic (black bars) and anaerobic (grey bars) conditions, respectively. At OD$_{600}$ ~0.2, cultures were treated with either 0.5 µg/ml or 1.0 µg/ml Gm alone or in combination with 75 µg/ml AGXX® for four hours. Cells were subsequently plated onto LB agar for 20 hours at 37°C to enumerate surviving colonies (*n=3, ± S.D.*). **(B)** Overnight cultures of PA14 wildtype and mutant strains with MrT7 transposon insertions in *oxyR, katA, katB* and *dps* were diluted into MOPSg to an OD$_{600}$=0.01 and grown under aerobic conditions until OD$_{600}$=0.1. Cultures were either left untreated (grey bars) or treated with a combination of 0.25 µg/ml tobramycin and 25 µg/ml AGXX720C® (white bars). Bacterial survival was quantified after two hours by serially diluting cells in PBS and plating on LB agar for 20 hours at 37°C (*n=3, ± S.D*; Student t-test, * *p* < 0.05).

FIG 6 The synergy between AGXX® and aminoglycosides on PA14 killing is in parts mediated by a disruption in iron homeostasis. Overnight PA14 cultures were diluted into MOPSg and incubated under aerobic conditions until exponential phase was reached. **(A)** Cells were left untreated or treated with 100 µg/ml AGXX720C®, 0.6 µg/ml Gm, or the combination thereof for 1 hour. Aconitase activities were determined in crude extracts (*n=4, ± S.D*). One-way ANOVA, Dunnett's posttest; *ns p>0.05, ** p < 0.01*. **(B)** Cultures were either left untreated (white square) or treated with 0.25 µg/ml Gm (white diamond), 50µg/ml AGXX720C® (white circle), or the combination thereof (white triangle) for three hours. Survival was determined each hour by serially diluting samples in PBS and plating onto LB agar for overnight growth. The impact of free iron on the increased killing by AGXX®/Gm cotreatments was tested by the absence (white triangle) and presence (black diamonds) of 125 µM 2',2' bipyridyl (*n=3, ± S.D.*). Student's t test, * *p<0.05*).

FIG 7 Combined AGXX® and aminoglycoside treatment induces significant membrane damage. PA14 cells grown to mid-log phase in MOPSg media were left untreated or treated with sublethal concentrations of Gm (0.25 µg/ml), AGXX720C® (50 µg/ml), and the combination thereof. Cells were harvested after 1 hour of treatment, washed in PBS, and stained with **(A)** 10 µM N-phenyl-1-naphthylamine (NPN) dye, and **(B)** 0.5 µM propidium iodide (PI). Fluorescence intensities were determined at excitation/emission wavelengths of **(A)** 350/420 nm and **(B)** 535/617 nm, respectively, (*n=4, ± S.D.*). One-way ANOVA, Dunnett's posttest; *ns p>0.05,* * *p* < 0.05, **** *p* < 0.0001. **(C)** Samples were washed in PBS, stained with PI/Syto9, incubated in the dark for 15 minutes at room temperature, mounted onto a glass slide with 1% agarose, and imaged at 63x magnification via inverted confocal microscopy. One representative of three independent experiments with similar outcomes.

FIG 8 AGXX® increases aminoglycoside uptake and lethality through increased activity of the proton motive force (PMF). (A) Mid-log PA14 cells were treated with 1.0 µg/ml TR-Gm, 1.0 µg/ml TR-Gm + 50 µg/ml AGXX720C®, and 1.0 µg/ml TR-Gm + 2.0 µg/ml polymyxin B (PMB) for 1 hour, respectively. TR-Gm uptake was measured via flow cytometry. **(B)** Mid-log phase PA14 were left untreated (control) or exposed to 0.25 µg/ml Gm, 50 µg/ml AGXX720C® or 0.25 µg/ml Gm + 50 µg/ml AGXX720C® for 3 hours. Samples were serial diluted, plated on LB agar, and incubated for 20 hours for CFU counts. To test the impact of the PMF on the killing of a combination of AGXX® and aminoglycosides, PA14 were pretreated with or without 50 µM carbonyl cyanide *m*-chlorophenyl hydrazone (CCCP) prior to AGXX®/Gm exposure *(n=3, ± S.D.)*. Student's t test, ** $p<0.01$.

◆AGXX® ● Aminoglycoside

FIG 9 Proposed model for the synergy between AGXX® and aminoglycoside antibiotics. The antimicrobial action of AGXX® is mediated by superoxide (O_2^-) and hydrogen peroxide (H_2O_2), which are generated in a redox cycle between silver (Ag) and ruthenium (Ru^{x+1}). The combination of AGXX® and aminoglycosides concertedly increases endogenous ROS levels. As such anaerobic growth suppressed the synergistic interaction between the two antimicrobials at lower Gm concentrations. Potentially facilitated by a release of silver ions, the increased ROS level may disrupt iron-sulfur clusters in metabolic enzymes such as aconitase, resulting in the release of free iron, which ultimately triggers hydroxyl radical (•OH) formation in a Fenton reaction. Increasing ROS levels can inflict macromolecule damage such as DNA damage and protein aggregation, contributing to increased killing as observed for *P. aeruginosa* that were treated with a combination of AGXX® and aminoglycosides. The synergistic effect

between AGXX® and aminoglycosides is also mediated by an increased uptake and the cellular accumulation of aminoglycosides, which can be attenuated by disrupting the bacterial membrane potential with ionophores such as CCCP.