Mating systems and recombination landscape strongly shape genetic diversity and selection in wheat relatives

Concetta Burgarella[1,2,3]*, Marie-Fleur Brémaud[2], Gesa Von Hirschheydt[4], Veronique Viader[2], Morgane Ardisson[2], Sylvain Santoni[2], Vincent Ranwez[2], Miguel Navascués[5], Jacques David[2] and Sylvain Glémin[6,7]*

[1] CNRS, Univ Montpellier, ISEM– UMR 5554, 34095 Montpellier, France.

[2] AGAP Institut, Univ Montpellier, CIRAD, INRAE, Institut Agro, Montpellier, France.

[3] Department of Organismal Biology, Evolutionary Biology Center, Uppsala University, Uppsala, Sweden.

[4] Swiss Federal Research Institute WSL, Zürcherstrasse 111 8903 Birmensdorf, Switzerland.

[5] UMR CBGP, Univ Montpellier, CIRAD, INRAE, Institut Agro, Montpellier, France.

[6] CNRS, Univ. Rennes, ECOBIO - UMR 6553, F-35000 Rennes, France.

[7] Department of Ecology and Evolution, Evolutionary Biology Center, Uppsala University, Uppsala, Sweden.

*Corresponding authors

Concetta Burgarella: AGAP Institut, Univ Montpellier, CIRAD, INRAE, Institut Agro, Montpellier, France. E-mail: concetta.burgarella@inrae.fr

24 Sylvain Glémin: CNRS, Univ. Rennes, ECOBIO - UMR 6553, F-35000 Rennes, France.

25 Department of Ecology and Evolution, Evolutionary Biology Center, Uppsala University,

26 Uppsala, Sweden. E-mail: sylvain.glemin@univ-rennes.fr

29 **Keywords**: self-fertilization, polymorphism, linked selection, fitness effect of mutations,

30 selfing-syndrome

Abstract

How and why genetic diversity varies among species is a long-standing question in evolutionary biology. Life history traits have been shown to explain a large part of observed diversity. Among them, mating systems have one of the strongest impacts on genetic diversity, with selfing species usually exhibiting much lower diversity than outcrossing relatives. Theory predicts that a high rate of selfing amplifies selection at linked sites, reducing genetic diversity genome wide, but frequent bottlenecks and rapid population turn-over could also explain low genetic diversity in selfers. However, how linked selection varies with mating systems and whether it is sufficient to explain the observed difference between selfers and outcrossers has never been tested. Here, we used the *Aegilops*/*Triticum* grass species, a group characterized by contrasted mating systems (from obligate out-crossing to high selfing) and marked recombination rate variation across the genome, to quantify the effects of mating system and linked selection on patterns of neutral and selected polymorphism. By analyzing phenotypic and transcriptomic data of 13 species, we show that selfing strongly affects genetic diversity and the efficacy of selection by amplifying the intensity of linked selection genome wide. In particular, signatures of adaptation were only found in the highly recombining regions in outcrossing species. These results bear implications for the evolution of mating systems and more generally for our understanding of the fundamental drivers of genetic diversity.

Introduction

How and why genetic diversity varies among species is a central and long-standing question in evolutionary biology, dating back from the 1960's (Ellegren and Galtier, 2016). For neutral variation, patterns of genetic diversity depend on the balance between mutation and genetic drift, characterized by the effective size of a population, N_e, and also on the efficacy of selection for functional regions of the genome. Recently, thanks to the availability of population genomic data in many non-model species, several studies have explored the ecological correlates of diversity levels, usually measured as nucleotide polymorphism, π. These surveys have shown that life history traits (LHTs), especially life-span and reproductive mode, can explain a large part of the observed variation in genetic diversity among species (Romiguier *et al.*, 2014; Chen *et al.*, 2017; Mackintosh *et al.*, 2019; Muyle *et al.*, 2021). LHTs may reflect long-term effective population size, which depends on current population size and past fluctuations across generations (e.g. Romiguier *et al.*, 2014; Mackintosh *et al.*, 2019). N_e can also depend on selection at linked sites, i.e. the hitch-hiking effect of the fixation of beneficial or the removal of deleterious mutations on linked neutral variation (Cutter and Payseur, 2013), which also affects long-term N_e and seems rather pervasive across genomes (Corbett-Detig *et al.*, 2015; Mackintosh *et al.*, 2019; Chen *et al.*, 2020; Buffalo, 2021).

Among LHTs, mating systems deeply affect the genetic and ecological functioning of a species and are predicted to strongly impact both demographic outcomes and the response to selection. Thanks to the ability to produce seeds under limited mate availability, the capacity of autonomous selfing provides reproductive assurance and can be an ecologically successful strategy, allowing colonizing new habitats and increasing species range (Grossenbacher *et al.*, 2015), which should be associated with a large census population size. However, being able to reproduce alone implies a much higher demographic

stochasticity due to recurrent bottlenecks and colonization-extinction dynamics, which

strongly reduces genetic diversity, not only at the population scale but also at the whole

species scale (Pannell and Charlesworth, 1999; Ingvarsson, 2002). Moreover, the dynamics

of range expansions, which can be associated with the evolution of selfing, can also

unintuitively lead to the loss of diversity, especially on the expansion front (Excoffier et $al.$,

2009). So, despite possible large species range and census population size, the specific

ecology of selfing species may lead to a reduction in N_e. In addition to these demographic

effects, selfing also has direct genetic effects that can reduce N_e. Non-independent gamete

sampling during mating automatically increases genetic drift and reduces genetic mixing,

which generates genome-wide genetic linkage disequilibrium enhancing the effect of linked

selection (Agrawal and Hartfield, 2016; Roze, 2016; Hartfield and Bataillon, 2020).

So far, striking differences in genetic diversity have already been observed between

outcrossing and selfing relatives (e.g., Hazzouri et $al.$, 2013; Slotte et $al.$, 2013; Burgarella et

$al.$, 2015; Teterina et al., 2023) and the underlying causes (linked-selection, demographic

instability) have been often discussed and studied from a theoretical point of view

(Charlesworth et $al.$, 1993; Barrett et $al.$, 2014), but, to our knowledge, attempts at a direct

quantification with empirical data are recent and only partial (see the comparison between

two outcrossing and selfing $Caenorhabditis$ species in Teterina et al., 2023). Yet, how the

intensity of linked selection varies with mating system and whether it can be sufficient to

explain the observed difference between outcrossing and selfing species remains to be

quantified. Beyond a genomewide reduction in polymorphism and selection efficacy with

increasing selfing rates, theory also predicts that genomic patterns across chromosomes

should vary with the interaction between recombination and selfing rates. We expect a clear

positive relationship between genetic diversity and recombination in outcrossers, but an

increasingly flatter relationship in species with increasing selfing rates. We also expect that

deleterious mutations should accumulate mainly in lowly recombining regions whereas

adaptation should be prevalent in highly recombining regions in outcrossing species, in

contrast to selfing species where signatures of high deleterious load and low adaptation

should be more evenly distributed along the genome.

Here, we tested these hypotheses by comparing species with a large range of mating

systems occurring within a single genus, which allows strong genetic contrast among

otherwise similar species, an advocated sampling design (Leffler *et al.*, 2012; Cutter and

Payseur, 2013). We used the *Aegilops*/*Triticum* grass species as a study system. This group

of Mediterranean and Western/Central Asian grasses belongs to the Triticeae tribe

(Poaceae) and includes wheat and its wild relatives. The *Aegilops*/*Triticum* genus forms a

monophyletic group with 13 diploid and about 17 polyploid species that likely diversified ~4-7

millions years ago (Huang *et al.*, 2002; Marcussen *et al.*, 2014; Glémin *et al.*, 2019b). All

species are characterized by similar life-history traits (wind-pollinated, annual, herbaceous

species) and ecology (open landscapes, warm-temperate climate), but present a large

diversity of mating systems, spanning from obligate out-crossing to highly selfing species

(van Slageren, 1994; Kilian *et al.*, 2011) (**Fig. 1**). Triticeae genomes are large, with markedly

U-shaped recombination patterns along chromosomes conserved across species: most

recombination is located in the distal parts whereas no or very low recombination occurs in

their central part (Brazier and Glémin, 2022). Marked differences in both mating systems

among species and recombination rate within genomes make the group an ideal model to

unravel the role of selection on species genetic diversity.

Through a comparative population genomic approach, we assessed the expectation

that genetic diversity and selection efficacy decreases with higher selfing rate and with lower

recombination rate within each species, by controlling for species range that varies among

species (**Supplementary Fig. S1**). We then explicitly tested whether the effect of linked

selection was stronger in selfing species as predicted by population genetics theory. We

found that selfing strongly affects genetic diversity and the efficacy of selection by amplifying

the intensity of linked selection genome wide, while species range plays a minor role. We

also showed that genomic patterns remarkably matched the gradient of mating systems

across species, while models and empirical evidence so far suggested that only extreme

mating systems left clear signatures in the genomes (Agrawal and Hardfield 2016, Roze

2016). These results have multiple implications for both the evolution of mating systems and

our understanding of the fundamental drivers of genetic diversity.

Material and methods

Plant material

We analyzed molecular and morphological data of the 13 extant wild diploid species

of the *Aegilops*/*Triticum* genus (Triticeae tribe, Kilian *et al.*, 2011) and 3 outgroup species

(*Taeniatherum caput-medusae*, *Hordeum spontaneum* and *Secale strictum*) for a total of 98

accessions. We targeted 7-20 individuals from each of five focal species, including the two

self-incompatible outcrossing *Ae. speltoides* and *Ae. mutica* and the three predominantly

selfing *Ae. tauschii*, *T. urartu* and *T. monococcum*. We included 2-4 accessions from each of

the other *Aegilops*/*Triticum* species and 1-3 from each of the outgroups. The accessions

were obtained from several international seed banks and donor researchers. The list of

accessions per species with their passport information is provided in **Supplementary Table**

S1.

Morphological data

To finely characterize the selfing syndrome of each species, we measured several

morphofunctional traits describing the reproductive organs and function (Friedman and

Harder, 2005; Escobar *et al.*, 2010 and references therein). Around 3-4 grains per accession

were sown in January 2014 in a glass house. After emergence, the seedlings were

submitted to 4°C for 6 weeks to assure vernalization requirements. Only one seedling per

accession was kept after vernalization. The first two spikes of each plant were closed in

paper bags to prevent cross fertilization. We collected the two bagged spikes, one open

mature spike, mature anthers, stigmas and ovaries for measurements. Anthers, stigmas and

ovaries were preserved in Carnoy fixative solution. For each of 5-7 accessions per

Aegilops/*Triticum* species and 1-3 accessions per outgroup species, we measured a mature

spike (length, spikelet number, grain number), three spikelets per spike (spikelet length) and

three flowers from one spikelet (length of palea and lemma). All flowers of the three spikelets

were classified as fertile (if the presence of grain was observed), female (only stigma

observed), male (anthers observed), or sterile. For each accession, we measured 6 anthers

(length and width) and 3 stigmas and ovaries (length of each organ). For anthers and stigma

and ovaries, each organ was measured five times and the mean value of these replicates

was used in further analysis. Measures were manually recorded on millimeter paper or taken

on photographs with the software analySIS (Soft Imaging System GmbH 2002; see

Supplementary Fig. S12 for an example).

Missing data (31%) on the directly observed measures were imputed with the

missMDA package (Josse and Husson, 2016) under the R environment (R Core Team,

2018). Parameters were set by default and the optimal number of components retained for

imputation were estimated with the cross-validation method (ncp=3). Raw and imputed

measures are provided in **Supplementary tables S2** and **S3**, respectively and the list of

measured traits is provided in **Table S4**.

Additional variables were calculated on imputed measures as follows. The mean

values of anther and stigma dimensions were calculated per accession and standardized by

dividing by the flower length. Following Escobar *et al*. (2010), the autonomous seed set was

estimated   as   self-fertilised_seed_number/(self-

fertilized_spikelet_number*number_fertile_flower/spikelet), corresponding to the number of

seeds per fertile flower. Spikelet compactness was calculated as the ratio (mean flower

length* number_fertile_flower)/mean spikelet length. Male investment was calculated as the

ratio of the mean anther length and the mean ovary length. We used these additional

variables to summarize the selfing syndrome with a synthetic measure corresponding to the

first axis of a Principal Component Analysis (PCA) (**Supplementary Table S3** and **S4**). The

PCA was performed with the ade4 package (Dray and Dufour, 2007) under the R

environment.

Species range

190   We expect that species with bigger census sizes also harbor higher genetic diversity,

a relationship that could mask or interact with the effect of the mating system. To control for

this potential effect, we used species range as a proxy for census size, since, to our

knowledge, there are no direct estimates of census size for wild *Aegilops*/*Triticum* species.

To estimate species range, we retrieved occurrence data from the Global Biodiversity

Information Facility (http://www.gbif.org) for each species. We manually cleaned the data set

to remove single occurrences outside species range, which can be due either to

identification errors or recent introductions. Cleaned data were mapped on the world map

(focusing on western Eurasia and North Africa) on which we applied a grid with cell size of

one decimal degree square (~10,000 km^2). We estimated species range as the number of

cells occupied by a species time 10,000 km^2 (**Supplementary Figure S1 and Table S5**).

Sequencing

We added 48 new sequences to the dataset used for the phylogenomic analysis of Glémin *et al*. (2019b) and Clément *et al*. (2017), for a total of 98 sequences for 13 *Aegilops*/*Triticum* species (n=2-21) and 3 outgroup species (n=1-3).

We performed full transcriptome sequencing following the procedure described in Sarah *et al*. (2017) and Glémin *et al*. (2019b). Briefly, RNAs were extracted and prepared separately for leaves and inflorescence tissues, and mixed subsequently in 20% and 80% proportions, respectively. RNA was extracted using a Spectrum Plant Total RNA kit (Sigma-Aldrich, USA) with a DNAse treatment. RNA concentration was measured with two methods, a NanoDrop ND-1000 Spectrophotometer and the Quant-iT™ RiboGreen® (Invitrogen, USA) protocol. RNA quality was assessed on the RNA 6000 Pico chip on a Bioanalyzer 2100 (Agilent Technologies, USA). Following the Illumina TruSeq mRNA protocol, we kept samples with an RNA Integrity Number (RIN) value greater than eight. Libraries were prepared with a modified protocol of the TruSeq Stranded mRNA Library Prep Kit (Illumina, USA) to obtain library fragments of 250-300 bp. Modification details and amplification conditions are available in Glémin *et al*. (2019b). After verifying and quantifying each indexed cDNA library using a DNA 100 Chip on a Bioanalyzer 2100, pooled libraries were made of twelve, equally represented, genotypes. Each final pooled library was quantified by qPCR with the KAPA Library Quantification Kit (KAPA Biosystems, USA) and sequenced using the Illumina paired-end protocol on a HiSeq3000 sequencer by the Get-PlaGe core facility (GenoToul platform, INRA Toulouse, France http://www.genotoul.fr).

Transcriptome assembly, mapping and genotype calling

Reads cleaning and assembly were performed with the pipeline described in Sarah

et al. (2017). Adapters were removed with cutadapt (Martin, 2011). Reads were trimmed at

the end, removing sequences with low quality scores (parameter -q 20), and we retained

only reads with a minimum length of 35 bp and a mean quality higher than 30. Orphan reads

were then discarded using a homemade script. Retained reads were assembled with ABySS

(Simpson *et al.*, 2009), using the paired-end option with a kmer value of 60, followed by one

step of Cap3 (Huang and Madan, 1999) run with the default parameters, 40 bases of overlap

and 90% percentage of identity. To predict the CDS embedded in our contigs, we used the

prot4est program (Wasmuth and Blaxter, 2004). We provided three gene datasets: the

output of a *Rapsearch* (Ye *et al.*, 2011) similarity analysis, *Oryza* matrix model for de-novo

based predictions and the codon usage bias observed in *T. monococcum*. We run

Rapsearch to identify protein sequences similar to our contigs in either plant species of

Uniprot swissprot (http://www.uniprot.org) or in the Monocotyledon species of greenphyl

(http://www.greenphyl.org/cgi-bin/index.cgi). For the individual used as mapping reference

within each species (see below), we discarded predicted CDS with less than 250bp.

For each species, mapping was done following Sarah *et al*. (2017), except for the

use of the bwa (Li and Durbin, 2009) option –mem (instead of –aln) more adapted for reads

of 100 bp. Reads were mapped on the sequences of the individual with the highest coverage

or with the highest number of annotated contigs. The list of samples used as reference

sequences and the total number of contigs per reference is given in **Supplementary Table**

S5.

For each individual, diploid genotypes were called with reads2snps v. 2.0.64

(Tsagkogeorga *et al.*, 2012; Gayral *et al.*, 2013) (available at

https://kimura.univ-montp2.fr/PopPhyl/index.php?section=tools). This tool is specifically

designed to analyze transcriptome data for population genomics of non-model species. The

method first calculates the posterior probability of each possible genotype in the maximum-

likelihood framework, after estimating the sequencing error rate. Genotypes supported with

probability higher than a given threshold (here 0.95) are retained, otherwise missing data are

called. We required a minimum coverage of 10X per position and per individual to call a

genotype. SNPs are then filtered for possible hidden paralogs (duplicated genes) using a

likelihood ratio test based on explicit modeling of paralogy ("paraclean" option embedded in

the reads2snps software, (Gayral *et al.*, 2013)). First, genotype and SNPs were called

assuming panmixia (heterozygote deficiency $F = 0$), and F was estimated on the retained

SNPs. As we have species-wide samples, F is equivalent to a F_{IT} and mainly corresponds to

F_{IS} for selfing species and to F_{ST} for outcrossing ones. As the assumed expected

heterozygosity can affect genotype calling and paralog filtering, reads2snps were run a

second time for each species using the F estimated after the first step. For the outgroup

species with sample size n=1 (*T. caput-medusae*) we kept the initial genotype calling and

filtering procedure.

Open-reading frames (ORFs) were predicted using the program ORF_extractor.pl

(available at https://kimura.univ-montp2.fr/PopPhyl/index.php?section=tools). Gene length

and number of SNPs in the final data set per species is provided in **Supplementary Table**

S5.

Orthologous pairs of ORFs, hereafter called genes, from the 5 focal and one

outgroup species (*T. caput-medusae*) were identified using reciprocal best hits on BLASTn

results, a hit being considered as valid when *e* value was below e-50. Outgroup sequences

were added to within-focal species alignments using MACSE v. 1.2 (Ranwez *et al.*, 2011), a

program dedicated to the alignment of coding sequences and the detection of frameshifts.

Genes were only retained if no frameshift was identified by MACSE, and if the predicted

ORF in the focal species was longer than 100 codons.

Chromosome patterns and recombination map

We wanted to analyze polymorphism patterns across chromosomes and as a

function of recombination rates. Unfortunately, there is neither reference genome nor genetic

map available for every species. Among the high-quality recombination maps available (see

Brazier and Glémin 2022), we first used the recombination map of *Hordeum vulgare* as a

reference for all species. The synteny is well conserved at the scale of Triticeae (Mayer *et*

al., 2011; but see Parisod and Badaeva, 2020) and, as *H. vulgare* is an outgroup, there

should not be a specific bias for one species or another. For comparison, we also used the

three constitutive genomes AA, BB and DD of the hexaploid wheat, *Trticum aestivum*, which

correspond to the three main lineages in the *Aegilops*/*Triticum* phylogeny (Glémin et al. 2019

and **Fig. 1**). These genomes are closer to the focal species but the phylogenetic distance

depends on the genome and the species.

To build the recombination maps, we built genetic versus physical distance maps

(Marey's maps). For the barley genome, we used the genetic SNP markers from Comadran

et al. (2012) which was initially mapped on version 082214v1 of the barley genome. We thus

used the coordinate correspondence between this first version and the new reference

genome assembly (Hv_IBSC_PGSB_v21) to locate the SNP markers on this reference

genome. After visual inspection of aberrant markers, we kept a total of 3590 markers (on

average ~513 markers per chromosome). Recombination rates were computed with the

MareyMap R package (Rezvoy *et al.*, 2007) by fitting a loess function with a second-degree

polynomial on sliding windows containing 20% of the markers of a chromosome. This led to

a rather smooth recombination map, which is sufficient for our purpose of capturing large-

scale patterns and reducing noise. For the bread wheat genome, we used the Marey maps

built in Brazier and Glémin (2022). Genetic distances were then interpolated between

markers using the fitted function so that a genetic distance could be attributed to each

annotated gene of the *Hordeum* genome. For each gene we computed the local

recombination rate by taking the local derivative of the fitted loess function as implemented

in MareyMap (recombination maps are provided in **Supplementary Fig. S13**).

For each assembled transcript of each focal species, we searched for its orthologous

sequence in the *H. vulgare* genome (H) using reciprocal best blast and retaining pairs when

e value was below e-50. Then the two sequences were aligned with MACSE v.1.2 and the

synonymous divergence, D_S, was computed using *codeml* from the PAML software (Yang,

2007). The D_S distribution was clearly bimodal for all species, and gene pairs showing too

high divergence ($D_S > 0.35$) were discarded as they likely corresponded to paralogues. For

each transcript in each focal species, we attributed the same genetic distance and local

recombination rate as its ortholog in *H. vulgare*. We applied the same procedure for the

three *T. aestivum* subgenomes (A, B and D) separately, except that we did not filtered on D_S

that cannot be used as an homogeneous criterium for all species as the distance depends

on the subgenome and the focal species, contrary to *Hordeum*, which as the same expected

distance with all focal species.

For each focal species, to assess the similarity among genomes, we counted how

many genes had an orthologue on the same chromosome of the four reference genomes (H,

A, B and D) and whether it corresponded to the same category of recombination (low: below

the median, or high: above the median).

Sequence polymorphism analysis

Polymorphism and divergence statistics were calculated with dNdSpNpS v.3.

(available at https://kimura.univ-montp2.fr/PopPhyl/index.php?section=tools) that rely on the

Bio++ libraries (Guéguen *et al.*, 2013). Further filters were applied to the data sets. Positions

at which a genotype could be called in less than five individuals for species with sample size

n ≥ 5 and in less than n/2 for species with n < 5 were discarded. Genes with less than 10

codons were discarded. For each gene, the following statistics were calculated: per-site

synonymous (π_S) and nonsynonymous (π_N) mean pairwise nucleotide diversity, heterozygote

deficiency (F), number of synonymous (S_S) and nonsynonymous (S_N) segregating sites,

number of synonymous (D_S) and nonsynonymous (D_N) fixed differences between focal and

outgroup species. These statistics were computed from complete, biallelic sites only, i.e.,

sites showing no missing data after alignment cleaning, and no more than two distinct states.

For each species, statistics were averaged across genes weighting by the number of

complete sites per gene, thus giving equal weight to every SNP. For π_N/π_S and D_N/D_S, we

first computed the averages of π_N, π_S, D_N, and D_S and subsequently the ratios of averages.

Confidence intervals were obtained by 10,000 bootstraps over genes. For the focal selfing

species *Ae. tauschii*, *T. monococcum* and *T. urartu*, all statistics were calculated on n/2

alleles, by randomly drawing one haploid sequence per gene and individual

(**Supplementary Table S5**).

Fit of a linked-selection model

To go further, we fitted a linked-selection model, following Corbett-Detig *et al*. (2015)

and Elyashiv *et al*. (2016) but including the effect of partial selfing. To simplify the model,

and because we did not have information about substitutions across the genome, we only

considered background selection.

The genome was split into genomic windows. For each region where $\pi_S(i)$ has been

estimated on n_i positions, we assumed that n_i x $\pi_S(i)$ followed a binomial distribution with

parameter n_i and p_i given by:

$$p_i = \pi_{max}\, e^{-\sum_{j}^{\square} u_j \frac{s}{(r_{ij}+s)^2}} \qquad (2)$$

where s is the mean selection coefficient against deleterious mutations, r_{ij} represents the

probability of recombination between the focal region i and any other region of the genome, j

containing L_j coding positions so that $u_j = u\, L_j$, and where u is the rate of deleterious

mutations. To improve the fit to the data we considered a distribution of fitness effects of

mutations. We used a simple discrete distribution with three categories characterized by

their mutation rates and selection coefficients: u_1, u_2, u_3, and s_1, s_2, s_3. Equation (2) can thus

generalized as:

357
$$p_i = \pi_{max}\, e^{-\sum_j^{\square} L_j \left(u1 \frac{s1}{(r_{ij}+s1)^2} + u2 \frac{s2}{(r_{ij}+s2)^2} + u3 \frac{s3}{(r_{ij}+s3)^2} \right)} \quad (3)$$

Note that the s values correspond in fact to $(h + F - hF)s$ where h is the dominance

coefficient. However, because our aim is not to estimate and compare deleterious mutation

parameters we did not fit h and s separately. r_{ij} is obtained from the genetic distance, d_{ij},

using Haldane's mapping function: $r_{ij} = (1 - \exp(-2d_{ij}))/2$. To take partial selfing into account,

we rescaled r_{ij} by $r_{ij}(1 - F)$ (see Nordborg, 2000). This rescaling is correct only for low

recombination rates and a more accurate expression of background selection was obtained

by Roze (2016). However, the simple rescaling provides a good approximation and is much

simpler to handle than the full expression (Roze, 2016). Note that we took the sum on all

genomic regions on the same chromosome of the focal region but also on other

chromosomes (so with $r_{ij} = \frac{1}{2}$). This is especially important under partial selfing as selection

on one chromosome can also affect the other chromosomes. Under outcrossing it boils

down to the effect of the additive variance in fitness that reduces effective population size

(Roze, 2016).

We had seven parameters to estimate: u_1, u_2, u_3, s_1, s_2, s_3 and π_{max}. Because estimates of F

were not very precise, we run the model by letting F free and being estimated jointly with the

other parameters (so eight parameters in total). The log likelihood function was optimized

with the *optim* function in *R* using the "L-BFGS-B" method and the constraint: $u_1 > u_2 > u_3$.

Recombination is very heterogeneous along chromosomes (U-shaped) and large

central regions are strongly linked. Instead of splitting the genome into regions of equal

physical size (in Mb) we split it into regions of equal genetic size (in cM) based on the Marey

map interpolation. Genomic regions on the telomeric parts of the chromosomes were thus

shorter than centromeric regions, where recombination is very low. We chose a window size

of 1 cM and we discarded regions with less than 300 bp to avoid too noisy data. To obtain

confidence intervals on parameter estimations we bootstrapped data 100 times and rerun

the model.

385   Linear models and phylogenetic correction

We looked at the relation of polymorphism (π_S, π_N/π_S and π_{max}) and *F* with mating

system (PC1 of reproductive morphology) and species geographical range using simple

unweighted linear regressions of the form $y \sim x$ (*F* ~ PC1, $\log(\pi_S)$ ~ PC1, π_N/π_S ~ PC1,

$\log(\pi_S)$ ~ range, π_N/π_S ~ range, π_N/π_S ~ *F*). To evaluate the joint effects of mating system and

species range on polymorphism (π_S and π_N/π_S), we performed multiple linear regressions in

the form $\log(\pi_S)$ ~ mating_system + species range and π_N/π_S ~ mating_system + species

range. We then represented the residuals, after removing the effect of the mating system, as

a function of species range. All linear models were run with the function *lm* under the R

environment.

We also applied a correction to take into account the phylogenetic relationships

among species using the ultrametric tree retrieved from Glémin *et al*. (2019). For this, we

computed the phylogenetically independent contrasts using the method of Felsenstein

(1985) with the package *ape* version 5.6-2 (Paradis and Schliep, 2019). The function *pic* was

applied to the *y* and *x* vectors with default parameters, then the *lm* analysis was repeated

with the contrast values obtained instead of the raw values.

Estimation of the distribution of fitness effect of mutations

In the two self-incompatible species, *Ae. mutica* and *Ae. speltoides*, and in two highly

selfing species, *Ae. tauschii* and *T. urartu*, we estimated the distribution of fitness effect of

mutations using the *PolyDFE* method (Tataru *et al.*, 2017; Tataru and Bataillon, 2019). In

brief, this method used the unfolded site frequency spectrum (uSFS) for both synonymous

and non-synonymous mutations to fit the distribution of fitness effect (DFE) of mutations

modelled by the mix of a gamma distribution for deleterious mutations and an exponential

distribution for beneficial mutations. Demography is taken into account by adding and fitting

noise parameters that distort uSFS from the equilibrium expectation following Eyre-Walker

et al. (2006). Because uSFS are sensitive to polarization errors, which can give spurious

signatures of beneficial mutations, a probability of mis-polarization is also added and fitted in

the model. This yields a set of four related models: with and without beneficial mutations,

and with and without polarization errors. Instead of choosing the best model to estimate

parameters, we run all four and used a model averaging procedure (as in Muyle *et al.*,

2021): each parameter estimate was averaged using Akaike weights $e^{-\frac{1}{\Delta AIC_i}}$ with

$\Delta AIC_i = AIC_i - AIC_{min}$ where AIC_{min} is the AIC of the best model. Confidence intervals were

obtained by bootstrapping SNPs 1000 times.

For the four species, uSFS were polarized using *Taeniatherum caput-medusae* as an

outgroup. We performed three analyses: on the whole set of SNPs or by splitting the dataset

into two subsets: SNPs from genes in the high or low recombining regions. GC-biased gene

conversion (gBGC), a recombination associated processes mimicking selection in favor of G

and C nucleotides, is known to be active in grasses (Rodgers-Melnick *et al.*, 2016; Clément

et al., 2017) and could lead to spurious signatures of positive selection in highly recombining

regions. To test for the possible effect of gBGC, we also re-run the analyses on three

categories of SNPs: AT→GC, GC→AT, and G←→C + A←→T.

Simulations

To understand how different selfing rates affect our ability to detect the effects of

linked selection on polymorphism landscape and distribution of fitness effects, we run

forward time individual-based evolutionary simulations in SLiM v.3.3 (Haller and Messer

2019). We simulated a population of N = 10,000 individuals with five different selfing rates: 0,

0.5, 0.9, 0.99 and 1. We considered a genome of three Mb with a single chromosome

composed of 1000 genes of 1000 bp separated by intergenic regions of 2000 bp.

Recombination decreased exponentially from 60 cM/Mb at the tips to 6 cM/Mb in the center

of the chromosome, corresponding to a total genetic of 3.24 Morgan (so an average of three

crossovers per chromosome per meiosis, which is in the range of one to three/four that is

observed in plants, Brazier and Glémin 2022). We assumed a mutation rate of 10^{-6}, with ⅔ of

mutations being neutral, corresponding to an expected genetic diversity of $4Nu$ = 0.027, of

the order of magnitude of what we observed in the outcrossing species, and corresponding

to an average r/u close to one. The other third of mutations were considered deleterious with

a dominance level of h = 0.25 and deleterious effects in homozygotes drawn in a gamma

distribution with mean = 0.01 and shape = 0.5. This corresponds to a genomic deleterious

mutation rate of U = 0.33. After a burn-in period of $10N$ generations we recorded the

genome sequence of 15 individuals. We run ten replicates for each selfing rate.

We used the simulated data to assess the effect of selfing and linked selection on the

estimation of the DFE. Importantly, polyDFE estimates the shape and the population-scaled

mean of the DFE: $S = 4N_e(h + F - hF)s$. However, N_e is not set nor fixed in the model,

contrary to h, s and F, but depends on the intensity of linked selection. We thus used the

observed π_S divided by $4u$ to get the predicted N_e, hence the predicted S.

Results

Mating system widely varies in *Aegilops*/*Triticum* genus

We analyzed phenotypic and transcriptomic diversity in 98 accessions from the 13

diploid *Aegilops*/*Triticum* species and three close outgroup species *Taenatherium caput-*

medusae, *Hordeum vulgare* and *Secale vavilovii* (**Fig. 1, Table S1** and **Fig. S1**). Individuals

were sampled over the whole geographic range to assess genetic diversity at the species

scale. For some species the mating system was already well known, including the self-

incompatible *Ae. speltoides* and *Ae. mutica* and the highly selfing *T. urartu*, *T. monococcum*

and *Ae. tauschii* (Dvořák *et al.*, 1998; Escobar *et al.*, 2010), but for others it was poorly

documented (Kilian *et al.*, 2011). We thus characterized the mating system of each species

by quantifying six floral and reproductive traits, including the size of female and male

reproductive organs (anthers, stigmas), male investment, spikelet compactness, and the

autonomous seed set (Escobar *et al.*, 2010) (**Supplementary Tables S2, S3 and S4**).

Building on previous work (Escobar *et al.*, 2010), we considered these traits as indicative of

the selfing syndrome, i.e. the specific changes in flower morphology and function that are

expected to occur following the evolution of self-fertilization, especially for anemophilous

species (Escobar *et al.*, 2010; Sicard and Lenhard, 2011). The autonomous seed set

provided a verification that self-incompatible species *Ae. mutica* and *Ae. speltoides*

produced almost no seeds under imposed self-fertilization in the greenhouse (bagged

spikes), while all the other species were able to produce seeds (**Supplementary Fig. S2**).

All the other traits were significantly negatively (anther and stigma size, male investment) or

positively (spikelet compactness) correlated with the autonomous seeds set

(**Supplementary Fig. S3**), indicating that the selected traits are good indicators of the

mating system of each species.

We summarized this reproductive morpho-functional diversity with a multivariate

approach, principal component analysis (PCA). The PCA first axis reflected the differences

in mating systems within the *Aegilops/Triticum* genus (**Fig. 1B** and **Supplementary Fig. S4**).

On one extreme of PC1 there were self-incompatible *Ae. mutica* and *Ae. speltoides*, which

showed bigger anthers and stigmas, higher male/female investment, lower spikelet

compactness and lower (null) autonomous seed set. On the other extreme there were the

predominantly selfing *Ae. tauschii*, *T. urartu* and *T. monococcum*, with opposite states of the

traits (**Fig. 1A and 1B**). The other *Aegilops* species showed intermediate values of the multi-

trait statistic. This result still held when outgroups were included in the analysis

(**Supplementary Fig. S5**).

Species morphology measured by PC1 explained well the genome-wide estimate of

inbreeding coefficient, *F,* calculated on the whole transcriptome dataset (R^2=0.83, p-

value=1.68e-05, **Fig. 1C**), confirming that our phenotypic data represented a good proxy of

the mating system. Thus, in the following analyses, we used PC1 to describe the selfing

syndrome, which avoids using genomic data for both characterizing the mating systems and

their genomic consequences.

These findings are in agreement with previous knowledge (Dvořák *et al.*, 1998;

Escobar *et al.*, 2010) and allowed us to characterize the mating system for the species that

lacked outcrossing rate estimations. They also showed that the effects of selfing might be

gradual, with species exhibiting mixed mating strategies described by intermediate values of

F and of the phenotypic selfing syndrome. Mapping mating systems on the phylogeny

suggested that self-incompatibility was likely ancestral and may have broken only once as all

species are self-compatible, except the two external ones (*Ae. mutica* and *Ae. speltoides*).

However, several breakdowns of self-incompatibility cannot be excluded. For example, high

selfing could have evolved four times independently, in the branches leading to *Ae. tauschii*,

Ae. searsii, *Ae. uniaristata* and *Triticum* species (**Fig. 1A**).

Polymorphism strongly correlates with mating systems

Genetic diversity was estimated for each species from whole transcriptome

sequencing data. Sequences of 48 accessions were generated and *de novo* assembled in

this study and were added to the datasets of Glémin *et al*. (2019b) and Clément *et al*. (2017)

(**Supplementary Table S1**). Between 19,518 and 28,834 coding sequences were obtained

per species. After filtering, genotype calling was performed on a number of contigs varying

from 7,083 (*Ae. searsii*) to 21,706 (*T. caput-medusae*) (**Supplementary Table S5**).

Selfing is expected to reduce neutral genetic diversity (Pollak, 1987; Schoen and

Brown, 1991; Jarne, 1995; Nordborg, 2000; Ingvarsson, 2002), here estimated as

synonymous polymorphism, π_S. Across species, π_S varied more than one order of

magnitude, from 0.0011 for the self-compatible *Ae. searsii* to 0.02 for the self-incompatible

Ae. speltoides. According to expectations, genetic diversity decreases with increasing selfing

rate. Neutral genetic diversity, π_S and PC1 were significantly correlated across species (R^2 =

0.75, p-value = 0.00014), indicating a gradient in which stronger selfing syndrome

corresponds to lower genome-wide neutral diversity (**Fig. 2A**). The correlation was still

significant after phylogenetic control (R^2 = 0.67, p-value = 0.00111). Interestingly, this

relationship is more or less log-linear (**Fig. 2A**) with the main difference being observed

between the two self-incompatible (*Ae. speltoides* and *Ae. mutica*) and the self-compatible

species (all the others).

Selfing is also expected to reduce the efficacy of selection, thus leading to higher

accumulation of segregating slightly deleterious mutations in selfing than in outcrossing

species (Glémin, 2007). In agreement with this prediction, the efficacy of purifying selection

estimated by the ratio of non-synonymous to synonymous polymorphism π_N/π_S (Kimura,

1983) was lower for selfing species (max π_N/π_S value 0.21 for *T. urartu*) than for outcrossing

ones (0.09 for *Ae. speltoides*). Similarly to neutral diversity, the efficacy of selection was also

significantly explained by the selfing syndrome (R^2 = 0.48, p-value = 0.0083; with

phylogenetic control R^2 = 0.35, p-value = 0.041; **Fig. 2B**). We further verified that both

polymorphism statistics, π_S and π_N/π_S, also correlated with *F* estimates (**Supplementary Fig.**

S6).

We also tested whether species range, used as a proxy of census population size,

also correlated with genetic diversity, with widespread species predicted to be more

polymorphic than species with restricted geographic distribution. In contrast to the mating

system, species range was not correlated with either π_S (**Fig. 2C)** or π_N/π_S (**Fig. 2D)**. Such a

correlation could be masked by the strong effect of selfing, which is expected to favor

species range expansions. For example, *Ae. tauschii* is highly selfing and has by far the

largest species range (**Supplementary Fig. S1**). However, a linear model with the two

effects showed that the mating system still significantly explained both π_S (p-value=0.00004)

and π_N/π_S (p-value=0.0157) whereas species range did not (**Supplementary Fig. S7**). Yet,

the effect of species range on π_S is barely significant (p-value = 0.063), so it is still possible

that there is a weak effect that we could not detect with only thirteen species. Overall, these

results suggest that the mating system is the main driver of genetic diversity in

Aegilops/*Triticum* species and overwhelms potential effects of recent population history.

The effect of linked selection depends on the mating system

We tested the hypothesis that selfing increases the effect of linked selection by

comparing polymorphism patterns and recombination along chromosomes. Species-specific

recombination maps were not available for most species, so we used the recombination map

of the outgroup species *Hordeum vulgare*, which we compared to the recombination maps of

the three diploid subgenomes of bread wheat (A, B and D, corresponding to the wild parents

T. urartu, *Ae. speltoides/mutica*, *Ae. tauschii*). For each focal species, we found that 99% or

orthologs with *H. vulgare* mapped on the same chromosome as at least one of the three

subgenomes of *T. aestivum*, and 96% to 97% as all three of them. Similarly, 97 to 99% of

orthologs with *H. vulgare* belonged to the same recombination category as at least one of

the three subgenomes, and 78 to 79% as all three of them (**Supplementary Table S6**).

In what follows, we only show the results with *H. vulgare*, since it provides the further

advantage that the outgroup has (on average) the same phylogenetic distance to every

Aegilops/*Triticum* species, ensuring an unbiased analysis that does not favor species closer

to the reference. For comparison, some additional results using the *T. aestivum*

subgenomes as reference are given in supplementary material (**Supplementary Table S6**

and **Fig. S9**).

In all species, synonymous polymorphism was strongly correlated with recombination

and presented a U-shaped pattern along chromosomes more or less mirroring the

recombination pattern (see **Fig. 3A** and **Supplementary Fig. S14**). However, the higher the

selfing rate, the flatter the relationship (**Fig. 3B**), suggesting a strong effect of the mating

system on the relationship between polymorphism and recombination. We verified that the

positive relationship between diversity and recombination rate was not merely caused by the

mutagenic effect of recombination, by looking at the correlation between synonymous

divergence (D_S) with the outgroup (*H. vulgare*) and recombination rate (Kulathinal *et al.*,

2008). We found that the magnitude of D_S variation (factor 1.5, **Supplementary Fig. S8**) is

much lower than the range of variation observed for polymorphism along the genome (factor

5 to 80, depending on the species, **Fig. 3B**). If a mutagenic effect of recombination cannot

be ruled out, it is clearly insufficient to explain the magnitude of the correlation between π_S

and recombination. This mere observation suggested that linked selection could strongly

reduce π_S by at least one or two orders of magnitude.

To quantify more directly the effect of linked selection, we fitted a model similar to

Corbett-Detig *et al.* (2015) and Elyashiv *et al.* (2016) that we adapted to partially selfing

species but only considering background selection (see **Supplementary Table S6** for full

results). From the fit of the model, we obtained the maximum π_S that could be reached in the

absence of linked selection, π_{max}, which ranged between 0.028 to 5.82 (**Fig. 4**). Note that,

here $\pi_{max} = 4N_{e\_max}u$, so can be higher than 1. We estimated that linked selection reduced π_S

by 3.5 in *Ae. speltoides* and 5.6 in *Ae. mutica*, the two self-incompatible species. For other

species, π_S was reduced by a few tens or even a few hundreds (from 7 to 888), but without a

clear relationship with the mating system. In contrast to π_S, π_{max} did not correlate with PC1.

Surprisingly, it correlated negatively with species range, but the correlation was mainly

driven by the species of the *Sitopsis* section and was no longer significant after phylogenetic

correction (**Fig. 4**). It can be difficult to properly fit a realistic linked selection model for selfing

species and the results can be sensitive to the fact that we did not use the reference

genome of each species. In particular, some π_{max} values were very high and could be

overestimated, but fitting the model using the three subgenomes A, B and D of *T. aestivum*

gave similar results (**Supplementary Fig. S9**). Overall, although they must be viewed with

caution, the results strongly suggested that linked selection is a main driver of the effect of

selfing on genetic diversity whereas species range has only a minor effect, and if any, not in

the predicted direction. When recombination maps will be available in all species, it will be

possible to re-assess this result.

Deleterious mutations accumulate and adaptation is reduced under selfing and low recombination

Another central prediction of the effect of genetic linkage is that selection should be less efficient in genomic regions of low recombination, which can extend genome wide in highly selfing species. In agreement with this expectation, the efficacy of purifying selection at the genome wide level clearly decreased with the selfing rate (**Fig. 2B**). All species also showed a negative relationship between recombination rates and the π_N/π_S ratio (**Fig. 3B**), indicating that purifying selection was more efficient in highly recombining regions. More precisely, the π_N/π_S ratio sharply dropped with increasing recombination in outcrossing species but more and more smoothly with increasing selfing rate, which supports the prediction that reduced selection efficacy extended to larger genomic regions in selfing species.

The π_N/π_S ratio is a rather crude proxy for the efficacy of purifying selection, and can be affected by several factors such as non-equilibrium population dynamics that can lead to spurious signature of relaxed selection (Brandvain and Wright 2016). To better characterize how selection efficacy varies with mating system and recombination, we estimated the full distribution of fitness effects (DFE) of mutations - i.e. including both deleterious and beneficial mutations - using the *polyDFE* method (Tataru *et al.*, 2017). This approach leverages information from unfolded synonymous and non-synonymous site frequency spectra to infer the DFE of each species. It takes into account factors that can distort the SFS such as non-equilibrium demography and linked selection, in addition to potential polarization errors. The method requires a sufficient number of chromosomes sampled (say >10), so we applied it only to the four species with the largest sample sizes, which correspond to the extreme mating systems used to calibrate the selfing syndrome: the two

self-incompatible *Ae. speltoïdes* and *Ae. mutica* and two highly selfing *Ae. tauschii* and *T. urartu*. In agreement with the π_N/π_S ratio analysis, we found that the two selfers suffered from a higher load than the two outcrossers, with 22% to 25% of mutations not being efficiently selected against (-10 < $Ne\ s$ < 0), versus only 9 to 15% in the outcrossers (**Fig. 5**). We also found a strong difference between regions of low and high recombination (higher vs lower than the median) for all species. However, the difference was stronger in outcrossers (more than two-fold) than in selfers (30-60% difference only) (**Fig. 5**). Interestingly, purifying selection appeared as efficient in low-recombination regions of outcrossing genomes than in high-recombination regions of highly selfing genomes (**Fig. 5**).

Another striking result is that we estimated an adaptive substitution rate not different from zero in the two selfers, but a quite high value in the two outcrossers and only in highly recombining regions (**Fig. 5**). We verified that this signature of positive selection was not due to the spurious effect of GC-biased gene conversion (**Supplementary Fig. S10**), which is known to happen in recombining regions of grass genomes (Muyle *et al.* 2011). Overall, selection appeared to be much less efficient, both on beneficial and against deleterious mutations, in low-recombination regions of outcrossing species, and throughout the genome of highly selfing ones. When the proportion of weakly deleterious mutations is estimated with the DFE-alpha method (Keightley and Eyre-Walker 2007) it was shown that selfing could overestimate it (Gilbert et al. 2022). Here, we used polyDFE, which was claimed to be less sensitive to linked selection effects (Tataru et al. 2017). As a control, we run simulations with linked selection and varying degree of selfing and applied polyDFE. In contrast with DFE-alpha (see Gilbert et al. 2022), we found that polyDFE tended to underestimate the proportion of weakly deleterious mutations (**Supplementary Fig. S11).** We also found that the method did not estimate spurious signatures of beneficial mutations in outcrossers. Overall, the results were conservative to the effects of linked selection and selfing.

Discussion

We compared the patterns of genetic diversity and selection across the genome in

thirteen species with contrasted mating systems. We found far less polymorphism and far

less selection efficacy in selfing than in outcrossing species, as observed in previous studies

(e.g., Glémin *et al.*, 2006; Hazzouri *et al.*, 2013; Slotte *et al.*, 2013; Barrett *et al.*, 2014;

Burgarella *et al.*, 2015; Chen *et al.*, 2017; Laenen *et al.*, 2018). We also showed that these

genomic effects depend on the interplay between linked selection and mating systems, and

vary with self-fertilization rates. For this, we leveraged a study design tailored to go beyond

global patterns and decipher their underlying causes. First, differently from previous general

comparisons among plant species (Glémin *et al.*, 2006; Chen *et al.*, 2017), we compared

related species with similar life history traits, ecology and genomic features, which allows

more direct testing of the effect of the mating system. Second, we investigated all species of

a clade covering a large range of mating systems, including intermediate mixed mating

species, whereas previous studies addressing sister (or closely related) species mainly

focused on extreme outcrossing vs selfing comparisons (e.g., Hazzouri *et al.*, 2013; Slotte *et*

al., 2013; Burgarella *et al.*, 2015; Teterina *et al.*, 2023). Third, by using recombination maps

we quantified the effect of linked selection in a comparative way.

Linked selection appears to be a main mechanism shaping levels of diversity in wild

wheats, as it can reduce polymorphism to three to five-fold in outcrossing species (**Fig. 3B**)

and to one or two orders of magnitude in selfing species. This is in agreement with but

higher than observed by Corbett-Detig *et al*. (2015), who found a quantitatively limited effect

of linked selection except in selfing species (see the re-analysis of Coop 2016). These

quantitative values must be viewed with caution because it is difficult to properly fit a model

of linked selection in selfing species. In addition to the complexity of the interaction between

recombination, selfing and selection, it is not clear whether the species scale or a more local

population scale is the most relevant. Another expected consequence of linkage and selfing

is a reduction in selection efficacy, both against deleterious and in favor of beneficial

mutations. We observed a striking contrast between regions of high and low recombination

in self-incompatible species, with twice more weakly selected deleterious mutations and no

beneficial mutation expected to fix in regions of low recombination (**Fig. 5, Supplementary**

Fig. S10). In highly selfing species, instead, recombination had a weaker effect and only for

deleterious mutations (**Fig. 5)**. Interestingly, the regions of high recombination in selfers

exhibited a similar amount of weakly deleterious mutations as regions of low recombination

in outcrossers, and we detected no signature of adaptive evolution at all in the two highly

selfing species. Simulations showed that these results were not artefactual and may even

underestimate the effect of selfing (**Supplementary Fig. S11**). Overall, the cross

comparison between mating systems and recombination levels clearly showed that the main

quantitative effect of selfing is due to high linkage and linked selection.

Although our studied species share life history traits and have similar ecology, other

factors than the mating system could affect genetic diversity, for example, factors generating

contrasted geographic ranges unrelated to mating systems. However, we showed that the

geographic range has no effect on polymorphism patterns (**Fig. 2, Supplementary Fig. S7**).

More generally, we cannot exclude that other factors could play a role, but they could hardly

generate the very strong relationship between the mating system and genetic diversity we

observed. Strikingly, this strong relationship holds despite the use of an indirect measure of

the selfing rates through phenotypic proxies.

Our results help better understand the evolution of selfing species. In the short term,

selfing is known to recurrently evolve from outcrossing, depending on reproductive

assurance and gene transmission advantage balanced by inbreeding depression, which can

be partly purged during such transitions. In the long run, selfing lineages tend to diversify

less than outcrossing ones and selfing is considered an evolutionary dead-end, likely

because of higher probability of extinction (Stebbins, 1957; Igic *et al.*, 2008; Goldberg and

Igić, 2012). The very causes of higher extinction rates in selfers remain unclear, but

increased load and loss of genetic diversity and adaptive potential are possible drivers.

However, the pace at which the effects of selfing manifest is still poorly known, although it is

likely rapid as in the selfing *Capsella rubella* recently derived from the self-incompatible *C.*

grandiflora (Slotte *et al.*, 2013) or within the species *Arabis alpina* among populations with

contrasted mating systems (Laenen *et al.*, 2018).

In the *Aegilops*/*Triticum* genus, several instances of evolution towards different

degrees of selfing likely occurred in a short evolutionary time period, which manifested by

wide variations in floral traits associated with specific genetic diversity patterns. At the short

phylogenetically scale we studied, we thus found a clear signature of the joint evolution of

morphological traits and population genomic patterns, suggesting that the negative effects of

selfing manifests rapidly. It is tempting to propose that the strong and rapid deleterious

effects of selfing we detected will accelerate the extinction of the most selfing lineages.

However, so far there is no approach to properly test this hypothesis (Wright *et al.*, 2013;

Glémin *et al.*, 2019a). Genomic degradation certainly accompanies the transition towards

selfing, but we still do not know whether it is the ultimate cause of selfing lineages extinction.

723  Finally, our results also bear more general implications about the central question of
724 the determinants of genetic diversity, beyond the case of selfing species. In line with
725 previous comparative analyses of polymorphism at the genome scale (Romiguier *et al.*,
726 2014; Chen *et al.*, 2017), we showed that life history traits, here the mating system, are a
727 much stronger determinant of genetic diversity than proxies for census population sizes,
728 here species range. Species range had no effect either after globally controlling for the
729 mating system (**Supplementary Fig. S7),** or after removing specifically the effect of linked
730 selection (no significant correlation between species range and π_{max} or, if so, in the
731 unpredicted direction). In contrast to previous studies, however, the range of genetic
732 diversity is particularly large despite species being closely related and recently diverged
733 (about 6 MYA), with similar genomes and life-history traits (wind-pollinated annual herbs)
734 except mating system. In these wild wheats, species nucleotide polymorphism varies with a
735 factor 20 (from 0.0011 to 0.022). For comparison, a clade of selfing and outcrossing
736 *Caenorhabditis* nematodes species diverged less than 30 MYA show even wider disparities
737 in polymorphism, up to a factor 80 (Cutter 2008; Li *et al*. 2014). In contrast, in a butterfly
738 family that diverged around 120 MYA, with four-fold variation in body mass and two-fold
739 variation in chromosome numbers, only a factor ten was observed (from 0.0044 to 0.043)
740 (Mackintosh *et al.*, 2019). Similarly, among 28 species widely covering the seed-plant
741 phylogeny and life forms (from annual herbs to trees), the observed range was only slightly
742 higher than in wild wheats, with a factor 28 (from 0.00064 to 0.018) (Chen *et al.*, 2017). This
743 points to mating systems as a main determinant of variation in genetic diversity among
744 species.

745  However, variation in genetic diversity is still narrower than predicted from variation in
746 species range, which varies by a factor 500 here. This is in line with the "Lewontin's
747 paradox", the general observation that genetic diversity varies much less across species

than census size does (Lewontin, 1974; Charlesworth and Jensen, 2022). Variation in π_{max} is

around 60, so higher than variation in π_S, around 20, but only by a factor three. Despite its

strong effect, linked selection is thus unlikely to explain alone the limited range of variation in

π_S, in agreement with previous results (Corbett-Detig *et al.*, 2015; Coop, 2016; Buffalo, 2021;

Charlesworth and Jensen, 2022).

References

Agrawal, A.F. & Hartfield, M. (2016) Coalescence with Background and Balancing Selection

in Systems with Bi- and Uniparental Reproduction: Contrasting Partial Asexuality and

Selfing. *Genetics*, **202**, 313–326.

Barrett, S.C.H., Arunkumar, R. & Wright, S.I. (2014) The demography and population

genomics of evolutionary transitions to self-fertilization in plants. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, **369**, 20130344.

Brandvain, Y. & Wright, S.I. (2016) The Limits of Natural Selection in a Nonequilibrium

World. *Trends in Genetics*, **32**, 201–210.

Brazier, T. & Glémin, S. (2022) Diversity and determinants of recombination landscapes in

flowering plants. *PLOS Genetics*, **18**, e1010141.

Buffalo, V. (2021) Quantifying the relationship between genetic diversity and population size

suggests natural selection cannot explain Lewontin's Paradox. *eLife*, **10**, e67509.

Burgarella, C., Gayral, P., Ballenghien, M., Bernard, A., David, P., Jarne, P., *et al.* (2015)

Molecular Evolution of Freshwater Snails with Contrasting Mating Systems. *Molecular*

Biology and Evolution, **32**, 2403–2416.

Burgarella, C. & Glémin, S. (2017) Population Genetics and Genome Evolution of Selfing

Species. In *eLS* (ed. by John Wiley and Sons Ltd). John Wiley and Sons, Ltd, Chichester,

UK, pp. 1–8.

Charlesworth, B. (2012) The Effects of Deleterious Mutations on Evolution at Linked Sites.

Genetics, **190**, 5–22.

Charlesworth, B. & Jensen, J.D. (2021) Effects of Selection at Linked Sites on Patterns of

Genetic Variability. *Annual Review of Ecology, Evolution, and Systematics*, **52**, 177–197.

Charlesworth, B. & Jensen, J.D. (2022) How Can We Resolve Lewontin's Paradox?

Genome Biology and Evolution, **14**, evac096.

Charlesworth, D., Morgan, M.T. and Charlesworth, B. (1993) Mutation accumulation in finite

outbreeding and inbreeding populations. *Genetics Research*, **61**, 39–56.

Chen, J., Glémin, S. & Lascoux, M. (2017) Genetic Diversity and the Efficacy of Purifying

Selection across Plant and Animal Species. *Molecular Biology and Evolution*, **34**, 1417–

1428.

Chen, J., Glémin, S. & Lascoux, M. (2020) From Drift to Draft: How Much Do Beneficial

Mutations Actually Contribute to Predictions of Ohta's Slightly Deleterious Model of

Molecular Evolution? *Genetics*, **214**, 1005–1018.

Clément, Y., Sarah, G., Holtz, Y., Homa, F., Pointet, S., Contreras, S., *et al.* (2017)

Evolutionary forces affecting synonymous variations in plant genomes. *PLOS Genetics*,

13, e1006799.

Comadran, J., Kilian, B., Russell, J., Ramsay, L., Stein, N., Ganal, M., *et al.* (2012) Natural

variation in a homolog of Antirrhinum CENTRORADIALIS contributed to spring growth

habit and environmental adaptation in cultivated barley. *Nature Genetics*, **44**, 1388–1392.

Coop, G. (2016) Does linked selection explain the narrow range of genetic diversity across

species? *bioRxiv*, 042598.

Corbett-Detig, R.B., Hartl, D.L. & Sackton, T.B. (2015) Natural Selection Constrains Neutral

Diversity across A Wide Range of Species. *PLOS Biology*, **13**, e1002112.

Cutter, A.D., 2008. Divergence Times in Caenorhabditis and Drosophila Inferred from Direct

Estimates of the Neutral Mutation Rate. *Mol. Biol. Evol*. **25**, 778–786.

Cutter, A.D. & Payseur, B.A. (2013) Genomic signatures of selection at linked sites: unifying

the disparity among species. *Nature Reviews. Genetics*, **14**, 262–274.

Dray, S. & Dufour, A.-B. (2007) The ade4 Package: Implementing the Duality Diagram for

Ecologists. *Journal of Statistical Software*, **22**, 1–20.

Dvořák, J., Luo, M.-C. & Yang, Z.-L. (1998) Restriction Fragment Length Polymorphism and

Divergence in the Genomic Regions of High and Low Recombination in Self-Fertilizing

and Cross-Fertilizing Aegilops Species. *Genetics*, **148**, 423–434.

Ellegren, H. & Galtier, N. (2016) Determinants of genetic diversity. *Nature Reviews.*

Genetics, **17**, 422–433.

Elyashiv, E., Sattath, S., Hu, T.T., Strutsovsky, A., McVicker, G., Andolfatto, P., *et al.* (2016)

A Genomic Map of the Effects of Linked Selection in Drosophila. *PLOS Genet*, **12**,

e1006130.

Escobar, J.S., Cenci, A., Bolognini, J., Haudry, A., Laurent, S., David, J., *et al.* (2010) An

integrative test of the dead-end hypothesis of selfing evolution in Triticeae (poaceae).

Evolution, **64**, 2855–2872.

Excoffier, L., Foll, M. & Petit, R.J. (2009) Genetic Consequences of Range Expansions.

Annual Review of Ecology, Evolution, and Systematics, **40**, 481–501.

Eyre-Walker, A., Woolfit, M. & Phelps, T. (2006) The distribution of fitness effects of new

deleterious amino acid mutations in humans. *Genetics*, **173**, 891–900.

Felsenstein, J. (1985) Phylogenies and the comparative method. *American Naturalist*, **125**,

1--15.

Friedman, J. & Harder, L.D. (2005) Functional associations of floret and inflorescence traits

among grass species. *American Journal of Botany*, **92**, 1862–1870.

Gayral, P., Melo-Ferreira, J., Glémin, S., Bierne, N., Carneiro, M., Nabholz, B., *et al.* (2013) Reference-free population genomics from next-generation transcriptome data and the vertebrate–invertebrate gap. *PLoS Genetics*, **9**, e1003457.

Gilbert, K.J., Zdraljevic, S., Cook, D.E., Cutter, A.D., Andersen, E.C., Baer, C.F., 2022. The distribution of mutational effects on fitness in Caenorhabditis elegans inferred from standing genetic variation. *Genetics* 220, iyab166.

Glémin, S. (2007) Mating systems and the efficacy of selection at the molecular level. *Genetics*, **177**, 905–916.

Glémin, S., Bazin, E. & Charlesworth, D. (2006) Impact of mating systems on patterns of sequence polymorphism in flowering plants. *Proceedings of the Royal Society B: Biological Sciences*, **273**, 3011–3019.

Glémin, S., François, C.M. & Galtier, N. (2019a) Genome Evolution in Outcrossing vs. Selfing vs. Asexual Species. In *Evolutionary Genomics: Statistical and Computational Methods*, Methods in Molecular Biology (ed. by Anisimova, M.). Springer, New York, NY, pp. 331–369.

Glémin, S., Scornavacca, C., Dainat, J., Burgarella, C., Viader, V., Ardisson, M., *et al.* (2019b) Pervasive hybridizations in the history of wheat relatives. *Science Advances*, **5**, eaav9188.

Goldberg, E.E. & Igić, B. (2012) Tempo and Mode in Plant Breeding System Evolution. *Evolution*, **66**, 3701–3709.

Grossenbacher, D., Runquist, R.B., Goldberg, E.E. and Brandvain, Y. (2015) Geographic range size is predicted by plant mating system. *Ecology Letters*, **18**, 706–713.

Guéguen, L., Gaillard, S., Boussau, B., Gouy, M., Groussin, M., Rochette, N.C., *et al.* (2013) Bio++: Efficient Extensible Libraries and Tools for Computational Molecular Evolution.

Molecular Biology and Evolution, **30**, 1745–1750.

Haller, B.C. & Messer, P.W. (2019) SLiM 3: Forward Genetic Simulations Beyond the

Wright–Fisher Model. *Molecular Biology and Evolution*, **36**, 632–637.

Hartfield, M. & Bataillon, T. (2020) Selective Sweeps Under Dominance and Inbreeding. *G3*

Genes|Genomes|Genetics, **10**, 1063–1075.

Hazzouri, K.M., Escobar, J.S., Ness, R.W., Killian Newman, L., Randle, A.M., Kalisz, S., *et*

al. (2013) Comparative population genomics in Collinsia sister species reveals Evidence

for reduced effective population size, relaxed selection, and evolution of biased gene

conversion with an ongoing mating system shift. *Evolution*, **67**, 1263–1278.

Huang, S., Sirikhachornkit, A., Su, X., Faris, J., Gill, B., Haselkorn, R., *et al.* (2002) Genes

encoding plastid acetyl-CoA carboxylase and 3-phosphoglycerate kinase of the

Triticum/Aegilops complex and the evolutionary history of polyploid wheat. *Proceedings*

of the National Academy of Sciences, **99**, 8133–8138.

Huang, X. & Madan, A. (1999) CAP3: A DNA Sequence Assembly Program. *Genome*

Research, **9**, 868–877.

Igic, B., Lande, R. and Kohn, J.R. (2008) Loss of Self-Incompatibility and Its Evolutionary

Consequences. *International Journal of Plant Sciences*, **169**, 93–104.

Ingvarsson, Pärk. (2002) A metapopulation perspective on genetic diversity and

differentiation in partially self-fertilizing plants. *Evolution*, **56**, 2368–2373.

Jarne, P. (1995) Mating system, bottlenecks and genetic polymorphism in hermaphroditic

animals. *Genetics Research*, **65**, 193–207.

Josse, J. & Husson, F. (2016) missMDA: A Package for Handling Missing Values in

Multivariate Data Analysis. *Journal of Statistical Software*, **70**, 1–31.

Kilian, B., Mammen, K., Millet, E., Sharma, R., Graner, A., Salamini, F., *et al.* (2011)

Aegilops. In *Wild Crop Relatives: Genomic and Breeding Resources: Cereals* (ed. by

Kole, C.). Springer, Berlin, Heidelberg, pp. 1–76.

Kimura, M. (1983) Rare variant alleles in the light of the neutral theory. *Molecular Biology*

and Evolution, **1**, 84–93.

Kulathinal, R.J., Bennett, S.M., Fitzpatrick, C.L. and Noor, M.A.F. (2008) Fine-scale mapping

of recombination rate in Drosophila refines its correlation to diversity and divergence.

Proceedings of the National Academy of Sciences, **105**, 10051–10056.

Laenen, B., Tedder, A., Nowak, M.D., Toräng, P., Wunder, J., Wötzel, S., *et al.* (2018)

Demography and mating system shape the genome-wide impact of purifying selection in

Arabis alpina. *Proceedings of the National Academy of Sciences*, **115**, 816–821.

Leffler, E.M., Bullaughey, K., Matute, D.R., Meyer, W.K., Ségurel, L., Venkat, A., *et al.*

(2012) Revisiting an Old Riddle: What Determines Genetic Diversity Levels within

Species? *PLoS Biol*, **10**, e1001388.

Lewontin, R.C. (1974) *The genetic basis of evolutionary change*. Columbia University Press.

Li, H. and Durbin, R. (2009) Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics (Oxford, England)*, **25**, 1754–1760.

Li, S., Jovelin, R., Yoshiga, T., Tanaka, R., Cutter, A.D., 2014. Specialist versus generalist

life histories and nucleotide diversity in Caenorhabditis nematodes. *Proc. R. Soc. B Biol.*

Sci. **281**, 20132858.

Mackintosh, A., Laetsch, D.R., Hayward, A., Charlesworth, B., Waterfall, M., Vila, R., *et al.*

(2019) The determinants of genetic diversity in butterflies. *Nature Communications*, **10**,

3466.

Marcussen, T., Sandve, S.R., Heier, L., Spannagl, M., Pfeifer, M., The International Wheat

Genome Sequencing Consortium, *et al.* (2014) Ancient hybridizations among the

ancestral genomes of bread wheat. *Science*, **345**, 1250092.

Martin, M. (2011) Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet.journal*, **17**, 10–12.

Mayer, K.F.X., Martis, M., Hedley, P.E., Šimková, H., Liu, H., Morris, J.A., *et al.* (2011)

Unlocking the Barley Genome by Chromosomal and Comparative Genomics. *The Plant*

Cell, **23**, 1249–1263.

Muyle, A., Martin, H., Zemp, N., Mollion, M., Gallina, S., Tavares, R., *et al.* (2021) Dioecy Is

Associated with High Genetic Diversity and Adaptation Rates in the Plant Genus Silene.

Molecular Biology and Evolution, **38**, 805–818.

Muyle, A., Serres-Giardi, L., Ressayre, A., Escobar, J. and Glémin, S. (2011) GC-biased

gene conversion and selection affect GC content in the Oryza genus (rice). *Molecular*

Biology and Evolution, **28**, 2695–2706.

Nordborg, M. (2000) Linkage disequilibrium, gene trees and selfing: an Ancestral

recombination graph with partial self-fertilization. *Genetics*, **154**, 923–929.

Pannell, J.R. and Charlesworth, B. (1999) Neutral genetic diversity in a metapopulaiton with

recurrent local extinction and recolonization. *Evolution*, **53**, 664–676.

Paradis, E. & Schliep, K. (2019) ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. *Bioinformatics*, **35**, 526–528.

Parisod, C. & Badaeva, E.D. (2020) Chromosome restructuring among hybridizing wild

wheats. *New Phytologist*, **226**, 1263–1273.

Pollak, E. (1987) On the theory of partially inbreeding finite populations. I. Partial selfing.

Genetics, **117**, 353–360.

R Core Team. (2018) *R: A language and environment for statistical computing*. R

Foundation for Statistical Computing, Vienna, Austria.

Ranwez, V., Harispe, S., Delsuc, F. & Douzery, E.J.P. (2011) MACSE: Multiple Alignment of

Coding SEquences accounting for frameshifts and stop codons. *PLoS ONE*, **6**, e22594.

Rezvoy, C., Charif, D., Guéguen, L. & Marais, G.A.B. (2007) MareyMap: an R-based tool

with graphical interface for estimating recombination rates. *Bioinformatics*, **23**, 2188–

2189.

Rodgers-Melnick, E., Vera, D.L., Bass, H.W. and Buckler, E.S. (2016) Open chromatin

reveals the functional maize genome. *Proceedings of the National Academy of Sciences*,

113, E3177–E3184.

Romiguier, J., Gayral, P., Ballenghien, M., Bernard, A., Cahais, V., Chenuil, A., *et al.* (2014)

Comparative population genomics in animals uncovers the determinants of genetic

diversity. *Nature*, **515**, 261–263.

Roze, D. (2016) Background Selection in Partially Selfing Populations. *Genetics*, **203**, 937–

957.

Sarah, G., Homa, F., Pointet, S., Contreras, S., Sabot, F., Nabholz, B., *et al.* (2017) A large

set of 26 new reference transcriptomes dedicated to comparative population genomics in

crops and wild relatives. *Molecular Ecology Resources*, **17**, 565–580.

Schoen, D.J. & Brown, A.H. (1991) Intraspecific variation in population gene diversity and

effective population size correlates with the mating system in plants. *Proceedings of the*

National Academy of Sciences, **88**, 4494–4497.

Sicard, A. & Lenhard, M. (2011) The selfing syndrome: a model for studying the genetic and

evolutionary basis of morphological adaptation in plants. *Annals of Botany*, **107**, 1433–

1443.

Simpson, J.T., Wong, K., Jackman, S.D., Schein, J.E., Jones, S.J.M. and Birol, İ. (2009)

ABySS: A parallel assembler for short read sequence data. *Genome Research*, **19**,

1117–1123.

Slageren, M.W. van. (1994) Wild wheats: a monograph of Aegilops L. and Amblyopyrum

(Jaub. and Spach) Eig (Poaceae). *Wageningen Agricultural University Papers*, **94–7**.

Slotte, T., Hazzouri, K.M., Ågren, J.A., Koenig, D., Maumus, F., Guo, Y.-L., *et al.* (2013) The

Capsella rubella genome and the genomic consequences of rapid mating system

evolution. *Nature Genetics*, **45**, 831–835.

Stebbins, G.L. (1957) Self fertilization and population variability in the higher plants.

American Naturalist, **91**, 337–54.

Tataru, P. & Bataillon, T. (2019) polyDFEv2.0: testing for invariance of the distribution of

fitness effects within and across species. *Bioinformatics*, **35**, 2868–2869.

Tataru, P., Mollion, M., Glémin, S. & Bataillon, T. (2017) Inference of Distribution of Fitness

Effects and Proportion of Adaptive Substitutions from Polymorphism Data. *Genetics*, **207**,

1103–1119.

Teterina, A.A., Willis, J.H., Lukac, M., Jovelin, R., Cutter, A.D., Phillips, P.C. (2023) Genomic

diversity landscapes in outcrossing and selfing *Caenorhabditis* nematodes. *PLOS*

Genetics **19**, e1010879.

Tsagkogeorga, G., Cahais, V. & Galtier, N. (2012) The population genomics of a fast

evolver: high levels of diversity, functional constraint, and molecular adaptation in the

tunicate Ciona intestinalis. *Genome Biology and Evolution*, **4**, 852–861.

Wasmuth, J.D. & Blaxter, M.L. (2004) prot4EST: Translating Expressed Sequence Tags

from neglected genomes. *BMC Bioinformatics*, **5**, 187.

Wright, S.I., Kalisz, S. & Slotte, T. (2013) Evolutionary consequences of self-fertilization in

plants. *Proceedings of the Royal Society B: Biological Sciences*, **280**, 20130133.

Yang, Z. (2007) PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Molecular Biology*

and Evolution, **24**, 1586–1591.

Ye, Y., Choi, J.-H. & Tang, H. (2011) RAPSearch: a fast protein similarity search tool for

short reads. *BMC Bioinformatics*, **12**, 159.

Acknowledgements

We thank G. Sarah, Y. Holtz and P. Joncour for help with bioinformatic analyses and T. Bataillon, M. Lascoux and D. Schoen for helpful discussions and suggestions on the manuscript. The work was funded by the French Agence Nationale de la Recherche (ANR) (ANR-11-BSV7-013-03). CB has also received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 839643. The authors declare no conflicts of interest.

Data and code availability

Custom R and bash codes used for the analyses are available on https://github.com/sylvainglemin/ms-rec-triticeae along with input files. Software for genotype calling (reads2snps v. 2.0.64, ORF_extractor.pl) and polymorphism estimates (dNdSpNpS v.3) are available at https://kimura.univ-montp2.fr/PopPhyl/index.php?section=tools. Morphological traits measures are provided as Supplementary Tables S2 and S3.

Filtered and cleaned sequence alignments to perform polymorphism analyses are available at https://bioweb.supagro.inra.fr/WheatRelativeHistory/index.php?menu=downloadMating. Raw data are deposited at the Sequence Read Archives (SRA) under project PRJNA945064 (submission number SUB12943046).

Supplementary Information

Figures S1 to S14

Supplementary tables:

Supplementary Table S1. Passport information for the accessions of Triticum/Aegilops

species and outgroups used in this study.

Supplementary Table S2. Dataset 1: Morphofunctional traits linked to reproduction

measured in 13 Triticum/Aegilops species and three outgroups.

Supplementary Table S3. Dataset 2: Morphofunctional traits linked to reproduction

measured in 13 Triticum/Aegilops species and three outgroups where missing data were

imputed with package missMDA under R environment (see Materials and Methods).

Supplementary Table S4. List of morphological traits measured and used for the analysis.

Supplementary Table S5. Sequencing and genome summary statistics per species (13

Triticum/Aegilops species and three outgroups).

Supplementary Table S6. Comparison of genes mapped on the Hordeum genome with the

characteristics of the three Triticum sub-genomes (A, B and D).

Supplementary Table S7. Results of the linked-selection model.

1010

1011

1012

1013

Figure 1. **Mating system diversity in the genus *Aegilops/Triticum***. (A) Phylogenetic

relationships among the 13 diploid species of the genus, following Glémin *et al*. (2019b).

Colors have been assigned to represent a gradient of self fertilization: *Ae. speltoides* and

Ae. mutica have previously been described as self-incompatible (SI, blue tone colors), *Ae.*

tauschii and *Triticum* sp. are known to be prevalently self-fertilizing species (pink-purple

colors), while the remaining species are self-compatible (SC, yellow-green colors). The

arrow indicates when self-compatibility may have appeared, if we assume that it appeared

once according to a parsimony approach. Pictures illustrate the phenotypes associated with

different mating systems: SI *Ae. speltoides*, highly selfing *Ae. tauschii* and self-compatible

Ae. caudata. (B) Principal Component Analysis of morphological measures of reproductive

organs. PC1 resumes the morphological selfing syndrome. On one side there are the self-

incompatible species, which show bigger anther and stigmas, higher male investment, lower

spikelet compactness and lower autonomous seed set. On the other side there are

predominately selfing species, with opposite states of the traits. (C) Positive correlation

between PC1 of reproductive traits and the inbreeding coefficient *F* estimated on the

genomic data shows that species with stronger morphological selfing syndrome have also

higher *F* values.

1031

1032 **Figure 2. Mating systems traits and global polymorphism patterns**. Neutral genetic

1033 diversity π_S is negatively correlated with the first axis of a PCA of morphofunctional traits

1034 associated with reproduction (A), while the efficacy of selection estimated with π_N/π_S ratio is

1035 positively correlated with the same PC1 axis (B). Species geographical range estimated from

1036 GBIF occurrence data cannot explain the variation observed neither in π_S (C) nor π_N/π_S (D).

1037

Figure 3. **Recombination and patterns of genetic polymorphism across the genome.**

(A) Mean synonymous diversity (π_S) along chromosome 3 for three species with contrasted

mating system: self-incompatible (*Ae. mutica*), mixed-mating (*Ae. sharonensis*) and highly

selfing (*T. urartu*); same scales for the three species. Each point corresponds to a contig

mapped on the *Hordeum vulgare* genome. The blue line is the loess fitting function

(degree=2, span=0.1). The dashed black line indicates *H. vulgare* recombination map (in

cM/Mb). (B) π_S and π_N/π_S as a function of recombination rate. Contigs have been grouped in

20 quantiles of recombination. The value associated with each species corresponds to the

ratio between the highest and the lowest value among the quantiles: from about 5 to 80 for

π_S and from 2.5 to 33 for π_N/π_S. Curves correspond to loess fitting functions (degree = 2,

span = 0.2).

1049

1050 **Figure 4**. **Relation of estimated maximum polymorphism in absence of linked-**

1051 **selection and mating system.** π_{max} does not correlate with the first axis of the PCA of

1052 morphofunctional traits (left). π_{max} correlates negatively with species range (right), but the

1053 correlation is mainly driven by the species of the *Sitopsis* section and is no longer significant

1054 after phylogenetic correction.

1055

Figure 5. **Mating systems, recombination and the efficacy of selection**. The distribution

of fitness effect of mutations was estimated in two self-incompatible species (*Ae. mutica* and

Ae. speltoides) and in two highly selfing species (*A. tauschii* and *T. urartu*) at the whole

genome scale (blue) and for regions of low (green) and high (red) recombination. (A)

Proportion of weakly selected deleterious mutations ($-10 < 2N_eS < 0$). (B) Rate of adaptive

substitution, i.e. the expected D_N/D_S ratio due to the flux of beneficial mutations.

(A)

outgroups

Ae. mutica (Amu)
Ae. speltoides (Asp)   SI

Ae. umbellulata (Aum)
Ae. caudata (Aca)
Ae. comosa (Aco)
Ae. uniaristata (Aun)
Ae. bicornis (Abi)   SC
Ae. longissima (Alo)
Ae. sharonensis (Ash)
Ae. searsii (Ase)
Ae. tauschii (Ata)
T. monococcum (Tmo)   Highly Selfing
T. urartu (Tur)

6   0 MYA

2 cm   *Ae. speltoides*

2 cm   *Ae. caudata*

2 cm   *Ae. tauschii*

(B)

PC2 (21%)

Out-crossing   Selfing

PC1 (53%)

(C)

F

$R^2 = 0.83$ ***
(With phylogenetic correction $R^2 = 0.78$***)

PC1