Employing RNA editing to engineer personalized tumor-specific neoantigens (editopes)

Authors

Riccardo Pecori[1,†,*], Beatrice Casati[1,†], Rona Merdler-Rabinowicz[2,3†], Netanel Landesman[3],

Khwab Sanghvi[4], Stefan Zens[5,6], Kai Kipfstuhl[5,6], Veronica Pinamonti[7,8,9], Annette Arnold[1], John

M. Lindner[8], Michael Platten[4,10,11,12,13,14,15], Rienk Offringa[5,6], Rafael Carretero[5,16], Eytan

Ruppin[2,*], Erez Y. Levanon[3,*], Fotini Nina Papavasiliou[1,*]

Affiliations
[1]Division of Immune Diversity, German Cancer Research Centre (DKFZ), Heidelberg, 69120,
Germany.
[2]Cancer Data Science Lab, Center for Cancer Research, National Cancer Institute, National
Institutes of Health, Bethesda, MD, 20892, United States.
[3]Mina and Everard Goodman Faculty of Life Sciences, Bar-Ilan University, Ramat
Gan, 5290002, Israel.
[4]Division of Neuroimmunology and Brain Tumor Immunology, German Cancer Research Centre
(DKFZ), Heidelberg, 69120, Germany.
[5]Division of Molecular Oncology of Gastrointestinal Tumors, German Cancer Research Centre
(DKFZ), Heidelberg, 69120, Germany.
[6]DKFZ Drug Discovery Lab (D3Lab), German Cancer Research Centre (DKFZ), Heidelberg, 69120,
Germany.
[7]Faculty of Biosciences, Heidelberg University, 69120 Heidelberg, Germany.
[8]BioMed X Institute, 69120 Heidelberg, Germany.
[9]Current affiliation: Department of Experimental Oncology, European Institute of Oncology
IRCCS, Milan, Italy.
[10]DKTK (German Cancer Consortium) Clinical Cooperation Unit (CCU) Neuroimmunology and
Brain Tumor Immunology, German Cancer Research Center (DKFZ), Heidelberg, Germany.
[11]Department of Neurology, Medical Faculty Mannheim, MCTN, University of Heidelberg,
Mannheim, Germany.
[12]German Cancer Consortium (DKTK), DKFZ, Core Center Heidelberg, Germany.
[13]Hertie Network of Excellence in Clinical Neuroscience.
[14]DKFZ-Hector Cancer Institute at University Medical Center Mannheim, Mannheim, Germany.
[15]Helmholtz Institute for Translational Oncology (HI-TRON) Mainz, German Cancer Research
Center, Mainz, Germany.
[16]Department of Biochemistry, molecular Biology 3 and immunology. University of Granada,
Spain.
37
38   [*]Corresponding authors
39   [†]These authors contributed equally to this work.
40

Abstract

Increasing the quantity and immunogenicity of neoantigens in tumors is essential for advancing immunotherapy. However, engineering neoantigens remains challenging due to the need for precise, tumor-specific antigen modification without affecting normal cells. To tackle this challenge, we developed *Short Precise-Encodable ADAR Recruiting* (SPEAR) ADAR-engagers, an approach that uses short guide RNAs to engage the endogenous RNA editor ADAR1 and direct it to regions of mRNA targets known to encode MHC-presented peptides. By precisely editing adenosine-to-inosine (A-to-I) in these contexts, we effectively mutate specific epitopes into neoepitopes (which we now term "editopes"). As proof of concept, we targeted the known antigen MART-1 (Melanoma-Associated Antigen Recognized by T cells-1), and demonstrated that guided ADAR1 editing can generate immunogenic epitopes that activate T cells and promote tumor cell elimination. Building on this concept, we developed a computational pipeline to identify tumor-specific somatic mutations suitable for SPEAR-mediated editing. This strategy enables selective neoantigen generation in cancer cells, effectively increasing their apparent tumor mutational burden and potentially enhancing their susceptibility to immunotherapy.

Introduction

In recent years, immunotherapy has emerged as a revolutionary approach in cancer therapy, demonstrating remarkable clinical response in numerous cases with acceptable patient risk[1]. This success depends on the ability to trigger a robust immune response against the tumor, and requires both the presence of cancer-specific antigens on tumor cells and T lymphocytes bearing antigen-specific T cell receptors (TCR) that recognize and kill them. Responsiveness to immunotherapy is positively correlated with a tumor's mutational burden (TMB)[2–4]. Since only a subset of somatic mutations give rise to neoepitopes that are processed and presented as non-self by the major histocompatibility complex (MHC), tumors with a high TMB are more likely to produce immunogenic neoantigens and to respond favorably to immunotherapy[5].

Neoantigens arise from non-synonymous DNA mutations, but also from aberrant post-transcriptional and post-translational modifications, including aberrant mRNA translation, alternative splicing, and RNA editing[6–8]. However, many cancers harbor an insufficient repertoire of neoantigens capable of eliciting a strong immune reaction[9] and are therefore poorly responsive to immunotherapy. To address this challenge, strategies aimed at artificially enhancing neoantigen production have emerged, such as modulation of alternative splicing[10]. Yet, these approaches are not tumor-specific and can often also affect healthy tissues, potentially leading to adverse side effects, such as autoimmunity. Effective methods to generate specific neoantigens exclusively within tumor cells are still lacking.

Here, we engineer tumor-specific neoantigens by employing ADAR-recruiting gRNAs (reviewed in *REFs*[11–13]) to introduce precise and controlled RNA edits specifically in transcripts exclusively present in cancer cells, thereby generating immunogenic peptides selectively in malignant tissues. Physiologically, Adenosine Deaminase Acting on RNA 1 (ADAR1) is an enzyme that recognizes long double-stranded RNA (dsRNA) structures and converts adenosines (A) to inosines (I), which are interpreted as guanosines (G) by the cellular translation machineries[14,15] (an attribute that has enabled the development of guided editing through the engagement of the endogenous enzyme by provision of an antisense guide RNA - an "ADAR engager"). ADAR1

activity, which is already prolific across the mammalian transcriptome[14], increases in almost all

cancers[16], contributing to transcriptomic diversity and neoantigen generation[7].

While ADAR engagers have been explored in the clinic for the correction of disease-

causing mutations such as premature stop codons, we hypothesized that this same mechanism

could be repurposed to deliberately introduce precise nucleotide substitutions into cancer cell

mRNAs to generate tumor-specific neoantigens. To this end, we set out to develop Short

Precise-Encodable ADAR Recruiting (SPEAR) ADAR-engagers; compact, genetically encodable

guide RNAs specifically designed to recruit endogenous ADAR1 and drive A-to-I RNA editing in a

highly controlled manner. To evaluate the feasibility of this approach, we selected a well-

characterized tumor antigen, MART-1 (Melanoma-Associated Antigen Recognized by T cells-1).

We engineered a non-immunogenic variant of it as a model system to assess whether targeted

A-to-I editing could restore T cell recognition and cytotoxicity, thereby demonstrating the

potential of RNA editing to induce functional neoantigen formation directly from endogenous

transcripts.

Building on this conceptual framework, we also designed a computational pipeline to

systematically identify tumor-specific somatic mutations that could serve as substrates for

SPEAR-mediated editing. Because these editing events are designed to target only transcripts

harboring tumor-specific mutations, the strategy offers the potential for tumor-selective

neoantigen formation with minimal risk to healthy tissues. This approach provides a

personalized and safe avenue for enhancing tumor immunogenicity and expanding the

repertoire of actionable neoantigens for cancer immunotherapy.

Results

Optimization of Short Precise-Encodable ADAR1 Recruiting (SPEAR) gRNAs

Several methods have enabled precise site-directed RNA editing using engineered or

endogenous editing enzymes. A particularly useful approach relies on ADAR-engagers that

base-pair with target transcripts to form dsRNA structures capable of recruiting endogenous

ADAR1 to specific RNA sites[17]. This method offers several therapeutic advantages: it eliminates the need for exogenous protein (these programmed oligonucleotides can be delivered directly into cells), reduces the molecular payload (ADAR-engagers consisting of merely a few dozen nucleotides (nt) prove effective variants[18–21], and minimizes the risk of an anti-drug immune response (allowing for repeat dosing).

ADAR-engagers can be delivered as chemically modified antisense oligonucleotides (ASOs), as genetically encoded constructs via plasmids or viral particles, or as *in vitro* transcribed RNA moieties. Upon hybridization with target mRNAs, gRNAs form duplexes that recruit endogenous ADAR enzymes to catalyze A-to-I editing[19–27]. These gRNAs generally fall into two main categories: (1) long gRNAs (>100 nt) with extended antisense "specificity domains" for target pairing[20,22,24], and (2) shorter, structured gRNAs that include a compact specificity domain along with a "recruitment domain" RNA motif to enhance ADAR recruitment[19,21,23,28]. Long gRNAs are prone to extensive bystander off-target editing, requiring time-consuming, target-specific optimization[20,22,24]. In contrast, short, structured gRNAs exhibit high precision and editing efficiency when chemically modified and delivered as ASOs[18,21], but show poor performance when genetically encoded[29]. More complex designs using clustered ADAR recruitment motifs offer improved specificity, though they involve intricate design rules and require non-intuitive computational tools[19,23]. To overcome these challenges, we sought to optimize a genetically encodable gRNA that could efficiently and precisely recruit endogenous ADAR1. Our goal was to minimize the gRNA structure required for efficient ADAR1 recruitment, thereby developing a simple and flexible platform for therapeutic deployment. Such optimized gRNAs could be delivered as plasmid- or virus-encoded constructs, or alternatively as chemically stabilized RNA molecules for direct administration.

To evaluate the capability of such gRNAs to recruit endogenous ADARs, we generated a dual-reporter HEK293T cell line stably expressing mCherry and a mutated eGFP containing a premature stop codon (W58X) linked by a P2A self-cleaving peptide[30]. In the absence of editing, only mCherry is expressed. Successful recruitment of ADAR to the UAG codon converts this stop codon to UIG, which is interpreted as a tryptophan codon (UGG), thereby restoring eGFP

fluorescence (**Figure 1A** and *REF*[31]). This system enabled rapid, quantitative assessment of gRNA

activity via flow cytometry. Additionally, HEK293T cells express ADAR1 but not ADAR2[20],

allowing us to specifically assess ADAR1-dependent editing. Notably, the use of a stably

integrated reporter mimics the behavior of an endogenous transcript more accurately than co-

transfection of guide and target RNAs, thereby offering a more realistic and relevant testing

environment.

We began by examining the impact of three structural features of encoded gRNAs: (1)

specificity domain length, (2) the distance between the target adenosine and the recruitment

domain, and (3) the recruitment domain itself (**Figure S1A**). Consistent with prior work, we

confirmed that long, unstructured gRNAs were more effective than structured gRNAs when

expressed from plasmids (**Figure S1B**), and that editing efficiency improved upon inclusion of a

4-nt mismatch placed 35 nt downstream of the target adenosine[32] (**Figure S1B,C**). However,

amplicon sequencing revealed that longer gRNAs incurred greater off-target editing across

adjacent adenosines, validating concerns about precision and bystander activity (**Figure S1D-E**).

This analysis also confirmed the flow cytometry results for on-target editing, validating the

robustness of our dual-reporter cell line for gRNA screening (**Figure S1D**).

To systematically optimize gRNA architecture, we designed a series of truncated gRNAs

based on the long LEAPER construct with a 4-nt mismatch. We varied the length of the 5▯ and

3▯ arms independently (**Figure 1B,C**). Reducing the 3▯ arm from 75 to 16 nt resulted in a 1.5-

fold increase in mCherry⁺eGFP⁺ cells. Truncating the 5▯ arm (externally of the 4-nt mismatch)

from 38 to 20 nt resulted in a modest gain in editing, but further shortening to 10 nt drastically

reduced activity (**Figure 1C**). These findings suggest that the duplex formed by the 5' arm of the

gRNA is essential for ADAR1 recruitment and positioning.

Combining both optimizations, we generated a compact 74-nt-long gRNA with reduced

5▯ and 3▯ arms, half the size of the original 151-nt LEAPER. This shortened design yielded the

highest editing efficiency, as measured by fluorescence restoration (**Figure 1C**), and exhibited

reduced off-target editing, as determined by amplicon sequencing (**Figure 1D,E**). Collectively,

these results demonstrate that the efficiency and precision of plasmid-encoded ADAR-engagers
can be significantly improved through rational design of duplex geometry. Specifically, we
identify the length of the duplex region downstream of the 4 nt-mismatch as a key determinant
of ADAR1 recruitment. We refer to this optimized design as *Short Precise-Encodable ADAR*
Recruiting (SPEAR) gRNAs.

SPEAR-mediated RNA editing restores T cell activation

To test whether endogenous ADAR1 RNA editing could be leveraged with SPEAR gRNAs
to generate functional *editopes* capable of activating antigen-specific T cells, we designed a
proof-of-concept experiment based on the well-characterized melanoma antigen MART-1
(Melanoma Antigen Recognized by T cells-1)[33,34]. MART-1 is encoded by the *MLANA* gene and is
a clinically relevant immunogenic target in melanoma. Critically for our purposes, the specific T
cell receptor (TCR) recognizing the MART-1 epitope (EAAGIGLTV) is well characterized. For this,
we used the DMF5 TCR, known for its high avidity against MART-1-expressing cells[35].

To increase T cell sensitivity and stability of the peptide-MHC complex, we used a MART-
1 peptide variant (E**L**AGIGILTV)[36] (**Figure 2A, S2A**). We then introduced a point mutation in the
MART-1 coding sequence to generate the G31S mutant (MART-1^{G31S}; ELAGI**S**ILTV), substituting
glycine (**G**GC) with serine (**A**GC) at position 31. This mutation disrupted the epitope, rendering
it incapable of activating DMF5 T cells (**Figure 2A, B, S2A**). To reverse this effect via RNA editing,
we designed a SPEAR gRNA targeting the mutant *MART-1^{G31S}* transcript to recruit endogenous
ADAR1 and direct A-to-I editing at the first base of the serine codon (**A**GC), converting it to IGC
(interpreted by the ribosome as GGC) at the RNA level and restoring the original epitope (**Figure**
2A, S2A). This system models our broader goal of generating immunogenic *editopes* from non-
immunogenic epitopes through precise RNA editing.

To evaluate whether SPEAR-driven ADAR1 editing could restore antigenicity and T cell
activation, we engineered HEK293T and TCR-deficient Jurkat cell lines to stably express either
MART-1^{WT} or *MART-1^{G31S}*, and the DMF5 TCR, respectively. HEK293T cells were also expressing
either a non-targeting (NT) or targeting (T) SPEAR gRNA (**Figure S2B, C**). Sanger sequencing of

PCR amplicons from genomic DNA and cDNA in *MART-1*^{G31S} cells expressing the T-gRNA

confirmed A-to-I editing at the intended site, with an editing efficiency of approximately 20%

(**Figure 2B**). We then co-cultured the engineered HEK293T and Jurkat cells and assessed T cell

activation via flow cytometry using CD2 as a T cell marker and CD69 as an activation marker. As

expected, cells expressing *MART-1*^{WT} with a NT-gRNA elicited robust T cell activation (~65%

CD2$^+$CD69$^+$ cells), while *MART-1*^{G31S} cells with NT-gRNA showed minimal activation (~5%).

Importantly, co-cultures with *MART-1*^{G31S} cells expressing the T-gRNA led to a significant

restoration of T cell activation (~30% CD2$^+$CD69$^+$ cells) (**Figure 2C, D**).

Despite the "moderate" level of editing, recovery of T cell activation was substantial

(~50%) relative to the wild-type epitope, suggesting that even modest RNA editing rates (<30%)

may be sufficient to elicit strong T cell responses and potentially therapeutic immune

activation. These results demonstrate that SPEAR gRNAs can efficiently harness endogenous

ADAR1 to generate immunogenic peptides capable of activating a specific T cell response. Our

findings establish a mechanistic foundation for using programmable RNA editing to engineer

editope formation in tumor cells.

Programmable RNA editing drives T cell-mediated tumor rejection in an *in vivo* preclinical

model

Having demonstrated that SPEAR-mediated RNA editing can restore cognate T cell

activation *in vitro*, we next evaluated its therapeutic potential *in vivo* using a xenograft mouse

model. For this, we employed the human melanoma cell line 624-mel, which endogenously

expresses MART-1. We first introduced the same G-to-A mutation described above into the

endogenous *MLANA* gene using DNA base editing, resulting in expression of the mutant MART-

1^{G31S} from the endogenous locus (**Figure S3A**). These cells were further engineered to stably

express either a targeting (T) or a non-targeting (NT) SPEAR gRNA designed to recruit

endogenous ADAR1 to MART-1^{G31S} (**Figure S3B-D**). We initially assessed editing efficiency in

these modified cell lines and observed ~10% A-to-I editing at the endogenous target site in the

cell line expressing MART-1^{G31S} together with the T-gRNA, while no editing was detected in cells

expressing the NT-gRNA (**Figure 3A, B**). To test whether this level of editing was sufficient to

induce T cell activation, we performed an *in vitro* T cell-mediated cytotoxicity assay using the

xCELLigence system. As expected, 624-mel MART-1^{WT} cells grew normally in the absence of T

cells, but the addition of MART-1-specific T cells resulted in a strong antitumoral effect (**Figure**

3C, green lines). In contrast, MART-1^{G31S} cells with no or the control NT-gRNA were unaffected

by T cell addition, indicating a lack of recognition by the latter (red and gray lines). Notably,

MART-1^{G31S} cells expressing the T-gRNA showed robust killing by T cells, with a cytotoxicity

profile similar to MART-1^{WT} cells (**Figure 3C**, orange dotted line). Despite only ~10% RNA editing

at the G31S site, the resulting partial correction was sufficient to generate a functional

immunogenic *editope*, again underscoring that even modest editing can restore effective T cell

recognition (**Figure 3A-C**).

We next evaluated this approach *in vivo* using a melanoma xenograft mouse model.

Immunodeficient mice were injected subcutaneously with either MART-1^{WT} or MART-1^{G31S} cells

to induce tumor formation. After tumors were established, mice received adoptive T cell

therapy (ACT) with MART-1 specific T cells (**Figure S3E**). As expected, tumors derived from

MART-1^{WT} cells were completely eliminated following two rounds of ACT, while tumors in

untreated controls progressed to the experimental endpoint (**Figure S3F**), validating the

working parameters of the system. To assess the effect of RNA editing in this context, we

repeated the experiment, but now injecting MART-1^{G31S} cells expressing either the NT- or T-

SPEAR gRNA into immunodeficient mice (**Figure 3D**). Mice injected with MART-1^{G31S} cells

expressing the NT-gRNA failed to respond to ACT, and tumors continued to grow. In contrast,

tumors derived from MART-1^{G31S} cells expressing the T-gRNA showed marked regression

following ACT (**Figure 3E, S3G**). Furthermore, we also observed prolonged survival for mice

carrying tumors derived from MART-1^{G31S} cells expressing the T-gRNA relative to those

expressing the NT-gRNA (**Figure 3F**).

Together, these findings demonstrate that SPEAR gRNAs can successfully reprogram

tumor cells by recruiting endogenous ADAR1 to generate immunogenic neoepitopes, or

editopes, enabling precise T cell targeting and tumor rejection.

A pipeline for the generation of patient-specific *editopes* across seven cancer types

To extend our proof of concept findings beyond single antigen models and into a broader oncologic context, we developed and applied our neoADARgen algorithm (explained in detail in Methods) to genomic and transcriptomic data from The Cancer Genome Atlas (TCGA). We focused on seven major cancer types: breast cancer (BRCA), skin melanoma (SKCM), lung adenocarcinoma (LUAD), lung squamous cell carcinoma (LUSC), glioblastoma (GBM), ovarian cancer (OV) and liver hepatocellular carcinoma (LIHC). For each of the 3,224 patients in these cohorts, we integrated somatic mutation profiles, HLA haplotype and gene expression data to predict the likelihood that individual mutations could result in neoantigen formation. In total, 808,297 somatic mutations were analyzed, resulting in 19,311 predicted novel neoantigens across all HLA haplotypes. Prior to applying NeoADARgen, patients harbored on average of 4.3 predicted *epitopes*. Following the algorithm's application, this number increased to an average of 10.3 epitopes per patient, demonstrating a consistent enhancement in predicted immunogenicity across all cancer types (**Table 1**).

TCGA project	Total number of patients included	Epitopes prior to NeoADARgen algorithm (total / avg. per patient)	Patients with novel epitopes identified	Novel epitopes predicted (total / avg. per patient)	Shared novel epitopes
BRCA	965	2431/2.5	801	3675/4.6	39
SKCM	467	3089/6.6	434	4642/10.7	41
OV	289	3323/11.5	273	3190/11.7	6
GBM	147	188/1.3	107	289/2.7	4
LIHC	362	702/1.9	267	1029/3.9	4
LUAD	507	1999/3.9	474	3185/6.7	27
LUSC	487	2029/4.2	466	3301/7.1	15

Table 1: Summary of NeoADARgen-predicted neoantigens across seven TCGA cancer cohorts. Application of the NeoADARgen algorithm to 3,224 TCGA patients revealed a consistent increase in predicted neoantigens across all analyzed cancer types. The table summarizes cohort sizes, baseline epitope counts, and the number of novel epitopes identified following RNA-editing–based prediction.

For instance, in breast cancer (BRCA), patients initially exhibited an average of 2.5

predicted neoantigens. After applying our algorithm, 801 patients were predicted to generate a

total of 3,675 *editopes*, corresponding to an average of about 6.8 neoantigens per patient).

These *editopes* could, in principle, be generated by the delivery of multiple SPEAR ADAR-

engagers, thereby increasing the number of recognizable antigens and enhancing the likelihood

of effective immune targeting in low-TMB tumors.

Engineering shared neoepitopes between individuals

ADAR-engager based therapies, whether aimed at correcting pathogenic variants or

generating personalized *editopes*, are inherently limited in scalability. To broaden the

applicability of our editope-generation approach to a wider patient population, we sought to

identify a limited set of shared *editopes* (and their corresponding SPEAR-gRNAs) that could

cover a large fraction of cancer cases. To this end, we analyzed the 50 most frequent somatic

mutations documented in the NCI Genomic Data Commons (GDC) database[37] across all cancer

types, along with the 50 most common HLA alleles represented in the same database. We then

applied our algorithm to evaluate each possible mutation-HLA combination and identify

potential *editopes* predicted to be strong neoantigens.

For each combination, we first used a peptide-HLA binding prediction tool to calculate

the best possible rank before editing[38]. If no high-affinity neoantigen was predicted at a

particular site, we ran our algorithm to search for nearby adenosines that could be targeted for

editing to generate a neoantigenic peptide. Only combinations predicted to yield a potent

neoantigen after editing were retained as viable (**Figure 4**). As an illustrative example, KRAS,

one of the most frequently mutated oncogenes, harbors mutations at codon 12, where

substitution of glycine is a common driver event in pancreatic, colorectal, and lung cancers.

These mutations lead to the constitutive activation of the KRAS protein, promoting oncogenic

signaling and uncontrolled proliferation[21]. According to the GDC database, KRAS^{G12} mutations

occur in approximately 7% of all cancers.

Focusing on this hotspot, our algorithm identified a candidate *editope* for four common

HLA types by targeting an adenosine located eleven bases downstream of the mutation site.

The corresponding mRNA sequence is:

CUUGUGGUAGUUGGAGCUG**X**UGGCGUAGGC*****AGAGUGCCU, where **X** represents the mutation

site (as multiple SNVs at this hotspot are known KRAS mutations), and the asterisk (*) denotes

the adenosine predicted for editing. For this target, the SPEAR gRNA sequence would be

UUGGAUCAUAUUCGUCCACA<u>UUUA</u>GAUUCUGAAUUAGCUGUA

UCGUCAAGGCACUCU<u>C</u>GCCUACGCCA**Y**CAGCU, where the underlined bases are mismatched to

the mRNA sequence for editing purposes and **Y** indicates the base complementary to **X**. The

resulting neoantigen achieved a predicted rank of 0.01% and a binding affinity of 6.35 nM.

Notably, this same *editope* can also be generated for tumors carrying mutations in KRAS[G13], as

well as NRAS[G12] or NRAS[G13] due to gene homology, requiring only minor adjustments to the

designed gRNA, potentially extending applicability to an additional 2% of cancers. This example

highlights how our approach can exploit recurrent hotspot mutations to generate *editopes*

across multiple patients. Such shared targets could enable the development of off-the-shelf

SPEAR gRNAs for tumors with common mutational profiles, substantially increasing the clinical

and pharmaceutical utility of the editope generation strategy.

Discussion

Enhancing the number and immunogenicity of neoantigens within tumors is a critical

step for improving the efficacy of immunotherapy. This is especially relevant for cancer

vaccines, where the ability to design potent, tumor-specific neoantigens enables the patient to

direct an immune response against these engineered neoantigens. However, the selective

generation of neoantigens remains challenging.

Here, we introduce a novel approach for neoantigen engineering based on the

recruitment of endogenous ADAR1 via SPEAR gRNAs. This technology enables programmable

RNA editing through gRNA delivery and without the need for exogenous enzymes. Furthermore,

base editing of RNA rather than DNA enhances safety by reducing the risk of permanent

genomic alterations. In our proof of concept, we demonstrated that ADAR1-mediated RNA

editing can modulate the immunogenicity of a tumor epitope by altering its RNA sequence

while maintaining its translation, processing, and presentation by MHC-I molecules. This RNA-

edited epitope, termed *editope*, can be recognized by and activate antigen-specific T cells, inducing T cell-mediated cytotoxicity.

After establishing and demonstrating the feasibility of *editope* generation both *in vitro* and *in vivo*, we developed a computational pipeline capable of designing SPEAR gRNAs tailored to each patient's HLA genotype but derived from the mutational landscape of a population of tumors. This personalization ensures that edits are introduced exclusively in tumor cells— anchored to somatic mutations—and that the resulting peptides are highly likely to bind MHC-I and be presented as neoantigens.

Applying our computational algorithm to thousands of tumor samples across seven cancer types, we demonstrated that our approach could nearly triple the number of predicted potent neoantigens for the majority of patients. We also showcased several editing candidates derived from prevalent somatic mutations that can serve as entry points for broad clinical applications of oncologic patients.

While our *in silico* predictions provide a compelling roadmap, we acknowledge the complexity of immune recognition. We relied on an established MHC-binding prediction tool that estimates the likelihood of a peptide binding to a specific MHC molecule based on the top percentile scores obtained from random natural peptides[38]. Additionally, we incorporated gene expression levels into our scoring. However, accurate modeling of neoantigen presentation remains challenging due to the involvement of multiple cellular processes, including antigen processing, peptide trimming, proteasomal cleavage, and peptide loading. Currently, modeling these processes computationally remains highly complex[39]. Of course, there are general mechanisms that enable tumor cells to escape immune surveillance, notably the somatic loss of HLA-I heterozygosity, which is a known factor in immune evasion[40]. This is a risk that our platform shares with neoepitope vaccines; however, recent data support substantial efficacy for such vaccines[41], which theoretically our platform should also share.

A general limitation of ADAR engagers is editing efficiency. Previous studies have reported variable editing levels across targets and delivery formats[19–27]. In our context, high editing efficiency is not necessary: the generation of even a small number of edited transcripts

appears to suffice to trigger T cell recognition, as we observed both *in vitro* and *in vivo*. Unlike

conventional base editing applications that seek full protein restoration, our goal is to create

peptides that are processed and presented as *non-self*, enabling immune activation by a

relatively small fraction of edited mRNAs and for this, even low editing efficiency is sufficient.

A final general limitation of targeted ADAR1 recruitment lies in the delivery of the gRNAs to the

tissues of interest. This is less applicable to our technology: our focus on *editopes* that are

translated from tissue/HLA specific tumour exclusive transcripts, already supports a level of

selectivity even to systemic delivery. However as delivery technologies continue to advance, we

anticipate that the innovative approach presented here could serve as a valuable therapeutic

adjunct for immunotherapy. In all, the approach we present, harnessing programmable RNA

editing via SPEAR gRNAs and endogenous ADAR1, offers a safer, precise, and adaptable

platform for generating tumor-specific neoantigens or *editopes*. This technology has the

potential to strongly enhance immunogenicity in poorly immunogenic ("cold") tumors and to

serve as a powerful complement to current immunotherapeutic strategies, including checkpoint

inhibitors and personalized cancer vaccines.

Acknowledgements and funding

We thank the Flow Cytometry unit of the Imaging and Cytometry Core Facility, German Cancer

Research Center (DKFZ), for providing excellent sorting services. We also acknowledge Dr.

Florian Grünschläger for his support and guidance in data visualization using R. This work was

supported by the Israel Science Foundation grants 2039/20 and 231/21 to E.Y.L., the HI-TRON

Kick-Start Seed Funding Program 2021 awarded to RP, and by the Deutsche

Forschungsgemeinschaft (DFG, German Research Foundation; SVB TRR319-RMaP) to F.N.P. as

well as core funds to F.N.P. by the Helmholtz Association.

Author contributions

R.P. and F.N.P. conceived the project and supervised the research. R.M.R., N.L., E.R., and E.Y.L.

conceived and developed the computational algorithm. R.P. and A.A. conceived, performed,

and analyzed targeted RNA editing experiments. V.P. and J.M.L. contributed to the co-culture

assays. R.P., B.C., K.S., S.Z., K.K., R.C., M.P., and R.O. conceived, performed, and analyzed
experiments in mice. R.P., B.C., and F.N.P. interpreted the overall results of the study. R.P., B.C.,
and R.N.R. prepared the data visualizations. R.P., B.C., and R.N.R. wrote the manuscript with
input from all authors. All authors reviewed and approved the final version of the manuscript.
385

Data Availability
The source code used to produce the results and analyses presented in this manuscript are
available on a GitHub repository at: https://github.com/landsboy/neoADARgen.git

Competing interests

R.P. and A.A. are inventors on a patent application related to the design of ADAR-recruiting
guide RNAs or ASOs (PCT/EP2024/051033) and R.P., B.C., A.A. and F.N.P. on a separate
application covering the use of *editopes*. All other authors declare no competing interests.

References

1. Esfahani, K. *et al.* A review of cancer immunotherapy: from the past, to the present, to the
future. *Curr Oncol* **27**, S87–S97 (2020).
2. Chan, T. A. *et al.* Development of tumor mutation burden as an immunotherapy biomarker:
utility for the oncology clinic. *Ann Oncol* **30**, 44–56 (2019).
3. Yarchoan, M., Hopkins, A. & Jaffee, E. M. Tumor Mutational Burden and Response Rate to
PD-1 Inhibition. *N Engl J Med* **377**, 2500–2501 (2017).
4. Samstein, R. M. *et al.* Tumor mutational load predicts survival after immunotherapy across
multiple cancer types. *Nat Genet* **51**, 202–206 (2019).
5. Schumacher, T. N. & Schreiber, R. D. Neoantigens in cancer immunotherapy. *Science* **348**,
69–74 (2015).

6. Nagel, R., Pataskar, A., Champagne, J. & Agami, R. Boosting Antitumor Immunity with an

Expanded Neoepitope Landscape. *Cancer Res* **82**, 3637–3649 (2022).

7. Zhang, M. *et al.* RNA editing derived epitopes function as cancer antigens to elicit immune

responses. *Nature Communications* https://doi.org/10.1038/s41467-018-06405-9 (2018)

doi:10.1038/s41467-018-06405-9.

8. Smart, A. C. *et al.* Intron retention is a source of neoepitopes in cancer. *Nature*

Biotechnology https://doi.org/10.1038/nbt.4239 (2018) doi:10.1038/nbt.4239.

9. Bonaventura, P. *et al.* Cold Tumors: A Therapeutic Challenge for Immunotherapy. *Front*

Immunol **10**, 168 (2019).

10. Lu, S. X. *et al.* Pharmacologic modulation of RNA splicing enhances anti-tumor immunity.

Cell **184**, 4032-4047.e31 (2021).

11. Casati, B., Stamkopoulou, D., Tasakis, R. N. & Pecori, R. ADAR-Mediated RNA Editing

and Its Therapeutic Potentials. in *Epitranscriptomics* (eds Jurga, S. & Barciszewski, J.) 471–

503 (Springer International Publishing, Cham, 2021). doi:10.1007/978-3-030-71612-7_18.

12. Aquino-Jarquin, G. Novel Engineered Programmable Systems for ADAR-Mediated RNA

Editing. *Mol Ther Nucleic Acids* **19**, 1065–1072 (2020).

13. Vogel, P. & Stafforst, T. Critical review on engineering deaminases for site-directed RNA

editing. *Current opinion in biotechnology* **55**, 74–80 (2018).

14. Tan, M. H. *et al.* Dynamic landscape and regulation of RNA editing in mammals. *Nature*

550, 249–254 (2017).

15. Porath, H. T., Knisbacher, B. A., Eisenberg, E. & Levanon, E. Y. Massive A-to-I RNA

editing is common across the Metazoa and correlates with dsRNA abundance. *Genome*

Biology **18**, 185 (2017).

16. Paz-Yaacov, N. *et al.* Elevated RNA Editing Activity Is a Major Contributor to

Transcriptomic Diversity in Tumors. *Cell Reports* **13**, 267–276 (2015).

17. Booth, B. J. *et al.* RNA editing: Expanding the potential of RNA therapeutics. *Mol Ther* **31**,

1533–1549 (2023).

18. Monian, P. *et al.* Endogenous ADAR-mediated RNA editing in non-human primates using

stereopure chemically modified oligonucleotides. *Nat Biotechnol* **40**, 1093–1102 (2022).

19. Reautschnig, P. *et al.* CLUSTER guide RNAs enable precise and efficient RNA editing with

endogenous ADAR enzymes in vivo. *Nat Biotechnol* **40**, 759–768 (2022).

20. Qu, L. *et al.* Programmable RNA editing by recruiting endogenous ADAR using engineered

RNAs. *Nat Biotechnol* **37**, 1059–1069 (2019).

21. Merkle, T. *et al.* Precise RNA editing by recruiting endogenous ADARs with antisense

oligonucleotides. *Nat Biotechnol* **37**, 133–138 (2019).

22. Katrekar, D. *et al.* Efficient in vitro and in vivo RNA editing via recruitment of endogenous

ADARs using circular guide RNAs. *Nat Biotechnol* **40**, 938–945 (2022).

23. Reautschnig, P. *et al.* Precise in vivo RNA base editing with a wobble-enhanced circular

CLUSTER guide RNA. *Nat Biotechnol* **43**, 545–557 (2025).

24. Yi, Z. *et al.* Engineered circular ADAR-recruiting RNAs increase the efficiency and fidelity

of RNA editing in vitro and in vivo. *Nat Biotechnol* **40**, 946–955 (2022).

25. Byrne, S. M. *et al.* An engineered U7 small nuclear RNA scaffold greatly increases ADAR-

mediated programmable RNA base editing. *Nat Commun* **16**, 4860 (2025).

26. Smargon, A. A. *et al.* Enhancing RNA base editing on mammalian transcripts with small

nuclear RNAs. *Nat Chem Biol* https://doi.org/10.1038/s41589-025-02026-8 (2025)

doi:10.1038/s41589-025-02026-8.

27. Sun, Y. *et al.* Improved RNA base editing with guide RNAs mimicking highly edited

endogenous ADAR substrates. *Nat Biotechnol* https://doi.org/10.1038/s41587-025-02628-6

(2025) doi:10.1038/s41587-025-02628-6.

28. Katrekar, D. *et al.* In vivo RNA editing of point mutations via RNA-guided adenosine

deaminases. *Nat Methods* **16**, 239–242 (2019).

29. Wettengel, J., Reautschnig, P., Geisler, S., Kahle, P. J. & Stafforst, T. Harnessing human

ADAR2 for RNA repair - Recoding a PINK1 mutation rescues mitophagy. *Nucleic acids*

research **45**, 2797–2808 (2017).

30. Liu, Z. *et al.* Systematic comparison of 2A peptides for cloning multi-genes in a

polycistronic vector. *Sci Rep* **7**, 2193 (2017).

31. Montiel-González, M. F., Vallecillo-Viejo, I. C. & Rosenthal, J. J. C. An efficient system for

selectively altering genetic information within mRNAs. *Nucleic Acids Res* **44**, (2016).

32. Uzonyi, A. *et al.* Deciphering the principles of the RNA editing code via large-scale

systematic probing. *Mol Cell* **81**, 2374-2387.e3 (2021).

33. Kawakami, Y. *et al.* Cloning of the gene coding for a shared human melanoma antigen

recognized by autologous T cells infiltrating into tumor. *Proceedings of the National*

Academy of Sciences **91**, 3515–3519 (1994).

34. Coulie, P. G. *et al.* A new gene coding for a differentiation antigen recognized by autologous

cytolytic T lymphocytes on HLA-A2 melanomas. *Journal of Experimental Medicine* **180**,

35–42 (1994).

35. Johnson, L. A. *et al.* Gene transfer of tumor-reactive TCR confers both high avidity and

tumor reactivity to nonreactive peripheral blood mononuclear cells and tumor-infiltrating

lymphocytes. *J Immunol* **177**, 6548–6559 (2006).

36. Valmori, D. *et al.* Enhanced generation of specific tumor-reactive CTL in vitro by selected

Melan-A/MART-1 immunodominant peptide analogues. *J Immunol* **160**, 1750–1758 (1998).

37. Grossman, R. L. *et al.* Toward a Shared Vision for Cancer Genomic Data. *N Engl J Med* **375**,

1109–1112 (2016).

38. Reynisson, B., Alvarez, B., Paul, S., Peters, B. & Nielsen, M. NetMHCpan-4.1 and

NetMHCIIpan-4.0: improved predictions of MHC antigen presentation by concurrent motif

deconvolution and integration of MS MHC eluted ligand data. *Nucleic Acids Res* **48**, W449–

W454 (2020).

39. Gartner, J. J. *et al.* A machine learning model for ranking candidate HLA class I neoantigens

based on known neoepitopes from multiple human tumor types. *Nat Cancer* **2**, 563–574

(2021).

40. Garrido, F. & Algarra, I. MHC antigens and tumor escape from immune surveillance. *Adv*

Cancer Res **83**, 117–158 (2001).

41. Rojas, L. A. *et al.* Personalized RNA neoantigen vaccines stimulate T cells in pancreatic

cancer. *Nature* **618**, 144–150 (2023).

42. Kluesner, M. *et al.* MultiEditR: The first tool for detection and quantification of multiple

RNA editing sites from Sanger sequencing demonstrates comparable fidelity to RNA-seq.

Molecular Therapy - Nucleic Acids https://doi.org/10.1016/j.omtn.2021.07.008 (2021)

doi:10.1016/j.omtn.2021.07.008.

43. Arakawa, H., Lodygin, D. & Buerstedde, J. M. Mutant loxP vectors for selectable marker

recycle and conditional knock-outs. *BMC Biotechnol* **1**, 7 (2001).

44. Cui, Z., Geurts, A. M., Liu, G., Kaufman, C. D. & Hackett, P. B. Structure–Function

Analysis of the Inverted Terminal Repeats of the *Sleeping Beauty* Transposon. *Journal of*

Molecular Biology **318**, 1221–1235 (2002).

45. Chan, A. S. L. *et al.* Titration of RAS alters senescent state and influences tumour initiation.

Nature **633**, 678–685 (2024).

46. Doman, J. L., Raguram, A., Newby, G. A. & Liu, D. R. Evaluation and minimization of

Cas9-independent off-target DNA editing by cytosine base editors. *Nat Biotechnol* **38**, 620–

628 (2020).

47. Walton, R. T., Christie, K. A., Whittaker, M. N. & Kleinstiver, B. P. Unconstrained genome

targeting with near-PAMless engineered CRISPR-Cas9 variants. *Science* **368**, 290–296

(2020).

48. Fu, Y. *et al.* High-frequency off-target mutagenesis induced by CRISPR-Cas nucleases in

human cells. *Nat Biotechnol* **31**, 822–826 (2013).

49. Ye, J. *et al.* Primer-BLAST: A tool to design target-specific primers for polymerase chain

reaction. *BMC Bioinformatics* **13**, (2012).

50. Jullien, N. AmplifX.

51. Davis, M. W. & Jorgensen, E. M. ApE, A Plasmid Editor: A Freely Available DNA

Manipulation and Visualization Program. *Front Bioinform* **2**, 818619 (2022).

52. Cetin, M. *et al.* T-FINDER: A highly sensitive, pan-HLA platform for functional T cell

receptor and ligand discovery. *Science Advances* **10**, eadk3060 (2024).

53. Castro, F. *et al.* High-throughput SNP-based authentication of human cell lines. *Int J Cancer*

132, 308–314 (2013).

54. Schmitt, M. & Pawlita, M. High-throughput detection and multiplex identification of cell

contaminations. *Nucleic Acids Res* **37**, e119 (2009).

55. Clement, K. *et al.* CRISPResso2 provides accurate and rapid genome editing sequence

analysis. *Nat Biotechnol* **37**, 224–226 (2019).

56. Peper, J. K. *et al.* An impedance-based cytotoxicity assay for real-time and label-free

assessment of T-cell-mediated killing of adherent cells. *Journal of Immunological Methods*

405, 192–198 (2014).

Figure legends

Figure 1. Short Precise Encodable ADAR Recruiting (SPEAR) gRNAs enable efficient and precise
site-directed RNA editing. (**A**) Schematic of the dual reporter cell line and the experimental
setup used to screen different genetically encoded gRNAs. (**B**) Diagram of the eGFP W58X
target mRNA and the corresponding targeting gRNA used in the experiment shown in panel **C**.
(**C**) Flow cytometry analysis of mCherry and eGFP double-positive cells, presented as a bar plot
for the tested gRNAs. (**D**) Schematic of the sequencing window where A-to-I edits were
assessed, as shown in panel **E**. (**E**) Heatmap of A-to-I editing from NGS amplicon sequencing,
showing on-target editing (in blue) and off-target editing (in red). The region targeted by each
gRNA is indicated with a red rectangle. For panel C, bars represent the mean for three biological
replicates; error bars indicate the standard deviation.

Figure 2. SPEAR gRNAs can generate functional epitopes and activate specific T cells. (**A**)
Schematic of the proof-of-principle experiment: HEK293T cells expressing either wild-type (WT)
or G31S mutant MART-1 are co-cultured with DMF5-expressing T cells, followed by assessment
of T cell activation via flow cytometry. SPEAR gRNAs are used to restore the original MART-1
coding sequence at the RNA level, thereby rescuing antigen-specific T cell activation. (**B**)
Representative Sanger sequencing traces at the G31S codon from both genomic DNA and cDNA

of HEK293T MART-1 G31S cells expressing the SPEAR gRNA. (**C**) Representative flow cytometry

dot plots showing CD69 expression as a marker of T cell activation under the three

experimental conditions. (**D**) Bar plot showing average T cell activation, measured as the

percentage of CD69$^+$ CD2$^+$ double-positive cells, across three biological replicates for each

condition (adjusted p-values: **** < 0.0001, one-way ANOVA with multiple comparisons). Bars

represent the mean for three biological replicates; error bars indicate the standard deviation.

Figure 3. SPEAR gRNA editing induces T cell activation and tumor cell elimination *in vivo*. (A)

Bar plot showing average A-to-I editing percentages at the MART-1 G31S site, quantified from

Sanger sequencing traces using MultiEditR[42], in 624-mel MART-1^{G31S} cell lines expressing either

a non-targeting (NT) or targeting (T) gRNA (p-value: **** < 0.0001, two-tailed unpaired t-test).

(**B**) Representative Sanger sequencing traces at the G31S codon from both genomic DNA and

cDNA of 624-mel MART-1^{G31S} cells expressing NT- or T-gRNAs. Bars represent the mean for

three biological replicates; error bars indicate the standard deviation. (**C**) xCELLigence real-time

analysis of the cytotoxicity of MART-1-specific human T cells co-cultured with 624-mel cells

expressing MART-1, MART-1^{G31S}, or mutant MART-1^{G31S} with NT- or T-gRNAs. For MART-1 or

MART-1^{G31S} cells, each curve represents the mean of three technical replicates. For cells

expressing NT- or T-gRNAs, each curve represents the mean from three biological replicates,

each of those with three technical replicates. T cells were added after 24 hours; the cell index is

normalized to the time point of T cell addition. (**D**) Schematic of the *in vivo* experimental setup.

(**E**) *In vivo* tumor growth analysis of MART-1^{G31S} cells expressing the T- or NT-gRNA with or

without adoptive cell therapy (ACT) using MART-1-specific human-derived T cells. Each curve

represents the mean of 6 biological replicates. Ordinary one-way ANOVA was applied: * (P

value: 0.0493). (**F**) Kaplan-Meier survival analysis curves of mice injected with MART-1^{G31S} cells

expressing the T- or NT-gRNA with or without adoptive cell therapy (ACT) using MART-1-specific

human-derived T cells. n = 6, Logrank (Mantel-Cox) test: ** (P value: 0.0055).

Figure 4. Neoantigens that can be generated next to prevalent hot-spot somatic mutations

A heatmap of the 50 most common somatic mutations across all cancer types and the 50 most

prevalent HLA types, illustrating all possible combinations that can generate novel potent

neoantigens. Only rows and columns in which a result was detected are shown.

Materials and Methods

Plasmids

pcDNA3.1(-) mCherry-P2A-eGFP^{W58X} puro: This plasmid was generated by inserting a

puromycin resistance cassette into the pcDNA3.1(-) mCherry-2A-FLAG-eGFP^{W58X} backbone,

which was linearized with BglII. The parental plasmid was a kind gift from Dr. Joshua Rosenthal

(University of Chicago) and previously described by REF[31]. The puromycin selection cassette was

excised from plasmid pLoxPuro[43] via BamHI digestion and ligated into the linearized vector

using the Rapid DNA Ligation kit (Merck, Cat# 11635379001).

Plasmids encoding ADAR1-recruiting gRNAs for eGFP^{W58X}: Guide RNA-encoding DNA fragments

were generated by inserting single-stranded DNA (ssDNA) oligonucleotides into a pENTR-U6

backbone (a kind gift of Dr. Joshua Rosenthal, University of Chicago), which was linearized via

PCR using Q5® High-Fidelity DNA Polymerase (NEB, Cat# M0491) and primers #1 and #2 (see

Supplementary Table 1). One or two ssDNA oligos or a dsDNA fragment were inserted into the

PCR-amplified vector using NEBuilder® HiFi DNA Assembly Master Mix (NEB, Cat# E2621). All

ssDNA oligonucleotides were ordered by Sigma-Aldrich, and the dsDNA fragments by Integrated

DNA Technologies (IDT). Nucleotide sequences are provided below, with NEBuilder-compatible

overlapping regions shown in lowercase. All plasmids were verified by Sanger sequencing.

Detailed plasmid maps are available in the Supplementary Information as text files.

The following gRNA-expressing plasmids were generated for eGFP^{W58X} (**Figure S1B and 1C**):

• **pENTR-U6 GluRv1-16** (eGFP^{W58X}) – inserted ssDNA oligo: tatatcttgtggaaaggacgaaacacc

GGTGAATAGTATAACAATATGCTAAATGTTGTTATAGTATCCACCGTTGGCCATGGAACAGttttt

ctagacccagctttcttgtac

• **pENTR-U6 GluRv4-16** (eGFP^{W58X}) – inserted ssDNA oligo: tatatcttgtggaaaggacgaaacacc

GGTGAAGAGGAGAACAATATGCTAAATGTTGTTCTCGTCTCCACCGTTGGCCATGGAACAGttttt

tctagacccagctttcttgtac

• **pENTR-U6 GluRv9.4-16** (eGFP^{W58X}) – inserted ssDNA oligo: tcttgtggaaaggacgaaacacc

GGTGTCGAGAAGAGGAGAACAATATGCTAAATGTTGTTCTCGTCTCCTCGACACCGTTGGCCAT

GGAACAGttttttctagacccagctttctt

• **pENTR-U6 GluRv9.4-38** (eGFP^{W58X}) – inserted ssDNA oligos: (1) tcttgtggaaaggacgaaacacc

GGTGTCGAGAAGAGGAGAACAATATGCTAAATGTTGTTCTCGTCTCCTCGACACCGTATGTCAG

GGTAGTGACAA; (2) acaagaaagctgggtctagaaaaaaCTGTTCCATGGCCAACACTTGTCACTACCC

TGACATACGGTGTCGAGGAGACGAGAACAACAT

• **pENTR-U6 LEAPER-70** (eGFP^{W58X}) – inserted ssDNA oligo: tcttgtggaaaggacgaaacaccGGAT

TGCACGCCGTATGTCAGGGTAGTGACAAGTGTTGGCCATGGAACAGGTAGTTTTCCAGTAGTG

AAATttttttctagacccagctttctt

• **pENTR-U6 LEAPER-151 (Qu et al 2019)** (eGFP^{W58X}) – inserted ssDNA oligos: (1)

tatcttgtgga

aaggacgaaacaccGGTCGTGCTGTTTCATGTGGTCGGGGTACCTGCTGAAGCATTGCACGCCGTA

TGTCAGGGTAGTGACAAGTGTTGGCCATGGAACAGGTAGTTTT; (2) acaagaaagctgggtctagaaa

aaaAAGGAGAGGGCGATGCCACCTATGGCAAGCTGACCCTGAAGTTTATTTGCACTACTGGAAA

ACTACCTGTTCCATGGCCAACACTTGTCACTACC

• **pENTR-U6 LEAPER-151 (38-4ntMM-33-C-75; Uzonyi et al 2021)** (eGFP^{W58X}) – inserted

ssDNA oligos: (1) tatcttgtggaaaggacgaaacaccGGTCGTGCTGTTTCATGTGGTCGGGGTACCTG

CTGAAGCAAACGACGCCGTATGTCAGGGTAGTGACAAGTGTTGGCCATGGAACAGGTAGTTTT;

(2) acaagaaagctgggtctagaaaaaaAAGGAGAGGGCGATGCCACCTATGGCAAGCTGACCCTGAA

GTTTATTTGCACTACTGGAAAACTACCTGTTCCATGGCCAACACTTGTCACTACC

• **pENTR-U6 38-4ntMM-33-C-31 (sdl 107)** (eGFP^{W58X}) – inserted ssDNA oligos: (1)

tatcttgtgg

aaaggacgaaacaccGGTCGTGCTGTTTCATGTGGTCGGGGTACCTGCTGAAGCAAACGACGCCGT

ATGTCAGGGTAGTGACAAGTGTTGGCCATGGAACAGGTAGTTTT;   (2)

acaagaaagctgggtctagaa

aaaaATTTGCACTACTGGAAAACTACCTGTTCCATGGCCAACACTTGTCACTACC

- **pENTR-U6 38-4ntMM-33-C-16 (sdl 92)** (eGFP^{W58X}) – inserted ssDNA oligos: (1)
  tatcttgtgga

  aaggacgaaacaccGGTCGTGCTGTTTCATGTGGTCGGGGTACCTGCTGAAGCAAACGACGCCGTA

  TGTCAGGGTAGTGACAAGTGTTGGCCATGGAACAGGTAGTTTT; (2) acaagaaagctgggtctagaaa

  aaaAAACTACCTGTTCCATGGCCAACACTTGTCACTACC

- **pENTR-U6 20-4ntMM-33-C-75 (sdl 133)** (eGFP^{W58X}) – inserted ssDNA oligos: (1)
  tatcttgtgg

  aaaggacgaaacaccGGTCGGGGTACCTGCTGAAGCAAACGACGCCGTATGTCAGGGTAGTGACA

  AGTGTTGGCCATGGAACAGGTAGTTTT; (2) acaagaaagctgggtctagaaaaaaAAGGAGAGGGCG

  ATGCCACCTATGGCAAGCTGACCCTGAAGTTTATTTGCACTACTGGAAAACTACCTGTTCCATGG

  CCAACACTTGTCACTACC

- **pENTR-U6 10-4ntMM-33-C-75 (sdl 133)** (eGFP^{W58X}) – inserted ssDNA oligos: (1)
  tatcttgtgg

  aaaggacgaaacaccGGTGCTGAAGCAAACGACGCCGTATGTCAGGGTAGTGACAAGTGTTGGCC

  ATGGAACAGGTAGTTTT; (2) acaagaaagctgggtctagaaaaaaAAGGAGAGGGCGATGCCACCTAT

  GGCAAGCTGACCCTGAAGTTTATTTGCACTACTGGAAAACTACCTGTTCCATGGCCAACACTTGT

  CACTACC

- **pENTR-U6 20-4ntMM-33-C-16 (sdl 74, SPEAR)** (eGFP^{W58X}) – inserted ssDNA oligo:
  cttgtgga

  aaggacgaaacaccGGTCGGGGTACCTGCTGAAGCAAACGACGCCGTATGTCAGGGTAGTGACAA

  GTGTTGGCCATGGAACAGGTAGTTTtttttttctagacccagctttct

Plasmids for *in vitro* T cell activation assay:

The plasmids used for the proof-of-concept T cell activation assay (**Figure 2, S2**) contain a dual-

expression cassette structure: a hU6-driven gRNA expression cassette, followed by a hPGK1-

driven cassette encoding a codon optimized MART-1 (coMART-1), eBFP, and the blasticidin

resistance gene (BSD), separated by two different self-cleaving 2A peptides[30] (**Figure S2B**). This

design enables both antibiotic selection and monitoring of MART-1 expression via eBFP

fluorescence. We began by ordering a custom plasmid backbone from VectorBuilder containing

the hU6-gRNA and hPGK1-BSD expression cassettes (VectorBuilder ID: VB210525-1302qyy; this

ID can be used to retrieve detailed information about the vector on vectorbuilder.com). The

vector was linearized using SpeI-HF (NEB, Cat# R3133) and BsrGI-HF (NEB, Cat# R3575). The

individual components—hU6-gRNA cassette, hPGK-MART-1^{WT}- or MART-1^{G31S}-T2A, and eBFP-

P2A-BSD—were amplified separately by PCR using Q5® High-Fidelity DNA Polymerase (NEB,

Cat# M0491) from existing plasmids, and assembled into the linearized backbone using

NEBuilder® HiFi DNA Assembly Master Mix (NEB, Cat# E2621). The resulting plasmids express

either a non-targeting (NT) or targeting (T) SPEAR gRNA as follows:

- **NT-SPEAR gRNA** (control): TCGGGGTACCTGCTGAAGCAAACGACGCCGTATGTCAGGGTAGT
GACAAGTGTTGGCCATGGAACAGGTAGTTT

- **T-SPEAR gRNA** (coMART-1^{G31S}): TCCGGCAGTACCAGCAGCCGTAGTGCAGCAGCACGCCTAG
GATCACTGTCAGGAGGCCGATCCCAGCAAGCTCC

Plasmids for *in vivo* T cell activation and tumor elimination assay:

The plasmids used in the *in vivo* experiments (**Figure 3, S3**) were also designed with a dual-

expression cassette structure comprising a hU6-driven gRNA expression cassette, followed by

an hPGK1-driven cassette encoding eBFP and the blasticidin resistance gene (BSD), separated

by a self-cleaving 2A peptide (**Figure S3B**). This configuration enables antibiotic selection and

allows monitoring of plasmid integration through eBFP fluorescence. To enable stable genomic

integration, the dual-expression cassette was cloned into a pT vector containing the necessary

inverted terminal repeats (ITRs) for Sleeping Beauty transposon system-mediated integration[44].

The vector was linearized using NsiI-HF (NEB, Cat# R3127) and XhoI (NEB, Cat# R0146). The

individual elements—hU6-gRNA cassette and hPGK-eBFP-P2A-BSD—were amplified separately

by PCR using Q5® High-Fidelity DNA Polymerase (NEB, Cat# M0491) from existing plasmids, and

assembled into the linearized backbone using NEBuilder® HiFi DNA Assembly Master Mix (NEB,

Cat# E2621). The resulting plasmids express either a non-targeting (NT) or targeting (T) SPEAR

gRNA as follows:

- **NT-SPEAR gRNA** (control): TCGGGGTACCTGCTGAAGCAAACGACGCCGTATGTCAGGGTAGT
  GACAAGTGTTGGCCATGGAACAGGTAGTTT
- **T-SPEAR gRNA** (MART-1[G31S]): TTCTACAATACCAACAGCCGTACTGCAGTAAGACTCCCAGGAT
  CACTGTCAGGATGCCGATCCCAGCGGCCTCT

687 For stable integration, these plasmids were co-transfected with pPGK-SB13, a Sleeping Beauty

688 transposase expression plasmid driven by a PGK1 promoter. This plasmid was a gift from

689 Masashi Narita (Addgene plasmid #236078; http://n2t.net/addgene:236078; RRID:

690 Addgene_236078)[45].

Plasmids for genome base editing and single guide RNA (sgRNA) expression:

The plasmid AALN-BE4max was a gift from David Liu (Addgene plasmid #138161;

http://n2t.net/addgene:138161; RRID: Addgene_138161) and encodes a modified cytosine

base editor consisting of rAPOBEC1(AALN)-nSpCas9-UGI-UGI, in which APOBEC1 has been

engineered to reduce off-target activity[46]. To facilitate the identification of transfected cells, we

inserted an eGFP reporter into this plasmid. Specifically, plasmid AALN-BE4max was digested

with AgeI-HF (NEB, Cat# R3552) to linearize it between the UGI-SV40_NLS and 6xHis tag

regions. The insert, encoding P2A-EGFP-SV40_NLS-6xHis, was PCR-amplified from pCMV-T7-

ABEmax(7.10)-SpCas9-NG-P2A-EGFP (Addgene plasmid #140005; gift from Benjamin Kleinstiver;

http://n2t.net/addgene:140005; RRID: Addgene_140005)[47] using Q5® High-Fidelity DNA

Polymerase (NEB, Cat# M0491) and primers #3 and #4 (see Supplementary Table 1). The insert

was assembled into the linearized backbone using NEBuilder® HiFi DNA Assembly Master Mix

(NEB, Cat# E5520).

For sgRNA expression, we used the vector pFYF1320 EGFP Site#1, a gift from Keith Joung

(Addgene plasmid #47511; http://n2t.net/addgene:47511; RRID: Addgene_47511)[48], which

originally encoded a sgRNA targeting EGFP. This sequence was replaced with a sgRNA targeting

the *MART-1* gene as follows. The backbone was amplified using Q5® PCR to exclude the EGFP-

targeting region using primers #5 and #6 (see Supplementary Table 1). A ssDNA oligo containing

the   protospacer   for   the   *MART-1*-targeting   sgRNA   (atatatcttgtggaaag

gacgaaacaccggGATGCCGATCCCAGCGGCCCgttttagagctagaaatagcaagtt) was assembled into the

linearized backbone using NEBuilder® HiFi DNA Assembly Master Mix (NEB, Cat# E5520).

DNA oligonucleotides

All primers used in this study can be found in (Supplementary Table S1) and were designed

using Primer-BLAST[49], AmplifX 2.1.1[50], or ApE[51].

Cell culture and cell lines generation

HEK293T cells (ATCC, Cat# CRL-3216, RRID: CVCL_0063) were cultured at 37°C in a humidified

incubator with 5% CO_2 in high-glucose DMEM (Sigma-Aldrich, Cat# D6429), supplemented with

5% FBS (PAN Biotech, Cat# P40-37100) and 1% penicillin/streptomycin (Sigma-Aldrich,

Cat#P4333).

The stable dual-reporter cell line expressing mCherry-P2A-eGFP W58X-puro was generated by

transfecting the plasmid into HEK293T cells using Lipofectamine 2000 (ThermoFisher, Cat#

11668019) according to the manufacturer's protocol. 48 h after transfection, cells were plated

by limiting dilution into 96-well plates in medium supplemented with puromycin (1.5 µg/ml).

Clonal populations were identified by microscopic inspection, and single clones were screened

via flow cytometry analysis to select a single clone stably expressing the dual fluorescence

reporter plasmid with no detectable EGFP background.

HEK293T cells stably expressing the constructs for the *in vitro* T cell activation assay (**Figure 2,**

S2), which included hU6-gRNA (T- or NT-) and hPGK1-coMART-1$^{\text{WT or G31S}}$-2A-eBFP-2A-BSD were

generated similarly. Cells were transfected using Lipofectamine 2000, expanded for 48 hours,

and then selected with blasticidin (15 µg/ml). As soon as non-transfected cells died, BFP⁺ cells

were sorted by flow cytometry. The expression of MART-1$^{WT \text{ or } G31S}$ and the presence of SPEAR-

mediated editing were confirmed by RT-PCR and Sanger sequencing.

Jurkat cells expressing the DMF5 TCR were cultured at 37°C and 5% CO_2 in RPMI 1640 (Sigma-

Aldrich, Cat# R8758) supplemented with 10% FBS (PAN Biotech, Cat# P40-37100), 2 mM L-

glutamine (Gibco, Cat# 25030-024) and 1X penicillin/streptomycin (Sigma-Aldrich, Cat#P4333).

Jurkat-DMF5 cells were generated as previously described[52] using a lentivirally-encoded, single-

ORF TCR sequence separated by a furin-T2A cleavage sequence, followed by 12 days of

puromycin selection for the transgenic TCR.

624-mel cells (RRID:CVCL_8054) were cultured at 37°C and 5% CO_2 in high-glucose DMEM

(Sigma-Aldrich, Cat# D6429) supplemented with 5% FBS (PAN Biotech, Cat# P40-37100) and 1%

penicillin/streptomycin (Sigma-Aldrich, Cat#P4333).

The 624-mel MART-1^{G31S} cell line was generated via DNA base editing (see below). For *in vivo*

experiments (**Figure 3, S3**), 624-mel MART-1^{G31S} cells were transfected with plasmids expressing

either a T- or NT-gRNA together with pPGK-SB13 (Sleeping Beauty transposase) in a 2:1 ratio,

using Lipofectamine 2000 (ThermoFisher, Cat# 11668019). After 48 hours, cells were expanded,

and selected with blasticidin (5 μg/ml). Once selection was complete, BFP$^+$ cells were sorted by

flow cytometry. Successful expression of MART-1^{G31S} and gRNA-dependent RNA editing were

confirmed by RT-PCR and Sanger sequencing.

All cell lines were authenticated using Multiplex Cell Authentication by Multiplexion

(Heidelberg, Germany) as described recently[53]. Additionally, the purity of cell lines was

validated using the Multiplex cell Contamination Test by Multiplexion (Heidelberg, Germany) as

described recently[54]. No Mycoplasma, SMRV or interspecies contamination was detected.

Genome base editing with a cytosine base editor (CBE)

10^5 624-mel cells were seeded in 500 μl complete growth medium (DMEM, FBS 10%, P/S 1%) in

a flat-bottom 24-well plate. After 18-24 hours, the complete growth medium was replaced with

250 μl antibiotic-free reduced serum medium (OptiMEM, gibco, Cat #31985062). Cells were

transfected with a mixture of 2 µl lipofectamine 2000, 500 ng sgRNA vector and 1 µg base

editor vector diluted in 100 µl OptiMEM following the manufacturer's instructions. After 6

hours, the transfection medium was replaced with 500 µl complete growth medium. After 72

hours, the medium was removed, and cells were washed with 250 µl PBS. To detach the cells,

100 µl trypsin were added, and cells were incubated for 1 minute at 37°C. Trypsin was

inactivated by adding 400 µl pre-warmed complete growth medium. Detached cells were

filtered and transferred to a 5 ml round bottom polystyrene test tube with cell strainer snap

cap (Corning, Cat #352235). GFP+ cells were single-cell sorted into flat-bottom 96-well plates.

Single-cell clones were incubated for at least 2 weeks at 37°C until they were ready to be

screened. When ready, medium was removed, and clones were washed once with 100 µl PBS.

Cells were detached by adding 20 µl trypsin and resuspended in 200 µl complete growth

medium. 100 µl of each clone were transferred into a new flat-bottom 96-well plate already

prepared with 100 µl medium per well to keep and further expand the selected clones. The

remaining 100 µl were transferred into another flat-bottom 96-well plate already prepared with

100 µl medium per well for screening the next day. DNA was extracted with QuickExtract™ DNA

Extraction Kit (Lucigen, Cat #QE0905T). Briefly, medium was removed from the screening plate

and cells were washed once with 100 µl PBS. 100 µl Quick-Extract DNA solution were added,

cells were detached by thoroughly pipetting up and down and transferred into a 96-well PCR

plate (or PCR strips). Cells were incubated at 65°C for 15 min in a thermal cycler, vortexed for

15 sec, and incubated at 98°C for 5 min a thermal cycler. The extracted DNA was used for PCR

and   then   stored   at   -80°C.

PCR was performed on the extracted DNA with Q5 High-Fidelity DNA Polymerase from NEB

(#M0491S) according to the manufacturer's instructions and the following conditions. 5-20 ng

of DNA (2.5 µl) were added to 31 µl of RNase-free water, 10 µl of Q5 Buffer, 1 µl of dNTP mix

(10 mM each), 2.5 µl of each primer (10 µM) and 0.5 µl of Q5 polymerase in a final reaction

volume of 50 µl. *MART-1* amplification was obtained using primers #7 and #8 (Supplementary

Table 1). The PCR program was run as follows: denaturation at 98°C for 30 sec, followed by 35

cycles of denaturation at 98°C for 10 sec, annealing at 65°C for 15 sec and elongation at 72°C

for 10 sec. In the end, a final elongation step at 72°C was run for 2 min. The correct size of the

PCR products (245 bp) was checked on a 1.5% agarose gel. 5 µl of each product diluted with 10

µl nuclease-free water was sent for Sanger sequencing to Eurofins Genomics Germany GmbH.

Co-culture and related flow cytometry analysis

In the co-culture experiment using HEK293T overexpressing either MART-1 or MART-1^{G31S}

(**Figure 2, S2**), we used the following antibodies: PE anti-human CD69 (BioLegend, Cat# 310906)

and APC anti-human CD2 (BioLegend, Cat# 300214). We additionally assessed viability of cells

using 7AAD staining (Invitrogen, Cat# A1310). 2.5 x 10^4 HEK293T stably expressing either MART-

1 or MART-1^{G31S} and the SPEAR gRNAs (targeting or not-targeting MART-1^{G31S}) were seeded in a

96-wells flat bottom plate in 40 µl of supplemented DMEM. 24 h post seeding, 1 x 10^5 DMF5$^+$

Jurkat cells in 35 µl of supplemented RPMI were added to the same wells for a total volume of

75 µl per well. The cells were maintained in co-culture for 16 h at 37°C and 5% CO_2. Samples

were washed twice in FACS buffer (2 mM EDTA and 2% FBS in PBS) and cells were stained using

50 µl of FACS buffer containing the antibodies diluted 1:200. After 20 minutes of incubation at

4°C, samples were washed twice and resuspended in a final volume of 200 µl of FACS buffer.

Flow cytometer acquisition was done using a BD Canto II (BD Biosciences), and data were

analyzed with FlowJo.

Adenosine-to-inosine (A-to-I) editing sites quantification

For A-to-I editing site quantification, RNA was extracted using the RNeasy Mini Kit (Qiagen, Cat#

74106) and treated with DNase (Invitrogen, Cat# AM1907). One-step RT-PCRs were performed

using the One-step RT-PCR kit (Qiagen, Cat# 210212) using primers #9-10 (for *coMART-1*), #11-

12 (for endogenous *MART-1*) or #13-14 for eGFP^{W58X}. Additionally, when needed the same PCR

was performed on genomic DNA (extracted using the All Prep DNA/RNA Kit, Qiagen, Cat#

80204) with Q5 High-Fidelity DNA Polymerase (NEB, Cat# M0491) as a no-editing control. All

the PCR products were purified (Macherey-Nagel, Cat# 740609) and analyzed by Sanger

sequencing. Quantification of editing was performed directly from the Sanger traces using

MultiEditR[42].

On-target and bystander off-target editing analysis via amplicon-seq

To evaluate in detail on-target and bystander off-target editing on the eGFP W58X reporter, a

second PCR with primers containing adaptor sequences (#15-16, Supplementary Table 1) for

MiSeq amplicon sequencing (Eurofins Genomics, GATC services, Germany) was performed using

Q5® High-Fidelity DNA Polymerase (NEB, Cat# M0491). Before performing the second PCR, the

amplicons from three independent replicates were mixed in a 1:1:1 ratio. Analysis of the

amplicon-seq data was performed using CRISPResso2[55].

xCELLigence cytotoxicity assay

MART-1 ELAGIGILTV/A2 TCR (DMF5) was transduced into human PBMC. DMF5 positive cells

were expanded in the presence of PBMCs feeder cell loaded with MART-1 peptide (ProImmune)

and IL2 (70 IU/ml) for 14 days. >90% of T cells were DMF5 positive after the procedure. Cell-

mediated cytotoxicity was analyzed in an impedance-based label-free cytotoxicity assay

(xCELLigence) system[56]. 2×10^4 624-mel cells (each cell line in 3 technical replicates; 3 biological

replicates of the MART-1^{G31S} cells expressing either the NT- or T-SPEAR gRNA were compared)

were seeded in 50 µl of complete growth medium (DMEM FBS 10% P/S 1%) in a 96-Well E-plate

(E-Plate 96 PET, Agilent). The attachment of the cells changes the electrical impedance of these

electrodes, which is monitored as an increase of the dimensionless "cell index". After 24 hours,

4×10^4 MART-1 specific T cells were added in 50 µl of complete growth medium in half of the

samples (condition "+ T cells"), while the other half only received 50 µl of fresh complete

growth medium. Lysis of the tumor cells by the T cells causes their detachment from the

electrodes, which changes the impedance of the wells, resulting in a decrease in the cell index.

The tumor cell growth was continuously monitored from the seeding of the tumor cells until 48

hours after addition of the T cells. The co-culture of tumor cells and T cells was incubated for 48

h at 37 °C 5% CO_2 and 95% humidity.

Mice

Immunodeficient NXG mice came from the Janvier Laboratory and were bred at the DKFZ

animal facility. 10- to 16-week-old mice were assigned to age-matched experimental groups.

Mice were housed under Specific and Opportunistic Pathogen Free (SOPF) conditions and a 12-

hour day/night cycle. Animal procedures were performed in accordance with all relevant ethical

regulations for animal testing and research, and were approved by the governmental

authorities (Regional Administrative Authority Karlsruhe, Germany; permit number 35-

9185.81/G-11/22).

***In vivo* tumor growth and Adoptive T cell Therapy (ATC)**

On day 1, 500.000 melanoma cells in 100 μl were mixed in a 1:1 volume with 100 μl Matrigel

Basement Membrane Matrix (Corning, Cat# 356234) and kept strictly on ice till injection in

mice. The cell-Matrigel mix was subcutaneously injected on the right flank of NXG mice. Each

cell line was injected into 12 mice. Tumor size was measured every other day starting on day 6.

On days 12 and 19, 5×10^6 MART-1 specific T cells in 50 μl PBS were administered

intratumorally to half of the mice. The other half received PBS as negative control. On day 23,

mice were sacrificed, tumors were isolated (if any) and analyzed via flow cytometry and ELISA

to search for markers of T cell activation.

Statistical analysis and data visualization

Data were analyzed and plotted using GraphPad Prism (version 10.5.0). Specific information

about data presentation is provided in each figure caption throughout the manuscript. Multiple

illustrations were created with BioRender.

Algorithm

Here we present the NeoADARgen – an algorithm that designs a patient-specific gRNA tailored

to the individual's unique tumor genetic characteristics. Owing to its precise design, using this

gRNA as the main component of an Endogenous-ADAR will induce A-to-I(G) edits at the tumor

RNA level upon delivery to the tissue. The gRNA interacts exclusively with tumor cells, as it is

designed to base-pair with a region containing somatic mutations that are absent in normal

cells. Our algorithm identifies the optimal points for editing, which in turn alter the translated

peptide within the cancer cell, creating a novel potent neoantigen that takes into account the

individual's HLA typing. We utilize a list of somatic mutations sequenced from the tumor and

the patient's HLA typing as basic inputs. For increased accuracy, incorporating tumor gene

expression levels is also feasible.

<u>Step 1: Ensuring binding to tumor cells while excluding normal cells.</u>

To generate neoantigens in tumor-cells but not in normal-cells, the initial step involves

targeting a somatic mutation within the cancerous cell. This mutation can manifest in any

possible variant type, as long as it modifies the RNA sequence in comparison to the wildtype.

The preliminary gRNA is programmed to bind by base pairing to a sequence comprising the

mutation along with the 20 nucleotides upstream and the 20 nucleotides downstream, totaling

in a sequence of ~40 nucleotides. Due to the presence of the somatic mutation at the core of

the sequence, the gRNA binds to the cancerous cells but not to the normal cells upon

approaching the tissue, as normal cells typically lack the tumor somatic variants. This step

guarantees binding specificity exclusively to the tumor cells (**Figure S4**).

<u>Step2: Identifying the optimal adenosines in the region for editing.</u>

The following step involves scanning the gRNA sequence for adenosines which upon

conversion to guanosines will modify the peptide in a manner predicted to generate a potent

neoantigen. For this purpose, we evaluate each adenosine within the sequence to determine

the effect of editing it to guanosine on the resulting protein. Given the natural ADAR's ability to

edit multiple adenosines in clusters, we also explore the impact of editing two adenosines at a

time on the resulting protein. Note that adenosines preceded by guanosines are not

considered, as they do not meet the criteria for the ADAR motif, which requires the absence of

a guanosine 5' to the edited adenosine. (While this constraint can be overcome by manipulating

the gRNA to include an additional mismatch 5' to the edited adenosine, we prefer not to do so

in order to maintain ideal complementarity to the cancerous sequence). Additionally, the

examination excludes the first and last nucleotides of the sequence.

To identify the optimal alteration capable of generating a neoantigen, we analyzed all

possible 9-mer sub-sequences derived from the gRNA sequence, where one or two adenosines

are substituted with guanosines (**Figure S5**). To assess the effect on the resultant peptide, we

employed the external tool NetMHCpan4.1[38] which utilizes artificial neural networks to predict

peptide binding for a given sequence by a given HLA. We submitted each combination of

altered sub-sequences along with each HLA class I type of the patient to the prediction tool,

examining the output parameters for each combination. By default, thresholds of %Rank < 0.5%

and %Rank < 2% are used to identify strong MHC binders (SBs) and weak MHC binders (WBs)

for class I, respectively[38]. In our analysis, we used even more conservative parameters: %Rank <

0.1% and affinity < 50 as indicators for SB. Additionally, if gene expression data were available,

we set a threshold of TPM above 15 as an additional mandatory criterion for robustness. This

ensures adequate transcription of the gene for RNA editors to function effectively. If more than

one combination met these criteria, we prioritized the sub-sequence with the lowest detected

rank.

<u>Step3: Presenting the best results</u>

At the end, the algorithm presents all potential gRNA sequences, marking the optimal

adenosines to edit with an asterisk (*). These gRNAs are the optimal sequences tailored for a

given tumor sample, to be utilized with Endogenous-ADAR base editors. Upon delivery to the

tissue, they will bind to a region encompassing a somatic mutation expressed in this tumor to

ensure cancer cell specificity and induce single or double A-to-I editing leading to precise

peptide alterations. These alterations are predicted to generate neoantigens according to the

patient's HLA typing.

Cancer databases

The computational results presented are based on cancer data extracted from the Tumor

Cancer Genome Atlas (TCGA). To apply our algorithm to a variety of cancer samples, we utilized

the Genomic Data Commons (GDC) data portal [https://portal.gdc.cancer.gov/]. We

downloaded the data for seven cancer TCGA projects: BRCA, SKCM, OV, GBM, LIHC, LUAD, and

LUSC. The first five were downloaded in September 2023 and the last two in January 2024. For

each patient, we used the DNA sample as sequenced from the primary tumor or from the

metastatic tumor if the primary sample was not available. In case the DNA file was not

available, we used the whole genome sequence file (Supplementary Table 2). For each patient,

we also used the gene expression quantification file as sequenced from its tumor. Lastly, we

used The Cancer Immunome Atlas (TCIA) data portal [https://tcia.at/home] to extract the HLA

typing of each individual as reported by its unique TCGA ID. Only individuals with access to all

data were included in our analysis.

To apply our algorithm to the most common cancer hotspots, we extracted a list of the

most common somatic mutations and their reported prevalence across all cancer types as

presented in GDC. We also downloaded all TCGA samples available in TCIA to aggregate the

most common HLA types among them. Both lists were downloaded in September 2023. For this

analysis gene expression could not be considered by the algorithm, as the calculation was not

done on an individual basis.

Figure 1

A

CMV — mCherry — P2A — eGFP — genomic DNA

W58X

↓ transcription

5' — A — 3' RNA

↓ transfection

plasmid-encoded gRNA

5' — A/C — 3'
ADAR1

↓ A-to-I editing

5' — I — 3'

B

target mRNA (eGFP W58X)
5'- |||||||||||| UAG ||||||||||| -3'
X58   4 nt mismatch

gRNA
3'- |||||||||||| ACC ||||||||||| -5'

3' arm ← 33 nt → 5' arm

specificity domain length (sdl)

C

sdl (nt)	5' arm (nt)	3' arm (nt)	reference
151	38	75	Uzonyi et al 2021
107	38	31	
92	38	16	
133	20	75	
123	10	75	
74	20	16	

mCherry+eGFP+ cells (%) 0 20 40 60

D

eGFP W58X mRNA

5'- ...GAA GGA GAG GGC GAU GCC ACC UAU GGC AAG CUG ACC CUG AAG UUU AUU UGC ACU ACU GGA AAA CUA CCU GUU CCA UAG CCA ACA CUU GUC ACU ACC CUG ACA UAC GGC GUG CAA

(positions) 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32

UGC UUC AGC AGG UAC CCC GAC CAC AUG AAA CAG CAC GAU...-3'

(positions) 33 34 35 36 37 38 39 40 41 42 43 44

E

A position

sdl (nt)	5' arm (nt)	3' arm (nt)
151	38	75
74	20	16

off-target editing (%) 2 4 6 8

on-target editing (%) 30 40 50

Figure 2

A

Mutagenesis → RNA editing

	MART-1	MART-1^G31S	MART-1^G31S + T-gRNA
DNA	GAGCTTGCTGGGATCGGCATCCTGACAGTG	GAGCTTGCTGGGATCAGCATCCTGACAGTG	GAGCTTGCTGGGATCAGCATCCTGACAGTG
mRNA	GAGCUUGCUGGGAUCGGCAUCCUGACAGUG	GAGCUUGCUGGGAUCAGCAUCCUGACAGUG	GAGCUUGCUGGGAUCIGCAUCCUGACAGUG
peptide	E L A G I G I L T V	E L A G I S I L T V	E L A G I G I L T V

B

DNA

cDNA — 18%

C

MART-1 + NT-gRNA   MART-1^G31S + NT-gRNA   MART-1^G31S + T-gRNA

CD69 (Activated T cells) vs CD2 (T cells)

D

CD2⁺CD69⁺ cells (%)

| MART1 | WT | G31S | G31S |
| gRNA | NT- | NT- | T- |

Figure 3

Figure 4