Bamboozle: A bioinformatic tool for identification and quantification of intraspecific barcodes

Matthew I M Pinder[†a]*, Björn Andersson[†a], Karin Rengefors[b], Hannah Blossom[b,c], Marie Svensson[b], and Mats Töpel[ad]

[†]These authors contributed equally to the manuscript

[a]Department of Marine Sciences, University of Gothenburg, Göteborg, Sweden
[b]Department of Biology, Lund University, Lund, Sweden
[c]Bigelow Laboratory for Ocean Sciences, Maine, USA
[d]IVL Swedish Environmental Research Institute, Göteborg, Sweden
*Corresponding author: mats.topel@ivl.se

Keywords: Metabarcoding, Population genomics, microbial evolution, microalgae, polymorphism,

Funding sources
This project was funded by the Swedish Research Council VR grant number 2018-04555 (MT), and the Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning (FORMAS) grant numbers 2017-00466 (MT) 2016-00594 (Anna Godhe), and the Oscar and Lili Lamm Foundation grant number FO2018-0042 (Helena L. Filipsson).

Competing Interest Statement: The authors declare no competing financial interests.

Abstract

Evolutionary changes in populations of microbes, such as microalgae, cannot be traced using

conventional metabarcoding loci as they lack intraspecific resolution. Consequently, selection

and competition processes amongst strains of the same species cannot be resolved without

elaborate isolation, culturing, and genotyping efforts. Bamboozle, a new bioinformatic tool

introduced here, scans a species' entire genome and identifies allele-rich barcodes that enable

direct identification of different strains from a common population, and a single DNA sample,

using amplicon sequencing. We demonstrate its usefulness by identifying hypervariable

barcoding loci (<500 bp) from genomic data in two microalgal species, the diploid diatom

Skeletonema marinoi, and the haploid chlorophyte *Chlamydomonas reinhardtii*. Across the

genomes, only 26 loci capable of resolving all available strains' genotypes were identified, all of

which are within protein-coding genes of variable metabolic function. Single nucleotide

polymorphisms (SNPs) provided the most reliable genetic markers, and amongst 55 strains of *S.*

marinoi, three 500 bp loci contained, on average, 46 SNPs, 103 unique alleles, and displayed

100% heterozygosity. The prevalence of heterozygosity was identified as a novel opportunity to

improve strain quantification and detect false positive artefacts during denoising of amplicon

sequences. Finally, we illustrate how metabarcoding of a single genetic locus can be used to

track strain abundances of 58 strains of *S. marinoi* in an artificial selection experiment. As future

genomics datasets become available and DNA sequencing technologies develop, Bamboozle has

flexible user settings enabling optimal barcodes to be designed for other species and applications.

Introduction

Microalgae are a diverse paraphyletic group of aquatic microbes responsible for half of global primary production (Falkowski et al. 1998; Field et al. 1998). Many species have enormous population sizes compared with multicellular organisms and contain some of the highest intraspecific genetic diversity observed within eukaryotes (Flowers et al. 2015). Despite their ecological importance, knowledge about individual species' functional diversity and evolutionary potential is limited (Godhe and Rynearson 2017). Microalgal blooms can contain thousands to millions of different clones (Sassenhagen et al. 2021), but designs of evolution experiments are currently constrained to only one or a few strains as they generally utilise clonal cultures, e.g., (Lohbeck et al. 2012; Schaum et al. 2017; Schaum et al. 2018; Sefbom et al. 2015; Wolf et al. 2019). If these strains are co-cultured in competition or *in situ*, as needed for many evolutionary and ecological questions, the individual strains cannot readily be identified or quantified using existing microscopic or genotyping technologies. Currently, genotyping of microalgae relies heavily on microsatellites (Rengefors et al. 2017), although newer sequencing methods such as RAD-seq provide a method for fine-scale genotyping and genomic information (Rengefors et al. 2021). Although suitable for population genetic and genomic studies, these methods still rely on single DNA samples from each strain for genotype identification. This severely restricts these methods' usefulness in quantitative evolutionary studies, where strain isolation is time-consuming or even impossible if the species is difficult to culture (Rengefors et al. 2021), or if the population's clonal diversity is large (Schaum et al. 2017; Scheinin et al. 2015).

An alternative approach to track cellular abundances involves using massively parallel amplicon sequencing of a diverse locus. The main advantage of such an approach is that millions of individual DNA molecules, and by extension the cells they originate from, can be quantified

from a single DNA sample. This simple principle - metabarcoding - has revolutionised the field

of microbial ecology in the past decade (Hebert et al. 2003; Taberlet et al. 2012). Depending on

the taxonomic group and the resolution required, various highly conserved genes are commonly

used in metabarcoding, including ribosomal RNA genes (16S, 18S, 23S, or 28S), mitochondrial

cytochrome c-oxidase subunit 1 (COI), and the chloroplastic ribulose-1,5-bisphosphate

carboxylase/oxygenase large subunit (*rbcL)* (Guo et al. 2015; Pujari et al. 2019; Yamada et al.

2017). Metabarcoding enables a relatively simple and standardised quantification of ecosystem-

wide parameters, such as diversity, presence of low abundance or cryptic taxa, and community

composition. However, the conserved nature of the available metabarcoding loci restricts their

usefulness as intraspecific markers (Canesi and Rynearson 2016; Godhe et al. 2006; Guo et al.

2015).

With a sufficiently diverse intraspecific barcode marker it should be possible to track the

abundances and quantify fitness of individual strains from a mixed microbial population. Such

an approach would be analogous to 'barcoding analysis by sequencing', or Bar-seq experiments,

where unique barcodes are used to replace genes in knock-out mutants (Robinson et al. 2014).

Bar-seq experiments then uses amplicon sequencing to quantify fitness changes in thousands of

non-lethal gene deletion mutants simultaneously, using a co-culture experimental design (Kim et

al. 2010; Li et al. 2019; Robinson et al. 2014; Smith et al. 2009), and it has accelerated the

functional annotations of several microbial species' genomes (Dent et al. 2005; Giaever and

Nislow 2014; Han et al. 2010). Access to 'intraspecific' DNA barcode loci in microalgae and

other microbes would provide a powerful tool to address fundamental questions about their

evolution and population structures, analogous to how metabarcoding and Bar-seq have radically

changed our understanding of microbial ecology and phenotypic functions of genes in the last

decade.

Genome sequencing projects are beginning to resolve the genetic diversity between species of

microalgae (Armbrust et al. 2004; Bowler et al. 2008; Merchant et al. 2007; Mock et al. 2017;

Read et al. 2013; Worden et al. 2009), and population genomic datasets are becoming

increasingly available (Blanc-Mathieu et al. 2017; Flowers et al. 2015; Osuna-Cruz et al. 2020;

Rastogi et al. 2020; von Dassow et al. 2015), providing information about intraspecific genetic

diversity. From this data, it should be possible to identify novel barcoding loci with intraspecific

resolution. Here we introduce Bamboozle, a bioinformatic approach that uses whole genome

sequencing data to identify intraspecific barcodes by scanning a species' entire genome for the

most suitable sites. As a proof-of-concept, we identify and design primers for a set of barcodes in

the diploid marine diatom *Skeletonema marinoi* and the haploid model chlorophyte

Chlamydomonas reinhardtii. Furthermore, we demonstrate how one single barcode locus can be

used to track strain selection in an artificial evolution experiment incorporating 58 environmental

strains of *S. marinoi* from two Baltic Sea inlets on Sweden's east coast. Based on the data from

this experiment we identify opportunities, as well as challenges, in employing amplicon

sequencing to address questions regarding evolutionary processes in microbes. Although we see

obvious applications in microalgae and other microbes, Bamboozle can identify intraspecific

barcodes for both haploid and diploid species when provided with a reference genome and whole

genome sequencing data from multiple individuals.

New approaches

Our aim with the Bamboozle tool was to enable identification of novel intraspecific barcoding loci with resolution down to the level of genotypes originating from one common population. To this end, we needed to identify loci fulfilling three criteria that render them suitable as barcoding loci, as outlined by Kress and Erickson (2008): 1) they should be short enough to allow amplification and sequencing with current technologies, 2) they should have significant interspecific (in our case intraspecific) variability, and 3) they should have conserved flanking regions that allow the binding of primers. Our approach builds on existing approaches for identifying novel barcoding loci by scanning the entire genome (Angers-Loustau et al. 2016; Paracchini et al. 2017). In contrast to the aforementioned studies, we focus on identification of barcodes with intraspecific resolution, implement several filters to optimise allele richness, exclude unsuitable genomic regions, and minimise the amount of manual screening needed.

The steps of the Bamboozle approach for barcode identification are outlined in Fig. 1 and described in detail below. The required input data includes a reference genome for the organism of interest, and whole genome sequencing (WGS) data from the strains to be analysed. Throughout the paper we use the strain concept (Lakeman et al. 2009), which for most practical purposes is synonymous with genotype in multicellular organisms. The WGS data is required in two formats, one pair per strain: 1) a sorted BAM file generated by aligning the WGS reads to the reference, and 2) single nucleotide polymorphism (SNP) data in VCF format generated from the aforementioned BAM file using GATK (Van der Auwera and O'Connor 2020).

In the case of diploid species, phased BAM and VCF files (i.e., where data from each allele is in a separate file) are also required to analyse the sequences of individual alleles. The main

Bamboozle tool, written and tested in Python version 3.7.10, consists of six main steps, detailed below.

1. *Identification of regions of unusual coverage.* When mapping whole genome shotgun sequencing reads to a reference genome, one expects a similar read coverage across the entirety of the reference. While short regions deviating from this expectation could result from potentially informative deletions or insertions in the mapped strain versus the reference, longer such regions could be due to misassemblies in the reference genome, repeat regions with significant length differences between strains, or gene duplications. As these phenomena could be problematic in identifying appropriate barcoding loci and downstream amplicon sequencing, SAMtools (Danecek et al. 2021) and Bedtools (Quinlan and Hall 2010) are called by Bamboozle and used to determine the median read depth of each contig in each strain (`samtools depth`), and to identify regions which deviate heavily from this median value (`bedtools genomecov`). By default, regions with coverage less than 50% or more than 200% of the contig median in that strain are flagged, in order to filter out large heterozygous deletions and gene duplications, respectively.

2. *Identification of variant positions.* Using a custom algorithm written in Python version 3.7.10, the locations of all variants in each strain are extracted from their respective VCF files and consolidated.

3. *Genome screening based on conserved sites.* As one of the core requirements for a suitable barcode is conserved flanking regions for primer binding, windows containing such conserved regions are identified in this step. Bamboozle traverses each contig/chromosome (step size 1; window size defined by the user), saving for further analysis those windows with

conserved regions (i.e., regions without any SNP's) at each end, based on the variant list

from step 2. The size of the window and the conserved end regions can be defined by the

user with the `--window_size` and `--primer_size` options, respectively, to account for

different amplification and sequencing strategies.

4. Merging of overlapping windows. Where both conserved end regions of two windows

overlap, these two are merged. For example, a 500 bp window with conserved regions from

contig positions 1-21 and 480-500, and a second such window with conserved regions from

contig positions 2-22 and 481-501, would be merged to form a single 501 bp window with

conserved regions from contig positions 1-22 and 480-501. This is done to reduce the

number of overlapping windows to be screened manually after Bamboozle outputs the end

results. For a visual example, see step 4 of Fig. 1.

5. Exclusion of windows with unusual read coverage. The merged windows are now compared

to the coverage data generated in step 1, and windows that overlap with a region of read

coverage fluctuation by ten or more bases are excluded. As length variations can be

potentially valuable in terms of differentiating between strains, allowing short (<10 bp)

stretches of irregular coverage (indicative of short indel regions) is intended as a compromise

between the potential for informative windows and the complications such regions may

introduce, as noted in step 1.

6. Identifying informative loci. As a unique sequence is required to identify and quantify each

strain in, for example, a strain selection or phenotyping experiment, SAMtools (`samtools`

`faidx`) and BCFtools (`bcftools consensus`) (Danecek et al. 2021) are applied to each

remaining window to generate consensus sequences for each strain (and each allele, in the

case of a diploid organism) using the reference genome and the respective VCF files. The

individual allele sequences are then analysed to determine if each strain in the dataset

contains at least one unique allele, in which case the window is reported to the user as a

potentially suitable barcoding locus.

As output, Bamboozle reports the identified barcoding loci in 1) a tab-separated file, giving

coordinates and metadata for each locus, 2) a BED file of locus coordinates, intended for easy

visualisation, such as in a genome browser, and 3) a multi-FASTA file for each locus, containing

the sequence of each allele in each strain. This enables quick and easy manual quality control of

the output, and design of primers for PCR tests and amplicon sequencing. As a proof-of-concept,

we performed amplicon sequencing on one locus identified in the diploid *S. marinoi* and used a

custom-made script that merges amplicons from paired-end Illumina data using BBMerge

(Bushnell et al. 2017), denoises the data using DADA2 (Callahan et al. 2016), and takes

advantage of heterozygosity to quality control the output and translate allele abundances into

counts of individual strains. Bamboozle offers a complete pipeline from WGS data through to

amplicon sequencing analysis of the barcoding loci, and its performance has been fine-tuned and

benchmarked using the diploid diatom *S. marinoi*.

Results

Performance of the bioinformatic pipeline: millions of potential barcoding loci reduced to

twenty-six

As expected, common metabarcoding loci lacked intraspecific resolution in *S. marinoi*, with 80-

100% of strains having the same single allele (Table S1). We applied Bamboozle to WGS data

from 54 strains of *S. marinoi*, to identify novel barcoding loci for use with 2x301 bp paired-end

Illumina MiSeq sequencing. Two of the initial 56 datasets were excluded from the analysis, one

due to low WGS read coverage, and one for being a clone of another strain in the study. Initial

attempts to identify suitable barcoding loci of 300-400 bp failed in this species (data not shown),

so we scanned for loci of 500 bp. Within the *S. marinoi* genome, most (99.5%) of the 54,753,029

possible 500 bp windows were filtered out due to either coverage fluctuations (main problem,

Table 1) or unconserved flanking regions. Of the remaining 263,000 windows, the average

frequency of SNP positions was 12±11, and the average number of alleles was 14 (Table 1). Four

potential barcoding loci were identified as containing at least one unique allele per strain and

passing all filters (Table 1). Three of these, plus an additional hypervariable locus (*Sm_C2W24*)

from a previous iteration of Bamboozle that did not include a coverage filter (step 1, Fig. 1),

were selected for further evaluation. Compared to other 500 bp windows passing the read depth

and conserved flanking region filters, three barcodes proved especially rich in terms of SNPs and

unique alleles, falling in the 97.6th to 99.9th percentiles, respectively (Fig. 2A and B).

Sm_C2W24 was predicted to contain an even higher SNP frequency (Fig. 2B). The *Sm_C2W24*

locus was retained through the analysis to assess the performance of Bamboozle in identifying

loci without coverage filter implementation.

The Bamboozle tool was also applied to seventeen haploid *C. reinhardtii* strains to identify loci

of ~300 bp. While the initial analysis on all strains returned no suitable barcoding loci,

separation by mating type (eight mt+ strains and nine mt- strains) was successful. From a

reference genome of 107,613,365 bp [nuclear genome, mitochondrion, plastid, and mt- locus

(Merchant et al. 2007)], with 107,161,335 potential 300 bp windows, only 137,000 loci (0.13%)

contained less than two-fold coverage deviation compared with the contig median (again the

main issue, Table 1) and had conserved flanking regions. About 40% of remaining windows

contained only one allele in the population, and less than 1% contained more than seven (Fig. 2C

and E). Only 22 of these loci were identified by Bamboozle as having one unique allele for all

strain genomes - fifteen for mt+ and seven for mt- (Table S2). In addition to being significantly

more diverse in allelic richness, these Bamboozle-identified loci also contained a much higher

number of SNPs than other loci across the genome (three out of four ranked in the 99.8th

percentile; Fig. 2D and F).

In silico annotation of intraspecific barcodes and common features

All 26 barcodes identified in *S. marinoi* and *C. reinhardtii* were inside predicted protein-coding

genes (Table S2). Several barcodes were also in close physical proximity within the same, or

adjacent genes, e.g., *Sm_C12W1*, *-2*, and *-3*, located within a 2,197 bp region containing two

gene models (Sm_t00009768-RA and Sm_t00009769-RA), as well as *Cr_Chr9W-12* and *-13*

inside *CHLRE_09g389134v5*, and *Cr_Chr11W-16, -17,* and *-18,* inside *CHLRE_11g467528v5*.

The predicted function of the proteins did not have a consistent pattern across the two species,

and ranged from ribosomal proteins (Sm_t00008465-RA [ribosome biogenesis protein WDR12]

and   *CHLRE_10g447800v5*   [60kDa   SS-A/Ro   ribonucleoprotein]),   to   enzymes

(*CHLRE_03g207250v5*   [putative   glutamine   synthetase]   and   *CHLRE_13g592050v5*

[allantoinase]) and ion transporters (*CHLRE_11g467528v5* [calcium channel]). Eleven barcodes

were located inside genes without conserved domains or with conserved Domains of Unknown

Function (DUF).

Four barcodes of each species were selected for primer design and further barcode development

(Fig. 3). Among these, no part of the ~500 bp barcoding loci was intronic in *S. marinoi*. Instead,

three out of four barcodes spanned domains with repetitive amino acid regions (*Sm_C2W24*,

Sm_C12W1, and *Sm_C16W4*) with the conserved primer sites anchored inside the same or

flanking conserved domains (Fig. 3A). In contrast, for *C. reinhardtii*, the major part of the 300

bp barcoding loci was intronic, while the conserved primer sites were located either in introns or

exons (Fig. 3C). A more detailed annotation of the barcode-containing genes is provided in the

Supplemental Information.

As future barcode applications may involve co-cultivation with other species, or *in situ*

experiments in natural microalgal communities, we evaluated to what extent the primers were

species-specific. In the target species, primers could be designed to amplify the selected loci

around the 21 bp conserved regions (Fig. 3B and D), although strain amplification appeared

uneven between strains for *Sm_C12W2, Sm_C16W4, Cr_Chr3W1*, and *Cr_Chr1W3*, suggesting

possible PCR amplification bias. Homologous genes for the *S. marinoi* barcode loci were

identified in three other species of centric diatoms within the order *Thalassiosirales*

(*Thalassiosira pseudonana* and *Thalassiosira oceanica,* and *Skeletonema subsalsum*) (Table S3).

In the two *Thalassiosira* species, each locus's primer sites contained mismatches in at least three

positions. However, in *S. subsalsum* the *Sm_C12W2* locus contained only one mismatch, and this

was also the only locus that yielded a PCR product using *S. subsalsum* DNA as template. None

of the primer pairs amplified in eleven other Baltic Sea phytoplankton species (Fig. S1A and B).

Although the 54 strains of *S. marinoi* all came from a remote population of *S. marinoi* inside the

Baltic Sea, the primer sites were conserved also in three strains from the American Atlantic

Coast and the Mediterranean Sea (Table S3), suggesting that the barcodes will function outside

the local Baltic Sea population. PCRs of the *C. reinhardtii* barcode loci *Cr_Chr9W12* and

Cr_Chr3W10 did not amplify in two other chlorophytes, including another *Chlamydomonas* sp.

strain, suggesting they could also be species-specific (Fig. S1C).

Accuracy of bioinformatic predictions and amplicon sequencing of barcodes in S. marinoi

The allele sequences predicted by Bamboozle were validated using amplicon sequencing of

individual strains' PCR products, or Sanger sequencing for the haploid *C. reinhardtii*. This was

done for two reasons: firstly, to generate the accurate allele sequences in the strains and contrast

this with Bamboozle's predictions based on genomic shotgun sequences, and secondly, to assess

what PCR and sequencing artefacts were prevalent.

Although *Sm_C2W24* was identified by an earlier version of Bamboozle that did not employ a

coverage filter, we retained this locus throughout the analysis to illustrate issues that micro- or

minisatellite-like loci can produce in the pipeline. Bamboozle predicted that *Sm_C2W24*

contained 95 SNPs and five indels within our studied population, but manual inspection

indicated that some strains had low sequencing coverage across the most variable region.

Amplicon sequencing revealed length differences of 198 bp amongst alleles of *Sm_C2W24* in the

S. marinoi population, caused by variable copy number (2 to 24 copies) of a 9 bp repeat

encoding Alanine- Asparagine- Glutamic acid (AN(E)) (length difference also confirmed by gel-

electrophoresis, Fig. 3B). The variant calling algorithm of GATK, and by extension Bamboozle,

had misinterpreted these repeats as a high abundance of SNPs (Table 2). Consequently, almost

all allele-sequences in the different strains were incorrectly predicted, and per-base accuracy

across the locus was only 82.6%. Hence, as outlined in Fig. 1, adjustments were made to

Bamboozle to filter out similar regions, and *Sm_C2W24,* together with 90-99% of the genomes

(Table 1), was disregarded in subsequent analysis.

All strain-specific sequences of the three *S. marinoi* barcoding loci, as well as *Cr_Chr9W12* and

Cr_Chr3W10 in *C. reinhardtii*, identified with the coverage filter were predicted with >99.9%

accuracy (Table 2). Looking into the reason for the remaining discrepancies, we discovered that

some of the strains were triploid (GP2-4_54 and VG1-2_78; Fig. 4), confounding the software's

expectations of diploidy. In other cases, the inaccuracies were caused by insufficient filtering of

the variant calling data, resulting in spurious base-calling errors.

Once the inaccuracies in the reference alleles were corrected using the confirmed sequences,

merged amplicons that still did not match the expected allele sequences were investigated to

determine the sources of error. This artifact search was limited to *S. marinoi* genotype samples

with coverage >100 amplicons per locus. A detailed description of the analysis and its findings

can be found in the Supplemental Information (Fig. S2, S3 and text). Briefly, 61-89% of

amplicons contained sequencing-errors, with 39-76% failing to merge. 11 to 40% of merged

amplicons mapped perfectly to known alleles, while remaining sequences were explained by

erroneous base-calling (10-40%), off-target amplification (0.1-12%), and PCR chimeras (1-3%).

Most amplicon sequence-errors could be either corrected or filtered out using the denoising

algorithm of DADA2 (Callahan et al. 2016). We finetuned the settings for the 523 bp

Sm_C12W1 locus to minimize false negative (alleles not predicted as real Amplicon Sequence

Variants [ASV]) and false positive (any ASVs not corresponding to a biological allele)

observations (Table S6). From mixed DNA samples, the most stringent settings evaluated failed

to predict three out of 110 alleles, while 16 artefact sequences were retained as ASV, although

they comprised only 0.03% of all merged sequences. By analysing the patterns of the remaining

false positive ASVs across these samples, searching for linked alleles, we were able to iteratively

identify the heterozygous alleles of three strains for which we lacked genotype sequencing data

(GP2-4_32, VG1-2_65, VG1-2_99). The remaining false positives were annotated as either

chimeras that DADA2's *de novo* chimera filter failed to remove, or one bp sequencing errors

from an abundant allele. Relaxing filtering settings, resulted in higher false positive ASVs (346,

totalling 1.2% of all amplicons), but detection of all known alleles as ASVs (Table S6). Using

the later settings on a mixed DNA sample, the *Sm_C12W1* alleles could be used to enumerate the

abundance of 58 strains accurately (Fig. 4).

Implementation of an intraspecific metabarcoding method in a microbial evolution experiment

The quantitative performance of *Sm_C12W1* as a metabarcoding locus was further evaluated in

an artificial evolution experiment using *S. marinoi*. In the artificial evolution experiment, 30

strains from Gåsfjärden (VG) and 28 strains from Gropviken (GP) were density normalised

(relative cell abundance of each strain was 3.4±0.89% based on cell counts using microscopy),

pooled separately for each population, and subjected to 42 days (50-100 generations) selection

with and without toxic copper stress. This data was used to evaluate the quantitative performance

of *Sm_C12W1*'s ability to track strain abundances.

Despite our expectations of diploidy in the *S. marinoi* strains, there were signs of polyploidy in

the experimental data. Some strains exhibited a skewed allele ratio closer to 1:3 (GP2-4_43,

GP2-4_55, GP2-4_63, VG1-2_45, and VG1-2_61; Fig. 5) suggesting that they could be

polyploids or that the PCR reaction amplified alleles with slightly different efficiencies. The

combined allele abundance of strains was sometimes over-represented two- to three-fold (e.g.,

GP2-4_57, VG1-2_45, VG1-2_90; Fig. 4) while others were underrepresented at a similar

magnitude (VG1-2_103, VG1-2_105, VG1-2_56; Fig. 4). This could not be explained by

differences in cell density (R^2=0.0003, slope=-0.0059, N=58), again suggesting ploidy

differences, or different DNA extraction efficiencies between cells of strains. Despite these

subtle deviations from expectations, the barcode *Sm_C12W1* could be used to identify all strains

at close to the expected relative abundances, most often with strains two alleles at close to a 1:1

ratio (Fig. 4). As the populations evolved through selection on strain diversity, one or two strains

ultimately became dominant in each treatment, and during this process, their allelic ratio was

conserved at 1:1 (Fig. 5). These observations show that relative changes in allele frequencies

should reflect changes in the relative abundance of strains.

Throughout the experiment, most strains experienced negative selection. At around 10 ASV

observations per sample, DADA2's algorithm started filtering out observations (compare Fig. 5,

with non-denoised sequence matches in Fig. S4), indicating that false-negative artefacts were

created. However, DADA2 effectively removed all but four false-positive observations in the

experiment (allele of GP population seen in the VG population, and vice versa: Tables S6). The

remaining four false positives appeared to be chimeric, as did allele GP2-4_44#1, which

displayed a skewed allelic ratio representing up to 5% (>1,000 copies) of amplicons in several

replicate samples, while GP2-4_44#2 was not detected at all (Fig. 5). As the first 316 bp of this

allele are identical with GP2-4_27#1, and the last 354 bp with GP2-4_27#2 (the two alleles of

the most dominant strain in the same saples), there is a central 147 bp region where a PCR

chimera from these two alleles would produce GP2-4_44#1. In addition to enabling chimera

detection in situations like the one above, heterozygosity provided an additional analytical

advantage, enabling differential equations to accurately partition alleles that were shared between

two or three strains (Fig. 4 and 5). These results show that heterozygosity can be used to both

filter out chimeric observations, and to enable improved quantification of strains, even if they

lack a unique allelic marker.

Discussion

By scanning the entire genome for suitable loci, we illustrate how the Bamboozle approach can identify hyper-variable barcodes with intraspecific resolution. In the case of both *S. marinoi* and *C. reinhardtii*, most of the genome (>90%) was not suitable for barcoding and was effectively filtered out by the algorithm. Of the remaining loci fulfilling the criteria for barcodes (Kress and Erickson 2008), only a fraction contained sufficient allelic richness to provide allele-rich markers for strain quantification, highlighting the power of a whole-genome scan approach for development of novel intraspecific barcodes. Using a combination of original (54) and published [17: (Flowers et al. 2015; Ness et al. 2016)] WGS datasets, we illustrate how such intraspecific barcodes can be designed from either publicly available sequencing data, or through the generation of new data for a species or population of interest.

Within our amplicon sequencing data, many amplicons (~40%) did not correspond to any known allele in our database. Therefore, we attempted to identify the sources of these artefacts, and how to address them. This was done in order to make recommendations for future studies using workflows similar to ours.

First, we note that the 523 bp *Sm_C12W1* locus, paired with Illumina MiSeq's 2×300 bp sequencing technology, creates some challenges that may not be encountered in shorter loci. For example, we could not use DADA2 as a stand-alone tool to process reads since it truncated 100% of all amplicons during merging (Table S6). BBMerge performed better but still failed to merge 78% of all reads, due to the short overlapping region and low base-call quality at the 3' end of reads. By pairing BBMerge with DADA2, we found a flexible approach that could correct 99-99.97% of all sequencing errors in reads that had merged, with only spurious false-positive

allelic observations (Fig. 5). Although a shorter locus is preferable and will likely perform better

than this, our analysis shows that the tools available to make robust biological inferences from

noisy amplicon data extend to 500 bp barcode loci, albeit with a loss of four-fifths of the

sequence data. If future studies cannot identify window sizes smaller than 500 bp, accurate long-

read sequencing of barcodes (e.g., PacBio's Circular Consensus Sequences technique (Rhoads

and Au 2015)), or paired end 300 bp sequences from the newer platform Illumina NovaSeq 6000

could be options worth considering. Long-read sequencing, in combination with DADA2's error

correction, has enabled differentiation between pathogenic and benign *E. coli* strains using single

SNP markers and metabarcoding of the whole ~1450 bp 16S rRNA gene (Callahan et al. 2019).

Such a sequencing approach opens up for loci of several kb to be used as intra-specific barcodes.

In contrast to sequencing errors, PCR chimeras provided a more problematic type of artefact. We

found that PCR chimeras could not be removed effectively using software built for processing

metabarcoding data, such as Mothur (Rognes et al. 2016; Schloss et al. 2009), UCHIME2 (Edgar

2016), or DADA2s chimera removal algorithm (Callahan et al. 2016). Exacerbating this issue is

the potential for 'perfect fake' sequences (chimeras identical to another, non-chimeric sequence in

the dataset (Edgar 2016)). Out of the 110 alleles in *Sm_C12W1,* we identified several cases of

such 'perfect fakes' (e.g., allele 1 in GP2-4_44; Fig. 5), and these instances should become

progressively more numerous if more strains and alleles are included in the experiments and

analysis. However, we could identify these PCR chimeras in our experimental design primarily

by looking at deviation from the expected heterozygous allelic ratios of individual strain (Fig. 5).

Depending on the research question and experimental design, chimeras can be more or less

effectively filtered out using similar approaches in future studies. If chimeras risk causing

erroneous biological interpretations from the data, approaches to reduce the number of chimeras

produced includes optimising a low PCR cycle threshold (Smyth et al. 2010) or using chimera-

free PCR kits, such as emulsion PCR (Williams et al. 2006).

The allele phasing step (upstream of Bamboozle) also created some problems, as it could not

handle cases of triploidy (Fig. 4). Haplotype phasing of polyploids is non-trivial and benefits

greatly from long-read data (Abou Saada et al. 2021). However, as long as polyploidy is

accurately identified (e.g., by amplicon sequencing of each strain's barcode individually), it

should not provide any issues with downstream analysis of strain abundances and may even aid

in providing increased allelic diversity for quantification purposes. Consequently our evaluation

of Bamboozle as both a tool to identify intra-specific barcodes suggest it can accurately locate

such hypervariable regions. Our quantitative analyse show that with appropriate choice of

barcode lengths, PCR kits, sequencing platforms and denoising strategies, amplicon sequencing

of intra-specific barcodes can be used to track the abundances of large numbers of strains from

mixed DNA samples.

413
Conclusions

We built and employed the Bamboozle tool to develop metabarcoding loci that provide

intraspecific resolution in the diploid diatom *S. marinoi*. This allowed us to identify and quantify

the abundance of 58 environmental strains from mixed DNA sample. We illustrated the

usefulness of such an intraspecific barcode locus in an artificial evolution experiment where we

tracked the abundance of all strains during co-cultivation. Although further tests are needed to

assay performance in new populations and environmental samples, as well as the amplicon

sequencing performance of the *C. reinhardtii* loci, our preliminary tests suggest that intra-

specific metabarcoding can simplify or enable novel studies of microbial ecology and evolution

in natural and artificial settings. This is not limited to artificial selection experiments like ours

but possibly also to determine genotype diversity in natural populations, studies of ancient DNA

from sediment cores (Adams et al. 2019; Härnström et al. 2011), mesocosm incubations of

natural phytoplankton communities (Scheinin et al. 2015; Tatters et al. 2013), effects of

predation (Sjöqvist et al. 2014), nutrient competition within (Collins 2011) and between species

(Descamps-Julien and Gonzalez 2005), or other drivers that are challenging or impossible to

observe in mono-culture experiments (Baert et al. 2016; Collins and Schaum 2021). Using the

haploid chlorophyte *C. reinhardtii*, we illustrate how the Bamboozle tool should, if possible, be

able to identify loci in both haploid and diploid organisms with a reference genome and whole

genome sequencing data from multiple genotypes. Our comparative analysis of barcoding loci

identified in *S. marinoi* and *C. reinhardtii* suggests that much of the genome is unsuitable for

barcode design, and that different species have different optimal barcode loci. Consequently,

Bamboozle's strategy of scanning the entire genome of a species is a suitable approach to

identify novel barcodes for evolutionary studies.

Acknowledgments

The authors would like to thank André Soares and Vilma Canfjorden who have developed

additional functionality of the Bamboozle tool. Olga Kourtchenko, Lara Hoepfner, and Nora

Klasen, assisted in experiments or DNA sample preparation. The manuscript benefited from

revision suggestions and thoughtful comments by Kerstin Johannesson.

Data availability

The code for Bamboozle is available at https://github.com/topel-research-group/Bamboozle and

the WGS and amplicon sequencing data has been deposited at NCBI under BioProject

PRJNA939970.

Materials and Methods

Strain retrieval and whole genome sequencing

Individual strains of *S. marinoi* were germinated and isolated from sediment resting stages using standard micropipetting techniques (Härnström et al. 2011). Surface sediment was collected from two semi-enclosed inlets of the brackish Baltic Sea, where one (Gåsfjärden [VG]; 57°34.35'N 16°34.98'E) has been exposed to historical copper mining (Söderhielm and Sundblad 1996) while the other (Gropviken [GP]; 58°19.92'N 16°42.35'E) has not. A total of 69 and 55 strains were isolated from VG and GP, with 88% and 94% survival, respectively. Strains were cultured in locally-sourced seawater (salinity 7), which had been sterile-filtered (Sarstedt's [Helsingborg, Sweden] 0.2 mm polyethersulfone membrane filter), and amended with f/2 nutrients (Guillard 1975) and 106 µM SiO_2. Cultures were continuously screened for contamination by other microalgae, and auxospore formation or bimodal cell sizes, which indicate sexual inbreeding in *S. marinoi* (Ferrante et al. 2019), and such cultures were discarded.

DNA extraction was performed within one month of revival from resting stages in a total of 28 strains from GP and 30 from VG. Cultures were made temporarily axenic via a combination of mechanical cleaning (triple washing in 20 µg mL[-1] Triton X-100 media and cell collection on a 3.0-µm polycarbonate filter), followed by a five-day antibiotic cocktail treatment (90 µg mL[-1] Paromomycin, ten µg mL[-1] Ciprofloxacin, and 40 µg mL[-1] Cefotaxime). On day six, cells were collected via centrifugation in 50 mL Falcon tubes (10 000 x g, for ten min.), flash-frozen in liquid nitrogen, and stored at -80ºC. DNA was extracted using a CTAB-phenol–chloroform protocol as described in (Godhe et al. 2001), but with additional RNA digestion during cell lysis (65ºC for 60 min. using 1 mg RNaseA mL[-1] CTAB buffer). DNA yield was quantified using

Qubit (Thermo Fisher Scientific). In two subsequential attempts, strains VG1-2_65 and VG1-

2_99 did not survive the antibiotic treatment, and extractions yielded insufficient amounts of

DNA for PCRs.

Whole genome sequencing was performed on the remaining 56 *S. marinoi* strains using ½ lane

of an Illumina NovaSeq S4 flowcell. Library preparation was done using Nextera's PCR-based

protocol with enzymatic fragmentation. This resulted in 722.82 Mreads (2x150 bp paired-end).

Data preparation for the pipeline – S. marinoi

Trimming of reads was performed on the whole genome sequencing data to remove the first 15

bases of each read. The trimmed reads were mapped to the *S. marinoi* reference genome version

1.1.2 (https://albiorix.bioenv.gu.se/Skeletonema_marinoi.html) using Bowtie2 version 2.3.4.3

(Langmead and Salzberg 2012), and the resultant SAM files were converted to sorted BAM files

and indexed using SAMtools version 1.9 (Danecek et al. 2021). Variant calling was performed

using GATK version 4.1.8.0 (Van der Auwera and O'Connor 2020), following the Best Practices

recommendations on the tool's website, and the resultant VCF files were indexed using

BCFtools version 1.10.2 (Danecek et al. 2021). In addition, SAMtools version 1.10 was used to

phase the BAM files, and GATK was re-run on each of these phased BAM files individually.

These phased files were indexed as described above.

Data preparation for the pipeline – C. reinhardtii

Sequencing data from the strains of interest was downloaded from either the NCBI Sequence

Read Archive (SRA; downloaded using the `prefetch` and `fastq-dump` tools from the SRA

toolkit version 2.9.6 (Leinonen et al. 2010)) or the European Nucleotide Archive (ENA;

downloaded using the `enaDataGet` tool from enaBrowserTools version 1.6

[https://github.com/enasequence/enaBrowserTools]) (Table S4; retrieved 22 February 2021).

PCR duplicates were removed using the filterPCRdupl Perl script version 2.3

(https://github.com/linneas/condetri), then trimmed with Cutadapt version 3.2 (Martin 2011),

using a quality threshold of 30 and a post-trimming length threshold of either 90 (ENA reads) or

45 (SRA reads).

The trimmed reads were mapped to a reference consisting of the following sequences from

C. reinhardtii: the contig-level assembly of the reference nuclear genome version 5.5 (accession

no. ABCN00000000.2); the mitochondrial genome (accession no. NC_001638.1); the plastid

genome (accession no. NC_005353.1); and the mating type minus locus (accession no.

GU814015.1). Mapping was performed using Bowtie2 version 2.3.4.3 (Langmead and Salzberg

2012), with the resultant SAM files being converted to sorted BAM files and indexed using

SAMtools version 1.10 (Danecek et al. 2021). Variant calling and VCF indexing were performed

as described above for *S. marinoi.*

Bamboozle workflow

Barcodes for *S. marinoi* were obtained using commit 75467da of the Bamboozle main branch

(except for *Sm_C2W24*, which was obtained using commit ae28b1b). Barcodes for *C. reinhardtii*

were obtained using commit db64632 of the `matt_improvement` branch. The latter commit has

the fourth and fifth steps of the pipeline reversed (i.e., windows with irregular coverage are

excluded prior to window merging), along with other minor improvements.

The input files (reference genome FASTA files, plus BAM and VCF files [and respective

indexes] for each strain) were prepared as described above.

Bamboozle was run in both species with a `-window_size` parameter of both 500 and 300, and a

`-primer_size` parameter of 21.

Development of the bioinformatic pipeline

The predicted barcodes were first visualised against the reference genomes using IGV version

2.6.3 (Robinson et al. 2011) to look for anomalies. In an early iteration of the pipeline (commit

ae28b1b) without coverage depth filters (steps 1 and 5, Fig. 1), many loci contained obvious

errors in the conserved primer sites, or spanned regions of low read coverage (e.g., *Sm_C2W24*)

suggesting the pipeline could not effectively process regions with large indels (e.g. micro- or

mini-satellites). This issue has largely been resolved by adding the coverage depth filter, as

evidenced by the reduction of identified barcodes in *S. marinoi* from 245 to 4.

Additional adjustments of the pipeline used for analysing both *S. marinoi* and *C. reinhardtii*,

included extending the code to allow analysis of haploid genomes, and steps 4 and 5 of the

pipeline were switched (i.e., windows with irregular coverage are excluded prior to window

merging), in order to avoid excluding potentially informative windows without coverage issues.

In silico annotation of intraspecific barcodes

The gene models of the predicted barcodes were retrieved from version 5.5 of the *C. reinhardtii*

reference genome (https://ensembl.gramene.org/Chlamydomonas_reinhardti) and version 1.1.2

of the *S. marinoi* reference genome (https://albiorix.bioenv.gu.se/Skeletonema_marinoi.html).

Functional domains were annotated using NCBI's CD-Search webtool (Marchler-Bauer and

Bryant 2004). When gene models lacked a functional annotation, putative functions were

assigned by a homology search using BLASTp against the NCBI database (Altschul et al. 1990),

and comparison of functional domain structure.

To assess primer specificity outside our Baltic Sea strains of *S. marinoi,* we mapped

transcriptional data from three strains from global culture collections (Keeling et al. 2014), and

manually looked for variants within the predicted conserved regions. A targeted BLASTn search

was used to identify orthologous genes in three species within the order *Thalassiosirales* with

reference genomes, namely: *Thalassiosira oceanica* (Lommer et al. 2012), *Thalassiosira*

pseudonana (Armbrust et al. 2004), and the closely related *Skeletonema subsalsum* (Sarno et al.

2005) using an in-house draft genome for the latter (Pinder et al. unpublished data).

Confirmation of the bioinformatic predictions

Illumina amplicon sequencing was used to assess the sequence accuracy of each of Bamboozle's

locus predictions in the diploid *S. marinoi,* and Sanger sequencing for the haploid *C. reinhardtii.*

Primers (standard de-salted oligos) were designed for the 21 bp conserved regions flanking the

variable region (Table S5) and extended with Illumina adapter sequences for *S. marinoi. rbcL*

was used as a positive control of amplification in all diatoms (Guo et al. 2015), and 16s or 18S

rRNA across other phytoplankton taxa (Stoeck et al. 2010; Sundberg et al. 2013). The barcodes

were amplified from 100 ng DNA and a final primer concentration of 0.1 µM using the

Phusion® High-Fidelity PCR Kit (Thermo Scientific) with individually optimised annealing

temperatures between 62 and 65ºC. A one-step PCR reaction, with adapter extended primers,

was run for 30 cycles, with 5 s denaturation (98ºC), 5 s annealing, 30 s extension (72ºC), and a

final 5 min extension. The evenness of amplification across strains was qualitatively assessed

using gel electrophoresis. Strains with poor barcode amplification were re-run with either

increased DNA concentration or up to 35 PCR cycles. Species specificity of the primers was

assessed using DNA from two strains of *Skeletonema subsalsum*, as well as a mixture of other

common phytoplankton species from the Baltic Sea (see Supplemental information). The *C.*

reinhardtii primers were only tested on two other green algae strains; *Chlamydomonas* sp and

Microglena sp.

For the 56 *S. marinoi* strains, the barcode alleles were sequenced individually using 2x301 bp

reads on 1/10 of a lane on the Illumina MiSeq platform with v3 chemistry (expected 1.8 million

reads in total). The four PCR reactions were run independently for each barcode, and the

products pooled based on volume. Further library preparation, including amplicon size selection

>300 bp, Nextera dual-indexing, and amplicon sequencing were performed at SciLifeLab (NGI

Stockholm), according to the manufacturer's instructions.

Identifying sources of error in the amplicon sequencing results

Paired end reads were first quality filtered and trimmed using Cutadapt version 3.2 to remove

adapter and primer sequences, with a quality threshold of 28 and a minimum read length of 180

bp. The resultant trimmed reads were merged using BBMerge version 38.86 (Bushnell et al.

2017). Amplicon sequences from the individually strains that did not match the two expected

allele sequences were investigated to determine the sources of error. Chimera detection was

initially performed using the `chimera.vsearch` function of Mothur version 1.47.0 (Schloss et al.

2009). However, as this highlighted fewer chimeras than we knew to exist through manual

inspection, we instead switched to the `uchime2_ref` function of Usearch version 11.0.667

(Edgar 2016). This was run with the options `'-strand plus -mode sensitive'` against a

database containing all alleles expected to appear in that strain.

Identification of truncated sequences, sequences containing Ns, off-targets, and localisation of

sequencing errors and SNPs along the amplicons, was performed using a custom Python script,

incorporating Bowtie2 version 2.3.4.3 (Langmead and Salzberg 2012), SAMtools version 1.12,

and BCFtools version 1.12 (Danecek et al. 2021).

Performance of intraspecific metabarcoding in a selection experiment

The quantitative performance of *Sm_C12W1* barcodes locus was validated in a selection

experiment using the 58 *S. marinoi* strains. A detailed description of this experimental design

and the results will be published elsewhere (Andersson et al, unpublished). Briefly, the single

strain cultures (non-axenic) were pooled back to their original populations (Gropviken: GP and

Gåsfjärden: VG) at even densities, and maintained in exponential growth phase via a semi-

continuous dilution scheme as outlined in Andersson et al. (2020) through dilutions every third

day for a total of 42 days. Five 100 mL replicates per population were grown with toxic copper

stress of 8.65 µM $CuSO_4$, and five replicates were grown without toxic stress. The dilution

bottleneck was ~10,000 chains of cells per replicate. Cultures were harvested at the start of the

experiment, after nine, and 42 days of selection. DNA was extracted from experimental samples

as for the WGS preparation outlined above, but without the axenic treatment. Barcodes were

PCR amplified from each replicate timepoint. For the single time 0 sample, two technical DNA

extractions replicates × two PCR replicates were processed and analysed in parallel (i.e., N=4

technical replicates). The selection experiment samples were indexed and pooled together with

the individual genotypes, on the same MiSeq run using 9/10 of the read capacity (i.e., an

expected 16.2 million reads).

The individual strains' genotype samples were used to create a database of all strains' alleles,

and amplicons from the selection experiment samples were compared against this database

(script available at: https://github.com/topel-research-group/Live2Tell). As the experimental

samples contained mixtures of strains at variable densities, and allele calling was expected to

require single SNP resolution across long amplicons (up to 523 bp), we explored avenues to

denoise the data. Initial attempts to use the nf-core/ampliseq pipeline version 2.4.0 (Straub et al.

2020) failed to detect all expected alleles. Running DADA2 version 1.16.0 (Callahan et al. 2016)

– a component of the nf-core/ampliseq pipeline – as a standalone program for identifying

amplicon sequence variants (ASVs) proved more successful. Different settings and combinations

of steps were iteratively explored (described in detail in Supplemental information). We used

four criteria to evaluated the quality of the output including: 1) how many known allele

sequences were <u>not </u>identified as ASVs (false negatives); 2) how many unexpected ASVs were

outputted (false positives); 3) what fraction of total raw reads assembled into amplicons

matching either known alleles or false positives; and 4) if false positive or false negative

observations risked affecting the biological interpretation of the artificial evolution experiment.

The relative abundances of strains were quantified based on the number of amplicons matching

its two alleles. In this step, a set of differential equations were used to parse out the observations

of alleles shared between strains. In general, the proportion of shared allele i_1 belonging to a

specific strain x, y, and z was computed individually for each sample based on strains second

unique allele (x_2, y_2, and z_2):

Eq. 1: Reads of i_1 belonging to strain $x = i_1 \times \dfrac{\text{strain } x \text{ allele } x_2}{(\text{strain } x \text{ allele } x_2 + \text{strain } y \text{ allele } y_2 + \text{strain } z \text{ allele } z_2)}$

The two heterozygous allele counts were then summed up to represent the abundance of each

strain, with the third allele of triploid strains omitted. Amplicon counts per strain were

simultaneously normalized to relative abundance (RA) as:

Eq. 2: $$\text{RA strain } x = \frac{\text{counts allele } x_1 + counts \text{ allele } x_2}{\sum counts\ allels_{a-z}}$$

To validate the approach, we assessed how closely the barcode enumeration of strains matched

with microscopic cell counts in the time zero samples and how closely the abundance of each

strain's two alleles correlated throughout the selection experiment.

References

Abou Saada, O., A. Tsouris, C. Eberlein, A. Friedrich, and J. Schacherer. 2021. nPhase: an accurate and contiguous phasing method for polyploids. Genome Biol. **22:** 1-27. doi:10.1186/s13059-021-02342-x

Adams, C. I., M. Knapp, N. J. Gemmell, G.-J. Jeunen, M. Bunce, M. D. Lamare, and H. R. Taylor. 2019. Beyond biodiversity: can environmental DNA (eDNA) cut it as a population genetics tool? Genes **10:** 192. doi:10.3390/genes10030192

Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment search tool. J. Mol. Biol. **215:** 403-410. doi:10.1016/S0022-2836(05)80360-2

Andersson, B., A. Godhe, H. L. Filipsson, K. Rengefors, and O. Berglund. 2020. Differences in metal tolerance among strains, populations, and species of marine diatoms-importance of exponential growth for quantification. Aquat. Toxicol. **226:** 105551. doi:10.1016/j.aquatox.2020.105551.

Angers-Loustau, A., M. Petrillo, V. Paracchini, D. M. Kagkli, P. E. Rischitor, A. Puertas Gallardo, A. Patak and others 2016. Towards plant species identification in complex samples: a bioinformatics pipeline for the identification of novel nuclear barcode candidates. PLoS one **11:** e0147692. doi:10.1371/journal.pone.0147692

Armbrust, E. V., J. A. Berges, C. Bowler, B. R. Green, D. Martinez, N. H. Putnam, S. Zhou and others 2004. The genome of the diatom *Thalassiosira pseudonana*: ecology, evolution, and metabolism. Science **306:** 79-86. doi:10.1126/science.1101156

Baert, J. M., F. De Laender, K. Sabbe, and C. R. Janssen. 2016. Biodiversity increases functional and compositional resistance, but decreases resilience in phytoplankton communities. Ecology **97:** 3433-3440. doi:10.1002/ecy.1601

Blanc-Mathieu, R., M. Krasovec, M. Hebrard, S. Yau, E. Desgranges, J. Martin, W. Schackwitz and others 2017. Population genomics of picophytoplankton unveils novel chromosome hypervariability. Sci. Adv. **3:** e1700239. doi:10.1126/sciadv.1700239

Bowler, C., A. E. Allen, J. H. Badger, J. Grimwood, K. Jabbari, A. Kuo, U. Maheswari and others 2008. The *Phaeodactylum* genome reveals the evolutionary history of diatom genomes. Nature **456:** 239-244. doi:10.1038/nature07410

Bushnell, B., J. Rood, and E. Singer. 2017. BBMerge–accurate paired shotgun read merging via overlap. PloS one **12:** e0185056. doi:10.1371/journal.pone.0185056

Callahan, B. J., P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, and S. P. Holmes. 2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nat. Methods **13:** 581-583. doi:10.1038/nmeth.3869

Callahan, B. J., J. Wong, C. Heiner, S. Oh, C. M. Theriot, A. S. Gulati, S. K. McGill and others 2019. High-throughput amplicon sequencing of the full-length 16S rRNA gene with single-nucleotide resolution. Nucleic Acids Res. **47:** e103-e103. doi:10.1093/nar/gkz569

Canesi, K. L., and T. A. Rynearson. 2016. Temporal variation of *Skeletonema* community composition from a long-term time series in Narragansett Bay identified using high-throughput DNA sequencing. Mar. Ecol. Prog. Ser. **556:** 1-16. doi:10.3354/meps11843

Collins, S. 2011. Competition limits adaptation and productivity in a photosynthetic alga at elevated CO2. Proc. R. Soc. B. **278:** 247-255. doi:10.1098/rspb.2010.1173

Collins, S., and C. E. Schaum. 2021. Growth strategies of a model picoplankter depend on social milieu and p CO2. Proc. R. Soc. B **288:** 20211154.

670 Danecek, P., J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M. O. Pollard, A. Whitwham and
671    others 2021. Twelve years of SAMtools and BCFtools. Gigascience **10:** giab008.
672    doi:10.1093/gigascience/giab008
673 Dent, R. M., C. M. Haglund, B. L. Chin, M. C. Kobayashi, and K. K. Niyogi. 2005. Functional
674    genomics of eukaryotic photosynthesis using insertional mutagenesis of *Chlamydomonas*
675    *reinhardtii*. Plant Physiol. **137:** 545-556. doi:10.1104/pp.104.055244
676 Descamps-Julien, B., and A. Gonzalez. 2005. Stable coexistence in a fluctuating environment: an
677    experimental demonstration. Ecology **86:** 2815-2824. doi:10.1890/04-1700
678 Edgar, R. C. 2016. UCHIME2: improved chimera prediction for amplicon sequencing. BioRxiv**:**
679    074252. doi:10.1101/074252
680 Falkowski, P. G., R. T. Barber, and V. V. Smetacek. 1998. Biogeochemical Controls and
681    Feedbacks on Ocean Primary Production. Science **281:** 200-207.
682    doi:10.1126/science.281.5374.200
683 Ferrante, M. I., L. Entrambasaguas, M. Johansson, M. Töpel, A. Kremp, M. Montresor, and A.
684    Godhe. 2019. Exploring Molecular Signs of Sex in the Marine Diatom *Skeletonema*
685    *marinoi*. Genes **10:** 494. doi:10.3390/genes10070494
686 Field, C. B., M. J. Behrenfeld, J. T. Randerson, and P. Falkowski. 1998. Primary production of
687    the biosphere: integrating terrestrial and oceanic components. Science **281:** 237-240.
688    doi:10.1126/science.281.5374.237
689 Flowers, J. M., K. M. Hazzouri, G. M. Pham, U. Rosas, T. Bahmani, B. Khraiwesh, D. R. Nelson
690    and others 2015. Whole-genome resequencing reveals extensive natural variation in the
691    model green alga *Chlamydomonas reinhardtii*. Plant Cell **27:** 2353-2369.
692    doi:10.1105/tpc.15.00492
693 Giaever, G., and C. Nislow. 2014. The yeast deletion collection: a decade of functional
694    genomics. Genetics **197:** 451-465. doi:10.1534/genetics.114.161620
695 Godhe, A., M. R. McQuoid, I. Karunasagar, I. Karunasagar, and A. S. Rehnstam☐Holm. 2006.
696    Comparison of three common molecular tools for distinguishing among geographically
697    separated clones of the diatom *Skeletonema marinoi* Sarno et Zingone
698    (Bacillariophyceae) 1. J. Phycol. **42:** 280-291. doi:10.1111/j.1529-8817.2006.00197.x
699 Godhe, A., S. K. Otta, A. S. Rehnstam-Holm, I. Karunasagar, and I. Karunasagar. 2001.
700    Polymerase chain reaction in detection of *Gymnodinium mikimotoi* and *Alexandrium*
701    *minutum* in field samples from southwest India. Mar. Biotechnol. **3:** 152-162.
702    doi:10.1007/s101260000052
703 Godhe, A., and T. Rynearson. 2017. The role of intraspecific variation in the ecological and
704    evolutionary success of diatoms in changing environments. Phil. Trans. R. Soc. B. **372:**
705    20160399. doi:10.1098/rstb.2016.0399
706 Guillard, R. R. L. 1975. Culture of phytoplankton for feeding marine invertebrates., p. 29–60. *In*
707    W. L. Smith and M. H. Chanley [eds.], *Culture of marine invertebrate animals.* Springer.
708 Guo, L., Z. Sui, S. Zhang, Y. Ren, and Y. Liu. 2015. Comparison of potential diatom
709    'barcode' genes (the 18S rRNA gene and ITS, COI, rbcL) and their effectiveness in
710    discriminating and determining species taxonomy in the Bacillariophyta. Int. J. Syst.
711    Evol. Microbiol. **65:** 1369-1380. doi:10.1099/ijs.0.000076
712 Han, T. X., X.-Y. Xu, M.-J. Zhang, X. Peng, and L.-L. Du. 2010. Global fitness profiling of
713    fission yeast deletion strains by barcode sequencing. Genome Biol. **11:** 1-13.
714    doi:10.1186/gb-2010-11-6-r60

Hebert, P. D. N., A. Cywinska, S. L. Ball, and J. R. deWaard. 2003. Biological identifications
through DNA barcodes. Proc. R. Soc. B. **270:** 313-321. doi:10.1098/rspb.2002.2218
Härnström, K., M. Ellegaard, T. J. Andersen, and A. Godhe. 2011. Hundred years of genetic
structure in a sediment revived diatom population. Proc. Natl. Acad. Sci. USA **108:**
4252–4257. doi:10.1073/pnas.1013528108
Keeling, P. J., F. Burki, H. M. Wilcox, B. Allam, E. E. Allen, L. A. Amaral-Zettler, E. V.
Armbrust and others 2014. The Marine Microbial Eukaryote Transcriptome Sequencing
Project (MMETSP): illuminating the functional diversity of eukaryotic life in the oceans
through transcriptome sequencing. PLoS Biol. **12:** e1001889.
doi:10.1371/journal.pbio.1001889
Kim, D.-U., J. Hayles, D. Kim, V. Wood, H.-O. Park, M. Won, H.-S. Yoo and others 2010.
Analysis of a genome-wide set of gene deletions in the fission yeast
Schizosaccharomyces pombe. Nat. Biotechnol. **28:** 617-623. doi:10.1038/nbt.1628
Kress, W. J., and D. L. Erickson. 2008. DNA barcodes: genes, genomics, and bioinformatics.
Proc. Natl. Acad. Sci. USA **105:** 2761-2762. doi:10.1073/pnas.0800476105
Lakeman, M. B., P. von Dassow, and R. A. Cattolico. 2009. The strain concept in phytoplankton
ecology. Harmful Algae **8:** 746-758. doi:10.1016/j.hal.2008.11.011
Langmead, B., and S. L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. Nat.
Methods **9:** 357-359. doi:10.1038/nmeth.1923
Leinonen, R., H. Sugawara, M. Shumway, and I. N. S. D. Collaboration. 2010. The sequence
read archive. Nucleic Acids Res. **39:** D19-D21. doi:10.1093/nar/gkq1019
Li, X., W. Patena, F. Fauser, R. E. Jinkerson, S. Saroussi, M. T. Meyer, N. Ivanova and others
2019. A genome-wide algal mutant library and functional screen identifies genes required
for eukaryotic photosynthesis. Nat. Genet. **51:** 627-635. doi:10.1038/s41588-019-0370-6
Lohbeck, K. T., U. Riebesell, and T. B. Reusch. 2012. Adaptive evolution of a key
phytoplankton species to ocean acidification. Nat. Geosci. **5:** 346. doi:10.1038/ngeo1441
Lommer, M., M. Specht, A. S. Roy, L. Kraemer, R. Andreson, M. A. Gutowska, J. Wolf and
others 2012. Genome and low-iron response of an oceanic diatom adapted to chronic iron
limitation. Genome Biol. **13:** R66. doi:10.1186/gb-2012-13-7-r66
Marchler-Bauer, A., and S. H. Bryant. 2004. CD-Search: protein domain annotations on the fly.
Nucleic Acids Res. **32:** W327-W331. doi:10.1093/nar/gkh454
Martin, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet. journal **17:** 10-12. doi:10.14806/ej.17.1.200
Merchant, S. S., S. E. Prochnik, O. Vallon, E. H. Harris, S. J. Karpowicz, G. B. Witman, A.
Terry and others 2007. The *Chlamydomonas* genome reveals the evolution of key animal
and plant functions. Science **318:** 245-250. doi:10.1126/science.1143609
Mock, T., R. P. Otillar, J. Strauss, M. McMullan, P. Paajanen, J. Schmutz, A. Salamov and
others 2017. Evolutionary genomics of the cold-adapted diatom *Fragilariopsis cylindrus*.
Nature **541:** 536-540. doi:10.1038/nature20803
Ness, R. W., S. A. Kraemer, N. Colegrave, and P. D. Keightley. 2016. Direct estimate of the
spontaneous mutation rate uncovers the effects of drift and recombination in the
Chlamydomonas reinhardtii plastid genome. Mol. Biol. Evol. **33:** 800-808.
doi:10.1093/molbev/msv272
Osuna-Cruz, C. M., G. Bilcke, E. Vancaester, S. De Decker, A. M. Bones, P. Winge, N. Poulsen
and others 2020. The *Seminavis robusta* genome provides insights into the evolutionary

adaptations of benthic diatoms. Nat. Commun. **11:** 3320. doi:10.1038/s41467-020-17191-
8

Paracchini, V., M. Petrillo, A. Lievens, A. P. Gallardo, J. T. Martinsohn, J. Hofherr, A. Maquet
and others 2017. Novel nuclear barcode regions for the identification of flatfish species.
Food Control **79:** 297-308. doi:10.1016/j.foodcont.2017.04.009

Pujari, L., C. Wu, J. Kan, N. Li, X. Wang, G. Zhang, X. Shang and others 2019. Diversity and
Spatial Distribution of Chromophytic Phytoplankton in the Bay of Bengal Revealed by
RuBisCO Genes (rbcL). Front. Microbiol. **10:** 1501. doi:10.3389/fmicb.2019.01501

Quinlan, A. R., and I. M. Hall. 2010. BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics **26:** 841-842. doi:10.1093/bioinformatics/btq033

Rastogi, A., F. R. J. Vieira, A.-F. Deton-Cabanillas, A. Veluchamy, C. Cantrel, G. Wang, P.
Vanormelingen and others 2020. A genomics approach reveals the global genetic
polymorphism, structure, and functional diversity of ten accessions of the marine model
diatom *Phaeodactylum tricornutum*. ISME J **14:** 347-363. doi:10.1038/s41396-019-0528-
3

Read, B. A., J. Kegel, M. J. Klute, A. Kuo, S. C. Lefebvre, F. Maumus, C. Mayer and others
2013. Pan genome of the phytoplankton *Emiliania* underpins its global distribution.
Nature **499:** 209-213. doi:10.1038/nature12221

Rengefors, K., R. Gollnisch, I. Sassenhagen, K. Harnstrom Aloisi, M. Svensson, K. Lebret, D.
Certnerova and others 2021. Genome-wide single nucleotide polymorphism markers
reveal population structure and dispersal direction of an expanding nuisance algal bloom
species. Mol. Ecol. **30:** 912-925. doi:10.1111/mec.15787

Rengefors, K., A. Kremp, T. B. Reusch, and A. M. Wood. 2017. Genetic diversity and evolution
in eukaryotic phytoplankton: revelations from population genetic studies. J. Plankton
Res. **39:** 165-179. doi:10.1093/plankt/fbw098

Rhoads, A., and K. F. Au. 2015. PacBio Sequencing and Its Applications. Genom. Proteom.
Bioinform. **13:** 278-289. doi:10.1016/j.gpb.2015.08.002

Robinson, D. G., W. Chen, J. D. Storey, and D. Gresham. 2014. Design and analysis of Bar-seq
experiments. G3-Genes Genom. Genet. **4:** 11-18. doi:10.1534/g3.113.008565

Robinson, J. T., H. Thorvaldsdóttir, W. Winckler, M. Guttman, E. S. Lander, G. Getz, and J. P.
Mesirov. 2011. Integrative genomics viewer. Nat. Biotechnol. **29:** 24-26.
doi:10.1038/nbt.1754

Rognes, T., T. Flouri, B. Nichols, C. Quince, and F. Mahé. 2016. VSEARCH: a versatile open
source tool for metagenomics. PeerJ **4:** e2584. doi:10.7717/peerj.2584

Sarno, D., W. H. Kooistra, L. K. Medlin, I. Percopo, and A. Zingone. 2005. Diversity in the
genus *Skeletonema* (Bacillariophyceae). II. An assessment of the taxonomy of *S.*
costatum□like species with the description of four new species 1. J. Phycol. **41:** 151-176.
doi:10.1111/j.1529-8817.2005.04067.x

Sassenhagen, I., D. Erdner, B. Lougheed, M. Richlen, and C. Sjöqvist. 2021. Estimating
proportion of clones and genotype richness in aquatic microalgae. Authorea Preprints.
doi:10.22541/au.161876383.32271114/v1

Schaum, C.-E., S. Barton, E. Bestion, A. Buckling, B. Garcia-Carreras, P. Lopez, C. Lowe and
others 2017. Adaptation of phytoplankton to a decade of experimental warming linked to
increased photosynthesis. Nat. Ecol. Evol. **1:** 1-7. doi:10.1038/s41559-017-0094

Schaum, C.-E., A. Buckling, N. Smirnoff, D. Studholme, and G. Yvon-Durocher. 2018.
Environmental fluctuations accelerate molecular evolution of thermal tolerance in a
marine diatom. Nat. Commun. **9:** 1719. doi:10.1038/s41467-018-03906-5

Scheinin, M., U. Riebesell, T. A. Rynearson, K. T. Lohbeck, and S. Collins. 2015. Experimental
evolution gone wild. J. R. Soc. Interface **12:** 20150056. doi:10.1098/rsif.2015.0056

Schloss, P. D., S. L. Westcott, T. Ryabin, J. R. Hall, M. Hartmann, E. B. Hollister, R. A.
Lesniewski and others 2009. Introducing mothur: open-source, platform-independent,
community-supported software for describing and comparing microbial communities.
Appl. Environ. Microbiol. **75:** 7537-7541. doi:10.1128/AEM.01541-09

Sefbom, J., I. Sassenhagen, K. Rengefors, and A. Godhe. 2015. Priority effects in a planktonic
bloom-forming marine diatom. Biol. Lett. **11:** 20150184. doi:10.1098/rsbl.2015.0184

Sjöqvist, C., A. Kremp, E. Lindehoff, U. Båmstedt, J. Egardt, S. Gross, M. Jönsson and others
2014. Effects of grazer presence on genetic structure of a phenotypically diverse diatom
population. Microb. Ecol. **67:** 83-95. doi:10.1007/s00248-013-0327-8

Smith, A. M., L. E. Heisler, J. Mellor, F. Kaper, M. J. Thompson, M. Chee, F. P. Roth and others
2009. Quantitative phenotyping via deep barcode sequencing. Genome Res. **19:** 1836-
1842. doi:10.1101/gr.093955.109

Smyth, R. P., T. E. Schlub, A. Grimm, V. Venturi, A. Chopra, S. Mallal, M. P. Davenport and
others 2010. Reducing chimera formation during PCR amplification to ensure accurate
genotyping. Gene **469:** 45-51. doi:10.1016/j.gene.2010.08.009

Stoeck, T., D. Bass, M. Nebel, R. Christen, M. D. Jones, H. W. Breiner, and T. A. Richards.
2010. Multiple marker parallel tag environmental DNA sequencing reveals a highly
complex eukaryotic community in marine anoxic water. Mol. Ecol. **19:** 21-31.
doi:10.1111/j.1365-294X.2009.04480.x

Straub, D., N. Blackwell, A. Langarica-Fuentes, A. Peltzer, S. Nahnsen, and S. Kleindienst.
2020. Interpretations of Environmental Microbial Community Studies Are Biased by the
Selected 16S rRNA (Gene) Amplicon Sequencing Pipeline. Front. Microbiol. **11**.
doi:10.3389/fmicb.2020.550420

Sundberg, C., W. A. Al-Soud, M. Larsson, E. Alm, S. S. Yekta, B. H. Svensson, S. J. Sørensen
and others 2013. 454 pyrosequencing analyses of bacterial and archaeal richness in 21
full-scale biogas digesters. FEMS Microbiol. Ecol. **85:** 612-626. doi:10.1111/1574-
6941.12148

Söderhielm, J., and K. Sundblad. 1996. The Solstad Cu☐Co☐Au mineralization and its relation
to post☐Svecofennian regional shear zones in southeastern Sweden. GFF **118:** 47-47.
doi:10.1080/11035899609546323

Taberlet, P., E. Coissac, F. Pompanon, C. Brochmann, and E. Willerslev. 2012. Towards
next☐generation biodiversity assessment using DNA metabarcoding. Mol. Ecol. **21:**
2045-2050. doi:10.1111/j.1365-294X.2012.05470.x

Tatters, A. O., M. Y. Roleda, A. Schnetzer, F. Fu, C. L. Hurd, P. W. Boyd, D. A. Caron and
others 2013. Short-and long-term conditioning of a temperate marine diatom community
to acidification and warming. Phil. Trans. R. Soc. B. **368:** 20120437.
doi:10.1098/rstb.2012.0437

Van der Auwera, G. A., and B. D. O'Connor. 2020. Genomics in the cloud: using Docker,
GATK, and WDL in Terra, 1st ed. Sebastopol, CA: O'Reilly Media Inc.

von Dassow, P., U. John, H. Ogata, I. Probert, M. Bendif el, J. U. Kegel, S. Audic and others
2015. Life-cycle modification in open oceans accounts for genome variability in a
cosmopolitan phytoplankton. ISME J **9:** 1365-1377. doi:10.1038/ismej.2014.221
Williams, R., S. G. Peisajovich, O. J. Miller, S. Magdassi, D. S. Tawfik, and A. D. Griffiths.
2006. Amplification of complex gene libraries by emulsion PCR. Nat. Methods **3:** 545-
550. doi:10.1038/nmeth896
Wolf, K. K., E. Romanelli, B. Rost, U. John, S. Collins, H. Weigand, and C. J. Hoppe. 2019.
Company matters: The presence of other genotypes alters traits and intraspecific selection
in an Arctic diatom under climate change. Global Change Biol. **25:** 2869-2884.
doi:10.1111/gcb.14675
Worden, A. Z., J.-H. Lee, T. Mock, P. Rouzé, M. P. Simmons, A. L. Aerts, A. E. Allen and
others 2009. Green evolution and dynamic adaptations revealed by genomes of the
marine picoeukaryotes *Micromonas*. Science **324:** 268-272. doi:10.1126/science.1167222
Yamada, M., M. Otsubo, Y. Tsutsumi, C. Mizota, Y. Nakamura, K. Takahashi, and M. Iwataki.
2017. Utility of mitochondrial☐encoded cytochrome c oxidase I gene for phylogenetic
analysis and species identification of the planktonic diatom genus *Skeletonema*. Phycol.
Res. **65:** 217-225. doi:10.1111/pre.12179
865
866

Table 1. Statistical parameters from Bamboozles genomic scan for barcode windows. Shown are

average (standard deviation) for the variable barcoding loci identified by Bamboozle (selected

windows), with regard to the number of predicted SNPs, indels, and unique alleles per locus, in

contrast to average statistics for the genome as a whole, obtained using a sliding-window

approach (other windows). Where two sets of values are given, the first gives statistics are for

those windows where only the coverage filter is applied, and the second (in square brackets)

gives statistics for those windows were both the coverage filter and the conserved end region

filter are applied.

	S. marinoi	*C. reinhardtii* Mating type +	Mating type -
Reference genome size (nuclear, mitochondrion, and plastid)	54,794,446 bp	107,613,365 bp	
Number of strains included	54	8	9
Length of genome conforming to coverage filtering	4,740,972	630,835	464,867
SNP density of selected windows	45.0 (9.13)[†]	17.4 (3.25)	22.86 (4.74)
SNP density of other windows	26.26 (14.53) [12.36 (10.83)]	5.30 (6.02) [2.17 (3.42)]	4.68 (5.87) [1.92 (2.95)]
Indel density of selected windows	2.75 (4.27)	1.47 (0.83)	1.71 (1.98)
Indel density of other windows	3.27 (4.55) [1.38 (2.76)]	0.53 (1.03) [0.27 (0.73)]	0.46 (0.98) [0.23 (0.67)]
Number of unique alleles in selected windows	100 (4.27)	8 (0)[††]	9 (0)[††]
Number of unique alleles in other windows	30.56 (20.41) [14.30 (13.89)]	2.87 (1.58) [2.08 (1.23)]	2.99 (1.85) [2.11 (1.31)]

876 [†]Note that in the case of *S. marinoi*, these statistics exclude *Sm_C2W24* (identified using an older version
877 of the pipeline) but include *Sm_C12W3* (identified in the more recent version).
878 [††]When comparing all *C. reinhardtii* selected windows among all 17 strains (i.e., not separated by mating
879 type), there are 12.55 (1.79) alleles per locus.
880

Table 2: Accuracy of Bamboozle's allele sequence predictions for selected barcodes in *S.*
marinoi and *C. reinhardtii*. 'Predictions' corresponds to Bamboozle's output sequences based on
WGS data, and 'observations' are confirmations based on targeted PCR-based sequencing. *C.*
reinhardtii loci with poor amplification and lack of Sanger sequencing data are shown as n/a.
Sm_C2W24 results marked with * indicate results from an earlier iteration of the pipeline, which
didn't take coverage or allele phasing into account, and was variant-called using BCFtools (cf.
GATK in the most recent iteration).

Skeletonema marinoi[†]	*Sm_C2W24*	*Sm_C12W1*	*Sm_C12W2*	*Sm_C16W4*
Gene model	Sm_t00004275-RA	Sm_t00009768-RA	Sm_t00009769-RA	Sm_t00008465-RA
Barcode length in reference (2 × alleles in ref. genome)	471 (471/471)	523 (523/523)	494 (494/494)	504 (501/504)
Predicted SNP positions	95*	37	44	58
Predicted indel positions	5*	2	0	9
Predicted unique alleles	56*	98	99	106
Observed SNP positions	19	38	44	54
Observed indel positions	198	0	0	18
Mean WGS sequencing depth (range, N=55)	49 (13-101)	60 (17-140)	59 (17-131)	63 (19-131)
Mean amplicon sequencing depth (range, N=55)	5,391 (5-35,434)	4,156 (15-10,934)	1,693 (5-6,960)	3,591 (6-19,143)
Accuracy of prediction across all strains and bases	82.65%	99.91%	99.93%	99.88%

Chlamydomonas reinhardtii[††]	mt+			mt-
	Cr_Chr1W3	Cr_Chr3W8	Cr_Chr3W10	Cr_Chr9W12
Gene model	CHLRE_01g040900v5	CHLRE_03g181100v5	CHLRE_03g207250v5	CHLRE_09g389134v5
Barcode length	298	297	299	290
Predicted SNP positions	21	23	28	32
Predicted indel positions	4	2	8	3
Observed SNP positions	n/a	n/a	27	32
Observed indel positions	n/a	n/a	8	3
Mean WGS coverage (range N=19)	29 (11-48)	37 (17-58)	32 (16-52)	33 (14-53)
Accuracy of prediction across all strains and bases	n/a	n/a	99.98	99.88

890 [†]One strain (GP2-4_32) was excluded from this analysis due to a failed sequencing.
891 [††]One *strain* (CC-2938) was excluded from this analysis due to failed DNA extraction.
892

Figure 1: Schematic summarising the
steps of the Bamboozle pipeline, as
well as input and output files. Each
step is described in more detail in the
New Approaches section.

Figure 2: Histograms showing the predicted number of unique alleles (panels A, C, and E), and

predicted frequency of SNPs and indels (panels B, D, and F), at each window of 500 bp (*S.*

marinoi) or 300 bp (*C. reinhardtii*). Shown are data from all genomic windows that passed

Bamboozle's filters for coverage depth and flanking region conservation, as described in the New

Approaches section. In the case of *S. marinoi*, only nuclear contigs are included. Panels A and B

depict figures from *S. marinoi*; panels C and D depict figures from *C. reinhardtii* mating type +;

panels E and F depict figures from *C. reinhardtii* mating type -. Arrows indicate the relative

locations of the selected barcoding loci in terms of predicted SNP (red) and indel (blue)

frequency, and unique alleles (green). Note that *Sm_C2W24* was predicted with an earlier

version of Bamboozle, resulting in deviant values compared to other loci.

908
909

Figure 3: Genomic location of the four barcoding loci identified in *S. marinoi* and selected four

loci in *C. reinhardtii,* corresponding to those that primers were designed and evaluated for (for

complete list of all loci, and brief gene model annotations, see Table S2). A and C) Gene models

with functional domains, primer sites, and variable regions in *S. marinoi* and *C. reinhardtii,*

respectively. B and D) Gel electrophoresis of PCR products in two strains of *S. marinoi* and one

strain each of mating type + and - in *C. reinhardtii,* of loci with good amplification results.

Abbreviations – SUMO: Small Ubiquitin-like Modifier; AN(E): the three amino acids Alanine,

Asparagine, and Glutamic acid [occasionally replaced by Glycine]; PPR: pentatricopeptide

[RNA binding]; YehB: Uncharacterised membrane protein [partial, UPF0754 family]; NLE:

NUC135 domain located in N terminal of WD40 repeats; WD40: domain involved in various

cellular function including signal transduction, pre-mRNA processing and cytoskeleton

assembly; DUF563: Domain of Unknown Function; EBNA-3B: PHA03378 superfamily domain

(located on the reverse strand). Glutamine synthetase is represented by its PLN02284

superfamily domain and overlap with EBNA-3B is shown in green. Note that the primer symbols

are not drawn to scale but the tip of each symbol indicates the exact location where they end and

the variable region begins.

Department of Biology, Lund University, Lund, Sweden

927
928

Figure 4: Metabarcoded relative abundance of strains alleles using the barcode loci *Sm_C12W1*.

Shown are two mixed populations of *S. marinoi*. A) a mixture of 28 strains from Gropviken

(GP), and B) 30 strains from Gåsfjärden (VG). Legend numbers indicate whether this is the first,

second, or third allele in each strain, with colours/asterisks indicating if alleles are homologous

in two (*) or three (**) other strains. For asterisk-marked alleles, the abundance has been

partitioned between strains using differential equations. The horizontal line indicates the

expected fraction of amplicons (allele 1+2) per strain, assuming even cell densities, Error bars

show standard deviation, with N=3 technical PCR replicates for panel A, and N=4 for B. Data

has been denoised using the settings for 'Relaxed DADA2 on amplicons' as outlined in the

Supplementary Information.

Figure 5: Correlations between heterozygous allele of 58 strains of *S. marinoi* across the 42-day

selection experiment. For visibility, data has been transformed (x +1) with removal of 0+0 cases.

Shared homologous alleles are indicated by alphabetical indexes in the corner of each allele (i.e.,

lower right equals allele one, upper left allele two), and their abundances have been allocated

between strains using differential equations. All such equations assume a 1:1 ratio between

alleles, except allele two in GP2-4_43, where a ratio of 0.15 is used based on the genotype

sample. The black line across panels corresponds to the expected 1:1 ratio between heterozygous

alleles. The blue line is a second order polynomial function fitted to the data, with shaded area

corresponding to 95% confidence interval. Data has been denoised using the settings for

'Relaxed DADA2 on amplicons' as outlined in the Supplemental Information. Asterisks

highlight false-positive observations that the denoising pipeline failed to remove. Fig. S4 shows

the same figure without denoising (i.e. 'exact matches' settings in Table S6).

Read count allele 1