1

Loss Of Chromodomain of Male-Specific Lethal 3 (MSL3) Does Not Affect Spermatogenesis in Rodents

Mitchell T.A.[1,2,3], Lin J.M. [1,2], Hicks S.M. [1,2,3], James J.R. [1,2], McCarthy A. [1,2], Rangan P.[4,5] and Forni P.E. [1,2,3]*

[1]Department of Biological Sciences, University at Albany, State University of New York, Albany, NY 12222, USA.

[2]The RNA Institute, University at Albany, State University of New York, Albany, NY 12222, USA.

[3]The Center for Neuroscience Research, University at Albany, State University of New York, Albany, NY 12222, USA.

[4] Black Family Stem Cell Institute, Department of Cell, Developmental, and Regenerative Biology, Icahn School of Medicine at Mount Sinai, 1 Gustave L. Levy Place, New York, NY 10029, USA

*Corresponding author: pforni@albany.edu

Abstract:

Male-specific lethal (MSL3) is a member of the Dosage Compensation Complex in *Drosophila*. It is required for transcriptional upregulation of genes on the X chromosome in males to equal that of females. Even though the dosage complex is carried out differently in mammals, Msl3 is conserved in humans. Intriguingly, Msl3 is expressed in the undifferentiated cells from Drosophila to humans, including in spermatogonia of macaque and humans. During *Drosophila* oogenesis, Msl3 is required for meiotic entry. However, its role in meiotic entry in other organisms has not been explored. Using mouse spermatogenesis as a model system, we probed for the role of Msl3 in the meiotic entry. We found that MSL3 is expressed in mouse testes in meiotic cells in contrast to flies, primates, and humans. Further, using a newly generated MSL3 conditional knock-out mouse line, we found no spermatogenesis defects within the seminiferous tubules of the KOs.MSL3 mutants were also viable and fertile, suggesting that MSL3 is dispensable for rodent gametogenesis.

Introduction:

Gametes ensure the survival and propagation of all sexually reproducing organisms. Gametes arise from germ cells that differentiate and undergo meiosis to produce haploid oocytes and sperms. Meiosis is essential for reductional cell division to generate haploid gametes and genetic diversity by promoting recombination. Thus, regulation of meiosis is critical for gametogenesis and fertility. However, the regulation of meiotic entry remains incompletely understood.

Mammalian spermatogenesis is an excellent model system to determine regulators of meiotic entry due to the presence of spermatogonial stem cells (SSCs) that self-renew and differentiate, undergoing meiosis throughout the animal's life [1,2]. This is in contrast to oogenesis, where germ cells directly differentiate into oocytes during embryonic development that then meiotically arrest [3]. Spermatogenesis in mouse initiates with SSC, are also called Single spermatogonia (A_s), which give rise to A paired spermatogonia (A_{paired}) and A aligned spermatogonia ($A_{aligned}$), which make chains of 4, 8 and 16 Type A spermatogonia which eventually gives rise to type B differentiated spermatogonia through mitotic divisions. Type B spermatogonia enter meiosis prophase I when they initiate recombination. These type B spermatogonia then progress to a second round of meiotic divisions to yield round spermatids. Round spermatids then go through spermiogenesis, in which morphological changes occur, including the trading of histones for protamines, to get elongated spermatids and eventually mature spermatozoa [4-6].

In mammals, the entry into meiosis is promoted by steroid signaling during oogenesis and spermatogenesis. Retinoic acid (RA) from somatic cells surrounding the spermatogonia activates the transcription factor *Stimulated by retinoic acid 8* (*Stra8*) in the germline. During spermatogenesis, Stra8 fosters meiotic entry by promoting the transcription of a broad gene expression program. However, *Stra8* is not sufficient to induce meiosis, suggesting a cell type-specific chromatin landscape and/or factors that cooperate with STRA8 [7].

Male-specific lethal 3 (MSL3) is a member of the Male-specific lethal (MSL) complex which is necessary for dosage compensation in *Drosophila*. The MSL complex promotes transcriptional upregulation of genes on the X chromosome in males to equal the two copies from the two X chromosomes in females. This transcriptional upregulation is essential to the survival of male *Drosophila*. The complex is made up of several components, including MSL1, MSL2, MSL3, male-absent on first (MOF), maleless (MLE), and two roX noncoding RNAs. MSL3's chromodomain has been shown to be important for binding to X-linked genes by recognizing Histone 3 lysine 36 trimethylation marks (H3K36me3). Conversely, the MRG domain is important for interaction with

the Dosage Compensation Complex [8-10]. Thus, Msl3 and its Chromodomain are critical for
Dosage compensation.

It was thought that some MSL complex components were not expressed in the females, thus
precluding the formation of the MSL complex. However, Msl3 is expressed in a temporal manner
during *Drosophila* oogenesis from the undifferentiated germline stem cells (GSC) to the
differentiating cysts stages when the germ cell enters meiosis. It was found to be part of a
transcriptional axis that promotes GSC differentiation and meiotic entry by recognizing H3K36me3
marks, presumably via its chromodomain. Intriguing, it has been reported that Msl3 is expressed
in undifferentiated spermatogonia of primates and humans, suggesting it could have a potential
function in the germline in mammals. In addition, large-scale analyses of the X chromosome in
infertile men discovered Msl3 to be one of the genes associated with spermatogenic failure. Taken
together, this suggested that Msl3 may be playing a role in the meiotic entry in mammals [11-13].

Here, using published single-cell (sc) RNA-sequencing data [14] we found that MSL3 is expressed
in the mouse testes but specifically in meiotic cells. This contrasts with flies, primates, and
humans, where MSL3 is expressed in the undifferentiated cells before meiotic entry [11-13]. By
generating a new MSL3 conditional knock-out consisting of loxP sites flanking the chromodomain
of the MSL3 gene that is critical for its function. We found that conditional loss of MSL3 during
mouse spermatogenesis is not sufficient to perturb spermatogenesis and fertility. In addition,
MSL3 mutants were also viable and fertile, suggesting that MSL3 is dispensable for
gametogenesis in general. Thus, unlike in Drosophila, Msl3 is dispensable for meiotic entry in
rodents.

Materials & Methods

Animals

Mice used in this study were as follows: MSL3 floxed, Stra8-iCre and R26tdTomato. All mice
described were maintained on a C57BL/6J background. The MSL3 floxed mouse line was
designed using CRISPR/Cas9 technology by Cyagen. The MSL3 gene consists of 13 exons.
Exons 2-4 of the chromodomain of MSL3 were targeted using CRISPR/Cas9. Lox*P* inserts flank
exons 2-4, which upon Cre recombinase activity results in their removal from the gene and
disruption of the chromodomain. The targeting vector was engineered with homology arms and
the conditional knock-out region generated by BAC clone RP24-103L17 from the C57BL/6N

library as a template. Cas9 mRNA was injected into fertilized eggs along with MSL3 gRNA and
the targeting vector. Founder mice were identified by PCR and bred to C57BL/6N to test germline
transmission. The Stra8-iCre (stock #017490) and R26tdTomato (stock #007914) mouse lines
are commercially available from JAX [15,16]. All animal experiments were conducted in
accordance with the guidelines of the Animal Care and Use Committee (IUACUC) at the
University at Albany, SUNY. Genotyping of the mouse lines was conducted by PCR using tail
genomic DNA. **iCre Primers:** *Fwd*-AGA TGC CAG GAC ATC AGG AAC CTG, *Rev*-ATC AGC
CAC ACC AGA CAC AGA GAT, *Fwd control*-CTA GGC CAC AGA ATT GAA AGA TCT, *Rev*
control-GTA GGT GGA AAT TCT AGC ATC ATC C. **MSL3^{flx} Primers:** *Fwd*-GAA TGC ATG GCT
ATT AAG GGG AGG, *Rev*-GAA GTC ATT GGT CGG TAG ATT TGT C. **MSL3 KO Primers:**
MSL3   WT   Fwd-GCAGTAAGCAGCACCTGTCAT,   *MSL3   WT   Rev*-
AGGAAAACTTGCAACATTGGTCT, *MSL3 KO Fwd*-TCCCGTCCAGAGGCTCTCAT, *MSL3 KO*
Rev-GGAAAACTTGCAACATTGGTCTGA.   **R26tdTomato   Primers:**   *IMR0920*-
AAGGGAGCTGCAGTGGAGTA,   *IMR09021*-CCGAAAATCTGTGGGAAGTC,   *IMR9103*-
GGCATTAAAGCAGCGTATCC, *IMR9105*-CTGTTCCTGTACGGCATGG.

Surface spreading and Immunofluorescence

Testes were excised from male mice euthanized at P17-20. Excised testes were placed in ice-
cold PBS. Testes were then weighed to determine incubation time in hypotonic extraction buffer
(HEB). **30 mg or less for 30 min, tubules from a pair of testes weighing 31 to 50 mg for 30 -**
45 minutes, and larger testes for 60 minutes. Testes were then placed back in ice cold PBS to
remove the tunica albuginea and extract the seminiferous tubules. Seminiferous tubules were
placed in HEB consisting of **30 mM Tris pH 8.2, 50 2 mM Sucrose, 17 mM Trisodium Citrate**
Dihydrate, 5 mM EDTA and incubated according to their weight. Seminiferous tubules were then
transferred back into a petri dish and separated into small clumps. Individual clumps were placed
in 60 ul droplets of 100 mM sucrose pH 8.2 and pipetted repeatedly until the solution became
turbid. Droplets of seminiferous tubules were placed in the corner of uncharged slides previously
soaked in a fixative solution of 1% PFA and 0.15% triton X-100 pH 9.2 with a droplet of fixative
remaining on the slide [17,18]. The droplet was allowed to spread along the slide then left to dry
overnight in a humidified chamber at 4°C. Slides were then stored at -80°C until use.

For immunofluorescence analysis, the spreading nuclei were immunostained with mouse anti-
SYCP3 (1:500), rabbit anti-SYCP1 (1:500) and mouse anti-γH2AX (1:500). Using SYCP3 to label
axial/lateral element characteristics and DAPI-stained heterochromatin pattern, spermatocytes in
spreading were classified as leptotene, zygotene, pachytene and diplotene.

Histological and immunohistochemical analysis

Testes were fixed in 4% formaldehyde embedded in OCT and sectioned, then stored at -80°C until use. For immunofluorescence analysis, the following primary antibodies were used: Rabbit anti-RFP (1:500 Rockland), Rabbit anti-Stra8 (1:250 ABCAM), mouse anti-SCP3 (1:500 ABCAM).

Testes weight and Seminiferous Tubule Diameter Measurements

Average weights were calculated by weighing testes (n=3 for each genotype) immediately after excision and averaged for each genotype. Average seminiferous tubule diameter measurements were calculated by measuring 10 seminiferous tubules per animal per genotype and calculating the average diameter of the 10 seminiferous tubules. This was repeated for a total of 3 animals per genotype. The average of the 3 averages was taken for each genotype as the representative weight.

RNA-sequencing

For whole RNA sequencing, testes were collected, and RNA extraction was performed using the Thermo Fisher RNA PureLink Mini Kit. Sequencing was performed using Illumina Nextseq 500.

For Single-cell RNA sequencing, we utilized previously published data by Green at al. [14] available through NCBI's Gene Expression Omnibus, GEO series accession number GSE112393 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE112393). Files were imported into the R package *Seurat 4.0.5* [19] for filtering. After filtering, mitochondrial genes and RNA features, individual files were merged into one file representing all germ cell types. Dimensionality reduction was performed using principal component analysis (PCA).

Gene Ontology Term Analysis

Gene Ontology term analysis was performed using the Panther Classification System.

Results

MSL3 is expressed in mouse testes during meiotic stages via ScSeq

To determine when MSL3 is expressed in mouse testes, we turned to single-cell RNA-sequencing (scRNA-seq) analysis. Using previously published drop-seq data by Green et al. [14], we reproduced a PCA clustering of testes germ cells. Markers used to identify spermatogonia were as follows: Zbtb16, Uchl1, Stra8, and Kit [20-23]. Markers used to identify spermatocytes were as

follows: Sycp3, Tex101, Spag6 and H2afx [24-27]. Markers used to identify round/elongating
spermatids were as follows: Acrv1, Tssk1, Tnp1, Prm1 [28-31] (Fig. 2A,B). As continuous
transitions occur between cell states during spermatogenesis, they are represented by the
changes in expression patterns of marker genes. (Fig. 1B). After identifying germ cell populations,
we probed for MSL3 expression. We found that MSL3 is expressed in spermatocytes and
continues through round spermatids (fig.1C). MSL3 expression correlates with spermatocyte
markers Tbpl1 and Spag6 confirming its expression in this cell population [32] (fig.1D). Thus, Msl3
is expressed in meiotic stages during mouse spermatogenesis.

iCre successfully recombines in Stra8+ cells

To test iCre recombination efficacy in the testes, we mated Stra8iCre female mice with R26tdTom
males and looked for tdTomato expression in Stra8-positive cells. We collected the testes of
R26tdTomato/Stra8iCre males at P30. Testes were fixed, sectioned, and immunostained for
tdTomato and Stra8. Immunostaining showed colocalization between Stra8+ spermatogonia and
tdTomato+ cells in Stra8iCre/R26tdtomato mice but not R26tdTomato controls (fig. 2). This means
tdTomato/Rosa26 is only being expressed in Stra8iCre spermatogonia and that iCre successfully
recombines in Stra8+ cells. By counting the number of Stra8+ and tdTomato+ cells, we conclude
that Stra8iCre has a 99.4% efficacy in the testes.

MSL3 mutant mice show normal testes morphology

To determine if MSL3 has a role in mouse spermatogenesis, we generated a conditional Knock
out of MSL3 by targeting exons 2-4 of the Msl3 chromodomain with *loxP* upstream of exon 2 and
downstream of exon 4 (Fig.3). Under the Stra8 promoter, iCre expression results in excision of
exons 2-4 of Msl3 resulting in a nonfunctional chromodomain.
We first examined whether MSL3cKO (Stra8iCre$^{+/-}$; MSL3$^{flox/flox}$) mice displayed any gross
testicular morphological defects. Testes from MSL3 cKO and control littermates were removed at
P45, when males are sexually mature, and all cell types should be present in the testes. By
dissecting the testes, we found no significant difference in size, shape, or weight of the testes
between controls (0.056 g +/- 0.003) and mutants (0.059 g +/- 0.002) (Fig. 4A,B).
Microscopic examination of testes sections showed no morphological defects between mutants
and controls. Moreover, we found no significant difference between the size of seminiferous
tubules as indicated by diameter measurements between controls (114.6um +/-4.6) and mutants
(115.0um +/-6.7) (Fig. 4C). Both mutants and controls had rounded seminiferous tubules with

similar cellular organization and all cell types present from primary spermatogonia to elongated
spermatids based on the morphology of cells present within the seminiferous tubules.

Formation of synaptonemal complex in the absence of functional MSL3

scRNA-Seq analysis showed that MSL3 is expressed in primary spermatocytes (Fig.1), and
previous work suggested that MSL3 is necessary for transcriptional upregulation of synaptonemal
complex proteins [10]. To determine if Msl3 is required for meiosis, we examined the meiotic
prophase-I using marker for synapsis that is promoted by synaptonemal complex proteins. We
prepared Meiotic chromosome spreads from both control and MSL3 cKO testes from P17-21 male
mutants. Using an antibody against SYCP3, one of the lateral elements of the synaptonemal
complex, we characterized primary spermatocytes as in leptonema, zygonema, pachynema or
diplonema by way of immunofluorescent staining. Chromosome spreads from both controls and
MSL3 cKO showed spermatocytes at all stages of prophase I, suggesting that both genotypes
can complete prophase-I. Counts of 100 spermatocytes in both mutants and controls also showed
that there were no significant differences between the distribution of spermatocytes between
genotypes. (Fig. 5). This was determined by the change in localization of SYCP3 from a more
diffuse/widespread expression in leptonema to a more condensed expression with strong puncta
at the centromeric region in diplonema. Taken together this suggested that MSL3 is not required
for the formation of the synaptonemal complex.

MSL3 mutants are fertile and produce normal litter sizes

Histological analysis of MSL3 cKO seminiferous tubules showed the presence of all cell types,
from primary spermatogonia to elongated spermatids. The presence of elongated spermatids
means cells can initiate spermiogenesis or the differentiation of spermatids leading to mature
spermatozoa. MSL3 cKO males were thus mated with WT females to examine their fertility along
with littermate controls. Each MSL3 cKO male was paired with 2 C57B/6J females and allowed
to mate for 2 weeks. MSL3 cKO males successfully impregnated both females and produced
litters of comparable sizes to that of controls. MSL3 cKO offspring were also viable (Table 1).
Thus, we conclude that MSL3 is not necessary for spermatogenesis.

Loss of Msl3 shows no gross disruptions in the transcriptome

As MSL3 mutants show no defects in the spermatogenic process, we asked if upregulation of
other genes may compensate for a potential function of MSL3. To do this, we performed whole
RNA-sequencing on MSL3 cKO mutants and wildtype controls (n=3 for each genotype). We
performed differential gene expression (DGE) analysis on the sequenced testes to check for any
differentially expressed genes between MSL3 cKOs and controls.

Read counts for MSL3 between genotypes were visualized using the Interactive Genomics
Viewer. We found few reads that mapped for exons 2-4 in our MSL3 cKOs (Fig. 6A) consistent
with the deletion of the chromo domain. Notably few read counts were observed in our cKOs,
these may reflect MSL3 expression in Sertoli cells or in some spermatogonial cells that escaped
recombination.

Differential expression analysis between control and mutant samples reveled the upregulation of
seven genes and downregulation of 44 genes in MSL3 cKOs (Fig. 6B) (significant genes were
determined by $\log_2$ fold change >0.5 and <-0.5 and a p-adj. value <0.05). Of the 7 upregulated
genes, 3 of them with the highest $\log_2$ fold change (0.5 or greater) (Cyp2c55, A1CF, Pyroxd2)
have enzymatic and RNA metabolic functions; however, there is not known function for these
genes in the testes or spermatogenesis specifically. Of the 44 significantly down-regulated genes,
there were 18 with a $\log_2$ fold change of -0.5 or lower. GO term analysis on genes with a $\log_2$
fold change of -0.5 or lower and 0.5 or greater did not reveal convergence on any commonly
known pathway. Thus, based on our analysis, the absence of functional chromodomain in MSL3
does not alter the meiotic program during mouse spermatogenesis.

Discussion

Our efforts to uncover a functional role of Msl3 in murine spermatogenesis revealed it's
dispensability in this context. Notably, there are expression pattern differences between rodents
and primates and humans. In humans, Msl3 was shown to be a marker of spermatogonial stem
cells (SSC), particularly State 0 SSCs, in fetal-infant testes [12]. Similarly, in macaque monkeys,
Msl3 is expressed in Spermatogonia state 1 (SPG1) [13]. Our study suggests that, in mice, Msl3
expression is mostly restricted to the spermatocyte stage and only in a small portion of
spermatogonia (Figure 1C). In this study, we ablated Msl3 downstream of Stra8, which marks the

differentiation phase of spermatogonia; we cannot exclude that Msl3 may be potentially critical
prior to the differentiation of spermatogonia.

While MSL3 is important for transcriptionally upregulating synaptonemal complex components
during *Drosophila* oogenesis, it does not appear to have any effect on synaptonemal complex
formation during mouse spermatogenesis. Using MSL3 conditional knock-out mouse model, we
observed no noticeable defects or interruptions in spermatogenesis. The MSL3 gene consists of
a chromodomain and an MRG domain. Our conditional knockout has exons 2-4 of the
chromodomain removed upon Cre recombinase activity. It is possible that the two domains serve
redundant functions resulting in the lack of any observable phenotype in our mouse models. The
MRG gene family, consisting of MRG15 and MRGX, is involved in cellular proliferation and cycle
progression [33]. In *Drosophila*, the MRG domain mediates MSL3's incorporation into the Dosage
Compensation Complex [34]. The chromatin-binding protein MORF-related gene on chromosome
15 (MRG15) is involved in pre-mRNA splicing at the round spermatid stage [35]. The diverse
involvement of MRG genes leaves open this possibility of a role of the MRG domain in MSL3
function. It may be responsible for chromatin binding and HAT activity in transcriptionally active
genes. However, our current mouse model does not allow us to explore this idea. To further
confirm that MSL3is a dispensable gene for rodent spermatogenesis and rule out a redundant
function between the two domains, it would be beneficial to design a new MSL3 mouse model
with a functionally inactive chromodomain and MRG domains

Funding: P.E.F. is funded by NIDCD (R01 DC017149) (PEF) and NICHD (R01 HD097331);
P.R. is funded by NIH/NIGMS (RO1GM11177 and RO1GM135628)

273
274
275

References

1. Clermont Y. Kinetics of spermatogenesis in mammals: seminiferous epithelium cycle and spermatogonial renewal. Physiol Rev. 1972 Jan;52(1):198-236. doi: 10.1152/physrev.1972.52.1.198. PMID: 4621362.

2. de Rooij DG. Proliferation and differentiation of spermatogonial stem cells. Reproduction. 2001 Mar;121(3):347-54. doi: 10.1530/rep.0.1210347. PMID: 11226060.

3. Arur S. (2017). Signaling-Mediated Regulation of Meiotic Prophase I and Transition During Oogenesis. *Results and problems in cell differentiation*, *59*, 101–123. https://doi.org/10.1007/978-3-319-44820-6_4

4. de Rooij DG, Russell LD. All you wanted to know about spermatogonia but were afraid to ask. J Androl. 2000 Nov-Dec;21(6):776-98. PMID: 11105904.

5. Zickler D, Kleckner N. Meiotic chromosomes: integrating structure and function. Annu Rev Genet. 1999;33:603-754. doi: 10.1146/annurev.genet.33.1.603. PMID: 10690419.

6. Govin J, Caron C, Lestrat C, Rousseaux S, Khochbin S. The role of histones in chromatin remodelling during mammalian spermiogenesis. Eur J Biochem. 2004 Sep;271(17):3459-69. doi: 10.1111/j.1432-1033.2004.04266.x. PMID: 15317581.

7. Endo, T., Mikedis, M. M., Nicholls, P. K., Page, D. C., & de Rooij, D. G. (2019). Retinoic Acid and Germ Cell Development in the Ovary and Testis. *Biomolecules*, *9*(12), 775. https://doi.org/10.3390/biom9120775

8. Lucchesi, J. C., & Kuroda, M. I. (2015). Dosage compensation in Drosophila. *Cold Spring Harbor perspectives in biology*, 7(5), a019398. https://doi.org/10.1101/cshperspect.a019398

9. Sural, T. H., Peng, S., Li, B., Workman, J. L., Park, P. J., & Kuroda, M. I. (2008). The MSL3 chromodomain directs a key targeting step for dosage compensation of the Drosophila melanogaster X chromosome. *Nature structural & molecular biology*, *15*(12), 1318–1325. https://doi.org/10.1038/nsmb.1520

10. McCarthy, A., Sarkar, K., Martin, E. T., Upadhyay, M., Jang, S., Williams, N. D., Forni, P. E., Buszczak, M., & Rangan, P. (2021). MSL3 promotes germline stem cell differentiation in female Drosophila. *Development (Cambridge, England)*, dev.199625. Advance online publication. https://doi.org/10.1242/dev.199625

11. Guo, J., Grow, E. J., Mlcochova, H., Maher, G. J., Lindskog, C., Nie, X., Guo, Y., Takei,
Y., Yun, J., Cai, L., Kim, R., Carrell, D. T., Goriely, A., Hotaling, J. M., & Cairns, B. R.
(2018). The adult human testis transcriptional cell atlas. *Cell research*, *28*(12), 1141–
1157. https://doi.org/10.1038/s41422-018-0099-2
12. Guo, J., Sosa, E., Chitiashvili, T., Nie, X., Rojas, E. J., Oliver, E., DonorConnect, Plath,
K., Hotaling, J. M., Stukenborg, J. B., Clark, A. T., & Cairns, B. R. (2021). Single-cell
analysis of the developing human testis reveals somatic niche cell specification and fetal
germline stem cell establishment. *Cell stem cell*, *28*(4), 764–778.e4.
https://doi.org/10.1016/j.stem.2020.12.004
13. Shami AN, Zheng X, Munyoki SK, Ma Q, Manske GL, Green CD, Sukhwani M, Orwig
KE, Li JZ, Hammoud SS. Single-Cell RNA Sequencing of Human, Macaque, and Mouse
Testes Uncovers Conserved and Divergent Features of Mammalian Spermatogenesis.
Dev Cell. 2020 Aug 24;54(4):529-547.e12. doi: 10.1016/j.devcel.2020.05.010. Epub
2020 Jun 5. PMID: 32504559; PMCID: PMC7879256.
14. Green, C. D., Ma, Q., Manske, G. L., Shami, A. N., Zheng, X., Marini, S., Moritz, L., Sultan,
C., Gurczynski, S. J., Moore, B. B., Tallquist, M. D., Li, J. Z., & Hammoud, S. S. (2018). A
Comprehensive Roadmap of Murine Spermatogenesis Defined by Single-Cell RNA-
Seq. *Developmental   cell*, *46*(5),   651–667.e10.
https://doi.org/10.1016/j.devcel.2018.07.025
15. Sadate-Ngatchou, P. I., Payne, C. J., Dearth, A. T., & Braun, R. E. (2008). Cre
recombinase activity specific to postnatal, premeiotic male germ cells in transgenic
mice. *Genesis (New York, N.Y. : 2000)*, *46*(12), 738–742.
https://doi.org/10.1002/dvg.20437
16. Madisen, L., Zwingman, T. A., Sunkin, S. M., Oh, S. W., Zariwala, H. A., Gu, H., Ng, L.
L., Palmiter, R. D., Hawrylycz, M. J., Jones, A. R., Lein, E. S., & Zeng, H. (2010). A
robust and high-throughput Cre reporting and characterization system for the whole
mouse brain. *Nature neuroscience*, *13*(1), 133–140. https://doi.org/10.1038/nn.2467
17. Dia, F., Strange, T., Liang, J., Hamilton, J., & Berkowitz, K. M. (2017). Preparation of
Meiotic Chromosome Spreads from Mouse Spermatocytes. *Journal of visualized
experiments : JoVE*, (129), 55378. https://doi.org/10.3791/55378
18. Gopinathan, L., Szmyd, R., Low, D., Diril, M. K., Chang, H. Y., Coppola, V., Liu, K.,
Tessarollo, L., Guccione, E., van Pelt, A., & Kaldis, P. (2017). Emi2 Is Essential for
Mouse Spermatogenesis. *Cell reports*, *20*(3), 697–708.
https://doi.org/10.1016/j.celrep.2017.06.033

19. Satija, R., Farrell, J. A., Gennert, D., Schier, A. F., & Regev, A. (2015). Spatial reconstruction of single-cell gene expression data. *Nature biotechnology*, *33*(5), 495–502. https://doi.org/10.1038/nbt.3192

20. Kwon, J., Wang, Y. L., Setsuie, R., Sekiguchi, S., Sakurai, M., Sato, Y., Lee, W. W., Ishii, Y., Kyuwa, S., Noda, M., Wada, K., & Yoshikawa, Y. (2004). Developmental regulation of ubiquitin C-terminal hydrolase isozyme expression during spermatogenesis in mice. *Biology of reproduction*, *71*(2), 515–521. https://doi.org/10.1095/biolreprod.104.027565

21. Suliman, B. A., Xu, D., & Williams, B. R. (2012). The promyelocytic leukemia zinc finger protein: two decades of molecular oncology. *Frontiers in oncology*, *2*, 74. https://doi.org/10.3389/fonc.2012.00074

22. Oulad-Abdelghani, M., Bouillet, P., Décimo, D., Gansmuller, A., Heyberger, S., Dollé, P., Bronner, S., Lutz, Y., & Chambon, P. (1996). Characterization of a premeiotic germ cell-specific cytoplasmic protein encoded by Stra8, a novel retinoic acid-responsive gene. *The Journal of cell biology*, *135*(2), 469–477. https://doi.org/10.1083/jcb.135.2.469

23. Yoshinaga, K., Nishikawa, S., Ogawa, M., Hayashi, S., Kunisada, T., Fujimoto, T., & Nishikawa, S. (1991). Role of c-kit in mouse spermatogenesis: identification of spermatogonia as a specific site of c-kit expression and function. *Development (Cambridge, England)*, *113*(2), 689–699. https://doi.org/10.1242/dev.113.2.689

24. Yuan, L., Liu, J. G., Zhao, J., Brundell, E., Daneholt, B., & Höög, C. (2000). The murine SCP3 gene is required for synaptonemal complex assembly, chromosome synapsis, and male fertility. *Molecular cell*, *5*(1), 73–83. https://doi.org/10.1016/s1097-2765(00)80404-9

25. Takayama, T., Mishima, T., Mori, M., Jin, H., Tsukamoto, H., Takahashi, K., Takizawa, T., Kinoshita, K., Suzuki, M., Sato, I., Matsubara, S., Araki, Y., & Takizawa, T. (2005). Sexually dimorphic expression of the novel germ cell antigen TEX101 during mouse gonad development. *Biology of reproduction*, *72*(6), 1315–1323. https://doi.org/10.1095/biolreprod.104.038810

26. Liu, Y., Zhang, L., Li, W., Huang, Q., Yuan, S., Li, Y., Liu, J., Zhang, S., Pin, G., Song, S., Ray, P. F., Arnoult, C., Cho, C., Garcia-Reyes, B., Knippschild, U., Strauss, J. F., & Zhang, Z. (2019). The sperm-associated antigen 6 interactome and its role in spermatogenesis. *Reproduction (Cambridge, England)*, *158*(2), 181–197. https://doi.org/10.1530/REP-18-0522

27. Hamer, G., Roepers-Gajadien, H. L., van Duyn-Goedhart, A., Gademan, I. S., Kal, H. B., van Buul, P. P., & de Rooij, D. G. (2003). DNA double-strand breaks and gamma-H2AX

signaling in the testis. *Biology of reproduction*, *68*(2), 628–634.
https://doi.org/10.1095/biolreprod.102.008672

28. Reddi, P. P., Flickinger, C. J., & Herr, J. C. (1999). Round spermatid-specific
transcription of the mouse SP-10 gene is mediated by a 294-base pair proximal
promoter. *Biology of reproduction*, *61*(5), 1256–1266.
https://doi.org/10.1095/biolreprod61.5.1256

29. Shang, P., Baarends, W. M., Hoogerbrugge, J., Ooms, M. P., van Cappellen, W. A., de
Jong, A. A., Dohle, G. R., van Eenennaam, H., Gossen, J. A., & Grootegoed, J. A.
(2010). Functional transformation of the chromatoid body in mouse spermatids requires
testis-specific serine/threonine kinases. *Journal of cell science*, *123*(Pt 3), 331–339.
https://doi.org/10.1242/jcs.059949

30. Meistrich, M. L., Mohapatra, B., Shirley, C. R., & Zhao, M. (2003). Roles of transition
nuclear proteins in spermiogenesis. *Chromosoma*, *111*(8), 483–488.
https://doi.org/10.1007/s00412-002-0227-z

31. Matsumoto, M., Kurata, S., Fujimoto, H., & Hoshi, M. (1993). Haploid specific activations
of protamine 1 and hsc70t genes in mouse spermatogenesis. *Biochimica et biophysica
acta*, *1174*(3), 274–278. https://doi.org/10.1016/0167-4781(93)90196-k

32. Zhang, D., Penttila, T. L., Morris, P. L., & Roeder, R. G. (2001). Cell- and stage-specific
high-level expression of TBP-related factor 2 (TRF2) during mouse
spermatogenesis. *Mechanisms of development*, *106*(1-2), 203–205.
https://doi.org/10.1016/s0925-4773(01)00439-7

33. Pena, A. N., & Pereira-Smith, O. M. (2007). The role of the MORF/MRG family of genes
in cell growth, differentiation, DNA repair, and thereby aging. *Annals of the New York
Academy of Sciences*, *1100*, 299–305. https://doi.org/10.1196/annals.1395.031

34. Morales, V., Regnard, C., Izzo, A., Vetter, I., & Becker, P. B. (2005). The MRG domain
mediates the functional integration of MSL3 into the dosage compensation
complex. *Molecular and cellular biology*, *25*(14), 5947–5954.
https://doi.org/10.1128/MCB.25.14.5947-5954.2005

35. Iwamori, N., Tominaga, K., Sato, T., Riehle, K., Iwamori, T., Ohkawa, Y., Coarfa, C., Ono,
E., & Matzuk, M. M. (2016). MRG15 is required for pre-mRNA splicing and
spermatogenesis. *Proceedings of the National Academy of Sciences of the United States
of America*, *113*(37), E5408–E5415. https://doi.org/10.1073/pnas.1611995113

Figure 1. Single-cell RNA sequencing of mouse testes. A) PCA plot of different germ cell population markers for spermatogonia, spermatocytes and round/elongating spermatids to identify germ cell type of each cluster. B) Clusters grouped into spermatogonia, transition states, spermatocytes and spermatids based on the marker genes they expressed to establish trajectory of spermatogenesis. C) MSL3 is expressed in spermatocytes and continues into spermatids then ceases as spermatids transition into elongating spermatids. D) Correlation map of different marker genes showing that MSL3 expression most closely correlates with that of Tbpl1 and Spag6 which are markers of spermatocytes.

Figure 2. Testes Immunofluorescence staining for Stra8 and tdTomato. Stra8 in green, tdTomato in red. Immunofluorescence staining shows that in Stra8iCre+/-R26tdTomato+/- mutants, stra8 and tdTomato staining colocalize, while in tdTomato+/- controls, Stra8 positive cells are negative for tdTomato. This confirms that tdTomato is only expressed in Stra8iCre cells. Scale bar = 50μm.

437 **Figure 3.** Design strategy adopted or MSL3 conditional knock-out mouse line generation.

438 Crispr/Cas9 was used to insert loxP sites upstream of exon 2 and downstream of exon 4 of the

439 MSL3 chromodomain. Upon Cre recombinase activity, exons 2-4 are excised from the MSL3 gene

440 generating a constitutive knockout allele.

448   **A.) Figure 4.** Image of MSL3 wildtype (WT) (left) and MSL3 cKO (right) testes at P45. Image shows

449   that they are of similar size and shape. B.) Bar graph showing the weight of pairs of testes for

450   MSL3 WT and MSL3 cKO mice. C.) Bar graph showing the average diameter of the seminiferous

451   tubules for MSL3 WT and MSL3 cKO mice. Average diameter of 10 seminiferous tubules for

452   each genotype is represented by each dot. n=3 for each genotype

461 **Figure 5. A.) Meiotic chromosome spread of spermatocytes** from P17-21 MSL3 WT and cKO testes

462 stained with antibody against SYCP3. Meiosis prophase I was examined between mutants and controls.

463 Spermatocytes were characterized as in leptonema, zygonema, pachynema or diplonema. Scale

464 bar=25um B.) 209 spermatocytes were counted for WT controls and 195 for Msl3 cKO. Subphases

465 of meiosis prophase I show no significant differences between MSL3 WT and MSL3 cKO.

Table 1

Genotype	# of Breeders	# of Litters	Avg. litter size
MSL3 WT	3	6	8.5 ± 1.05
MSL3 cKO	3	6	8.17 ± 1

Table 1.) MSL3 cKO males and littermate controls were mated with 2 C57B/6 WT females each. Males
were given 2 weeks to impregnate females and removed at the conclusion of the 2 week period. All males
successfully impregnated respective female. The average litter size of MSL3 cKO males (8.17± 1) was not
significantly different from that of littermate MSL3 WT controls (8.5± 1.05) (p=0.5826).

Figure 6 A) Read counts between MSL3 WT and MSL3 cKO show little to no reads for exons 2-4,

validating their excision from the gene in our cKO model. B.) Volcano plot showing significantly

upregulated and downregulated genes in our MSL3 cKO model.

487