Chemodiversity affects preference for _Tanacetum vulgare_ chemotypes in two aphid

species

Annika Neuhaus-Harr[1*], Lina Ojeda-Prieto[1*], Elisabeth Eilers[2,3], Caroline Müller[2], Wolfgang

W. Weisser[1], Robin Heinen[1]

[1]Technical University of Munich, Department of Life Science Systems, School of Life

Sciences, D-85354 Freising

[2]Department of Chemical Ecology, Bielefeld University, 33615 Bielefeld, Germany

[3]CTL GmbH Bielefeld, Krackser Str. 12, 33659 Bielefeld, Germany

*These authors share first authorship and contributed equally to the study.

Author for correspondence: Annika Neuhaus-Harr annika.neuhaus@tum.de

Abstract

Plants of the same species often strongly differ in morphological traits, as well as in the

abundance and composition of specialized metabolite profiles. Specialized metabolites can

act as mediators of interactions on plants, and partially explain insect presence and

abundance in the field. However, how specialized chemistry shapes plant attractiveness to

herbivorous insects is not fully understood. Here we used common tansy (_Tanacetum_

vulgare L., Asteraceae) – a perennial plant that is highly diverse in terpenoid composition

and is known to have variable chemotypes – to test whether 1) plants with different

chemotype profiles differ in attractiveness to two specialized aphids, _Macrosiphoniella_

tanacetaria and _Uroleucon tanaceti_, in pairwise choice assays. Furthermore, we tested

whether 2) the diversity of the terpenoid blend affects aphid attractiveness. Lastly, we tested

how 3) plant chemical traits relate to plant morphological traits, and which best explain aphid

preference. We found that *M. tanacetaria* preferred two out of five chemotypes, dominated by

α-thujone/β-thujone and β-trans-chrysanthenyl acetate, respectively, while avoiding a

chemotype dominated by α-pinene/sabinene. *U. tanaceti* showed no clear preference

towards chemotypes, but when given the choice between chemotypes dominated by α-

thujone/β-thujone and by α-pinene/sabinene, they preferred the former. Importantly, plant

attractiveness to aphids tended to be negatively correlated chemodiversity, i.e. the number of

terpenoid compounds, in *M. tanacetaria*, but not in *U. tanaceti*. Interestingly, the approximate

concentration and number of terpenoid compounds was generally higher in larger and

bushier plants. Hence, we did not observe a trade-off between plant growth and defence. We

conclude that plant chemical composition affects plant attractiveness to aphids and hence

may contribute to variation in natural aphid colonization patterns on plants of the same

species.

Keywords: aphids; attractiveness; choice assays; genotypic variation; intraspecific

phytochemical diversity; plant diversity; terpenes; terpenoids

Introduction

Understanding relationships between plants and herbivores is an important goal in ecology. How variation in plant diversity shapes herbivory has been a subject of study for many decades (Scherber et al. 2010, Weisser et al. 2017). For a long time between-species diversity was believed to be more important than within-species diversity as a driver of ecosystem processes (Des Roches et al. 2018). Intraspecific diversity - which includes the richness and abundance of genotypes and phenotypes within a species - has only more recently gained attention as an important driver of ecological processes (Raffard et al. 2019). It has been shown that intraspecific variation in plant geno- and phenotypes can structure food webs, such as arthropod-plant interactions (Crutsinger et al. 2006, Poelman et al. 2008, Bálint et al. 2016). As plants of the same species can differ strongly in various traits, understanding how this variation contributes to shaping interactions between plants and interaction partners is currently an important goal in plant ecology.

One important dimension of intraspecific variation in plants is chemodiversity; the variation in chemical composition between individuals of plants of the same species (Wetzel & Whitehead 2020). Primary metabolites are important as the building blocks of plants, and these compounds are comparatively similar across the plant kingdom (Weng, Philippe & Noel 2012). Specialized metabolites on the other hand, play critical roles as mediators of interactions and plants have evolved a staggering inter- and intraspecific diversity in these specialized compounds (Pichersky & Gang 2000; Iason, Dicke & Hartley 2012). Such chemical compounds can either be stored, emitted, or emitted after a stress event (Clancy et al. 2016; Holopainen and Gershenzon 2010). Intraspecific variation in specialized metabolite profiles is known to affect the structure of associated herbivore communities including phloem-feeding insects such as aphids (Schmitz 2008, Poelman et al. 2008, Richards et al. 2015, Bálint et al. 2016, Bustos-Segura et al. 2017, Salazar et al. 2018, Volf et al. 2019, Singh et al. 2021, Whitehead et al. 2021) , but few studies have disentangled the role of

chemodiversity on the forces that structure herbivore abundance and community structure in

natural conditions, such as plant attractiveness, bottom-up and top-down forces.

Chemodiversity may mediate presence and abundance of insect herbivores on a plant in the

field, which likely results as a combination of i) attractiveness of the plant for the insect

herbivore, ii) plant quality (i.e., bottom-up effects), and iii) the influence of higher trophic

levels on the insect herbivore through predation (i.e., top-down effects). Many specialized

compounds likely evolved to deter and repel herbivores (Herms & Mattson 1992, Kessler &

Baldwin 2001, Whitehead et al. 2021), but particularly specialized herbivore species may use

particular compounds as host-finding cues (Nishida 2014, Wink 2018). As such, insects can

be repelled or attracted, particularly by emitted volatile organic compounds, or VOCs (Clancy

et al. 2016, Jakobs & Müller 2019). Once an insect herbivore has arrived on the plant, stored

and emitted compounds can affect herbivore performance, for instance by feeding

deterrence, influencing metabolic processes or attracting herbivore enemies, but these

processes largely depend on metabolite-specific modes of action (Unsicker et al. 2009,

McCormick et al. 2012, Jakobs, Schweiger & Müller 2019). Emission of VOCs can also

attract natural enemies of herbivores, thereby acting as an indirect layer of defense for the

plant via top-down regulation (Dicke 2009, Unsicker et al. 2009, McCormick et al. 2012,

Rzanny et al. 2013). As these three processes act on plant individuals simultaneously, their

independent weights cannot reliably be inferred from natural colonization data collected from

the field. It is therefore important to understand how intraspecific chemical variation mediates

processes driving herbivore preference and performance in a controlled environment.

Besides intraspecific variation in chemical traits, individual plants of the same species may

also differ strongly in morphological traits. Various studies point out that traits related to

growth or to structural defenses may play an important role in driving interactions between

plants and insects (Herms & Mattson 1992; Agrawal & Fischbein 2006). For instance, in a

study on wheat plants, Batyrshina et al. (2020) found higher numbers of aphids on fast-

maturing than on slow-maturing wheat plants, and trade-offs between plant growth and

defense are common in nature (Coley et al. 1985; Herms & Mattson 1992). Furthermore,

plants with stronger mechanical defenses tend to be better defended against insect herbivory

(Caldwell, Read & Samson 2015). It is plausible that chemical traits will be linked with other

morphological traits, together driving insect preference. For instance, in common tansy,

(*Tanacetum vulgare* L., Asteraceae), plants with a higher storage of camphor were found to

have taller shoots than those with lower amounts, while plants containing davadone-D or

artemisia ketone developed more flower heads, taller corymbs, and delayed flowering

compared to plants with a lower content of these compounds (Keskitalo et al. 2001).

Furthermore, *T. vulgare* from different origins (e.g. North America and Europe) have been

found to differ in morphological and chemical traits, and exhibit negative correlations

between reproductive biomass and terpene concentrations (Wolf et al. 2011). Moreover,

Hayashi, Tahara and Ohgushi (2005) found that pubescence density and leaf mass per area

differed between the two chemotypes of *Salix sachalinensis*. The interplay of morphological

and chemical plant traits are plausible drivers of insect communities, but their relative

contribution as drivers is not fully understood.

Tanacetum vulgare is a perennial, aromatic plant that has a large geographic distribution and

is associated with a complex herbivore community including mono-, oligo- and polyphagous

aphids (Schmitz 1998, Keskitalo et al. 2001, Kleine & Müller 2011). Tansy is rich in different

mono- and sesquiterpenoids and plants can be divided into chemotype groups based on the

composition of terpenoids (Keskitalo et al. 2001, Kleine & Müller 2011). Previous studies and

breeding experiments showed that terpenoid composition has a genetic basis in tansy

(Keskitalo et al. 2001). Specialized aphids are thought to be adapted to harmful metabolites

in tansy and to use emitted volatiles for finding host plants (Schoonhoven et al. 2005, Jakobs

& Müller 2019). Aphid colonization, growth rate and survival, and even the structure of aphid

genotypes were found to be partly explained by tansy chemotypes under natural colonization

(Senft et al. 2017, 2019, Zytynska et al. 2019, Clancy et al. 2018). Specialized aphids show

preferences towards specific chemotypes (Jakobs & Müller 2018), but whether this is driven

by chemotype composition or concentration of compounds when several chemotypes are on

offer, still needs investigation.

In this study, we used six tansy chemotypes to investigate how differences between plant

individuals in chemical traits shapes attractiveness to the specialized aphids

Macrosiphoniella tanacetaria and *Uroleucon tanaceti* in factorial pairwise choice-assays

under lab conditions. We hypothesized that 1) the two aphid species would show species-

specific attraction to distinct tansy chemotypes, as previously shown by Jakobs and Müller

(2018, 2019) for two chemotypes. Furthermore, we hypothesized that 2) chemodiversity

negatively correlates with attractiveness, under the assumption that most specialized

metabolites repel antagonistic organisms. We specifically investigated whether approximate

concentrations of individual compounds predicted attractiveness. We also investigated how

chemical composition relates to morphological traits. We hypothesized that 3) growth-related

traits would trade-off with chemodiversity, under the assumption that maintaining

chemodiversity is costly and limits available resources for growth. We specifically explored

whether chemical or morphological traits are more important in driving insect preference

towards specific plant chemotypes.

Materials and methods

Chemotypic characterization of tansy lines

In 2019, leaf and seed samples of 27 tansy plants (hereafter: mothers) were collected in

Jena, Germany (50.93°N, 11.58°E), and chemotyped based on their terpenoid profiles (see

Chemical analysis of plant material). Unsupervised hierarchical clustering analysis on the

mother leaf terpenoid profiles clustered the plants in six chemotype clusters. Two mothers

were randomly selected per cluster and the collected seeds were mass-sown in seedling

trays in November 2020. Three weeks later, ten healthy seedlings were selected from each

mother plant lineage and transplanted to 10-cm pots filled with standard potting substrate

(Stender potting substrate C 700 coarse structure, 1 kg NPK minerals/m³, pH 5.5–6.0) and

transferred to a greenhouse compartment at the greenhouse facilities at TUM Plant

Technology Centre (with 16h:8h L:D and supplemental lighting that was turned off when

outdoor light was > 40 klx). In total, this resulted in a set of 120 plants (hereafter: daughters).

After seven weeks of growth in the greenhouse, the ultimate 3-4 leaflets from the youngest

fully expanded pinnate leaf from the top of each plant were harvested. Samples were flash-

frozen in liquid nitrogen, and subsequently freeze-dried for chemotyping.

A second unsupervised hierarchical clustering analysis was performed on the daughter

terpenoid profiles, again indicating six separate clusters. Daughter lineages did not typically

resemble mother lineages in their terpenoid profiles, and therefore the daughter clusters did

not reflect the mother lineages (Fig. 1). From each cluster, we selected three daughters from

the same mother lineage. When more than three daughters were available from a mother in

a cluster, we selected the three daughters clustering closest together, resulting in six

maternal chemotypes with three replicate daughters.

All daughters were grown under the same greenhouse conditions until July 2021, after which

they were transplanted to a common garden on-site and watered well until fully established.

Chemical analysis of plant material

Terpenoids were analysed as in Kleine & Müller (2011) and Wolf et al. (2012). Slight

modifications were made as in Eilers et al. (2021). The leaf material was freeze-dried,

homogenised, weighed and extracted in heptane, adding 1-bromodecane as internal

standard. Extracts were centrifuged and the supernatants analyzed using gas

chromatography coupled with mass spectrometry (GC-MS; GC 2010plus – MS QP2020,

Shimadzu, Kyoto, Japan) on a semi-polar column (VF-5 MS, 30 m length, 0.2 mm ID, 10 m

guard column, Varian, Lake Forest, United States) in electron impact ionisation mode at 70

eV, with helium as carrier gas. Samples were injected at 240 °C with a 1:10 split. A starting

temperature of 50 °C was kept for 5 min, ramping up to 250 °C at 10 °C min^{-1}, then

increasing with 30 °C min^{-1} to a final temperature of 280 °C, hold for 5 min. An alkane

standard mix (C7–C40, Sigma Aldrich, Taufkirchen, Germany) was measured regularly

between samples. Terpenoids were identified based on their retention indices and by

comparing spectra to various reference sources. Terpenoids were semi-quantified based on

the peak area of the total ion chromatogram, related to sample dry mass and the internal

standard. As we did not have calibrated response factors, the total concentration is always

an approximate total concentration.

Morphological traits

Morphological traits were measured non-destructively ten weeks after sowing for each of the

120 daughter plants in February 2021. The number of stems, leaves, and total number of

nodes per plant were counted, height was recorded in cm, and internode length was derived

by dividing height (in cm) by the number of nodes. From the youngest fully expanded leaf,

the total number of leaflets from one leaf, leaf length, and leaflets density (number of leaflets

/ petiole length) were assessed, and chlorophyll content was measured using a chlorophyll

meter (Konica Minolta SPAD-502Plus, Tokyo, Japan). The leaf was harvested and

photographed in a flattened position with a reference mark for determination of leaf surface

area using ImageJ®. The leaves were dried for 72 hours at 60°C to determine dry weight,

specific leaf area, and specific leaf mass.

Propagation of plant material for aphid choice assays

In August 2021, fresh plant material was taken from the 18 selected daughters, and shoot

cuttings were prepared by cutting stem parts 1-2 cm below and 4-5 cm above a leaf node.

The leaf size was reduced by roughly 60%, to reduce evapotranspiration. The stem cuttings

were pressed in seedling trays filled with potting substrate that was soaked. Cuttings were

covered by a transparent plastic hood that was gradually opened after the plants established

roots and showed first signs of shoot formation after three weeks. After acclimatizing the

cuttings for two weeks, they were repotted to 10 cm pots, and later to 17 cm pots, to avoid

pot limitation before their use in choice assays in December 2021. Plants were bottom-

watered automatically and fertilized with Universol Blue fertilizer (18% N - 11% P - 18% K;

ICL Deutschland, Nordhorn, Germany), target electrical conductivity of 1.0. No chemical

insecticides or fungicides were used during plant propagation. Due to low propagation

success of the Aue41 lineage, only the three daughters of the remaining five chemotypes

were used in pair-wise aphid choice assays.

Aphid rearing

Adults and nymphs of *Macrosiphoniella tanacetaria* and *Uroleucon tanaceti* were collected in

August 2021 tansy plants in a field in Dürnast (48.4049230,11.6898638) and kept in plastic

cages at room temperature with supplemental light at long day regimes (16h:8h L:D). Each

aphid colony was provided with two to four tansy plants at a time. Plants for feeding the

aphid colonies were collected from different locations in the vicinity of Technical University of

Munich campus in Freising, Germany, and were unrelated to the used chemotypes for this

study to avoid systemic bias in preferences in the aphid colonies. For each testing round,

roughly 100 unwinged adult aphids of each species were placed on individual fresh plants.

After 48 h, adult aphids were removed, and all nymphs kept on the plant. Subsequently, the

cohorts were kept in a Fitotron® standard growth chamber 120 (21/16°C, 60% RH, Weiss

Technik, Giessen, Germany) for eight days. After that the aphids were starved in a petri dish

with a small piece of wet tissue for 24 h before the start of choice assays.

Pairwise choice assays

Five *T. vulgare* maternal chemotypes with three replicate daughters were used for choice

assays. *A priori*, a full-factorial series of pairwise choice assays was designed, in which

aphids could choose between two different chemotypes (Suppl. Table S1). Within each

replicate series, all chemotypes were tested against each other using randomly picked

daughters. However, we ensured that all possible chemotype-daughter combinations were

used in multiple replicate rounds. Choice assays were conducted in three rounds on three

different days with different aphid cohorts that were standardized by age (9-10 days old), but

within one week (9th - 16th December 2021). Multiple replicate rounds were performed in

each experimental day. In total we performed 23 replicate rounds for *M. tanacetaria,* and 13

for *U. tanaceti*, as their numbers were substantially lower. *T. vulgare* daughters were

assigned beforehand to ensure that each daughter was tested against each other daughter

from the other chemotypes. However, as *M. tanacetaria* was used in five rounds which had

been originally assigned to *U. tanaceti*, numbers of combinations of tansy daughters differed

(for more details see Suppl. Table S1).

For the choice assay setup, the second and third youngest fully expanded leaf of a plant of

the respective chemotype were selected, the first three leaflets discarded, and the following

leaflets clipped with scissors. Plant leaflets were cut between 9:00 and 10:00 and placed in

petri dishes. In petri dishes (14.5 cm diameter), two different leaflets were placed at equal

distances from the centre, 8 cm apart from each other. The sides for each replicate

alternated between replicate rounds to account for external bias. After preparing all the

leaflets, one starved aphid was placed in the centre of the petri dish, and the petri dishes

were sealed with Parafilm®, to prevent the leaves from drying quickly and placed in a TUM

Model EcoSystem Analyser (TUM*mesa*) climate chamber. The LED lighting system in these

chambers generates homogenous light conditions, aimed at reducing spatial effects. Petri

dishes were placed onto a stainless-steel table and left for 24 h (16h:8h L:D photoperiod,

21°C, 60% RH). Aphid preference was recorded after two, five and 24 h. Each aphid and leaf

were only used once. Dead aphids were excluded from further analyses.

Statistical analysis

All statistical analyses were performed in R (R Core Team 2021). Plant chemical profiles

were clustered using unsupervised hierarchical k-means clustering with the hclust() function,

performed on the dissimilarity matrix that was obtained from the approximate absolute

terpenoid concentrations using the dist() function. Absolute values were used here, as

attractiveness of a chemical profile is likely to be a combination of blend composition and

approximate total concentration, the latter of which would be taken out by normalizing the

terpenoid profiles. The number of clusters, *k*, was obtained using the elbow method. As for

mother and daughter profiles the elbow or the optimal k differed, we selected a $k = 7$ for

mothers, of which six were used for further chemotype selection. For the daughter profile

clustering, the elbow k was at 5 - 7 clusters, which is why $k = 6$ clusters was used for

daughters.

To test our first hypothesis, whether aphids show species-specific attraction to chemotypes,

we conducted binomial tests to test if aphids preferred one chemotype over another for each

pairwise combination. We tested whether the number of choices made for one chemotype in

a specific combination was significantly different from what would be expected in a random

choice. Binomial tests were also used to rule out spatial (left/right side of the petri dish)

effects on aphid preference. We ruled out observer effects on the choices, as well as on the

proportion of choice/no-choice using Chi-square tests. Clogit models were used to

investigate the attractiveness of specific tansy chemotypes and daughters for each aphid

species, across all combined pairwise tests with all replicates (see Table 2 & 3), using the

survival package (Therneau 2021). We used the z-values obtained from the model for each

tansy daughter as a proxy for attractiveness, with daughters with more positive z-values

being more and negative z-values being less attractive to aphids.

To test our second hypothesis, whether chemodiversity negatively correlated with

attractiveness, we calculated plant chemodiversity metrics (compound richness, Shannon

diversity, evenness, approximate total concentration) from the absolute terpenoid profiles,

using the vegan package (Oksanen et al. 2020) and tested them against aphid preference.

We tested for relationships between the approximate total concentrations of individual

compounds in a plant and attractiveness of the plant separately for each of the two aphid

species. We used Spearman correlations, as changes in chemical compounds within

individual profiles are potentially related to one another as they might occur on the same

synthetic pathways, but between individuals these relationships might not be linear. We

analyzed correlations of terpenoids and aphid preference at the tansy daughter level (n = 18,

note: we did not have data for the three daughters from Aue41 regarding aphid preference

due to propagation issues). We used unadjusted correlation plots and Holm-adjusted plots

for multiple correlations, using the RcmdrMisc package (Fox 2022) and present both for

visualization purposes. For verification purposes, we ran a multiple regression model testing

the effect of all compounds on attractiveness. As the models were limited by available

degrees of freedom, we removed all compounds that were measured in less than two-thirds

of all samples. Using step() in combination with variance inflation factors to address

multicollinearity, we reduced the model to the minimum adequate version, which only

included individual compounds and proved out to be non-significant.

To test our third hypothesis, whether there were growth-related trade-offs with

chemodiversity, we analyzed plant morphological and chemodiversity variables, by using

one-way ANOVA with chemotype as a fixed factor (six chemotypes, with n = 3 daughters).

We tested the height, number of stems, number of leaves, leaflets density, specific leaf area

and chlorophyll against terpenoid richness, terpenoid Shannon diversity, terpenoid Shannon

evenness and approximate total terpenoid concentration (R-code can be found in the Suppl.

1).

Results

Cluster analysis

Hierarchical k-means clustering of terpenoid profiles resulted in six large clusters, without

any small outlier clusters driven by strongly deviating individual profiles (Fig. 1). Chemotype

selection resulted in six maternal chemotype lines with three daughter replicates for each

cluster (Fig. 1). Selected chemotypes differed based on their terpenoid blend composition

(Fig. 2a), as well as terpenoid diversity, evenness, richness and approximate total

concentration (Fig. 2b-e). Specifically, Aue41 and Schienen3 were both dominated by β-

thujone. Schie5 had both, α- and β-thujone as prevalent compounds. Most of the terpenoid

blend in Saal2 was made up by β-trans-chrysanthenyl acetate. In the chemotype Spar11B,

sabinene and α-pinene were made up of 39% and 20% of the terpenoid concentration, but

no terpenoid was clearly dominating. The chemical profile of Cam7 was very mixed with no

compound clearly dominating. Cam7 and Spar11B had significantly higher terpenoid

Shannon diversity and terpenoid evenness than other chemotypes (Fig. 2b-c). The

chemotype did not significantly affect chemotype richness (Fig. 2d). Spar11B and Aue41 had

significantly lower approximate total terpenoid concentrations compared to the other

chemotypes (Fig. 2e).

Choice experiments

After two hours, seven *M. tanacetaria* and five *U. tanaceti* individuals were found dead and

were excluded from the analyses. 178 *M. tanacetaria* and 69 *U. tanaceti* had chosen a

specific chemotype pinna. *M. tanacetaria* aphids tended to marginally prefer leaflets from

Schie5 over leaflets from Spar11B and Schienen3 (Table 1, Fig. 3a). After five hours of

observation, *M. tanacetaria* showed a significant preference for Schie5 over Spar11B and

tended to prefer Saal2 over Cam7 and Spar11B (Table 1, Fig. 3b). After two hours, *U.*

tanaceti aphids significantly preferred Schie5 over Spar11B (Table 1, Fig. 3c), but after five

hours no more significant preferences were observed (Table 1, Fig. 3d).

Considering all aphid choices made across all pairwise comparisons in a clogit model, *M.*

tanacetaria aphids showed a significant attraction to Saal2 and a marginally significant

attraction to Schie5 chemotypes (Table 2, Fig. 4a), while *U. tanaceti* aphids did not show a

clear preference for any chemotype (Table 2, Fig. 4b). However, we found that aphids also

exhibited preferences at the individual plant daughter level, with *M. tanacetaria* showing a

significant preference for Saal2-95 and Saal2-100, and Schie5-55 and Schie5-56, whereas

U. tanaceti did not show significant preferences to any tansy daughters (Table 3).

For *M. tanacetaria*, we observed a trend of decreased attraction to plants with higher

diversity (Fig. 4c) and more evenly distributed terpenoid blends (Fig. 4d), but a trend of

increased attraction to plants with a higher approximate total terpenoid concentration (Fig.

4e). For *U. tanaceti*, we observed no relationships between attractiveness and

chemodiversity. Unadjusted correlation plots revealed that various relationships existed

between individual compounds and attractiveness of aphid species, but when Holm-adjusted

for multiple correlations were applied, this reduced the relationships between attractiveness

and individual terpenoids to only a weak negative relationship between 1-terpinen-4-yl

acetate and tansy attractiveness to *M. tanacetaria* (Fig. 4f). This was further verified with

simplified multiple regression models corrected for collinearity, which reduced the models to

single compound factors, which did not significantly affected attractiveness.

Correlation between chemical diversity and morphological plant traits

Significant positive relationships were observed between the number of stems per tansy

plant and terpenoid Shannon diversity (Table 4, Fig. 5a), tansy height and terpenoid

Shannon diversity (Fig. 5b), and tansy height and approximate total terpenoid concentration

(Fig. 5c).

Discussion

In this study, we used five distinct maternal chemotype lines to test how leaf chemical

profiles affected aphid preference in pairwise choice assays. We found that two aphid

species specialized on tansy showed species-specific preferences in some of the pairwise

choices. Across all pairwise combinations *Macrosiphoniella tanacetaria* showed stronger

preferences and patterns of attraction than *Uroleucon tanaceti*. In line with our expectations,

we observed a trend of higher attractiveness of plants with a higher approximate total

terpenoid concentration and lower diversity. These patterns were observed for *M.*

tanacetaria, but not for *U. tanaceti*. Interestingly, attraction of aphids could not be clearly

linked to individual terpenoids.

Supporting our first hypothesis, we show that aphids preferred different chemotypes in

pairwise choice assays. *M. tanacetaria* preferred leaves from plants with α-thujone/β-thujone

(Schie5) and β-trans-chrysanthenyl acetate (Saal2) chemotype, with preference becoming

more pronounced after five hours. Previous studies found contradicting results, with higher

numbers of *M. tanacetaria* on plants with β-thujone compared to trans-carvyl acetate in a

climate chamber experiment (Jakobs & Müller 2018), but a higher abundance of *M.*

tanacetaria on plants with camphor instead of β-thujone as dominant compound in a field

study (Benedek et al. 2019a). Similarly, *U. tanaceti* was significantly attracted towards

Schie5 (α-thujone/β-thujone), with preference becoming less pronounced over time. This is

interesting, as previously negative effects of β-thujone on *U. tanaceti* numbers have been

found in the field (Benedek et al. 2019b) although no clear preference was observed towards

chemotypes for this species in another common garden study (Kleine & Müller 2011). Under

field conditions different factors (such as insect preference, bottom-up and top-down

processes) affect aphid communities simultaneously, and this may explain some of the

discrepancies between field observations and choice assays, asin 'snapshot' field

observations, these factors can be hard to disentangle. Therefore studies under laboratory

conditions are necessary to understand the influence of these single factors on aphid

behaviour. Our results indicate that aphid preference for plants might not always reflect how

they perform on plants.

In our experiment, 70 - 91 % of *M. tanacetaria* individuals made a choice after five hours for

each combination, compared to only 30 - 61% of *U. tanaceti* individuals. These species-

specific differences between the aphids can possibly be explained by their life history and the

difference in their preferred niche (Jakobs, Schweiger & Müller 2019), and aligns with our

observations. We observed in various field and greenhouse experiments that these aphid

species showed different behaviours. For instance, *M. tanacetaria* tends to be more mobile

than *U. tanaceti*, and readily searches for new host plants when its current host deteriorates.

Uroleucon tanaceti typically remains on leaves until yellowing, to then move upward to a new

leaf, often directly above the one they were on. We found more deaths and fewer choices

made at later time points (i.e., after 24 h), which could be because individual leaflets do dry

out after some time. However, as individual choices were already apparent and similar after

two and five hours, we believe that deteriorating quality did not affect choices made after two
hours.

Across all experimental choice combinations, we found significant effects of chemotype on
attractiveness of those chemotypes for *M. tanacetaria*, but not for *U. tanaceti*. This is in line
with previous research from Kleine & Müller (2011), who found that *M. tanacetaria* did, but *U.*
tanaceti did not exhibit distinct preferences towards specific chemotypes. To understand
what the drivers of the attractiveness of a plant to *M. tanacetaria* may be, we investigated
relationships between tansy individual chemical properties and their level of attractiveness.
We observed that diversity of the terpenoid blend negatively correlated with attractiveness.
This supports ecological theory on the evolution of chemodiversity in plants, and particularly
the evolution of a breadth of specialized compounds to repel antagonists (Wetzel &
Whitehead 2020). Furthermore, we found that terpenoid evenness marginally negatively
correlated with attractiveness, indicating that blends that were more evenly distributed in
terpenoid composition were less attractive, than those dominated by one or several
compounds. A plausible explanation might be that having some highly dominant compounds
could serve as strong cues for plant recognition (Kleine & Müller 2011).Subsequent aphid
performance on tansy likely is affected more strongly by phloem composition (Jakobs &
Müller 2019). We also found a positive effect of approximate total terpenoid concentration on
attractiveness to *M. tanacetaria*. This might be an indication that although chemodiversity
can have a deterrent effect (Whitehead et al. 2021), the approximate concentration (as a
proxy for potential emission) of terpenoids can be an important cue for host finding. A role of
specialized metabolite concentration in attraction and repellence has been found in other
studies (reviewed in Macel 2011). For instance, contrasting effects have been observed in
the specialized aphid *Aphis jacobaea* on *Jacobaea vulgaris*, where plants high in
pyrrolizidine alkaloids hosted fewer aphids than plants with low concentrations, although this
could not be related to preference alone (Vrieling et al. 1991). Our results align with those
from a recent study that shows that chemodiversity is an important driver of dietary
specialization in insects (Leong et al. 2022). The relationships between *M. tanacetaria* and

approximate total terpenoid concentration, terpenoid diversity and evenness were marginally

significant, which was likely a result of the limiting number of chemotype lines (15) included

in our current study. Our study was limited by the poor propagation success of one of the

chemotype lines with replicates. It is likely that including a larger number of diverse

chemotypes would further strengthen these findings, and future studies on a range of

terpenoid diversity profiles would be meaningful.

Our analyses of the role of individual compounds in driving attractiveness indicated that the

effect of individual compounds as drivers of tansy attractiveness are minimal. After

corrections for multiple correlations, we observed a weak negative effect of 1-terpinen-4-yl

acetate on tansy attractiveness to *M. tanacetaria*. Several studies have shown relationships

between chemotypes, or dominant compounds and aphid abundance in the field. For

instance, β-thujone has been related to decreased colony distribution and colony numbers in

U. tanaceti and abundances in *M. tanacetaria* in field studies (Bálint et al. 2016, Benedek et

al. 2019b). Other studies have found an increased density of *Metopeurum fuscoviride* on

chemotypes with high concentrations of borneol or camphor (Bálint et al. 2016, Senft et al.

2019), while plants with high amounts of α-thujone, (E)-dihydrocarvone, α-copaene and β-

cubebene were colonized earlier (Clancy et al. 2016). However, importantly, all these studies

investigated presence and abundance of aphids in the field, and did not take into account

aphid preference. Hence these studies are limited in the information they can provide about

aphid choice. Although we find only limited evidence that individual compounds drive aphid

attractiveness, it cannot be ruled out that these compounds play a role in feeding deterrence

when aphids are on the plant, or attraction of herbivore antagonists. Importantly, our study

included terpenoids only, but other volatile compounds (e.g., flavonoids) that were not

detected here, may affect the preference behavior of the aphids. This calls for future studies

disentangling the role of broader chemodiversity in driving aphid performance predator

attraction.

It is important to disentangle the relation between chemical and growth traits, as both traits

have been shown to influence plant-insect interactions, including preference. While growth-

defence trade-offs are a fundamental principle in plant ecology (Herms & Mattson 1992;

Karasov et al. 2017), we did not observe such a trade-off in our model system. Instead, we

found a synergistic effect between plant growth and approximate total terpenoid

concentration and diversity in our plants. It is commonly assumed that plants have to partition

their resources between growth and defence traits, leading to either smaller and better

defended plants, or *vice versa* (Coley at al. 1985, Herms & Mattson 1992, He et al. 2022).

Terpenoids are more expensive to build than many other metabolites, as they require a wide

array of different enzymes, hence posing substantial production and storage costs

(Gershenzon 1994).   However, contrary to these hypotheses, we found that larger and

bushier plants have a higher approximate total concentration and Shannon diversity of

terpenoid compounds. Similarly, Barton in 2007 found a positive relationship of growth and

defense in two *Plantago* species. One plausible explanation for our findings may be that

larger plants photosynthesize more, and have a larger energy budget, which can be used for

elevated and diversified local terpenoid synthesis. Furthermore, it is vital for plants to grow

and defend to optimize their fitness within a dynamic environment (Huot et al. 2014), and

trade-offs might also change with different life ontogeny stages (Boege & Marquis 2005).

Recent work has shown strong effects of maternal "chemo-genotypes" on plant chemical

composition in tansy (Dussarrat et al. 2023). Given that our used chemotypes were

confounded by their maternal origin (chemo-genotypes), this may also affect chemical and

morphological traits, and future work should investigate whether this also influences aphid

attraction and performance.

In this study, we used unwinged aphid morphs. Although in early colonization in the field,

winged morphs are generally more likely to choose host plants than unwinged morphs

(Mehrpavar et al. 2014), dispersal to other plants is commonly observed for unwinged aphids

in this system, both in experimental colonies and the field (the authors, pers. obs.). Although

performing choice assays with winged early colonizers would be an important next step, such

early wing morphs are difficult to obtain experimentally. In a choice assay study, Mehrparvar

and colleagues (2014) found that winged dispersal morphs of *M. tanacetaria* displayed

preferences for plants with specific herbivore infestation history, while unwinged aphids did

not show any such host plant preferences. Our results indicate that preferences and choices

in unwinged morphs do occur at the chemotype level, which likely represents a stronger

chemical contrast for aphids than the herbivory history studied in previous work.

Even though aphids preferred specific chemotypes, these might not necessarily be the most

suitable chemotypes for colony development. There has been a whole debate regarding the

"mother knows best" principle which is connected to the preference-performance hypothesis,

which posits that females insects prefer to oviposit on host plants on which their offspring

performs best (Martínez et al 2013, Gripenberg et al 2010). In the beetle *Gonioctena*

linnaeana for example, it has been found that mothers preferred host plants on which the

survival rate of their larvae was significantly higher (Wennström et al. 2010), while females of

the root-feeding vine weevil *Otiorhynchus sulcatus* preferred to oviposit on plants with small

root systems, which decreased offspring performance (Clark et al. 2011). Whether

preference overlaps with performance differs between species (Valladares & Lawton 1991,

García-Robledo & Horvitz 2012). For *U. tanaceti* the "mother knows best" principle seems to

apply, as they chose the chemotype and plant parts in a choice-study, on which their

offspring performed best, while in *M. tanacetaria* this was not the case (Jakobs & Müller

2018).

Conclusion

In this study, we show that plant chemodiversity can be an important driver for aphid host

plant preference, which is in line with other studies (Wolf et al. 2012), strengthening our

understanding of the role of chemodiversity in plants. We found that different chemotypes

have distinct attraction patterns, and that these can be explained in part by diversity metrics

of the terpenoid blend, as well as the approximate total concentration. Insects use chemical

cues, including host plant metabolites, to inform their decisions. It is eminent that we develop

a better understanding of how intraspecific plant chemodiversity shapes the various aspects

of the (herbivorous) insect life cycle, including development, survival, defense, and overall

fitness.

Acknowledgements

We thank the staff at TUM PTC Dürnast, particularly Sabine Zuber and Petra Scheuerer, for

providing excellent care during plant propagation and preparation phase.

Contribution of authors

ANH, LOP, WWW and RH conceived and designed the experiment. LOP and RH

propagated plant material. EE and CM chemotyped plant material. RH and LOP performed

morphological trait measurements. ANH and LOP performed the choice assays. ANH, LOP

and RH analyzed the data. ANH and RH wrote the paper, with substantial input from LOP. All

authors read and approved the manuscript before submission.

References

Agrawal, A. A., & Fishbein, M. (2006). Plant defense syndromes. *Ecology*, *87*(sp7), S132–S149.

https://doi.org/10.1890/0012-9658(2006)87[132:PDS]2.0.CO;2

Bálint, J., Zytynska, S. E., Salamon, R. V., Mehrparvar, M., Weisser, W. W., Schmitz, O. J., Benedek, K.,

& Balog, A. (2016). Intraspecific differences in plant chemotype determine the structure of arthropod

food webs. *Oecologia*, *180*(3), 797–807. https://doi.org/10.1007/s00442-015-3508-y

Barton, K. E. (2007). Early ontogenetic patterns in chemical defense in Plantago (Plantaginaceae):

genetic variation and trade-offs. American Journal of Botany, 94(1), 56-66.Batyrshina, Z. S., Cna'ani,

A., Rozenberg, T., Seifan, M., & Tzin, V. (2020). The combined impacts of wheat spatial position and

phenology on cereal aphid abundance. *PeerJ*, *8*, e9142. https://doi.org/10.7717/peerj.9142

Benedek, K., Bálint, J., Máthé, I., Mara, G., Felföldi, T., Szabó, A., Fazakas, C., Albert, C., Buchkowski,

R. W., Schmitz, O. J., & Balog, A. (2019a). Linking intraspecific variation in plant chemical defence

with arthropod and soil bacterial community structure and N allocation. *Plant and Soil*, *444*(1–2), 383–

397. https://doi.org/10.1007/s11104-019-04284-7

Benedek, K., Mara, G., Mehrparvar, M., Bálint, J., Loxdale, H. D., & Balog, A. (2019b). Near-regular

distribution of adult crimson tansy aphids, *Uroleucon tanaceti* (L.), increases aposematic signal

honesty on different tansy plant chemotypes. *Biological Journal of the Linnean Society*, *126*(2), 315–

326. https://doi.org/10.1093/biolinnean/bly180

Boege, K., & Marquis, R. J. (2005). Facing herbivory as you grow up: the ontogeny of resistance in

plants. Trends in Ecology & Evolution, 20(8), 441–448. https://doi.org/10.1016/j.tree.2005.05.001

Bustos-Segura, C., Poelman, E. H., Reichelt, M., Gershenzon, J., & Gols, R. (2017). Intraspecific

chemical diversity among neighbouring plants correlates positively with plant size and herbivore load

but negatively with herbivore damage. *Ecology Letters*, *20*(1), 87–97.

https://doi.org/10.1111/ele.12713

Caldwell, E., Read, J., & Sanson, G. D. (2015). Which leaf mechanical traits correlate with insect

herbivory among feeding guilds? *Annals of Botany*, mcv178. https://doi.org/10.1093/aob/mcv178

Clancy, M. V., Zytynska, S. E., Senft, M., Weisser, W. W., & Schnitzler, J.-P. (2016). Chemotypic

variation in terpenes emitted from storage pools influences early aphid colonisation on tansy. *Scientific*

Reports, *6*(1), 38087. https://doi.org/10.1038/srep38087

Clancy, M. V., Zytynska, S. E., Moritz, F., Witting, M., Schmitt-Kopplin, P., Weisser, W. W., & Schnitzler,

J.-P. (2018). Metabotype variation in a field population of tansy plants influences aphid host selection:

Plant chemical diversity in a plant-aphid system. *Plant, Cell & Environment*, *41*(12), 2791–2805.

https://doi.org/10.1111/pce.13407

Clark, K. E., Hartley, S. E., & Johnson, S. N. (2011). Does mother know best? The preference-

performance hypothesis and parent-offspring conflict in aboveground-belowground herbivore life

cycles. *Ecological Entomology*, *36*(2), 117–124. https://doi.org/10.1111/j.1365-2311.2010.01248.x

Coley, P. D., Bryant, J. P., & Chapin, F. S. (1985). Resource availability and plant anti-herbivore defense.

Science, *230*(4728), 895–899. https://doi.org/10.1126/science.230.4728.895

Crutsinger, G. M., Collins, M. D., Fordyce, J. A., Gompert, Z., Nice, C. C., & Sanders, N. J. (2006). Plant

genotypic diversity predicts community structure and governs an ecosystem process. *Science*,

313(5789), 966–968. https://doi.org/10.1126/science.1128326

Dicke, M. (2009). Behavioural and community ecology of plants that cry for help. Plant, Cell &

Environment, 32(6), 654–665. https://doi.org/10.1111/j.1365-3040.2008.01913.x

Des Roches, S., Post, D. M., Turley, N. E., Bailey, J. K., Hendry, A. P., Kinnison, M. T., Schweitzer, J. A.,

& Palkovacs, E. P. (2018). The ecological importance of intraspecific variation. *Nature Ecology &*

Evolution, *2*(1), 57–64. https://doi.org/10.1038/s41559-017-0402-5

Dussarrat, T., Schweiger, R., Ziaja, D., Nguyen, T. T. N., Krause, L., Jakobs, R., Eilers, E. J., & Müller,

C. (2023). Influences of chemotype and parental genotype on metabolic fingerprints of tansy plants

uncovered by predictive metabolomics [Preprint]. Ecology. https://doi.org/10.1101/2023.02.16.528607

Eilers, E. J., Kleine, S., Eckert, S., Waldherr, S., & Müller, C. (2021). Flower Production, Headspace

Volatiles, Pollen Nutrients, and Florivory in Tanacetum vulgare Chemotypes. Frontiers in Plant

Science, 11, 611877. https://doi.org/10.3389/fpls.2020.611877

Fox, J. (2022). RcmdrMisc: R Commander Miscellaneous Functions. R package version 2.7-2.

https://CRAN.R-project.org/package=RcmdrMisc

García-Robledo, C., & Horvitz, C. C. (2012). Parent-offspring conflicts, "optimal bad motherhood" and the

"mother knows best" principles in insect herbivores colonizing novel host plants: Parent-Offspring

Conflicts in Insect Herbivores. *Ecology and Evolution*, *2*(7), 1446–1457.

https://doi.org/10.1002/ece3.267

Gershenzon, J. (1994). Metabolic costs of terpenoid accumulation in higher plants. *Journal of Chemical*

Ecology, *20*(6), 1281–1328. https://doi.org/10.1007/BF02059810

Gripenberg, S., Mayhew, P. J., Parnell, M., & Roslin, T. (2010). A meta-analysis of preference-

performance relationships in phytophagous insects. *Ecology Letters*, *13*(3), 383–393.

https://doi.org/10.1111/j.1461-0248.2009.01433.x

Hayashi, T., Tahara, S., & Ohgushi, T. (2005). Genetically-controlled leaf traits in two chemotypes of

Salix sachalinensis Fr. Schm (Salicaceae). *Biochemical Systematics and Ecology*, *33*(1), 27–38.

https://doi.org/10.1016/j.bse.2004.07.001

He, Z., Webster, S., & He, S. Y. (2022). Growth–defense trade-offs in plants. *Current Biology*, *32*(12),

R634–R639. https://doi.org/10.1016/j.cub.2022.04.070

Herms, D. A., & Mattson, W. J. (1992). The dilemma of plants: To grow or defend. *The Quarterly Review*

of Biology, *67*(3), 283–335. https://doi.org/10.1086/417659

Holopainen, J. K., & Gershenzon, J. (2010). Multiple stress factors and the emission of plant VOCs.

Trends in Plant Science, *15*(3), 176–184. https://doi.org/10.1016/j.tplants.2010.01.006

Huot, B., Yao, J., Montgomery, B. L., & He, S. Y. (2014). Growth–Defense Tradeoffs in Plants: A

Balancing Act to Optimize Fitness. *Molecular Plant*, *7*(8), 1267–1287.

https://doi.org/10.1093/mp/ssu049

Iason, G. R., Dicke, M., & Hartley, S. E. (2012). The Ecology of Plant Secondary Metabolites: From

Genes to Global Processes. *Cambridge University Press*. ISBN: 1107375703

Jakobs, R., & Müller, C. (2018). Effects of intraspecific and intra-individual differences in plant quality on

preference and performance of monophagous aphid species. Oecologia, 186(1), 173–184.

https://doi.org/10.1007/s00442-017-3998-x

Jakobs, R., & Müller, C. (2019). Volatile, stored and phloem exudate-located compounds represent

different appearance levels affecting aphid niche choice. Phytochemistry, 159, 1–10.

https://doi.org/10.1016/j.phytochem.2018.11.018

Jakobs, R., Schweiger, R., & Müller, C. (2019). Aphid infestation leads to plant part-specific changes in

phloem sap chemistry, which may indicate niche construction. *New Phytologist*, *221*(1), 503–514.

https://doi.org/10.1111/nph.15335

Karasov, T. L., Chae, E., Herman, J. J., & Bergelson, J. (2017). Mechanisms to mitigate the trade-off

between growth and defense. *The Plant Cell*, *29*(4), 666–680. https://doi.org/10.1105/tpc.16.00931

Keskitalo, M., Pehu, E., & Simon, J. E. (2001). Variation in volatile compounds from tansy (*Tanacetum*

vulgare L.) related to genetic and morphological differences of genotypes. *Biochemical Systematics*

and Ecology, *29*(3), 267–285. https://doi.org/10.1016/S0305-1978(00)00056-9

Kessler, A., & Baldwin, I. T. (2001). Defensive function of herbivore-induced plant volatile emissions in

nature. *Science*, *291*(5511), 2141–2144. https://doi.org/10.1126/science.291.5511.2141

Kleine, S., & Müller, C. (2011). Intraspecific plant chemical diversity and its relation to herbivory.

Oecologia, *166*(1), 175–186. https://doi.org/10.1007/s00442-010-1827-6

Leong, J. V., Jorge, L. R., Seifert, C. L., & Volf, M. (2022). Quantity and specialisation matter: Effects of

quantitative and qualitative variation in willow chemistry on resource preference in leaf-chewing

insects. *Insect Conservation and Diversity*, *15*(4), 453–460. https://doi.org/10.1111/icad.12559

Macel, M. (2011). Attract and deter: a dual role for pyrrolizidine alkaloids in plant–insect interactions.

Phytochemistry Reviews, *10*(1), 75–82. https://doi.org/10.1007/s11101-010-9181-1

Martínez, G., Soler, R., & Dicke, M. (2013). Behavioral ecology of oviposition-site selection in

herbivorous true bugs. In *Advances in the Study of Behavior* (Vol. 45, pp. 175–207). Elsevier.

https://doi.org/10.1016/B978-0-12-407186-5.00004-5

McCormick, A. C., Unsicker, S. B., & Gershenzon, J. (2012). The specificity of herbivore-induced plant

volatiles in attracting herbivore enemies. *Trends in Plant Science*, *17*(5), 303–310.

https://doi.org/10.1016/j.tplants.2012.03.012

Mehrparvar, M., Mansouri, S. M., & Weisser, W. W. (2014). Mechanisms of species-sorting: effect of

habitat occupancy on aphids' host plant selection: Host selection by competing herbivores. *Ecological*

Entomology, *39*(3), 281–289. https://doi.org/10.1111/een.12096

Nishida, R. (2014). Chemical ecology of insect–plant interactions: ecological significance of plant

secondary metabolites. *Bioscience, Biotechnology, and Biochemistry*, *78*(1), 1–13.

https://doi.org/10.1080/09168451.2014.877836

Oksanen, J. F., Blanchet, G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara,

R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E. & Wagner, H. (2020). vegan:

Community Ecology Package. R package version 2.5-7. https://CRAN.R-project.org/package=vegan

Pichersky, E., & Gang, D. R. (2000). Genetics and biochemistry of secondary metabolites in plants: an

evolutionary perspective. *Trends in Plant Science*, *5*(10), 439–445. https://doi.org/10.1016/S1360-

1385(00)01741-6

Poelman, E. H., van Loon, J. J. A., & Dicke, M. (2008). Consequences of variation in plant defense for

biodiversity at higher trophic levels. *Trends in Plant Science*, *13*(10), 534–541.

https://doi.org/10.1016/j.tplants.2008.08.003

R Core Team (2021). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Raffard, A., Santoul, F., Cucherousset, J., & Blanchet, S. (2019). The community and ecosystem

consequences of intraspecific diversity: a meta-analysis. *Biological Reviews*, *94*(2), 648–661.

https://doi.org/10.1111/brv.12472

Richards, L. A., Dyer, L. A., Forister, M. L., Smilanich, A. M., Dodson, C. D., Leonard, M. D., & Jeffrey, C.

S. (2015). Phytochemical diversity drives plant–insect community diversity. *Proceedings of the*

National Academy of Sciences, *112*(35), 10973–10978. https://doi.org/10.1073/pnas.1504977112

Rzanny, M., Kuu, A., & Voigt, W. (2013). Bottom-up and top-down forces structuring consumer

communities in an experimental grassland. *Oikos*, *122*(7), 967–976. https://doi.org/10.1111/j.1600-

0706.2012.00114.x

Salazar, D., Lokvam, J., Mesones, I., Vásquez Pilco, M., Ayarza Zuñiga, J. M., de Valpine, P., & Fine, P.

V. A. (2018). Origin and maintenance of chemical diversity in a species-rich tropical tree lineage.

Nature Ecology & Evolution, *2*(6), 983–990. https://doi.org/10.1038/s41559-018-0552-0

Scherber, C., Eisenhauer, N., Weisser, W. W., Schmid, B., Voigt, W., Fischer, M., Schulze, E.-D.,

Roscher, C., Weigelt, A., Allan, E., Beßler, H., Bonkowski, M., Buchmann, N., Buscot, F., Clement, L.

W., Ebeling, A., Engels, C., Halle, S., Kertscher, I., Klein, A.-M., Koller, R., König, S., Kowalski, E.,

Kummer, V., Kuu, A., Lange, M., Lauterbach, D., Middelhoff, C., Migunova, V. D., Milcu, A., Müller, R.,

Partsch, S., Petermann, J. S., Renker, C., Rottstock, T., Sabais, A., Scheu, S., Schumacher, J.,

Temperton, V., Tscharntke, T. (2010). Bottom-up effects of plant diversity on multitrophic interactions

in a biodiversity experiment. Nature, 468(7323), 553–556. https://doi.org/10.1038/nature09492

Schmitz, G. (1998): The phytophagous insect fauna of Tanacetum vulgare L. (Asteraceae) in Central

Europe. Contributions to Entomology 48(1): 219–235 - doi: 10.21248/contrib.entomol.48.1.219-235

Schmitz, O. J. (2008). Herbivory from Individuals to Ecosystems. *Annual Review of Ecology, Evolution,*

and Systematics, *39*, 133–152. https://doi.org/10.1146/annurev.ecolsys.39.110707.173418

Schoonhoven, L. M., Loon, J. J. A. van, & Dicke, M. (2005). Insect-plant biology (2nd ed). *Oxford*

University Press. ISBN: 0198525948

Senft, M., Weisser, W. W., & Zytynska, S. E. (2017). Habitat variation, mutualism and predation shape

the spatio-temporal dynamics of tansy aphids: Spatio-temporal dynamics of tansy aphids. *Ecological*

Entomology, *42*(4), 389–401. https://doi.org/10.1111/een.12396

Senft, M., Clancy, M. V., Weisser, W. W., Schnitzler, J., & Zytynska, S. E. (2019). Additive effects of plant

chemotype, mutualistic ants and predators on aphid performance and survival. *Functional Ecology*,

33(1), 139–151. https://doi.org/10.1111/1365-2435.13227

Singh, A., Dilkes, B., Sela, H., & Tzin, V. (2021). The effectiveness of physical and chemical defense

responses of wild emmer wheat against aphids depends on leaf position and genotype. *Frontiers in*

Plant Science, *12*, 667820. https://doi.org/10.3389/fpls.2021.667820

Therneau, T. (2021). A Package for Survival Analysis in R. R package version 3.2-13, <URL:

https://CRAN.R-project.org/package=survival>

Unsicker, S. B., Kunert, G., & Gershenzon, J. (2009). Protective perfumes: the role of vegetative volatiles

in plant defense against herbivores. *Current Opinion in Plant Biology*, *12*(4), 479–485.

https://doi.org/10.1016/j.pbi.2009.04.001

Valladares, G., & Lawton, J. H. (1991). Host-plant selection in the holly leaf-miner: does mother know

best? *The Journal of Animal Ecology*, *60*(1), 227. https://doi.org/10.2307/5456

Volf, M., Salminen, J.-P., & Segar, S. T. (2019). Evolution of defences in large tropical plant genera:

perspectives for exploring insect diversity in a tri-trophic context. *Current Opinion in Insect Science*,

32, 91–97. https://doi.org/10.1016/j.cois.2018.12.005

Vrieling, K., Smit, W., & van der Meijden, E. (1991). Tritrophic interactions between aphids (*Aphis*

jacobaeae Schrank), ant species, *Tyria jacobaeae* L., and *Senecio jacobaea* L. lead to maintenance of

genetic variation in pyrrolizidine alkaloid concentration. *Oecologia*, *86*(2), 177–182.

https://doi.org/10.1007/BF00317529

Weisser, W. W., Roscher, C., Meyer, S. T., Ebeling, A., Luo, G., Allan, E., Beßler, H., Barnard, R. L.,

Buchmann, N., Buscot, F., Engels, C., Fischer, C., Fischer, M., Gessler, A., Gleixner, G., Halle, S.,

Hildebrandt, A., Hillebrand, H., De Kroon, H., Lange, M., Leimer, S., Le Roux, X., Milcu, A., Mommer,

L., Niklaus, P. A., Oelmann, Y., Proulx, R., Roy, J., Scherber, C., Scherer-Lorenzen, M., Scheu, S.,

Tscharntke, T., Wachendorf, M., Wagg, C., Weigelt, A., Wilcke, W., Wirth, C., Schulze, E.-D., Schmid,

B., Eisenhauer, N. (2017). Biodiversity effects on ecosystem functioning in a 15-year grassland

experiment: Patterns, mechanisms, and open questions. Basic and Applied Ecology, 23, 1–73.

https://doi.org/10.1016/j.baae.2017.06.002

Weng, J.-K., Philippe, R. N., & Noel, J. P. (2012). The rise of chemodiversity in plants. *Science*,

336(6089), 1667–1670. https://doi.org/10.1126/science.1217411

Wennström, A., Niemi Hjulström, L., Hjältén, J., & Julkunen-Tiitto, R. (2010). Mother really knows best:

Host choice of adult phytophagous insect females reflects a within-host variation in suitability as larval

food. *Chemoecology*, *20*(1), 35–42. https://doi.org/10.1007/s00049-010-0040-8

Wetzel, W. C., & Whitehead, S. R. (2020). The many dimensions of phytochemical diversity: Linking

theory to practice. *Ecology Letters*, *23*(1), 16–32. https://doi.org/10.1111/ele.13422

Whitehead, S. R., Bass, E., Corrigan, A., Kessler, A., & Poveda, K. (2021). Interaction diversity explains

the maintenance of phytochemical diversity. *Ecology Letters*, *24*(6), 1205–1214.

https://doi.org/10.1111/ele.13736

Wink, M. (2018). Plant secondary metabolites modulate insect behavior-steps toward addiction. *Frontiers*

in Physiology, 9, 364. https://doi.org/10.3389/fphys.2018.00364

Wolf, V. C., Berger, U., Gassmann, A., & Müller, C. (2011). High chemical diversity of a plant species is

accompanied by increased chemical defence in invasive populations. Biological Invasions, 13(9),

2091–2102. https://doi.org/10.1007/s10530-011-0028-5

Wolf, V. C., Gassmann, A., & Müller, C. (2012). Choice behaviour and performance of Cassida

stigmatica on various chemotypes of Tanacetum vulgare and implications for biocontrol: Insect

responses to different plant chemotypes. *Entomologia Experimentalis et Applicata*, *144*(1), 78–85.

https://doi.org/10.1111/j.1570-7458.2012.01242.x

Ziaja D, Müller C (2022) Intraspecific chemodiversity provides plant individual- and neighbourhood-

mediated associational resistance towards aphids. BioRxiv,

https://www.biorxiv.org/content/10.1101/2022.12.21.521353v1

Zytynska, S. E., Guenay, Y., Sturm, S., Clancy, M. V., Senft, M., Schnitzler, J., Dilip Pophaly, S.,

Wurmser, C., & Weisser, W. W. (2019). Effect of plant chemical variation and mutualistic ants on the

local population genetic structure of an aphid herbivore. *Journal of Animal Ecology*, *88*(7), 1089–1099.

https://doi.org/10.1111/1365-2656.12995

Figures

712   Figure 1

714   Figure 1: Hierarchical clustering tree for all 120 daughters terpenoid profiles. Clusters (k = 6)

715   are indicated in red boxes. For the selection, in each cluster a different mother lineage was

716   chosen, and three daughters were manually selected and highlighted. The final selection of

717   daughters of each cluster are indicated in blue boxes.

Figure 2

Figure 2: a) Stacked bar chart chemical profiles for the selected six maternal chemotypes

with three daughter replicates. Box plots show the interquartile ranges of b) terpenoid

Shannon diversity, c) terpenoid evenness, d) terpenoid richness, and e) approximate total

terpenoid concentration for each chemotype. The lower hinge corresponds to the first quartile

(25th percentile) and the upper hinge depicts the third quartile (75th percentile). Whiskers

extend to the 5% and 95% percentiles; solid lines represent the medians. Chemotype effect

is indicated in panels, letters above bars indicate significant differences ($p < 0.05$) between

chemotypes based on posthoc Tukey tests.

Figure 3

M. tanacetaria

U. tanaceti

Figure 3: Mean percentage of choices by a) *Macrosiphoniella tanacetaria*, and b) *Uroleucon tanaceti* after two hours for all possible pairwise combinations of chemotypes, with error bars indicating standard errors. Effective sample size is indicated for each pairwise comparison, and no-choice replicates were excluded. Note that chemotype Aue41 was not included in pairwise comparisons due to propagation difficulties. P-values next to bars indicate (marginally) significant preferences (for test statistics see Table 1).

Figure 4

Figure 4: Relationships between tansy attractiveness and chemodiversity for both aphid

species. Attractiveness (reflected by the z-value) was computed by summarizing all

decisions made towards a specific chemotype over all combinations and using a clogit model

to test whether a certain chemotype was chosen more often than expected compared to a

random choice. a) mean attractiveness per chemotype for *M. tanacetaria.* b) Mean

attractiveness per chemotype for *U. tanaceti*. (For detailed model output for a) and b) see

Table 2) c) Relationship between tansy terpenoid Shannon diversity and attractiveness to *M.*

tanacetaria. d) Relationship between tansy terpenoid evenness and attractiveness to *M.*

tanacetaria. e) Relationship between tansy terpenoid abundance and attractiveness to *M.*

tanacetaria; in c) to e) dots represent the mean z-value (attractiveness) for each of the 15

daughters (5 Chemotypes * 3 daughters). f) Correlation plots showing relationship between

individual terpenoid compounds and attractiveness to both aphid species. The size of the dot

depicts the strength of the correlation. Blue colors indicate positive and red colors indicate

negative correlations. The left correlation plot shows all unadjusted correlations with a

significant p-value. The right correlation plot shows all correlations with a significant p-value

after using Holm-adjusted for multiple correlations.

    Figure 5

Figure 5: Relationships between chemical and morphological parameters. Each dot

represents one of the 18 daughters (6 chemotypes * 3 daughters). a) Relationship between

the number of stems per tansy plant and terpenoid Shannon diversity. b) Relationship

between tansy height and terpenoid Shannon diversity. c) Relationship between tansy height

and approximate total terpenoid concentration.

Tables

Table 1: Influence of tansy chemotypes on their attractiveness to *M. tanacetaria* and *U.*

tanaceti, when given pairwise choices. A binomial test was used to obtain credible intervals

and p-values. Significant values are highlighted in bold, marginally significant values in

italics.

			M. tanacetaria			*U. tanaceti*		
Time	Chemotype 1	Chemotype 2	n	95% CI	p-value	n	95% CI	p-value
2 h	Schie5	Spar11B	18	*0.465, 0.903*	*0.096*	9	**0.518, 0.997**	**0.039**
2 h	Schie5	Schienen3	19	*0.488, 0.909*	*0.064*	9	0.212, 0.863	1
2 h	Schie5	Cam7	20	0.231, 0.685	0.824	7	0.184, 0.901	1
2 h	Schie5	Saal2	17	0.230, 0.722	1	6	0.118, 0.882	1
2 h	Spar11B	Saal2	16	0.110, 0.587	0.210	7	0.184, 0.901	1
2 h	Cam7	Saal2	15	0.078, 0.51	0.119	6	0.043, 0.778	0.688
2 h	Schienen3	Saal2	18	0.133, 0.590	0.238	6	0.043, 0.778	0.688
2 h	Schienen3	Spar11B	17	0.330, 0.816	0.629	8	0.085, 0.755	0.727
2 h	Schienen3	Cam7	19	0.435, 0.874	0.167	6	0.043, 0.778	0.688
2 h	Spar11B	Cam7	18	0.215, 0.692	0.815	5	0.284, 0.995	0.375
5 h	Schie5	Spar11B	17	**0.566, 0.962**	**0.013**	5	0.147, 0.947	1
5 h	Schie5	Schienen3	18	0.410, 0.867	0.238	8	0.157, 0.843	1
5 h	Schie5	Cam7	19	0.335, 0.798	0.648	4	0.194, 0.994	0.625
5 h	Schie5	Saal2	16	0.247, 0.754	1	7	0.099, 0.816	1
5 h	Spar11B	Saal2	*18*	*0.097, 0.535*	*0.096*	6	0.043, 0.778	0.688
5 h	Cam7	Saal2	*19*	*0.092, 0.512*	*0.064*	5	0.147, 0.947	1
5 h	Schienen3	Saal2	16	0.247, 0.754	1	6	0.118, 0.882	1
5 h	Schienen3	Spar11B	19	0.289, 0.756	1	6	0.118, 0.882	1
5 h	Schienen3	Cam7	21	0.430, 0.854	0.189	8	0.085, 0.755	0.727
5 h	Spar11B	Cam7	18	0.215, 0.692	0.815	5	0.147, 0.947	1

772   Table 2: Influence of tansy chemotypes on their attractiveness to *M. tanacetaria* and *U.*
773   *tanaceti* based on all pairwise combinations of the choice assays after two hours of
774   observation. A c-logit model, including effects of chemotype (five chemotypes) on aphid
775   choices after two hours, was used to obtain z- and p-values (for visualization see figure 4).
776   Significant values are highlighted in bold, marginally significant values in italics.

Chemotype	*M. tanacetaria*		*U. tanaceti*	
	z-value	p-value	z-value	p-value
Cam7	-0.669	0.503	-1.226	0.220
Saal2	**2.331**	**0.020**	0.783	0.434
Schie5	*1.730*	*0.084*	1.206	0.228
Schienen3	0.340	0.734	-0.448	0.654
Spar11B	-0.752	0.452	0.404	0.686

778   Table 3: Influence of tansy daughters on their attractiveness to *M. tanacetaria* and *U.*
779   *tanaceti* based on all pairwise combinations of the choice assays after two hours of
780   observation. A c-logit model, including effects of daughter (five chemotypes x three
781   daughters) on aphid choices after two hours, was used to obtain z- and p-values. Significant
782   values are highlighted in bold, marginally significant values in italics.

Daughter	*M. tanacetaria*		*U. tanaceti*	
	z-value	p-value	z-value	p-value
Cam7_21	-0.792	0.429	-1.203	0.229
Cam7_23	1.064	0.288	-1.447	0.148
Cam7_30	1.602	0.110	0.506	0.613
Saal2_100	**2.171**	**0.030**	0.302	0.762
Saal2_94	1.589	0.112	-0.659	0.510
Saal2_95	**2.439**	**0.015**	0.442	0.658
Schie5_55	**2.183**	**0.029**	-0.207	0.836
Schie5_56	**1.983**	**0.047**	0.100	0.921
Schie5_57	1.119	0.263	0.467	0.640

Schienen3_11	1.058	0.290	-1.597	0.110
Schienen3_14	1.451	0.147	-1.103	0.270
Schienen3_17	0.858	0.391	0.434	0.664
Spar11B_64	-0.246	0.806	-0.292	0.770
Spar11B_67	0.666	0.505	-0.446	0.656
Spar11B_68	1.075	0.283	-0.297	0.766

784 Table 4: Model output of multiple linear regression models testing the effects of plant

785 morphological characteristics on plant attractiveness for *M. tanacetaria* and *U. tanaceti*, and

786 on plant chemodiversity characteristics. Models were checked for collinearity by variance

787 inflation factors and simplified via stepwise model selection. Dashes indicate variables were

788 removed from the model.

Explanatory variable	Attractiveness *M. tanacetaria*	Attractiveness *U. tanaceti*	Terpenoid richness	Terpenoid Shannon diversity	Terpenoid Shannon evenness	Terpenoid concentration
	F (p-value)	F (p-value)	F (p-value)	F (p-value)	F (p-value)	F (p-value)
Height	-	-	0.20 (0.665)	**6.03 (0.028)**	**6.44 (0.024)**	**7.05 (0.017)**
# Stems	**10.20 (0.010)**	-	0.11 (0.741)	**8.33 (0.012)**	**9.12 (0.009)**	-
# Leaves	**5.02 (0.049)**	-	0.07 (0.801)	-	-	-
Leaflets density	3.14 (0.107)	1.85 (0.199)	-	-	-	-
SLA	-	-	-	-	-	-
Chlorophyll	**6.94 (0.0250)**	2.42 (0.146)	-	2.55 (0.133)	2.54 (0.133)	-

789