Sex differences in anxiety and threat avoidance in GAD65 knock-out mice

Michelle Ulrich[1], Evangelia Pollali[2,3], Gürsel Çalışkan[2,3,4] and Oliver Stork[2,3,5],

Anne Albrecht[1,3,5]*

[1] Institute of Anatomy, Otto-von-Guericke-University, Leipziger Str. 44, 39120 Magdeburg, Germany; michelle.ulrich@st.ovgu.de; anne.albrecht@med.ovgu.de

[2] Institute of Biology, Otto-von-Guericke-University Leipziger Str. 44, 39120 Magdeburg, Germany; evangelia.pollali@ovgu.de; guersel.caliskan@ovgu.de; oliver.stork@ovgu.de

[3] Center for Behavioral Brain Sciences (CBBS), Universitätsplatz 2, 39106 Magdeburg, Germany

[4] Research Group "Synapto-Oscillopathies", Institute of Biology, Otto-von-Guericke-University, Magdeburg, Germany

[5] Center for Intervention and Research on adaptive and maladaptive brain Circuits underlying mental health (C-I-R-C), Jena-Magdeburg-Halle, Germany

corresponding author: Anne Albrecht

Institute of Anatomy, Otto-von-Guericke-University, Leipziger Str. 44, 39120 Magdeburg, Germany; anne.albrecht@med.ovgu.de, Tel.: +49 391 6713604, Fax: +49 391 6713604

Number of figures: 5

Number of tables: 0

Number of supplementary figures: 3

Highlights

- GAD65 -/- mice demonstrate a role of GABA in sex-specific anxiety endophenotypes
- Enhanced *in vitro* gamma-beta oscillations in ACC slices of GAD65-/- mice
- Increased parvalbumin(+) interneuron counts in ACC slices of GAD65-/- mice
- impaired active avoidance learning in male but not female GAD65-/- mice
- reduced somatostatin(+) interneuron counts in dorsal DG and BLA of male GAD65-/- mice

Graphical abstract

Abstract

Anxiety disorders have been linked to a disbalance of excitation and inhibition in a network of brain structures comprising frontal cortical regions, the amygdala and the hippocampus, among others. While recent imaging studies in humans suggest sex differences in the activation of such anxiety network structures in processing emotional information, the neuronal basis of activation shifts and their relation to anxiety endophenotypes is less well studied. Rodent models with genetically altered ɣ-amino butyric acid (GABA) neurotransmission may provide a targeted approach for addressing such questions, but to date sex effects have rarely been addressed in such models. Using mice with a null mutation of the GABA synthetizing enzyme glutamate decarboxylase 65 (GAD65-/-), we started to compare anxiety-like behavior, discrimination and avoidance learning in male vs. female GAD65-/- mice and their wildtype littermates. Our behavioral test battery revealed increased activity on GAD65-/- mice in the open field and an adaptation of anxiety-like behavior especially in male GAD65-/- mice. Male GAD65-/- mice further displayed a deficit in avoidance learning, while Female GAD65 -/- mice performed well in this task. To gain insights into interneuron function in anxiety and threat perception network structures, fast oscillations (10-45 Hz) were measured in *ex vivo* slice preparations of the ACC. GAD65-/- mice of both sexes displayed increased gamma power in the ACC that was further accompanied by higher densities of PV-positive interneurons, which are crucial for generating such rhythmic activity. In addition, male GAD65-/- mice had lower numbers of somatostatin-positive interneurons in the basolateral amygdala and dorsal dentate gyrus, two key regions important for anxiety and active avoidance responses. Together, male mice appeared more challenged by GAD65 deficiency in avoidance learning tasks and adapted their anxiety-like response differently, which was associated with an altered interneuron composition and function in an ACC-amygdala-hippocampal network.

Keywords: ACC, anxiety, avoidance, sex differences, GAD65, inhibition, oscillations

Abbreviations: AA: active avoidance; ACC: anterior cingulate cortex; BLA: basolateral amygdala; DAPI: 4, 6-diamidino-2-phenylindole; dDG: dorsal dentate gyrus; EPM: elevated plus maze; F: female; LFP: local field potentials; GAD65: glutamic acid decarboxylating

enzyme, 65 kDa isoform; GAD67: glutamic acid decarboxylating enzyme, 67 kDa isoform; M: male; NOR: novel object recognition; OF: open field; PV: parvalbumin; SI: social interaction; SST: somatostatin; vCA1+3: ventral cornu ammonis 1+3.

1. Introduction

Anxiety disorders are among the psychopathologies with the highest prevalence in western societies and cause enormous individual and socioeconomic burden (Lépine, 2002; Steel et al., 2014). They are characterized by an excessive and uncontrollable worry that interferes with social or occupational functions and is accompanied by sleep disturbances, irritability, restlessness or fatigue (American Psychiatric Association, 2013). Anxiety endophenotypes and cognitive bias to negative interpretations of ambiguous stimuli are strong predisposing factors for developing an anxiety disorder (Ouimet et al, 2009; van der Heiden et al., 2010; Blanco et al., 2014). Further epidemiological studies frequently reveal sex differences (McLean & Anderson, 2009; Vesga-Lopez et al., 2008), indicating a two-times higher prevalence of anxiety disorders in women.

Differential activation of frontal cortical areas such as the medial prefrontal cortex (mPFC) and the anterior cingulate cortex (ACC) is evident between men and women depending on the emotional valence of presented stimuli, induced stress levels or prior adverse childhood experiences (Thomas et al., 2019; Javanbakht et al., 2016; Seo et al., 2017; Stevens & Hamann 2012). Accordingly, a dysfunctional activation of those cortical regions and their interaction with the amygdala and the hippocampal formation are observed during emotion regulation in anxiety patients (Mochkovotich et al., 2014; Goossen et al., 2019; Kim & Kim, 2021). This matches the basic function of these areas and their homologous structures in rodents regulating fear and anxiety as well as emotional affect, attention and decision making (Admon et al., 2013; Tovote et al., 2015; van Heukelum et al., 2020; Chen et al., 2022).

Interactions of cortical frontal areas, amygdala and hippocampus are coordinated through oscillatory rhythms that are under tight control of local inhibitory interneurons (Çalışkan & Stork, 2019). Such GABAergic interneurons control the local excitability and information processing within these regions. Drugs that modulate GABAergic neurotransmission are powerful anxiolytics (Roy-Byrne, 2005) and mice lacking the 65kDa isoform of the GABA-

synthetizing enzyme glutamic acid decarboxylase (GAD65) demonstrate a phenotype reminiscent of core symptoms of anxiety disorders, including increased anxiety, hyperarousal, fear generalization and an altered fear extinction (Müller et al., 2014). Different subpopulations of GABAergic interneurons can be described based on their morphological, physiological and neurochemical characteristics. Interneurons containing parvalbumin (PV) are essential for generation of hippocampal oscillations at the gamma frequency range and can modulate anxiety-like behavior and fear persistence (Winkelmann et al., 2014; Çalışkan et al., 2016). Experimental silencing of GABAergic neurotransmission in another subpopulation, somatostatin (SST)- positive interneurons, increases anxiety-like behavior (Miyata et al., 2019; Fee et al., 2021), suggesting a strong contribution of such inhibitory neurons to anxiety endophenotypes.

GAD65-/- mouse have been established as model for studying the neurobiological basis of anxiety phenotypes (Kash et al., 1999; Stork et al., 2000), but strikingly sex effects have not been yet addressed in these animals. In a previous study we were able to relate a polymorphism in the GAD65 gene to an increased GABA/Glutamate ratios in the ACC, associated with increased anxiety endophenotype and harm avoidance in female but not male participants (Colic et al., 2018). In the current study, we therefore set out to investigate sex differences in anxiety-related behavioral tasks and avoidance behavior in a two-way shuttle box in GAD65-/- mice. In addition to the characterization of anxiety-related endophenotypes we investigated beta/ slow gamma oscillations in the ACC as a function of interneuron activity and examined the configuration of GABAergic networks by immunohistochemical labelling of PV- and SST-positive interneurons in frontal cortical, hippocampal and amygdala subregions.

2. Material and Methods

2.1 Animals

Male and female homozygous GAD65 knock-out (-/-) mice and their wild type (+/+) littermates were obtained from a heterozygous breeding and genotyped by allele-specific polymerase chain reaction (PCR) as described previously (Stork et al., 2000). All mice were bred and raised in the animal facility at the Institute of Biology, Otto-von-Guericke University Magdeburg, and kept in groups of 2–6 of the same sex on an inverse 12 h light/dark cycle (lights on at 7 pm with a 30 min dawn phase) with food and water *ad libitum*. Animal housing and animal experiments were in accordance with European regulations for animal experiments and

approved by the Landesverwaltungsamt Saxony-Anhalt (Permission Nr. 42502-2-1177 and -1516). All experiments were performed with young adult female and male mice (age 10-16 weeks) during the dark, active circadian phase with handling under red light illumination.

2.2 Experiment 1: Behavioral test battery

Female GAD65-/- mice (n=12) and female GAD65+/+ littermates (n=12) as well as male GAD65-/- mice (n=13) and their male GAD65+/+ littermates (n=12) underwent a behavioral test battery consisting of open field (OF) on day 1, elevated plus maze (EPM) on day 2, novel object recognition (NOR) on day 3, social interaction (SI) on day 4 and five days of active avoidance training (AA) on day 7-11. The order of the behavioral tests was chosen according to rising stress levels induced by the single tests.

Open field. Mice were placed in the center (25 x 25 cm) of a square arena (50 x 50 cm, bright light illumination, 200 lux) and allowed to explore freely for 20 min. Total distance and distance in center were recorded using the ANY-maze™ Video Tracking System (version 4.50, Stoelting Co., Wood Dale, USA).

Elevated plus maze. Mice were placed for 5 min in maze consisting of two closed arms with 15cm high plastic walls and two orthogonally positioned open arms without walls (arm length 35 cm, width 5cm) elevated 110 cm above the floor under low-light conditions (10 lux). Total distance and distance on open arms were recorded using the ANY-maze™ Video Tracking System (version 4.50, Stoelting Co., Wood Dale, USA).

Novel object recognition. Following EPM, mice were familiarized to a Y-shaped maze consisting of 3 arms with walls (length 35 cm, width 6cm, height 14 cm) arranged at an 120° angle for 10 min. The next day, two wooden toy blocks were placed at the end of two arms and the test animal was placed in the empty arm to start an 8 min free explorations session under low light conditions (10 lux). The third arm stayed empty and served as starting point for the 8 min exploration. After a 3 h interval in the home cage, one of the wooden toy blocks was randomly interchanged with a Lego plastic ensemble serving as a novel object. All sessions were recorded via ANY-maze™ Video Tracking System (version 4.50, Stoelting Co., Wood Dale, USA) and contact times to objects were scored manually. A discrimination index was calculated as [(contact time novel object/total object contact time) - (contact time familiar object/total object contact time)], with an index close to 1 reflecting strong preference for a novel object, 0 no preference and close to -1 avoidance of a novel object.

Social interaction. Social preference and memory were tested in a three-compartment-chamber (chamber: width 40 cm x length 20 cm x height 40 cm) with grid tubes (diameter 8 cm x height 15 cm) in each outer chamber. Mice were first habituated to the arena by allowing free exploration for 5 min, then they were briefly restricted to the central compartment and a same sex interaction partner was introduced to one of the grid tubes. After 5 min of free exploration another new interaction partner was introduced to the second tube to test for social memory in 1 min session. Animals were video tracked using ANY-maze™ software and contact times to tube were scored manually offline. Discrimination indices were calculated for social preference [(contact time mouse tube/ total tube contact time) - (contact time empty tube/ total tube contact time)] and for social memory [(contact time new mouse tube/ total tube contact time) - (contact time familiar mouse tube/ total tube contact time)].

Active Avoidance. Animals were placed in a sound isolation cubicle containing a 16x32x20cm acrylic glass arena with a grid floor split in two compartments joined by a door, loudspeaker and ventilation fan (background noise 70dB SPL, light intensity <10 lux; TSE, Bad Homburg, Germany) and received 5 training days. Each training started with a 60 s habituation session to the box, followed by 50 trials consisting of a tone (85 dB, 10 kHz, max. 10 s) followed by footshock (0.4 mA) and a variable intertrial interval of 20 – 40s. Movement of animals was recognized by light beams and shuttling to another compartment during the tone prevented the footshock and was counted as successful avoidance, while shuttling after onset of the footshock was registered as an escape response. Failures, i.e. absence of any shuttling during a trial, was only occasionally observed (<1% of all trials) and not further analyzed. Number of defecation boli as a measure of arousal were counted after each trial.

On day 15, brains were prepared for a ***post-training gene expression analysis*** of somatostatin and parvalbumin in the dorsal ACC, dDG and BLA via qPCR. See Suppl. Material, Fig. S2 for details.

2.3. Experiment 2: Oscillations

In a different set of naive animals of both sexes, slices that contain the anterior cingulate cortex (ACC) were prepared (female GAD65+/+ animals N=8/ slices n=47; female GAD65-/- animals N=6/ slices n=36; male GAD65+/+ animals N=6/ slices n=30; male – GAD65-/- animals N=6/ slices n=30). Mice were decapitated under isoflurane anesthesia and their brains were quickly placed in ice-cold (-4 °C) standard artificial cerebrospinal fluid (aCSF; in mM: NaCl, 129;

NaH2PO4, 1.25; glucose, 10; MgCl2, 1; CaCl2, 2; KCl, 3; NaHCO3, 21; bubbled with 95% O2 / 5% CO2, pH≈7.4). Using a vibratome (Campden Instruments), coronal slices of 400 μm were obtained between 1.42 mm and -0.22 mm from Bregma (Paxinos and Franklin, 2001) and transferred to an interface chamber for 1 h incubation at 32 °C perfused with standard aCSF. Then, a modified aCSF containing 50 μM carbachol (CCh), 0.8 μM kainate (KA) and increased K^+ (~5 mM) concentration was perfused to induce fast local field potential (LFP) oscillations in the range of beta/gamma band, as described previously (Steullet et al., 2014). Forty-five min later borosilicate glass electrodes of ~1 MΩ resistance, filled with ACSF, were placed at the dorsal ACC (dACC) of layer 5 and the induced oscillatory activity was recorded for at least 10 min.

The signal was low-pass filtered at 3 kHz with the use of an extracellular amplifier (EXT-02F - Extracellular Amplifier; npi Electronic GmbH, Tamm, Germany). Data were recorded using the Spike2 software with a sampling rate of 5 kHz (CED-Micro1401; Cambridge Electronic Design, Cambridge, UK) and stored on a hard disc for offline analysis.

Artifact-free data from the last two minutes of the recordings were analyzed via computing fast Fourier transformation with a frequency resolution of 2.441 Hz revealing peaks in two frequency ranges at 10-25 Hz (beta oscillations) and 26-45 Hz (slow gamma oscillations). Two oscillation parameters were analyzed: 1) Peak Frequency (Hz): The frequency with the highest power at the corresponding frequency range (10-25 Hz and 26-45 Hz), 2) Integrated Power (mV^2/Hz): Sum of the binned power values at the corresponding frequency ranges (10-25 Hz and 26-45 Hz). For statistical comparison, only slices showing signals with peak powers higher than 5 μV^2 and peak frequencies higher than 10 Hz were included.

In an additional set of animals, gamma oscillations were induced in ventral-to-intermediate hippocampal-entorhinal slices via bath-application of 5 μM CCh and recorded in the CA3 and CA1 subregions (see Suppl. Material, Fig. S2 for methodological details).

2.4. Experiment 3: Interneuron density

Behaviorally naïve female and male GAD65-/- mice as well as their +/+ littermates of both sexes (n=6 per group) were transcardially perfused with phosphate buffered 4% paraformaldehyde solution (PFA). After overnight incubation in the same fixative and immersion in 30% sucrose solution at 4°C brains were frozen in liquid nitrogen-cooled methyl butane and stored at -20°C. Brains were then cryo sliced into 30 μm serial coronar sections at

the level of PFC (1.94- 1.54 mm from Bregma), posterior ACC (1.10 - -0.22 mm from Bregma) dorsal hippocampus / amygdala until ventral hippocampus (-1.46 - -2.70 mm from Bregma; Paxinos and Franklin, 2001). Free-floating sections were either incubated with a rabbit monoclonal anti-SST antibody (1:250, Santa Cruz Biotechnology #sc13099, Dallas, Texas, USA) or a combination of a rabbit monoclonal anti-PV antibody (1:500, abcam ab11427, Amsterdam, Netherlands) and a mouse monoclonal anti-GAD67 antibody (1:250; Millipore (Merck KgaA) MAB5406, Darmstadt, Germany) over night. A secondary antibody against rabbit (Alexa-Fluor 488, donkey anti rabbit IgG, 1:1000, Invitrogen (Thermofisher) A21206, Waltham, Massachusetts, USA) was used to label SST and PV, respectively and a biotinylated goat anti-mouse antibody (1:200, Jackson ImmunoResearch, Ely, UK), linked to Cy3 via Streptavidin incubation (1:1000, Jackson ImmunoResearch, Ely, UK), was used to label GAD67. Nuclei were visualized with DAPI (300 nM, Thermofischer, Waltham, MA, USA). Images of the immunostainings were acquired with a DMI6000 epifluorescence microscope (Leica, Wetzlar, Germany) and SST-, PV- and GAD67-positive cells were counted manually within selected brain areas using the software ImageJ. The cell number was normalized to area size of each region of interest (measured with imageJ). The cell counts per target area were averaged from 2 slices per animal covering both hemispheres and the averaged values per individual were used for statistical comparison between groups.

2.5. Statistics

Statistical analysis was performed using SPSS 22.0 (IBM Germany GmbH, Ehningen, Germany) and GraphPad Prism (version 9.1.2; GraphPad Software, CA, USA). Effects of sex or genotype and their interaction were analyzed by a two-way analysis of variance (ANOVA) in normally distributed data sets (tested by Shapiro–Wilk test), while in non-normal distributed samples a Mann–Whitney U-test was performed for the main factors sex and genotype. In case of significant interactions, a paired comparison of subgroups was performed using Student's T-test or Mann-Whitney-U-test. For data sets dependent measures of time and training trials (e.g. Open Field, Active Avoidance), repeated measure ANOVAs (RMA) were performed with sex and genotype as between-subject factors. Here, in case of sphericity issues Greenhouse-Geisser (Epsilon<0.8 at Mauchly's test of sphericity) or Huynh-Feldt corrections (Epsilon>0.8) were used. To assess successful avoidance learning, a one-sample T-test against chance level

of avoidances (50%) was performed. Differences were considered statistically significant if $p \leq 0.05$.

3. Results

3.1 Sex-specific effects of GAD65 deficiency on anxiety, social preference and active avoidance

Mice first performed an open field test under bright light conditions. The 20 min test phase was analyzed in 5 min bins via repeated measure ANOVA, revealing a significant interaction of time bin and genotype for total distance covered (Fig. 1A; RMA with Huynh-Feldt correction: $F(2.633,118.494)=8.783$, $p<0.001$), with GAD65-/- mice travelling higher distances towards the end of the test compared to their wildtype littermates, without additional effects of sex (RMA time bin x sex: $F(2.633,118.494)=0.896$, $p=0.434$) or interactions of sex and genotype (RMA time bin x sex x genotype: $F(2.633,118.494)=1.029$, $p=0.376$). Analyses of the distance covered only in the center revealed interactions of time bin x genotype (Fig. 1B; RMA: $F(3,135)=6.22$, $p=0.001$) and of time bin x sex x genotype ($F(3,135)=4.026$, $p=0.009$). Paired comparisons for the separate time bins (Fig. 1C) revealed then that especially male GAD65-/- vs. GAD65+/+ mice showed differences in the time course of anxiety-related behavior. While the relative distance in center was increased in male GAD65+/+ compared to male GAD65-/- mice at the beginning of the session ($T(23)=2.472$, $p=0.021$), male GAD65-/- mice subsequently increased their exploration of the center area and covered significantly more distance in the center towards the end of the test session ($T(23)=-2.785$, $p=0.011$ compared to male GAD65+/+. This differential response in males over the time course of the test drives also the significant main effect of sex observed regardless of time bins ($F(1,45) = 4.928$, $p=0.032$). In female mice, GAD65-/- and GAD65+/+ mice displayed comparable levels of activity in the during the first 5 min (paired comparison for genotype in females: $T(22)=0.571$, $p=0.574$) and the last 5 min ($T(22)=-0.326$, $p=0.748$) of the open field session.

During a 5 min session on the EPM under low light conditions following the next day, no genotype effects were observed. Male GAD65+/+ and GAD65-/- mice were generally less active than their female counterparts (Fig. 1D, total distance, two-way-ANOVA for effect of sex: $F(1,45)=7.65$, $p=0.008$; genotype: $F(1,45)=0.175$, $p=0.678$; sex x genotype: $F(1,45)=0.152$,

p=0.699). No significant differences in the distance travelled in the open arms was observed for sex or genotype (Fig. 1E two-way-ANOVA for effect of sex: $F_{(1,45)}=0.875$, p=0.355; genotype: $F_{(1,45)}=0.001$, p=0.982; sex x genotype: $F_{(1,45)}=0.958$, p=0.333).

Figure 1: Sex-specific effects of GAD65 deficiency on anxiety. (A) In a brightly lit open field**,** GAD65-/- mice of both sexes showed increased activity over time compared to their wildtype littermates. **(B)** For the activity in the center region, interactions of genotype and sex were observed in addition to an effect of genotype. **(C)** The detailed analysis of the first and last 5 min of the test revealed that male GAD65+/+ mice showed higher activity at the beginning of the open field session, while male GAD65-/- increased their center activity towards the end of the session. Over all time bins, female GAD65+/+ and GAD65-/- mice appear to explore the center less than their male counterparts. **(D)** During an elevated plus maze test under low light conditions, male mice of both genotypes were less active than their female counterparts. **(E)** No significant differences between genotypes and sexes were observed for the activity in the open arms. All data presented as mean ± SD and mean + SD with superimposed individual values for specific analysis time points. Two-way ANOVA: *** significant effect of genotype, p<0.001; # significant effect of sex, p<0.05; $$ significant interaction of sex x genotype, p<0.01.

Mice were then subjected to a novel object recognition task under low light conditions. Here, no significant effects of genotype or sex were observed, but male mice of both genotypes tended to avoid the novel object (Fig. 2A, two-way ANOVA, discrimination index: $F_{(1,45)}=3.965$, $p=0.053$). Discrimination within a social interaction task was achieved well by mice of all groups, with male mice showing higher social preference than females (mouse vs. empty tube; Fig. 2B; two-way-ANOVA for effect of sex: $F_{(1,45)}=8.928$, $p = 0.005$). Additionally, GAD65-/- mice showed a greater social preference than their wildtype littermates (two-way-ANOVA for effect of gene: $F_{(1,45)}=4.254$, $p=0.045$; effect of sex x genotype: $F_{(1,45)}=0.236$, $p=0.630$). Concordantly, the highest social preference was expressed by male GAD65-/- mice. When analyzing social memory by introducing an unfamiliar second interaction partner, male mice tended to prefer the unfamiliar partner more than females did (Fig. 2B; two-Way-ANOVA for effect of sex: $F_{(1,45)}=3.735$, $p=0.06$; effect of genotype: $F_{(1,45)}=0.973$, $p=0.329$; effect of sex x genotype: $F_{(1,45)}=0.067$, $p=0.797$).

Figure 2: Sex-specific effects of GAD65 deficiency on social preference. (A) In the novel object recognition task (NOR) mice explored two similar wooden toy blocks within a Y-maze. After a 3h interval, one of the objects was replaced with a novel object assembled from Lego. Regardless of sex and genotype, all mice did not prefer the novel object, male mice even tended to avoid the unfamiliar object as indicated by a negative discrimination index. **(B)** However, when mice had to discriminate between test tubes containing an unfamiliar mouse vs. an empty grid tube in the three-chamber social interaction task, all mice spent more time with the social interaction partner. Such social preference was increased in male compared to female mice and GAD65-/- mice compared to their GAD65+/+ littermates regardless of sex. Social memory tested by discriminating between a familiar and unfamiliar social interaction partner was neither affected by sex nor by genotype nor their interaction. All data presented as mean +/- SD with superimposed individual values. Two-way ANOVA: * significant effect of genotype, p<0.05; ## significant effect of sex, p < 0.01.

A significant effect of sex was evident during avoidance learning in the shuttle box. Female mice had lower numbers of escapes (Fig. 3A; RMA for escapes with Huynh-Feldt correction with Huynh-Feldt correction: main effect of sex: $F_{(1,43)}=4.066$, $p=0.05$; training day x sex: $F_{(3.767,161.978)}=1.678$, $p=0.159$; training day x genotype: $F_{(3.767,161.978)}=1.584$, $p=0.184$;

training day x sex x genotype: $F(3.767,161.978)=1.206$, $p=0.311$) and showed a trend to more successful avoidances than males (Fig. 3B; RMA for avoidances with Hyunh-Feldt correction: main effect of sex: $F(43)=3.883$; $p=0.055$; training day x sex: $F(3.731,160.436)=1.709$, $p=0.155$; training day x genotype: $F(3.731,160.436)=1.66$, $p=0.166$; training day x sex x genotype: $F(3.731,160.436)=1.296$, $p=0.275$). On the third training day the majority of female mice of both genotypes as well as of male GAD65+/+ mice performed above chance level and completed more than 50% of the trials with successful avoidances (Fig. 3B; one-sample T-test against chance level on day 3: F – GAD+/+: $T(11)=3.489$, $p=0.005$; F – GAD65-/-: $T(11)=2.916$, $p=0.014$; M – GAD65+/+: $T(11)=2.894$, $p=0.013$), but male GAD65-/- mice failed to reach the learning criterion on day 3 ($T(11) =-0.432$, $p=0.674$) and their poor performance persisted until day 5 ($T(119=1,833$, $p=0.098$). In addition, the number of defecation boli as an indicator of arousal and stress showed significant interactions of training day and sex (Fig. 3D; RMA: $F(4,168)=2.58$, $p = 0.039$) and of training day x sex x genotype ($F(4,168)=2.468$, $p=0.047$), but not of genotype alone (training day x genotype: $F(4,168)=1,076$, $p=0.370$). Paired comparison revealed that only in females a genotype effect was evident with increased boli counts in female GAD65-/- mice on day 1 (Fig. 3D; Student's T-test: $T(22)=-2.587$, $p=0.017$ vs. F – GAD65+/+). On day 5, boli counts were similarly lowered in female GAD65-/- and +/+ mice ($T(22)=-0.777$, $p=0.445$). Interestingly, paired comparison in GAD65+/+ only confirmed sex differences, with significantly higher boli counts in male compared to females wildtype mice on day 1 (Fig. 3D; Student's T-test: $T(23=-2.2.99$, $p=0.031$) and day 5 ($T(23)=-2.179$, $p=0.04$).

Figure 3: Sex-specific effects of GAD65 deficiency on active avoidance in a 2-way shuttle box. (A) Female mice reduced the number of escapes (shuttling after footshock) over five days of training days faster than males. **(B)** In trend (p=0.055 for effect of sex), the number of avoidances (successful shuttling during tone presentation before footshock onset) increased more in females than males. **(C)** Accordingly, on day 3 of the training, female GAD65+/+ and GAD65-/- mice as well as male GAD65+/+ mice performed significantly above chance level, while male GAD65-/- mice did not reach this criterion even on day 5. **(D)** The number of defecation boli after each session as a measure of arousal and perceived stress varied as function of the training days especially for female GAD65-/- mice, resulting in a significant effect of genotype and an interaction of sex x genotype. **(E)** A detailed analysis of boli counts on day 1 revealed lowest counts in female GAD65+/+ mice compared to female GAD65-/- mice and to male GAD65+/+ mice. On day 5, female GAD65-/- mice displayed comparable boli counts to their female GAD65+/+ littermates, while no reduction was observed in male mice. All data presented as mean ± SD and mean + SD with superimposed individual values for specific analysis time points. Two-way ANOVA: # significant effect of sex, p < 0.05; $ significant interaction of sex x genotype, p<0.05. One-sided T-test against chance level of avoidances (color-coded for respective group): + p<0.05; Paired comparison (T-Test): * (in grey) significant effect of genotype within the female subgroup, p<0.05; # (in grey) significant effect of sex within the GAD65+/+ subgroup, p<0.05.

Taken together, general sex differences were evident in mildly ambiguous environments like a low lit elevated plus maze and in a novel object recognition task, with male mice generally showing less activity and a tendency to avoid novel objects. In more challenging situations like a brightly lit open field or in an active avoidance task, sex dependent effects of GAD65 deficiency became evident. In the open field, male GAD65-/- mice showed a protracted reduction of anxiety-like behavior and the performed less well in the active avoidance task compared to all other groups. Female GAD65-/- showed no modulation of anxiety like-behavior in the open field, but performed well in the active avoidance task and even adapted their arousal response as indicated by boli counts.

3.2. Augmented beta/gamma oscillations in the ACC of GAD65-/- mice

In a separate cohort of animal *ex vivo* CCh-and KA-induced oscillations were assessed in the ACC (Fig. 4A-C) at two frequency ranges: 10-25 Hz (beta oscillations) and 26-45 Hz (slow gamma oscillations). A two-way ANOVA analysis revealed significant interaction of sex x genotype on the peak frequency at the beta range (Fig. 4D; two-way ANOVA for 10-25 Hz: interaction of sex x genotype: $F_{(1,139)}=5.088$, $p=0.0256$; effect of sex: $F_{(1,139)}=1.308$, $p=0.2547$; effect of genotype: $F_{(1,139)}=0.01773$, $p=0.8943$), but a paired comparison for each sex did not demonstrate significant differences between female GAD65+/+ and GAD65-/- mice ($T_{(81)}=1.699$, $p=0.0931$) or between male GAD65+/+ and GAD65-/- mice ($T_{(58)}=1.595$, $p=0.116$). At the slow gamma range, 26-45 Hz, a significant effect of sex was observed with slightly increased peak frequencies for male compared to female mice (Fig. 4D; two-way ANOVA for 26-45 Hz: effect of sex: $F_{(1,139)}=4.016$, $p=0.047$; effect of genotype: $F_{(1,139)}=1.022$, $p=0.314$; interaction of sex x genotype: $F_{(1,139)}=1.526$, $p=0.219$).

The integrated power of oscillations at 10-25 Hz and 26-45 Hz was increased in GAD65-/- mice of both sexes (Fig. 4E; genotype effect for 10-25 Hz: $F_{(1,139)}=11.42$, $p=0.0009$; 26-45 Hz: $F_{(1,139)}=4.594$, $p=0.034$), without further effects of sex (10-25 Hz: $F_{(1,139)}=0.3055$, $p=0.581$; 26-45 Hz: $F_{(1,139)}=0.9057$, $p=0.343$) or interactions of sex x genotype (10-25 Hz: $F_{(1,139)}=0.3618$, $p=0.5485$; 26-45 Hz: $F_{(1,139)}=0.311$, $p=0.578$).

Figure 4: Increased propensity for induced fast oscillations in the ACC of GAD65-/- mice. (A) Scheme of an ACC slice and electrode positions for network oscillation recording induced by carbachol (50 µM) and kainate (0.8 µM) administration. **(B)** Representative traces of unprocessed data (5 kHz), low-pass filtered (<25 Hz) for illustration of beta-dominant LFP and band-pass filtered (25-45 Hz) for illustration of slow gamma oscillations *in vitro*. **(C)** Example power spectra of female and male mice across oscillation frequencies with two main peak frequencies in the LFP signal; approx. at 15 Hz and 35 Hz. **(D)** A significant interaction of sex x genotype was observed for the peak frequency at 10-25 Hz, while a sex effect was evident for the peak frequency at 26-45 Hz. **(E)** The integrated power of fast oscillations at the 10-25 Hz and 26-45 Hz frequency ranges were increased in GAD65-/- mice compared to their wildtype littermates, irrespective of their sex. All data presented as mean ± SD with superimposed individual values. Two-way ANOVA: $ significant interaction of sex x genotype, p<0.05; # significant effect of sex, p<0.05; * significant effect of genotype, p<0.05; *** p<0.001.

Ex vivo gamma oscillations induced by CCh (5 µM) were further assessed in slices containing the ventral-to-intermediate CA3 and CA1 subregions (see supplementary material, Fig. S1). Here, the peak frequency of gamma oscillations (20-80 Hz) was significantly modulated by an interaction of sex x genotype in vCA3 (Fig. S1C; two-way-ANOVA: interaction of sex x genotype: $F_{(1,70)}=6.19$, p=0.015), with a significant effect of genotype ($F_{(1,70)}=4.676$, p=0.034), but not sex ($F_{(1,70)}=0.131$, p=0.719). Paired comparison revealed that male GAD65-/- mice showed higher gamma oscillation peak frequencies compared to male GAD65+/+ littermates (Fig. S1C; Student's T-test: T(31)=3.274, p=0.003), while females showed comparable levels (T(39)=0.235, p=0.815). The peak frequency in vCA1 was not significantly affected by sex (Fig. S1C; Two-way ANOVA: $F_{(1,70)}=0.1096$, p=0.742), genotype ($F_{(1,70)}=3.642$, p=0.0605) nor by the interaction of sex x genotype ($F_{(1,70)}=2.796$, p=0.099). A significant interaction of sex x genotype was also observed for the integrated power of gamma oscillations in vCA3 (Fig. S1D; two-way-ANOVA: interaction of sex x genotype:

F(1,70)=4.842, p=0.0311; effect of sex: F(1,70)=0.9391, p=0.336; effect of genotype: F(1,70)=0.0997, p=0.753), but a paired comparison of genotype effects within each sex did not reveal significant differences (Fig. S1D; Student's T-Test for genotype in females: T(39)=1.525, p=0.135; in males: T(31)=1.555, p=0.130). No significant effects were observed for the integrated power in vCA1 (Fig. S1D; two-way-ANOVA: interaction of sex x genotype: F(1,70)=3.329, p=0.072; effect of sex: F(1,70)= 1.933, p=0.169; effect of genotype: F(1,70)=1.871, p=0.176). Of note, the synchronization of gamma activity between vCA3 and vCA1 was significantly affected by sex (Fig. S1E; two-way-ANOVA, effect of sex: F(1,65)=9.598, p=0.003; effect of genotype: F(1,65)=0.1851, p=0.668; interaction of sex x genotype: F(1,65)=1.751, p=0.19).

Together, electrophysiological recordings demonstrate a rather genotype-mediated effect for ACC beta/gamma oscillations (increased oscillation power in both male and female GAD65 -/- mice) while hippocampal gamma oscillation properties appear driven by sex differences (increased CA3-CA1 gamma correlation in males). These results suggest a potentially altered interneuron function in these brain regions relevant for emotional behaviors due to their involvement in generation of fast LFP oscillations.

3.3. Altered interneuron density in anxiety-related brain areas of GAD65-/- mice

Oscillatory activity is shaped by the activity of different interneuron subpopulations, mainly by PV-positive interneurons (Gulyás et al., 2010). Therefore, the number of these interneurons as well as SST-positive cells for comparison were analyzed in brain areas relevant for emotional behavior. Two-way ANOVA revealed an increased number of PV-positive cells in the ACC of GAD65-/- mice regardless of sex (Fig. 5A; effect of genotype: F(1,20)=8.301, p=0.009; effect of sex: F(1,20)<0.001, p=0.988; interaction sex x genotype: F(1,20)=0.671, p=0.422). The number of GAD67-postive cells, another isoform of GAD found ubiquitously in the cytosol of interneurons, was reduced in GAD65-/- mice (Fig. 5A; Mann-Whitney for main effects for non-normally distributed data: for genotype: U=-2.57, p=0.008; for sex: U=0.346, p=0.755). However, no significant change in the number of SST-positive cells was found in this area (Fig. 5A; effect of genotype: F(1,20)=2.325, p=0.143; effect of sex: F(1,20)=2.453, p=0.133; interaction sex x genotype: F(1,20)=1.682, p=0.209).

In the dDG, by contrast, a main effect of sex (Fig. 5B; F(1,20)=6.64, p=0.018) and a significant interaction of genotype x sex (F(1,20)=5.715, p=0.027) were observed for the number of SST-

positive cells, without additional effect of genotype (F(1,20)=0.303, p=0.588). Paired comparison revealed that male GAD65-/- mice had significantly less SST-positive cells in the DG compared to female GAD65-/- mice (T(10)=2.983, p=0.014) and to their male wildtype littermates (T(10)=2.633, p=0.025). The number of PV-positive and GAD67-postive cells did not differ in this area (Fig. 5B; Mann-Whitney for main effects for non-normally distributed data: PV – sex: U=-1.501, p=0.143; PV – genotype: U=-0.289, p=0.799; GAD67 – sex: U=-0.808, p=0.443; GAD67 – genotype: U=0.404, p=0.713).

A main effect of genotype was also observed for the number of SST-positive interneurons in the BLA (Fig. 5C; F(1,20)=6.486, p=0.019), while main effects of sex only reached a trend for reduced cell numbers in males (F(1,20)=4.299, p=0.051) and no interactions of sex x genotype (F(1,20)=2.098, p=0.163) were observed. However, as demonstrated by a paired comparison, a reduction of SST-positive cells was evident only in male GAD65-/- mice (T(10)=3.978, p=0.003), but not in female GAD65-/- mice (T(10)=0.635, p=0.54) compared to their respective same-sex GAD65+/+ littermates. The number of PV-positive and GAD67-postive cells did not differ within the BLA (Fig. 5C; two-way ANOVA for PV: effect of sex: F(1,20)=0.791, p=0.384; genotype: F(1,20)=0.032, p=0.859; sex x genotype: F(1,20)=0.118, p=0.735; two-way ANOVA for GAD67: effect of sex: F(1,20)=2.108, p=0.162; genotype: F(1,20)=0.249, p=0.623; sex x genotype: F(1,20)=1.802, p=0.194).

Figure 5: Altered interneuron density in anxiety-related brain areas of GAD65-/- mice. (A) In the posterior part of the anterior cingulate cortex (ACC), the number of GAD67- and parvalbumin (PV)-positive cells was increased in GAD65-/- mice of both sexes. Representative images of PV and GAD67 double immunostainings in the posterior ACC are shown on the right side. **(B)** In the dorsal dentate gyrus (dDG) the number of somatostatin (SST)- positive interneurons was generally lower in males, but interactions of genotype x sex were revealed. Paired comparisons demonstrated lower cell counts especially in male GAD65-/- mice. Representative images of SST immunostainings in the dDG are shown on the right side**. (C)** In the basolateral amygdala (BLA), SST-positive interneurons were generally reduced in GAD65-/- mice, but paired comparison confirmed that this cell population was reduced in especially in male GAD65-/- mice as well. All data presented as mean ± SD with superimposed individual values. Two-way ANOVA: * significant effect of genotype, p<0.05, **p<0.01; # # significant effect of sex, ## p<0.01; $$ significant interaction of sex x genotype, p < 0.01. Paired comparison (T-Test): * (in grey) significant effect of genotype within the female subgroup, p<0.05, ** (in grey) p<0.01; # (in grey) significant effect of sex within the GAD65+/+ subgroup, p<0.05.

In addition to interneuron counts, mRNA expression levels for PV and SST were assessed in the same areas in mice that went through the behavioral testing battery (see Supplementary material, Fig. S2). Here, no mRNA expression differences were observed in the ACC, but the expression of PV mRNA was reduced in the dDG of males (Fig. S2B; two-way ANOVA, main

effect of sex: F(1,28)=6.973, p=0.013) as well as in the BLA of males (Fig. S2C; two-way ANOVA, main effect of sex: F(1,28)=5.112, p=0.036). In addition, expression of SST was increased in the BLA of male mice (Fig. S2C; Mann-Whitney for main effects for non-normally distributed data: main effect of sex U=2.569, p=0.009).

Together, the number of parvalbumin-positive cells was increased in the ACC of GAD65-/- mice regardless of genotype, whereas the number of somatostatin-positive interneurons was reduced in the dDG and BLA of male GAD65-/-. No significant effects of genotype, sex or their interaction were observed on the number of PV- and SST-positive interneurons in vCA1 and vCA3 (Supplementary material, Fig. S3).

4. Discussion

In this study we investigated sex-specific differences in emotional information processing related to anxiety and threat avoidance in GAD65-/- mice that display a postnatal GABA deficit. To date, mainly male GAD65-/- have been investigated, demonstrating increased anxiety, generalization of fear memory and deficits in fear extinction (see Müller et al., 2015 for review). However, epidemiological studies in human suggest sex differences with an increased incidence of anxiety disorders in women (McLean & Anderson, 2009; Vesga-Lopez et al., 2008). In accordance, in our rodent study, we found sex differences in anxiety, discrimination learning and avoidance that were modulated by GAD65 deficiency.

In an open field GAD65-/- mice of both sexes stayed more active, in line with previous results in males (Stork et al., 2000). Male GAD65-/- mice increased their activity in the center of the arena over time, suggesting an adaptive anxiety-like response. With a short exposure and less aversive conditions under low light in the EPM, no difference in anxiety-like behavior was revealed and locomotor activity was even higher in females than males, regardless of the genotype. Previous studies indeed report that bright lighting conditions increase anxiety-like behavior and may even induce hyperactivity in males (Strekalova, 2005), in line with our observation in male wildtypes. Increased anxiety can potentially interact with cognitive tasks such as the novel object recognition test. Here, male mice showed a trend for a negative discrimination and avoided novel objects. In these mice, extensive pre-habituation to objects and test environments may be required to overcome neophobia and avoid emotional bias when assessing cognition. When introducing social interaction partners instead of

unanimated objects, mice of both sexes and genotypes preferred the partner mouse, despite reduced overall interaction times in female GAD65-/- mice (data not shown). In line with previous studies (Stack et al., 2010), male mice showed greater social preference that may also relate to aggression and an intensified interaction with the partner mouse restricted to a test tube. Social preference was enhanced in GAD65-/- mice compared to their wildtype littermates. In previous, studies male GAD65-/- mice showed reduced aggressive encounters that could be linked to increased anxiety as well (Stork et al., 2000), but in our setting with restricted access to interaction partners GAD65-/- mice may adapt their anxiety-like response. To assess learning and memory in a more emotionally salient situation, a two-way avoidance paradigm in a shuttle box was used. Here, sex effects were revealed over training sessions. In accordance with previous studies females performed better in the active avoidance task (Dalla and Shors, 2009). A more detailed analysis of successful avoidances demonstrated that male GAD65-/- mice showed the lowest success rate of avoidances even after 5 days of training. Checking on fecal boli as a parameter for successful stress and anxiety adaptation due to mastering the cognitive task during training (Hadad-Ophir et al., 2017; Sturman, 2018), female GAD65+/+ mice had low boli counts throughout the training sessions and female GAD65-/- mice reduced the number of boli with successful training. Male mice of both genotypes showed no such adaptation. Thus, in contrast to the social memory or object discrimination tasks, females performed better in this aversely motivated learning paradigm. Male GAD65-/- mice, however, adapted their anxiety-like response in the open field test, but failed to adapt during this emotional learning task.

Adapting cognitive demands in emotional challenging situations relies on the function of a limbic network comprising the amygdala, the hippocampus and their interaction with frontal cortical areas. The ACC is an important part of the anxiety network and involved in the assessment of emotional situations and the appropriate choice of action, especially in stressful situations (Walton, 2007). It is active during threat perception (Straube, 2009) and a hyperactivity of this region, together with the amygdala and the ventral prefrontal cortex, is observed in patients with a generalized anxiety disorder (McClure et al., 2007). During attentional processes, oscillations at the gamma frequency range are registered in the frontal cortex and the hippocampus (Voloh et al., 2015; Rothé et al., 2011). To test whether oscillatory activity at gamma range is also altered in the ACC of male and female GAD65-/- and GAD65+/+ mice, we used a reductionist slice model that depends on activation of cholinergic and

glutamatergic transmission upon co-perfusion of carbachol and kainate (Steullet et al., 2014). *Ex vivo* ACC oscillations are sensitive to behavioral manipulations such as chronic stress (Ito et al., 2020) and were robustly induced at the beta (10-25 Hz) and slow gamma range (26-45 Hz) in slices of all groups (Fig. 4). Phasic inhibition provided by PV-positive interneurons play a fundamental role in sustaining and pacing *ex vivo* LFP oscillations in the hippocampus (Gulyas et al., 2010) but also in various cortical structures including ACC (Cabungcal et al., 2013; Steullet et al., 2014). Specifically, enhanced *ex vivo* beta/gamma power in the ACC was observed upon degradation of the perineuronal network that enwraps PV-positive interneurons (Steullet et al., 2014) and low PV-positive cell number was associated with reduction of *ex vivo* ACC beta/gamma oscillation power (Cabungcal et al., 2013). In accordance with these observations, we found an enhanced beta/gamma oscillation power and increased number of PV-positive interneurons in the ACC of GAD65-/- male and female mice. Interestingly, the observed reduction in the density of pan-interneuronal marker GAD67 - in the ACC of GAD65-/- mice may suggest a generally increased excitability via disinhibition of excitatory pyramidal neurons which could also contribute the enhanced LFP oscillations in the ACC. Thus, it appears that GAD67 as another isoform producing GABA in the cytosol of interneurons does not compensate for GAD65 deficiency, as indicated by the reduced GABA content in cortical areas of these mice (Stork et al, 2000). In the ventral hippocampus augmented gamma oscillations have been reported upon elevation of anxiety levels and aberrant gamma hypersynchrony is linked to symptom severity in post-traumatic stress disorder (PTSD) (Çalışkan and Stork, 2019; Dunkley et al., 2014). While we found only tendential effects for the increase of gamma power in the ventral CA3 region of male GAD65-/- mice, we observed an enhanced CA3-CA1 gamma hypersynchrony in male compared to female mice, regardless of a GAD65 deficiency. This may be linked to increased anxiety-like behavior of males in specific tests, such as decreased EPM activity and higher novel object avoidance. No sex-specific effects on the number of PV-positive cells were observed in the ventral CA3 or CA1 area. Sex-specific effects, however, were observed for another class of interneurons. The number of SST-positive was reduced in the BLA and dDG especially in male GAD65-/- mice. Indeed, the DG is relevant for active avoidance learning and anxiety-like behavior as well and a recent study using local knock down strategies support a role for GAD65 in these behaviors (Tripathi et al., 2021). Moreover, previous studies demonstrate anxiolytic actions of the peptide SST via the BLA (Gaskin et al., 2021) and analyzing mRNA levels four

days after the last active avoidance training (Fig. S2) revealed a small increase in SST levels in the BLA of male mice, paralleled by decreased PV mRNA in this region.

Finally, we identified an increase in peak frequency of ACC (males and females) and hippocampal CA3 (only males) gamma oscillations in GAD65-/- mice which indicates faster gamma oscillation cycles via altered excitatory-inhibitory interactions. Here, a more detailed cell-type-specific analysis of GABA-A receptors subtypes that determine the frequency of fast oscillations (Heistek et al., 2010) might help to further resolve cellular correlates of gamma peak frequency alterations in the ACC and hippocampal CA3.

5. Conclusions and further perspectives

In summary, our analysis of behavior and network activity as well as interneuron density in male and female GAD65 mice revealed adaptive anxiety-like responses in a brightly lid open field as well as deficits in avoidance learning especially in male GAD65-/- mice. Female GAD65 -/- mice appeared more similar to their wildtype littermates and performed well in the active avoidance task. Associated with their deficit in avoidance learning, male GAD65-/- mice had lower cell counts of SST-positive interneurons in the dDG and the BLA and, despite unchanged interneuron numbers, gamma frequencies in ventral CA3 were increased. GAD65-/- mice of both sexes displayed higher densities of PV-positive interneurons in the ACC, together with increased gamma power in this region. On the behavioral level, this genotype effect regardless of sex corresponded to the increased activity on GAD65-/- mice in the open field. On top of this, our study found several sex-specific effects regardless of the genotype, among them a decreased EPM activity, higher novel object avoidance and increased social preference in males compared to females, in association with an increased peak frequency of gamma oscillation in the ACC as well as a heightened correlation of gamma activity in the ventral CA3 and CA1 areas. This suggests that sex-dependent changes in anxiety-like behavior in ambiguous environments and active avoidance learning may rely on a GAD65-dependent network involving the BLA, dorsal and ventral hippocampal regions as well as their interaction with the ACC that may be driven by fast oscillatory activity.

Due to a possible stress induction by taking vaginal smear samples for estrus analysis in female mice, we refrained from correlating behavior, neuronal activity and expression values with estrous stages. Nevertheless, studies in cell culture and in vivo confirm that estrogen and its derivates can regulate the expression levels of GAD65 and other important co-

neuromodulators in interneurons such as NPY or PV activity (Nakamura et al., 2004; Nakamura & McEwen, 2005; Ikeda et al., 2015; Sandhu et al., 2015; Picard et al., 2019). Therefore, we cannot rule out that individual variability in females in the analyzed responses might stem from different stages of the estrous cycle. Future studies are required that systematically address estrous phase effects on interneuron activity in limbic brain regions and their impact on anxiety-like and avoidance behavior. In addition, while this study engaged transgenic mice with a brain-wide GAD65 deficiency, regional effects of altered GAD65 expression can be further studied with local viral knock down tools and current analysis in this direction should be expanded to females (e.g. Tripathi et al., 2021; Albrecht et al., 2022).

Nevertheless, using a transgenic GAD65 deficient mouse allows to study genetic variation in GABA levels and their relevance for the function of anxiety networks in emotional challenging situations. In humans various SNPs that affect GABA levels are described. One of those variants results in increased GABA levels especially in the ACC, as revealed by fMRI-SPECT, and was associated to increased harm avoidance behavior in females but not males as an anxiety-related trait (Colic, 2018). While functional imaging studies indeed pinpoint to the ACC as an important region for emotion regulation (Mochkovotich et al., 2014; Goossen et al., 2019; Kim & Kim, 2021), sex-specific regulation differences are not well studied. Interestingly in female, but not male healthy volunteers ACC volume and positive coping style predicted lower anxiety- or depression-related psychopathologies (Holz et al., 2016). In our animal model of lowered GABA levels due to GAD65 deficiency, we found that female GAD65-/- mice showed a more stable emotional response in challenging situations such as the brightly lid open field or the active avoidance challenge. Deficits, especially in avoidance behavior, were more pronounced in male GAD65-/- mice. Females may regulate their avoidance behavior differently from males in emotionally challenging situations, at least in part, via a GABAergic impact on their anxiety network structures. Further studies in healthy volunteers and anxiety patients as well as targeted intervention studies in relevant animal models are needed to investigate such sex-dependent biases in ACC function and its interaction with other structures such as the hippocampus and BLA and to investigate a possible modulation of such anxiety networks by GABAergic neurotransmission.

Acknowledgement

We thank Franziska Blitz, Antje Koffi von Hoff, Vivian Sand and Annika Lenuweit for excellent technical assistance.

Funding information

This work was supported by grants from the Leibniz Postdoctoral Network fellowship to Anne Albrecht (SAW-2015-LIN-3), from the German Research Foundation (Projects CRC779-B5 and 362321501/RTG 2413 SynAGE to Oliver Stork; Project-ID 425899996 – CRC 1436 to Anne Albrecht and Oliver Stork) and from the Center for Behavioural Brain Sciences Magdeburg - CBBS funded by the European funds for regional development (EFRE, Funding Nr ZS/2016/04/78113). Evangelia Pollali is a PhD student of ESF graduate school ABINEP (Funded by the federal state Saxony-Anhalt and the European Structural and Investment Funds, project number ZS/2016/08/80645 to Oliver Stork).

Declaration of interest

None

References

Admon, R., Milad, M. R., & Hendler, T. (2013). A causal model of post-traumatic stress disorder: disentangling predisposed from acquired neural abnormalities. *Trends in Cognitive Sciences*, *17*(7), 337–347. https://doi.org/10.1016/j.tics.2013.05.005

Albrecht, A., Müller, I., Weiglein, A., Pollali, E., Çalışkan, G. & Stork O. (2022). Choosing memory retrieval strategies: A critical role for inhibition in the dentate gyrus. *Neurobiol Stress*, 20:100474. https://doi.org/10.1016/j.ynstr.2022.100474

American Psychiatric Association (Ed.) (2013). *Diagnostic and Statistical Manual of Mental Disorders: Anxiety disorders* (5th). Washington DC.

Blanco, C., Rubio, J., Wall, M., Wang, S., Jiu, C. J., & Kendler, K. S. (2014). Risk factors for anxiety disorders: common and specific effects in a national sample. *Depression and Anxiety*, *31*(9), 756–764. https://doi.org/10.1002/da.22247

Cabungcal, J.-H., Steullet, P., Morishita, H., Kraftsik, R., Cuenod, M., Hensch, T. K., & Do, K. Q. (2013). Perineuronal nets protect fast-spiking interneurons against oxidative stress. *Proceedings of the National Academy of Sciences*, *110*(22), 9130–9135. https://doi.org/10.1073/pnas.1300454110

Çalışkan, G., Müller, I., Semtner, M., Winkelmann, A., Raza, A. S., Hollnagel, J. O., . . . Meier, J. C. (2016). Identification of parvalbumin interneurons as cellular substrate of fear memory persistence. *Cerebral Cortex*, *26*(5), 2325–2340. https://doi.org/10.1093/cercor/bhw001

Çalışkan, G., & Stork, O. (2019). Hippocampal network oscillations at the interplay between innate anxiety and learned fear. *Psychopharmacology*, *236*(1), 321–338. https://doi.org/10.1007/s00213-018-5109-z

Chen, Y., Hu, N., Yang, J., & Gao, T. (2022). Prefrontal cortical circuits in anxiety and fear: an overview. *Frontiers of Medicine*, 1–22. https://doi.org/10.1007/s11684-022-0941-2

Colic, L., Li, M., Demenescu, L. R., Li, S., Müller, I., Richter, A., . . . Schott, B. H. (2018). GAD65 promoter polymorphism rs2236418 modulates harm avoidance in women via inhibition/excitation balance in the rostral ACC. *Journal of Neuroscience*, *38*(22), 5067–5077. https://doi.org/10.1523/JNEUROSCI.1985-17.2018

Dalla, C., & Shors, T. J. (2009). Sex differences in learning processes of classical and operant conditioning. *Physiology & Behavior*, *97*(2), 229–238. https://doi.org/10.1016/j.physbeh.2009.02.035

Dunkley, B. T., Doesburg, S. M., Sedge, P. A., Grodecki, R. J., Shek, P. N., Pang, E. W., & Taylor, M. J. (2014). Resting-state hippocampal connectivity correlates with symptom severity in post-traumatic stress disorder. *NeuroImage: Clinical*, *5*, 377–384. https://doi.org/10.1016/j.nicl.2014.07.017

Fee, C., Prevot, T. D., Misquitta, K., Knutson, D. E., Li, G., Mondal, P., . . . Sibille, E. (2021). Behavioral deficits induced by somatostatin-positive GABA neuron silencing are rescued by alpha 5 GABA-A receptor potentiation. *International Journal of Neuropsychopharmacology*, *24*(6), 505–518. https://doi.org/10.1093/ijnp/pyab002

Goossen, B., van der Starre, J., & van der Heiden, C. (2019). A review of neuroimaging studies in generalized anxiety disorder:"So where do we stand?". *Journal of Neural Transmission*, *126*(9), 1203–1216. https://doi.org/10.1007/s00702-019-02024-w

Gulyás, A. I., Szabó, G. G., Ulbert, I., Holderith, N., Monyer, H., Erdélyi, F., . . . Hájos, N. (2010). Parvalbumin-containing fast-spiking basket cells generate the field potential oscillations induced by cholinergic receptor activation in the hippocampus. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *30*(45), 15134–15145. https://doi.org/10.1523/JNEUROSCI.4104-10.2010

Hadad-Ophir, O., Ardi, Z., Brande-Eilat, N., Kehat, O., Anunu, R., & Richter-Levin, G. (2017). Exposure to prolonged controllable or uncontrollable stress affects GABAergic function in sub-regions of the hippocampus and the amygdala. *Neurobiology of Learning and Memory*, *138*, 271–280. https://doi.org/10.1016/j.nlm.2016.06.009

Heistek, T. S., Timmerman, J., Spijker, S., Brussaard, A. B., & Mansvelder, H. D. (2010). GABAergic synapse properties may explain genetic variation in hippocampal network oscillations in mice. *Frontiers in Cellular Neuroscience*, 18. https://doi.org/10.3389/fncel.2010.00018

Holz, N. E., Boecker, R., Jennen-Steinmetz, C., Buchmann, A. F., Blomeyer, D., Baumeister, S., . . . Meyer-Lindenberg, A. (2016). Positive coping styles and perigenual ACC volume: two related mechanisms for conferring resilience? *Social Cognitive and Affective Neuroscience*, *11*(5), 813–820. https://doi.org/10.1093/scan/nsw005

Ikeda, T., Makino, Y., & Yamada, M. K. (2015). 17α-estradiol is generated locally in the male rat brain and can regulate GAD65 expression and anxiety. *Neuropharmacology*, *90*, 9–14. https://doi.org/10.1016/j.neuropharm.2014.10.019

Ito, R., Nakano, T., Hojo, Y., Hashizume, M., Koshiba, M., & Murakoshi, T. (2020). Chronic restraint stress affects network oscillations in the anterior cingulate cortex in mice. *Neuroscience*, *437*, 172–183. https://doi.org/10.1016/j.neuroscience.2020.04.021

Javanbakht, A., Kim, P., Swain, J. E., Evans, G. W., Phan, K. L., & Liberzon, I. (2016). Sex-specific effects of childhood poverty on neurocircuitry of processing of emotional cues: a neuroimaging study. *Behavioral Sciences*, *6*(4), 28. https://doi.org/10.3390/bs6040028

Kash, S. F., Tecott, L. H., Hodge, C., & Baekkeskov, S. (1999). Increased anxiety and altered responses to anxiolytics in mice deficient in the 65-kDa isoform of glutamic acid decarboxylase. *Proceedings of the National Academy of Sciences*, *96*(4), 1698–1703. https://doi.org/10.1073/pnas.96.4.1698

Kim, N., & Kim, M. J. (2021). Altered Task-Evoked Corticolimbic Responsivity in Generalized Anxiety Disorder. *International Journal of Molecular Sciences*, *22*(7), 3630. https://doi.org/10.3390/ijms22073630

Lépine, J.-P. (2002). The epidemiology of anxiety disorders: prevalence and societal costs. *Journal of Clinical Psychiatry*, *63*, 4–8.

McClure, E. B., Monk, C. M., Nelson, E. E., Parrish, J. M., Adler, A., Blair, J. R., . . . Pine, D. S. (2007). Abnormal Attention Modulation of Fear Circuit Function in Pediatric Generalized Anxiety Disorder. *Arch Gen Psychiatry*. Advance online publication. https://doi.org/10.1001/archpsyc.64.1.97

McLean, C. P., & Anderson, E. R. (2009). Brave men and timid women? A review of the gender differences in fear and anxiety. *Clinical Psychology Review*, *29*(6), 496–505. https://doi.org/10.1016/j.cpr.2009.05.003

Miyata, S., Kumagaya, R., Kakizaki, T., Fujihara, K., Wakamatsu, K., & Yanagawa, Y. (2019). Loss of glutamate decarboxylase 67 in somatostatin-expressing neurons leads to anxiety-like behavior and alteration in the Akt/GSK3β signaling pathway. *Frontiers in Behavioral Neuroscience*, *13*, 131. https://doi.org/10.3389/fnbeh.2019.00131

Mochcovitch, M. D., da Rocha Freire, Rafael Christophe, Garcia, R. F., & Nardi, A. E. (2014). A systematic review of fMRI studies in generalized anxiety disorder: evaluating its neural and cognitive basis. *Journal of Affective Disorders*, *167*, 336–342. https://doi.org/10.1016/j.jad.2014.06.041

Müller, I., Çalışkan, G., & Stork, O. (2015). The GAD65 knock out mouse–a model for GABAergic processes in fear- and stress-induced psychopathology. *Genes, Brain and Behavior*, *14*(1), 37–45. https://doi.org/10.1111/gbb.12188

Müller, I., Obata, K., Richter-Levin, G., & Stork, O. (2014). GAD65 haplodeficiency conveys resilience in animal models of stress-induced psychopathology. *Frontiers in Behavioral Neuroscience*, *8*, 265. https://doi.org/10.3389/fnbeh.2014.00265

Nakamura, N. H., & McEwen, B. S. (2005). Changes in interneuronal phenotypes regulated by estradiol in the adult rat hippocampus: a potential role for neuropeptide Y. *Neuroscience*, *136*(1), 357–369. https://doi.org/10.1016/j.neuroscience.2005.07.056

Nakamura, N. H., Rosell, D. R., Akama, K. T., & McEwen, B. S. (2004). Estrogen and ovariectomy regulate mRNA and protein of glutamic acid decarboxylases and cation-chloride cotransporters in the adult rat hippocampus. *Neuroendocrinology*, *80*(5), 308–323. https://doi.org/10.1159/000083657

Ouimet, A. J., Gawronski, B., & Dozois, D. J. A. (2009). Cognitive vulnerability to anxiety: A review and an integrative model. *Clinical Psychology Review*, *29*(6), 459–470. https://doi.org/10.1016/j.cpr.2009.05.004

Paxinos, G & Franklin, K.B.J. (2001). The mouse brain in stereotaxic coordinates. San Diego, CA: Academic Press

Picard, N., Takesian, A. E., Fagiolini, M., & Hensch, T. K. (2019). NMDA 2A receptors in parvalbumin cells mediate sex-specific rapid ketamine response on cortical activity. *Molecular Psychiatry*, *24*(6), 828–838. https://doi.org/10.1038/s41380-018-0341-9

Rothé, M., Quilodran, R., Sallet, J., & Procyk, E. (2011). Coordination of high gamma activity in anterior cingulate and lateral prefrontal cortical areas during adaptation. *Journal of Neuroscience*, *31*(31), 11110–11117. https://doi.org/10.1523/JNEUROSCI.1016-11.2011

Roy-Byrne, P. P. (2005). The GABA-benzodiazepine receptor complex: structure, function, and role in anxiety. *J Clin Psychiatry*, *66*(Suppl 2), 14–20.

Sandhu, K. V., Yanagawa, Y., & Stork, O. (2015). Transcriptional Regulation of Glutamic Acid Decarboxylase in the Male Mouse Amygdala by Dietary Phyto-Oestrogens. *Journal of Neuroendocrinology*, *27*(4), 285–292. https://doi.org/10.1111/jne.12262

Seo, D., Ahluwalia, A., Potenza, M. N., & Sinha, R. (2017). Gender differences in neural correlates of stress-induced anxiety. *Journal of Neuroscience Research*, *95*(1-2), 115–125. https://doi.org/10.1002/jnr.23926

Stack, A., Carrier, N., Dietz, D., Hollis, F., Sorenson, J., & Kabbaj, M. (2010). Sex differences in social interaction in rats: role of the immediate-early gene zif268. *Neuropsychopharmacology*, *35*(2), 570–580. https://doi.org/10.1038/npp.2009.163

Steel, Z., Marnane, C., Iranpour, C., Chey, T., Jackson, J. W., Patel, V., & Silove, D. (2014). The global prevalence of common mental disorders: a systematic review and meta-analysis 1980–2013. *International Journal of Epidemiology*, *43*(2), 476–493. https://doi.org/10.1093/ije/dyu038

Steullet, P., Cabungcal, J.-H., Cuénod, M., & Do, K. Q. (2014). Fast oscillatory activity in the anterior cingulate cortex: dopaminergic modulation and effect of perineuronal net loss. *Frontiers in Cellular Neuroscience*, *8*, 244. https://doi.org/10.3389/fncel.2014.00244

Stevens, J. S., & Hamann, S. (2012). Sex differences in brain activation to emotional stimuli: a meta-analysis of neuroimaging studies. *Neuropsychologia*, *50*(7), 1578–1593. https://doi.org/10.1016/j.neuropsychologia.2012.03.011

Stork, O., Ji, F.-Y., Kaneko, K., Stork, S., Yoshinobu, Y., Moriya, T., . . . Obata, K. (2000). Postnatal development of a GABA deficit and disturbance of neural functions in mice lacking GAD65. *Brain Research*, *865*(1), 45–58. https://doi.org/10.1016/S0006-8993(00)02206-X

Straube, T., Schmidt, S., Weiss, T., Mentzel, H.-J., & Miltner, W. H. R. (2009). Dynamic activation of the anterior cingulate cortex during anticipatory anxiety. *NeuroImage*, *44*(3), 975–981. https://doi.org/10.1016/j.neuroimage.2008.10.022

Strekalova, T., Spanagel, R., Dolgov, O., & Bartsch, D. (2005). Stress-induced hyperlocomotion as a confounding factor in anxiety and depression models in mice. *Behavioural Pharmacology*, *16*(3), 171–180. https://doi.org/10.1097/00008877-200505000-00006

Sturman, O., Germain, P.-L., & Bohacek, J. (2018). Exploratory rearing: a context-and stress-sensitive behavior recorded in the open-field test. *Stress*, *21*(5), 443–452. https://doi.org/10.1080/10253890.2018.1438405

Thomas, M., Gaudeau-Bosma, C., Bouaziz, N., Andrianisaina, P. S.-K., Benadhira, R., Januel, D., & Moulier, V. (2019). Gender-related brain activations during an emotional task: an fMRI study. *Neuropsychobiology*, *78*(3), 128–135. https://doi.org/10.1159/000499977

Tovote, P., Fadok, J. P., & Lüthi, A. (2015). Neuronal circuits for fear and anxiety. *Nature Reviews Neuroscience*, *16*(6), 317–331. https://doi.org/10.1038/nrn3945

Tripathi, K., Demiray, Y. E., Kliche, S., Jing, L., Hazra, S., Hazra, J. D., . . . Stork, O. (2021). Reducing glutamic acid decarboxylase in the dorsal dentate gyrus attenuates juvenile stress induced emotional and cognitive deficits. *Neurobiology of Stress*, *15*, 100350. https://doi.org/10.1016/j.ynstr.2021.100350

Van der Heiden, C., Melchior, K., Muris, P., Bouwmeester, S., Bos, A. E. R., & van der Molen, H. T. (2010). A hierarchical model for the relationships between general and specific vulnerability factors and symptom levels of generalized anxiety disorder. *Journal of Anxiety Disorders*, *24*(2), 284–289. https://doi.org/10.1016/j.janxdis.2009.12.005

Van Heukelum, S., Mars, R. B., Guthrie, M., Buitelaar, J. K., Beckmann, C. F., Tiesinga, P. H. E., . . . Havenith, M. N. (2020). Where is cingulate cortex? A cross-species view. *Trends in Neurosciences*, *43*(5), 285–299. https://doi.org/10.1016/j.tins.2020.03.007

Vesga-López, O., Schneier, F. R., Wang, S., Heimberg, R. G., Liu, S.-M., Hasin, D. S., & Blanco, C. (2008). Gender differences in generalized anxiety disorder: results from the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC). *Journal of Clinical Psychiatry*, *69*(10), 1606.

Voloh, B., Valiante, T. A., Everling, S., & Womelsdorf, T. (2015). Theta–gamma coordination between anterior cingulate and prefrontal cortex indexes correct attention shifts. *Proceedings of the National Academy of Sciences*, *112*(27), 8457–8462. https://doi.org/10.1073/pnas.150043811

Walton, M. E., Croxson, P. L., Behrens, T. E. J., Kennerley, S. W., & Rushworth, M. F. S. (2007). Adaptive decision making and value in the anterior cingulate cortex. *NeuroImage*, *36 Suppl 2*, T142-54. https://doi.org/10.1016/j.neuroimage.2007.03.029

Winkelmann, A., Maggio, N., Eller, J., Caliskan, G., Semtner, M., Häussler, U., . . . Kowalczyk, S. (2014). Changes in neural network homeostasis trigger neuropsychiatric symptoms. *The Journal of Clinical Investigation*, *124*(2), 696–711. https://doi.org/10.1172/JCI71472