Coupling of Slack and Na$_V$1.6 sensitizes Slack to quinidine blockade and guides anti-seizure strategy development

Tian Yuan[1#], Yifan Wang[1#], Yuchen Jin[1#], Hui Yang[1#], Shuai Xu[1], Heng Zhang[4], Qian Chen[1], Na Li[1], Xinyue Ma[1], Huifang Song[1], Chao Peng[1], Ze Geng[1], Jie Dong[1], Guifang Duan[1], Qi Sun[1], Yang Yang[2], Fan Yang[3,4] and Zhuo Huang[1,5*]

AFFILATIONS

[1]State Key Laboratory of Natural and Biomimetic Drugs, Department of Molecular and Cellular Pharmacology, School of Pharmaceutical Sciences, Peking University Health Science Center, Beijing, 100191, China;

[2]Department of Medicinal Chemistry and Molecular Pharmacology, College of Pharmacy, Purdue University, West Lafayette, IN 47907, USA;

[3]Department of Biophysics, Kidney Disease Center of the First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, Zhejiang, 310058, China;

[4]NHC and CAMS Key Laboratory of Medical Neurobiology, MOE Frontier Science Center for Brain Research and Brain-Machine Integration, School of Brain Science and Brain Medicine, Zhejiang University, Hangzhou, Zhejiang, 310058, China;

[5]IDG/McGovern Institute for Brain Research, Peking University, Beijing, 100871, China.

# These authors contribute equally to this work

*Correspondence: huangz@hsc.pku.edu.cn (Z.H.)

KEYWORDS

Slack, Nav1.6, quinidine, KCNT1-related epilepsy

GRAPHICAL ABSTRACT

ABSTRACT

Quinidine has been used as an anticonvulsant to treat patients with KCNT1-related

epilepsy by targeting gain-of-function KCNT1 pathogenic mutant variants. However,

the detailed mechanism underlying quinidine's blockade against KCNT1 (Slack)

remains elusive. Here, we report a functional and physical coupling of the voltage-gated

sodium channel Nav1.6 and Slack. Nav1.6 binds to and highly sensitizes Slack to

quinidine blockade. Homozygous knockout of Nav1.6 reduces the sensitivity of native

sodium-activated potassium currents to quinidine blockade. Nav1.6-mediated

sensitization requires the involvement of Na$_V$1.6's N- and C-termini binding to Slack's

C-terminus, and is enhanced by transient sodium influx through Na$_V$1.6. Moreover,

disrupting the Slack-Na$_V$1.6 interaction by viral expression of Slack's C-terminus can

protect against Slack^{G269S}-induced seizures in mice. These insights about a Slack-

Na$_V$1.6 complex challenge the traditional view of "Slack as an isolated target" for anti-

epileptic drug discovery efforts, and can guide the development of innovative

therapeutic strategies for KCNT1-related epilepsy.

INTRODUCTION

The sodium-activated potassium (K$_{Na}$) channels Slack and Slick were first identified in

guinea pig cardiomyocytes, and were subsequently found to be encoded by two genes

of the *Slo2* family[1,2]. Slack channels encoded by the *Slo2.2* (*KCNT1*) gene are gated by

Na$^+$, while Slick channels encoded by the *Slo2.1* (*KCNT2*) gene are more sensitive to

Cl$^-$ than Na$^+$. Slack channels are expressed at high levels in the central nervous system

(CNS), especially the cortex and brainstem[3-5]. Activation of Slack channels by

intracellular sodium ions forms delayed outward currents in neurons, contributes to

slow afterhyperpolarization (AHP) following repeated action potentials, and modulates

the firing frequency of neurons[6,7].

Mutations in the *KCNT1* gene have been implicated in a wide spectrum of epileptic

disorders, including early-onset epilepsy (*e.g.*, epilepsy of infancy with migrating focal

seizures (EIMFS), non-EIMFS developmental and epileptic encephalopathies, and

autosomal dominant or sporadic sleep-related hypermotor epilepsy (ADSHE))[8-11]. Over

50 mutations related to seizure disorders have been identified, typically displaying a

gain-of-function (GOF) phenotype in heterologous expression systems[11,12]. The

prescribed antiarrhythmic drug, quinidine, has emerged as a precision therapy for

KCNT1-related epilepsy by blocking Slack mutant variants *in vitro* and conferring

decreased seizure frequency and improved psychomotor development in clinical

treatment[13-16]. However, clinical quinidine therapy has shown limited success and

contradictory therapeutic effects, probably due to poor blood–brain barrier penetration,

dose-limiting off-target effects, phenotype-genotype associations, and rational

therapeutic schedule[16-19].

Slack requires high intracellular free Na^+ concentrations ($[Na^+]_{in}$) for its activation in

neurons (K_d of ~66 mM)[20]. However, previous investigations have shown that the

$[Na^+]_{in}$ at resting states (~10 mM) is much lower than the $[Na^+]_{in}$ needed for effective

Slack activation (*e.g.*, K_d value)[1,20,21]. Therefore, native Slack channels need to localize

with Na^+ sources within a nanodomain to be activated and exert physiological function.

Slack channels are known to be functionally coupled with sodium-permeable ion

channels in neurons, such as voltage-gated sodium (Na_V) channels and AMPA

receptors [21,22]. A question arises as to whether these known Na^+ sources modulate

Slack's sensitivity to quinidine blockade.

Here, we found that $Na_V1.6$ sensitizes Slack to quinidine blockade. Slack and $Na_V1.6$

form a complex that functions in $Na_V1.6$-mediated transient sodium influx to sensitize

Slack to quinidine blockade in HEK293 cells and in primary cortical neurons. The

widespread expression of these channel proteins in cerebral cortex, hippocampus, and

cerebellum supports that the Nav1.6-Slack complex is essential for the function of a

wide range of electrically excitable neurons, and moreover, that this complex can be

viewed as a vulnerable target for drug development to treat KCNT1-related disorders.

RESULTS

Nav1.6 sensitizes Slack to quinidine blockade

Slack currents are activated by sodium entry through voltage-gated sodium (Nav)

channels and ionotropic glutamate receptors (*e.g.* AMPA receptors)[21-23]. To investigate

potential modulators of Slack's sensitivity to quinidine blockade, we initially focused

on the known Na^+ sources of Slack. Working in HEK293 cells, we co-expressed Slack

with AMPA receptor subunits (GluA1, GluA2, GluA3, or GluA4) or Nav channel α

subunits (Nav1.1, Nav1.2, Nav1.3, or Nav1.6), which are highly expressed in the central

nervous system[24-27]. The sensitivity of Slack to quinidine blockade was assessed based

on the detected inhibitory effects of quinidine on delayed outward potassium

currents[13,28]. Interestingly, all neuronal Nav channels significantly sensitized Slack to

30 μM quinidine blockade, whereas no effect was observed upon co-expression of

Slack with GluA1, GluA2, GluA3, or GluA4 (Fig. S1).

When co-expressing Slack with Nav1.6 in HEK293 cells, Nav1.6 sensitized Slack to

quinidine blockade by nearly 100-fold (IC_{50} = 85.13 μM for Slack expressed alone and

an $IC_{50} = 0.87$ μM for Slack upon co-expression with $Na_V1.6$) (Fig.1A,B,D,F). $Na_V1.6$

also exhibited >10-fold selectivity in sensitizing Slack to quinidine blockade against

$Na_V1.1$, $Na_V1.2$, and $Na_V1.3$ (Fig. 1C,F, and Supplementary Table 1). When we co-

expressed the cardiac sodium channel $Na_V1.5$ with Slack, we observed only ~3-fold

sensitization of Slack to quinidine blockade (Fig.1E,F). These results together indicate

that the apparent functional coupling between Slack and Na_V channels is Na_V-channel-

subtype-specific, with $Na_V1.6$ being particularly impactful in Slack's responsivity to

quinidine blockade.

Slack and Slick are both K_{Na} channels (with 74% sequence identity) and adopt similar

structures[29], and Slick is also blocked by quinidine[30]. We next assessed whether $Na_V1.6$

sensitizes Slick to quinidine blockade and observed that, similar to Slack, co-expression

of Slick and $Na_V1.6$ in HEK293 cells resulted in a sensitization of Slick to quinidine

blockade (7-fold) (Fig. S2A,B). These results support that $Na_V1.6$ regulates both Slack

and Slick and that $Na_V1.6$ can sensitize K_{Na} channels to quinidine blockade *in vitro*.

We also asked whether $Na_V1.6$ sensitizes native K_{Na} channels to quinidine blockade in

neurons. We performed whole-cell patch-clamp recordings in primary cortical neurons

from postnatal (P0-P1) homozygous $Na_V1.6$ knockout C3HeB/FeJ mice and the wild-

type littermate controls. As Lithium is a much weaker activator of K_{Na} channels than

sodium[29], K_{Na} currents (I_{KNa}) were isolated by replacing sodium ions with equivalent

lithium ions in the bath solution (Fig. 1G,H) as previously described[22]. The amplitudes

of I_{KNa} remained unaffected by the homozygous knockout of $Na_V1.6$ (Fig. S3A).

Notably, 3 µM quinidine significantly inhibited native I_{KNa} (44%) in wild-type neurons

(Fig. 1I,K) and this inhibitory effect was not limited to targeting the larger I_{KNa} (Fig.

S3B). Conversely, the same concentration of quinidine had no significant effect on I_{KNa}

in $Na_V1.6$-knockout ($Na_V1.6$-KO) neurons (Fig. 1J,K). These results support that

$Na_V1.6$ is required for the observed high sensitivity of native K_{Na} channels to quinidine

blockade.

Transient sodium influx through $Na_V1.6$ enhances $Na_V1.6$-mediated sensitization

of Slack to quinidine blockade

We next investigated the biomolecular mechanism underlying $Na_V1.6$-mediated

sensitization of Slack to quinidine blockade. Considering that Slack currents are

activated by sodium influx[22,29], we initially assessed the effects of $Na_V1.6$-mediated

sodium influx on sensitizing Slack to quinidine blockade. We used 100 nM tetrodotoxin

(TTX) to block $Na_V1.6$-mediated sodium influx (Fig. S4A,B)[31]. In HEK293 cells

expressing Slack alone, 100 nM TTX did not affect Slack currents; nor did it affect

Slack's sensitivity to quinidine blockade ($IC_{50} = 83.27$ µM) (Fig. 2A,B and Fig. S4C,D).

In contrast, upon co-expression of Slack and $Na_V1.6$ in HEK293 cells, bath-application

of 100nM TTX significantly reduced the effects of $Na_V1.6$ in sensitizing Slack to

quinidine blockade ($IC_{50} = 25.04$ µM) (Fig. 2A,B). These findings support that sodium

influx through $Na_V1.6$ contributes to $Na_V1.6$-mediated sensitization of Slack to

quinidine blockade.

It is known that Nav1.6-mediated sodium influx involves a transient inward flux that

reaches a peak before subsequently decaying to the baseline within a few milliseconds;

this is termed a transient sodium current (I_{NaT})[32]. A small fraction of Nav1.6 currents

are known to persist after the rapid decay of I_{NaT}, and these are termed persistent sodium

currents (I_{NaP})[33]. We isolated I_{NaT} and I_{NaP} to explore their potential contributions in

sensitizing Slack to quinidine blockade. We selectively inactivated I_{NaT} using a

depolarized prepulse of -40 mV (Fig. S5A) and selectively blocked I_{NaP} by bath-

application of 20 μM riluzole, which is a relatively specific I_{NaP} blocker that is known

to stabilize inactivated-state Nav channels and delay recovery from inactivation[34,35].

Our findings ultimately confirmed that the 20 μM riluzole selectively blocked I_{NaP}

compared to I_{NaT} in HEK293 cells co-expressing Slack and Nav1.6 (Fig. S5B) and that

20 μM riluzole had no effect on Slack currents when expressed alone (Fig. S5C,D).

Consistent with previous investigations[21,22], inactivating I_{NaT} reduced whole-cell Slack

currents by 20%, and blocking I_{NaP} reduced Slack currents by 40% at +100 mV (Fig.

2C,D,F,G), supporting that Nav1.6-mediated sodium influx activates Slack.

Interestingly, inactivating I_{NaT} resulted in a > 20-fold decrease in Slack's sensitization

to quinidine blockade (IC_{50} = 22.26 μM) (Fig. 2C,E). In contrast, blocking I_{NaP} had no

effect on Slack's sensitization to quinidine blockade (IC_{50} = 1.60 μM) (Fig. 2F,H).

These findings indicate that Nav1.6 sensitizes Slack to quinidine blockade via I_{NaT} but

not I_{NaP}.

Given that Slack current amplitudes are sensitive to sodium influx, and considering that

quinidine is a sodium channel blocker, we examined whether $Na_V1.6$ has higher

sensitivity to quinidine blockade than other Na_V channel subtypes, which could

plausibly explain the observed increased strength of sensitization. We used whole-cell

patch-clamping to assess the sensitivity of $Na_V1.1$, $Na_V1.2$, $Na_V1.3$, $Na_V1.5$, and $Na_V1.6$

to quinidine blockade. These sodium channels exhibited similar levels of quinidine

sensitivity (IC_{50} values in the range of 35.61-129.84 μM) (Fig. 2I,J and Supplementary

Table 2), all of which were at least 40-fold lower than the $Na_V1.6$-mediated

sensitization of Slack to quinidine blockade (Fig. 1F). Additionally, co-expressing

Slack with $Na_V1.1$, $Na_V1.2$, $Na_V1.3$, $Na_V1.5$, or $Na_V1.6$ in HEK293 cells did not change

the sensitivity of these Na_V channel subtypes to quinidine blockade (Fig. S6 and

Supplementary Table 2). Thus, differential quinidine affinity for specific Na_V channel

subtypes cannot explain the large observed $Na_V1.6$-mediated sensitization of Slack to

quinidine blockade. Moreover, it is clear that $Na_V1.6$-mediated sensitization of Slack to

quinidine blockade is directly mediated by I_{NaT}, rather than through some secondary

effects related to $Na_V1.6$'s higher sensitivity to quinidine blockade.

Slack physically interacts with $Na_V1.6$

We found that the specific voltage-gated sodium channel blocker TTX did not

completely abolish the effects of $Na_V1.6$ on sensitizing Slack to quinidine blockade

(Fig. 2B), so it appears that a sodium-influx-independent mechanism is involved in the

observed $Na_V1.6$-mediated sensitization of Slack to quinidine blockade. We therefore

investigated a potential physical interaction between Slack and Na$_V$1.6. We initially

assessed the cellular distribution of Na$_V$1.2, Na$_V$1.6, and Slack in the hippocampus and

the neocortex of mouse. Consistent with previous studies[36,37], Na$_V$1.2 and Na$_V$1.6 were

localized to the axonal initial segment (AIS) of neurons, evident as the co-localization

of Na$_V$ and AnkG, a sodium channel-associated protein known to accumulate at the AIS

(Fig. 3A). Slack channels were also localized to the AIS of these neurons (Fig. 3A),

indicating that Slack channels are located in close proximity to Na$_V$1.6 channels, and

supporting their possible interaction. Moreover, Na$_V$1.6 was co-immunoprecipitated

with Slack in homogenates from mouse cortical and hippocampal tissues and from

HEK293T cells co-transfected with Slack and Na$_V$1.6 (Fig. 3B,C), supporting that

Slack and Na$_V$1.6 form protein complexes in mouse brains.

To assess the interaction between Slack and Na$_V$1.6 inside living cells, we performed a

FRET assay in transfected HEK293T cells[38]. Briefly, we genetically fused mTFP1 and

mVenus to the C-terminal regions of Slack and Na$_V$1.6, respectively (Fig. 3D). Upon

imaging the emission spectra cells co-expressing Na$_V$1.6-mVenus and Slack-mTFP1

(measured at the plasma membrane region) (Fig. 3E), we detected positive FRET

signals, indicating a Slack-Na$_V$1.6 interaction (Fig. 3F,H). The plasma membrane

regions from HEK293T cells co-transfected with Na$_V$1.6 and Slack showed FRET

efficiency values much larger than a negative control (in which standalone mVenus and

mTFP1 proteins were co-expressed) (Fig. 3G,H), indicating that Slack channels reside

in close spatial proximity (less than 10 nm) to Na$_V$1.6 channels in living cells.

We next characterized the consequences of the Slack-Na$_V$1.6 interaction in HEK293

cells using whole-cell recordings. Slack increased the rate of recovery from fast

inactivation of Na$_V$1.6 (Fig. S7E), with no significant effects on the steady-state

activation, steady-state fast inactivation, or ramp currents (Fig. S7C,D,F and

Supplementary Table 3). Additionally, we found that Na$_V$1.6 had no significant effects

on the activation rate or the current-voltage (I-V) relationship of Slack currents (Fig.

S7A,B). These results indicate that the physical interaction between Slack and Na$_V$1.6

produces functional consequences. Taken together, these findings support functional

and physical coupling of Slack and Na$_V$1.6.

Na$_V$1.6's N- and C-termini bind to Slack's C-terminus and sensitize Slack to

quinidine blockade

To explore whether the physical interaction between Slack and Na$_V$1.6 is required for

sodium-influx-mediated sensitization of Slack to quinidine blockade, we performed

inside-out patch-clamp recordings on HEK293 cells transfected with Slack alone or co-

transfected with Slack and Na$_V$1.6. Note that in these experiments the intracellular

sodium concentration ($[Na^+]_{in}$) was raised to 140 mM (a concentration at which most

Slack channels can be activated[20]), seeking to mimic the increased intracellular sodium

concentration upon sodium influx. When expressing Slack alone, increasing the sodium

concentration did not sensitize Slack to quinidine blockade (IC$_{50}$ = 120.42 μM) (Fig.

4a). However, upon co-expression of Na$_V$1.6 and Slack, Na$_V$1.6 significantly sensitized

Slack to quinidine blockade (IC$_{50}$ = 2.91 μM) (Fig. 4A). These results support that

physical interaction between Slack and $Na_V1.6$ is a prerequisite for sodium-influx-

mediated sensitization of Slack to quinidine blockade.

To investigate which interacting domains mediate $Na_V1.6$'s sensitization of Slack to

quinidine blockade, we focused on $Na_V1.6$'s cytoplasmic fragments, including its N-

terminus, inter domain linkers, and C-terminus (Fig. 4B). Whole-cell recordings from

HEK293 cells co-expressing Slack with these $Na_V1.6$ fragments revealed that the N-

terminus and C-terminus of $Na_V1.6$ significantly enhanced the sensitivity of Slack to

quinidine blockade ($IC_{50} = 31.59$ μM for Slack upon co-expression with $Na_V1.6$'s N-

terminus and $IC_{50} = 43.70$ μM for Slack upon co-expression with $Na_V1.6$'s C-terminus);

note that the inter domain linkers had no effect (Fig. 4C).

Subsequent co-immunoprecipitation and glutathione S-transferase (GST) pull down

assays of HEK293T cell lysates experimentally confirmed that $Na_V1.6$'s N- and C-

termini each interact with Slack (Fig. 4D and Fig. S8). Additionally, whole-cell

recordings using an $[Na^+]_{in}$ of 5 mM again showed that $Na_V1.6$'s N-terminus and

$Na_V1.6$'s C-terminus sensitize Slack to quinidine blockade ($IC_{50} = 27.87$ μM) (Fig. 4E).

And inside-out recordings using an $[Na^+]_{in}$ of 140 mM showed that co-expression of

Slack, $Na_V1.6$'s N-terminus, and $Na_V1.6$'s C-terminus resulted in obvious sensitization

of Slack to quinidine blockade, with an IC_{50} of 2.57 μM (Fig. 4F)—thus fully

mimicking the aforementioned effects of full-length $Na_V1.6$ ($IC_{50} = 2.91$ μM) (Fig. 4A).

These findings support that the binding of $Na_V1.6$'s N- and C-termini to Slack is

required for $Na_V1.6$'s sensitization of Slack to quinidine blockade.

Recalling that Nav1.5 had the least pronounced effect in sensitizing Slack to quinidine

blockade among all examined Nav channels (IC_{50} = 29.46 μM) (Fig. 1F and

Supplementary Table 1), we initially compared the sodium currents of Nav channel

subtypes co-expressed with Slack. The results revealed no significant differences in

activation time constants (Fig. S9A). It's worth noting that Nav1.5 exhibited

significantly larger current amplitudes than Nav1.6 (Fig. S9B). Subsequently, we

constructed Nav1.5-1.6 chimeras to test the roles of Nav1.6's N- and C-termini in

sensitizing Slack to quinidine blockade. The replacement of both the N-terminus

[residues 1-131] and C-terminus [residues 1772-2016] of Nav1.5 with Nav1.6's N-

terminus [residues 1-132] and C-terminus [residues 1766-1980] (namely Nav1.5/6^{NC})

fully mimicked effects of Nav1.6 in sensitizing Slack to quinidine blockade (IC_{50} = 1.13

μM) (Fig. 4G-I). We also found that replacement of Nav1.5's N-terminus [residues 1-

131] with Nav1.6's N-terminus [residues 1-132] (namely Nav1.5/6^{N}) fully mimicked

the effects of Nav1.6 (IC_{50} = 1.18 μM) (Fig. 4G-I).

Notably, despite both Nav1.5 and Nav1.5/6^{N} exhibited much larger current amplitudes

compared to Nav1.6 (Fig. S9B), only Nav1.5/6^{N} replicated the effect of Nav1.6 in

sensitizing Slack to quinidine blockade (Fig. 4H-I). These results suggest that the

observed differences between Nav1.5 and Nav1.6 in sensitizing Slack are unlikely to

be attributed to Nav1.6's smaller sodium current amplitudes but involve Nav1.6's N-

terminus. Consistently, Nav1.5's C-terminus sensitized Slack to quinidine blockade

(IC_{50} = 37.59 μM), whereas Nav1.5's N-terminus had no effect on Slack sensitization

(Fig. 4C). These findings support that Na$_V$1.6's N-terminus is essential for sensitizing

Slack to quinidine blockade.

Having demonstrated that Na$_V$1.6 sensitizes Slack via Na$_V$1.6's cytoplasmic N- and C-

termini, we investigated which domains of Slack interact with Na$_V$1.6 and focused on

Slack's cytoplasmic fragments, including Slack's N-terminus and C-terminus (Fig.

5A). In HEK293 cells co-expressing Na$_V$ and Slack, Na$_V$-mediated sensitization of

Slack to quinidine blockade was significantly attenuated upon the additional expression

of Slack's C-terminus, but not of Slack's N-terminus (Fig. 5B,C), suggesting that

Slack's C-terminus can disrupt the Slack-Na$_V$1.6 interaction by competing with Slack

for binding to Na$_V$1.6. Consistently, Slack's C-terminus co-immunoprecipitated with

Na$_V$1.6's N- and C-termini in HEK293T cell lysates (Fig. 5D). Together, these results

support that Slack's C-terminus physically interacts with Na$_V$1.6 and that this

interaction is required for Na$_V$1.6's sensitization of Slack to quinidine blockade.

Na$_V$1.6 binds to and sensitizes epilepsy-related Slack mutant variants to quinidine

blockade

Over 50 mutations in KCNT1 (Slack) have been identified and related to seizure

disorders[11]. Having established that Na$_V$1.6 can sensitize wild-type Slack to quinidine

blockade, we next investigated whether Na$_V$1.6 also sensitizes epilepsy-related Slack

mutant variants to quinidine blockade. We chose 3 Slack pathogenic mutant variants

(K629N, R950Q, and K985N) initially detected in patients with KCNT1-related

epilepsy[15,39,40]. Considering that these 3 mutations are located in Slack's C-terminus,

and recalling that Slack's C-terminus interacts with $Na_V1.6$ (Fig. 5D), we first used co-

immunoprecipitation assays and successfully confirmed that each of these Slack mutant

variants interacts with $Na_V1.6$ in HEK293T cell lysates (Fig. 6A).

Subsequently, whole-cell recordings revealed that the examined Slack mutant variants

exhibited no discernible impact on the amplitudes of $Na_V1.6$ currents (Fig. S10A),

while pharmacologically blocking $Na_V1.6$ currents using bath-application of 100 nM

TTX significantly reduced the amplitudes of Slack mutant variant ($Slack^{R950Q}$) currents

(Fig. S10B), suggesting a Na^+-mediated functional coupling between Slack mutant

variant and $Na_V1.6$ currents. Notably, $Na_V1.6$ significantly sensitized all of the

examined Slack mutant variants to quinidine blockade, with IC_{50} values ranging from

0.26 to 2.41 μM (Fig. 6B-D and Supplementary Table 4). These results support that

$Na_V1.6$ interacts with examined Slack mutant variants and sensitizes them to quinidine

blockade. It is plausible that the Slack-$Na_V1.6$ interaction contributes to the

therapeutical role of quinidine in the treatment of KCNT1-related epilepsy.

Viral expression of Slack's C-terminus prevents $Slack^{G269S}$-induced seizures

Having established that blocking $Na_V1.6$-mediated sodium influx significantly reduced

Slack current amplitudes (Fig. 2D,G and Fig. S10B), we found that the heterozygous

knockout of $Na_V1.6$ significantly reduced the afterhyperpolarization amplitude in

murine hippocampal neurons (Fig. S11), together indicating that $Na_V1.6$ activates

native Slack through providing Na^+. We therefore assumed that disruption of the Slack-

$Na_V1.6$ interaction should reduce the amount of Na^+ in the close vicinity of epilepsy-

related Slack mutant variants and thereby counter the increased current amplitudes of

these Slack mutant variants. Pursuing this, we disrupted the Slack-Nav interaction by

overexpressing Slack's C-terminus (to compete with Slack) and measured whole-cell

current densities. In HEK293 cells co-expressing epilepsy-related Slack mutant variants

(G288S, R398Q)[41,42] and Nav1.5/6[NC], expression of Slack's C-terminus significantly

reduced whole-cell current densities of Slack[G269S] and Slack[R398Q] (Fig. 7A,B),

supporting that disrupting the Slack-Nav1.6 interaction can indeed reduce current

amplitudes of Slack mutant variants, which may protect against seizures induced by

Slack mutant variants.

We next induced an *in vivo* epilepsy model by introducing a Slack G269S variant into

C57BL/6N mice using adeno-associated virus (AAV) injection to mimic the human

Slack mutation G288S[41]. Specifically, we delivered stereotactic injections of AAV9

containing expression cassettes for Slack[G269S] (or GFP negative controls) into the

hippocampal CA1 region of 3-week-old C57BL/6N mice (Fig. 7C,D). At 3-5 week

intervals after AAV injection, we quantified the seizure susceptibility of mice upon the

induction of a classic kainic acid (KA) model of temporal lobe epilepsy[43,44]. In this

model, seizures with stage IV or higher (as defined by a modified Racine, Pinal, and

Rovner scale[45]) are induced in rodents by intraperitoneal administration of 28 mg/kg

KA.

We assessed a time course of KA-induced seizure stages at 10-min intervals and found

that viral expression of Slack[G269S] resulted in faster seizure progression in mice

compared to control GFP expression (Fig. 7E). We calculated the total seizure score

per mouse to assess seizure severity[46]. Slack[G269S]-expressing mice showed significantly

higher seizure severity than GFP-expressing mice (Fig. 7F). The percentage of mice

with stage VI~IX seizures also increased, from 9.1% in GFP-expressing control mice

to 58.3% in the Slack[G269S]-expressing mice (Fig. 7G). These results support that viral

expression of Slack[G269S] significantly increases seizure susceptibility in mice.

To evaluate the potential therapeutic effects of disrupting the Slack-Na$_V$1.6 interaction,

we delivered two AAV9s (one for Slack[G269S] and one for Slack's C-terminus [residues

326-1238]) into the CA1 region of mice (Fig. 7D). Viral expression of Slack's C-

terminus in Slack[G269S]-expressing mice significantly decreased seizure progression,

seizure severity, and the percentage of mice experiencing stage VI~IX seizures (Fig.

7E-G). These results support that viral expression of Slack's C-terminus can prevent

Slack[G269S]-induced seizures in mice, thus showcasing that using Slack's C-terminus to

disrupt the Slack-Na$_V$1.6 interaction is a promising therapeutic intervention to treat

KCNT1-related epilepsy.

DISCUSSION

We here found that Na$_V$1.6's N- and C- termini bind to Slack's C-terminus and sensitize

Slack to quinidine blockade via Na$_V$1.6-mediated transient sodium currents. These

results suggest that the pharmacological blocking effects of a channel blocker are not

exclusively mediated by the channel *per.se.*, but modulated by channel's interacting

proteins. Moreover, we show that viral expression of Slack's C-terminus can rescue the

increased seizure susceptibility and confer protection against Slack[G269S]-induced

seizures in mice.

At resting membrane potential, the intracellular sodium concentration ($[Na^+]_{in}$) in

neurons (equal to ~10 mM) is too low to effectively activate Slack (K_d of 66 mM)[1,20].

Slack is functionally coupled to sodium influx, which is known to be mediated by ion

channels and receptors, including Na_V, AMPARs, and NMDARs[21-23,47]. Such Na^+

sources can provide both the membrane depolarization and the Na^+ entry known to be

required for Slack activation, enabling Slack to contribute both to action potential

repolarization during neuronal high-frequency firing[7,48] and to regulating excitatory

postsynaptic potential (EPSP) at post-synaptic neurons[23]. Our results support that Slack

and $Na_V1.6$ form a channel complex, while also implying that $Na_V1.6$-mediated sodium

influx increases the Na^+ concentration in the close vicinity of Slack to activate Slack.

As a low threshold Na_V channel subtype, $Na_V1.6$ has been reported to dominate the

initiation and propagation of action potentials in axon initial segments (AIS) in

excitatory neurons[36,49-51]. The activation of Slack by $Na_V1.6$ at AIS has multiple

impacts, including ensuring the timing of the fast-activated component of Slack

currents, regulating the action potential amplitude, and apparently contributing to

intrinsic neuronal excitability[29]. The use of HEK cells and cultured primary cortical

neurons in this study may not fully capture the complexity of native Slack-$Na_V1.6$

interaction. Future studies employing more intact systems, such as *in vivo* models, could

provide a more comprehensive understanding of the physiological relevance of the

native Slack-Nav1.6 interaction.

An interesting question arises from our observation that Slack's sensitivity to quinidine

blockade is enhanced by I_{NaT} but not I_{NaP}: what can explain the distinct I_{NaT} and I_{NaP}

contributions? We speculate that with physical modulation by Nav1.6, I_{NaT} may elicit a

specific open conformation of Slack that brings its quinidine binding pocket into a high

affinity state, which could lead to a substantial increase in Slack's sensitivity to

quinidine blockade. The possibility of this hypothetical open conformation is supported

by previous reports of the presence of subconductance states detected in single-channel

recordings of *Xenopus* oocytes expressing Slack channels[52], which implies that there

are multiple open conformations of Slack. Although only one open conformation of

Slack has been observed in cryo-EM, the gap between maximum conformational open

probability (~1.0) and maximum functional open probability (~0.7) implies a subclass

within this class of open channels[53]. Additionally, it's worth noting that blocking of I_{NaP}

by application of riluzole has its limitation on selectivity. Riluzole has been reported to

bind to Slack channels with low affinity[54]. Therefore, the dual binding of riluzole to

both Slack and Nav1.6 may stabilize or inhibit the Slack-Nav1.6 interaction,

consequently influencing the sensitization of Slack to quinidine blockade.

As previously mentioned, gain-of-function Slack mutant variants have been linked to a

broad spectrum of epileptic disorders that are accompanied by intellectual disabilities

and both psychomotor and developmental defects[18,55]. Given that many patients are

refractory or non-responsive to conventional anticonvulsants[12,55,56], and considering the

limited success of quinidine in clinical treatment of KCNT1-related epilepsy, inhibitors

targeting Slack are needed urgently[18]. Several small-molecule inhibitors against Slack

have been reported, providing informative starting points for drug development efforts

with KCNT1-related epilepsy[28,57,58]. However, our discovery of the Slack-Nav1.6

complex challenges the traditional view that Slack acts as an isolated target in KCNT1-

related epilepsy[18]. Indeed, our study supports that co-expression of Slack and Nav1.6

in heterologous cell models should be performed when analyzing clinically relevant

Slack mutations and when screening anti-epileptic drugs for use in treating KCNT1-

related disorders.

Genotype-phenotype analysis has shown that ADSHE-related mutations are clustered

in the regulator of conductance of K^+ (RCK2) domain; while EIMFS-related mutations

do not show a particular pattern of distribution[11]. All functionally tested Slack mutant

variants show gain-of-function phenotypes, with increased Slack currents [9,11,18].

Further, the epilepsy-related Slack mutant variants confer their gain-of-function

phenotypes through two molecular mechanisms: increasing maximal channel open

probability (P_{max}) or increasing sodium sensitivity (K_d) of Slack[59]. Both P_{max} and K_d are

highly sensitive to $[Na^+]_{in}$; these are respectively analogous to the efficacy and potency

of $[Na^+]_{in}$ on Slack currents[59]. Notably, several Slack mutant variants show gain-of-

function phenotypes only at high $[Na^+]_{in}$ (*e.g.* 80 mM)[59]. These results indicate that the

gain-of-function phenotype of epilepsy-related Slack mutant variants is aggravated by

high $[Na^+]_{in}$. Our discovery of functional coupling between $Na_V1.6$ and Slack presents

a plausible basis for how $Na_V1.6$-mediated sodium influx can increase $[Na^+]_{in}$ and thus

apparently aggravate the gain-of-function phenotype of Slack mutant variants.

Therefore, it makes sense that disruption of the Slack-$Na_V1.6$ interaction by

overexpressing Slack's C-terminus reduces the current amplitudes of gain-of-function

Slack mutant variants (Fig. 7A,B). Our successful demonstration that viral expression

of Slack's C-terminus prevents epilepsy-related Slack^{G269S}-induced seizures in mice

(Fig. 7E-G) warrants further translational evaluation for developing therapeutic

interventions to treat KCNT1-related epilepsy.

METHODS

Animals

C57BL/6 mice were purchased from Charles River Laboratories. $Na_V1.6$ knockout

C3HeB/FeJ mice were generous gifts from Professor Yousheng Shu at Fudan

University. All animals were housed on a 12-hour light/dark cycle with *ad libitum*

access to food and water. All procedures related to animal care and treatment were

approved by the Peking University Institutional Animal Care and Use Committee and

met the guidelines of the National Institute of Health Guide for the Care and Use of

Laboratory Animals. Each effort was made to minimize animal suffering and the

number of animals used. The experiments were blind to viral treatment condition during

behavioral testing.

Antibodies and Reagents

Commercial antibodies used were: anti-AnkG (Santa Cruz), anti-Slack (NeuroMab), anti-Nav1.2 (Alomone), anti-Nav1.6 (Alomone), anti-HA (Abbkine), anti-Flag (Abbkine), anti-β-Actin (Biodragon), HRP Goat Anti-Mouse IgG LCS (Abbkine), HRP Mouse Anti-Rabbit IgG LCS (Abbkine), Alexa Fluor 488-AffinityPure Fab Fragment Donkey anti-rabbit IgG (Jackson), Alexa Fluor 594 Donkey anti-mouse IgG (Yeason). GPCR Extraction Reagent was from Pierce, NP40 lysis buffer was from Beyotime, protease inhibitor mixture cocktail was from Roche Applied Science, rabbit IgG and mouse IgG was from Santa Cruz, and Protein G Dynabeads were from Invitrogen. Tetrodotoxin was from Absin Bioscience. Quinidine was from Macklin, and riluzole was from Meilunbio. All other reagents were purchased from Sigma-Aldrich.

Molecular cloning

The ion channels used are: Slack-B (Ref Seq: NM_020822.3), Slick (Ref Seq: NM_198503.5), Nav1.1 (Ref Seq: NM_001165963.4), Nav1.2 (Ref Seq: NM_001040142.2), Nav1.3 (Ref Seq: NM_006922.4), Nav1.5 (Ref Seq: NM_198056.3), Nav1.6 (Ref Seq: NM_014191.4), GluA1 (Ref Seq: XM_032913972.1), GluA2 (Ref Seq: NM_017261.2), GluA3 (Ref Seq: NM_007325.5), and GluA4 (Ref Seq: NM_000829.4). Human Slack-B and Nav1.6 were subcloned into the modified pcDNA3.1(+) vector using Gibson assembly. All mutations and chimeras of ion channels were also constructed using Gibson assembly. For GST pull down assay, the segments of Slack and Nav1.6 were subcloned into

447   pCDNA3.1(+) and pGEX-4T-1 vector, respectively. For FRET experiments, mVenus-

448   tag was fused to the C-terminus of Na$_V$1.6 sequence, and mTFP1-tag was also fused to

449   the C-terminus of Slack sequence.

Immunoprecipitation

The brain tissues or HEK293T cells co-expressing full-length or fragments of Slack

and Na$_V$1.6 were homogenized and lysed in GPCR Extraction Reagent (Pierce) with

cocktail for 30 min at 4 °C. The homogenate was centrifuged for 20 min at 16 000 g

and 4 °C to remove cell debris and then supernatant was incubated with 5 μg Slack

antibody (NeuroMab) or Na$_V$1.6 antibody (Alomone) for 12 h at 4 °C with constant

rotation. 40 μl of protein G Dynabeads (Invitrogen) was then added and the incubation

was continued until the next day. Beads were then washed three times with NP40 lysis.

Between washes, the beads were collected by DynaMag. The remaining proteins were

eluted from the beads by re-suspending the beads in 1×SDS-PAGE loading buffer and

incubating for 30 min at 37 °C. The resultant materials from immunoprecipitation or

lysates were then subjected to western blot analysis. And the co-immunoprecipitation

assays for Slack and Na$_V$1.6 were repeated three times independently.

Western blot analysis

Proteins suspended in 1×SDS-PAGE loading buffer were denatured for 30 min at 37 °C.

Then proteins were loaded on 6% or 8% sodium dodecyl sulphate–polyacrylamide gel

electrophoresis and transferred onto nitrocellulose filter membrane (PALL). Non-

specific binding sites were blocked with Tris-buffered saline-Tween (0.02 M Tris,

0.137 M NaCl and 0.1% Tween 20) containing 5% non-fat dried milk. Subsequently,

proteins of interest were probed with primary antibodies for overnight at 4 °C. After

incubation with a secondary antibody, immunoreactive bands were visualized using

HRP Substrate Peroxide Solution (Millipore) according to the manufacturer's

recommendation.

GST pull down assay

Plasmids encoding GST-fused Nav1.6 segments were transformed into BL21(DE3).

After expressing recombinant proteins induced by overnight application of isopropyl-

1-thio-β-D-galactopyranoside (IPTG) (0.1mM) at 25°C, the bacteria were collected,

lysed and incubated with GSH beads using BeaverBeads GSH kit (Beaver) according

to the manufacturer's instruction.

Plasmids encoding KCNT1 channel or HA-tagged KCNT1 segments were transfected

into HEK293T cells using lipofectamine 2000 (Introvigen). 40 h after transfection, cells

were lysed in NP40 lysis buffer with inhibitor cocktail, and then centrifugated at 4 °C,

15000 g for 20 min. The supernatants were incubated with protein-bound beads. The

protein-bound beads were washed by washing buffer (50 mM Tris-HCl pH = 7.4, 120

mM NaCl, 2 mM EGTA, 2 mM DTT, 0.1% triton X-100, and 0.2% Tween-20) for 5

min 3 times and then denatured with 1×SDS-PAGE loading buffer and incubating for

30 min at 37 °C. The resultant materials were subjected to western blot analysis.

Cell culture

The human embryonic kidney cells (HEK293 and HEK293T) were maintained in Dulbecco's Modified Eagle Medium (DMEM, Gibco) supplemented with 15% Fetal Bovine Serum (FBS, PAN-Biotech) at 37ºC and 5% CO_2. Primary cortical neurons were prepared from either sex of postnatal (P0-P1) homozygous $Na_V1.6$ knockout C3HeB/FeJ mice and the wild-type littermate controls. After the mice were decapitated, the cortices were removed and separated from the meninges and surrounding tissue. Tissues were digested in 2 mg/mL Papain (Aladdin) containing 2 µg/mL DNase I (Psaitong) for 30 min followed by centrifugation and resuspension. Subsequently, the cells were plated on poly-D-lysine (0.05 mg/mL) pre-coated glass coverslips in plating medium (DMEM containing 15% FBS), at a density of 4×10^5 cells/ml, and cultivated at 37°C and 5% CO_2 in a humidified incubator. 5 h after plating, the medium was replaced with Neurobasal Plus medium (Invitrogen) containing 2% v/v B-27 supplement (Invitrogen), 2 mM Glutamax (Invitrogen), 50 U/mL penicillin and streptomycin (Life Technologies). The primary neurons were grown 6-10 days before electrophysiological recordings with half of the media replaced every three days.

Voltage-clamp recordings

The plasmids expressing full-length or fragments of Slack and Na_V channels (excluding full-length $Na_V1.6$) were co-transfected into HEK293 cells using lipofectamine 2000 (Introvigen). To co-express Slack with $Na_V1.6$, the plasmid expressing Slack was transfected into a stable HEK293 cell line expressing $Na_V1.6$. 18-36 h after transfection,

voltage-clamp recordings were obtained using a HEKA EPC-10 patch-clamp amplifier

(HEKA Electronic) and PatchMaster software (HEKA Electronic). For all whole-cell

patch clamp experiments in HEK cells except for data presented in Fig. S7D, the

extracellular recording solution contained (in mM): 140 NaCl, 3 KCl, 1 $CaCl_2$, 1 MgCl2,

10 glucose, 10 HEPES, 1 Tetraethylammonium chloride (310 mOsm/L, pH 7.30 with

NaOH). The recording pipette intracellular solution (5 mM Na) contained (in mM): 100

K-gluconate, 30 KCl, 15 Choline-Cl, 5 NaCl, 10 glucose,5 EGTA, 10 HEPES (300

mOsm/L, pH 7.30 with KOH). For data presented in Fig. S7D, the extracellular

recording solution remains the same as above, the pipette intracellular solution

contained (in mM): 140 CsF, 10 NaCl, 5 EGTA, 10 HEPES (pH 7.30 with NaOH),

indicating that the unusual right shifts (15~20 mV) in voltage dependence of $Na_V1.6$

were induced by components in pipette solution, not recording system errors. For

primary cortical neurons, intracellular solution (0 mM Na) was used to prevent the

activation of sodium-activated potassium channels by basal intracellular sodium ions.

NaCl in the intracellular solution was replaced with choline chloride in an equimolar

concentration. For inside out patch-clamps, the bath solution contained (in mM): 140

NaCl, 1 EDTA, 10 HEPES and 2 $MgCl_2$ (310 mOsm/L, pH 7.30 with NaOH). Pipette

solution contained (in mM): 130 KCl, 1 EDTA, 10 HEPES and 2 $MgCl_2$ (300 mOsm/L,

pH 7.30 with KOH). The pipettes were fabricated by a DMZ Universal Electrode puller

(Zeitz Instruments) using borosilicate glass, with a resistance of 1.5-3.5 MΩ for whole-

cell patch clamp recordings and 8.0-10.0 MΩ for inside-out patch clamp recordings.

All experiments were performed at room temperature. The cells were exposed to the

bath solution with quinidine for about one minute before applying voltage protocols.

The concentration-response curves were fitted to four-parameter Hill equation:

$Y = Bottom + (Top - Bottom)/(1 + 10^{(lgIC_{50}-X)*H})$ (1)

where Y is the value of $I_{Quinidine}/I_{Control}$, Top is the maximum response, Bottom is the

minimum response, X is the lg of concentration, IC_{50} is the drug concentration

producing the half-maximum response, and H is the Hill coefficient. Significance of

fitted IC_{50} values compared to control was analyzed using extra sum-of-squares F test.

For data presented in Fig. S7, cells were excluded from analysis if series resistance > 5

MΩ and series resistance compensation was set to 70%~90%.

The time constants (τ) of activation were fitted with a single exponential equation:

$I(t) = Offset + A * exp(-\frac{t}{\tau})$ (2)

where I is the current amplitude, t is time, offset represents the asymptote of the fit, and

A represents the amplitude for the activation or inactivation.

Steady-state fast inactivation (I-V) and conductance-voltage (G-V) relationships were

fitting with Boltzmann equations:

$I/I_{max} = 1/(1 + \exp(\frac{V_m - V_{1/2}}{k}))$ (3)

$G/G_{max} = 1/(1 + \exp(\frac{V_m - V_{1/2}}{-k}))$ (4)

$G = I/(V_m - E_{Na})$ (5)

where I is the peak current, G is conductance, V_m is the stimulus potential, $V_{1/2}$ is the

midpoint voltage, E_{Na} is the equilibrium potential, and k is the slope factor. Significance

of fitted $V_{1/2}$ compared to control was analyzed using extra sum-of-squares F test.

Recovery from fast inactivation data were fitted with a single exponential equation:

$$I/I_{max} = A * (1 - exp\left(-\frac{t}{\tau}\right)) \qquad (6)$$

Where I is the peak current of test pulse, I_{max} is the peak current of first pulse, A is the

proportional coefficient, t is the delay time between the two pulses, and τ is the time

constant of recovery from fast inactivation.

The activation time constants of NaV channels were fitted with a single exponential

equation:

$$I = B + A * exp\left(\frac{t}{\tau}\right) \qquad (7)$$

Where I is the current amplitude, A and B are the partition coefficients, t is the time,

and τ is the time constant of activation.

Acute slice preparation and current-clamp recordings

Horizontal slices containing hippocampus were obtained from 6-8 weeks old

heterozygous Nav1.6 knockout ($Scn8a^{+/-}$) C3HeB/FeJ mice and the wild-type littermate

controls. In brief, animals were anesthetized and perfused intracardially with ice-cold

modified "cutting solution" containing (in mM): 110 choline chloride, 2.5 KCl, 0.5

$CaCl_2$, 7 $MgCl_2$, 25 $NaHCO_3$, 1.25 NaH_2PO_4, 10 glucose; bubbled continuously with

95%O_2/5%CO_2 to maintain PH at 7.2. The brain was then removed and submerged in

ice-cold "cutting solution". Next, the brain was cut into 300 μm slices with a vibratome

(WPI). Slices were incubated in oxygenated (95% O_2 and 5% CO_2) "recording solution"

containing (in mM): 125 NaCl, 2.5 KCl, 2 $CaCl_2$, 2 $MgCl_2$, 25 $NaHCO_3$, 1.25 NaH_2PO_4,

10 glucose (315 mOsm/L, PH 7.4, 37°C) for 30 min, and stored at room temperature.

Slices were subsequently transferred to a submerged chamber containing "recording

solution" maintained at 34-36 °C. Whole-cell recordings were obtained from

hippocampal CA1 neurons under a ×60 water-immersion objective of an Olympus

BX51WI microscope (Olympus). Pipettes had resistances of 5-8 MΩ. For current-

clamp recordings, the external solution (unless otherwise noted) was supplemented

with 0.05 mM (2R)-amino-5-phosphonovaleric acid (APV), 0.01 mM 6-cyano-7-nitro-

quinoxaline-2,3-dione, 0.01 mM bicuculline and 0.001 mM CGP 55845, and internal

pipette solution containing (in mM): 118 $KMeSO_4$, 15 KCl, 10 HEPES, 2 $MgCl_2$, 0.2

EGTA, and 4 Na_2ATP, 0.3 Tris-GTP, 14 Tris-phosphocreatinin (pH 7.3 with KOH).

In our whole-cell current-clamp recordings with an Axon 700B amplifier (Molecular

Devices), we initially applied a 100 ms, 20 pA test pulse to the recording neurons right

after breaking into the whole-cell configuration. A fast-rising component and a slow-

rising component of voltage response were clearly visible. Then we zoomed into the

fast-rising component of the voltage responses and turned up the pipette capacitance

neutralization slowly to shorten the rise time of the fast-rising component until the

oscillations of voltage responses appeared. Subsequently, we decreased the capacitance

compensation just until the oscillations disappeared. For bridge balance of current-

clamp recordings, we increased the value of bridge balance slowly until the fast

component of the voltage response disappeared, and the slow-rising component

appeared to rise directly from the baseline. Series resistance and pipette capacitance

were compensated using the bridge balance and pipette capacitance neutralization

options in the Multiclamp 700B command software (Molecular Devices). The bridge

balance value was between 20 and 30 MΩ and the pipette capacitance neutralization

value was between 3 and 5 pF. For in vitro experiments, the cells were selected by

criteria based on hippocampal CA1 cell morphology and electrophysiological

properties in the slices. Electrophysiological recordings were made using a Multiclamp

700B amplifier (Molecular Devices). Recordings were filtered at 10 kHz and sampled

at 50 kHz. Data were acquired and analyzed using pClamp10.0 (Molecular Devices).

Series resistance was in the order of 10–30 MΩ and was approximately 60–80%

compensated. Recordings were discarded if the series resistance increased by more than

20% during the time course of the recordings.

Fluorescence Imaging and FRET Quantification

The spectroscopic imaging was built upon a Nikon TE2000-U microscope. The

excitation light was generated by an Ar laser. The fluorescent protein mVenus fused to

Na$_V$1.6 and mTFP1 fused to Slack were excited by laser line at 500 and 400-440 nm,

respectively. The duration of light exposure was controlled by a computer-driven

mechanical shutter (Uniblitz). A spectrograph (Acton SpectraPro 2150i) was used in

conjunction with a charge coupled device (CCD) camera (Roper Cascade 128B). In this

recording mode two filter cubes (Chroma) were used to collect spectroscopic images

from each cell (excitation, dichroic): cube I, D436/20, 455dclp; cube II, HQ500/20,

Q515lp. No emission filter was used in these cubes. Under the experimental conditions,

auto fluorescence from untransfected cells was negligible. Fluorescence imaging and

analysis were done using the MetaMorph software (Universal Imaging). User-designed

macros were used for automatic collection of the bright field cell image, the

fluorescence cell image, and the spectroscopic image. Emission spectra were collected

from the plasma membrane of the cell by positioning the spectrograph slit across a cell

and recording the fluorescence intensity at the position corresponding to the membrane

region (Fig. 2e, dotted lines in red); the same slit position applied to both the spectrum

taken with the mTFP1 excitation and the spectrum taken with the mVenus excitation.

Using this approach, the spectral and positional information are well preserved, thus

allowing reliable quantification of FRET efficiency specifically from the cell

membrane. Spectra were corrected for background light, which was estimated from the

blank region of the same image.

FRET data was quantified in two ways. First, the FRET ratio was calculated from the

increase in mVenus emission due to energy transfer as described in the previous study.[60]

Briefly, mTFP1 emission was separated from mVenus emission by fitting of standard

spectra acquired from cells expressiring only mVenus or mTFP1. The fraction of

mVenus-tagged molecules that are associated with mTFP1-tagged molecules, Ab, is

calculated as

$Ab = 1/(1 + K_D/[\text{Dfree}])$ (7)

where K_D is the dissociation constant and [Dfree] is the concentration of free donor

molecules. Note that

$FRET\ Ratio = 1 + Ab * (FRET\ Ratio_{max} - 1)$ (8)

Regression analysis was used to estimate Ab in individual cells. From each cell, the

FRET ratio$_{exp}$ was experimentally determined. The predicted Ab value was then

computed by adjusting two parameters, FRET Ratio$_{max}$ and apparent K_D. Ab was in turn

used to give a predicted FRET ratiopredicted. By minimizing the squared errors (FRET

ratio$_{exp}$ – FRET ratio$_{predicted}$)^2, K_D was determined.

Second, apparent FRET efficiency was also calculated from the enhancement of

mVenus fluorescence emission due to energy transfer[60-63] using a method as previously

described.[64] Briefly, Ratio A$_0$ and Ratio A were measured to calculate FRET efficiency.

Ratio A$_0$ represents the ratio between tetramethylrhodamine maleimide emission

intensities (in the absence of fluorescein maleimide) upon excitation at the donor and

acceptor excitation wavelengths[63-65], and was calculated in the present study at the

mVenus peak emission wavelength. A particular advantage of quantifying Ratio A$_0$ for

FRET measurement is that changes in fluorescence intensity caused by many

experimental factors can be cancelled out by the ratiometric measurement. A similar

ratio, termed Ratio A, was determined in the presence of mTFP1 in the same way as

Ratio A$_0$. If FRET occurred, the Ratio A value should be higher than Ratio A$_0$; the

difference between Ratio A and Ratio A_0 was directly proportional to the FRET

efficiency by the factor of extinction coefficient ratio of mTFP1 and mVenus.[63-65]

Immunostaining

After deep anesthesia with sodium pentobarbital, mice were sacrificed by perfusion

with 0.5% paraformaldehyde and 0.5% sucrose (wt/vol) in 0.1 M phosphate buffer (pH

7.4). The brain was removed and post-fixed in the same fixative for 2 h, and

subsequently immersed in 30% sucrose in 0.1 M phosphate buffer for 48 h. Cryostat

coronal sections (20 µm) were obtained using a freezing microtome (Leica). The

sections were rinsed in 0.01 M phosphate-buffered saline (PBS, pH 7.4), permeabilized

in 0.5% Triton X-100 in PBS for 30 min, and incubated in a blocking solution (5% BSA,

0.1% Triton X-100 in PBS, vol/vol) at 20–25 °C for 2 h, followed by overnight

incubation at 4 °C with primary antibody to AnkG (1:100, Santa Cruz, sc-31778), Slack

(1:100, NeuroMab, 73-051), $Na_V1.2$ (1:200, Alomone, ASC-002), $Na_V1.6$ (1:200,

Alomone, ASC-006), Flag (1:500, Abbkine, ABT2010), and HA (1:500, Abbkine,

ABT2040) in blocking solution. After a complete wash in PBS, the sections were

incubated in Alexa 488-conjugated doney anti-rabbit IgG and Alexa 594 donkey anti-

mouse IgG in blocking solution at 20–25 °C for 2 h. The sections were subsequently

washed and rinsed in Dapi solution. Images were taken in the linear range of the

photomultiplier with a laser scanning confocal microscope (ZEISS LSM 510 META

NLO).

Adeno-associated virus construction and injection

The adeno-associated viruses (AAVs) and the negative GFP control were from Shanghai GeneChem. Co., Ltd. The full-length Slack$_{G269S}$ sequence (1-1238aa) was ligated into modified CV232 (CAG-MCS-HA-Poly A) adeno-associated viral vector. The Slack's C-terminus sequence (326-1238aa) and the negative control were ligated into GV634 (CAG-MCS-3×Flag-T2A-EGFP-SV40-Poly A) adeno-associated viral vector. The viruses (>10[11] TU/ml) were used in the present study.

For dorsal CA1 viral injection, C57BL/6J mice aged 3 weeks were anesthetized with isoflurane and placed in a stereotaxic apparatus (RWD Life Science Co., Ltd.). Using a 5 μL micro syringe (Hamilton) with a 30 gauge needle (RWD Life Science Co., Ltd.), 600 nL of the viruses was delivered at 10 nL/min by a micro-syringe pump (RWD Life Science Co., Ltd.) at the following site in each of the bilateral CA1 regions, using the stereotaxic coordinates: 2.5 mm (anterior-posterior) from bregma, 2 mm (medio-lateral), ± 1.5 mm (dorsal-ventral)[66]. The syringe was left in place for 5 min after each injection and withdrawn slowly. The exposed skin was closed by surgical sutures and returned to home cage for recovery. All the experiments were conducted after at least 3 weeks of recovery. All the mice were sacrificed after experiments to confirm the injection sites and the viral trans-infection effects by checking EGFP under a fluorescence microscope (ZEISS LSM 510 META NLO).

Kainic acid-induced status epilepticus

KA (Sigma-Aldrich) was intraperitoneally administered to produce seizures with stage IV or higher. The dose of kainic acid used was 28 mg/kg for mice (6–8 weeks)[67,68]. To

assess epilepsy susceptibility, seizures were rated using a modified Racine, Pinal, and

Rovner scale[45,69]: (1) Facial movements; (2) head nodding; (3) forelimb clonus; (4)

dorsal extension (rearing); (5) Loss of balance and falling; (6) Repeated rearing and

failing; (7) Violent jumping and running; (8) Stage 7 with periods of tonus; (9) Dead.

Seizures was terminated 2 h after onset with the use of sodium pentobarbital (30 mg/kg;

Sigma-Aldrich).

Statistical analysis

For in vitro experiments, the cells were evenly suspended and then randomly distributed

in each well tested. For in vivo experiments, the animals were distributed into various

treatment groups randomly. Statistical analyses were performed using GraphPad Prism

9 (GraphPad Software) and SPSS 26.0 software (SPSS Inc.). Before statistical analysis,

variation within each group of data and the assumptions of the tests were checked.

Comparisons between two independent groups were made using unpaired Student's

two-tailed t test. Comparisons among nonlinear fitted values were made using extra

sum-of-squares F test. Comparisons among three or more groups were made using one-

or two-way analysis of variance followed by Bonferroni's post hoc test. No statistical

methods were used to predetermine sample sizes but our sample sizes are similar to

those reported previously in the field[37,70]. All experiments and analysis of data were

performed in a blinded manner by investigators who were unaware of the genotype or

manipulation. $* p < 0.05$, $** p < 0.01$, $*** p < 0.001$, $**** p < 0.0001$. All data are

presented as mean $\pm$ SEM.

AUTHOR CONTRIBUTIONS

T.Y., Y.J., and H.Y. performed and analyzed voltage-clamp recordings. Y.W. performed and analyzed Western blotting, immunoprecipitation. S.X., C.P., and G.D. performed and analyzed pull down assay. Q.C. performed the immunostaining. H.Z. and F.Y. performed and analyzed the FRET imaging. H.S., N.L., and X.M. performed and analyzed the current clamp recordings. T.Y., H.Y., Z.G., and J.D. performed the molecular cloning. Z.H., T.Y., and Y.W. designed the experiments. Z.H., T.Y., and Y.J. wrote the manuscript. Z.H., F.Y., Y.Y., and Q.S. reviewed the manuscript.

ACKNOWLEDGMENTS

This work was supported by Chinese National Programs for Brain Science and Brain-like intelligence technology No.2021ZD0202102 to Z.H.; National Natural Science Foundation of China Grant (Nos. 31871083 and 81371432 to Z.H.; Nos. 32000674 to G.D.). We also thank Professor Yousheng Shu at Fudan University for providing the Nav1.6-knockout mice.

DATA AVAILABILITY

The data that support the findings of this study are available from the corresponding author upon reasonable request.

REFERENCES

1. Kameyama M, Kakei M, Sato R, Shibasaki T, Matsuda H, Irisawa H. Intracellular

Na+ activates a K+ channel in mammalian cardiac cells. *Nature*. May 24-30 1984;309(5966):354-6. doi:10.1038/309354a0

2. Yuan A, Santi CM, Wei A, et al. The sodium-activated potassium channel is encoded by a member of the Slo gene family. *Neuron*. 2003;37(5):765-773.

3. Joiner WJ, Tang MD, Wang LY, et al. Formation of intermediate-conductance calcium-activated potassium channels by interaction of Slack and Slo subunits. *Nat Neurosci*. Oct 1998;1(6):462-9. doi:10.1038/2176

4. Rizzi S, Knaus HG, Schwarzer C. Differential distribution of the sodium-activated potassium channels slick and slack in mouse brain. *J Comp Neurol*. Jul 1 2016;524(10):2093-116. doi:10.1002/cne.23934

5. Bhattacharjee A, Gan L, Kaczmarek LK. Localization of the Slack potassium channel in the rat central nervous system. *J Comp Neurol*. Dec 16 2002;454(3):241-54. doi:10.1002/cne.10439

6. Yang B, Desai R, Kaczmarek LK. Slack and Slick K(Na) channels regulate the accuracy of timing of auditory neurons. *J Neurosci*. Mar 7 2007;27(10):2617-27. doi:10.1523/JNEUROSCI.5308-06.2007

7. Wallen P, Robertson B, Cangiano L, et al. Sodium-dependent potassium channels of a Slack-like subtype contribute to the slow afterhyperpolarization in lamprey spinal neurons. *J Physiol*. Nov 15 2007;585(Pt 1):75-90. doi:10.1113/jphysiol.2007.138156

8. Heron SE, Smith KR, Bahlo M, et al. Missense mutations in the sodium-gated potassium channel gene KCNT1 cause severe autosomal dominant nocturnal frontal

lobe epilepsy. *Nat Genet*. Nov 2012;44(11):1188-90. doi:10.1038/ng.2440

9.   Barcia G, Fleming MR, Deligniere A, et al. De novo gain-of-function KCNT1

channel mutations cause malignant migrating partial seizures of infancy. *Nature*

genetics. 2012;44(11):1255-1259.

10.   Kingwell K. Mutations in potassium channel KCNT1—a novel driver of epilepsy

pathogenesis. *Nature Reviews Neurology*. 2012;8(12):658-658.

11.   Bonardi CM, Heyne HO, Fiannacca M, et al. KCNT1-related epilepsies and

epileptic encephalopathies: phenotypic and mutational spectrum. *Brain*. Dec 31

2021;144(12):3635-3650. doi:10.1093/brain/awab219

12.   McTague A, Nair U, Malhotra S, et al. Clinical and molecular characterization of

KCNT1-related severe early-onset epilepsy. *Neurology*. Jan 2 2018;90(1):e55-e66.

doi:10.1212/WNL.0000000000004762

13.   Milligan CJ, Li M, Gazina EV, et al. KCNT1 gain of function in 2 epilepsy

phenotypes is reversed by quinidine. *Annals of neurology*. 2014;75(4):581-590.

14.   Bearden D, Strong A, Ehnot J, DiGiovine M, Dlugos D, Goldberg EM. Targeted

treatment of migrating partial seizures of infancy with quinidine. *Ann Neurol*. Sep

2014;76(3):457-61. doi:10.1002/ana.24229

15.   Mikati MA, Jiang YH, Carboni M, et al. Quinidine in the treatment of KCNT1-

positive epilepsies. *Ann Neurol*. Dec 2015;78(6):995-9. doi:10.1002/ana.24520

16.   Numis AL, Nair U, Datta AN, et al. Lack of response to quinidine in KCNT 1‐

related neonatal epilepsy. *Epilepsia*. 2018;59(10):1889-1898.

17. Liu R, Sun L, Wang Y, Wang Q, Wu J. New use for an old drug: quinidine in

KCNT1-related epilepsy therapy. *Neurol Sci*. Nov 28 2022;doi:10.1007/s10072-022-

06521-x

18. Cole BA, Clapcote SJ, Muench SP, Lippiat JD. Targeting KNa1.1 channels in

KCNT1-associated epilepsy. *Trends Pharmacol Sci*. Aug 2021;42(8):700-713.

doi:10.1016/j.tips.2021.05.003

19. Xu D, Chen S, Yang J, Wang X, Fang Z, Li M. Precision therapy with quinidine of

KCNT1-related epileptic disorders: A systematic review. *Br J Clin Pharmacol*. Dec

2022;88(12):5096-5112. doi:10.1111/bcp.15479

20. Zhang Z, Rosenhouse-Dantsker A, Tang QY, Noskov S, Logothetis DE. The RCK2

domain uses a coordination site present in Kir channels to confer sodium sensitivity to

Slo2.2 channels. *J Neurosci*. Jun 2 2010;30(22):7554-62.

doi:10.1523/JNEUROSCI.0525-10.2010

21. Budelli G, Hage TA, Wei A, et al. Na+-activated K+ channels express a large

delayed outward current in neurons during normal physiology. *Nat Neurosci*. Jun

2009;12(6):745-50. doi:10.1038/nn.2313

22. Hage TA, Salkoff L. Sodium-activated potassium channels are functionally

coupled to persistent sodium currents. *J Neurosci*. Feb 22 2012;32(8):2714-21.

doi:10.1523/JNEUROSCI.5088-11.2012

23. Nanou E, Kyriakatos A, Bhattacharjee A, Kaczmarek LK, Paratcha G, El Manira

A. Na+-mediated coupling between AMPA receptors and KNa channels shapes synaptic

transmission. *Proceedings of the National Academy of Sciences*. 2008;105(52):20941-

20946.

24. Goldin AL. Resurgence of sodium channel research. *Annu Rev Physiol*.

2001;63:871-94. doi:10.1146/annurev.physiol.63.1.871

25. Trimmer JS, Rhodes KJ. Localization of voltage-gated ion channels in mammalian

brain.   *Annu*   *Rev*   *Physiol*.   2004;66:477-519.

doi:10.1146/annurev.physiol.66.032102.113328

26. Lai HC, Jan LY. The distribution and targeting of neuronal voltage-gated ion

channels. *Nat Rev Neurosci*. Jul 2006;7(7):548-62. doi:10.1038/nrn1938

27. Pickard L, Noel J, Henley JM, Collingridge GL, Molnar E. Developmental changes

in synaptic AMPA and NMDA receptor distribution and AMPA receptor subunit

composition in living hippocampal neurons. *J Neurosci*. Nov 1 2000;20(21):7922-31.

28. Cole BA, Johnson RM, Dejakaisaya H, et al. Structure-Based Identification and

Characterization of Inhibitors of the Epilepsy-Associated KNa1.1 (KCNT1) Potassium

Channel. *iScience*. May 22 2020;23(5):101100. doi:10.1016/j.isci.2020.101100

29. Kaczmarek LK. Slack, Slick and Sodium-Activated Potassium Channels. *ISRN*

Neurosci. Apr 18 2013;2013(2013)doi:10.1155/2013/354262

30. Bhattacharjee A, Joiner WJ, Wu M, Yang Y, Sigworth FJ, Kaczmarek LK. Slick

(Slo2.1), a rapidly-gating sodium-activated potassium channel inhibited by ATP. *J*

Neurosci. Dec 17 2003;23(37):11681-91.

31. Rosker C, Lohberger B, Hofer D, Steinecker B, Quasthoff S, Schreibmayer W. The

TTX metabolite 4, 9-anhydro-TTX is a highly specific blocker of the Nav1. 6 voltage-dependent sodium channel. *American Journal of Physiology-Cell Physiology*. 2007;293(2):C783-C789.

32. Rush AM, Dib-Hajj SD, Waxman SG. Electrophysiological properties of two axonal sodium channels, Nav1.2 and Nav1.6, expressed in mouse spinal sensory neurones. *J Physiol*. May 1 2005;564(Pt 3):803-15. doi:10.1113/jphysiol.2005.083089

33. Chatelier A, Zhao J, Bois P, Chahine M. Biophysical characterisation of the persistent sodium current of the Nav1.6 neuronal sodium channel: a single-channel analysis. *Pflugers Arch*. Jun 2010;460(1):77-86. doi:10.1007/s00424-010-0801-9

34. Urbani A, Belluzzi O. Riluzole inhibits the persistent sodium current in mammalian CNS neurons. *Eur J Neurosci*. Oct 2000;12(10):3567-74. doi:10.1046/j.1460-9568.2000.00242.x

35. Lukacs P, Foldi MC, Valanszki L, et al. Non-blocking modulation contributes to sodium channel inhibition by a covalently attached photoreactive riluzole analog. *Sci Rep*. May 25 2018;8(1):8110. doi:10.1038/s41598-018-26444-y

36. Hu W, Tian C, Li T, Yang M, Hou H, Shu Y. Distinct contributions of Na(v)1.6 and Na(v)1.2 in action potential initiation and backpropagation. *Nat Neurosci*. Aug 2009;12(8):996-1002. doi:10.1038/nn.2359

37. Liu Y, Lai S, Ma W, et al. CDYL suppresses epileptogenesis in mice through repression of axonal Nav1.6 sodium channel expression. *Nat Commun*. Aug 25 2017;8(1):355. doi:10.1038/s41467-017-00368-z

38. Takanishi CL, Bykova EA, Cheng W, Zheng J. GFP-based FRET analysis in live

cells. *Brain Res*. May 26 2006;1091(1):132-9. doi:10.1016/j.brainres.2006.01.119

39. Dilena R, DiFrancesco JC, Soldovieri MV, et al. Early Treatment with Quinidine

in 2 Patients with Epilepsy of Infancy with Migrating Focal Seizures (EIMFS) Due to

Gain-of-Function KCNT1 Mutations: Functional Studies, Clinical Responses, and

Critical Issues for Personalized Therapy. *Neurotherapeutics*. Oct 2018;15(4):1112-1126.

doi:10.1007/s13311-018-0657-9

40. Abdelnour E, Gallentine W, McDonald M, Sachdev M, Jiang Y-H, Mikati MA.

Does age affect response to quinidine in patients with KCNT1 mutations? Report of

three new cases and review of the literature. *Seizure*. 2018;55:1-3.

41. Rizzo F, Ambrosino P, Guacci A, et al. Characterization of two de novoKCNT1

mutations in children with malignant migrating partial seizures in infancy. *Mol Cell

Neurosci*. Apr 2016;72:54-63. doi:10.1016/j.mcn.2016.01.004

42. Barcia G, Chemaly N, Kuchenbuch M, et al. Epilepsy with migrating focal seizures:

KCNT1 mutation hotspots and phenotype variability. *Neurol Genet*. Dec

2019;5(6):e363. doi:10.1212/NXG.0000000000000363

43. Nadler JV. Kainic acid as a tool for the study of temporal lobe epilepsy. *Life

sciences*. 1981;29(20):2031-2042.

44. Levesque M, Avoli M. The kainic acid model of temporal lobe epilepsy. *Neurosci

Biobehav R*. Dec 2013;37(10):2887-2899. doi:10.1016/j.neubiorev.2013.10.011

45. Pinel JP, Rovner LI. Electrode placement and kindling-induced experimental

epilepsy. *Exp Neurol*. Jan 15 1978;58(2):335-46. doi:10.1016/0014-4886(78)90145-0

46. Kim EC, Zhang J, Tang AY, et al. Spontaneous seizure and memory loss in mice expressing an epileptic encephalopathy variant in the calmodulin-binding domain of Kv7.2. *Proc Natl Acad Sci U S A*. Dec 21 2021;118(51)doi:10.1073/pnas.2021265118

47. Ehinger R, Kuret A, Matt L, et al. Slack K(+) channels attenuate NMDA-induced excitotoxic brain damage and neuronal cell death. *FASEB J*. May 2021;35(5):e21568. doi:10.1096/fj.202002308RR

48. Markham MR, Kaczmarek LK, Zakon HH. A sodium-activated potassium channel supports high-frequency firing and reduces energetic costs during rapid modulations of action potential amplitude. *J Neurophysiol*. Apr 2013;109(7):1713-23. doi:10.1152/jn.00875.2012

49. Kole MH, Ilschner SU, Kampa BM, Williams SR, Ruben PC, Stuart GJ. Action potential generation requires a high sodium channel density in the axon initial segment. *Nat Neurosci*. Feb 2008;11(2):178-86. doi:10.1038/nn2040

50. Lazarov E, Dannemeyer M, Feulner B, et al. An axon initial segment is required for temporal precision in action potential encoding by neuronal populations. *Science advances*. 2018;4(11):eaau8621.

51. Katz E, Stoler O, Scheller A, et al. Role of sodium channel subtype in action potential generation by neocortical pyramidal neurons. *Proc Natl Acad Sci U S A*. Jul 24 2018;115(30):E7184-E7192. doi:10.1073/pnas.1720493115

52. Brown MR, Kronengold J, Gazula VR, et al. Amino-termini isoforms of the Slack

K+ channel, regulated by alternative promoters, differentially modulate rhythmic firing and adaptation. *J Physiol*. Nov 1 2008;586(21):5161-79. doi:10.1113/jphysiol.2008.160861

53. Hite RK, MacKinnon R. Structural Titration of Slo2.2, a Na(+)-Dependent K(+) Channel. *Cell*. Jan 26 2017;168(3):390-399 e11. doi:10.1016/j.cell.2016.12.030

54. Biton B, Sethuramanujam S, Picchione KE, et al. The antipsychotic drug loxapine is an opener of the sodium-activated potassium channel slack (Slo2.2). *J Pharmacol Exp Ther*. Mar 2012;340(3):706-15. doi:10.1124/jpet.111.184622

55. Gertler T, Bearden D, Bhattacharjee A, Carvill G. KCNT1-Related Epilepsy. In: Adam MP, Ardinger HH, Pagon RA, et al, eds. *GeneReviews((R))*. 1993.

56. Fitzgerald MP, Fiannacca M, Smith DM, et al. Treatment Responsiveness in KCNT1-Related Epilepsy. *Neurotherapeutics*. Jul 2019;16(3):848-857. doi:10.1007/s13311-019-00739-y

57. Griffin AM, Kahlig KM, Hatch RJ, et al. Discovery of the First Orally Available, Selective KNa1.1 Inhibitor: In Vitro and In Vivo Activity of an Oxadiazole Series. *ACS Med Chem Lett*. Apr 8 2021;12(4):593-602. doi:10.1021/acsmedchemlett.0c00675

58. Spitznagel BD, Mishra NM, Qunies AM, et al. VU0606170, a Selective Slack Channels Inhibitor, Decreases Calcium Oscillations in Cultured Cortical Neurons. *ACS Chem Neurosci*. Nov 4 2020;11(21):3658-3671. doi:10.1021/acschemneuro.0c00583

59. Tang QY, Zhang FF, Xu J, et al. Epilepsy-Related Slack Channel Mutants Lead to Channel Over-Activity by Two Different Mechanisms. *Cell Rep*. Jan 5 2016;14(1):129-

139. doi:10.1016/j.celrep.2015.12.019

60. Qiu S, Hua YL, Yang F, Chen YZ, Luo JH. Subunit assembly of N-methyl-d-aspartate receptors analyzed by fluorescence resonance energy transfer. *J Biol Chem*. Jul 1 2005;280(26):24923-30. doi:10.1074/jbc.M413915200

61. Cheng W, Yang F, Takanishi CL, Zheng J. Thermosensitive TRPV channel subunits coassemble into heteromeric channels with intermediate conductance and gating properties. *Journal of General Physiology*. 2007;129(3):191-207.

62. Zou L, Peng Q, Wang P, Zhou B. Progress in Research and Application of HIV-1 TAT-Derived Cell-Penetrating Peptide. *J Membr Biol*. Apr 2017;250(2):115-122. doi:10.1007/s00232-016-9940-z

63. Zheng J, Trudeau MC, Zagotta WN. Rod cyclic nucleotide-gated channels have a stoichiometry of three CNGA1 subunits and one CNGB1 subunit. *Neuron*. 2002;36(5):891-896.

64. Yang F, Cui Y, Wang K, Zheng J. Thermosensitive TRP channel pore turret is part of the temperature activation pathway. *Proceedings of the National Academy of Sciences*. 2010;107(15):7083-7088.

65. Erickson MG, Alseikhan BA, Peterson BZ, Yue DT. Preassociation of calmodulin with voltage-gated Ca2+ channels revealed by FRET in single living cells. *Neuron*. 2001;31(6):973-985.

66. Chai AP, Chen XF, Xu XS, et al. A Temporal Activity of CA1 Neurons Underlying Short-Term Memory for Social Recognition Altered in PTEN Mouse Models of Autism

Spectrum Disorder. *Front Cell Neurosci*. 2021;15:699315.

doi:10.3389/fncel.2021.699315

67. Huang Z, Walker MC, Shah MM. Loss of dendritic HCN1 subunits enhances

cortical excitability and epileptogenesis. *J Neurosci*. Sep 2 2009;29(35):10979-88.

doi:10.1523/JNEUROSCI.1531-09.2009

68. He X-P, Kotloski R, Nef S, Luikart BW, Parada LF, McNamara JO. Conditional

deletion of TrkB but not BDNF prevents epileptogenesis in the kindling model. *Neuron*.

2004;43(1):31-42.

69. Racine RJ. Modification of seizure activity by electrical stimulation. II. Motor

seizure. *Electroencephalogr Clin Neurophysiol*. Mar 1972;32(3):281-94.

doi:10.1016/0013-4694(72)90177-0

70. Huang Z, Lujan R, Martinez-Hernandez J, Lewis AS, Chetkovich DM, Shah MM.

TRIP8b-independent trafficking and plasticity of adult cortical presynaptic HCN1

channels. *J Neurosci*. Oct 17 2012;32(42):14835-48. doi:10.1523/JNEUROSCI.1544-

12.2012

Main Figures for

Coupling of Slack and Nav1.6 sensitizes Slack to quinidine blockade and guides

anti-seizure strategy development

Tian Yuan *et al.*

*Corresponding author. Email: huangz@hsc.pku.edu.cn

This PDF file includes:

Figure. 1 to 7

Figure 1. **Nav1.6 specifically sensitizes Slack to quinidine blockade.** (**A**) The voltage

protocol and current traces from control (non-transfected) HEK293 cells. The arrows

on the voltage protocol indicate the onset of inward sodium currents through Nav

channels and delayed outward potassium currents through Slack channels. The currents

were evoked by applying 600-ms step pulses to voltages varying from -120 mV to +100 mV in 10 mV increments, with a holding potential of -90 mV and a stimulus frequency of 0.20 Hz. (**B**) Example current traces from HEK293 cells expressing Slack alone. The left traces show the family of control currents; the right traces show Slack currents remaining after application of 30 μM quinidine in the bath solution. (**C-E**) Example current traces from HEK293 cells co-expressing Slack with $Na_V1.2$ (**C**), $Na_V1.6$ (**D**), or $Na_V1.5$ (**E**) channels before and after application of 30 μM quinidine. (**F**) The concentration-response curves for blocking of Slack by quinidine at +100 mV upon expression of Slack alone (n = 6) and co-expression of Slack with $Na_V1.1$ (n = 7), $Na_V1.2$ (n = 10), $Na_V1.3$ (n = 13), $Na_V1.5$ (n = 9), or $Na_V1.6$ (n = 19). Please refer to Supplementary Table 1 for IC_{50} values. (**G-H**) Delayed outward currents in primary cortical neurons from postnatal (P0-P1) homozygous $Na_V1.6$ knockout C3HeB/FeJ mice ($Na_V1.6$-KO) (**H**) and the wild-type littermate controls (WT) (**G**). Current traces were elicited by 600-ms step pulses to voltages varying from -120 mV to +100 mV in 20 mV increments, with a holding potential of -70 mV, and recorded with different bath solutions in the following order: Na^+-based bath solution ($I_{Control}$), replacement of external Na^+ with Li^+ in equivalent concentration (I_{Li}), washout of quinidine by Na^+-based bath solution (I_{Wash}), Na^+-based bath solution with 3 μM quinidine (I_{Quid}), Li^+-based bath solution with 3 μM quinidine ($I_{Li+Quid}$). The removal and subsequent replacement of extracellular Na^+ revealed the I_{KNa} in neurons. (**I-J**), The sensitivity of native sodium-activated potassium currents (I_{KNa}) to 3 μM quinidine blockade in WT (**I**) and $Na_V1.6$-KO (**J**) neurons. I_{KNa} before application of quinidine was obtained from the subtraction of $I_{Control}$ and I_{Li}. Maintained I_{KNa} after application of 3 μM quinidine was obtained from the subtraction of I_{Quid} and $I_{Li+Quid}$. (**K**) Summarized amplitudes of I_{KNa} before and after application of 3 μM quinidine in the bath solution in WT (black,

44 n = 12) and Nav1.6-KO (red, n = 10) primary cortical neurons. ****$p < 0.0001$, Two-

45 way repeated measures ANOVA followed by Bonferroni's post hoc test.

Figure 2. Blocking transient sodium influx through Nav1.6 reduces Nav1.6-

mediated sensitization of Slack to quinidine blockade. (**A**) Example current traces

from HEK293 cells expressing Slack alone (top) and co-expressing Slack with Nav1.6

(bottom), with 100nM TTX in the bath solution. The left traces show the family of

control currents; the right traces show Slack currents remaining after application of

quinidine. The presented concentrations of quinidine were chosen to be near the IC_{50}

values. (**B**) The concentration-response curves for blocking of Slack by quinidine at

+100 mV upon expression of Slack alone (n = 3) and co-expression of Slack with

$Na_V1.6$ (n = 7), with 100 nM TTX in the bath solution. (**C**) Top, example current traces

recorded from a HEK293 cell co-expressing Slack with $Na_V1.6$ and evoked from a 100-

ms prepulse (pre) of -90 mV, with the same voltage protocol as in Fig. 1D. Bottom,

example current traces recorded from the same cell but evoked from a 100-ms prepulse

of -40 mV, before and after application of quinidine. (**D**) I-V curves of Slack upon co-

expression with $Na_V1.6$. The currents were evoked from a prepulse of -90 mV (black)

or -40 mV (blue). (**E**) The concentration-response curves for blocking of Slack by

quinidine at +100 mV with a prepulse of -90 mV (black, n = 19) or -40 mV (blue, n =

5). (**F**) Top, example current traces recorded from a HEK293 cell co-expressing Slack

and $Na_V1.6$ without riluzole in the bath solution. Bottom, example current traces

recorded from the same cell with 20 µM riluzole in the bath solution, before and after

application of quinidine. (**G**) I-V curves of Slack upon co-expression with $Na_V1.6$

before (black) and after (red) application of 20 µM riluzole into bath solution.The

concentration-response curves for blocking of Slack by quinidine upon co-expression

of Slack with $Na_V1.6$, without (n = 19) or with (n = 6) 20 µM riluzole in the bath

solution. (**I-J**) The sensitivity of Na_V channel subtypes to quinidine blockade upon

expression of Na_V alone in HEK293 cells. Example current traces (**I**) were evoked by

a 50-ms step depolarization to 0 mV from a holding potential of -90 mV. The

Concentration-response curves for blocking of Na_V channel subtypes by quinidine (**J**)

were shown on the right panel (n = 5 for Nav1.1, n = 3 for Nav1.2, n = 6 for Nav1.3, n

= 6 for Nav1.5, and n = 4 for Nav1.6).

Figure 3. Slack physically interacts with Nav1.6. (**A**) Immunofluorescence of Slack,

Nav1.2, Nav1.6 (green), and AnkG (red) in neocortex layer 5 (left) and hippocampal

CA1 pyramidal cell layer (right). Confocal microscopy images were obtained from

Coronal brain slices of C57BL/6 mice. The panels from top to bottom show the double

staining of Slack with AnkG, Nav1.2 with AnkG, and Nav1.6 with AnkG, respectively.

DAPI (blue) was used for nuclear counterstaining. (**B**) Coimmunoprecipitation (Co-IP)

of Slack and Nav1.6 in cell lysates from HEK293T cells co-transfected with Slack and

Nav1.6. (**C**) Co-IP of Slack and Nav1.6 in mouse brain tissue lysates. Input volume

corresponds to 10% of the total lysates for Co-IP. (**D**) A schematic diagram showing

the fluorescence-labeled Slack and Nav1.6. mTFP1 and mVenus were fused to the C-

terminal region of Slack (Slack-mTFP1) and Nav1.6 (Nav1.6-mVenus), respectively.

(**E**) FRET imaging of Slack-mTFP1 and Nav1.6-mVenus co-expressed in HEK293

cells. The emission spectra measured from the edge of cell (dotted arrows in red) are

used for FRET efficiency calculation. (**F**) The apparent FRET efficiency measured

from cells co-expressing the fluorophore-tagged ion channels or co-expressing the

fluorophores. **** $p < 0.0001$, Mann-Whitney test. (**G-H**) The FRET efficiency

measured from cells co-expressing the fluorophore-tagged ion channels (**G**), or from

cells co-expressing fluorophores (**H**). The efficiency value was plotted as a function of

the fluorescence intensity ratio between mTFP1 and mVenus (Fc/Fy). Each symbol

represents a single cell. The solid curve represents the FRET model that yields the best

fit; dotted curves represent models with 5% higher or lower FRET efficiencies.

Figure 4. Nav1.6's N- and C-termini interacting with Slack is a prerequisite for Nav1.6-mediated sensitization of Slack to quinidine blockade. (**A**) The sensitivity of Slack to quinidine blockade upon expression of Slack alone (n = 3) and co-expression

of Slack with Nav1.6 (n = 3) from excised inside-out patches. The pipette solution contained (in mM) 130 KCl, 1 EDTA, 10 HEPES and 2 MgCl$_2$ (pH 7.3); the bath solution contained (in mM) 140 NaCl, 1 EDTA, 10 HEPES and 2 MgCl$_2$ (pH 7.4). The membrane voltage was held at 0 mV and stepped to voltages varying from −100 mV to 0 mV in 10 mV increments. Example current traces were shown on the left panel. The concentration-response curves for blocking of Slack by quinidine were shown on the right panel. (**B**) Domain architecture of the human Nav channel pore-forming α subunit. (**C**) Calculated IC$_{50}$ values at +100 mV of quinidine on Slack upon co-expression with indicated cytoplasmic fragments from Nav channels. For Nav1.6, cytoplasmic fragments used include N-terminus (Nav1.6-N, residues 1-132), inter domain linkers (Domain I-II linker, residues 409-753; Domain II-III linker, residues 977-1199; Domain III-IV linker, residues 1461-1523), and C-terminus (Nav1.6-C, residues 1766-1980). For Nav1.5, cytoplasmic fragments used include N-terminus (Nav1.5-N, residues 1-131) and C-terminus (Nav1.5-C, residues 1772-2016). (**D**) Co-IP of Slack and terminal domains of Nav1.6 in cell lysates from HEK293T cells co-expressing 3×Flag-tagged Slack (Slack-3×Flag) and 3×HA-tagged termini of Nav1.6 (3×HA-Nav1.6-N or 3×HA-Nav1.6-C). The 3×Flag tag was fused to the C-terminal region of Slack and the 3×HA tag was fused to the N-terminal region of Nav1.6's fragments. (**E**) The sensitivity of Slack to quinidine blockade upon co-expression of Slack with GFP (n = 12) or N- and C-termini of Nav1.6 (n = 11), from whole-cell recordings. (**F**) The sensitivity of Slack to quinidine blockade upon co-expression of Slack with N- and C-termini of Nav1.6, from excised inside-out recordings (n = 10, using the same protocols as in Fig. 4A). Example current traces before and after application of quinidine were shown on the left panel. The concentration-response curves were shown on the right panel. (**G**) A schematic diagram of the Nav1.5-1.6 chimeric channels (Nav1.5/6[NC] and

130   Nav1.5/6^N) used in this study. (**H**) Example current traces recorded from HEK293 cells

131   co-expressing Slack and Nav1.5-1.6 chimeras before and after application of the

132   indicated concentration of quinidine. (**I**) The concentration-response curves for

133   blocking of Slack by quinidine upon co-expression of Slack with Nav1.5 (n = 9),

134   Nav1.6 (n = 19), Nav1.5/6^{NC} (n = 9), or Nav1.5/6^N (n = 9).

Figure 5. Slack's C-terminus is required for Nav1.6-mediated sensitization of Slack to quinidine blockade. (**A**) Domain architecture of the human Slack channel subunit. Slack's N-terminus (Slack-N, residues 1-116) and C-terminus (Slack-C, residues 345-1235) were shown in the blue boxes. (**B**) The concentration-response curves for blocking of Slack by quinidine upon additional expression of Slack's N- or C-terminus in HEK293T cells co-expressing Slack and Nav1.5/6^{NC}. (**C**) The concentration-response curves for blocking Slack by quinidine upon additional expression of Slack's C-terminus in HEK293 cells co-expressing Slack and Nav1.6. (**D**) Co-IP of Myc-tagged Slack's C-terminus (Slack-C-Myc) with 3×HA-tagged Nav1.6's termini (3×HA-Nav1.6-N or 3×HA-Nav1.6-C) in HEK293T cell lysates. The 3×HA tag was fused to the N-terminal region of Nav1.6's fragments, and the Myc tag was fused to the C-terminal region of Slack's fragment.

Figure 6. Na$_V$1.6 sensitizes epilepsy-related Slack mutant variants to quinidine blockade. (**A**) Co-IP of 3×Flag-tagged Slack or its mutations (Slack-3×Flag) with 3×HA-tagged Na$_V$1.6 (Na$_V$1.6-3×HA) in HEK293T cell lysates. The tags were all fused to the C-terminal region of wild-type or mutant ion channels. (**B-D**) The sensitivity of Slack mutant variants (Slack^{K629N} [**B**], Slack^{R950Q} [**C**], and Slack^{K985N} [**D**]) to quinidine blockade upon expression of Slack mutant variants alone and co-expression of Slack

mutant variants with Nav1.6. Left, example current traces recorded from HEK293 cells

expressing Slack mutant variants alone and co-expressing Slack mutant variants with

Nav1.6, before and after application of the indicated concentrations of quinidine. Right,

the concentration-response curves for blocking of Slack mutant variants by quinidine

upon expression of Slack mutant variants alone (n = 8 for Slack^{K629N}, n = 7 for

Slack^{R950Q}, and n = 5 for Slack^{K985N}) and co-expression of Slack mutant variants with

Nav1.6 (n = 8 for Slack^{K629N} upon co-expression with Nav1.6, n = 5 for Slack^{R950Q} upon

co-expression with Nav1.6, and n = 7 for Slack^{K985N} upon co-expression with Nav1.6).

Please refer to Supplementary Table 4 for IC$_{50}$ values.

Figure 7. Viral expression of Slack's C-terminus prevents Slack^{G269S}-induced

seizures. (**A-B**) The current densities of Slack mutant variants (Slack^{G288S} [**A**] and

Slack^{R398Q} [**B**]) upon co-expression with Nav1.5/6^{NC} in HEK293T cells were reduced

by additional expression of Slack's C-terminus. Left, example current traces from

HEK293T cells co-expressing Slack mutant variants and Nav1.5/6^{NC} or co-expressing

Slack mutant variants, Nav1.5/6^{NC}, and Slack's C-terminus. Right, summarized current

densities at +100 mV. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; one-way ANOVA

followed by Bonferroni's post hoc test. (**C**) Architecture for expression cassettes of

AAVs. (**D**) Top, study design and timeline for the stereotactic injection model. Bottom,

Immunofluorescence of HA-tagged Slack^{G269S} (green), 3×Flag-tagged Slack's C-

terminus (red), and DAPI (blue) in hippocampal CA1 pyramidal cell layer at 5 weeks

after viral injection of Slack^{G269S} with Slack's C-terminus into CA1 of mice. (**E**) Time-

course of KA-induced seizure stage changes at 10-min intervals based on a modified

Racine, Pinal, and Rovner scale (please refer to Methods for further details). The

number of mice used: "GFP" control group (n = 11), "Slack^{G269S}+GFP" group (n = 12),

"Slack^{G269S}+Slack-C" group (n = 12). "GFP" vs. "Slack^{G269S}+GFP": $F_{(1,21)} = 10.48$, p

$= 0.0040$, * $p < 0.05$, ** $p < 0.01$; "Slack^{G269S}+GFP" vs. "Slack^{G269S}+Slack-C": $F_{(1,22)}$

$= 10.30$, $p = 0.0040$, # $p < 0.05$, ## $p < 0.01$. "GFP" vs. "Slack^{G269S}+Slack-C": $F_{(1,21)}$

$= 0.09574$, $p = 0.7600$. Repeated two-way ANOVA followed by Bonferroni's post

hoc test. (**F**) Total seizure score per mouse over the 2 h after KA injection of these three

groups. * $p < 0.05$, ** $p < 0.01$; one-way ANOVA followed by Bonferroni's post hoc

test. (**G**) The percentage of mice with stage VI~IX seizures over the 2 h after KA

injection in each group. * $p < 0.05$; Fisher's exact test.