Autophagy functions in lung macrophages and dendritic cells to suppress T$_H$17 responses

and neutrophil accumulation during allergic airway inflammation

Neha Dubey[1], Reilly Woodson[1], Samuel R. McKee[1], Anne L. Rosen[2], Nicole Rivera-Espinal[1],

Darren Kreamalmeyer[1], Andrew L. Kau[2], Christina L. Stallings[1*]

[1]Department of Molecular Microbiology and Center for Women's Infectious Disease Research,

Washington University School of Medicine, St. Louis, MO 63110, USA

[2]Division of Allergy and Immunology, Department of Medicine and Center for Women's

Infectious Disease Research, Washington University School of Medicine, St. Louis, MO 63110,

USA

*Correspondence: stallings@wustl.edu

Keywords: asthma; allergy; autophagy; inflammation; myeloid cells; neutrophils; T cells; T$_H$17

ABSTRACT

Asthma affects nearly 260 million people worldwide, where severe asthma cases represent the

most difficult to treat due to corticosteroid insensitivity. Severe asthma is associated with higher

levels of T$_H$17 and T$_H$1 responses, accompanied by neutrophil dominated inflammation. Better

understanding of the immune responses to airway allergens that promote or protect against severe

asthma is critical for identifying ways to treat these patients. Single nucleotide polymorphisms in

the *ATG5* gene, which encodes for a protein required for the cellular recycling process of

autophagy, has been associated with higher risk for developing severe asthma. However, the exact

role for ATG5 during allergic inflammation and whether other autophagy proteins are involved

remains elusive. Using genetic tools to dissect the roles for ATG5 in innate immune cells in house

dust mite (HDM)-challenged mice, we have identified a specific role for ATG5 in CD11c$^+$ lung

macrophages and dendritic cells for suppressing T$_H$17 responses and neutrophil accumulation. We

found that this role for ATG5 in CD11c$^+$ cells to regulate neutrophil accumulation during allergic

airway inflammation also required other autophagy proteins but did not involve regulation of

inflammasome, despite higher levels of IL-1β and Caspase 1 in the lungs of mice lacking

autophagy in innate immune cells. Our data support a role for autophagy in CD11c[+] lung

macrophages and dendritic cells to promote an immune response to airway allergens that is

associated with less severe asthma by suppressing T_H17 responses and neutrophil accumulation in

the lung.

INTRODUCTION

Asthma affects nearly 260 million people worldwide[1] and involves a deleterious inflammatory

response that results in immune cell infiltration in the lungs, bronchoconstriction, airway

remodeling, and airway hyperresponsiveness (AHR)[2]. Asthma and allergic airway inflammation

occurs in response to a wide variety of substance called allergens that induce varying levels of

immune hypersensitivity[3] The most prevalent form of asthma is associated with T_H2 responses

signaled through airway epithelial cell-derived alarmins IL-33 and IL-25, which stimulate the

production of IL-4, IL-5, IL-13, and TGF-β cytokines by airway cells and an influx of eosinophils

and mast cells, along with other inflammatory cells, into the lungs[4–6]. Although this form of the

disease is usually less severe and responsive to conventional corticosteroid therapy[4], asthma can

also present with other distinct immunological and pathological features[7–9]. In particular, 5-15%

of the total asthma cases involve a T_H17-dominated response, leading to neutrophil recruitment

and accumulation in the lungs[10,11]. Increased accumulation of neutrophils is associated with severe

asthma[10,11], which is clinically defined by resistance to corticosteroid therapy[8,12–14]. In addition,

T_H1 responses mediated by interferon-γ (IFN-γ) production by CD4[+] T cells are also associated

with severe asthma cases[15,16]. The association of heightened T_H17 responses, T_H1 responses, and

neutrophil accumulation with AHR and corticosteroid resistance has been reported during severe

asthma in both mice and humans[14–17].

A better understanding of the immunological processes that contribute to severe asthma is

required to develop alternative therapies for when corticosteroids fail. The T_H17 polarization of T

cells that is associated with severe asthma is stimulated by airway epithelial cell derived cytokines

IL-6, IL-1β, TGF-β and IL33[12,18] that are produced upon activation of several pattern recognition

receptors in response to the allergen[19–23]. Simultaneously, IL-23 produced by macrophages and

dendritic cells (DCs) plays a critical role in differentiation of IL-17A (herein IL-17) producing

T_H17 cells[24–26]. IL-17 produced by T cells further induces neighboring cells, including bronchial

epithelial cells and fibroblasts, to produce CXCL8 and IL-8 (KC), thereby leading to the

recruitment of neutrophils to the lungs[27–29]. Studies in murine asthma models have shown that

depletion of neutrophils or blocking IL-17 signaling can reduce neutrophil accumulation in the

lungs, rescue the pathology associated with severe asthma, and restore the efficacy of

corticosteroids[14,15,30–32]. In addition, adoptive transfer of antigen specific T_H17 cells resulted in

excessive neutrophilic infiltration and higher AHR in ovalbumin (OVA) sensitized mice[13].

Together, these reports support a direct contribution of T_H17 responses and neutrophils to the

development of severe asthma, highlighting the need for a better understanding of how neutrophil

recruitment and accumulation during asthma is regulated.

Several studies attempting to map genetic associations with asthma have identified a link

between the *ATG5* gene and susceptibility to severe asthma. A single nucleotide polymorphism

(SNP) in the third intron of the *ATG5* gene (rs12212740, alleles G/A, minor variant: G) was

associated with lower pre-bronchodilator FEV1 (forced expiratory volume) and a risk of moderate

to severe asthma in non-Hispanic Caucasian children[33]. In a separate genome wide association

study, a SNP in the 5' untranslated region (UTR)/annotated promoter region of the *ATG5* gene

(rs510432, alleles G/A, minor variant: G) was associated with increased risk of asthma in

children[34]. The G variant allele of SNP rs510432 results in increased promoter activity and higher

expression of *ATG5* mRNA in nasal mucosal cells isolated from the children with acute asthma

compared to non-acute and healthy controls. Another study reported the association of the G

variant allele of SNP rs510432 with increased sputum neutrophil counts in adult asthma patients,

establishing a correlation between dysregulated *ATG5* expression and neutrophil recruitment

during asthma[35].

ATG5 is essential for the process of autophagy, which involves multiple protein complexes

that target cytosolic components and organelles to the lysosome for degradation[36–38]. The ULK1

complex (ULK1/ULK2, ATG13, FIP200, ATG101) and PI3 kinase complex (ATG14L, BECN1,

VPS15, and VPS34) initiate autophagy by generating a phagophore from the membranes of the

endoplasmic reticulum[39,40]. Elongation of the autophagosomal double membrane depends on two

ubiquitin-like conjugation systems. In the first system, ATG12 (ubiquitin-like protein) is activated

by ATG7 (E1-like), transferred to ATG10 (E2-like), and covalently attached to ATG5. In the

second system, LC3 (microtubule-associated protein 1 light chain 3) is conjugated to

phosphatidylethanolamine to generate the membrane bound form LC3-II through the actions of

ATG7 (E1-like), ATG3 (E2-like), and the ATG5-ATG12/ATG16L complex[40,41]. The

autophagosome membrane is then completed and targeted for fusion with the lysosome, where the

autophagosome cargo are degraded. Several studies have reported that the expression of genes

required for autophagy, including *ATG5*, are elevated in the lungs of asthma patients compared to

non-asthmatic individuals[42–44]. Furthermore, LC3-II expression is increased in the sputum

granulocytes from severe asthma patients compared to non-severe asthma patients[35,45]. Despite the

genetic associations of ATG5 and autophagy with severe asthma, the precise roles for autophagy

during asthma have remained elusive. Several experimental studies have implicated both

deleterious and protective functions of autophagy during asthma[46], creating uncertainty in the field.

There are likely multiple contributors to the discrepancies. For example, in most of the

experimental studies so far, the interpretations were largely made based on pleiotropic chemical

treatments that impact autophagy, among other pathways[47–53]. In addition, autophagy likely plays

cell-type specific roles during exposure to allergens, which may complicate studies that that use

chemical treatments or globally knockdown autophagy gene expression[14].

ATG5 has recently been shown to function both dependent and independent of autophagy

to regulate inflammatory responses in innate immune cells during infection[36,37,54–57]. Innate

immune cells are also a primary contributor to lung inflammation during asthma. However, studies

into the roles for autophagy specifically in innate immune cells during asthma and allergic airway

inflammation have been limited. In the one study that focused on ATG5 in innate immune cells,

loss of *Atg5* in hematopoietic cells was sufficient to lead to increased neutrophils and macrophages

in the bronchoalveolar lavage (BAL), increased AHR, and higher levels of IL-17 and IL-1β in

response to house dust mites (HDM) challenge in mice[14]. When *Atg5*-deficient bone marrow cells

were cultured in the presence of GM-CSF to generate a mixed culture of DCs, macrophages, and

granulocytes, and treated with HDM, the cells produced higher levels of IL-1α, IL-1β, IL-23, and

caspase 1 as compared to wild type (WT) GM-CSF cultures. *Atg5*$^{-/-}$ GM-CSF cultures also caused

a higher level of IL-17 production in a co-culture with CD4$^+$ T cells as compared to WT GM-CSF

cultures. Transfer of *Atg5*$^{-/-}$ GM-CSF cultured cells loaded with HDM into WT mice

intratracheally resulted in increased neutrophilic inflammation and IL-17 production as compared

to transfer of WT GM-CSF cultured cells, indicating that ATG5 is important to control neutrophil

recruitment in response to HDM exposure. These studies support a role for ATG5 in innate

immune cells to control neutrophilic inflammation during allergic challenge. However, it is still

unknown which innate immune cell types specifically require ATG5 and whether its role in

controlling neutrophilic inflammation during allergic challenge is dependent on autophagy, since

ATG5 and other autophagy proteins have been shown to function outside of autophagy to regulate

inflammation[36,38,56,57]. To address these gaps in understanding, we used genetic tools to dissect the

autophagy-dependent and independent roles for ATG5 in specific innate immune cells during

HDM challenge *in vivo* in mice. We find that autophagy is specifically required in CD11c$^+$ lung

macrophages and dendritic cells to suppress T$_H$17 responses and control the accumulation of

neutrophils during allergic airway inflammation. These findings highlight a protective role for

autophagy in lung resident innate immune cells during allergic challenge.

RESULTS

Loss of *Atg5* expression in LysM$^+$ cells results in increased neutrophil-dominated

inflammation upon HDM challenge

The role of ATG5 and autophagy in specific innate immune cell types during asthma has yet to be

tested since all prior work was done with general administration of pleiotropic and non-cell type

specific treatments[43,47,51,52] or using cell cultures containing a mixed population of cell types[14]. To

determine if ATG5 functions in innate immune cells during allergic airway inflammation, we used

Atg5[fl/fl]-LysM-Cre mice[36,54,58,59], which specifically delete *Atg5* in LysM$^+$ monocytes,

macrophages, neutrophils, and dendritic cells (DCs). Analysis of the immune cell populations in

the lungs of naïve *Atg5*[fl/fl] control mice and *Atg5*[fl/fl]*-LysM-Cre* mice by flow cytometry revealed no

significant differences **(Figure 1, S1)**, similar to previous reports[36]. We then challenged the *Atg5*[fl/fl]

and *Atg5*[fl/fl]*-LysM-Cre* mice with HDM intranasally on day 1, day 8 and day 15 before analyzing

lung inflammation on day 16 by flow cytometry **(Figure 1A)**, which was the same regimen and

timeline used in the prior study that explored the effects of transferring Atg5$^{-/-}$ GM-CSF cultured

cells into the lungs of mice[14]. HDM challenge resulted in an increase in the numbers of immune

cells in the lungs of mice from both genotypes compared to naïve mice **(Figure 1, S1)**, providing

evidence that we are achieving an allergic airway inflammatory response. The predominant innate

immune cell type recruited to the lungs in both *Atg5*[fl/fl] and *Atg5*[fl/fl]*-LysM-Cre* mice following HDM

challenge was neutrophils **(Figure 1B)**. HDM-challenge of *Atg5*[fl/fl]*-LysM-Cre* mice resulted in

significantly increased recruitment and accumulation of neutrophils, non-alveolar macrophages,

DCs, and inflammatory monocytes compared to *Atg5*[fl/fl] controls **(Figure 1B, C)**. There were no

differences in the numbers of eosinophils or alveolar macrophages in the lungs of HDM-
challenged *Atg5^fl/fl^-LysM-Cre* and *Atg5^fl/fl^* mice (**Figure 1C, S1).** Analysis of the lymphoid cell
repertoire revealed higher total cell counts for $CD4^+$ T cells, but no difference in the number of
$CD8^+$ T cells or B cells in the lungs of HDM-challenged *Atg5^fl/fl^-LysM-Cre* mice compared to
*Atg5^fl/fl^* controls (**Figure 1D, S1**). Further analysis of the $CD4^+$ T cell population revealed that the
lungs of HDM-challenged *Atg5^fl/fl^-LysM-Cre* mice had an increased number of activated $CD4^+$ T
cells ($CD4^+CD44^+CD62L^-$) and T regulatory (T_{reg}) cells ($CD4^+CD25^+$) compared to *Atg5^fl/fl^* mice
(**Figure 1E, F**). Together our results show that *Atg5* is required in $LysM^+$ innate immune cells to
control neutrophil, inflammatory monocyte, macrophage, and DC recruitment and accumulation
in the lungs of mice challenged with HDM, which is associated with higher numbers of activated
$CD4^+$ T cells and T_{reg} cells in the lungs.

Loss of *Atg5* in innate immune cells results in increased production of IL-17 from $CD4^+$ T
cells
The increased accumulation of neutrophils, inflammatory monocytes, and recruited macrophages
without a change in the numbers of eosinophils, along with the increase in activated $CD4^+$ T cells,
suggests that loss of ATG5 in innate immune cells results in increased T_H17 and/or T_H1 responses,
rather than the T_H2 response traditionally associated with asthma. The increased inflammation in
HDM-challenged *Atg5^fl/fl^-LysM-Cre* mice was dominated by neutrophils (**Figure 1B**), which is
particularly suggestive of increased T_H17 responses[10,28]. To determine if the higher levels of
neutrophil accumulation in *Atg5^fl/fl^-LysM-Cre* mice is accompanied by elevated T_H17 responses,
we crossed the *Atg5^fl/fl^-LysM-Cre* and *Atg5^fl/fl^* mice with an IL-17 reporter mouse[60] (*Il17a*^eGFP^,
#018472, Jackson Laboratory) that expresses GFP as a proxy of IL-17, generating *Atg5^fl/fl^-LysM-*
Cre/Il17a^eGFP^ and *Atg5^fl/fl^/Il17a*^eGFP^ mice. Following HDM challenge, we detected a similar
increase in the abundance of neutrophils and activated $CD4^+$ T cells in *Atg5^fl/fl^-LysM-Cre/Il17a*^eGFP^
mice compared to *Atg5^fl/fl^/Il17a*^eGFP^ controls (**Figure 2A, B**) as observed in the *Atg5^fl/fl^-LysM-Cre*
and *Atg5^fl/fl^* parent strains (**Figure 1B, E**). We monitored GFP expression from immune cells
following HDM challenge of *Atg5^fl/fl^-LysM-Cre/Il17a*^eGFP^ and *Atg5^fl/fl^/Il17a*^eGFP^ mice using flow
cytometry and found that the only cell types positive for GFP were $CD4^+$ T cells. A higher
percentage of total $CD4^+$ T cells, activated $CD4^+$ T cells, and Treg cells expressed GFP in HDM-
challenged *Atg5^fl/fl^-LysM-Cre/Il17a*^eGFP^ mice compared to *Atg5^fl/fl^/Il17a*^eGFP^ mice (**Figure 2C**),

which reflected a higher number of GFP$^+$ activated CD4$^+$ T cells and T$_{regs}$ in the lungs of HDM-

challenged *Atg5$^{fl/fl}$-LysM-Cre/Il17a*^{eGFP} mice compared to controls **(Figure 2D)**. Based on mean

fluorescence intensity, the CD4$^+$ T cells in *Atg5$^{fl/fl}$-LysM-Cre/Il17a*^{eGFP} mice also expressed more

GFP per cell than CD4$^+$ T cells from *Atg5$^{fl/fl}$/Il17a*^{eGFP} mice following HDM challenge **(Figure**

2E). These data indicate that loss of *Atg5* in LysM$^+$ cells results in increased IL-17 expression

from CD4$^+$ T cells, which could represent increased T$_H$17 responses that would promote neutrophil

recruitment during allergic airway inflammation. A higher T$_H$17/T$_{reg}$ ratio is also associated with

increased asthma severity, increased AHR, and refractory asthma[61–65]. Indeed, we observed an

increase in the ratio of T$_H$17/T$_{reg}$ cells in HDM-challenged *Atg5$^{fl/fl}$-LysM-Cre/Il17a*^{eGFP} mice

compared to *Atg5$^{fl/fl}$/Il17a*^{eGFP} controls **(Figure 2F)**, supporting that loss of *Atg5* in myeloid cells

results in a CD4$^+$ T cell imbalance that is associated with increased asthma severity[61,62,64].

Autophagy is required in innate immune cells to control neutrophil-dominated inflammation

during HDM-challenge

ATG5 has been shown to control inflammatory responses via both autophagy-dependent and

autophagy-independent pathways[36,56–58,66,67]. In particular, we previously reported that ATG5 can

function independent of autophagy in innate immune cells to suppress neutrophil inflammation

during pulmonary *Mycobacterium tuberculosis* infection in mice[36]. To determine whether the

regulation of neutrophil recruitment by ATG5 in HDM-challenged mice was dependent on other

autophagy proteins or represented an autophagy-independent role for ATG5, we monitored

neutrophil abundance in the lungs of mice lacking expression of the essential autophagy genes

Atg16l1 or *Becn1* in LysM$^+$ cells following HDM challenge by flow cytometry. BECN1 operates

during the nucleation step of the autophagosome membrane biogenesis, whereas ATG16L1

functions with ATG5 in the phagophore elongation complex[40]. In response to HDM, we observed

higher numbers of neutrophils and activated CD4$^+$ T cells in the lungs of both *Atg16l1$^{fl/fl}$-LysM-*

Cre and *Becn1$^{fl/fl}$-LysM-Cre* mice compared to the *Atg16l1$^{fl/fl}$* and *Becn1$^{fl/fl}$* controls, respectively

(Figure 3A, B). The lungs of HDM-challenged *Atg16l1$^{fl/fl}$-LysM-Cre* mice also accumulated

significantly higher numbers of DCs and T$_{reg}$ cells in comparison to *Atg16l1$^{fl/fl}$* mice **(Figure 3A,**

D), and the lungs of HDM-challenged *Becn1$^{fl/fl}$-LysM-Cre* mice accumulated a higher number of

inflammatory monocytes compared to Becn1$^{fl/fl}$ mice **(Figure 3A)**. Similar to *Atg5$^{fl/fl}$-LysM-Cre*

mice, we did not observe differences in eosinophils in the HDM-challenged *Atg16l1$^{fl/fl}$-LysM-Cre*

and *Becn1$^{fl/fl}$-LysM-Cre* mice compared to controls (**Figure 3B**), indicating that the increased

inflammation was representative of enhanced T$_H$17 or T$_H$1 responses and not T$_H$2 responses. Our

results indicate that ATG5 is required in an autophagy-dependent manner to control neutrophil

accumulation and activation of CD4$^+$ T cells in the lungs of mice during HDM challenge. Together,

we show that autophagy has a protective role in innate immune cells during HDM-challenge.

Autophagy regulates neutrophil accumulation during HDM challenge independent of

inflammasome activation

A key inflammatory process negatively regulated by autophagy is inflammasome activation[68–70].

Several reports have demonstrated a role for the NLRP3 (nucleotide-binding oligomerization

domain-like Receptor Family Pyrin Domain Containing 3) inflammasome pathway, which

catalyzes caspase-1 dependent release of pro-inflammatory cytokines IL-1β and IL-18, during

severe allergic inflammation[71–74]. Elevated levels of caspase-1, IL-1β and IL-18 have been

observed in patients and mice with severe asthma[68,72,73,75]. Genetic deletion of *Nlrp3* or treatment

with NLRP3 inhibitor in mice alleviates the immune cell filtration and mucus secretion in the lungs

of mice during asthma[68,71,72]. Considering that *Atg5$^{-/-}$* bone marrow cells cultured in GM-CSF

produced higher levels of IL-1β following exposure to HDM *in-vitro*[14], we sought to investigate

if the higher neutrophil abundance and inflammation following HDM challenge is a consequence

of inflammasome activation in the absence of autophagy in *Atg5$^{fl/fl}$-LysM-Cre* mice. Since caspase-

1 is the key mediator of the inflammatory consequences of the NLRP3 inflammasome, we crossed

the *Atg5$^{fl/fl}$-LysM-Cre* and *Atg5$^{fl/fl}$* mice with *Casp1/11$^{-/-}$* germline deletion mice, which are deleted

for the genes encoding caspase-1 and caspase-11 (Jax #016621, Jackson Laboratory). We

challenged *Atg5$^{fl/fl}$/Casp1/11$^{-/-}$* and *Atg5$^{fl/fl}$-LysM-Cre/Casp1/11$^{-/-}$* mice with HDM and analyzed

the lung immune cell populations. Similar to *Atg5$^{fl/fl}$-LysM-Cre* mice, HDM-challenged *Atg5$^{fl/fl}$-*

LysM-Cre/Casp1/11$^{-/-}$ mice accumulated more neutrophils, non-alveolar macrophages, DCs,

activated CD4$^+$ T cells, and T$_{reg}$ cells in their lungs than *Atg5$^{fl/fl}$/Casp1/11$^{-/-}$* mice (**Figure 4A, B**),

indicating that elevated caspase 1-dependent NLRP3 inflammasome activation in the absence of

autophagy was not responsible for the increased neutrophil dominated inflammation during HDM

challenge of *Atg5$^{fl/fl}$-LysM-Cre* mice. Therefore, the increased inflammasome activation reported

previously in myeloid cells exposed to HDM[14] is not responsible for the neutrophilic inflammation

in HDM-challenged *Atg5$^{fl/fl}$-LysM-Cre* mice.

Autophagy is required in CD11c[+] lung macrophages and dendritic cells to control neutrophil

recruitment during exposure to HDM

Atg5[fl/fl]-LysM-Cre mice delete *Atg5* in multiple different types of innate immune cells, including

neutrophils, macrophages, inflammatory monocytes, and some DCs. Autophagy has been shown

to function in many of these cell types, including neutrophils, macrophages, and DCs, to regulate

inflammation[36,37,59,66,76]. Therefore, to determine specifically which LysM[+] cell type requires

ATG5 to control neutrophil recruitment during allergic airway inflammation, we challenged

Atg5[fl/fl]-CD11c-Cre (delete *Atg5* in lung macrophages and DCs) and *Atg5[fl/fl]-MRP8-Cre* (delete

Atg5 in neutrophils) mice[36] with HDM and monitored lung inflammatory responses. Flow

cytometric analysis of the immune cell populations in the lung revealed a significantly higher

abundance of neutrophils, non-alveolar macrophages, DCs, inflammatory monocytes, activated

CD4[+] T cells, and T_{reg} cells in *Atg5[fl/fl]-CD11c-Cre* mice (**Figures 5A-D**) but not in *Atg5[fl/fl]-MRP8-*

Cre mice (**Figures 5E-H**) compared to respective littermate *Atg5[fl/fl]* controls. Therefore, loss of

Atg5 in CD11c[+] cells was sufficient to phenocopy loss of Atg5 in LysM[+] cells, demonstrating that

ATG5 functions in lung macrophages and DCs to control neutrophil recruitment during HDM

challenge. We did not observe differences in any of the immune cell populations in the lungs of

HDM-challenged *Atg5[fl/fl]-MRP8-Cre* and *Atg5[fl/fl]* littermate control mice, demonstrating that

ATG5 is dispensable in neutrophils to control inflammation during HDM challenge.

DISCUSSION

The association of genetic polymorphisms in the *ATG5* gene with susceptibility to asthma in

humans has drawn attention to the roles for autophagy in regulating inflammatory responses during

allergen challenge and asthma. In the present study, we directly investigate the roles for ATG5 and

autophagy in innate immune cells during asthma using a series of conditional knockout mouse

lines and an HDM challenge model[14]. Our data support a model where ATG5 functions in an

autophagy-dependent process in CD11c[+] lung macrophages and/or dendritic cells to suppress T_H17

responses and neutrophil recruitment to the lungs during HDM challenge. These studies establish

a protective role of autophagy in CD11c[+] cells in a murine model of allergic airway inflammation.

Our findings build off of the observations reported previously by Suzuki *et al.*, where
tamoxifen-inducible deletion of *Atg5* in HDM-challenged mice or intratracheal instillation of
HDM-primed *Atg5*[-/-] GM-CSF myeloid cell cultures led to neutrophil recruitment to the lungs,
increased AHR, and increased production of IL-1β and IL-17[14]. In our study, we determine that
ATG5 and other autophagy proteins are specifically required in lung macrophages and DCs to
suppress the activation of T_H17 cells and the accumulation of T_{reg} cells. In addition, despite the
higher levels of IL-1β and caspase-1 previously reported, we find that inflammasome activation is
not responsible for the increased accumulation of neutrophils in HDM-challenged *Atg5*[fl/fl]*-LysM-*
Cre mice.

There are multiple possible mechanisms by which autophagy could be functioning in
CD11c[+] DCs and lung macrophages to impact T cell polarization during HDM challenge. Both
DCs and lung macrophages are antigen-presenting cells (APCs), and allergen antigen presentation
by APCs is associated with CD4[+] T cell activation during asthma[77–79]. Autophagy plays a central
role in transferring antigen to vesicles that contain MHC class II molecules for antigen presentation
to CD4[+] T cells[80,81]. In addition to direct roles in MHC class II antigen presentation, autophagy
also regulates the production of cytokines that can impact the polarization of T cells during
activation. Loss of autophagy can result in increased production of IL-23 by macrophages and
DCs[82], and IL-23 can induce the differentiation of T cells into T_H17 cells[24,83]. Macrophage
polarization also impacts T cell responses, where M1-like macrophages produce T_H1 associated
cytokines (TNF-α, IL-6, IL-1β, and IL-23) and induce a T_H1/T_H17 type inflammatory response,
while M2-like macrophages produce type 2 cytokines (IL-5 and IL-13), activating a T_H2 type
asthmatic response[84]. M1 polarized macrophages also express higher levels of the transcription
factor IRF5, which was demonstrated to drive T_H1/T_H17 responses in culture[85]. In addition,
deletion of *Irf5* in mice causes a reduction in IFN-γ, IL-17, and neutrophil recruitment in HDM-
challenged mice[86]. Loss of autophagy in macrophages promotes pro-inflammatory M1-like
macrophage polarization in inflammatory conditions such as aging-associated hepatic injury and
obesity[66,87]. Therefore, autophagy in macrophages and DCs can drive differential T cell responses
via antigen presentation and the production of inflammatory signals. In addition, autophagy in
macrophages and DCs suppresses pro-inflammatory responses independent of T_H17 and
inflammasome activation, including through mitophagy[88–91], clearance of damage-associated
molecular patterns (DAMPs) and PAMPs[92–94], ER-phagy[95,96], as well as through direct interaction

with MyD88[97,98]. These roles for autophagy could also impact neutrophil recruitment and
accumulation in the lungs during HDM challenge. While we ruled out a neutrophil intrinsic role
for ATG5 in our study using the *Atg5[fl/fl]-MRP8-Cre* mice, it is likely that the neutrophils that
accumulate in the lungs also exacerbate the T_H17 response and inflammation. For example,
neutrophil-derived antimicrobial peptide cathelicidin can induce T_H17 differentiation in a TGF-β
dependent manner in purified cells and induce IL-17 production in mice[99].

Loss of autophagy in lung macrophages and DCs also resulted in the accumulation of T_{regs}
in the lungs following HDM challenge. Although T_{regs} are well-known to suppress the
inflammatory responses during asthma[100–103], T_{reg} accumulation is also reported to be increased
during severe asthma[104,105]. Higher T_{reg} accumulation during severe asthma is associated with
higher T_H17 responses, where the transfer of purified T_H17 cells promoted the expansion of T_{regs}
in mice[106]. Therefore, it is possible that the increased T_H17 responses directly contribute to the
increase in T_{regs} during HDM challenge of *Atg5[fl/fl]-LysM-Cre* mice. Loss of autophagy in DCs is
associated with dysregulated T_{reg} function in mice due to an accumulation of ADAM10
metalloprotease on the surface of DCs in the absence of autophagy[107]. Accumulation of ADAM10
leads to cleavage of ICOS-L, which is required for the efficient maturation of T_{regs} in steady state.
Moreover, in the absence of proper maturation, $FOXP3^+ CD25^+$ T_{regs} transition to $IL-17^+ CD4^+$ T
cells[107], which may explain the increased IL-17 expression in T_{regs} that we observed in HDM-
challenged *Atg5[fl/fl]-LysM-Cre* mice.

Together our studies highlight a protective role for autophagy in controlling neutrophil
accumulation during asthma, suggesting that inducing autophagy may hold promise as a
therapeutic for treating severe asthma cases. Given that autophagy can perform different functions
in different cell types, this will need to be tested directly, which is currently limited by the absence
of autophagy-specific inducers. Nonetheless, these studies establish the role for autophagy in lung
macrophages and DCs during exposure to an allergen, which may also be applicable to other lung
inflammatory diseases.

METHODS

<u>Mouse strains</u>

All flox mice (*Atg5[fl/fl], Becn1[fl/fl], Atg16l1[fl/fl]*) used in this study have been described
previously[54,59,108,109] and are maintained in an enhanced barrier facility. LysM-Cre (Jax #004781),

CD11c-Cre (Jax #007567), and MRP8-Cre (Jax #021614) mice from the Jackson Laboratory were

crossed to specific flox mice, respectively. *Atg5*$^{fl/fl}$ and *Atg5*$^{fl/fl}$*-LysM-Cre* mice were crossed with

Casp1/11$^{-/-}$ (Jax #016621) germline deletion mice to generate *Atg5*$^{fl/fl}$*/Casp1/11*$^{-/-}$ and *Atg5*$^{fl/fl}$-

LysM-Cre/Casp1/11$^{-/-}$ mice. *Atg5*$^{fl/fl}$ and *Atg5*$^{fl/fl}$*-LysM-Cre* mice were crossed with *Il17a*^{eGFP} (Jax

#018472) mice to generate *Atg5*$^{fl/fl}$*/Il17a*^{eGFP} and *Atg5*$^{fl/fl}$*-LysM-Cre/Il17a*^{eGFP} mice. Adult male

and female mice (aged 8–12 weeks) were used for the experiments. All the experiments were

performed with respective littermate flox controls. All the mice have been backcrossed to

C57BL/6J background. The mice were housed and bred at Washington University in St. Louis in

specific pathogen-free conditions in accordance with federal and university guidelines. All

procedures involving animals were conducted following National Institute of Health guidelines

and protocols were approved by the Institutional Animal Care and Use Committee of Washington

University in St. Louis. Animals were euthanized by CO_2 asphyxiation, approved by Euthanasia

of the American Veterinary Association.

<u>Asthma model</u>

Mice were challenged with HDM as described previously[14]. Briefly, mice were challenged

intranasally with HDM in PBS (Greer Laboratories). The first primary dose of HDM (200 µg) was

administered on Day 1 and two secondary (100 µg) doses were administered on day 8 and day 15

respectively. Mice were sacrificed on day 16 and the whole lung except the post caudal lobe was

used in flow cytometry experiments.

<u>Flow cytometry</u>

Mouse lungs were excised after PBS perfusion, placed in DMEM containing 10% FBS, minced

finely, and digested at 37ºC for an hour with mechanical disruption with a stir bar and enzymatic

digestion with collagenase D (Roche 11088875103) and DNase I (Sigma D4527). Digested tissue

was treated with ACK buffer to remove red blood cells and were passed through a 70 μm cell

strainer to generate single cell suspension in 1 ml of semi-facs buffer (1X PBS pH 7.4

supplemented with 2% FBS and 2 mM EDTA). 100 μl of cells were stained with Zombie-NIR

before staining with the antibody cocktail mix. Cells were resuspended in semi-facs buffer in the

presence of Fc receptor blocking antibody (BioLegend, 101302) and stained with antibodies at a

1:200 dilution with the myeloid or lymphoid antibody cocktails. The innate immune cell staining

antibody cocktail included the following antibodies: Ly6G-BV421 (Biolegend, 1A8), SiglecF-BV480 (Biolegend, E50-2440(RUO), CD45-BV570 (Biolegend, 30-F11), CD11b-BV650 (Biolegend, M1/70), CD103-BV750 (Biolegend, M290), Ly6C-BV785 (Biolegend, HK1.4), MHCII-SparkBlue550 (Biolegend, M5/114.15.2), CD64-PerCP-eFluor710 (Biolegend, X54-5/7.1), MerTK-PE-Cy7 (Biolegend, DS5MMER), CD206-eF660 (AF647) (eBiosciences, C068C2), CD11c-APC/Fire750 (Biolegend, N418). The lymphoid cell staining antibody cocktail included the following antibodies: CD8α-Pacific blue (Biolegend, 53-6.7), CD44-BV510 (Biolegend, IM7), CD45-BV570 (Biolegend, 30-F11), CD19-BV605 (Biolegend, 6D5), CD11b-BV650 (Biolegend, M1/70), NK1.1-BV785 (Biolegend, PK136), TCRβ-FITC (Biolegend, H57-597), NKp46-PerCP-eFluor710 (eBiosciences, 29A1.4), CD62L-PE/Cy5 (Biolegend, MEL-14), CD25-PE-Cy7 (Biolegend, PC61), and CD4-APC (Biolegend, 100516). Flow cytometric analysis was performed on an Cytek Aurora and data was analyzed with FlowJo software (Tree Star). The gating strategies are shown in **Figure S2**. Cell populations are represented as counts per total lung suspension (1 ml) excluding the post-caudal lobe.

Statistical analysis for biological experiments

All data from HDM-challenged mice are from at least two independent experiments. Samples represent biological (not technical) replicates of mice randomly sorted into each experimental group. No blinding was performed during animal experiments. If presented on a log scale, the data was log transformed before statistical analysis. Statistical differences were calculated using Prism 9.0 (GraphPad Software) using unpaired Welch's t-tests or one-way ANOVA as denoted in the figure legends. Sample sizes were sufficient to detect differences as small as 10% using the statistical methods described. When used, center values and error bars represent means ± S.E.M. $P < 0.05$ was considered significant. $P < 0.05$ was denoted *, ** for $P < 0.01$, *** $P < 0.001$, and **** $P < 0.0001$. Comparisons that were not significantly different are designated as ns.

AUTHOR CONTRIBUTIONS

N.D. and C.L.S. designed the project, analyzed the data, and wrote the manuscript. N.D., S.R.M., N.R.E., and R.W. performed the experiments. N.D., A.R., A.L.K, and C.L.S. designed the asthma protocols. D.K. performed mouse crossing, breeding, and housing. A.L.K. and C.L.S. advised project design. All authors read and edited the manuscript.

ACKNOWLEDGEMENTS

This work was supported by NIH grants R01 AI132697 and U19 AI142784, Burroughs Wellcome Fund Investigators in the Pathogenesis of Infectious Disease, and the Philip and Sima Needleman Center for Autophagy Therapeutics and Research to C.L.S. N.D. is supported by Alexander and Gertrude Berg fellowship from the Department of Molecular Microbiology at Washington University in St. Louis School of Medicine. This material is based upon work supported by the National Science Foundation Graduate Research Fellowship Program under Grant #DGE 2139839 to N.R.E. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

DISCLOSURE STATEMENT

No potential conflict of interest was reported by the authors.

REFERENCES

1. GBD 2019 Diseases and Injuries Collaborators. Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. Lancet 2020; 396:1204–22.

2. Papi A, Brightling C, Pedersen SE, Reddel HK. Asthma. Lancet 2018; 391:783–800.

3. Platts-Mills TAE, Woodfolk JA. Allergens and their role in the allergic immune response. Immunol Rev 2011; 242:51–68.

4. Walford HH, Doherty TA. Diagnosis and management of eosinophilic asthma: a US perspective. J Asthma Allergy 2014; 7:53–65.

5. Lambrecht BN, Hammad H, Fahy JV. The Cytokines of Asthma. Immunity 2019; 50:975–91.

6. Bartemes KR, Kita H. Dynamic role of epithelium-derived cytokines in asthma. Clin Immunol 2012; 143:222–35.

7. Haldar P, Pavord ID. Noneosinophilic asthma: a distinct clinical and pathologic phenotype. J Allergy Clin Immunol 2007; 119:1043–52; quiz 1053–4.

8. Moore WC, Meyers DA, Wenzel SE, Teague WG, Li H, Li X, D'Agostino R, Castro M, Curran-Everett D, Fitzpatrick AM, et al. Identification of asthma phenotypes using cluster

analysis in the Severe Asthma Research Program. Am J Respir Crit Care Med 2010;
181:315–23.

9. Robinson D, Humbert M, Buhl R, Cruz AA, Inoue H, Korom S, Hanania NA, Nair P.
Revisiting Type 2-high and Type 2-low airway inflammation in asthma: current knowledge
and therapeutic implications. Clin Exp Allergy 2017; 47:161–75.

10. Ray A, Kolls JK. Neutrophilic Inflammation in Asthma and Association with Disease
Severity. Trends Immunol 2017; 38:942–54.

11. Zhang X, Xu Z, Wen X, Huang G, Nian S, Li L, Guo X, Ye Y, Yuan Q. The onset,
development and pathogenesis of severe neutrophilic asthma. Immunol Cell Biol 2022;
100:144–59.

12. Newcomb DC, Peebles RS. Th17-mediated inflammation in asthma. Curr Opin Immunol
2013; 25:755–60.

13. McKinley L, Alcorn JF, Peterson A, Dupont RB, Kapadia S, Logar A, Henry A, Irvin CG,
Piganelli JD, Ray A, et al. TH17 cells mediate steroid-resistant airway inflammation and
airway hyperresponsiveness in mice. J Immunol 2008; 181:4089–97.

14. Suzuki Y, Maazi H, Sankaranarayanan I, Lam J, Khoo B, Soroosh P, Barbers RG, James
Ou J-H, Jung JU, Akbari O. Lack of autophagy induces steroid-resistant airway
inflammation. J Allergy Clin Immunol 2016; 137:1382-1389.e9.

15. Raundhal M, Morse C, Khare A, Oriss TB, Milosevic J, Trudeau J, Huff R, Pilewski J,
Holguin F, Kolls J, et al. High IFN-γ and low SLPI mark severe asthma in mice and
humans. J Clin Invest 2015; 125:3037–50.

16. Wisniewski JA, Muehling LM, Eccles JD, Capaldo BJ, Agrawal R, Shirley D-A, Patrie JT,
Workman LJ, Schuyler AJ, Lawrence MG, et al. TH1 signatures are present in the lower
airways of children with severe asthma, regardless of allergic status. J Allergy Clin
Immunol 2018; 141:2048-2060.e13.

17. Al-Ramli W, Préfontaine D, Chouiali F, Martin JG, Olivenstein R, Lemière C, Hamid Q.
T(H)17-associated cytokines (IL-17A and IL-17F) in severe asthma. J Allergy Clin
Immunol 2009; 123:1185–7.

18. Cho K-A, Suh JW, Sohn JH, Park JW, Lee H, Kang JL, Woo S-Y, Cho YJ. IL-33 induces
Th17-mediated airway inflammation via mast cells in ovalbumin-challenged mice. Am J
Physiol Lung Cell Mol Physiol 2012; 302:L429-440.

19. Whetstone CE, Ranjbar M, Omer H, Cusack RP, Gauvreau GM. The Role of Airway
Epithelial Cell Alarmins in Asthma. Cells 2022; 11:1105.

20. Trompette A, Divanovic S, Visintin A, Blanchard C, Hegde RS, Madan R, Thorne PS,
Wills-Karp M, Gioannini TL, Weiss JP, et al. Allergenicity resulting from functional
mimicry of a Toll-like receptor complex protein. Nature 2009; 457:585–8.

21. Wilson RH, Maruoka S, Whitehead GS, Foley JF, Flake GP, Sever ML, Zeldin DC, Kraft
M, Garantziotis S, Nakano H, et al. The Toll-like receptor 5 ligand flagellin promotes
asthma by priming allergic responses to indoor allergens. Nat Med 2012; 18:1705–10.

22. Redecke V, Häcker H, Datta SK, Fermin A, Pitha PM, Broide DH, Raz E. Cutting edge:
activation of Toll-like receptor 2 induces a Th2 immune response and promotes
experimental asthma. J Immunol 2004; 172:2739–43.

23. Willart MAM, Deswarte K, Pouliot P, Braun H, Beyaert R, Lambrecht BN, Hammad H.
Interleukin-1α controls allergic sensitization to inhaled house dust mite via the epithelial
release of GM-CSF and IL-33. J Exp Med 2012; 209:1505–17.

24. Cheng S, Chen H, Wang A, Bunjhoo H, Cao Y, Xie J, Xu Y, Xiong W. Blockade of IL-23
ameliorates allergic lung inflammation via decreasing the infiltration of Tc17 cells. Arch
Med Sci 2016; 12:1362–9.

25. Li Y, Hua S. Mechanisms of pathogenesis in allergic asthma: role of interleukin-23.
Respirology 2014; 19:663–9.

26. McGeachy MJ, Chen Y, Tato CM, Laurence A, Joyce-Shaikh B, Blumenschein WM,
McClanahan TK, O'Shea JJ, Cua DJ. The interleukin 23 receptor is essential for the
terminal differentiation of interleukin 17-producing effector T helper cells in vivo. Nat
Immunol 2009; 10:314–24.

27. Jones CE, Chan K. Interleukin-17 stimulates the expression of interleukin-8, growth-related
oncogene-alpha, and granulocyte-colony-stimulating factor by human airway epithelial
cells. Am J Respir Cell Mol Biol 2002; 26:748–53.

28. Laan M, Cui ZH, Hoshino H, Lötvall J, Sjöstrand M, Gruenert DC, Skoogh BE, Lindén A.
Neutrophil recruitment by human IL-17 via C-X-C chemokine release in the airways. J
Immunol 1999; 162:2347–52.

29. Souwer Y, Groot Kormelink T, Taanman-Kueter EW, Muller FJ, van Capel TMM, Varga
DV, Bar-Ephraim YE, Teunissen MBM, van Ham SM, Kuijpers TW, et al. Human
TH17 cell development requires processing of dendritic cell-derived CXCL8 by neutrophil
elastase. J Allergy Clin Immunol 2018; 141:2286-2289.e5.

30. Chenuet P, Fauconnier L, Madouri F, Marchiol T, Rouxel N, Ledru A, Mauny P, Lory R,
Uttenhove C, van Snick J, et al. Neutralization of either IL-17A or IL-17F is sufficient to
inhibit house dust mite induced allergic asthma in mice. Clin Sci (Lond) 2017; 131:2533–
48.

31. Manni ML, Trudeau JB, Scheller EV, Mandalapu S, Elloso MM, Kolls JK, Wenzel SE,
Alcorn JF. The complex relationship between inflammation and lung function in severe
asthma. Mucosal Immunol 2014; 7:1186–98.

32. Nabe T, Hosokawa F, Matsuya K, Morishita T, Ikedo A, Fujii M, Mizutani N, Yoshino S,
Chaplin DD. Important role of neutrophils in the late asthmatic response in mice. Life Sci
2011; 88:1127–35.

33. Poon AH, Chouiali F, Tse SM, Litonjua AA, Hussain SNA, Baglole CJ, Eidelman DH,
Olivenstein R, Martin JG, Weiss ST, et al. Genetic and histologic evidence for autophagy in
asthma pathogenesis. J Allergy Clin Immunol 2012; 129:569–71.

34. Martin LJ, Gupta J, Jyothula SSSK, Butsch Kovacic M, Biagini Myers JM, Patterson TL,
Ericksen MB, He H, Gibson AM, Baye TM, et al. Functional variant in the autophagy-
related 5 gene promotor is associated with childhood asthma. PLoS One 2012; 7:e33454.

35. Pham DL, Kim S-H, Losol P, Yang E-M, Shin YS, Ye Y-M, Park H-S. Association of
autophagy related gene polymorphisms with neutrophilic airway inflammation in adult
asthma. Korean J Intern Med 2016; 31:375–85.

36. Kimmey JM, Huynh JP, Weiss LA, Park S, Kambal A, Debnath J, Virgin HW, Stallings
CL. Unique role for ATG5 in neutrophil-mediated immunopathology during M.
tuberculosis infection. Nature 2015; 528:565–9.

37. Kinsella RL, Nehls EM, Stallings CL. Roles for Autophagy Proteins in Immunity and Host
Defense. Vet Pathol 2018; 55:366–73.

38. Kimmey JM, Stallings CL. Bacterial Pathogens versus Autophagy: Implications for
Therapeutic Interventions. Trends Mol Med 2016; 22:1060–76.

39. Lamb CA, Yoshimori T, Tooze SA. The autophagosome: origins unknown, biogenesis
complex. Nat Rev Mol Cell Biol 2013; 14:759–74.

40. Mizushima N, Noda T, Yoshimori T, Tanaka Y, Ishii T, George MD, Klionsky DJ, Ohsumi
M, Ohsumi Y. A protein conjugation system essential for autophagy. Nature 1998;
395:395–8.

41. Mizushima N, Yoshimori T, Ohsumi Y. The role of Atg proteins in autophagosome
formation. Annu Rev Cell Dev Biol 2011; 27:107–32.

42. Li BB, Chen YL, Pang F. MicroRNA-30a Targets ATG5 and Attenuates Airway Fibrosis in
Asthma by Suppressing Autophagy. Inflammation 2020; 43:44–53.

43. McAlinden KD, Deshpande DA, Ghavami S, Xenaki D, Sohal SS, Oliver BG, Haghi M,
Sharma P. Autophagy Activation in Asthma Airways Remodeling. Am J Respir Cell Mol
Biol 2019; 60:541–53.

44. Poon AH, Choy DF, Chouiali F, Ramakrishnan RK, Mahboub B, Audusseau S, Mogas A,
Harris JM, Arron JR, Laprise C, et al. Increased Autophagy-Related 5 Gene Expression Is
Associated with Collagen Expression in the Airways of Refractory Asthmatics. Front
Immunol 2017; 8:355.

45. Ban G-Y, Pham DL, Trinh THK, Lee S-I, Suh D-H, Yang E-M, Ye Y-M, Shin YS, Chwae
Y-J, Park H-S. Autophagy mechanisms in sputum and peripheral blood cells of patients
with severe asthma: a new therapeutic target. Clin Exp Allergy 2016; 46:48–59.

46. Theofani E, Xanthou G. Autophagy: A Friend or Foe in Allergic Asthma? Int J Mol Sci
2021; 22:6314.

47. Liu J-N, Suh D-H, Trinh HKT, Chwae Y-J, Park H-S, Shin YS. The role of autophagy in
allergic inflammation: a new target for severe asthma. Exp Mol Med 2016; 48:e243.

48. Schrezenmeier E, Dörner T. Mechanisms of action of hydroxychloroquine and chloroquine:
implications for rheumatology. Nat Rev Rheumatol 2020; 16:155–66.

49. Das M, Ram A, Ghosh B. Luteolin alleviates bronchoconstriction and airway
hyperreactivity in ovalbumin sensitized mice. Inflamm Res 2003; 52:101–6.

50. Kim S-H, Saba E, Kim B-K, Yang W-K, Park Y-C, Shin HJ, Han CK, Lee YC, Rhee MH.
Luteolin attenuates airway inflammation by inducing the transition of CD4+CD25- to
CD4+CD25+ regulatory T cells. Eur J Pharmacol 2018; 820:53–64.

51. Wang S, Wuniqiemu T, Tang W, Teng F, Bian Q, Yi L, Qin J, Zhu X, Wei Y, Dong J.
Luteolin inhibits autophagy in allergic asthma by activating PI3K/Akt/mTOR signaling and
inhibiting Beclin-1-PI3KC3 complex. Int Immunopharmacol 2021; 94:107460.

52. Gu W, Cui R, Ding T, Li X, Peng J, Xu W, Han F, Guo X. Simvastatin alleviates airway
inflammation and remodelling through up-regulation of autophagy in mouse models of
asthma. Respirology 2017; 22:533–41.

53. McKay A, Leung BP, McInnes IB, Thomson NC, Liew FY. A novel anti-inflammatory role
of simvastatin in a murine model of allergic asthma. J Immunol 2004; 172:2903–8.

54. Wang Y-T, Zaitsev K, Lu Q, Li S, Schaiff WT, Kim K-W, Droit L, Wilen CB, Desai C,
Balce DR, et al. Author Correction: Select autophagy genes maintain quiescence of tissue-
resident macrophages and increase susceptibility to Listeria monocytogenes. Nat Microbiol
2021; 6:972.

55. Lu Q, Yokoyama CC, Williams JW, Baldridge MT, Jin X, DesRochers B, Bricker T, Wilen
CB, Bagaitkar J, Loginicheva E, et al. Homeostatic Control of Innate Lung Inflammation
by Vici Syndrome Gene Epg5 and Additional Autophagy Genes Promotes Influenza
Pathogenesis. Cell Host Microbe 2016; 19:102–13.

56. Zhao Z, Fux B, Goodwin M, Dunay IR, Strong D, Miller BC, Cadwell K, Delgado MA,
Ponpuak M, Green KG, et al. Autophagosome-independent essential function for the
autophagy protein Atg5 in cellular immunity to intracellular pathogens. Cell Host Microbe
2008; 4:458–69.

57. Galluzzi L, Green DR. Autophagy-Independent Functions of the Autophagy Machinery.
Cell 2019; 177:1682–99.

58. Choi J, Park S, Biering SB, Selleck E, Liu CY, Zhang X, Fujita N, Saitoh T, Akira S,
Yoshimori T, et al. The parasitophorous vacuole membrane of Toxoplasma gondii is
targeted for disruption by ubiquitin-like conjugation systems of autophagy. Immunity 2014;
40:924–35.

59. Hara T, Nakamura K, Matsui M, Yamamoto A, Nakahara Y, Suzuki-Migishima R,
Yokoyama M, Mishima K, Saito I, Okano H, et al. Suppression of basal autophagy in
neural cells causes neurodegenerative disease in mice. Nature 2006; 441:885–9.

60. Li L, Huang L, Vergis AL, Ye H, Bajwa A, Narayan V, Strieter RM, Rosin DL, Okusa MD.
IL-17 produced by neutrophils regulates IFN-gamma-mediated neutrophil migration in
mouse kidney ischemia-reperfusion injury. J Clin Invest 2010; 120:331–42.

61. Shi Y, Shi G, Wan H, Jiang L, Ai X, Zhu H, Tang W, Ma J, Jin X, Zhang B. Coexistence of
Th1/Th2 and Th17/Treg imbalances in patients with allergic asthma. Chin Med J (Engl)
2011; 124:1951–6.

62. Zheng R, Wang F, Huang Y, Xiang Q, Dai H, Zhang W. Elevated Th17 cell frequencies
and Th17/Treg ratio are associated with airway hyperresponsiveness in asthmatic children.
J Asthma 2021; 58:707–16.

63. Huang F, Yin J-N, Wang H-B, Liu S-Y, Li Y-N. Association of imbalance of effector T
cells and regulatory cells with the severity of asthma and allergic rhinitis in children.
Allergy Asthma Proc 2017; 38:70–7.

64. Zhu J, Liu X, Wang W, Ouyang X, Zheng W, Wang Q. Altered expression of regulatory T
and Th17 cells in murine bronchial asthma. Exp Ther Med 2017; 14:714–22.

65. Zou X-L, Chen Z-G, Zhang T-T, Feng D-Y, Li H-T, Yang H-L. Th17/Treg homeostasis,
but not Th1/Th2 homeostasis, is implicated in exacerbation of human bronchial asthma.
Ther Clin Risk Manag 2018; 14:1627–36.

66. Liu K, Zhao E, Ilyas G, Lalazar G, Lin Y, Haseeb M, Tanaka KE, Czaja MJ. Impaired
macrophage autophagy increases the immune response in obese mice by promoting
proinflammatory macrophage polarization. Autophagy 2015; 11:271–84.

67. Ye X, Zhou X-J, Zhang H. Exploring the Role of Autophagy-Related Gene 5 (ATG5)
Yields Important Insights Into Autophagy in Autoimmune/Autoinflammatory Diseases.
Front Immunol 2018; 9:2334.

68. Theofani E, Semitekolou M, Samitas K, Mais A, Galani IE, Triantafyllia V, Lama J,
Morianos I, Stavropoulos A, Jeong S-J, et al. TFEB signaling attenuates NLRP3-driven
inflammatory responses in severe asthma. Allergy 2022; 77:2131–46.

69. Biasizzo M, Kopitar-Jerala N. Interplay Between NLRP3 Inflammasome and Autophagy.
Front Immunol 2020; 11:591803.

70. Sun Q, Fan J, Billiar TR, Scott MJ. Inflammasome and autophagy regulation - a two-way
street. Mol Med 2017; 23:188–95.

71. Kim RY, Pinkerton JW, Essilfie AT, Robertson AAB, Baines KJ, Brown AC, Mayall JR,
Ali MK, Starkey MR, Hansbro NG, et al. Role for NLRP3 Inflammasome-mediated, IL-1β-
Dependent Responses in Severe, Steroid-Resistant Asthma. Am J Respir Crit Care Med
2017; 196:283–97.

72. Ma M, Li G, Qi M, Jiang W, Zhou R. Inhibition of the Inflammasome Activity of NLRP3
Attenuates HDM-Induced Allergic Asthma. Front Immunol 2021; 12:718779.

73. Simpson JL, Phipps S, Baines KJ, Oreo KM, Gunawardhana L, Gibson PG. Elevated
expression of the NLRP3 inflammasome in neutrophilic asthma. Eur Respir J 2014;
43:1067–76.

74. Theofani E, Semitekolou M, Morianos I, Samitas K, Xanthou G. Targeting NLRP3
Inflammasome Activation in Severe Asthma. J Clin Med 2019; 8:E1615.

75. Rossios C, Pavlidis S, Hoda U, Kuo C-H, Wiegman C, Russell K, Sun K, Loza MJ,
Baribaud F, Durham AL, et al. Sputum transcriptomics reveal upregulation of IL-1 receptor
family members in patients with severe asthma. J Allergy Clin Immunol 2018; 141:560–70.

76. Inoue D, Kubo H, Taguchi K, Suzuki T, Komatsu M, Motohashi H, Yamamoto M.
Inducible disruption of autophagy in the lung causes airway hyper-responsiveness.
Biochem Biophys Res Commun 2011; 405:13–8.

77. Liu MC, Xiao HQ, Breslin LM, Bochner BS, Schroeder JT. Enhanced antigen presenting
and T cell functions during late-phase allergic responses in the lung. Clin Exp Allergy
2018; 48:334–42.

78. VanHook AM. Macrophages present antigens in allergic asthma. Sci Signal 2022;
15:adf2043.

79. Zhu X, Cui J, Yi L, Qin J, Tulake W, Teng F, Tang W, Wei Y, Dong J. The Role of T Cells
and Macrophages in Asthma Pathogenesis: A New Perspective on Mutual Crosstalk.
Mediators Inflamm 2020; 2020:7835284.

80. Ghislat G, Lawrence T. Autophagy in dendritic cells. Cell Mol Immunol 2018; 15:944–52.

81. Lee HK, Mattei LM, Steinberg BE, Alberts P, Lee YH, Chervonsky A, Mizushima N,
Grinstein S, Iwasaki A. In vivo requirement for Atg5 in antigen presentation by dendritic
cells. Immunity 2010; 32:227–39.

82. Peral de Castro C, Jones SA, Ní Cheallaigh C, Hearnden CA, Williams L, Winter J, Lavelle
EC, Mills KHG, Harris J. Autophagy regulates IL-23 secretion and innate T cell responses
through effects on IL-1 secretion. J Immunol 2012; 189:4144–53.

83.   Aggarwal S, Ghilardi N, Xie M-H, de Sauvage FJ, Gurney AL. Interleukin-23 promotes a
647   distinct CD4 T cell activation state characterized by the production of interleukin-17. J Biol
648   Chem 2003; 278:1910–4.

649   84.   Sinclair C, Bommakanti G, Gardinassi L, Loebbermann J, Johnson MJ, Hakimpour P,
650   Hagan T, Benitez L, Todor A, Machiah D, et al. mTOR regulates metabolic adaptation of
651   APCs in the lung and controls the outcome of allergic inflammation. Science 2017;
652   357:1014–21.

653   85.   Krausgruber T, Blazek K, Smallie T, Alzabin S, Lockstone H, Sahgal N, Hussell T,
654   Feldmann M, Udalova IA. IRF5 promotes inflammatory macrophage polarization and TH1-
655   TH17 responses. Nat Immunol 2011; 12:231–8.

656   86.   Oriss TB, Raundhal M, Morse C, Huff RE, Das S, Hannum R, Gauthier MC, Scholl KL,
657   Chakraborty K, Nouraie SM, et al. IRF5 distinguishes severe asthma in humans and drives
658   Th1 phenotype and airway hyperreactivity in mice. JCI Insight 2017; 2:e91019, 91019.

659   87.   Liu R, Cui J, Sun Y, Xu W, Wang Z, Wu M, Dong H, Yang C, Hong S, Yin S, et al.
660   Autophagy deficiency promotes M1 macrophage polarization to exacerbate acute liver
661   injury via ATG5 repression during aging. Cell Death Discov 2021; 7:397.

662   88.   Harris J, Deen N, Zamani S, Hasnat MA. Mitophagy and the release of inflammatory
663   cytokines. Mitochondrion 2018; 41:2–8.

664   89.   Lazarou M, Sliter DA, Kane LA, Sarraf SA, Wang C, Burman JL, Sideris DP, Fogel AI,
665   Youle RJ. The ubiquitin kinase PINK1 recruits autophagy receptors to induce mitophagy.
666   Nature 2015; 524:309–14.

667   90.   Song Y, Zhou Y, Zhou X. The role of mitophagy in innate immune responses triggered by
668   mitochondrial stress. Cell Commun Signal 2020; 18:186.

669   91.   Killackey SA, Philpott DJ, Girardin SE. Mitophagy pathways in health and disease. J Cell
670   Biol 2020; 219:e202004029.

671   92.   Pouwels SD, Heijink IH, van Oosterhout AJM, Nawijn MC. A specific DAMP profile
672   identifies susceptibility to smoke-induced airway inflammation. Eur Respir J 2014;
673   43:1183–6.

674   93.   Tang D, Kang R, Coyne CB, Zeh HJ, Lotze MT. PAMPs and DAMPs: signal 0s that spur
675   autophagy and immunity. Immunol Rev 2012; 249:158–75.

676   94.   Huang X, Tan X, Liang Y, Hou C, Qu D, Li M, Huang Q. Differential DAMP release was
677   observed in the sputum of COPD, asthma and asthma-COPD overlap (ACO) patients. Sci
678   Rep 2019; 9:19241.

679   95.   Chipurupalli S, Samavedam U, Robinson N. Crosstalk Between ER Stress, Autophagy and
680   Inflammation. Front Med (Lausanne) 2021; 8:758311.

96. Moretti J, Roy S, Bozec D, Martinez J, Chapman JR, Ueberheide B, Lamming DW, Chen
ZJ, Horng T, Yeretssian G, et al. STING Senses Microbial Viability to Orchestrate Stress-
Mediated Autophagy of the Endoplasmic Reticulum. Cell 2017; 171:809-823.e13.

97. Peng X, Wang Y, Li H, Fan J, Shen J, Yu X, Zhou Y, Mao H. ATG5-mediated autophagy
suppresses NF-κB signaling to limit epithelial inflammatory response to kidney injury. Cell
Death Dis 2019; 10:253.

98. Inomata M, Into T, Niida S, Murakami Y. Atg5 regulates formation of MyD88 condensed
structures and MyD88-dependent signal transduction. Biochem Biophys Res Commun
2013; 437:509–14.

99. Minns D, Smith KJ, Alessandrini V, Hardisty G, Melrose L, Jackson-Jones L, MacDonald
AS, Davidson DJ, Gwyer Findlay E. The neutrophil antimicrobial peptide cathelicidin
promotes Th17 differentiation. Nat Commun 2021; 12:1285.

100. Strickland DH, Stumbles PA, Zosky GR, Subrata LS, Thomas JA, Turner DJ, Sly PD, Holt
PG. Reversal of airway hyperresponsiveness by induction of airway mucosal CD4+CD25+
regulatory T cells. J Exp Med 2006; 203:2649–60.

101. Wilson MS, Pesce JT, Ramalingam TR, Thompson RW, Cheever A, Wynn TA.
Suppression of murine allergic airway disease by IL-2:anti-IL-2 monoclonal antibody-
induced regulatory T cells. J Immunol 2008; 181:6942–54.

102. Dembele M, Tao S, Massoud AH, Miah SMS, Lelias S, De Groot AS, Mazer BD.
Tregitopes Improve Asthma by Promoting Highly Suppressive and Antigen-Specific Tregs.
Front Immunol 2021; 12:634509.

103. Kearley J, Barker JE, Robinson DS, Lloyd CM. Resolution of airway inflammation and
hyperreactivity after in vivo transfer of CD4+CD25+ regulatory T cells is interleukin 10
dependent. J Exp Med 2005; 202:1539–47.

104. Smyth LJC, Eustace A, Kolsum U, Blaikely J, Singh D. Increased airway T regulatory cells
in asthmatic subjects. Chest 2010; 138:905–12.

105. Lee J-H, Yu H-H, Wang L-C, Yang Y-H, Lin Y-T, Chiang B-L. The levels of CD4+CD25+
regulatory T cells in paediatric patients with allergic rhinitis and bronchial asthma. Clin Exp
Immunol 2007; 148:53–63.

106. Zhou Q, Hu Y, Howard OMZ, Oppenheim JJ, Chen X. In vitro generated Th17 cells
support the expansion and phenotypic stability of CD4(+)Foxp3(+) regulatory T cells in
vivo. Cytokine 2014; 65:56–64.

107. Niven J, Madelon N, Page N, Caruso A, Harlé G, Lemeille S, Seemayer CA, Hugues S,
Gannagé M. Macroautophagy in Dendritic Cells Controls the Homeostasis and Stability of
Regulatory T Cells. Cell Rep 2019; 28:21-29.e6.

108. Hwang S, Maloney NS, Bruinsma MW, Goel G, Duan E, Zhang L, Shrestha B, Diamond
MS, Dani A, Sosnovtsev SV, et al. Nondegradative role of Atg5-Atg12/ Atg16L1
autophagy protein complex in antiviral activity of interferon gamma. Cell Host Microbe
2012; 11:397–409.

109. Gawriluk TR, Hale AN, Flaws JA, Dillon CP, Green DR, Rucker EB. Autophagy is a cell
survival program for female germ cells in the murine ovary. Reproduction 2011; 141:759–
65.

FIGURE LEGENDS

Figure 1. Loss of *Atg5* expression in LysM$^+$ cells results in increased neutrophil-dominated

inflammation upon HDM challenge

(A) Schematic of asthma model used in study, made in part with BioRender. Mice were challenged

with house dust mites intranasally. **(B)** Representative flow gating of neutrophils (left) and graph

of neutrophil numbers per lung (right) from *Atg5$^{fl/fl}$* and *Atg5$^{fl/fl}$-LysM-Cre* mice either before HDM

challenge (unchallenged) or post-HDM challenge. Number of **(C)** non-alveolar macrophages,

eosinophils, DCs, inflammatory monocytes, **(D)** CD4$^+$ T cells, **(E)** activated CD4$^+$ T cells, naïve

CD4$^+$ T cells, and **(F)** T$_{reg}$ cells per lung. The legend for **(B-F)** is shown in **(B)**. **(B-F)** Each

datapoint is from an individual mouse and the HDM samples are compiled from 2 independent

experiments. The bar graphs represent the mean +/- S.E.M. One-way ANOVA was used to

determine the statistical significance of differences between the groups, where only the differences

within a single condition are shown. * P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001.

Comparisons that were not significantly different are designated as ns.

Figure 2. Loss of *Atg5* in innate immune cells results in increased production of IL-17 from

CD4$^+$ T cells

(A-B) Number of **(A)** neutrophils and **(B)** activated CD4$^+$ T cells in *Atg5$^{fl/fl}$/Il17a^{eGFP}* (n=5) and

Atg5$^{fl/fl}$-LysM-Cre/Il17a^{eGFP} (n=10) mice following HDM challenge. **(C)** Percent or **(D)** number of

CD4$^+$ T cells, activated CD4$^+$ T cells, and T$_{regs}$ expressing IL-17 based on GFP expression in

HDM-challenged *Atg5$^{fl/fl}$/Il17a^{eGFP}* and *Atg5$^{fl/fl}$-LysM-Cre/Il17a^{eGFP}* mice. **(E)** Mean fluorescence

intensity for GFP (IL-17) in CD4$^+$ T cells, activated CD4$^+$ T cells, and T$_{regs}$. **(F)** Ratio of IL-17$^+$

CD4$^+$ T cell numbers to total T$_{reg}$ cell numbers. **(A-F)** The legend for all panels is shown at the top

of the figure. Each datapoint is from an individual mouse from at least 2 independent experiments.

The bar graphs represent the mean +/- S.E.M. Unpaired Welch's t-test was used to determine the

statistical significance of differences between the groups. * P < 0.05 and ** P < 0.01. Comparisons

that were not significantly different are designated as ns.

Figure 3. Autophagy is required in innate immune cells to control neutrophil-dominated

inflammation during HDM-challenge

Number of **(A)** neutrophils, non-alveolar macrophages, eosinophils, DCs, inflammatory

monocytes, **(B)** CD4$^+$ T cells, **(C)** activated CD4$^+$ T cells, naïve CD4$^+$ T cells, and **(D)** T$_{reg}$ cells

per lung in HDM-challenged *Atg16$^{fl/fl}$* (n=4), *Atg16l1$^{fl/fl}$-LysM-Cre* (n=5), *Becn1$^{fl/fl}$* (n=7), and

Becn1$^{fl/fl}$-LysM-Cre (n=10) mice. The legend for **(A-D)** is shown in **(A)**. Each datapoint is from an

individual mouse from at least 2 independent experiments. The bar graphs represent the mean +/-

S.E.M. Unpaired Welch's t-test was used to determine the statistical significance of differences

between the groups. * P < 0.05. Comparisons that were not significantly different are designated

as ns.

Figure 4. Autophagy regulates neutrophil accumulation during HDM challenge independent

of inflammasome activation

Number of **(A)** neutrophils, non-alveolar macrophages, eosinophils, DCs, inflammatory

monocytes, **(B)** CD4$^+$ T cells, **(C)** activated CD4$^+$ T cells, naïve CD4$^+$ T cells, and **(D)** T$_{reg}$ cells

per lung in HDM-challenged *Atg5$^{fl/fl}$/Casp1/11$^{-/-}$* (n=13) and *Atg5$^{fl/fl}$-LysM-Cre/Casp1/11$^{-/-}$* (n=17)

mice. The legend for **(B-D)** is shown above **(A)**. Each datapoint is from an individual mouse from

at least 2 independent experiments. The bar graphs represent the mean +/- S.E.M. Unpaired

Welch's t-test was used to determine the statistical significance of differences between the groups.

* P < 0.05, ** P < 0.01, *** P < 0.001. Comparisons that were not significantly different are

designated as ns.

Figure 5. Autophagy is required in CD11c$^+$ lung macrophages and DCs to control neutrophil

recruitment during exposure to HDM

(A-D) Number of **(A)** neutrophils, non-alveolar macrophages, eosinophils, DCs, inflammatory

monocytes, **(B)** CD4$^+$ T cells, **(C)** activated CD4$^+$ T cells, naïve CD4$^+$ T cells, and **(D)** T$_{reg}$ cells

per lung from HDM-challenged *Atg5$^{fl/fl}$* (n=10) and *Atg5$^{fl/fl}$-CD11c-Cre* (n=12) mice. The legend

above **(A)** is shared for **(A-D)**. **(E-H)** Number of **(E)** neutrophils, non-alveolar macrophages,

eosinophils, DCs, inflammatory monocytes, **(F)** CD4$^+$ T cells, **(G)** activated CD4$^+$ T cells, naïve

CD4$^+$ T cells, and **(H)** T$_{reg}$ cells per lung in HDM-challenged *Atg5$^{fl/fl}$* (n=6) and *Atg5$^{fl/fl}$-MRP8-Cre*

(n=7) mice. The legend above **(E)** is shared for **(E-H)**. Each datapoint is from an individual mouse

from at least 2 independent experiments. The bar graphs represent the mean +/- S.E.M. Unpaired

Welch's t-test was used to determine the statistical significance of differences between the groups.

* $P < 0.05$ and ** $P < 0.01$. Comparisons that were not significantly different are designated as ns.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5