1

A mechanistic model of primer synthesis from catalytic structures of DNA polymerase α–primase

Elwood A. Mullins[1,2], Lauren E. Salay[2,3,4], Clarissa L. Durie[5,6,7], Noah P. Bradley[1,2,8], Jane E. Jackman[9,10], Melanie D. Ohi[5,6], Walter J. Chazin[2,3,11,*], and Brandt F. Eichman[1,2,3,*]

[1]Department of Biological Sciences, Vanderbilt University, Nashville, Tennessee, USA

[2]Center for Structural Biology, Vanderbilt University, Nashville, Tennessee, USA

[3]Department of Biochemistry, Vanderbilt University School of Medicine, Nashville, Tennessee, USA

[4]Present address: Department of Biochemistry, University of Washington, Seattle, Washington, USA

[5]Department of Cell and Developmental Biology, University of Michigan School of Medicine, Ann Arbor, Michigan, USA

[6]Life Sciences Institute, University of Michigan, Ann Arbor, Michigan, USA

[7]Present address: Department of Biochemistry, University of Missouri, Columbia, Missouri, USA

[8]Presesnt address: Department of Molecular Biosciences, Northwestern University, Evanston, Illinois, USA

[9]Department of Chemistry and Biochemistry, The Ohio State University, Columbus, Ohio, USA

[10]Center for RNA Biology, The Ohio State University, Columbus, Ohio, USA

[11]Department of Chemistry, Vanderbilt University, Nashville, Tennessee, USA

*Address correspondence to these authors: walter.j.chazin@vanderbilt.edu, brandt.eichman@vanderbilt.edu

Abstract

The mechanism by which polymerase α–primase (polα–primase) synthesizes chimeric RNA-DNA primers of defined length and composition, necessary for replication fidelity and genome stability, is unknown. Here, we report cryo-EM structures of polα–primase in complex with primed templates representing various stages of DNA synthesis. Our data show how interaction of the primase regulatory subunit with the primer 5′-end facilitates handoff of the primer to polα and increases polα processivity, thereby regulating both RNA and DNA composition. The structures detail how flexibility within the heterotetramer enables synthesis across two active sites and provide evidence that termination of DNA synthesis is facilitated by reduction of polα and primase affinities for the varied conformations along the chimeric primer/template duplex. Together, these findings elucidate a critical catalytic step in replication initiation and provide a comprehensive model for primer synthesis by polα–primase.

Introduction

DNA polymerase α–primase (polα–primase) is unique among the replicative polymerases by virtue of its ability to perform *de novo* synthesis from a DNA template. Polα–primase initiates DNA synthesis by generating short (~30-nucleotide) primers [1-3], which are required for leading and lagging strand replication by polε and polδ [4]. Generation of primers is particularly important on the lagging strand, where discontinuous DNA synthesis is repeatedly initiated during genome replication and telomere elongation [5]. As such, polα–primase is essential for replication of the genome, and defects in priming have been implicated in cellular ageing and an impaired DNA damage response [6-10].

Polα–primase possesses both DNA-dependent RNA polymerase (primase) and DNA polymerase (polα) activities. Primase initiates synthesis from nucleoside triphosphates (NTPs) to generate a 7–10-ribonucleotide primer [11,12], which is extended by ~20 deoxyribonucleotides by polα (**Fig. 1A**) [1,2]. The resulting chimeric RNA-DNA primers are extended by polδ to generate each Okazaki fragment. Polα lacks proofreading capability and is therefore mutagenic [13]. The potentially deleterious effects of polα generated errors and incorporation of large amounts of RNA into the nuclear genome by primase are avoided by enzymatic removal of primers during Okazaki fragment maturation [14-19]. Thus, the composition and short length of the primer synthesized by polα–primase is an important factor in genomic integrity.

Polα–primase is a heterotetramer composed of two polymerase subunits—catalytic POLA1 (also referred to as p180) and regulatory POLA2 (p70, p68)—and two primase subunits—catalytic PRIM1 (p49, p48) and regulatory PRIM2 (p58) (**Fig. 1B**) [20]. The primase dimer, which is necessary for normal RNA synthesis [21], is formed by interactions between PRIM1 and the N-terminal domain of PRIM2 ($PRIM2_N$), with the C-terminal domain of PRIM2 ($PRIM2_C$) tethered by a flexible linker [22-29]. POLA1, which is functional on its own, relies on POLA2 for heterotetramer formation [30-33]. Structural analysis of the yeast enzyme revealed a bilobal structure in which the catalytic domain of POLA1 ($POLA1_{cat}$) is flexibly tethered via its C-terminal domain ($POLA1_C$) to the tetramer core (TC), consisting of PRIM1, $PRIM2_N$, $POLA1_C$, and the C-terminal domain of POLA2 ($POLA2_C$) [26,30]. In the absence of nucleic acid substrates or protein partners, the human enzyme has only been observed in a compact, non-catalytic ("auto-inhibitory" or AI) configuration in which the active site of $POLA1_{cat}$ is occluded by the tetramer core, as visualized by both X-ray crystallography and electron microscopy (EM) [34,35]. The human enzyme has also been visualized in catalytically accessible configurations

associated with the replisome [36] and the telomere maintenance CTC1–STN1–TEN1 (CST)

complex [37]. However, neither structure showed polα–primase engaged with primer/template.

Several models have emerged in recent years that largely attribute the counting ability of

polα–primase to the PRIM2$_C$ regulatory domain. Despite the catalytic center of primase residing

in PRIM1, PRIM2$_C$ is critical for all aspects of RNA synthesis, including regulation of RNA length

[23-25,38,39]. PRIM2$_C$ has also been implicated in transfer of the RNA primer to polα by virtue of its

interaction with the primer 5′-triphosphate and the template 3′-overhang [21,34,40,41]. These

interactions are believed to dictate RNA counting by spatially limiting the length of RNA

synthesized by PRIM1, either via the PRIM2 interdomain linker [41,42] or via steric collision

between PRIM2$_C$ and PRIM2$_N$ [33,34]. While these models explain the upper limit of RNA length

(i.e., 10 nucleotides), neither explains the range of RNA primers (7–10 nucleotides) extended by

polα [12]. PRIM2$_C$ and the primer 5′-triphosphate have also been shown to be important for DNA

synthesis [43,44], although the role of PRIM2$_C$ in the transition from RNA to DNA synthesis and in

DNA counting by polα is unknown. However, a recent model for DNA synthesis proposed a

similar spatial limitation imposed by the POLA1 and PRIM2 interdomain linkers [44]. Alternatively,

structures of POLA1$_{cat}$ bound to RNA/DNA and DNA/DNA duplexes afforded a model for DNA

counting whereby preferential binding to relatively A-form duplexes by polα leads to decreased

affinity and eventual dissociation from the primer/template as a result of a progressive increase

in B-form character as DNA is synthesized [32,45].

The molecular details by which polα–primase synthesizes chimeric primers of defined

length and composition have remained a matter of debate because there are few structures of

the heterotetramer bound to primer/template [46,47], and no structures showing the intermediate

and final stages of DNA synthesis. Here, we present cryo-EM structures of polα–primase alone

and bound to primed templates representing DNA synthesis at early, intermediate, and final

stages, beginning at the point immediately following transfer of the RNA primer to polα. Our

structures reveal that dissociation of POLA1$_{cat}$ and PRIM2$_C$ from the AI complex enables a

series of configurational and conformational changes that span a remarkable range of motion,

both in the absence and the presence of nucleic acid substrates. At the onset of DNA synthesis,

we observe a direct interaction between POLA1$_{cat}$ and PRIM2$_C$ that provides an explanation for

how polα can extend RNA primers ranging in length from 7–10 nucleotides, and that suggests

primer transfer is largely defined by competition between PRIM1 and POLA1$_{cat}$ for the primer 3′-

end. We also observe association of PRIM2$_C$ with chimeric RNA-DNA substrates corresponding

to intermediate and late stages of DNA synthesis. Complementary biochemical experiments

comparing the synthesis activities of the full polα–primase complex and various subcomplexes

are consistent with $PRIM2_C$ acting as a limited DNA processivity factor, reducing the proportion

of both shorter and longer primers [44]. Additionally, we provide direct structural evidence linking

primer composition, duplex conformation, and substrate affinity, with central roles for $POLA1_{cat}$

and $PRIM2_C$. Together, these findings establish a comprehensive model for chimeric primer

synthesis by polα–primase.

Results

<u>Polα–primase exists in an equilibrium between compact and extended configurations</u>

Prior structures in the absence of nucleic acid substrates or protein partners have shown

polα–primase in one of two states [26,34,35,37,47-49], suggesting the heterotetramer undergoes a

dramatic structural transition from a compact, auto-inhibitory state to an extended, catalytically

competent configuration prior to primer synthesis. To better understand the pre-existing

equilibrium between the two states, we first analyzed the protein alone using single-particle

cryo-EM. *Xenopus laevis* polα–primase was selected based on the stability and the quality of

the purified complex (**Fig. 2A**) and the high sequence conservation with human polα–primase

(69% identity and 90% similarity) (**Supplemental Figs. S1–S4**). We removed the N-terminal tail

of POLA1 (residues 1–334) to further improve the stability of the complex. In the absence of

nucleic acid substrates, 2D classification sorted 69% of particles into classes with density for all

four subunits in the compact AI configuration, 30% into classes lacking density for $POLA1_{cat}$,

and the remaining 1% into a class with density for only $POLA1_{cat}$ (**Fig. 2B and Supplemental**

Figs. S5 and S6). We initially refined a partial 3D reconstruction of the AI complex to a

resolution of 2.8 Å that exhibited poor density for PRIM1 (**Supplemental Figs. S5 and S7 and**

Supplemental Table S1). To improve the interpretability of the map in this region, we identified

a subset of particles that yielded a 3.2-Å reconstruction with clear density for all four subunits

(**Fig. 2C, Supplemental Figs. S5 and S7, and Supplemental Table S1**). Only the small N-

terminal domain of POLA2 ($POLA2_N$) was not well defined. As previously observed in lower-

resolution structures [34,35,47,48], the AI complex is stabilized by extensive contacts between

$POLA1_{cat}$ and every domain in the tetramer core, as well as $PRIM2_C$. Conversely, $PRIM2_C$

contacts only $POLA1_{cat}$, $PRIM2_N$, and the interdomain linkers in POLA1 (residues 1242–1275)

and PRIM2 (residues 262–278). These interactions sterically occlude substrate recognition by

both $POLA1_{cat}$ and $PRIM2_C$. To better understand the structural rearrangements necessary to

bind nucleic acid substrates, dissociation of $POLA1_{cat}$ and $PRIM2_C$ from the AI complex was

examined by 3D variability analysis, using all particles initially aligned on the AI complex during

preliminary 3D refinement (**Supplemental Figs. S5 and S8**). Two principal motions were

observed: (1) a hinging motion in $PRIM2_N$ that pulls PRIM1 away from $POLA1_{cat}$, and (2) a

bending/twisting motion in $POLA1_{cat}$ that pulls the exonuclease domain away from $POLA2_C$ and

that begins to pull the palm domain away from $PRIM2_N$. Interactions involving the thumb domain

were unchanged, as were all interactions involving $PRIM2_C$.

Without the conformational restrictions imposed by engagement of $POLA1_{cat}$ in the AI

complex, conformational heterogeneity is increased in the tetramer core. The subset of particles

grouped into 2D classes that lacked density for $POLA1_{cat}$ could be further divided into classes

with clear, partial, or no density for $PRIM2_C$ (**Fig. 2B and Supplemental Figs. S5 and S6**). 3D

reconstructions of the tetramer core were generated from particles selected through multiclass

ab initio reconstruction and heterogeneous refinement. These reconstructions revealed three

distinct conformations characterized by hinging motions in which the PRIM1 and

$POLA1_C/POLA2_C$ lobes move relative to one another by pivoting around flexible loops in

$PRIM2_N$ (residues 153–157 and 216–219) and $POLA1_C$ (residues 1439–1443), respectively

(**Fig. 2D,E and Supplemental Fig. S5**). Together, these data demonstrate that in the absence

of nucleic acid substrates, polα–primase exists predominantly in a compact auto-inhibitory state,

but also adopts a series of extended configurations in which $POLA1_{cat}$ and $PRIM2_C$ are

disengaged from the other domains in the complex. Based on 3D variability analysis of the AI

complex, along with inspection of the 2D class averages, we speculate that dissociation of

$POLA1_{cat}$ precedes dissociation of $PRIM2_C$, and initiates the structural transition away from an

auto-inhibitory state.

<u>$PRIM2_C$ remains bound to the primer during DNA synthesis</u>

Polα–primase possesses dual enzymatic activities that enable synthesis of chimeric RNA-

DNA primers. To investigate the molecular bases for the transition from RNA to DNA synthesis

and the low processivity of polα, we designed substrates to visualize polα–primase at two key

points during DNA synthesis—immediately after intramolecular handoff of the RNA primer from

primase to polα (DNA initiation, DI) and after polα has elongated the primer with DNA (DNA

elongation, DE). The DI primer is comprised of nine ribonucleotides and the chimeric DE primer

is comprised of nine ribonucleotides followed by 11 deoxyribonucleotides (**Fig. 3A,**

Supplemental Fig. S9, and Supplemental Table S2). Both primers include the critically

important 5′-triphosphate group that would be introduced by primase during RNA initiation.

Ternary complexes containing polα–primase, primer/template, and incoming nucleotide were

trapped using 2′,3′-dideoxycytidine to terminate the primer strand and prevent phosphodiester

bond formation with dGTP.

Relative to polα–primase in the absence of substrate, the 2D class averages determined

in the presence of the DI and DE substrates showed increased configurational heterogeneity

and dramatic configurational rearrangements (**Fig. 3B and Supplemental Figs. S10–S13**). The

vast majority of the class averages lacked clear density for part of the complex, indicating a

large degree of flexibility and conformational/configurational heterogeneity remains when

substrates are bound. These class averages could be divided into two groups: those with

density for POLA1$_{cat}$ and PRIM2$_C$ bound to substrate (hereafter referred to as the DNA

synthesis assembly, or DSA), and those with density for everything else (i.e., the tetramer core).

With the DI substrate, 2D classes showed 67% of particles with clear density for only the DNA

synthesis assembly, 26% for only the tetramer core, and ~6% for polα–primase in various

configurations. With the DE substrate, 2D classes showed a similar distribution, but with fewer

particles aligned on the DNA synthesis assembly and more particles aligned on the tetramer

core. Only 0 and 3% of particles with the DI and DE substrates, respectively, were found to be

in the AI configuration.

To better characterize interactions with the substrates, we performed focused 3D

refinement using masks around POLA1$_{cat}$, PRIM2$_C$, and the primer/template (**Fig. 3C,**

Supplemental Figs. S10, S12, S14, and S15, and Supplemental Tables S3 and S4). With

both the DI and DE substrates, the 3.3-Å and 4.4-Å reconstructions showed POLA1$_{cat}$ and

PRIM2$_C$ bound to opposing ends of the duplexes (**Fig. 3C**), indicating that interactions with

PRIM2$_C$ are maintained during transfer of the primer to polα and throughout DNA synthesis.

Focused 3D refinement of only POLA1$_{cat}$ and the DNA/DNA portion of the DE substrate

improved the resolution of the reconstruction to 3.5 Å (**Supplemental Figs. S12 and S15 and**

Supplemental Table S4). In both DSA structures, POLA1$_{cat}$ is in a closed ternary complex with

the primer 3′-end and the incoming dGTP, and PRIM2$_C$ is bound to the primer 5′-triphosphate.

With the DI substrate, POLA1$_{cat}$ and PRIM2$_C$ are in contact with one another, albeit with the

entire subunit interface consisting of a single interaction between two loops (**Fig. 3C and**

Supplemental Fig. S16). With the longer DE substrate, the two domains are separated by a full

turn of the RNA-DNA/DNA duplex. 3D variability analysis of these DI and DE subcomplexes

revealed dissociation of PRIM2$_C$ from a subset of the particles, as well as bending of the

oligonucleotide duplexes (**Supplemental Figs. S17 and S18**). With the shorter DI substrate,

this bending altered the protein-protein interface between POLA1$_{cat}$ and PRIM2$_C$.

Prompted by the seeming malleability and small size of the subunit interface, we modeled

POLA1$_{cat}$ and PRIM2$_C$ with shorter RNA primers (**Supplemental Fig. S16**). With 8-mer and 7-

mer primers, we observed only minor clashes between side chains, which appear unlikely to

prevent DNA initiation. Conversely, we saw severe steric overlap between POLA1$_{cat}$ and

PRIM2$_C$ with a 6-mer primer. These models explain the previous observation that polα can

extend RNA primers as short as seven nucleotides in length [12]. Additionally, binding of the

nascent primer by PRIM2$_C$ during DNA elongation explains the earlier observation that the

primer 5′-triphosphate affects not only RNA synthesis but also DNA synthesis [43], as well as the

recent finding that PRIM2$_C$ increases the processivity of polα–primase [44].

For simultaneous binding of the primer ends to be maintained throughout DNA synthesis,

as the primer increases in length, large-scale protein conformational changes are necessary.

Because of the extreme conformational and configurational heterogeneity inherent in the polα–

primase complex, we were unable to generate reconstructions of the full substrate-bound

complex by multiclass ab initio reconstruction and heterogeneous refinement. However, we

were able to generate reconstructions from subsets of particles manually selected from 2D class

averages (**Supplemental Figs. S19 and S20 and Supplemental Tables S3 and S4**). Rigid-

body docking of the DSA and TC subcomplexes into these low-resolution reconstructions

enabled us to model two configurations of the complete polα–primase complex bound to DI

substrate and two configurations bound to DE substrate (**Fig. 3D,E, Supplemental Figs. S19**

and S20, and Supplemental Tables S3 and S4). Comparison of the four substrate-bound

configurations showed that the DNA synthesis assembly rotates by as much as 180° with

respect to the tetramer core, with the pivot point occurring between PRIM2$_C$ and PRIM2$_N$ (**Fig.**

3E). In addition to this swiveling, the DNA synthesis assembly rotates by nearly 180° with

respect to the helical axis of the oligonucleotide duplexes. In all models of the four-subunit

complex, the substrate-bound configurations appear to be stabilized by interactions between the

DNA synthesis assembly and the tetramer core. However, the variability of these interactions,

as well as the large proportion of 2D class averages showing density for only one or the other

subcomplex, suggests these protein-protein contacts are weak and transient. Thus, in the

presence of primer/template substrates representing two key stages of DNA synthesis, polα–

primase adopts a wide range of configurations characterized by concerted movement of

POLA1$_{cat}$, PRIM2$_C$, and the oligonucleotide duplex relative to the tetramer core.

<u>Primase enhances an intrinsic counting ability in polα</u>

Whether counting of DNA nucleotides is an inherent property of polα or only a property of

the complete polα–primase complex is unclear [32,34,44,50,51]. Our finding that $PRIM2_C$ remains

bound to the primer after transfer to polα reveals a means by which primase might participate in

counting during DNA synthesis. We theorized that binding of the substrate by $PRIM2_C$ would

alter DNA elongation by indirectly tethering $POLA1_{cat}$ to the primer via the tetramer core. To

investigate this theory, we compared the RNA and DNA synthesis activities of the polα–primase

complex with a complex in which the interdomain linker between $POLA1_{cat}$ and $POLA1_C$ was

severed, constructed by mixing purified $POLA1_{cat}$ and $\Delta POLA1_{cat}$

(PRIM1/PRIM2/$POLA1_C$/POLA2) (**Fig. 4A and Supplemental Fig. S21**). As a control, we

measured the RNA and DNA synthesis activities of $POLA1_{cat}$ and $\Delta POLA1_{cat}$ separately. Primer

extension was performed by adding radiolabeled ATP or dATP and unlabeled NTPs and dNTPs

to a 5′-triphosphorylated 6-ribonucleotide primer annealed to a 99-deoxyribonucleotide template

(**Fig. 4B and Supplemental Table S2**). $POLA1_{cat}$ alone generated extension products of

predominantly 17–33 nucleotides, with the main product being 30 nucleotides in length (**Fig.**

4A,C). Similar to previous observations with primase alone [21,29], $\Delta POLA1_{cat}$ rapidly formed

products of nine nucleotides before abruptly terminating at a 10-mer primer (**Fig. 4A**). The

severed complex generated by combining $POLA1_{cat}$ and $\Delta POLA1_{cat}$ afforded products

representing an admixture of the individual control reactions (**Fig. 4A,C and Supplemental Fig.**

S22). In contrast, intact polα–primase generated fewer extension products outside of 30

nucleotides (**Fig. 4A,C**), and produced the 30-mer product at a dramatically faster rate (**Fig.**

4D). However, the total activity of polα–primase was notably less than that of the other

constructs. We speculate this reduced activity results from a substantial proportion of the polα–

primase molecules being in the AI configuration, and therefore unable to perform primer

elongation without undergoing a dramatic configurational transition. Collectively, these data

demonstrate an intrinsic—albeit approximate—counting ability in polα that is improved by the

physical linkage between $POLA1_{cat}$ and the remainder of the complex, reinforcing the

preference of polα–primase to generate a 30-nucleotide primer.

To better understand the approximate counting ability of polα, we performed binding

assays with $POLA1_{cat}$ and DNA/DNA, RNA/DNA, and two RNA-DNA/DNA duplexes (**Fig. 4E**

and Supplemental Table S5). Consistent with previous findings [32,45], we observed significantly

tighter binding of an RNA/DNA substrate than a corresponding DNA/DNA substrate. Addition of

11 nucleotides of DNA to the RNA primer, creating an RNA-DNA/DNA substrate, had no effect

on binding affinity. However, a further addition of nine nucleotides of DNA, producing a longer

RNA-DNA/DNA duplex, resulted in significantly weaker binding, approaching that of the

DNA/DNA substrate. For both RNA-DNA/DNA duplexes, the DNA/DNA regions were sufficiently

long to fully accommodate the binding footprint of the polymerase catalytic core. Thus, these

findings indicate the affinity of POLA1$_{cat}$ for chimeric primers is dependent on not only the length

of the DNA/DNA region of the duplex but also the proximity of the duplex end to the RNA/DNA

region, providing a basis for counting and the termination of DNA synthesis.

<u>Primer/template conformation modulates substrate recognition</u>

Consistent with the results from our binding assays, the intrinsic counting ability of polα

has been proposed to result from conformation-dependent recognition of RNA/DNA and

DNA/DNA duplexes [32,45]. Because of the relatively high resolutions of the DSA reconstructions,

we were able to determine the conformations of the 5′-triphosphorylated RNA/DNA and RNA-

DNA/DNA substrates in the DI and DE subcomplexes and to evaluate the details of their

recognition by POLA1$_{cat}$ and PRIM2$_C$. In both structures, POLA1$_{cat}$ was observed in a closed

conformation in which the fingers domain was fully engaged with an incoming dGTP and active-

site residues within the palm domain (Asp859, Phe860, and Asp1000) coordinated a catalytic

Mg^{2+} ion (**Fig. 5A,B and Supplemental Fig. S23**). The thumb domain was also fully engaged

with both the RNA/DNA and RNA-DNA/DNA duplexes (**Fig. 5A,C**). To our knowledge, the DE

subcomplex represents the first structure of POLA1$_{cat}$ with the thumb and fingers domains

engaged with a DNA/DNA duplex and an incoming dNTP (**Supplemental Fig. S24**). In both

subcomplexes, POLA1$_{cat}$ induces a local A-form conformation in the duplexes that positions and

orients the 2′,3′-dideoxycytidine at the 3′-end of the primers for phosphodiester bond formation

with dGTP (**Fig. 5B,E,F**). Similar catalytic arrangements have been observed in structures of

polδ and polε [52,53]. However, POLA1$_{cat}$ possesses a wider substrate-binding cleft that results

from structural differences in the thumb domain (**Supplemental Fig. S25**). This wider cleft has

been proposed to complement the wider diameter of an A-form RNA/DNA duplex [32,45]. In both of

our structures, we observed intermediate AB-form conformations [54,55], or nearly intermediate

AB-form conformations, over most of the region of the duplexes bound by POLA1$_{cat}$ (**Fig. 5E,F**).

Importantly, small but significant differences in the interactions between the thumb domain

and the DI and DE substrates illustrate how POLA1$_{cat}$ accommodates different minor groove

geometries (**Fig. 5A,C,E,F**). With the DI substrate, three aspartate residues (Asp998, Asp1079,

and Asp1144) form hydrogen-bonding interactions with 2′-hydroxyl groups along the RNA

primer. With the DE substrate, conformational changes at the tip of the thumb domain (residues

1142–1148) pull Asp1144 away from the chimeric primer while directing Lys1145 and Lys1146

toward the relatively narrow minor groove in the same DNA/DNA region of the duplex. These

new hydrogen-bonding interactions likely partially offset the loss of RNA-specific hydrogen

bonds with the DE substrate. While RNA/DNA and DNA/DNA duplexes generally adopt A-form

and B-form geometries [56], respectively, the intermediate AB-form conformations observed with

both the DI and DE substrates suggest POLA1$_{cat}$ remodels the primer/template during DNA

synthesis. The local propensity of the duplex to adopt this intermediate conformation is likely

correlated with substrate affinity and, consequently, with processivity [32,45].

PRIM2$_C$ has been shown to increase the processivity of polα–primase during DNA

synthesis [44]. The two DSA structures suggest that this increased processivity is the result of

continued association of PRIM2$_C$ with the primer/template during DNA elongation. In both the DI

and DE subcomplexes, interactions between PRIM2$_C$ and the primer strand are identical (**Fig.**

5A,D). Three positively charged residues (His306, Arg308, and Arg312) at the N-terminus of

helix α15 surround the 5′-triphosphate and offset the negative charge, and His309 stacks

against the 5′-terminal nucleobase. Surprisingly, interactions between PRIM2$_C$ and the template

strand are markedly different in the two subcomplexes, despite PRIM2$_C$ binding the same

RNA/DNA portion of both substrates. In the DI structure, the DNA template is positioned such

that only the last paired nucleotide, which is not coplanar with the 5′-terminal nucleotide in the

RNA primer, and the first unpaired nucleotide form extensive contacts with PRIM2$_C$. Rotation of

the phosphodeoxyribose backbone swings the second and third unpaired nucleotides away

from PRIM2$_C$ and toward POLA1$_{cat}$. In the DE structure, the DNA template is positioned such

that the last paired nucleotide, which is coplanar with the 5′-terminal nucleotide in the RNA

primer, and the first, second, and third unpaired nucleotides form extensive contacts with

PRIM2$_C$. Relative to the DI substrate, interactions with several residues (Tyr353, His357,

Lys364, and Tyr368) in the flexible template-binding loop (residues 359–372) are shifted by one

nucleotide toward the 5′-end of the template strand as a result of increased base pair inclination

in the DE substrate. Virtually identical interactions were observed in an X-ray crystal structure of

human PRIM2$_C$ bound to a short RNA/DNA duplex [34] (**Supplemental Fig. S26**).

The unique interactions that we observe with the DI substrate appear to result from local

duplex remodeling by POLA1$_{cat}$, which causes the RNA/DNA duplex to have less A-form

character, as quantified by the Z$_P$ parameter [54,55,57], than either the RNA/DNA portion of the DE

substrate or the RNA/DNA duplex in the crystal structure (**Fig. 5D,E,F and Supplemental Fig.**

S26). In addition to reduced A-form character, the DI substrate has lower base pair inclination,

which pulls the 3′-overhang away from PRIM2$_C$. The affinity of PRIM2$_C$ for the primer/template

was previously shown to depend on both the 5′-triphosphate and the 3′-overhang [34,41]. Thus,

tight binding of the substrate appears to require substantial inclination of the base pairs at the

end of duplex. Together, these data indicate that like POLA1$_{cat}$, the affinity of PRIM2$_C$ for the

primer/template is modulated by the nucleic acid conformation, which is imposed by both the

DNA composition of the primer and binding by the polymerase catalytic core.

<u>Loose association of full-length primers allows for reversion to the auto-inhibitory configuration</u>

The finding that substrate recognition by both POLA1$_{cat}$ and PRIM2$_C$ is dependent on

duplex conformation has implications for the mechanism of DNA counting. To investigate how

duplex conformation affects the preference of polα–primase to generate 30-nucleotide primers,

we designed a substrate to visualize the complex at the end of DNA synthesis (DNA

termination, DT). The DT primer is comprised of nine ribonucleotides followed by 20

deoxyribonucleotides (**Fig. 6A, Supplemental Fig. S9, and Supplemental Table S2**) and

includes the same 5′-triphosphate group as the DI and DE primers. A ternary DT complex

containing polα–primase, primer/template, and incoming nucleotide was trapped using 2′,3′-

dideoxycytidine to terminate the primer strand and prevent phosphodiester bond formation with

dGTP.

As with the DI and DE substrates, a substantial proportion of 2D class averages showed

density for only the DNA synthesis assembly or the tetramer core (**Fig. 6B and Supplemental**

Figs. S27 and S28), consistent with large-scale conformational and configurational dynamics.

Unlike with the shorter substrates, however, no class averages showed density for the complete

polα–primase complex bound to substrate. This level of heterogeneity suggests that the

interdomain linkers in POLA1 and PRIM2 remain slack when bound to a full-length primer, as

taught linkers would seemingly reduce the overall flexibility of the DT complex. Surprisingly,

63% of particles were grouped into 2D classes with density clearly representative of the AI

complex. Protein-protein contacts in this configuration sterically occlude substrate recognition by

both POLA1$_{cat}$ and PRIM2$_C$, indicating the DT substrate is not bound in most particles. With the

DI and DE substrates, only 0 and 3% of particles, respectively, were found to be in the AI

configuration, while in the absence of substrate, 69% of particles were found to be in this

configuration. While this type of comparative quantitation of cryo-EM data has the potential for

bias, this dramatic increase in AI particles in the presence of DT substrate suggests

substantially weaker binding of the longer chimeric primer by both POLA1$_{cat}$ and PRIM2$_C$.

To better understand the structural basis for weaker association of the DT substrate, we

performed focused 3D refinement of the DT subcomplex. Using a mask around POLA1$_{cat}$ and

the DNA/DNA region of the primer/template, we generated a 2.9-Å reconstruction with clear

density for POLA1$_{cat}$ in a closed conformation, with the thumb and fingers domains fully

engaged with the duplex and an incoming dGTP (**Fig. 6C, Supplemental Figs. S27 and S29,**

and Supplemental Table S6). 3D variability analysis of the complete DT subcomplex, including

POLA1$_{cat}$, PRIM2$_C$, and the full-length primer/template, revealed duplex bending similar to that

seen in the DE subcomplex, as well as dissociation of PRIM2$_C$ from the substrate

(**Supplemental Fig. S30**). However, the degree to which PRIM2$_C$ is disengaged from the

primer/template appears to be substantially greater with the DT substrate. Consistent with this

observation, 3D classification sorted most particles aligned on the DT subcomplex into classes

with little to no density for PRIM2$_C$ (**Supplemental Fig. S27**). Following classification, focused

3D refinement provided 3.8- and 3.6-Å reconstructions of the DT subcomplex with (DT

subcomplex 1, or DTS-1) and without (DT subcomplex 2, or DTS-2) PRIM2$_C$ engaged with the

substrate (**Fig. 6D, Supplemental Figs. S27 and S29, and Supplemental Table S6**). The

overall conformations of the duplexes in these structures, as quantified by the Z_P parameter, are

quite similar (**Fig. 6E**). However, the base pair inclination in the RNA/DNA portion of DTS-2

appears to be reduced relative to DTS-1 (**Fig. 6F and Supplemental Fig. S31**). While the

degree of certainty associated with this observation is limited by the local resolution of the

reconstructions (**Supplemental Fig. S29**), this conformational difference is consistent with our

conclusion that tight substrate binding by PRIM2$_C$ is dependent on the base pair inclination at

the end of the duplex. Conversely, the origins for weaker binding of the DT substrate by

POLA1$_{cat}$ are not apparent in the structures. In particular, all structures of the DNA synthesis

assembly reveal POLA1$_{cat}$ has remodeled the bound region of the duplex—whether it is

RNA/DNA or DNA/DNA—to achieve similar intermediate AB-form conformations

(**Supplemental Fig. S31**). However, remodeling of this type requires an input of free energy,

which in turn reduces the free energy associated with binding affinity. Thus, primer/template

duplexes predisposed to adopt intermediate AB-form conformations, like those in the remodeled

substrates, require less free energy to be remodeled, and therefore will be bound with higher

affinity.

Discussion

Multiple, seemingly conflicting models have been proposed to explain the counting abilities

of primase and polα [32-34,41,42,44-47]. Our structures of polα–primase create a framework to

reconcile and integrate these models, providing a comprehensive mechanistic understanding of

nucleotide counting (**Fig. 7A,B**). We show that in the absence of nucleic acid substrates or

protein partners, polα–primase exists primarily in a compact, auto-inhibitory state, but

spontaneous dissociation of POLA1$_{cat}$ and PRIM2$_C$ initiates a series of dramatic structural

rearrangements necessary for chimeric primer synthesis across two spatially separate active

sites. In all current models of RNA synthesis, primer formation is initiated by PRIM1 and PRIM2$_C$

bringing together two NTPs on a single-stranded DNA template. In the linker model, elongation

of the primer continues until halted by full extension of the PRIM2 interdomain linker [41,42,47]. In

the collision model, elongation is ceased by a steric clash between PRIM2$_N$ and PRIM2$_C$ [34]. As

previously discussed [34], the linker model is not consistent with domain arrangement in the

tetramer core. Conversely, the collision model is supported by structural data and structure-

based modeling [34], and provides a convincing rationale for termination of RNA synthesis, but

strictly at 9–10 nucleotides. Both models fail to explain previous observations showing extension

of RNA primers as short as seven nucleotides in length by polα [12].

Based on our structure of the DI subcomplex, as well as previous biochemical data, we

propose a competition model for RNA counting and primer transfer. PRIM2$_C$ has been

suggested to anchor the primer during its transfer between primase and polα active sites by

remaining bound to the primer 5′-end [34,37,44]. Moreover, the highly distributive nature of RNA and

DNA synthesis by PRIM1 and POLA1$_{cat}$ in the absence of PRIM2$_C$ [29,44,58] suggests that PRIM2$_C$

acts as a hub to facilitate re-association of PRIM1 and POLA1$_{cat}$ with the primer. Our structures

confirm that PRIM2$_C$ remains bound to the 5′-end of the primer after transfer to polα, and

provide evidence that a dynamic protein-protein interface allows PRIM2$_C$ and POLA1$_{cat}$ to

simultaneously bind RNA primers of seven or more nucleotides, while transfer of RNA primers

less than seven nucleotides in length to POLA1$_{cat}$ would be sterically blocked by PRIM2$_C$. Thus,

the likely driving factor in the primase-to-polymerase switch is a competition between PRIM1

and POLA1$_{cat}$ for the 3′-end of the primer, rather than termination of RNA synthesis at a

maximal length of 10 nucleotides. In this competition model, steric collision between PRIM2$_N$

and PRIM2$_C$ would serve only as a failsafe to terminate RNA synthesis, ensuring longer

stretches of RNA (> 10 nts) are not incorporated into the genome, should POLA1$_{cat}$ not

outcompete PRIM1 for the primer.

Our finding that PRIM2$_C$ remains bound to the primer after transfer to polα has far-

reaching implications for DNA synthesis, including the preference of polα–primase to generate

chimeric primers of ~30 nucleotides. Simultaneous association of both PRIM2$_C$ and POLA1$_{cat}$ is

a prerequisite for the linker model of DNA termination, which posits that the POLA1 and PRIM2

interdomain linkers prevent separation of PRIM2$_C$ and POLA1$_{cat}$ beyond the distance associated

with an ~30-mer primer [44,47]. However, we observed seemingly weaker binding of chimeric

primers by both POLA1$_{cat}$ and PRIM2$_C$ as the DNA composition of the primer increased (as

measured by direct binding assays using POLA1$_{cat}$ and implied by the proportion of particles in

the AI configuration). Moreover, if termination were defined only by the length of the interdomain

linkers in POLA1 and PRIM2, we would expect to observe a stable complex between polα–

primase and the DT substrate, likely with well-ordered, taught linkers. Additionally, the linker

model is unable to provide a rationale for the intrinsic counting ability of POLA1$_{cat}$, which we and

others have observed [32,50]. By contrast, the affinity model of DNA termination, which proposes

that POLA1$_{cat}$ preferentially recognizes A-form duplexes typical of RNA/DNA hybrids, and loses

affinity for the substrate as the duplex becomes more B-form throughout DNA synthesis [32,45],

provides a mechanistic explanation for the innate counting ability of the polymerase catalytic

core. Our data expand upon the affinity model by suggesting a role for preferential binding of A-

form duplexes by PRIM2$_C$, and by providing a possible structural basis for this specificity. Thus,

we propose that DNA termination is the result of both POLA1$_{cat}$ and PRIM2$_C$ losing affinity for

the primer/template as the B-form propensity of the duplex increases during DNA synthesis.

Because the linker model requires both POLA1$_{cat}$ and PRIM2$_C$ to remain bound to the substrate,

the interdomain linkers are likely to play, at most, a failsafe role in restricting primer length, and

thereby limiting the mutagenic potential of polα.

The dramatic structural rearrangements that we observe are consistent with the idea that

flexible tethering of POLA1$_{cat}$ and PRIM2$_C$ to the tetramer core is necessary for transfer of the

nascent primer from primase to polα [26,34]. As we demonstrate here, these rearrangements are

also necessary for simultaneous binding of the primer ends by POLA1$_{cat}$ and PRIM2$_C$ during

DNA elongation, as the primer increases in length. Presumably, this plasticity also facilitates

interaction with the many protein partners with which polα–primase is known to interact at

replication forks and telomeres. At replication forks, polα–primase interacts with several proteins

associated with the replisome, including MCM10, AND-1, RPA, RFC, polδ, and the CMG

helicase [19,36,59], as well as multiple nucleosome core proteins [60,61]. Despite this extensive

network of contacts, polα–primase appears to remain conformationally dynamic [62,63]. Moreover,

recent structures of polα–primase bound to a replisome complex revealed a manner of flexible

association [36] that is unlikely to restrict even the extraordinarily broad range of motion that we

observed in our structures.

Polα–primase appears to remain similarly dynamic at telomeres, as three distinct

configurations of the polα–primase/CST complex bound to single-stranded DNA have been

observed [37,48,49]. The first structure, termed a recruitment complex, showed polα–primase in the

AI configuration with CST primarily associated via interactions with POLA1$_C$ [48]. The second

structure, designated a pre-initiation complex, showed polα–primase in an extended

configuration in which POLA1$_{cat}$ and the tetramer core are scaffolded by CST [37]. In this

configuration, the PRIM1 and POLA1$_{cat}$ active sites are locked in opposing orientations, with

PRIM2$_C$ free to move between them. This arrangement was proposed to facilitate RNA

synthesis and transfer of the RNA primer from primase to polα. Consistent with this notion, CST

has been shown to enhance primer synthesis, without substantially affecting primer length

[37,64,65]. Moreover, docking of our DSA structures into the pre-initiation complex revealed

additional interactions between the thumb domain of POLA1$_{cat}$ and the CTC1 and STN1

subunits of CST (**Supplemental Fig. S32**), which may stabilize the closed, substrate-bound

conformation of the polymerase catalytic core, supporting POLA1$_{cat}$ in its competition with

PRIM1 for the RNA primer. The positions of our structures in these models, however, suggest

DNA elongation would lead to minor and then severe steric clashes between PRIM2$_C$ and the

tetramer core as the primer reaches lengths of 10 and 18 nucleotides, respectively. In this

configuration, these clashes are likely to prevent synthesis of a full-length primer. This

conclusion is supported by the third structure of polα–primase/CST, which showed POLA1$_{cat}$

associated with CST, as in the pre-initiation complex, but with the tetramer core disengaged [49].

Together, these observations suggest the tetramer core is not tightly bound to CST and

dissociates prior to or during DNA elongation by POLA1$_{cat}$. Based on the evolutionarily

relationship between CST and RPA [66], we believe RPA likely scaffolds polα–primase in a similar

manner, explaining RPA-dependent stimulation of primer synthesis by polα–primase [67]. How

these interactions with RPA and CST are coordinated with those of other proteins at replication

forks and telomeres, as well as the effects these interactions may have on the conformational

and configurational dynamics of polα–primase and the mechanism of chimeric primer synthesis,

remain to be determined.

Conclusion

Determining structures of polα–primase with multiple catalytic substrates, along with

performing structure-based biochemical and biophysical analyses, has enabled us to propose a

comprehensive model for primer synthesis. This model is built upon the large body of available

structural and biochemical data while unifying and refining prevailing understanding of the

mechanistic basis for polα–primase function. Our structural data reveal that polα–primase both

remodels and reads the conformation of its primer/template substrate, which explains its unique

ability to regulate primer length (i.e., count). These data also reveal that polα–primase retains a

remarkable degree of conformational and configurational flexibility that is crucial to the intricate

functions of this complex and dynamic protein machine. A greater appreciation of the type and

degree of intrinsic flexibility that we have observed for polα–primase is anticipated as active

eukaryotic replisomes and other cellular machinery come into view at ever increasing resolution.

Methods

Cloning and expression. The genes encoding *Xenopus laevis* PRIM1 (NCBI accession

NP_001080819.1), PRIM2 (NCBI accession XP_018121015.1), POLA1 (NCBI accession

NP_001082055.1), and POLA2 (NCBI accession NP_001086972.1) were initially cloned into

pFastBac-HTa or pFastBac-HTb expression vectors (Invitrogen). PRIM1 was then subcloned

into a modified pET-19b expression vector (Novagen) encoding an HRV 3C-cleavable

decahistidine fusion tag. PRIM2 was subcloned into a modified pRSFDuet-1 expression vector

(Novagen). And, POLA1 and POLA2 were subcloned into an extensively modified, polygenic

pFastBac-1 expression vector (Invitrogen) encoding TEV-cleavable StrepII-GFP (POLA1) and

hexahistidine-MBP (POLA2) fusion tags. Protein expression and purification were facilitated by

truncating the gene encoding POLA1 to remove the N-terminal tail (residues 1–334). To

generate a POLA1$_{cat}$ construct, the catalytic core of POLA1 (residues 335–1254) was subcloned

into a modified pET-27b(+) expression vector (Novagen) encoding an HRV 3C-cleavable

hexahistidine-SUMO fusion tag. To generate a ΔPOLA1$_{cat}$ construct, the C-terminal domain of

POLA1 (residues 1262–1458) and POLA2 were subcloned into a modified, polycistronic

pCDFDuet-1 expression vector (Novagen) encoding an HRV 3C-cleavable hexahistidine

(POLA1$_C$) fusion tag. Production of baculovirus and expression of POLA1/POLA2 in *Spodoptera*

frugiperda Sf9 cells were performed by the Expression and Microbiology Core at UC Berkeley.

PRIM1/PRIM2, POLA1$_{cat}$, and POLA1$_C$/POLA2 were expressed in *Escherichia coli* Rosetta

2(DE3) cells. Cultures were grown at 37°C in LB medium supplemented with 34 mg/L

chloramphenicol, 100 mg/L ampicillin (PRIM1/PRIM2), 30 mg/L kanamycin (PRIM1/PRIM2 and

POLA1$_{cat}$), and 70 mg/L streptomycin (POLA1$_C$/POLA2). Upon reaching mid-log phase, cultures

were further supplemented with 100 mg/L ammonium iron(III) citrate (PRIM1/PRIM2) and 5 μM

ZnSO$_4$ (PRIM1/PRIM2 and POLA1$_C$/POLA2) and cooled to 16°C. Protein overexpression was

then induced by addition of IPTG to a final concentration of 0.5 mM. After overnight incubation,

cells were harvested by centrifugation. Cell pellets were stored at −80°C. *Acanthamoeba*

castellanii TLP2 was expressed as previously described [68-70]. Protein constructs are depicted in

Supplemental Fig. S33.

Protein purification. Cell pellets were thawed in lukewarm water and resuspended in buffer L
[50 mM sodium phosphate pH 7.5, 1 M NaCl, 5% (v/v) glycerol, 1 mM MgCl$_2$, 10 mM imidazole,
and cOmplete EDTA-free protease inhibitor cocktail (Roche)]. Both insect cells and bacterial
cells were lysed by Dounce homogenization followed by gentle sonication. The two lysates were
then combined and cleared by centrifugation prior to application onto a Ni-NTA Agarose column
(Qiagen) equilibrated in buffer L lacking protease inhibitor cocktail. The column was washed
with 15 column volumes (cv) of buffer N1 [50 mM sodium phosphate pH 7.5, 1 M NaCl, 5% (v/v)
glycerol, 1 mM MgCl$_2$, and 20 mM imidazole], followed by 5 cv of buffer N2 [50 mM
HEPES•NaOH pH 7.5, 300 mM NaCl, 5% (v/v) glycerol, 1 mM MgCl$_2$, and 20 mM imidazole].
Protein was eluted by linearly transitioning over 15 cv to buffer N3 [50 mM HEPES•NaOH pH
7.5, 300 mM NaCl, 5% (v/v) glycerol, 1 mM MgCl$_2$, and 500 mM imidazole]. Fractions containing
polα–primase were pooled and supplemented with 1 mM TCEP before fusion tags were
removed by overnight cleavage at 4°C. The following day, protein was diluted three-fold in
buffer H1 [50 mM HEPES•NaOH pH 7.5, 5% (v/v) glycerol, 1 mM MgCl$_2$, and 1 mM TCEP] and
passed through a 0.22-µm filter before being applied to a heparin Sepharose column (Cytiva)
equilibrated in a 9:1 mixture of buffer H1 and buffer H2 [50 mM HEPES•NaOH pH 7.5, 1 M
NaCl, 5% (v/v) glycerol, 1 mM MgCl$_2$, and 1 mM TCEP]. The column was then washed with 5 cv
of the same 9:1 mixture before protein was eluted by linearly transitioning over 10 cv to only
buffer H2. Fractions containing polα–primase were pooled and again applied to a Ni-NTA
Agarose column equilibrated in buffer L lacking protease inhibitor cocktail. Protein was eluted in
10 cv of the same buffer, and fractions containing polα–primase were pooled and supplemented
with 1 mM TCEP. Protein was then concentrated by diafiltration, cleared by centrifugation, and
applied to a Superose 6 Increase column (Cytiva) equilibrated in buffer S [20 mM
HEPES•NaOH pH 7.5, 150 mM NaCl, 5 mM MgCl$_2$, and 1 mM TCEP]. Protein was eluted in 1.5
cv of the same buffer, and fractions containing polα–primase were pooled and concentrated by
diafiltration. Aliquots were used immediately or frozen in liquid nitrogen and stored at −80°C.
POLA1$_{cat}$ and ΔPOLA1$_{cat}$ were purified in the same manner as polα–primase. AcaTLP2 was
purified as previously described [68-70].

Oligonucleotide synthesis and purification. RNA, DNA, and chimeric RNA-DNA
oligonucleotides were purchased from Integrated DNA Technologies and resuspended in buffer
A (10 mM Mes•NaOH pH 6.5 and 40 mM NaCl). To generate 5′-triphosphorylated primers, 500
µM 5′-monophosphorylated primer and 550 µM template were annealed by heating to 85°C and

then slowly cooling to 22°C, producing a double-stranded substrate for the tRNA-modifying

enzyme AcaTLP2 [70,71]. AcaTLP2 reaction mixtures were prepared with 50 µM enzyme, 100 µM

primer/template, 150 µM ATP, 1 mM GTP, 25 mM Hepes•NaOH pH 7.5, 125 mM NaCl, 10 mM

$MgCl_2$, and 3 mM DTT and incubated at 22°C. After 1 h, an equal volume of quenching/loading

buffer containing 80% (v/v) formamide, 100 mM EDTA pH 8.0, and 220 µM competitor

oligonucleotide was added. Samples were then heated to 85°C and slowly cooled to 22°C to

denature the primer and the template. Oligonucleotides were separated by denaturing urea-

polyacrylamide gel electrophoresis (PAGE) as previously described [72], except electrophoresis

was performed at 30 W for 9 h. For each reaction, the band corresponding to the

triphosphorylated primer was located by UV-shadowing and cut from the gel. The primer was

then extracted into 24 mL of 300 mM NaOAc pH 5.5 using a crush-and-soak procedure in which

the sample was placed at −80°C until frozen, thawed in a water bath at 42°C, and then rocked

overnight at 4°C. The following day, the supernatant was decanted and overnight extraction was

repeated using a second 24-mL volume of 300 mM NaOAc pH 5.5. The two volumes were

combined, passed through a 0.22-µm filter, and split into four 12-mL aliquots. 36 mL of cold

100% ethanol was then added to each aliquot. Samples were thoroughly mixed, incubated at

−80°C for 2 h, warmed to −20°C, and centrifuged at 12,000×g and 4°C for 1 h. After decanting

the supernatant, pellets were air-dried for ~15 min and resuspended in 1 mL of 300 mM NaOAc

pH 5.5, using the same 1 mL of buffer for all pellets. The sample was then split into two 0.5-mL

aliquots, and 1.5 mL of cold 100% ethanol was added to each. Samples were again thoroughly

mixed, incubated at −80°C for 2 h, warmed to −20°C, and centrifuged at 12,000×g and 4°C for 1

h. After decanting the supernatant, pellets were air-dried for ~15 min and then gently rinsed with

200 µL of cold 95% (v/v) ethanol. The ethanol was removed and the pellet was air-dried for an

additional ~15 min. The triphosphorylated primer was then resuspended in 100 µL of buffer A,

using the same 100 µL of buffer for both pellets, and stored at −80°C. Oligonucleotide

sequences are provided in **Supplemental Table S2**.

Cryo-EM sample preparation and data collection. Polα–primase was concentrated to ≥ 3 µM

by diafiltration. To generate RNA-primed templates, 110 µM 5′-triphosphorylated RNA primer

and 100 µM DNA template were annealed by heating to 85°C and then slowly cooling to 4°C.

Substrate complexes were prepared by incubating 3 µM polα–primase, 3.6 µM primer/template,

and 1 mM dGTP on ice for 30 min. Using a Vitrobot Mark IV (FEI), 2.5 µL of sample was applied

to glow-discharged, holey-carbon R1.2/1.3 200-mesh copper grids (Quantifoil) at 4°C and 100%

humidity. Grids were immediately blotted with filter paper (Ted Pella) and flash-frozen in liquid

ethane. Grids prepared without substrate or with DT substrate were loaded into a Titan Krios G4

electron microscope (Thermo Scientific) operated at 300 kV for automated data collection using

SerialEM [73]. Movies of dose-fractionated frames were acquired with a K3 BioQuantum direct

electron detector (Gatan) functioning in counting mode. Each of the datasets was collected from

a single grid in a single imaging session. Grids prepared with DI or DE substrates were loaded

into a Talos Arctica electron microscope (Thermo Scientific) operated at 200 kV for automated

data collection using Leginon [74]. Movies of dose-fractionated frames were acquired with a K2

Summit direct electron detector (Gatan) functioning in counting mode. Each of the datasets was

collected in three to four imaging sessions, using a different grid for each session. Additional

details for each of the datasets are provided in **Supplemental Tables S1, S3, S4, and S6**.

Cryo-EM data processing. For pre-processing of all datasets, motion-correction of raw movies

was performed using MotionCor2 [75] and patch CTF estimation was performed for motion-

corrected micrographs using cryoSPARC [76]. All initial processing steps were also performed in

cryoSPARC. Particles from each dataset were picked by a combination of blob picking, template

picking, and neural-network picking using Topaz [77]. Templates for template picking were

selected from the highest-resolution 2D class averages resulting from blob picking, while Topaz

was trained from a small set of manually picked particles. Particles were extracted with a box

size of either 400 (substrate-free and DT datasets) or 360 (DI and DE datasets) pixels and then

downsampled to 120 pixels (pixel size of 2.73 Å) by Fourier truncation. 2D classification was

first performed separately for particles picked by each method. Selected particles were then

combined and duplicate and triplicate particles were removed before final 2D classification was

performed. For the DT dataset, particles were sorted manually before initial 3D reconstructions

were generated by ab initio reconstruction. For all other datasets, initial 3D reconstructions were

generated by multiclass ab initio reconstruction followed by heterogeneous refinement.

Satisfactory 3D reconstructions could only be obtained for the AI complex (substrate-free and

DT datasets), the DNA synthesis assembly (DI, DE, and DT datasets), and the tetramer core (all

datasets). Because of extreme configurational and conformational heterogeneity, satisfactory

3D reconstructions could not be obtained for the full polα–primase complex bound to a

primer/template substrate. For reconstructions of the tetramer core, conformational

heterogeneity was reduced through additional iterations of multiclass ab initio reconstruction

and heterogeneous refinement. For all 3D reconstructions, homogeneous refinement was

initially performed using downsampled particles and then repeated after particles were

recentered and re-extracted either without downsampling or with reduced downsampling. For

the AI complex from the substrate-free dataset and the DNA synthesis assembly from the DI,

DE, and DT datasets, all subsequent processing steps were performed in RELION-3.1 [78].

Particle alignments were first improved by focused 3D refinement, before heterogeneity was

reduced through focused, alignment-free 3D classification (following particle subtraction, as

necessary). CTF refinement was also performed to correct for per-particle defocus differences,

as well as for imperfect microscope alignment during data collection. Using refined CTF

parameters, focused 3D refinement was repeated for each final set of particles. To improve

high-resolution features in the reconstructions, despite local quality differences, maps were

sharpened using DeepEMhancer [79]. Data-processing workflows are outlined in **Supplemental**

Figs. S5, S10, S12, and S27.

A modified workflow was employed to obtain reconstructions of the full polα–primase

complex bound to primer/template substrates. Particles corresponding to configurations DI-1,

DI-2, DE-1, and DE-2 were manually selected by examination of 2D class averages and

downsampled to 64 pixels (pixel size of 5.12 Å) by Fourier truncation. Ab initio reconstruction

and non-uniform refinement were then performed to generate low-resolution 3D reconstructions.

For particles initially assigned to configuration DI-2, multiclass ab initio reconstruction and

heterogeneous refinement revealed a possible third configuration of the DI complex. All

processing steps in the modified workflow were performed in cryoSPARC. Modified data-

processing workflows are outlined in **Supplemental Figs. S19 and S20**.

Model building and refinement. Homology models of *X. laevis* polα–primase subunits were

generated from corresponding *Homo sapiens* X-ray crystal structures using SWISS-MODEL [80].

For the AI complex, POLA1, POLA2$_C$, and PRIM2 homology models (from PDB accession

5EXR) were docked as rigid bodies into the high-resolution partial reconstruction. Following

manual modification, atomic coordinates and isotropic temperature factors were refined against

the cryo-EM density for all non-hydrogen atoms using Ramachandran, secondary structure, and

rotamer restraints. Hydrogen atoms were placed in riding positions and were not refined against

the density. Both the refined model and a PRIM1 homology model (from PDB accession 4BPU)

were then docked as rigid bodies into the complete reconstruction. After manual adjustment,

atomic coordinates and isotropic temperature factors were refined as described above, except

reference restraints were applied based on the *X. laevis* PRIM2/POLA1/POLA2$_C$ structure and

the *H. sapiens* PRIM1 structure. For the DI subcomplex, POLA1$_{cat}$ and PRIM2$_C$ homology

models (from PDB accessions 4QCL and 5F0Q) were docked as rigid bodies into the

corresponding reconstruction. Following manual modification of the protein and de novo addition

of the substrate, atomic coordinates and isotropic temperature factors were refined as described

above, except base pairing and base stacking restraints were added for the oligonucleotide

duplex. For the DE subcomplex, the refined POLA1$_{cat}$ model from the DI subcomplex was

docked as a rigid body into the high-resolution partial reconstruction. After manual adjustment of

the protein and de novo addition of the substrate, atomic coordinates and isotropic temperature

factors were refined as described above for the DI subcomplex. The refined models of POLA1$_{cat}$

from the partial DE structure and PRIM2$_C$ from the DI structure were then docked as rigid bodies

into the complete reconstruction. The oligonucleotide substrate from the partial DE subcomplex

and the 3′-overhang and first base pair from the *H. sapiens* PRIM2$_C$ crystal structure (PDB

accession 5F0Q) were also docked as rigid bodies. Following manual modification of the

protein, including adjustment of the template-binding residues in PRIM2$_C$ to match those in the

H. sapiens crystal structure, and de novo addition of the intervening region of the substrate,

atomic coordinates and isotropic temperature factors were refined as described above for the DI

subcomplex, except reference restraints were applied based on the *X. laevis* POLA1$_{cat}$ and

PRIM2$_C$ structures. For the DT subcomplex, the refined POLA1$_{cat}$ model from the DE

subcomplex was docked as a rigid body into the high-resolution partial reconstruction. After

manual adjustment of the protein and de novo addition of the substrate, atomic coordinates and

isotropic temperature factors were refined as described above for the DI subcomplex. The

refined models of POLA1$_{cat}$ from the partial DT structure and PRIM2$_C$ from the complete DE

structure were then docked as rigid bodies into the reconstruction of DTS-1. The oligonucleotide

substrate from the partial DT subcomplex and the 3′-overhang and first nine base pairs from the

DE subcomplex were also docked as rigid bodies. Following manual modification of the protein

and de novo addition of the intervening region of the substrate, atomic coordinates and isotropic

temperature factors were refined as described above for the complete DE subcomplex.

Similarly, the refined model of POLA1$_{cat}$ from the partial DT structure, the oligonucleotide

substrate from the partial DT structure, and the 3′-overhang and first nine base pairs from the

complete DE structure were docked as rigid bodies into the reconstruction of DTS-2. Following

manual modification of the protein and de novo addition of the intervening region of the

substrate, atomic coordinates and isotropic temperature factors were refined as described

above for the complete DE subcomplex. For all structures, manual protein modification and de

novo substrate addition were performed in Coot [81], while docking and refinement were

performed in PHENIX [82]. If possible, rotamers were determined from the cryo-EM density. In

instances in which the correct conformation of the side chain was ambiguous, rotamers were

selected to match those in higher-resolution cryo-EM structures from this study or those in

previously published crystal structures. In instances in which both the correct conformation was

ambiguous and an appropriate reference structure was unavailable, rotamers were manually

selected to avoid steric clashes, and if possible, form favorable electrostatic interactions. Final

models were validated using MolProbity [83].

Structures of the complete polα–primase complex in configurations DI-1, DI-2, DE-1, and

DE-2 and of the TC subcomplex in configurations 1, 2, and 3 were generated from rigid bodies.

For all structures, the TC subcomplex was reconstituted from three rigid bodies taken from the

AI complex, while the DI and DE subcomplexes were each maintained as a single rigid body.

Docking into low-resolution reconstructions was performed in ChimeraX [84].

Refinement and validation statistics are provided in **Supplemental Figs. S7, S14, S15,**

S19, S20, and S29 and Supplemental Tables S1, S3, S4, and S6. Figures were prepared in

PyMOL (https://www.pymol.org), UCSF Chimera [85], or UCSF ChimeraX.

Primer elongation. To generate an RNA-primed template for elongation assays, 110 µM 5′-

triphosphorylated RNA primer and 100 µM DNA template were annealed by heating to 85°C

and then slowly cooling to 4°C. Reaction mixtures were then prepared with 50 nM POLA1$_{cat}$,

ΔPOLA1$_{cat}$, POLA1$_{cat}$/ΔPOLA1$_{cat}$, or polα–primase; 500 nM primer/template; 0.1 µM [α-^{32}P]-ATP

(3,000 Ci/mmol; PerkinElmer) or [α-^{32}P]-dATP (3,000 Ci/mmol; PerkinElmer); 9.9 µM ATP or

dATP; 200 µM GTP, CTP, and UTP; 20 µM dGTP, dCTP, and dTTP; 20 mM HEPES•NaOH pH

7.5; 150 mM NaCl; 5 mM MgCl$_2$; and 1 mM TCEP and incubated at 22°C. 8-µL aliquots were

periodically withdrawn, quenched with 2 µL of 100 mM EDTA pH 8.0, and incubated at 65°C for

15 min. To denature the primer and the template, 10 µL of loading buffer [80% (w/v) formamide,

10 mM EDTA pH 8.0, 1 mg/mL bromophenol blue, 1 mg/mL xylene cyanol, and 1.5 µM

competitor oligonucleotide] was added and samples were incubated at 85°C for 10 min and then

slowly cooled to 22°C. Elongation products were separated by denaturing urea-PAGE as

previously described [72], except 18% polyacrylamide (19:1 acrylamide/bisacrylamide) gels were

run at 20 W for 7 h and visualized by phosphorimaging using an Amersham Typhoon

biomolecular imager (Cytiva). Experiments were performed in triplicate.

Primer/template binding. Binding reactions were performed in 20 mM HEPES•NaOH pH 7.5,

150 mM NaCl, 5 mM MgCl$_2$, 1 mM TCEP, 20 µM dGTP, and 0.05% (v/v) Tween-20 with 10 nM

Cy5-labeled primer/template and 0.15–5,120 nM POLA1$_{cat}$. Immediately after mixing, samples

were centrifuged at 20,000×g and 4°C for 5 min, incubated on ice for 10 min, and then

incubated at 20°C for 10 min. Microscale thermophoresis data were collected using a Monolith

NT.115 instrument (NanoTemper). All experiments were performed using the red fluorescence

laser ($\lambda_{excitation}$ = 600–640 nm; $\lambda_{emission}$ = 660–720 nm) at 80% power and the infrared laser (λ =

1480 nm) at medium power. Temperature-related intensity changes were monitored at 20°C

with a pre-IR phase of 3 s, followed by an IR-on phase of 20 s, and a post-IR phase of 1 s. Data

quality was assessed by monitoring fluorescence quenching and photobleaching, as well as

protein aggregation and adsorption. Experiments were performed in triplicate. For each protein-

substrate binding titration, normalized fluorescence values [$F_{norm} = (F_{1\,(IR-hot)})/(F_{0\,(IR-cold)})$] were

tracked as parts per thousand and converted to ΔF_{norm} [$F_{norm\,(n)} - F_{norm\,(initial)}$]. ΔF_{norm} values for

individual isotherms were normalized to generate the fraction of bound protein (f_B). Dissociation

constants (K_d) were derived from fits to the data using a one-site binding model [$f_B = x/(K_d+x)$,

where x is the total concentration of POLA1$_{cat}$].

Data availability. Structures were deposited in the Protein Data Bank under accession codes

8G99 (AI complex, partial), 8G9F (AI complex, complete), 8G9L (DI subcomplex), 8G9N (DE

subcomplex, partial), 8G9O (DE subcomplex, complete), 8UCU (DT subcomplex, partial), 8UCV

(DT subcomplex 1), and 8UCW (DT subcomplex 2). Maps were deposited in the Electron

Microscopy Data Bank under accession codes EMD-29862 (AI complex, partial), EMD-29864

(AI complex, complete), EMD-29871 (DI subcomplex), EMD-29872 (DE subcomplex, partial),

EMD-29862 (DE subcomplex, complete), EMD-42140 (DT subcomplex, partial), EMD-42141

(DT subcomplex 1, complete), EMD-42142 (DT subcomplex 2, complete), EMD-29888 (TC

subcomplex, conformation 1), EMD-29889 (TC subcomplex, conformation 2), and EMD-29891

(TC subcomplex, conformation 3).

References

1.   Nethanel, T., Reisfeld, S., Dinter-Gottlieb, G. & Kaufmann, G. An Okazaki piece of simian virus
781   40 may be synthesized by ligation of shorter precursor chains. *J Virol* **62**, 2867-73 (1988).
782   2.   Bullock, P.A., Seo, Y.S. & Hurwitz, J. Initiation of simian virus 40 DNA synthesis in vitro. *Mol Cell*
783   *Biol* **11**, 2350-61 (1991).
784   3.   Eki, T., Matsumoto, T., Murakami, Y. & Hurwitz, J. The replication of DNA containing the simian
785   virus 40 origin by the monopolymerase and dipolymerase systems. *Journal of Biological*
786   *Chemistry* **267**, 7284-7294 (1992).
787   4.   Garg, P. & Burgers, P.M. DNA polymerases that propagate the eukaryotic DNA replication fork.
788   *Crit Rev Biochem Mol Biol* **40**, 115-28 (2005).
789   5.   Stodola, J.L. & Burgers, P.M. Mechanism of Lagging-Strand DNA Replication in Eukaryotes. *Adv*
790   *Exp Med Biol* **1042**, 117-133 (2017).
791   6.   Fukuda, M., Taguchi, T. & Ohashi, M. Age-dependent changes in DNA polymerase fidelity and
792   proofreading activity during cellular aging. *Mech Ageing Dev* **109**, 141-51 (1999).

7.   Levy, N. et al. XRCC1 interacts with the p58 subunit of DNA Pol alpha-primase and may
coordinate DNA repair and replication during S phase. *Nucleic Acids Res* **37**, 3177-88 (2009).
8.   Srivastava, V.K. & Busbee, D.L. Replicative enzymes and ageing: importance of DNA
polymerase alpha function to the events of cellular ageing. *Ageing Res Rev* **1**, 443-63 (2002).
9.   Taricani, L., Shanahan, F. & Parry, D. Replication stress activates DNA polymerase alpha-
associated Chk1. *Cell Cycle* **8**, 482-9 (2009).
10.   Yan, S. & Michael, W.M. TopBP1 and DNA polymerase alpha-mediated recruitment of the 9-1-1
complex to stalled replication forks: implications for a replication restart-based mechanism for
ATR checkpoint activation. *Cell Cycle* **8**, 2877-84 (2009).
11.   Kuchta, R.D., Reid, B. & Chang, L.M. DNA primase. Processivity and the primase to polymerase
alpha activity switch. *J Biol Chem* **265**, 16158-65 (1990).
12.   Sheaff, R.J. & Kuchta, R.D. Mechanism of calf thymus DNA primase: slow initiation, rapid
polymerization, and intelligent termination. *Biochemistry* **32**, 3027-37 (1993).
13.   Thomas, D.C. et al. Fidelity of mammalian DNA replication and replicative DNA polymerases.
Biochemistry **30**, 11751-9 (1991).
14.   Maga, G. et al. Okazaki fragment processing: modulation of the strand displacement activity of
DNA polymerase delta by the concerted action of replication protein A, proliferating cell nuclear
antigen, and flap endonuclease-1. *Proc Natl Acad Sci U S A* **98**, 14298-303 (2001).
15.   Pavlov, Y.I. et al. Evidence that errors made by DNA polymerase alpha are corrected by DNA
polymerase delta. *Curr Biol* **16**, 202-7 (2006).
16.   Rossi, M.L. & Bambara, R.A. Reconstituted Okazaki fragment processing indicates two pathways
of primer removal. *J Biol Chem* **281**, 26051-61 (2006).
17.   Stith, C.M., Sterling, J., Resnick, M.A., Gordenin, D.A. & Burgers, P.M. Flexibility of eukaryotic
Okazaki fragment maturation through regulated strand displacement synthesis. *J Biol Chem* **283**,
34129-40 (2008).
18.   Williams, J.S. & Kunkel, T.A. Ribonucleotides in DNA: origins, repair and consequences. *DNA*
Repair (Amst) **19**, 27-37 (2014).
19.   Burgers, P.M.J. & Kunkel, T.A. Eukaryotic DNA Replication Fork. *Annu Rev Biochem* **86**, 417-438
(2017).
20.   Pellegrini, L. The Pol alpha-primase complex. *Subcell Biochem* **62**, 157-69 (2012).
21.   Zerbe, L.K. & Kuchta, R.D. The p58 subunit of human DNA primase is important for primer
initiation, elongation, and counting. *Biochemistry* **41**, 4891-900 (2002).
22.   Lao-Sirieix, S.H., Nookala, R.K., Roversi, P., Bell, S.D. & Pellegrini, L. Structure of the
heterodimeric core primase. *Nat Struct Mol Biol* **12**, 1137-44 (2005).
23.   Vaithiyalingam, S., Warren, E.M., Eichman, B.F. & Chazin, W.J. Insights into eukaryotic DNA
priming from the structure and functional interactions of the 4Fe-4S cluster domain of human
DNA primase. *Proc Natl Acad Sci U S A* **107**, 13684-9 (2010).
24.   Sauguet, L., Klinge, S., Perera, R.L., Maman, J.D. & Pellegrini, L. Shared active site architecture
between the large subunit of eukaryotic primase and DNA photolyase. *PLoS One* **5**, e10083
(2010).
25.   Agarkar, V.B., Babayeva, N.D., Pavlov, Y.I. & Tahirov, T.H. Crystal structure of the C-terminal
domain of human DNA primase large subunit: implications for the mechanism of the primase-
polymerase alpha switch. *Cell Cycle* **10**, 926-31 (2011).
26.   Nunez-Ramirez, R. et al. Flexible tethering of primase and DNA Pol alpha in the eukaryotic
primosome. *Nucleic Acids Res* **39**, 8187-99 (2011).
27.   Kilkenny, M.L., Longo, M.A., Perera, R.L. & Pellegrini, L. Structures of human primase reveal
design of nucleotide elongation site and mode of Pol alpha tethering. *Proc Natl Acad Sci U S A*
110, 15961-6 (2013).
28.   Vaithiyalingam, S. et al. Insights into eukaryotic primer synthesis from structures of the p48
subunit of human DNA primase. *J Mol Biol* **426**, 558-69 (2014).
29.   Baranovskiy, A.G. et al. Crystal structure of the human primase. *J Biol Chem* **290**, 5635-46
(2015).
30.   Klinge, S., Nunez-Ramirez, R., Llorca, O. & Pellegrini, L. 3D architecture of DNA Pol alpha
reveals the functional core of multi-subunit replicative polymerases. *EMBO J* **28**, 1978-87 (2009).

31. Kilkenny, M.L., De Piccoli, G., Perera, R.L., Labib, K. & Pellegrini, L. A conserved motif in the C-
terminal tail of DNA polymerase alpha tethers primase to the eukaryotic replisome. *J Biol Chem*
287, 23740-7 (2012).
32. Perera, R.L. et al. Mechanism for priming DNA synthesis by yeast DNA polymerase alpha. *Elife*
2, e00482 (2013).
33. Attali, I., Botchan, M.R. & Berger, J.M. Structural Mechanisms for Replicating DNA in Eukaryotes.
Annual Review of Biochemistry **90**, 77-106 (2021).
34. Baranovskiy, A.G. et al. Mechanism of concerted RNA-DNA primer synthesis by the human
primosome. *J Biol Chem* **291**, 10006-10020 (2016).
35. Kilkenny, M.L. et al. Structural basis for the interaction of SARS-CoV-2 virulence factor nsp1 with
DNA polymerase α-primase. *Protein Sci* **31**, 333-344 (2022).
36. Jones, M.L., Aria, V., Baris, Y. & Yeeles, J.T.P. How Pol α-primase is targeted to replisomes to
prime eukaryotic DNA replication. *Molecular Cell* **83**, 2911-2924.e16 (2023).
37. He, Q. et al. Structures of the human CST-Polalpha-primase complex bound to telomere
templates. *Nature* (2022).
38. Klinge, S., Hirst, J., Maman, J.D., Krude, T. & Pellegrini, L. An iron-sulfur domain of the
eukaryotic primase is essential for RNA primer synthesis. *Nat Struct Mol Biol* **14**, 875-7 (2007).
39. Weiner, B.E. et al. An iron-sulfur cluster in the C-terminal domain of the p58 subunit of human
DNA primase. *J Biol Chem* **282**, 33444-33451 (2007).
40. Arezi, B., Kirk, B.W., Copeland, W.C. & Kuchta, R.D. Interactions of DNA with human DNA
primase monitored with photoactivatable cross-linking agents: implications for the role of the p58
subunit. *Biochemistry* **38**, 12899-907 (1999).
41. Baranovskiy, A.G. et al. Insight into the Human DNA Primase Interaction with Template-primer. *J*
Biol Chem (2016).
42. Xie, P. A model for dynamics of primer extension by eukaryotic DNA primase. *Eur Biophys J* **40**,
1157-65 (2011).
43. Thompson, H.C., Sheaff, R.J. & Kuchta, R.D. Interactions of calf thymus DNA polymerase alpha
with primer/templates. *Nucleic Acids Res* **23**, 4109-15 (1995).
44. Baranovskiy, A.G., Lisova, A.E., Morstadt, L.M., Babayeva, N.D. & Tahirov, T.H. Insight into
RNA-DNA primer length counting by human primosome. *Nucleic Acids Res* **50**, 6264-6270
(2022).
45. Coloma, J., Johnson, R.E., Prakash, L., Prakash, S. & Aggarwal, A.K. Human DNA polymerase
alpha in binary complex with a DNA:DNA template-primer. *Sci Rep* **6**, 23784 (2016).
46. He, Q. et al. Structures of human primosome elongation complexes. *Nature Structural &*
Molecular Biology **30**, 579-583 (2023).
47. Yuan, Z., Georgescu, R., Li, H. & O'Donnell, M.E. Molecular choreography of primer synthesis by
the eukaryotic Pol α-primase. *Nature Communications* **14**, 3697 (2023).
48. Cai, S.W. et al. Cryo-EM structure of the human CST-Polalpha/primase complex in a recruitment
state. *Nat Struct Mol Biol* **29**, 813-819 (2022).
49. He, Y. et al. Structure of Tetrahymena telomerase-bound CST with polymerase alpha-primase.
Nature (2022).
50. Zhang, Y., Baranovskiy, A.G., Tahirov, T.H. & Pavlov, Y.I. The C-terminal domain of the DNA
polymerase catalytic subunit regulates the primase and polymerase activities of the human DNA
polymerase α-primase complex. *J Biol Chem* **289**, 22021-34 (2014).
51. Zhang, Y., Baranovskiy, A.G., Tahirov, E.T., Tahirov, T.H. & Pavlov, Y.I. Divalent ions attenuate
DNA synthesis by human DNA polymerase alpha by changing the structure of the
template/primer or by perturbing the polymerase reaction. *DNA Repair (Amst)* **43**, 24-33 (2016).
52. Swan, M.K., Johnson, R.E., Prakash, L., Prakash, S. & Aggarwal, A.K. Structural basis of high-
fidelity DNA synthesis by yeast DNA polymerase delta. *Nat Struct Mol Biol* **16**, 979-86 (2009).
53. Hogg, M. et al. Structural basis for processive DNA synthesis by yeast DNA polymerase
varepsilon. *Nat Struct Mol Biol* **21**, 49-55 (2014).
54. El Hassan, M.A. & Calladine, C.R. Conformational characteristics of DNA: empirical
classifications and a hypothesis for the conformational behaviour of dinucleotide steps.
Philosophical Transactions of the Royal Society of London. Series A: Mathematical, Physical and
Engineering Sciences **355**, 43-100 (1997).

55. Lu, X.J. & Olson, W.K. 3DNA: a software package for the analysis, rebuilding and visualization of
three-dimensional nucleic acid structures. *Nucleic Acids Res* **31**, 5108-21 (2003).
56. Neidle, S. *Nucleic Acid Structure and Recognition*, (Oxford University Press, New York, 2002).
57. Lu, X.J., Shakked, Z. & Olson, W.K. A-form conformational motifs in ligand-bound DNA
structures. *J Mol Biol* **300**, 819-40 (2000).
58. Schneider, A. et al. Primase activity of human DNA polymerase alpha-primase. Divalent cations
stabilize the enzyme activity of the p48 subunit. *J Biol Chem* **273**, 21608-15 (1998).
59. Salay, L.E., Cordoba, J.J. & Chazin, W.J. DNA Recogntition/Processing | DNA Polymerase
Alpha-Primase: Biochemical and Structural Mechanisms. in *Encyclopedia of Biological Chemistry*
3rd edn Vol. III (ed. Jez, J.) 431-444 (2021).
60. Evrin, C., Maman, J.D., Diamante, A., Pellegrini, L. & Labib, K. Histone H2A-H2B binding by Pol
alpha in the eukaryotic replisome contributes to the maintenance of repressive chromatin. *EMBO*
J **37**(2018).
61. Li, Z. et al. DNA polymerase alpha interacts with H3-H4 and facilitates the transfer of parental
histones to lagging strands. *Sci Adv* **6**, eabb5820 (2020).
62. Sun, J. et al. The architecture of a eukaryotic replisome. *Nat Struct Mol Biol* **22**, 976-82 (2015).
63. Yuan, Z., Georgescu, R., Schauer, G.D., O'Donnell, M.E. & Li, H. Structure of the polymerase
epsilon holoenzyme and atomic model of the leading strand replisome. *Nat Commun* **11**, 3156
(2020).
64. Lue, N.F., Chan, J., Wright, W.E. & Hurwitz, J. The CDC13-STN1-TEN1 complex stimulates Pol α
activity by promoting RNA priming and primase-to-polymerase switch. *Nature Communications* **5**,
5762 (2014).
65. Zaug, A.J., Goodrich, K.J., Song, J.J., Sullivan, A.E. & Cech, T.R. Reconstitution of a telomeric
replicon organized by CST. *Nature* **608**, 819-825 (2022).
66. Lim, C.J. et al. The structure of human CST reveals a decameric assembly bound to telomeric
DNA. *Science* **368**, 1081-1085 (2020).
67. Braun, K.A., Lao, Y., He, Z., Ingles, C.J. & Wold, M.S. Role of protein-protein interactions in the
function of replication protein A (RPA): RPA modulates the activity of DNA polymerase alpha by
multiple mechanisms. *Biochemistry* **36**, 8443-54 (1997).
68. Abad, M.G., Rao, B.S. & Jackman, J.E. Template-dependent 3'-5' nucleotide addition is a shared
feature of tRNAHis guanylyltransferase enzymes from multiple domains of life. *Proc Natl Acad Sci*
U S A **107**, 674-9 (2010).
69. Abad, M.G. et al. A role for tRNA(His) guanylyltransferase (Thg1)-like proteins from Dictyostelium
discoideum in mitochondrial 5'-tRNA editing. *Rna* **17**, 613-23 (2011).
70. Rao, B.S., Mohammad, F., Gray, M.W. & Jackman, J.E. Absence of a universal element for
tRNAHis identity in Acanthamoeba castellanii. *Nucleic Acids Res* **41**, 1885-94 (2013).
71. Smith, B.A. & Jackman, J.E. Saccharomyces cerevisiae Thg1 uses 5'-pyrophosphate removal to
control addition of nucleotides to tRNA(His.). *Biochemistry* **53**, 1380-91 (2014).
72. Mullins, E.A. et al. An HPLC-tandem mass spectrometry method for simultaneous detection of
alkylated base excision repair products. *Methods* **64**, 59-66 (2013).
73. Mastronarde, D.N. Automated electron microscope tomography using robust prediction of
specimen movements. *J Struct Biol* **152**, 36-51 (2005).
74. Suloway, C. et al. Automated molecular microscopy: the new Leginon system. *J Struct Biol* **151**,
41-60 (2005).
75. Zheng, S.Q. et al. MotionCor2: anisotropic correction of beam-induced motion for improved cryo-
electron microscopy. *Nat Methods* **14**, 331-332 (2017).
76. Punjani, A., Rubinstein, J.L., Fleet, D.J. & Brubaker, M.A. cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296 (2017).
77. Bepler, T. et al. Positive-unlabeled convolutional neural networks for particle picking in cryo-
electron micrographs. *Nat Methods* **16**, 1153-1160 (2019).
78. Zivanov, J. et al. New tools for automated high-resolution cryo-EM structure determination in
RELION-3. *Elife* **7**(2018).
79. Sanchez-Garcia, R. et al. DeepEMhancer: a deep learning solution for cryo-EM volume post-
processing. *Commun Biol* **4**, 874 (2021).
80. Waterhouse, A. et al. SWISS-MODEL: homology modelling of protein structures and complexes.
Nucleic Acids Res **46**, W296-w303 (2018).

81.   Emsley, P., Lohkamp, B., Scott, W.G. & Cowtan, K. Features and development of Coot. *Acta*
959   *Crystallogr D Biol Crystallogr* **66**, 486-501 (2010).
960   82.   Adams, P.D. et al. PHENIX: a comprehensive Python-based system for macromolecular structure
961   solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-21 (2010).
962   83.   Davis, I.W. et al. MolProbity: all-atom contacts and structure validation for proteins and nucleic
963   acids. *Nucleic Acids Res* **35**, W375-83 (2007).
964   84.   Goddard, T.D. et al. UCSF ChimeraX: Meeting modern challenges in visualization and analysis.
965   *Protein Sci* **27**, 14-25 (2018).
966   85.   Pettersen, E.F. et al. UCSF Chimera--a visualization system for exploratory research and
967   analysis. *J Comput Chem* **25**, 1605-12 (2004).
968

Acknowledgments

This work was funded by National Institutes of Health grants R35GM136401 (B.F.E.), R35GM118089 (W.J.C.), R01GM087543 (J.E.J.), and PO1CA92584. We thank Miaw-Sheue Tsai and members of the Expression and Microbiology Core at UC Berkeley for baculovirus production and insect cell protein expression, and members of the Eichman and Chazin research groups for helpful discussions. Cryo-EM data were collected at the Center for Structural Biology Cryo-EM Facility at Vanderbilt University and the Life Sciences Institute Cryo-EM Facility at the University of Michigan. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Author contributions

B.F.E. conceived of the study; B.F.E. and W.J.C. oversaw the study; E.A.M., L.E.S., N.P.B., and B.F.E. designed experiments; J.E.J. provided reagents; E.A.M., L.E.S., C.L.D., and N.P.B. collected data; E.A.M., L.E.S., N.P.B., M.D.O., W.J.C., and B.F.E. analyzed data; E.A.M., L.E.S., N.P.B., W.J.C., and B.F.E. wrote the manuscript.

Competing interests

The authors declare no competing interests.

 Figure Legends

Figure 1. Chimeric primer synthesis. A. Stages of primer synthesis by polα–primase.

Triphosphate moieties are depicted as brown circles. Primase and polymerase α synthesize

RNA and DNA, respectively, to generate a short chimeric primer. **B**. Schematic of subunit and

domain organization in the heterotetrameric polα–primase complex.

Figure 2. Conformational and configurational dynamics. A. Characterization of purified

Xenopus laevis polα–primase by denaturing gel electrophoresis. Subunits are colored as in **Fig.**

1B (POLA1, blue; POLA2, cyan; PRIM1, dark green; PRIM2, light green). The POLA1 construct

lacks residues 1–334. **B**. Representative 2D class averages and corresponding subunit

schematics. The percentage of particles belonging to the AI complex or the TC subcomplex is

indicated below the class averages. Class averages were selected to show $PRIM2_C$ at various

degrees of association with the tetramer core. All final 2D class averages are shown in

Supplemental Fig. S6. **C**. 3D reconstruction of polα–primase in the auto-inhibitory

configuration. *Left*, cryo-EM density map. *Right*, molecular model refined against the density

map. $POLA1_{cat}$ is outlined in blue. **D**. Distinct conformations of the tetramer core. *Top*, cryo-EM

density maps. *Bottom*, molecular models docked into the density maps. Models were docked as

three rigid bodies extracted from the AI complex. The points at which the complex were split and

then reassembled are indicated with arrowheads. **E**. Schematic overlay aligned on the central

rigid body of the three tetramer core conformations shown in panel D. The range of motion

resulting from conformational dynamics is represented with curved arrows.

1011

Figure 3. Configurational rearrangements during DNA initiation and DNA elongation. In

each panel, DI and DE (sub)complexes are positioned on the left and the right, respectively. **A**.

Substrate schematics. Triphosphate moieties are depicted as brown circles. In both substrates,

the primers terminate in a 2′,3′-dideoxycytidine to prevent incorporation of dGTP. **B**.

Representative 2D class averages and corresponding subunit schematics. The percentage of

particles belonging to the DI or DE subcomplexes, the TC subcomplex, or one of the distinct

configurations of the DI and DE complexes (DI-1, DI-2, DE-1, or DE-2) is indicated below the

class averages. ~3% of particles from the DI dataset could not be reliably assigned to a group.

0% and 3% of particles from the DI and DE datasets, respectively, are consistent with the AI

complex. All final 2D class averages are shown in **Supplemental Figs. S11 and S13**. **C**. 3D

reconstructions and molecular models of $POLA1_{cat}$ and $PRIM2_C$ bound to the 3′- and 5′-ends of

the primers, respectively. The positions of the interdomain linkers are indicated with

arrowheads. **D,E**. Distinct configurations of the DI and DE complexes. Models were constructed

from four rigid bodies, comprised of the POLA1$_{cat}$/PRIM2$_C$ models shown in panel C and the

PRIM1/PRIM2$_N$/POLA1$_C$/POLA2$_C$ models shown in **Fig. 2**, and docked into the low-resolution

3D reconstructions shown in **Supplemental Figs. S19 and S20**. **D**. DI and DE complexes

oriented as in the 2D class averages. **E**. DI and DE complexes aligned on the tetramer core.

1029

Figure 4. Primer/template binding and elongation. A. Separation and detection of RNA and

RNA-DNA elongation products by denaturing gel electrophoresis and autoradiography.

Intermediate and doublet bands result from a mixture of RNA and DNA incorporation at

positions 7–10 (**Supplemental Fig. S22**). **B**. Substrate schematic. Triphosphate moieties are

depicted as brown circles. [α-^{32}P]-labelled ATP and [α-^{32}P]-labelled dATP are only incorporated

at position 7 in the primer. **C**. Quantitation of the 30-min time points from the gel in panel A. The

intensities of the elongation products are normalized to the intensity of the 30-mer. **D**. Kinetics of

30-mer formation. **E**. Binding of DNA/DNA, RNA/DNA, and RNA-DNA/DNA duplexes by

POLA1$_{cat}$. Data in panels D and E are presented as the mean ± SEM from three replicate

experiments.

1040

Figure 5. Substrate recognition and duplex remodeling. In each panel, the DI and DE

structures are positioned on the left and the right, respectively. **A**. Schematic of substrate-

binding contacts in POLA1$_{cat}$ (blue) and PRIM2$_C$ (green). **B**. Recognition of an incoming dGTP

by POLA1$_{cat}$. Phosphodiester bond formation is prevented by a 2′,3′-dideoxycytidine (ddC) at the

end of the primers. **C**. Recognition of RNA/DNA and DNA/DNA duplexes through substrate-

adaptive conformational changes in POLA1$_{cat}$. **D**. Recognition of distinct primer/template

conformations by PRIM2$_C$. **E**. RNA/DNA and RNA-DNA/DNA duplexes extracted from the DI

and DE subcomplexes. Major (M) and minor (m) grooves are indicated. Numbers to the left of

the duplexes indicate the base step starting from the 5′-end of the primers. **F**. Analysis of duplex

conformation. Z_P is the interstrand phosphate-phosphate distance after projection onto the

helical (Z) axis. The regions of the duplexes bound by POLA1$_{cat}$ (blue) and PRIM2$_C$ (green) are

shown below the plots.

1053

Figure 6. Configurational rearrangements upon DNA termination. A. Substrate schematic.

Triphosphate moieties are depicted as brown circles. The primer terminates in a 2′,3′-

dideoxycytidine to prevent incorporation of dGTP. **B**. Representative 2D class averages and

corresponding subunit schematics. The percentage of particles belonging to the DT

subcomplex, the TC subcomplex, or the AI complex is indicated below the class averages. No

2D class averages were observed with clear density for the complete DT complex. All final 2D

class averages are shown in **Supplemental Fig. S28**. **C**. Masked 3D reconstruction of POLA1$_{cat}$

bound to a full-length primer/template. **D**. 3D reconstructions and molecular models of POLA1$_{cat}$

bound to a full-length primer/template with (DT subcomplex 1, left) and without (DT subcomplex

2, right) simultaneous association of PRIM2$_C$. The positions of the interdomain linkers are

indicated with arrowheads. **E**. Analysis of duplex conformation. Z_P is the interstrand phosphate-

phosphate distance after projection onto the helical (Z) axis. The regions of the duplexes bound

by POLA1$_{cat}$ (blue) and PRIM2$_C$ (green) are shown below the plots. **F**. RNA-DNA/DNA duplexes

extracted from DT subcomplex 1 (left) and DT subcomplex 2 (right). Major (M) and minor (m)

grooves are indicated. Numbers to the left of the duplexes indicate the base step starting from

the 5′-end of the primers.

1070

Figure 7. Structural rearrangements during chimeric primer synthesis. A. Stages of RNA

and DNA synthesis. PRIM2$_C$ remains bound to the 5′-end of the primer during all stages of

synthesis. The transition from RNA elongation to DNA initiation occurs when the 3′-end of the

primer is transferred from PRIM1 to POLA1$_{cat}$. After generation of a full-length primer, DNA

elongation is terminated by dissociation of polα–primase from the primer/template. Protein-

protein contacts between PRIM2$_C$ and POLA1$_{cat}$ are represented by a dashed line. **B**. Subunit

schematics showing configurational rearrangements throughout the catalytic cycle of primer

synthesis. Schematics are aligned on the tetramer core (PRIM1/PRIM2$_N$/POLA1$_C$/POLA2$_C$). The

RNA initiation and RNA elongation configurations were derived from hypothetical models in

which the positions of POLA1$_{cat}$ are indeterminate and the positions of PRIM2$_C$ are dictated by

the geometric requirements of phosphodiester bond formation in the active site of PRIM1. The

DNA initiation and DNA elongation configurations correspond to DI-2 and DE-2, respectively

(**Fig. 3**). The DNA termination configuration was derived from a hypothetical model in which the

DE subcomplex in DE-2 was replaced by DT subcomplex 1 while retaining the position of

PRIM2$_C$. Loose association of POLA1$_{cat}$ and PRIM2$_C$ with the full-length primer/template is

indicated with wobble lines. At each stage of primer synthesis, the configuration depicted is

likely representative of a broad configurational/conformational distribution.

Figure 1

Figure 2

Figure 3

A

DNA   RNA

5′   (9 nt)   3′   dGTP

3′   (50 nt)   5′

5′   (9 nt)   (11 nt)   3′   dGTP

3′   (50 nt)   5′

B

| DI subcomplex | TC subcomplex | DI complex (DI-1) | DI complex (DI-2) | DE subcomplex | TC subcomplex | DE complex (DE-1) | DE complex (DE-2) |

Cartoon schematic

2D class average

POLA1cat   PRIM1   PRIM2C   PRIM2N   POLA1C   POLA2C

67%   26%   1%   ≤ 5%   48%   44%   1%   4%

C

POLA1cat   dGTP   5′   3′   5′   3′   PRIM2C   110°

D

DI-1   POLA1cat   PRIM2C   PRIM2N   POLA1C   PRIM1   POLA2C

DE-1   PRIM2C   PRIM2N   POLA1C   PRIM1   POLA2C   POLA1cat

DI-2   POLA1cat   POLA1C   PRIM2C   PRIM2N   PRIM1   POLA2C

DE-2   PRIM2C   POLA1C   PRIM2N   POLA1cat   PRIM1   POLA2C

DI complex   DE complex

E

DI-1   POLA1cat   PRIM2C   PRIM2N   POLA1C   PRIM1   POLA2C

DE-1   PRIM2N   POLA1C   PRIM2C   PRIM1   POLA1cat   POLA2C

DI-2   PRIM2N   PRIM2C   POLA1C   PRIM1   POLA2C   POLA1cat

DE-2   PRIM2N   POLA1cat   PRIM1   POLA1C   PRIM2C   POLA2C

DI complex   DE complex

Figure 4

Figure 5

Figure 6

Figure 7