Projection-specific integration of convergent thalamic and retrosplenial signals in the presubicular head direction cortex

Authors

Louis Richevaux[1]*, Dongkyun Lim[1], Mérie Nassar[1], Léa Dias Rodrigues[1], Constanze Mauthe[1], Ivan Cohen[3], Nathalie Sol-Foulon[2], Desdemona Fricker[1,4]*

[1]Université Paris Cité, CNRS, Integrative Neuroscience and Cognition Center, F-75006 Paris, France, [2]Sorbonne Université, INSERM, CNRS, Paris Brain Institute, ICM, Pitié-Salpêtrière Hospital, F-75013 Paris, France, [3]Sorbonne Université, INSERM, CNRS, Neuroscience Paris Seine, Institut de Biologie Paris Seine, F-75005 Paris, France, [4]Lead contact.

Correspondence: desdemona.fricker@parisdescartes.fr, louis.richevaux@parisdescartes.fr

Summary

Head-direction (HD) signals function as the brain's internal compass. They are organized as an attractor, and anchor to the environment via visual landmarks. Here we examine how thalamic HD signals and visual information from the retrosplenial cortex combine in the presubiculum. We find that monosynaptic excitatory connections from anterior thalamic nucleus and from retrosplenial cortex converge on single layer 3 pyramidal neurons in the dorsal portion of mouse presubiculum. Independent dual wavelength photostimulation of these inputs in slices leads to action potential generation preferentially for near-coincident inputs, indicating that layer 3 neurons can transmit a visually matched HD signal to medial entorhinal cortex. Layer 4 neurons, which innervate the lateral mammillary nucleus, form a second step in the association of HD and landmark signals. They receive little direct input from thalamic and retrosplenial axons. We show that layer 4 cells are excited di-synaptically, transforming regular spiking activity into bursts of action potentials, and that their firing is enhanced by cholinergic agonists. Thus, a coherent sense of orientation involves projection specific translaminar processing in the presubiculum, where neuromodulation facilitates landmark updating of HD signals in the lateral mammillary nucleus.

Keywords
Keywords: postsubiculum, presubiculum, head direction, multisensory, dendritic integration, circuit mechanism, patch clamp, brain slices, layer specific, projection specific

Introduction

The head direction (HD) system functions as the brain's compass system. It is distributed across several interconnected brain structures, and hierarchically organized from the brainstem to the lateral mammillary nucleus (LMN; Stackman and Taube, 1998), the anterior thalamic nuclei (ATN; Taube, 1995), and the dorsal Presubiculum (PrS; Ranck, 1984; Taube et al., 1990a). Modeling work suggests the HD system functions as a ring attractor, where the population of HD neurons is arranged in a one dimensional manifold (Blair and Sharp, 1995; McNaughton et al., 2006; Skaggs et al., 1995). The activity of thalamic and PrS HD neurons is correlated across different brain states, even during sleep, independently of sensory cues (Peyrache et al., 2015). The bump-like activity dynamics of an attractor network can maintain HD signals based on excitatory-excitatory or excitatory-inhibitory interactions (Knierim and Zhang, 2012; Simonnet et al., 2017). However, HD signals may drift in the dark (Taube et al., 1990b; Zugaro et al., 2003). For precise navigation, internally generated information on head direction must be combined with awareness of location in an environment. Mechanisms underlying the combination of egocentric head direction signals with anchoring to allocentric landmarks remain to be clarified.

Head direction and visual landmark signals may be integrated in the presubiculum (Jeffery et al., 2016; Yoder et al., 2019). Head direction cells are found in the superficial and deep layers of dorsal presubiculum, also termed postsubiculum (Boccara et al., 2010), and most layer 3 neurons in the dorsal presubiculum are head direction cells (Tukker et al., 2015; Preston-Ferrer et al., 2016). These cells receive monosynaptic head direction inputs from the anterior thalamus (Nassar et al., 2018; Peyrache et al., 2015; Balsamo et al., 2022). Lesions of the PrS impair the visual landmark control of a cell's preferred direction in ATN and in LMN (Goodridge and Taube, 1997; Yoder et al., 2015). Presubicular lesions also induce place field instability in the hippocampus (Calton et al., 2003), suggesting this region may be crucial to the anchoring of directionally modulated neurons to environmental landmarks.

Landmark based navigation depends on reliable visual cues. The PrS appears to receive direct projections from the primary visual cortex and indirect visual input via the retrosplenial cortex (Van Groen & Wyss 2003; Vogt & Miller 1983). The retrosplenial cortex encodes angular head velocity (Alexander & Nitz 2015; Keshavarzi et al. 2022) and visual landmark information (Auger et al., 2012; Clark et al., 2010; Sit and Goard, 2023), and supports an array of cognitive functions including memory, spatial and non-spatial context and navigation (Vann et al., 2009; Vedder et al., 2017). In particular, some neurons in the dysgranular retrosplenial cortex encode landmark-dominated head-direction signals (Jacob et al., 2017). Presubicular layer 3 cells receiving projections from both the anterior thalamus and the retrosplenial cortex (Kononenko and Witter 2012) could update the compass and bind landmarks to head direction signals. Presubicular neurons could then broadcast integrated HD-landmark signals directly to the MEC, via deep layers to the ADN and via layer 4 neurons to the LMN (Huang et al., 2017; Yoder et al., 2015; Yoder and Taube, 2011). The different target-specific presubicular projection neurons are well positioned to integrate anterior thalamic and retrosplenial inputs, but technical constraints have limited understanding of how known anatomical inputs are transformed into functional integrated output signals.

This work was therefore designed to examine the convergence of visual landmark
information, that is possibly integrated in the RSC, and vestibular based thalamic head
direction signals, in the PrS. Retrograde tracing was used to confirm inputs to the PrS from
the ATN and the RSC. The spatial distribution of ATN and RSC targets in the dorsal and
ventral PrS was investigated by stereotaxic injection of viral vectors inducing anterograde
expression of light-gated ion channels fused to fluorescent reporters. We found that
superficial layers of the dorsal PrS are major targets of ATN and RSC projections. We
analyzed functional convergence of these inputs in the PrS using dual-wavelength
optogenetic stimulations in ex vivo brain slices, while recording from layer 3 and 4 pyramidal
neurons. Both ATN and RSC projections made mono-synaptic excitatory connections with
single layer 3 principal cells and mostly di-synaptic ones with layer 4 cells. We show that
EPSPs induced with close to coincident timing by ATN and RSC fibers summed non-linearly in
layer 3 neurons. Layer 4 cell firing is facilitated by cholinergic activation. These data provide
insights into the integration of landmark and head direction inputs and their distribution to
downstream targets by PrS pyramidal cells.

Results

ATN and RSC send strong axonal projections to the dorsal Presubiculum

The origins of the main afferents projecting to the presubiculum were explored by
injecting retrogradely transported fluorescent beads into the PrS (Figure 1A). After 4 days,
coronal slices were prepared to examine the injection site and transport of the beads. Some
beads remained close to the PrS injection site, due in part to local projections. Strong bead
signals were detected within two of the subnuclei that form the anterior thalamus, the
anterodorsal (AD) and anteroventral (AV) part of ATN. Neurons in AD were labeled most
strongly in the medial portion of the AD, while labeled neurons in AV were found in its
lateral portion (Figure 1B). Many neurons in the RSC were labeled. Cells with somata in
layers 2 and 5 of the dysgranular dRSC were labeled, while mostly layer 5 cells of granular
gRSC contained beads (Figure 1C). Regions adjacent to the presubiculum including the
subiculum (Sub), parasubiculum (PaS) and the medial and lateral entorhinal cortices (MEC,
LEC) were labeled as was the contralateral PrS. Beads were also detected in the laterodorsal
thalamic nucleus (LD). Figure 1—figure supplement 1 shows an example of a series of
labeled coronal sections from one out of two mice, indicating some labeling also in the
visual cortices, perirhinal cortex, the nucleus reuniens of the thalamus, the dorsolateral
geniculate nucleus, and the claustrum. Some beads were observed in the CA1 region, which
has not previously been reported. Potentially though, this labeling could derive from a bead
leak into the nearby PaS which is innervated by CA1 (Van Groen et Wyss 1990a). In summary,
the ATN and RSC are the major sources of afferents projecting to the presubiculum with
lesser inputs from other sites.
Projections from the ATN or RSC to the presubiculum were explored by injecting at
these sites an anterogradely transported viral construct which expressed the modified
Channelrhodopsin Chronos fused to GFP (Figure 1D). After 4 weeks, horizontal slices were
prepared to verify the ATN injection site. Chronos-GFP labeling was mostly confined to the
ATN, occasionally extending to medial thalamic and reticular nuclei nearby, possibly as
bundles of projecting fibers (Figure 1E). After injection at the RSC site, coronal sections
(Figure 1F) showed strong expression in layer 5, especially deeper zones, and layer 6 of both

dysgranular and granular RSC, while labeled dendrites were evident in layer 1. Chronos-GFP
expression extended throughout the RSC, from -1.7 to -3.7 posterior to Bregma on the
antero-posterior axis.
Projections of the ATN to the presubiculum were examined in horizontal sections
(Figure 1G). ATN axons expressing the Chronos-GFP construct targeted layers 1 and 3 of the
presubiculum, mostly avoiding layer 2 (Simonnet et al., 2017; Nassar et al., 2018). Labeling
was precisely limited to the presubiculum with a sharp reduction of green fluorescence at
the posterior border with the parasubiculum, the anterior border with the subiculum and
the lateral border with deep layers. Axon terminals in layer 3 were not homogeneously
distributed. Patches of higher density were evident in deep layer 3 (Figure 1G, left panel) or
in the upper layer 3 (Figure 1G, forth panel, and Figure 1K). Along the dorso-ventral axis, ATN
axons projected from -2 to -4 mm ventral to Bregma. Fluorescence was not detected in
adjacent regions including the Sub, PaS, and DG. ATN axons avoided layer 2 of the PrS,
including regions with patches of calbindin positive neurons that project to contralateral PrS
(Preston-Ferrer et al. 2016; Figure 1K).
RSC projections to the presubiculum (Figure 1H), traced with the fluorescent
Chronos-GFP construct, also projected to layers 1 and 3. Unlike ATN axons, they were
restricted to dorsal regions of the presubiculum (from -2 to -3 mm ventrally to Bregma on
the dorso-ventral axis; Figure 1H, I, J). RSC axons also presented a patchy innervation pattern
in L1-3, and tended to avoid microdomains of high-density ATN projections (Figure 1G, K.
Figure 3B). Overall, these data show ATN and RSC afferents innervate overlapping dorsal
regions of the PrS, in layer 1 and parts of layer 3. Different spatial patterns of fiber
terminations suggest distinct local connectivities.
152

Layer 3 neurons are directly excited by both ATN and RSC afferents
We next examined physiological effects of ATN and RSC afferents on PrS pyramidal
cells (Rees et al., 2017). Responses to photo-stimulation of the Chronos-GFP construct
injected either in the thalamus or in the retrosplenial cortex (Figure 2A) were recorded in
layer 3 neurons (Figure 2B). These cells had robust regular firing patterns in response to
depolarizing current injections (Figure 2C). Intrinsic properties of layer 3 cells were rather
uniform (Figure 2—figure supplement 1) and an unsupervised cluster analysis suggested
they formed a single population (Figure 2D).
Blue light activation of either ATN or RSC axons expressing Chronos-GFP reliably
evoked synaptic responses in whole-cell patch-clamp records from layer 3 cells. High
intensity stimulation (1 - 3 mW) evoked EPSCs or EPSPs with a probability of 84% (70/83
neurons from 14 mice) for ATN fibers and 85% (104/123 neurons from 20 mice) for RSC
fibers (Figure 2E). Excitatory postsynaptic currents or potentials evoked by optical
stimulation of thalamic and retrosplenial inputs will be referred to as oEPSCs and oEPSPs.
Mean oEPSC amplitudes (Figure 2F) were similar for ATN (-305.0 ± 61.0 pA, n = 24) and RSC
fiber stimuli (-305.8 ± 59.1 pA, n = 27). Response latency (Figure 2G) to ATN axon stimulation
was 2.45 ± 0.19 ms (n = 24) and that for RSC axon stimulation was 2.93 ± 0.23 ms (n = 27).
The variability in latency of oEPSCs (Figure 2—figure supplement 2A, B) induced by RSC
afferent stimuli was 0.35 ± 0.09 ms (n = 27) and that for oEPSCs induced by ATN stimuli was
0.19 ± 0.06 ms (n = 24). These short oEPSC latencies suggest monosynaptic transmission. We
confirmed this point by showing that oEPSCs were maintained in presence of TTX (1μM) and
4AP (100μM) (Figure 2H, n = 2 and 12 for ATN and RSC respectively). EPSCs induced by
stimulation of ATN and RSC fibers were reduced in amplitude and decayed more quickly in

the presence of the NMDA receptor antagonist APV (50 µM) and were completely abolished by further application of the AMPA receptor antagonist NBQX (1 mM) (Figure 2I). The shape of oEPSCs induced by light stimulation of ATN and RSC fibers was comparable. Rise times were 1.38 ± 0.15 ms for ATN EPSCs (n = 15) and 1.38 ± 0.08 ms for RSC-mediated EPSCs (n = 15) and mean half-widths were 4.00 ± 1.02 ms for ATN-induced ESPCs (n = 15) and 4.34 ± 1.18 ms for RSC EPSCs (n = 15). The mean time constant (tau decay) for ATN-mediated EPSCs was 3.01 ± 0.21 (n = 15) and for RSC-induced EPSCs 3.17 ± 0.27 (n = 15) (Figure 2—figure supplement 2C-E.).

Optical stimulation of ATN and RSC fibers at lower light intensities elicited sub-threshold synaptic events in layer 3 cells. oEPSP amplitude for ATN mediated events was 4.09 ± 1.18 mV (n = 11) and for RSC events it was 4.80 ± 0.84 mV (n = 11). Maximal rising slopes were 2.93 ± 0.82 mV/ms for ATN-mediated events (n = 11) and 3.02 ± 0.53 mV/ms for RSC EPSPs (n = 11), and decay time constants were 81.4 ± 11.6 ms for ATN EPSPs (n = 11) and 86.9 ± 11.2 ms for RSC EPSPs (n = 11). Rise times were shorter for subthreshold EPSPs induced by ATN afferents at 3.00 ± 0.24 ms (n = 11) while those for RSC initiated EPSPs were 3.59 ± 0.34 (n = 11, p = 0.042, Wilcoxon test; Figure 2—figure supplement 2F-J). These data show that ATN and RSC axons make monosynaptic, glutamatergic excitatory contacts on layer 3 PrS cells.

The dynamics of responses to repetitive stimulation of layer 3 PrS cells were tested using 20 Hz trains of light stimuli to activate either ATN or RSC afferents. oEPSCs for both inputs followed depressing kinetics. Amplitude decreased for both inputs (Figure 2J, K), significantly after the fourth pulse (Friedman's test and Dunn's multiple comparison test). The *10/1* ratio was significantly lower than the paired-pulse ratio (PPR) in both cases (ATN *10/1* 0.56 ± 0.05 vs PPR 0.86 ± 0.05, n = 15, p < 0.0001, Wilcoxon test, RSC *10/1* 0.64 ± 0.05 vs PPR 0.83 ± 0.03, n = 15, Wilcoxon test) (Fig 2L). ATN fiber stimulation induced PrS cell firing in current-clamp with a high probability for initial stimuli and lower stable probabilities for later stimuli in the train. Similarly, 20 Hz stimulation of RSC afferents evoked spikes with high probability for initial stimuli and decreasing probabilities for later stimuli (on fourth and 7[th]-10[th] pulses, Friedman's and Dunn's multiple comparison tests; Figure 2M, N). Responses to repeated subthreshold stimuli had similar dynamics (Figure 2O). Stimulation with lower light intensities (< 1 mW) to initiate sub-threshold oEPSPs elicited different dynamic responses. ATN afferent trains elicited responses with depressing dynamics: the amplitude of the fifth oEPSP was significantly less than the first one (3.18 ± 0.88 vs 5.11 ± 1.64 mV, p = 0.0185, Friedman's and Dunn's tests). In contrast, subthreshold 20 Hz stimulation of RSC afferents evoked little or no oEPSP depression (Figure 2—figure supplement 2K-M). These data reveal distinct patterns of integration of repetitive activity in glutamatergic ATN and RSC afferents terminating on PrS layer 3 pyramidal cells.

Convergence of ATN and RSC inputs on single layer 3 neurons

Are single layer 3 pyramidal cells innervated by both ATN and RSC afferents? It seemed likely since the probability of connection to a given cell was 84% for ATN fibers and 85% for RSC afferents, and cluster analysis (Figure 2D) provided no evidence for subpopulations of layer 3 cells in terms of intrinsic electrophysiological properties (see also Balsamo et al., 2022).

We tested the hypothesis using photostimulation at two different light frequencies to activate ATN and RSC afferents independently. However this approach may be compromised since there is an overlap in the excitation spectra of blue-light activated Chronos (400-600

nm) and red-light activated Chrimson (450-700 nm; Klapoetke et al., 2014). Excitation of
Chronos with high intensities of blue light might also excite Chrimson. Stimulating with red
light at 630 nm should elicit neurotransmitter release in Chrimson-containing but not
Chronos-expressing fibers, which we found was indeed the case. Blue light stimuli at 470 nm
and intensities higher than 0.01 mW elicited oEPSPs in layer 3 cells of animals with Chronos-
expressing fibers. In contrast, blue light intensities higher than 0.25 mW were needed to
induce oEPSPs in different animals with Chrimson-expressing axons (Figure 3—figure
supplement 1A-D). Moreover, the amplitude of these events was smaller in Chrimson-
injected than in Chronos-injected mice. Thus, in our experimental conditions, blue light
intensities up to 0.25 mW could be used to excite Chronos-positive axons with confidence
that Chrimson-expressing fibers would not be stimulated.

We also tested the inverse, red light component of the dual stimulus strategy.
Intensities were tested in mice injected either with AAV5-Chronos to label ATN fibers or
AAV5-Chrimson to label RSC afferents. The strategy was validated by showing that red light
induced oEPSPs in Chrimson-containing, but not Chronos-expressing fibers, while blue light
evoked synaptic events in Chronos-, but not Chrimson-expressing fibers (Figure 3—figure
supplement 1E-K).

Injection in the same animal of the Chronos construction to the thalamus and that
for Chrimson to the retrosplenial cortex let us visualize ATN and RSC afferents to the
presubiculum (Figure 3A, B). Axon terminal fields of ATN and RSC projections were evident in
superficial layers of dorsal PrS (cf. Figure 1). Layer specificities, and labeling inhomogeneities
were apparent as were high density ATN patches below layer 2 with no RSC axon innervation
(Figure 3B). Layer 3 pyramidal cells could be innervated by ATN and RSC contacts both with
apical dendrites in layer 1 and basilar dendrites of layer 3 (Figure 3D). We found optical
stimulation of ATN and RSC axons, evoked synaptic responses in 14 out of 17 layer 3
pyramidal cells tested (current-clamp and voltage-clamp; Figure 3E, F).

In order to quantify substrates of these innervations, we counted numbers of labeled
ATN or RSC terminals located at less than 1μm from dendrites of biocytin filled layer 3
pyramidal cells (n = 6; Figure 3G). The distribution of these potential contact sites on
dendritic trees of layer 3 neurons was diverse. Some neurons exhibited highly segregated
domains receiving either ATN or RSC inputs. For instance, cell 1 had a high proportion of
clustered RSC putative synapses on its apical tuft while basilar dendrites were mostly
surrounded by potential thalamic terminals. In cell 2, RSC terminals were clustered close to
apical dendrites and also together with ATN terminals close to basilar dendrites. In contrast,
many ATN terminals were located closer to apical dendrites of cell 3. Segregation of cortical
RSC inputs to apical dendrites and thalamic ATN inputs to basal dendrites might favor
supralinear EPSP summation. However, afferent terminal distributions differed and we found
no clear pattern of sub- or supralinear summation (cell 1, supralinear summation
(Dual/summed EPSP, 1.39); cell 2, roughly linear (0.91); cell 3, sublinear (0.69), Figure 3G, H).
Furthermore, we noted certain potential contacts were located with separations less than 20
μm on the same dendrite (Figure 3I). Such proximity might underly local dendritic
computations and supra-linearities such as NMDA-mediated dendritic spikes which are
suggested to contribute to non-linear summation of synaptic events (Larkum et al., 1999,
2007, 2009).

Supralinear integration and amplification of ATN and RSC excitatory postsynaptic potentials

We explored interactions between excitatory inputs to layer 3 neurons by stimulating ATN and RSC axons separately with blue or red light. Excitatory synaptic potentials evoked by ATN or RSC axon stimulation, were compared to responses elicited when both sets of axons were activated at short time intervals (dual; Figure 4A). The mean amplitude of dual oEPSPs (synchronous light onset, Figure 4A-C) was significantly larger than calculated summed amplitudes of single oEPSPs (n = 11 cells, dual/summed amplitude 2.06 ± 0.55 mV, p = 0.0068, Wilcoxon test). Charge transfer, quantified as the surface under the dual oEPSP, was greater than the summed values for single oEPSPs (Dual/summed surface 2.21 ± 0.54 mV.s, n = 11, p = 0.0098, Wilcoxon test). These mean values cover variability between individual neurons. Summation of responses to ATN and RSC axon stimuli was supralinear in 7 out of 11 cells tested, linear for 3 cells and sublinear for 1 cell.

Dynamic responses to repetitive stimulations were also transformed by dual photostimulation. Amplitudes of 5 dual oEPSPs elicited at 20 Hz were higher than for a linear summation (Amplitudes in mV, Summed vs. Dual: Stim 1, 7.7 ± 1.9 vs. 11.1 ± 1.9; Stim 2, 6.7 ± 1.6 vs. 9.5 ± 1.5; Stim 3, 6.5 ± 1.4 vs. 9.5 ± 1.5; Stim 4, 6.1 ± 1.3 vs. 9.2 ± 1.5; Stim 5, 5.9 ± 1.3 vs. 8.6 ± 1.4, p = 0.0117, Two-way ANOVA; Figure 4E) and charge transfer, quantified as the surface under the 5 oEPSPs, also typically summed supralinearly (20 Hz, Surface in mV.s, Summed vs. Dual: Stim 1, 136.9 ± 30.6 vs. 221.2 ± 41.5; Stim 2, 143.3 ± 25.7 vs. 242 ± 43.9; Stim 3, 147.3 ± 22.5 vs. 259.2 ± 44; Stim 4, 145.1 ± 22 vs. 263.5 ± 49.2; Stim 5, 145.6 ± 23.2 vs. 258.7 ± 45.2, p = 0.0128, Two-way ANOVA; Figure 4F). Trains of dual ATN and RSC fiber stimulations also showed increased oEPSP integrals, compared to the stimulation of either one set of fibers alone, indicating stronger excitation over time (Figure 4E, F). Taken together, these data show supralinear summation of ATN and RSC inputs in most single layer 3 pyramidal cells, and also a facilitation of dynamic responses to repetitive inputs at 20 Hz.

We next asked how efficiently action potentials were induced by combined ATN and RSC inputs. Firing probability of layer 3 cells was higher for dual than for single input stimulation at a membrane potential of -65 mV (ATN 0.008 vs Dual 0.268, n = 5, p = 0.0216, RSC 0.024 vs Dual 0.268, n = 5, p = 0.1195, Friedman's and Dunn's tests) and higher still at -55 mV (Figure 4G, H). Firing was induced over a narrow window of time delays between stimulation of the two inputs (Figure 4I, J). It occurred for delays of -2 to +5 ms (RSC preceding ATN stimulus) reaching a maximum at a delay of +1 ms.

Supralinear summed responses of excitatory synaptic events may be mediated via the activation of NMDA receptors or voltage dependent intrinsic currents that amplify EPSPs (Fricker and Miles, 2000), especially if synaptic inhibition is reduced. We assessed these mechanisms in records made from layer 3 pyramidal neurons using a Cesium-based internal solution to favor cationic amplifying currents, and also containing QX314 to suppress action potentials (Figure 5). In 3 cells (from 2 mice) photostimulation of ATN fibers evoked oEPSPs of amplitude 10.9 ± 2.0 mV and RSC fiber stimulation induced events of 13.7 ± 2.5 mV. The response dynamics to 5 light pulses at 20 Hz were moderately facilitating (2nd vs 1st amplitude ratio: ATN 1.39, RSC 1.17; 5th vs 1st: ATN 1.18, RSC 1.20; n = 3). Coincident activation of ATN and RSC afferents induced supralinear oEPSP summation. Furthermore, repetitive stimulation generated large all-or-none depolarizations on the second or third stimulus at 20 Hz (2nd vs 1st amplitude ratio: 1.52; 5th vs 1st: 2.90; n = 3 cells, Figure 5A). It may be mediated by VGCC or a QX-314 resistant Na+ inward current (Fricker et al., 2009). This component appeared with higher probability and shorter latency in the presence of

GABA$_A$ receptor blocker gabazine (Figure 5B). The NMDA receptor antagonist APV largely
abolished the dual EPSP amplification initially (Figure 5C, black trace; 2nd vs 1st integral
ratio: 1.22; 5th vs 1st: 1.32), although amplification was partially restored by increasing RSC
stimulus intensity (pale pink trace; "amplification" refers to the shape of the plateau-like
prolongation of the peak, most pronounced on the second EPSP, indicated with an arrow).
NMDA receptor activation thus assists depolarization towards the threshold of a voltage-
dependent process contributing to supra-linear EPSP summation, but is not the only charge
carrier involved.
323

Cholinergic modulation and recruitment of presubicular layer 4 neurons by ATN and RSC
afferents

Presubicular layer 4 neurons are intrinsic bursting pyramidal neurons that project to
the lateral mammillary nucleus (Huang et al., 2017). This pathway is critical for the
coordinated stable landmark control of HD cells in the thalamus and throughout the HD cell
circuit (Yoder et al., 2015; Yoder et al., 2017) To investigate the input connectivity of layer 4
principal cells, we recorded responses of these neurons to stimulation of ATN and RSC
afferents.
Layer 4 neurons labeled by retrograde tracers injected in the lateral mammillary
nucleus were located in the *lamina dissecans* of the presubiculum, below layer 3, where
thalamic axons ramify (Figure 6A-D). At the proximal end of the presubiculum, and in its
continuity (Ishihara and Fukuda 2016), some subicular neurons projecting to the medial
portion of the mammillary bodies are also labeled (Figure 6B). The apical dendrites of layer 4
presubicular neurons extended towards presubicular layer 1 as previously described (Huang
et al., 2017), but tended to circumvent layer 3 and avoid thalamic afferents, by swerving
towards the subiculum. Apical dendrites of some neurons crossed layer 3 obliquely, while
others avoided thalamic afferents in layer 3 (Figure 6C, D, Figure 6—figure supplement 1).
Layer 4 neurons had a more depolarized resting membrane potential, lower input
resistance and time constant than layer 3 neurons (Figure 6—figure supplement 2 compares
active and passive properties). A characteristic voltage sag in responses to hyperpolarizing
steps, indicated the presence of an I$_h$ current. Layer 4 neurons discharged bursts of two or
three action potentials at the onset of a depolarizing step current injection and also after the
offset of hyperpolarizing steps (Figure 6E). These bursts were abolished by the T-type Ca^{2+}
channel blocker TTA-P2 (Figure 6F).
We next recorded responses of layer 4 neurons to optical activation of ATN or RSC
afferents together with effects of the same stimuli on layer 3 pyramidal cells (Figure 6G).
Overall, latencies of oEPSCs in layer 4 neurons were longer than for layer 3 (ATN layer 4, 6.2
± 0.6 ms, n = 8, vs. layer 3, 2.4 ± 0.2 ms, n = 24; RSC layer 4, 5.6 ± 0.4 ms, n = 9, vs. layer 3,
2.9 ± 0.2 ms, n = 27; Figure 6H), indicating possible polysynaptic excitation of layer 4
neurons. Bath application of TTX-4AP did not abolish oEPSPs entirely, leaving a low
amplitude component with short, potentially monosynaptic latencies (latency ATN 3.9 ± 0.6
ms, RSC 2.9 ± 0.2 ms, n = 5; Figure 6I).
Comparison of the timing of synaptic events and firing (Fig 7A-C) showed that oEPSP
onset in layer 4 neurons occurred after firing in the layer 3 neuron, following ATN afferent
stimuli, in 4 out of 5 cell pairs. We also observed this sequence when RSC input was
activated, in one tested pair. Depolarization of the layer 3 cell via the patch pipette, initiated
firing, but EPSPs were not elicited in any simultaneously recorded layer 4 neuron (0 out of six
cell pairs tested). These data suggest that excitation of layer 3 cells by ATN and likely by RSC

afferents is transmitted to layer 4 neurons, even in the absence of direct evidence for mono-
synaptic coupling between cell pairs.

In records from pairs of layer 3 and 4 neurons (n = 3; Figure 7D, E) layer 3 cells
responded with precisely timed action potentials to low intensity stimulation of ATN
afferents, while only very small oEPSPs were initiated in layer 4 neurons. Increasing the
excitatory drive by stimulating RSC afferents elicited larger oEPSPs more reliably in layer 4
neurons. Higher intensity stimulation of both ATN and RSC axons, could evoke bursts of
action potentials, with the activation of an underlying Ca^{2+} current in layer 4 cells (Figure 7E).
We hypothesize that increasing the activity of layer 3 neurons by strong (and non-specific)
stimulation of ATN and RSC afferents is needed to induce discharges in layer 4 cells. We
should also note that strong photostimulation of a single set of afferent fibers could initiate
discharges in layer 4 pyramidal cells, as was shown in records from mice where only one
afferent brain area, ATN or RSC, expressed an opsin (Figure 7—figure supplement 1).

Overall, it seems probable that layer 3 neurons relay activity projected by ATN and
RSC inputs onto layer 4 pyramidal neurons. However, evidence on this point could be
improved. In n = 9 paired recordings, we did not detect functional synapses between layer 3
and layer 4 neurons. Anatomically, putative synaptic contacts between filled axons of layer 3
and dendrites of layer 4 cells, were not evident (n = 5 filled layer 4 cells; see also Peng et al.,
2017). We did find a possible synaptic connection between two neighboring L4 neurons in
one case (Figure 6—figure supplement 1D), with a close apposition between axon and
dendrite. Recurrent excitation may promote bursting in a positive feedback loop in layer 4
neurons, and NMDA receptor related EPSP amplification favors burst firing (Figure 7F).
384
Neuromodulatory factors may help drive layer 4 cells to fire action potentials. As
Presubiculum is rich in acetylcholinesterase (Slomianka and Geneser 1991), we examined
excitability changes in the presence of a broad cholinergic agonist. The application of
Carbachol (10 µM; Figure 7G) led to a depolarization of the recorded layer 4 neurons'
membrane potential by 13 ± 4 mV, with increased action potential firing during a positive
step current injection, and from the baseline (Figure 7Gi,ii).
391
392

393 # Discussion

394
Understanding the anchoring of HD signals to environmental landmarks provides a
useful framework to ask how external information is integrated to revise and improve brain
representations. Here we show that thalamic ATN afferents signaling HD data and cortical
RSC projections that carry landmark information converge on layer 3 pyramidal cells of the
dorsal presubiculum. Independent optogenetic stimulation of these two glutamatergic
afferents in slices was used to define mechanisms of their integration in the presubiculum.
Most layer 3 cells were innervated by both ATN and RSC fibers, sometimes close on the same
dendritic branches, sometimes on different dendrites. Nearly coincident EPSPs (2-5 ms)
induced independently by ATN and RSC afferents, evoked non-linear membrane responses
to trigger layer 3 cell firing, transmitted to the MEC. In a second processing step, layer 3
neurons also excite layer 4 pyramidal cells which receive little direct ATN and RSC
innervation. These burst firing neurons may project a distinct, di-synaptically mediated,
visually-updated HD signal to the LMN. Our data thus suggest that the dorsal presubiculum

integrates HD and landmark signals producing two distinct output signals which are
transmitted to different regions. Cholinergic modulation may facilitate responses to salient
stimuli, for flexible anchoring to landmarks.
411
Anatomical convergence of ATN and RSC projections in the dorsal Presubiculum
Retrograde tracing in this study confirmed strong projections to the presubiculum
from the ATN and the RSC to the presubiculum (Figure 1). Anterior thalamic fibers
(Van Groen et Wyss 1990c; 1990a; 1990b; 1992; Shibata et Honda 2012; Vogt et Miller 1983)
ramify in and delimit the anatomical borders of the presubiculum (Liu et al., 2021; Simonnet
et al., 2017). HD signals from HD cells of the anterior thalamus project to the presubiculum
(Goodridge and Taube, 1997) where they form synapses directly with layer 3 pyramidal cells
(Nassar et al., 2018). The retrosplenial cortex was the most strongly labeled cortical region
innervating the presubiculum. We detected retrogradely transported beads in cells of layers
2 and 5 of dysgranular RSC, and layer 5 of the granular RSC across its antero-posterior axis
(Sugar and Witter, 2016).
We examined the anatomy and physiology of these two afferent systems to ask how
visual landmark signals from the RSC are combined with HD signals relayed via the ATN.
Anterograde fiber tracing, with an AAV5 construct expressing Chronos-GFP, confirmed that
axons from both regions project to superficial layers of the dorsal presubiculum. No RSC
projections were found in ventral PrS for injections in rostral RSC, as previously noted (Jones
and Witter 2007; Kononenko and Witter 2012). Both RSC and ATN innervated superficial
presubicular layers 1 and 3, while sparing layer 2 containing somata of calbindin positive
neurons (Figure 1K and cf. Balsamo et al. 2022). We detected microzones containing a high
density of ATN-positive fibers, but no RSC fibers, in upper layer 3. It remains to be shown
how these anatomical inhomogeneities may translate to functional modularity.
433
Pathway specific functional connectivity onto PrS layer 3 neurons.
Photostimulation of GFP-Chronos expressing axons let us compare synaptic events
initiated in layer 3 pyramidal cells by fibers from the ATN or the RSC in dorsal presubiculum
slices (Figure 2). Both ATN and RSC afferents formed mono-synaptic glutamatergic
connections with components mediated by NMDA and AMPA receptors. The amplitudes of
synaptic currents varied between neurons, with an overall similar amplitude distribution for
ATN and RSC stimulation. This may be due to variable expression levels (Hooks et al. 2015),
or may imply variations in coupling weights across cells. ATN and RSC fiber mediated EPSCs
depressed during repetitive activation at 20 Hz. EPSPs induced firing early during repetitive
stimulation. RSC inputs tended to produce more sustained EPSP dynamics and slower rise
times than ATN afferents for low intensity stimulation. ATN synapses with presubicular
pyramidal cells resemble those made by thalamic afferents in the somatosensory cortex -
high release with depressing dynamics, (Gil et al., 1999) possibly due to presynaptic
expression of VGLut2 (Liu et al., 2021).
Dual wavelength optogenetic stimulation was used for independent activation of
intermingled afferents expressing blue-light sensitive Chronos in ATN fibers and red-shifted
Chrimson in RSC fibers (Klapoetke et al., 2014). Precautions were taken to avoid cross-
stimulation since all channelrhodopsin variants are somewhat sensitive to blue light. With
the fast, sensitive opsin Chronos expressed in ATN fibers, synaptic events were induced by
very short (0.5 ms), low intensity, 0.25 mW, stimuli. For RSC fibers expressing Chrimson,
using light stimuli of duration 2 ms, adjusting intensity up to 2 mW initiated synaptic events

of comparable amplitude. Calibration experiments (Figure 3—figure supplement 1) provided
strong evidence for the independence of responses. While Chrimson has slower dynamics
than Chronos (Klapoetke et al., 2014), synaptic events induced by stimulating either opsin
were similar, as we showed by swapping the two opsin variants injected in ATN and RSC
fibers respectively (Figure 4).
This dual opsin approach permitted independent stimulation with blue and red light
pulses. Most (76%) recorded layer 3 neurons generated synaptic events in response to
stimulation of both ATN and RSC fibers providing a substrate for integration of landmark
information from the RSC with thalamic HD signals. Layer 3 cell firing was most effectively
triggered by nearly coincident inputs (-2 to +5 ms separation, Figure 4I, J). This is of interest
since HD representations in the AD and RSC are highly coherent (Fallahnezhad et al., 2023),
with short intervals between spikes in these two regions (< 5ms; van der Goes et al. 2022).
Converging inputs from thalamic and retrosplenial axons can therefore excite common
postsynaptic presubicular layer 3 neurons with very short delays, such that coincidence
detection by these cells will tend to enhance HD signals. Response dynamics to combined
stimulation of both inputs at 20 Hz were maintained or facilitating, in contrast to the
depressing dynamics of repetitive stimulation of one set of afferent fibers. Combined and
temporally precise inputs from ATN and RSC may thus help maintain HD signaling during
immobility. Our findings may well underlie the recent in vivo observation by Siegenthaler et
al (2024) that visual objects refine HD coding, and that presubicular HD cells whose
preferred firing direction corresponds to a visual landmark respond with higher firing rates.
476
Nonlinear signal integration in layer 3 thalamo-recipient neurons and physiological
significance for HD-to-landmark anchoring
Convergence of ATN and RSC axons onto single layer 3 pyramidal cells provides the
anatomical basis of synaptic integration. Putative synaptic contacts from both afferent fiber
systems were found on the basal dendrites of pyramidal cells (Figure 3), sometimes on the
same branch. Photostimulation centered on the soma of recorded neurons predominantly
activated synapses on basal dendrites. Supralinear summation could result from local spike-
generating mechanisms if the activated synapses were located on a same dendritic branch
(Makarov et al., 2023; Poirazi and Papoutsi, 2020; Polsky et al., 2004). Clustered synapses
could also guide the formation of new spines and synapses during integration of landmark
information in the HD signal. Learning might bind new inputs into functional synaptic
clusters (Hedrick et al., 2022). Our small sample of layer 3 pyramidal cells suggest that both
ATN and RSC axons target basal dendrites, and more RSC than ATN axons contact apical
dendrites (Figure 3). Tests on the effects of precise, near coincident activation of basal and
apical synapses could be revealing but were not technically possible in this study. Distinct
dendritic inputs to layer 3 cells may improve spatial perception (Takahashi et al., 2016)
enhancing HD signal quality by integration with landmark information.
Our data show that EPSPs elicited by nearly coincident ATN and RSC inputs which
exceed a threshold are amplified by NMDA receptor activation and voltage gated inward
dendritic currents (Figure 5; Fricker et al. 2009). From a small sample of cells it appears that
EPSP amplification may be facilitated by a reduction in synaptic inhibition (n = 3; Figure 5),
indicating that disinhibition may be permissive for supralinearity and gate firing by dynamic
modulation of the balance between inhibition and excitation (Milstein et al., 2015). VIP
expressing interneurons, which are excited by cholinergic modulation could provide such
disinhibition of the presubicular microcircuit (Porter et al., 1999; Slomianka and Geneser,

1991). The excitation-inhibition balance in pyramidal cells may become tipped towards excitation in the case of coincident, co-tuned thalamic and retrosplenial input. A 'when-to-learn' signal for HD updating in the presubiculum might function analogously to the promotion by dopamine of associations between sensory cues and head direction cells in the fly (Fisher et al., 2022). It is however unclear whether LTP type learning takes place in the mammalian head direction circuit. Presubicular layer 3 cells do not express the GluR1 subunit of AMPA receptors (Martin et al 1993; Ishihara et al., 2016) that is critical for LTP expression (Boehm et al., 2006). The absence of GluR1 might indicate that the thalamo-presubicular synapses in layer 3 function without classical long-term synaptic plasticity. Matching directional with visual landmark information based on temporal coincidence may be sufficient. Algorithms for dynamic control of cognitive maps without synaptic plasticity (Whittington et al 2025 Neuron) propose that working memory is stored in neural attractor activity, and updated by recurrent connections, which might generalize to the HD system.

Layer 3 pyramidal cells may be described as multi-compartmental computational devices (Häusser and Mel, 2003; Mel, 1993; Poirazi et al., 2003; Spruston, 2008) which integrate HD and landmark information. ATN axons drive presubicular HD neurons (Peyrache et al., 2015) and RSC mixed selectivity neurons contribute visual landmark information and allocentric spatial references (Jacob et al., 2017; Mitchell et al., 2018; Vann et al., 2009). NMDA mediated dendritic spikes enhance tuning selectivity in visual cortex (Smith et al., 2013; Wilson et al., 2016) and barrel cortex (Lavzin et al., 2012). Dendritic events in Layer 3 PrS cells may enable binding of visual landmarks with HD tuning. It is tempting to speculate that nonlinear synaptic integration and inhibitory gating may be involved in flexibly updating the allocentric direction of HD cells based on the integration of visual landmark information to the current HD signal (Siegenthaler et al., 2024). In the primary sensory cortex, nonlinearities act to increase perceptual threshold of sensory information (Takahashi et al., 2016; 2020). The attractor network in the PrS could thus be either stabilized or flexibly reset to external spatial cues.

Alternative models of HD-to-landmark anchoring have focused on the RSC, which also treats HD and landmark signals (Page & Jeffrey, 2018; Yan, Burgess, Bicanski, 2021). Possibly the RSC registers a direction to a landmark rather than comparing it with the current HD (Sit & Goard, 2023). The integrated information would then reach PrS, which in contrast to RSC, is uniquely positioned to update the signal in the LMN (Yoder et al., 2011).

What is the neuronal and circuit mechanism for integrating the "right" set of landmarks to anchor the HD signal? In our slice work we are blind to the exact nature of the signal that is carried by ATN and RSC axons. Critical missing information to understand the logic of HD-to-landmark anchoring also includes the degree of divergence and convergence of connections from a single thalamic or retrosplenial neuron. Do HD tuned inputs from thalamus converge on similarly tuned HD neurons only? Is divergence greater for the retrosplenial inputs? If so, thalamic input might pre-select a range of HD neurons, and converging RSC input might narrow down the precise HD neurons that become active (Figure 7—figure supplement 2). In the future, activity dependent labeling strategies might help to tie together information on the tuning of pre-synaptic neurons, and their convergence or divergence onto functionally defined postsynaptic target cells.

Functional significance of two-layer processing preceding projection to LMN

Burst-firing layer 4 PrS cells project a distinct version of HD-landmark signals to the LMN. The dendrites of these LMN projecting neurons overlap little with direct ATN and RSC

inputs granting partial isolation from direct excitation, and permitting a second level of
integration of information from layer 3 neurons. Layer 3 axons may innervate layer 4
neurons basal dendrites, which may be driven to fire by sufficiently strong excitatory inputs
from layer 3, especially if combined with cholinergic activation.
There are advantages to segregating the integration of converging input signals from
the updating signal across layers. Segregation permits both a fast transmission of an
integrated signal to the medial entorhinal cortex, and the conditional transmission of an
updating signal mediated by burst firing to upstream HD circuit elements. The synaptic
threshold implicit in this two-stage system permits a gated updating. Functionally layer 4
neurons are uniquely positioned to update the HD signal in the LMN with visual landmark
information (Yoder et al., 2015). In this way, cell-type specific cholinergic facilitation may
help idiothetic cue based navigation (Yoder et al., 2017). These cellular and synaptic circuit
data support and expand the findings of Yoder et al. (2015 and 2017). They clarify how the
thalamic HD signal integrated with visual landmark information is relayed to the grid cell
system in the medial entorhinal cortex and to the lateral mammillary nuclei.
564
Limitations
Burst firing may play a role for learning in hierarchical circuits (Naud et al., 2023;
Payeur et al., 2021). In the HD circuit, landmark information contained in bursts might reset
or anchor the HD attractor in the LMN and beyond. The effects of burst firing signals
transmitted to the LMN remain to be assessed. Potentially HD signals may be updated with
visual signals at several sites including the PrS, but PrS uniquely provides a feed-back
projection to the lateral mammillary nucleus. Further work on layer 3 to layer 4 transmission
is warranted, and the link to spatial perception and behavioural updating needs to be
strengthened.
574

575 # Methods

576
Animals
Experiments were performed on male and female wild-type and transgenic (Ai14
reporter line - Jax n007914) C57Bl6 mice of age 35-70 days, housed on a 12-hours light/dark
cycle with food and water available *ad libitum*. Animal care and use conformed to the
European Community Council Directive (2010/63/EU) and French law (87/848). Our study
was approved by the local ethics committee (CEEA - 34) and the French Ministry for
Research 01025.02.

584
Viral vectors and beads
Projecting neurons were labeled with retrograde fluorescent tracers (Retrobeads,
Lumafluor, and AAV2retro-CAG-tdTomato, Addgene 59462P). Fluorescent beads were stored
at 4°C before use. Channelrhodopsin expression was achieved by injecting Adeno-associated
viral constructions. AAV5.Syn.Chronos-GFP.WPRE.bGH (AAV5-Chronos, Penn Vector Core,
Addgene 59170P) was used to induce neuronal expression of the blue light induced
channelrhodopsin Chronos fused to the GFP marker and under the control of the Synapsin
promoter. AAV5.Syn.ChrimsonR-tdTomato.WPRE.bGH (AAV5-Chrimson, Penn Vector Core,
Addgene 59171P) induced neuronal expression of the red light gated channelrhodopsin
Chrimson fused to the tdTomato marker, under the control of the Synapsin promoter. Viral

vectors were stored at -80°C before use.
596

Stereotaxic surgery

Mice at ages of 4-5 weeks were anesthetized by intraperitoneal (i.p.) injection of a mixture of ketamine hydrochloride and xylazine (100 and 15 mg/kg respectively, in NaCl 0.9%). They were placed in a stereotaxic frame for injections. Fluorescent beads for retrograde tracing were injected (300-500 nl) into the PrS at the coordinates: -4.06 antero-posterior (AP), 2.00 medio-lateral (ML) and -2.15 mm dorso-ventral (DV) and into the LMN (-2.8 AP, 0.75 ML, -5.35 DV) with respect to the bregma. Note, due to the small size of LMN, some overflow to the medial mammillary nucleus (MMN) may have occurred.

Viral injections were performed unilaterally (Mathon et al., 2015; Richevaux et al., 2019) at the coordinates -0.82 AP, 0.75 ML and -3.2 mm DV for the ADN, and at -2.1 to -2.15 AP, 0.65 ML and -0.65 mm DV for the RSC. Volumes of 200 to 250 nl were injected with a 10 μL Hamilton syringe equipped with 33ga needle over a time of 10 min. The needle was slowly removed after a delay of 10 min to avoid leakage from the injection site. Best expression of AAV5 serotypes was achieved after 3 to 4 weeks.

Tissue fixation and slicing for retrograde tracing

Brains were removed for anatomy at 4 days after retrobead injection. Mice were anesthetized by i.p. injection of the ketamine/xylazine mixture. An intracardiac perfusion with 0.1M PBS was followed by perfusion with 4% paraformaldehyde. Brains were stored overnight in paraformaldehyde at 4°C and then washed in PBS. Coronal or horizontal sections were cut at 100 μm with a vibratome and stored in sucrose at 4°C.

Preparation of brain slices for physiology

Slices of the temporal lobe were prepared 3-4 weeks after injection of AAV5 viral constructions. Mice were anesthetized by i.p. injection of the ketamine/xylazine mixture. They were then perfused intracardially with a cutting solution containing (in mM): 125 NaCl, 25 sucrose, 2.5 KCl, 25 NaHCO$_3$, 1.25 NaH$_2$PO$_4$, 2.5 D-glucose, 0.1 CaCl$_2$, 7 MgCl$_2$, cooled to 4°C, and oxygenated with a 5% CO$_2$ / 95% O$_2$. The brain was removed and a vibratome was used to cut horizontal slices at 300 μm in the same solution. Slices were stored for 15 min at 34°C in an ACSF containing (in mM): 124 NaCl, 2.5 KCl, 26 NaHCO$_3$, 1 NaH$_2$PO$_4$, 2 CaCl$_2$, 2 MgCl$_2$, and 11 D-glucose, bubbled with 5% CO$_2$ / 95% O$_2$. They were then kept in the same solution at room temperature until recording.

Whole-cell patch-clamp recordings

Slices were transferred to a recording chamber perfused with oxygenated, warmed (~32 °C) ACSF mounted on an epifluorescence microscope. Patch-clamp records were made from neurons with borosilicate glass pipettes of external diameter 1.5 mm (Clark Capillary Glass, Harvard Apparatus) pulled with a Brown-Flaming electrode puller (Sutter Instruments). Electrodes, filled with a potassium-gluconate based solution containing (in mM): 135 K-gluconate, 1.2 KCl, 10 HEPES, 0.2 EGTA, 2 MgCl$_2$, 4 MgATP, 0.4 Tris-GTP and 10 Na$_2$-phosphocreatine, had a resistance of 4-8 MΩ. An alternative, cesium-gluconate based solution facilitated neuronal depolarization to examine synaptic inhibition, in Figure 5. It contained (in mM): 125 Cs-gluconate, 10 HEPES, 0.2 EGTA, 2 MgCl$_2$, 4 MgATP, 0.4 Tris-GTP and 10 Na$_2$-Phosphocreatine, together with 5 mM QX-314 to block Na$^+$ channels. Pipette solutions also contained 3 mM biocytin to reveal morphology after recording. They were

adjusted to pH 7.3 and osmolarity 290 mOsm. Whole-cell, patch-clamp signals were filtered
at 3kHz, amplified with a MultiClamp 700B amplifier and acquired with pCLAMP software
(Molecular Devices). In a subset of experiments, the following drugs were used to modulate
the responses to optogenetic stimulations; the presence of these drugs is indicated in the
figure and figure legend, whenever applicable. Monosynaptic excitation induced by optical
stimulation was tested in the presence of TTX (1 µM) and 4-AP (100 µM). NBQX (10 µM) and
APV (100 µM) were used to block AMPA and NMDA receptors respectively. Gabazine (10
µM) was used to block $GABA_A$ receptors. Carbachol (10 µM) was used to activate
acetylcholine receptors. All drugs were bath applied.
651
Optical stimulation
LED illumination (Cairn Research, OptoLED) was used to visualize the fluorescent
reporters GFP and tdTomato, and to stimulate opsin-expressing axons, using a 470 nm LED
for Chronos and a 627 nm LED for Chrimson. Illuminated spots had a diameter of 200 µm
with a 60x objective and were centered on the recorded cell soma. Photostimulation thus
covered most of the basilar dendrites of layer 3 pyramidal neurons (typical distance from tip
of apical to tip of basilar dendrite <400µm). Stimuli consisted of light pulses of 0.5 to 5 ms
duration, repeated 5-10 times at 20 Hz.
A multiband filter allowed simultaneous stimulation by blue and red LEDs of axons
containing Chronos and Chrimson (Simonnet et al., 2021). Stimulus power intensity (set in
mV) was calibrated as light intensity. The response probability of layer 3 cells was calibrated
to blue or red illumination of ATN or RSC afferents expressing Chronos or Chrimson to avoid
stimulus overlap (Figure 3—figure supplement 1). Chronos was targeted to the ATN and
Chrimson to the RSC after testing the reverse configuration. Projections from the thalamus
are larger and Chronos is more sensitive to blue light, so this configuration assured reliable
activation of thalamic fibers at minimal blue-light intensities. For experiments investigating
the integration of ATN and RSC inputs, we aimed to give similar weights to both inputs:
Chrimson-expressing fibers were stimulated with light of intensity adjusted to initiate
optically evoked excitatory postsynaptic potentials (oEPSPs) of amplitude similar to those
induced by blue light Chronos-fiber stimuli.
672
Data analysis
ATN and RSC projections to the PrS, were analyzed from Chronos and Chrimson
expression, using the ImageJ Plot Profile plug-in to quantitate normalized plot profiles (2000
pixels) of horizontal presubicular sections. Dorso-ventral differences were derived by
dividing differences in labeling intensity between the PrS and the dentate gyrus (DG)
molecular layer by DG intensity in slices from 5 dorso-ventral PrS levels. Values from all
animals were averaged and then normalized.
Cells were qualified as synaptically 'connected' when they responded to light
stimulation of afferent axons with a delay < 8 ms. Slices with very low or absent expression
of the fluorescent reporter were excluded. Cells were qualified as 'non-connected'
synaptically, if they did not respond to light stimulation, but at least one neighboring cell in
the same slice did.
Intrinsic neuronal properties were analyzed with custom MATLAB routines to derive
15 electrophysiological parameters (Huang et al., 2017). Parameters were standardized and
unsupervised cluster analysis performed with MATLAB was used to compare different
neurons (Huang et al., 2017; Simonnet et al., 2013).

Axograph was used to analyze responses to optical stimulation of ATN and RSC fibers.
Layer 3 neurons were recorded at potentials near -65 mV. Responses to light were averaged
from 10 stimulus trains at 20 Hz. Amplitudes and latencies of initial light evoked EPSCs, of
latency shorter than 10ms, were quantified from voltage-clamp records. Latency was
measured from stimulus onset to 10% of the peak amplitude of the first optically induced
EPSC. Amplitude was measured with respect to the pre-stimulus baseline. Paired-pulse ratio
(PPR) was defined as the ratio of the amplitude of the second to the first EPSC and 10/1 ratio
as that between the 10th and the 1st EPSC, in responses to 20 Hz stimulations. Spike
probability was counted over 5 to 10 trials per cell and then averaged over all cells. EPSPs
induced by dual wavelength stimulation were analyzed using Axograph and a custom-made
software. Events evoked by light stimuli were detected in a window of 1-10 ms after
stimulation. EPSP amplitude and integrated area were calculated over 50 ms after
stimulation. Baseline suppression was applied using an average of membrane potential
during 50 ms before stimulation. Summation of ATN and RSC evoked EPSCs in layer 3
neurons was determined from the amplitude and integral of averaged events. Summation
was considered to be supralinear, if values were more than 10% higher than a linear
addition, linear for values within ± 10%, and sublinear for values more than 10% lower.
706
Biocytin revelation and morphology
Recorded neurons were filled with biocytin to visualize their morphology and
location. Slices were fixed in 4% PFA in 0.1 M PBS at 4°C overnight, washed 3 times in 0.1
PBS and cryoprotected in 30% sucrose. Cell membranes were permeabilized by three freeze-
thaw cycles over dry ice and rinsed in PBS. Slices were agitated in a saturation buffer (2%
milk, 1% Triton X-100 in 0.1 M PBS) for 2 hours at room temperature. They were then
incubated with Streptavidin-Cy5 conjugate (1:500) and DAPI (1:1000) overnight at 4°C.
Sections were washed 3 times with PBS and mounted on coverslips with ProLong gold
antifade mountant. For anatomical study they were mounted in Mowiol medium. Cell, bead
and virus expression were visualized with an epifluorescence microscope. Higher resolution
of morphology was obtained with confocal microscopy. Cell morphologies were
reconstructed using IMARIS.
719
Quantification and statistical analysis
Data was analyzed with AxoGraphX and custom-written software (MATLAB, The
MathWorks). Results are given as mean ± SEM. Statistical analysis was performed with Prism
(GraphPad Software). The Mann-Whitney unpaired t-test was used to compare two groups.
The Wilcoxon or Kruskal-Wallis test was used to compare paired groups. Evolution of
parameters evoked by optical stimulation (current or potential amplitudes, spike probability)
was analyzed with Friedman's test followed by multiple comparison Dunn's test. Šidák's
multiple comparison test was also used to compare linear and observed responses to ATN
and RSC stimulations. Significance levels are given as p values.
729
730 **Data availability**
731   Data are available upon request to the corresponding authors Desdemona Fricker
732 and Louis Richevaux. The Matlab code for analysis of intrinsic properties is available at
733 _https://github.com/schoki0710/Intrinsic_Properties_.
734
735

Acknowledgements

This work is supported by the Centre National de la Recherche Scientifique and the Université Paris Cité. DF received funding from the Agence Nationale de la Recherche (ANR-DFG Program, ANR-18-CE92-0051 BURST), from the ERA-NET NEURON Program (ANR-20-NEUR-0005 VELOSO), and the FLAG-ERA HBP Program (ANR-21-HBPR-0002 VIPattract). This work has benefited from support by the BioMedTech Facilities at Université Paris Cité (Institut National de la Santé et de la Recherche Médicale Unité S36/Unité Mixte de Service 2009). We thank Dr. Li-Wen Huang for help with analysis routines. We thank Dr. Kate Jeffery and the Royal Society COST-share program, technical assistance from Fabrice Licata and Dr. Abdelali Jalil from the UPC Microscopy platform, Claire Lovo from ICM Quant, Dr. Boris Lamotte d'Incamps, and Dr. Richard Miles for helpful comments on the manuscript.

Author contributions

L.R. and D.F. conceived experiments, designed study, and interpreted data. D.F. acquired funding, L.R. and D.F. managed the project. L.R., D.L., M.N., C.M., L.D-R. and N.S-F. performed experiments and analyzed data, I.C. helped with analysis software, L.R. and D.L. prepared figures, L.R. and D.F. wrote the manuscript.

Declaration of Interests

The authors declare no competing financial interests.

Figure titles and legends

Figure 1 with 2 supplements.
ATN and RSC send strong axonal projections to the superficial layers of dorsal Presubiculum
A, Retrograde labeling of cortical and subcortical regions projecting to the presubiculum (PrS) with retrobeads.
B, Ipsilateral anterodorsal and anteroventral thalamic nuclei are labeled with beads.
C, Ipsilateral granular (gRSC) and dysgranular (dRSC) retrosplenial cortex labeling. Scale bars B, C, 200 µm.
D, Anterograde labeling of thalamic and retrosplenial projections to PrS with AAV-Chronos-GFP.
E, F, Injection sites in ATN (E) and RSC (F). AD: anterodorsal thalamic nucleus, AV: anteroventral thalamic nucleus, CA1: field of the hippocampus, cc: corpus callosum, cg: cingulum, CPu: caudate putamen, DG: dentate gyrus, MD: thalamic medial dorsal nucleus, MEC: medial entorhinal cortex, PaS: parasubiculum, PVA: paraventricular thalamic nucleus, anterior, Rt: thalamic reticular nucleus, sm: stria medullaris, Sub: subiculum, VL: thalamic ventrolateral nucleus.
G, H, Five sequential 100-µm horizontal slices of the parahippocampal region show ATN (G) and RSC (H) axons expressing Chronos-GFP (green). Dashed lines show limits of the parahippocampal region to the left and of the dentate gyrus to the right. Numbers show

dorsoventral level with respect to bregma. Scale bars 100 µm.

I, Normalized profiles of fluorescent intensity for ATN and RSC projections to the
presubiculum, from white matter (WM) to pia (PIA). Mean (green) ± SEM (grey), n = 8 slices
each.

J, Ventral (V) to dorsal (D) normalized distribution of ATN and RSC projections to the
presubiculum, n = 3 mice.

K, ATN axon labeling (green) is segregated from calbindin labeling (white) in the PrS. Scale
bar 100 µm.

See also Figure 1—figure supplement 1.

Figure 1—figure supplement 1. Brain regions providing input to the Presubiculum.
Semi-quantitative estimate of retrobeads labelling in serial coronal slices following injection
in the presubiculum. Color intensity represents beads density (mouse #69).

- Thalamic nuclei: anterodorsal (AD) +++, anteroventral (AV) +++, laterodorsal (LD)
+++, lateroposterior ++, dorsal lateral geniculate ++, reuniens ++, anteromedial -,
ventrolateral -
- Retrosplenial cortex: dysgranular (dRSC) +++, granular +++, dysgranular
contralateral ++
- Visual cortex: primary (V1) ++, secondary (V2L) +++
- Entorhinal cortex: lateral (LEnt) +++, medial (MEnt) +++
- Hippocampus/parahippocampus: parasubiculum (PaS) +++, subiculum (Sub) +++,
controlateral PrS ++, CA1 ++, CA3 -, dentate gyrus –
- Other: claustrum ++, perirhinal cortex ++, temporal association cortex ++

ADN: anterodorsal thalamic nucleus; AM: anteromedial thalamic nucleus; APT(V): anterior
pretectal nucleus (ventral part); AVDM: anteroventral thalamic nucleus, dorsomedial part;
AVVL: anteroventral thalamic nucleus, ventrolateral part; CA: cornu ammonis; cc: corpus
callosum; Cl: Claustrum; Cg1,2: cingular cortex area 1,2; coll. sup: colliculus superior; CPu:
caudate putamen (striatum); dhc: dorsal hippocampal commissure; DG: dentate gyrus; DLG:
dorsal lateral geniculate nucleus; LEnt: lateral entorhinal cortex; LDDM: laterodorsal thalamic
nucleus, dorsomedial part; LDVL: laterodorsal thalamic nucleus, ventrolateral part; LMN:
lateral mammillary nucleus; LS: lateral septal nucleus; M1,2: primary, secondary motor
cortex; MEnt: medial entorhinal cortex; MMN: medial mammillary nucleus; PAG:
periaqueductal gray; PaS: Parasubiculum; PF: parafascicular thalamic nucleus; PH: posterior
hypothalamic area; PMCo: posteromedial cortical amygdaloid nucleus; Po: posterior
thalamic nuclear group; PRh: perirhinal cortex; PrS: presubiculum; PVA: paraventricular
thalamic nucleus, anterior part; S1BF, S1FL, S1ULp, S1Tr: primary somatosensory cortex,
barrel field, forelimb region, upper lip region, trunk region; S2: secondary somatosensory
cortex; sm: stria medullaris of the thalamus; Sub: subiculum; Subst. Nigra: substantia nigra;
Re/Reuniens: reuniens thalamic nucleus; RSA = dRSC, dysgranular retrosplenial cortex; RSC:
retrosplenial cortex; RSG = gRSC: granular retrosplenial cortex; Rt: reticular thalamic nucleus;
V1: primary visual cortex; V2M: secondary visual cortex, medial part; V2L: secondary visual
cortex, lateral part; VA: ventral anterior thalamic nucleus; VDB: nucleus of the vertical limb
of the diagonal band; VL: ventrolateral thalamic nucleus; VPL: ventral posterolateral thalamic
nucleus; VPM: ventral posteromedial thalamic nucleus

Figure 1 - Supplement 2. Calbindin immunostaining and GFP-expressing ATN axons in
horizontal sections.
A, Subicular (Sub), Parasubicular (PaS), medial entorhinal cortical (MEC), and presubicular
(PrS) borders are outlined, as well as the dentate gyrus (DG). Patches of high density of ATN
axonal ramifications in L3 are indicated as "ATN microdomains".
B, Territories devoid of ATN axons in PrS L2 stain positive for calbindin.
831
Figure 2 with 2 supplements.
Responses of layer 3 presubicular cells to photoactivation of ATN and RSC fibers.
A, Expression of AAV5-Chronos in ATN or RSC.
B, Biocytin labeled layer 3 cells (white) and GFP positive axons (green) from the ATN or RSC.
Scale bar 10 μm.
C, Firing pattern of two layer 3 cells receiving inputs from the ATN (*cerulean*) and the RSC
(*purple*). Insets show current commands.
D, Cluster analysis of physiological parameters for cells tested by stimulating ATN (n = 29
cells /14 mice) or RSC (n = 43 cells/20 mice) fibers.
E, Representative EPSCs evoked in layer 3 cells by light stimulation (blue bar) of ATN or RSC
inputs. Right, proportion of cells receiving ATN or RSC inputs.
F, Average amplitudes of ATN and RSC induced synaptic currents. Mann-Whitney U test
revealed no significant difference (p = 0.95).
G, Latency of EPSCs evoked by light-stimulation of ATN (n = 24 cells/12 mice) or RSC fibers (n
= 27 cells/14 mice). Mann-Whitney U test revealed no significant difference (p = 0.13).
H, EPSPs induced in layer 3 cells (single traces) by stimulating ATN or RSC inputs in absence
and presence of 1 μM TTX and 100 μM 4-AP.
I, EPSCs induced by stimulating ATN or RSC fibers in the absence and presence of 100 μM
APV and APV + 10 μM NBQX. Holding potential +40 mV.
J, Voltage-clamp responses of layer 3 cells to 20 Hz train stimulations of ATN (*left*) and RSC
(*right*) inputs. Insets show EPSCs in response to the first two stimuli.
K, oEPSC amplitudes for 10 ATN or RSC fiber stimuli at 20 Hz (ATN, n = 15 cells/8 mice, RSC, n
= 15 cells/11 mice). Short middle line, mean; Min to max and quartiles.
L, Paired-pulse ratio (PPR) and 10/1 ratio (ratio between 10^{th} and 1^{st} event amplitudes) for
ATN or RSC inputs. Wilcoxon matched-pairs signed rank test: ATN PPR vs 10/1 **** p <
0.0001, RSC PPR vs 10/1 *** p = 0.0001.
M, Current clamp traces showing action potentials and EPSPs evoked by 10 stimuli at 20 Hz.
N, Spiking probability during trains of 10 stimuli. ATN, n = 6 cells/4 mice, RSC, n = 12 cells/9
mice. Full line, median; short line, mean; Min to max and quartiles. In K and N, * p < 0.05, **
p < 0.01, *** p < 0.001, **** p < 0.0001 from Friedman's test followed by Dunn's *post-hoc*
test.
O, oEPSP amplitudes for trains of 5 stimuli at 20 Hz.
See also Figure 2—figure supplements 1 and 2.
865
866
867 **Figure 2—figure supplement 1. Table showing the intrinsic electrophysiological**
868 **physiological properties of layer 3 cells tested for responses to stimulating ATN or RSC**
869 **fibers.**
870

Figure 2—figure supplement 2. Comparison of evoked synaptic events in presubicular layer
3 neurons following photostimulations of ATN or RSC axons.
A, Variance of latency (jitter) in oEPSP onset following Chronos-expressing ATN or RSC fiber
stimuli. RSC input stimulation gives similar variation in latency as ATN inputs. Mann-Whitney
U test revealed no significant difference ($p = 0.26$).
B, Jitter as a function of latency for ATN and RSC stimuli shows a similar coefficient of
variation.
C, Decay time constants of EPSCs evoked in layer 3 neurons by stimulating ATN or RSC inputs.
Mann-Whitney U test revealed no significant difference ($p = 0.87$).
D, E, First oEPSC amplitude as a function of oEPSC half-width (D) and oEPSC rise time (E) for
both inputs. We found no clear segregation of oEPSC shapes according to the origin of
afferent fibers.
F, oEPSPs in a presubicular layer 3 cell, induced by blue or red light stimulation of Chronos-
expressing ATN or Chrimson-expressing RSC afferents, respectively.
G-J, Comparison of oEPSPs evoked by stimulating ATN or RSC axons, recorded from a same
postsynaptic layer 3 presubicular neuron. G, Mean amplitude. H, Maximum rising slope of
oEPSPs. I, oEPSPs rise time. $p < 0.05$, Wilcoxon test. J, oEPSPs decay time constants. oEPSPs
following ATN and RSC fiber stimulation recorded from the same cell are connected by lines.
Dashed lines indicate reversed experimental design (expressing Chronos in RSC and
Chrimson in ATN).
K, Responses to 10 light stimuli at 20 Hz for ATN and RSC inputs to the same cell.
L, Amplitudes of oEPSPs evoked by trains of ATN (left) or RSC (right) fiber stimuli for 11 cells.
Dashed lines indicate the reversed experimental design (expressing Chronos in RSC and
Chrimson in ATN).
M, Summary data from L. On average, the amplitudes of oEPSPs evoked by ATN fiber
stimulation were smaller for the fifth than for the first stimulus. oEPSPs evoked by RSC fiber
stimulation were similar across five stimuli. Full line, median; Short line, mean; Min to max
and quartiles. * $p < 0.05$, Friedman's test and post-hoc Dunn's test comparing each stimulus
with the first.
900
901
Figure 3 with 1 supplement.
ATN and RSC axons converge in dorsal presubiculum and contact single layer 3 pyramidal
neurons.
A, Expression of the blue light sensitive opsin Chronos in ATN and the red light sensitive
opsin Chrimson in RSC.
B, Axons from ATN (GFP labeled, green) and RSC (tdTomato labeled, red) overlap in layer 1
and 3 of dorsal presubiculum. Layer 2 possesses patches of axon-dense and axon-poor
zones. RSC fibers avoid axon-dense microstructures formed by ATN fibers in upper layer 3.
Scale bar, 200 μm.
C, Independent dual wavelength optogenetic stimulation of light sensitive afferent fibers in
presubicular slices.
D, Two biocytin labeled PrS layer 3 pyramidal cells surrounded by ATN (green) and RSC (red)
axons.
E, Patch clamp recording from a layer 3 neuron shows optical EPSCs following
photostimulation of ATN axons (blue light) and RSC axons (red light).

F, 76% of layer 3 pyramidal neurons tested (n = 17 cells/4 mice) received both ATN and RSC
input.
G, Distributions of putative synaptic contacts from ATN (green) and RSC (red) on the
dendrites of three layer 3 neurons. Scale bar 50 μm. The boxed area on Cell 2 is shown in
panel I.
H, Normalized number of green and red spots for 6 neurons as a function of the distance
from soma. Paired values are indicated by dotted lines for the 3 cells in G.
I, Examples of ATN-labeled (left), RSC-labeled (middle) and both (right) synapses closely
apposed to dendrites of a biocytin-filled layer 3 pyramidal cell. Scale bar 20 μm. Insets show
representative high-magnification images. Scale bar 2 μm.
See also Figure 3—figure supplement 1.
928

Figure 3—figure supplement 1. Calibration of blue and red light stimulation of Chronos
and Chrimson.
A, Blue light stimulation of ATN Chronos-expressing fibers evoked oEPSPs in presubicular
layer 3 neurons at an intensity of 0.25mW, and 0.5 ms duration of illumination.
B, Blue light stimulation at the same intensity and duration did not evoke oEPSPs from RSC
Chrimson-expressing fibers.
C, D, Response probabilities of presubicular layer 3 cells to stimulation of fibers expressing
Chronos or Chrimson with red (C) or blue (D, left) light.
E, Experimental design. Injection of AAV5-Chronos-GFP (green) in the ATN, followed by
photostimulation combined with patch recording from a presubicular layer 3 cell.
F, Amplitudes of oEPSCs evoked in layer 3 cells by blue (0.5 ms, 0.25 mW) or red light (2 ms,
0.6 mW) stimuli.
G, Representative responses to light stimulation of ATN inputs.
H, Injection of AAV5-Chrimson-tdTomato (red) in the RSC, followed by photostimulation
combined with patch recording from a presubicular layer 3 cell.
I, Amplitudes of oEPSCs evoked in layer 3 cells by blue (0.5 ms, 0.25 mW) or red light (2 ms,
0.6 mW) stimuli.
J, Representative responses to light stimulation of RSC inputs
K, Values and statistical analysis for different stimulation configurations (Panels F and I).

Figure 4.
Supralinear summation of EPSPs and action potential firing following photostimulation of
ATN and RSC axons.
A, Optical EPSPs in layer 3 neurons in response to blue light activation of ATN axons
(cerulean traces), red light activation of RSC axons (purple traces). Supralinear summation of
EPSPs following coincident activation of both ATN and RSC axons (black traces). The pink
broken line indicates the calculated linear sum of oEPSPs evoked by stimulation of either
ATN or RSC axons. Records from cell 1, Fig 3D.
B, Amplitudes of dual ATN and RSC oEPSPs plotted as a function of the sum of separate ATN
and RSC stimulations (*left*). Each circle is a cell (n = 11 cells/4 mice), Pink line (± 10%)
indicates linearity. Pie charts give the number of tested layer 3 neurons with supralinear
(grey), linear (pink) or sublinear (white) summation. oEPSPs normalized to linear sum
(*bottom right*). Solid circle, Chronos in ATN/Chrimson in RSC, empty circle, Chronos in
RSC/Chrimson in ATN. * p < 0.05, ** p < 0.01, from Wilcoxon test.

C, As in B, for dual ATN and RSC oEPSP integrals.

D, oEPSPs induced by 20 Hz stimulation of ATN (cerulean), RSC (purple) or both (black). The
pink broken line indicates the calculated linear sum of oEPSPs evoked by separate ATN and
RSC photostimulation.

E, Amplitudes of dual oEPSPs compared to those of the sum of ATN and RSC oEPSPs show
that supralinearity increases across 5 stimulations. Two-way ANOVA, $p = 0.0117$. * $p < 0.05$,
**** $p < 0.0001$, Šidák's multiple comparison test. # $p < 0.05$, ## $p < 0.01$, Friedman's and
post-hoc Dunn's test.

F, As in E, for dual ATN and RSC oEPSP integrals. Two-way ANOVA, $p = 0.0128$.

G, Excitatory postsynaptic responses to photostimulation of ATN (blue light, cerulean traces)
or RSC axons (red light, purple traces) or both (blue and red light, black traces) at resting
membrane potential (-65 mV) or at a depolarized holding potential (-55 mV). Synaptic
excitation led to action potentials when dual ATN and RSC stimuli reached firing threshold.

H, Action potential (AP) probability for either or both stimuli at -65 and -55 mV. Data are
presented as mean ± SEM.

I, Action potentials were induced in presubicular layer 3 neurons by near coincident
activation of ATN axons (red light) and RSC axons (blue light). Time delays varied from -50 to
+50 ms.

J, Firing probability was maximal for short delays between -2 to +5 ms (RSC preceding ATN).

Figure 5. EPSP amplifications in layer 3 pyramidal neurons.

A, Optical EPSPs evoked in layer 3 pyramidal neurons following the stimulation of ATN axons
(blue light, cerulean traces), RSC axons (red light, purple traces) or both (blue and red light,
black traces). Recording pipette contained a cesium gluconate based internal solution and
the Na^+ channel blocker QX-314. A large all–or-none EPSP amplification occurred for dual
stimuli at 20 Hz, on some trials.

B, In the presence of the $GABA_A$ receptor antagonist gabazine (10 µM), dual EPSP
amplification occurred earlier in the train.

C, The additional presence of the NMDA receptor antagonist APV (100 µM) abolished dual
EPSP amplification (black trace). EPSP amplification was partially restored by increasing red
light intensity x2 (pale pink traces).

Figure 6 with 2 supplements.

Presubicular LMN-projecting layer 4 neurons avoid thalamo-recipient layer 3 and receive
little direct input from ATN and RSC.

A, Expression of Chronos-GFP in ATN and retrograde labeling of neurons that target LMN.

B, Thalamic axons (green) in superficial layers 1 and 3 of presubiculum. Retrobeads label cell
bodies of presubicular layer 4 cells (red).

C, Retrograde rAAV2-tdTomato label cell bodies and dendrites of layer 4 LMN projecting
neurons (red). Apical dendrites of layer 4 pyramidal neurons avoid layer 3 where thalamic
axons ramify.

D, Presubicular slice containing two layer 3 and two layer 4 neurons filled with biocytin
(white) and GFP expressing thalamic axons (green). Scale bar 100 µm. Inset, retrobeads (red)
in the soma of a biocytin filled LMN-projecting layer 4 neuron. Scale bar 10 µm.

E, Layer 3 neurons are regular spiking and layer 4 neurons are burst firing, initially and at
rebound, in reponse to current injection. Black trace, rheobase.

F, T-type Ca^{2+} channel blocker TTA-P2 (1 µM) suppressed burst firing in presubicular layer 4

neurons, while single action potentials were preserved.
G. Representative oEPSCs in layer 3 (left) and layer 4 (right) pyramidal cells, in response to
stimulation of ATN (cerulean) or RSC (purple) inputs.
H, oEPSC latencies in layer 3 and layer 4 cells, for ATN inputs (left, cerulean), or RSC inputs
(right, purple). Each dot is a cell. Same layer 4 cells are indicated by connecting lines, to
show the difference in latency for direct and indirect synaptic responses.
I, oEPSPs in layer 4 neurons in response to stimulation of ATN (cerulean) or RSC (purple)
inputs in control and in the presence of TTX (1 µM) and 4-AP (100 µM). Dashed lines indicate
the timing of the large disynaptic component of the responses (i, indirect), and the small
monosynaptic response (d, direct), isolated in TTX-4AP.
See also Figure 6—figure supplements 1 and 2.
1021
1022

Figure 6—figure supplement 1. Apical dendrites of presubicular layer 4 neurons avoid the
thalamorecipient layer 3.
A, Retrograde AAV2retro-tdTom was injected in LMN.
B, PrS L4 neuron cell bodies and dendrites were labeled. Scale bar, 50 µm. Full arrowheads,
L4 neurons apical dendrites crossing L3; Empty arrowheads, L4 neurons apical dendrites
skirting L3.
C, Anterograde AAV5-Chronos-GFP was injected in ATN to label thalamic axons, delimiting
PrS layer 3.
D, Examples of biocytin filled layer 4 presubicular neurons. (1) The apical dendrite of this
neuron avoided the densely packed ATN afferents and it didn't show responses to thalamic
stimulations. (2) This neuron's apical dendrite crossed through a short stretch of layer 3
containing ATN afferents, and it responded with low amplitude indirect events. (3) The apical
dendrite of this neuron crossed layer 3 and the zone containing ATN afferents, and was
directly recruited. Scale bar 10µm.
E, AAV-Chronos-GFP injected in ATN, and AAV-Chrimson-tdTom injected in RSC.
F, Image from Figure 7 panel D. Scale bar 100µm.
G, (Left) Enlarged view of dotted area in F, with two labelled L4 neurons. The apical dendrite
of top neuron makes a U-turn to ramify in deep layers and seems to cross path with the axon
of the bottom neuron, potentially forming a synaptic contact. (Right) The dendrite of the top
neuron and axon of bottom neuron are overlayed with pink and green lines respectively.
Scale bar 20µm.
1044

Figure 6—figure supplement 2. Electrophysiological passive and active intrinsic properties
of layer 3 vs. layer 4 neurons.
Data for layer 3 are summarized in Figure 2—figure supplement 1 and are shown here for
comparison. Statistical differences are indicated as levels of p values (* $p < 0.05$, ** $p < 0.01$,
*** $p < 0.001$, **** $p < 0.0001$) obtained from Mann-Whitney tests between layer 4 (n = 19)
and layer 3 (n = 33) for each parameter. Box plots indicate maximum and minimum values,
quartiles, full line is the median, short line is the mean.
1052

Figure 7 with 2 supplements.
Cross-laminar activation of LMN-projecting layer 4 bursting neurons.
A, Simultaneous records of a layer 4 and a layer 3 neuron during photostimulation of ATN
afferents. EPSP onset was delayed in the layer 4 neuron. Right panel, expanded view of the

boxed area.

B, Simultaneous records of a layer 4 and a layer 3 neuron during photostimulation of RSC
afferents. As in A, layer 4 neurons responded with a delay. Right panel, expanded view of the
boxed area.

C, Latencies of synaptic activation indicated by the dotted lines in A, B (ATN stimulation, n =
4 cells; RSC stimulation, n = 1). EPSP onset and AP peak from layer 3 neurons (n = 5). Dotted
lines link to the EPSP onset in simultaneously recorded layer 4 neurons (n = 5). $p < 0.05$,
Kruskal-Wallis multiple comparison test.

D, Biocytin labeled layer 3 and layer 4 pyramidal neurons, in a presubicular slice containing
thalamic (green) and retrosplenial (red) axons. The apical dendrite of one layer 4 neuron
makes a U-turn (arrowhead), away from layer 3, where the thalamic axons ramify. The
neighboring layer 4 neuron's apical dendrite crosses the thalamo-recipient layer 3 for a short
distance before arborizing outside of ATN targeted area, towards the subiculum, on the
right. Another biocytin filled layer 4 neuron's dendrites extend toward the deep layers. Scale
bar, 100µm.

E, Simultaneous records from a layer 3 and a layer 4 cell to ATN input stimulation (top, 0.25
mW, blue light), ATN and RSC input stimulation (middle, 0.25 mW blue and 0.25 mW red
light; light intensities compatible with independent photostimulation) or non-specific ATN
and RSC input stimulation (bottom, 3.25 mW blue and 0.25 mW red light). ATN fibers
expressed Chronos-GFP (green) and RSC fibers expressed Chrimson-tdTomato (red).

F, Top, oEPSPs and bursts of action potentials in a layer 4 neuron, evoked by dual wavelength
stimulation of ATN and RSC afferents at 20 Hz. Amplifications of dual oEPSPs led to firing.
Bottom, the NMDA receptor antagonist APV (100 µM) reduced EPSP amplification and
prevented action potential firing.

G. Layer 4 bursting neurons are sensitive to the acetylcholine receptor agonist carbachol (10
µM). Action potential firing in response to step current injections in control (black), in the
presence of carbachol (grey), and after wash-out (black). Bottom graph, membrane
potential depolarization during a 2-minute carbachol application. The number of action
potentials increased during the depolarizing steps (i) and on the baseline (ii).

See also Figure 7—figure supplement 1.

1087

1088

1089 **Figure 7—figure supplement 1. Regular firing layer 3 vs. intrinsically bursting layer 4**
1090 **neurons responded to high-intensity light stimulations of ATN or RSC afferents.**

1091 Layer 3 neurons respond with single spikes following each light stimulation while layer 4
1092 neurons respond with bursts of action potentials at the beginning of the stimulation train.

1093

Figure 7 – figure supplement 2. Schematic models for landmark anchoring of HD signals.

A, different scenarios with varying degrees of divergence and convergence of the two input
sources. We note that matching directional with visual landmark information does not
necessarily require synaptic plasticity, and might rely on temporal coincidence.

B, subcellular arrangements of synaptic contacts are important.

C, Inhibitory connectivity is important.

1100

1101

1102 # References

1103

Alexander AS, Nitz DA. 2015. Retrosplenial cortex maps the conjunction of internal and
external spaces. Nat Neurosci 18:1143–1151.

Auger SD, Mullally SL, Maguire EA. 2012. Retrosplenial Cortex Codes for Permanent
Landmarks. PLOS ONE 7:e43620.

Balsamo G, Blanco-Hernández E, Liang F, Naumann RK, Coletta S, Burgalossi A, Preston-
Ferrer P. 2022. Modular microcircuit organization of the presubicular head-direction
map. Cell Rep 39, 110684.

Blair HT, Sharp PE. 1995. Anticipatory head direction signals in anterior thalamus: evidence
for a thalamocortical circuit that integrates angular head motion to compute head
direction. J Neurosci 15:6260–6270.

Boccara CN, Sargolini F, Thoresen VH, Solstad T, Witter MP, Moser EI, Moser M-B. 2010. Grid
cells in pre- and parasubiculum. Nat Neurosci 13:987–994.

Boehm J, Kang M-G, Johnson RC, Esteban J, Huganir RL, Malinow R. 2006. Synaptic
incorporation of AMPA receptors during LTP is controlled by a PKC phosphorylation
site on GluR1. Neuron. 51:213–225.

Calton JL, Stackman RW, Goodridge JP, Archey WB, Dudchenko PA, Taube JS. 2003.
Hippocampal place cell instability after lesions of the head direction cell network. J
Neurosci 23:9719–9731.

Clark BJ, Bassett JP, Wang SS, Taube JS. 2010. Impaired Head Direction Cell Representation in
the Anterodorsal Thalamus after Lesions of the Retrosplenial Cortex. J Neurosci
30:5289–5302.

Fallahnezhad M, Le Mero J, Zenelaj X, Vincent J, Rochefort C, Rondi-Reig L. 2023. Cerebellar
control of a unitary head direction sense. Proc Natl Acad Sci U S A 120:e2214539120.

Fisher YE, Marquis M, D'Alessandro I, Wilson RI. 2022. Dopamine promotes head direction
plasticity during orienting movements. Nature 612:316–322.

Fricker D, Miles R. 2000. EPSP amplification and the precision of spike timing in hippocampal
neurons. Neuron. 28:559–569.

Fricker D, Dinocourt C, Eugène E, Wood JN, Wood J, Miles R. 2009. Pyramidal cells of rodent
presubiculum express a tetrodotoxin-insensitive Na+ current. J Physiol 587:4249–
4264.

Gil Z, Connors BW, Amitai Y. 1999. Efficacy of thalamocortical and intracortical synaptic
connections: quanta, innervation, and reliability. Neuron 23:385–397.

Goodridge JP, Taube JS. 1997. Interaction between the postsubiculum and anterior thalamus
in the generation of head direction cell activity. J Neurosci 17:9315–9330.

Häusser M, Mel B. 2003. Dendrites: bug or feature? Current Opinion in Neurobiology.
13:372–383.

Hedrick NG, Lu Z, Bushong E, Singhi S, Nguyen P, Magaña Y, Jilani S, Lim BK, Ellisman M,
Komiyama T. 2022. Learning binds new inputs into functional synaptic clusters via
spinogenesis. Nat Neurosci 25:726–737.

Hooks BM, Lin JY, Guo C, Svoboda K. 2015. Dual-channel circuit mapping reveals
sensorimotor convergence in the primary motor cortex. Journal of Neuroscience.
35:4418–4426.

Huang L-W, Simonnet J, Nassar M, Richevaux L, Lofredi R, Fricker D. 2017. Laminar
Localization and Projection-Specific Properties of Presubicular Neurons Targeting the
Lateral Mammillary Nucleus, Thalamus, or Medial Entorhinal Cortex. eneuro
4:ENEURO.0370-16.2017.

Ishihara Y, Fukuda T. 2016. Immunohistochemical investigation of the internal structure of
the mouse subiculum. NSC. 337:1–25.

Jacob P-Y, Casali G, Spieser L, Page H, Overington D, Jeffery K. 2017. An independent,
landmark-dominated head-direction signal in dysgranular retrosplenial cortex. Nat
Neurosci 20:173–175.

Jeffery KJ, Page HJI, Stringer SM. 2016. Optimal cue combination and landmark-stability
learning in the head direction system. J Physiol. Lond). 594:6527–6534.

Jones BF, Witter MP. 2007. Cingulate cortex projections to the parahippocampal region and
hippocampal formation in the rat. Hippocampus 17:957–976.

Keshavarzi S, Bracey EF, Faville RA, Campagner D, Tyson AL, Lenzi SC, Branco T, Margrie TW.
2022. Multisensory coding of angular head velocity in the retrosplenial cortex.
Neuron 110:532-543.e9.

Klapoetke NC et al. 2014. Independent optical excitation of distinct neural populations. Nat
Methods 11:338–346.

Knierim JJ, Zhang K. 2012. Attractor dynamics of spatially correlated neural activity in the
limbic system. Annu Rev Neurosci 35:267–285.

Kononenko NL, Witter MP. 2012. Presubiculum layer III conveys retrosplenial input to the
medial entorhinal cortex. Hippocampus 22:881–895.

Larkum ME, Zhu JJ, Sakmann B. 1999. A new cellular mechanism for coupling inputs arriving
at different cortical layers. Nature 398, 338–341.

Larkum ME, Waters J, Sakmann B, Helmchen F. 2007. Dendritic spikes in apical dendrites of
neocortical layer 2/3 pyramidal neurons. J. Neurosci. 27, 8999–9008.

Larkum ME, Nevian T, Sandler M, Polsky A, Schiller J. 2009. Synaptic Integration in Tuft
Dendrites of Layer 5 Pyramidal Neurons: A New Unifying Principle. Science 325, 756–
760.

Lavzin M, Rapoport S, Polsky A, Garion L, Schiller J. 2012. Nonlinear dendritic processing
determines angular tuning of barrel cortex neurons in vivo. Nature. 490:397–401.

Liu J, Kashima T, Morikawa S, Noguchi A, Ikegaya Y, Matsumoto N. 2021. Molecular
characterization of superficial layers of the presubiculum during development. Front
Neuroanat 15.

Makarov R, Pagkalos M, Poirazi P. 2023. Dendrites and Efficiency: Optimizing Performance
and Resource Utilization. https://arxiv.org/abs/2306.07101.

Martin LJ, Blackstone CD, Levey AI, Huganir RL, Price DL. 1993. AMPA glutamate receptor
subunits are differentially distributed in rat brain. NSC. 53:327–358.

Mathon B, Nassar M, Simonnet J, Le Duigou C, Clemenceau S, Miles R, Fricker D. 2015.
Increasing the effectiveness of intracerebral injections in adult and neonatal mice: a
neurosurgical point of view. Neurosci Bull 31:685–696.

McNaughton BL, Battaglia FP, Jensen O, Moser EI, Moser M-B. 2006. Path integration and the
neural basis of the "cognitive map." Nat Rev Neurosci 7:663–678.

Mel BW. Synaptic integration in an excitable dendritic tree. 1993. J Neurophysiol.
70(3):1086-101.

Milstein AD, Bloss EB, Apostolides PF, Vaidya SP, Dilly GA, Zemelman BV, Magee JC. 2015.
Inhibitory Gating of Input Comparison in the CA1 Microcircuit. Neuron 87:1274–
1289.

Mitchell AS, Czajkowski R, Zhang N, Jeffery KJ, Nelson AJD. 2018. Retrosplenial cortex and its
role in spatial cognition. Brain and Neuroscience Advances. 8:586.

Nassar M, Simonnet J, Huang L-W, Mathon B, Cohen I, Bendels MHK, Beraneck M, Miles R,
Fricker D. 2018. Anterior Thalamic Excitation and Feedforward Inhibition of
Presubicular Neurons Projecting to Medial Entorhinal Cortex. J Neurosci 38:6411–
6425.

Naud R, Friedenberger Z, Toth K. 2023. Silences, Spikes and Bursts: Three-Part Knot of the
Neural Code. arXiv:2302.07206v1. 1–15.

Page HJI, Jeffery KJ. Landmark-Based Updating of the Head Direction System by Retrosplenial
Cortex: A Computational Model. Front Cell Neurosci. 2018 Jul 13;12:191. doi:
10.3389/fncel.2018.00191.

Payeur A, Guerguiev J, Zenke F, Richards BA, Naud R. 2021. Burst-dependent synaptic
plasticity can coordinate learning in hierarchical circuits. Nat Neurosci 24:1010–1019.

Peng Y, Barreda Tomás FJ, Klisch C, Vida I, Geiger JRP. 2017. Layer-Specific Organization of
Local Excitatory and Inhibitory Synaptic Connectivity in the Rat Presubiculum. Cereb
Cortex. 27:2435-2452.

Peyrache A, Lacroix MM, Petersen PC, Buzsáki G. 2015. Internally organized mechanisms of
the head direction sense. Nat Neurosci 18:569–575.

Poirazi P, Brannon T, Mel BW. 2003. Arithmetic of subthreshold synaptic summation in a
model CA1 pyramidal cell. Neuron. 37(6):977-87.

Poirazi P, Papoutsi A. 2020. Illuminating dendritic function with computational models. Nat
Rev Neurosci 1–19.

Polsky A, Mel BW, Schiller J. 2004. Computational subunits in thin dendrites of pyramidal
cells. Nat Neurosci 7:621–627.

Porter JT, Cauli B, Tsuzuki K, Lambolez B, Rossier J, Audinat E. 1999. Selective excitation of
subtypes of neocortical interneurons by nicotinic receptors. J Neurosci 19:5228–
5235.

Preston-Ferrer P, Coletta S, Frey M, Burgalossi A. 2016. Anatomical organization of
presubicular head-direction circuits. eLife 5.

Ranck JB. 1984. Head direction cells in the deep layer of dorsal presubiculum in freely
moving rats. Soc Neurosci Abstr.

Rees CL, Moradi K, Ascoli GA. 2017. Weighing the Evidence in Peters' Rule: Does Neuronal
Morphology Predict Connectivity? Trends Neurosci 40:63–71.

Richevaux L, Schenberg L, Beraneck M, Fricker D. 2019. In Vivo Intracerebral Stereotaxic
Injections for Optogenetic Stimulation of Long-Range Inputs in Mouse Brain Slices.
JoVE J Vis Exp e59534.

Shibata H, Honda Y. 2012. Thalamocortical projections of the anterodorsal thalamic nucleus
in the rabbit. J Comp Neurol 520:2647–2656.

Siegenthaler D, Denny H, Skromne Carrasco S, Mayer JL, Levenstein D, Peyrache A, Trenholm
S, Macé E. 2024. Visual objects refine head direction coding. bioRxiv
2024.10.21.619417; doi: https://doi.org/10.1101/2024.10.21.619417

Simonnet J, Eugène E, Cohen I, Miles R, Fricker D. 2013. Cellular neuroanatomy of rat
presubiculum. Eur J Neurosci 37:583–597.

Simonnet J, Nassar M, Stella F, Cohen I, Mathon B, Boccara CN, Miles R, Fricker D. 2017.
Activity dependent feedback inhibition may maintain head direction signals in mouse
presubiculum. Nat Commun 8:16032.

Simonnet J, Richevaux L, Fricker D. 2021. Single or Double Patch-Clamp Recordings In Ex Vivo
Slice Preparation: Functional Connectivity, Synapse Dynamics, and Optogenetics.
Methods Mol Biol Clifton NJ 2188:285–309.

Sit KK, Goard MJ. 2023. Coregistration of heading to visual cues in retrosplenial cortex. Nat
Commun 14:1992.

Skaggs WE, Knierim JJ, Kudrimoti HS, McNaughton BL. 1995. A model of the neural basis of
the rat's sense of direction. Adv Neural Inf Process Syst 7:173–180.

Slomianka L, Geneser FA. 1991. Distribution of acetylcholinesterase in the hippocampal
region of the mouse: I. Entorhinal area, parasubiculum, retrosplenial area, and
presubiculum. J Comp Neurol 303:339–354.

Smith SL, Smith IT, Branco T, Häusser M. 2013. Dendritic spikes enhance stimulus selectivity
in cortical neurons in vivo. Nature. 503:115–120.

Spruston N. Pyramidal neurons: dendritic structure and synaptic integration. 2008. Nat Rev
Neurosci. 9(3):206-21.

Stackman RW, Taube JS. 1998. Firing properties of rat lateral mammillary single units: head
direction, head pitch, and angular head velocity. J Neurosci 18:9020–9037.

Sugar J, Witter MP. 2016. Postnatal development of retrosplenial projections to the
parahippocampal region of the rat. eLife 5.

Takahashi N, Oertner TG, Hegemann P, Larkum ME. 2016. Active cortical dendrites modulate
perception. Science 354:1587–1590.

Takahashi N, Ebner C, Sigl-Glöckner J, Moberg S, Nierwetberg S, Larkum ME. 2020. Active
dendritic currents gate descending cortical outputs in perception. Nat Neurosci.
54:677–679.

Taube JS. 1995. Head direction cells recorded in the anterior thalamic nuclei of freely moving
rats. J Neurosci 15:70–86.

Taube JS, Muller RU, Ranck JB. 1990a. Head-direction cells recorded from the postsubiculum
in freely moving rats. I. Description and quantitative analysis. J Neurosci 10:420–435.

Taube JS, Muller RU, Ranck JB. 1990b. Head-direction cells recorded from the postsubiculum
in freely moving rats. II. Effects of environmental manipulations. J Neurosci 10:436–
447.

Tukker JJ, Tang Q, Burgalossi A, Brecht M. 2015. Head-Directional Tuning and Theta
Modulation of Anatomically Identified Neurons in the Presubiculum. Journal of
Neuroscience. 35:15391–15395.

Van der Goes M-SH, Voigts J, Newman JP, Toloza EHS, Brown NJ, Murugan P, Harnett MT.
2022. Coordinated Head Direction Representations in Mouse Anterodorsal Thalamic

Nucleus and Retrosplenial Cortex. bioRxiv doi.org/10.1101/2022.08.20.504604.

Van Groen T, Wyss JM. 2003. Connections of the retrosplenial granular b cortex in the rat. J Comp Neurol 463:249–263.

Van Groen T, Wyss JM. 1992. Connections of the retrosplenial dysgranular cortex in the rat. J Comp Neurol 315:200–216.

Van Groen T, Wyss JM. 1990a. The connections of presubiculum and parasubiculum in the rat. Brain Res 518:227–243.

Van Groen T, Wyss JM. 1990b. Connections of the retrosplenial granular a cortex in the rat. J Comp Neurol 300:593–606.

Van Groen T, Wyss JM. 1990c. The postsubicular cortex in the rat: characterization of the fourth region of the subicular cortex and its connections. Brain Res 529:165–177.

Vann SD, Aggleton JP, Maguire EA. 2009. What does the retrosplenial cortex do? Nat Rev Neurosci. 10:792–802.

Vedder LC, Miller AMP, Harrison MB, Smith DM. 2017. Retrosplenial Cortical Neurons Encode Navigational Cues, Trajectories and Reward Locations During Goal Directed Navigation. Cereb Cortex. 2017; 27(7):3713-3723.

Vogt BA, Miller MW. 1983. Cortical connections between rat cingulate cortex and visual, motor, and postsubicular cortices. J Comp Neurol 216:192–210.

Whittington JCR, Dorrell W, Behrens TEJ, Ganguli S, El-Gaby M. 2025. A tale of two algorithms: Structured slots explain prefrontal sequence memory and are unified with hippocampal cognitive maps. Neuron, Volume 113, Issue 2, 321 - 333.e6.

Wilson DE, Whitney DE, Scholl B, Fitzpatrick D. 2016. Orientation selectivity and the functional clustering of synaptic inputs in primary visual cortex. Nat Neurosci. 19:1003–1009.

Yan Y, Burgess N, Bicanski A. A model of head direction and landmark coding in complex environments. PLoS Comput Biol. 2021 Sep 27;17(9):e1009434. doi: 10.1371/journal.pcbi.1009434.

Yoder RM, Peck JR, Taube JS. 2015. Visual landmark information gains control of the head direction signal at the lateral mammillary nuclei. J Neurosci 35:1354–1367.

Yoder RM, Taube JS. 2011. Projections to the anterodorsal thalamus and lateral mammillary nuclei arise from different cell populations within the postsubiculum: implications for the control of head direction cells. Hippocampus 21:1062–1073.

Yoder RM, Chan JHM, Taube JS. 2017. Acetylcholine Contributes to the Integration of Self-Movement Cues in Head Direction Cells. Behav Neurosci 131:312-324.

Yoder RM, Valerio S, Crego ACG, Clark BJ, Taube JS. 2019. Bilateral postsubiculum lesions impair visual and nonvisual homing performance in rats. Behav Neurosci 133:496–507.

Zugaro MB, Arleo A, Berthoz A, Wiener SI. 2003. Rapid spatial reorientation and head direction cells. J Neurosci 23:3478–3482.

A
AAV5-
Chrimson-
tdTomato
AAV5-
Chronos-
GFP

RSC
ATN
PrS

B
ATN axons
RSC axons
Merge

C
RSC
ATN
PrS

D
L1
L2
Cell 2
L3
Cell 1
L4
L3 P-cells
ATN axons
RSC axons

E
V_H = -65 mV
10 pA
10 ms

F
Connected cells (%)
100
50
0
76 %
Dual

G
Cell 1
Cell 2
Cell 3

H
Green spots
Red spots
Cell
1
2
3
Normalized number of spots
1.0
0.5
0.0
Distance from soma (µm)

I
ATN axons
RSC axons

A

CsGluc/QX-314

ATN
-65 mV

RSC
-65 mV

10 mV

200 ms

-65 mV

B

+ Gabazine

C

+ Gabazine + APV

RSC*2

A
Red beads   AAV5-Chronos-GFP
PrS   ATN
LMN

B
DAPI
Red beads
ATN axons

C
DAPI
rAAV
L3
L4

D
Biocytin
ATN axons
L3
L4
Biocytin
Red beads

E
Layer 3   Layer 4
-65mV
10 mV
100 ms

F
Layer 4
Control
TTA-P2
10 mV
100 ms
10 mV
10 ms

G
Layer 3   Layer 4
ATN
V_H = -65 mV
RSC
100 pA
5 ms

H
oEPSC latency (ms)
**** L3 L4i L4d
**** L3 L4i L4d

I
Control   Control
TTX-4AP   TTX-4AP
-65 mV
d i   d i
1 mV   2 mV
20 ms   20 ms

A ATN

L4
-65 mV

AP peak
EPSP onset | EPSP onset

2 mV
20 mV
5 ms

L3

50 ms

B RSC

1 mV
0.25 mV
10 mV
5 ms

50 ms

C

Event (ms)

L3_EPSP onset   L3_AP peak   L4_EPSP onset

D

Biocytin
RSC axons
ATN axons

L3

L4

E

L3
-65 mV

L4
-50 mV

20 mV
50 ms

0.25 mW   1 mV

0.25 mW
0.25 mW

20 mV

3.25 mW
0.25 mW

F

L4

-65 mV

10 mV
50 ms

+ APV

G

Control   Carbachol   Wash-out

75 pA

20 mV
800 ms

-40 mV

ΔVm (mV)

ctrl   carb   w-o

Time (min)

(i)

number of APs

**

ctrl   carb   w-o

(ii)

number of APs

**   **

ctrl   carb   w-o