Breaking the *C. elegans* invasion/proliferation dichotomy

Michael A. Q. Martinez[1], Chris Z. Zhao[1], Frances E. Q. Moore[1], Callista Yee[2], Wan Zhang[1],

Kang Shen[2], Benjamin L. Martin[1], David Q. Matus[1]

[1]Department of Biochemistry and Cell Biology, Stony Brook University, Stony Brook, NY 11794,

USA.

[2]Howard Hughes Medical Institute, Department of Biology, Stanford University, Stanford, CA

94305, USA.

*Author for correspondence (david.matus@stonybrook.edu)

KEY WORDS: *C. elegans*, invasion, proliferation, dichotomy, CKI-1

ABSTRACT

The acquisition of the post-mitotic state is crucial for the execution of many terminally

differentiated cell behaviors during organismal development. However, the mechanisms that

maintain the post-mitotic state in this context remain poorly understood. To gain insight into

these mechanisms, we used the genetically and visually accessible model of *C. elegans* anchor

cell (AC) invasion into the vulval epithelium. The AC is a terminally differentiated uterine cell that

must exit the cell cycle and enter a post-mitotic to initiate contact between the uterus and vulva

through a cell invasion event. Here, we set out to identify the negative cell cycle regulators that

maintain the AC in a post-mitotic, invasive state. Although our findings revealed a critical role for

CKI-1 (p21^{CIP1}/p27^{KIP1}) in maintaining the post-mitotic state of the AC, loss of CKI-1 alone or in

combination with other negative cell cycle regulators—including CKI-2 (p21^{CIP1}/p27^{KIP1}), LIN-35

(pRb/p107/p130), FZR-1 (Cdh1/Hct1), and LIN-23 (β-TrCP)—resulted in proliferating ACs that

retained their invasive abilities. Upon examination of the gene regulatory network controlling AC

invasion, we determined that proliferating, invasive ACs maintain pro-invasive gene expression.

We therefore report that maintenance of the post-mitotic state is not necessary for AC invasion,

breaking the previously established *C. elegans* invasion/proliferation dichotomy.

INTRODUCTION

The proper timing and coordination of terminal differentiation is critical for organismal development. In many cases, terminal differentiation requires cells to exit the cell cycle and enter a post-mitotic state (Soufi and Dalton, 2016). The general model for how terminally differentiated cells exit the cell cycle involves the repression of cyclin/CDK activity by negative cell cycle regulators such as CDK inhibitors (CKIs), retinoblastoma (Rb) family members, and E3 ubiquitin ligase complexes, including the anaphase-promoting complex/cyclosome (APC/C) and the Skp-Cullin-F-box protein (SCF) complex (Buttitta and Edgar, 2007; Ruijtenberg and van den Heuvel, 2016). Despite this generally accepted model, the mechanisms by which terminally differentiated cells maintain their post-mitotic state remain unclear.

Exit from the cell cycle following terminal differentiation is considered essential for the execution of specialized cell behaviors. According to the "go versus grow" dichotomy, a mutually exclusive relationship exists between these cell behaviors and cell proliferation (Kohrman and Matus, 2017). However, accumulating evidence has challenged this dichotomy in both development and cancer. For instance, in the developing mouse retina, genetic ablation of Rb paralogs leads to fully differentiated neurons that retain their ability to form neurites and synapses while proliferating (Ajioka et al., 2007). Similar results have been observed in mouse inner ear hair cells following loss of Rb (Sage et al., 2005). Furthermore, recent evidence from two-dimensional and three-dimensional *in vitro* assays using skin cancer cells found that these cells can simultaneously invade and proliferate (Haass et al., 2014; Vittadello et al., 2020).

Here, we used *C. elegans* anchor cell (AC) invasion as a genetically dissectible *in vivo* model of developmental cell invasion. Since the AC must exit the cell cycle and enter a post-mitotic state upon invasive differentiation (Deng et al., 2020; Matus et al., 2015; Medwig-Kinney et al., 2020; Smith et al., 2022), we aimed to identify the negative regulators of the AC cell cycle that maintain the post-mitotic state. While our findings reveal a critical role for the CDK inhibitor CKI-1 (p21^{CIP1}/p27^{KIP1}) in maintaining the post-mitotic state of the AC, loss of CKI-1 alone or in combination with other negative cell cycle regulators resulted in proliferating ACs that retained their invasive abilities. Using two reporters of the gene regulatory network that controls AC invasion, we show that proliferating, invasive ACs maintain pro-invasive gene expression. Thus, our work demonstrates that maintenance of the post-mitotic state is not required for AC invasion, breaking the invasion/proliferation dichotomy in *C. elegans*.

RESULTS AND DISCUSSION

Background

The *C. elegans* egg-laying system requires a direct connection between the uterus and the vulva. In the mid-L3 larval stage, a terminally differentiated post-mitotic uterine cell, called the anchor cell (AC), initiates the uterine-vulval connection by breaching the basement membrane (BM) that separates these tissues (Sherwood and Sternberg, 2003) (Fig. 1A, left). AC invasion occurs in synchrony with the divisions of P6.p, the underlying 1º-fated vulval precursor cell (VPC). By the P6.p 4-cell stage, BM dissolution is complete (Sherwood and Sternberg, 2003). Robust control of the AC cell cycle appears to be required for invasive differentiation, as loss of the post-mitotic state results in a complete loss of invasive capacity (Matus et al., 2015) (Fig. 1A, right). The SWI/SNF chromatin remodeling complex, specifically the evolutionarily conserved BAF assembly, along with three conserved transcription factors—EGL-43 (EVI1), HLH-2 (E/Daughterless), and NHR-67 (TLX/Tailless)—have all been implicated in the regulation of AC invasion because of their critical role in maintaining the post-mitotic state (Deng et al., 2020; Matus et al., 2015; Medwig-Kinney et al., 2020; Smith et al., 2022). However, the direct negative regulators of the AC cell cycle are largely unknown.

Uterine-specific depletion of CKI-1 rarely causes AC proliferation

In a genetic background where only the uterine cells are sensitive to RNAi (Hagedorn et al., 2009), loss of either EGL-43 or NHR-67 leads to AC proliferation without BM invasion in over 60% of animals (Matus et al., 2015; Medwig-Kinney et al., 2020). Conversely, uterine-specific RNAi-mediated depletion of CKI-1 ($p21^{CIP1}/p27^{KIP1}$), a canonical negative regulator of the *C. elegans* cell cycle (Feng et al., 1999; Fukuyama et al., 2003; Hong et al., 1998), rarely results in this phenotype, occurring in about 3% of animals (Matus et al., 2015).

In an attempt to increase the penetrance of the proliferative AC phenotype (≥2 AC), we first built a new RNAi construct for *cki-1* using the highly efficient RNAi vector, T444T (Sturm et al., 2018). Unlike the standard RNAi vector, L4440 (Timmons and Fire, 1998), T444T includes a pair of T7 terminator sequences to prevent transcription of the vector backbone. We then tested the T444T-based *cki-1(RNAi)* construct in a uterine-specific RNAi strain (*rrf-3(pk1426); rde-*

1(ne219); fos-1ap::rde-1) with fluorescent reporters to visualize the AC cell membrane and BM.

Though 5% of animals exposed to this construct had ≥2 ACs (Fig. S1A), this result only

represents a 2% increase in the proliferative AC phenotype when compared to the original *cki-*

1(RNAi) experiment (Matus et al., 2015). In each of these animals, BM invasion was observed.

This finding could be attributed to either a late RNAi effect or reestablishment of the post-mitotic,

invasive state following AC proliferation. Alternatively, this result raises the intriguing possibility

that ACs can simultaneously invade and proliferate, which we explore in the subsections below.

Systemic depletion of CKI-1 causes a moderate increase in AC proliferation

To better visualize the dichotomy between AC invasion and proliferation, we generated a strain

for three-color imaging of the AC, cell cycle, and BM (Fig. 1A-C). Specifically, this strain

contains an endogenous transcriptional reporter of *lag-2* (*Delta*), LAG-

2::P2A::H2B::mTurquoise2, to mark the AC nucleus (Medwig-Kinney et al., 2022), a ubiquitous

CDK activity sensor, DHB::2xmKate2, to visualize and quantify cell cycle state (Adikes et al.,

2020), and an endogenous translational reporter of *lam-2*, LAM-2::mNeonGreen, to label

laminin, a core BM component (Jayadev et al., 2019). Conveniently, the histone-based *lag-2*

transcriptional reporter also functions as a P6.p lineage tracker, facilitating the developmental

staging of AC invasion (Fig. 1A,C). The CDK activity sensor is a fluorescent fragment of human

DNA helicase B (DHB), which is predominantly nuclear in post-mitotic cells, such as the wild-

type AC, and becomes increasingly cytoplasmic in proliferating cells as they progress through

the cell cycle (Fig. 1B,C). Thus, the ratio cytoplasmic-to-nuclear of DHB fluorescence provides a

visual and quantitative readout of CDK activity, and more generally, cell cycle progression

(Adikes et al., 2020; Martinez and Matus, 2022; Spencer et al., 2013; van Rijnberk et al., 2017).

With the three-color imaging strain at our disposal, we first wanted to determine the penetrance

of the proliferative AC phenotype following RNAi-mediated depletion of CKI-1 in a wild-type

background. Unexpectedly, 100% of *control(RNAi)* and *cki-1(RNAi)* animals showed a post-

mitotic, invasive AC (Fig. S1B,C; Table 1). We next introduced the *rrf-3(pk1426)* null allele

(Simmer et al., 2002), which on its own confers RNAi hypersensitivity in many somatic tissues,

including the uterus (Matus et al., 2010). While nearly all *rrf-3(pk1426); control(RNAi)* animals

contained a post-mitotic, invasive AC (Fig. 1D,E), there was one animal where a BM gap was

not observed below a post-mitotic AC (Table 1). In comparison, 24% of *rrf-3(pk1426); cki-*

1(RNAi) animals possessed ≥2 ACs, which correlated with a significant increase in AC-specific

CDK activity when compared to *rrf-3(pk1426); control(RNAi)* animals (0.25 ± 0.06 (n = 30) vs.

0.73 ± 0.56 (n = 55)). In almost half of those animals with ≥2 ACs, BM invasion was observed

(Table 1). Taken together, these data demonstrate that systemic depletion of CKI-1 enhances

the penetrance of the proliferative AC phenotype.

Unfortunately, we were unable to assess the effect of the *cki-1(gk132)* null allele on AC

proliferation (The *C. elegans* Deletion Mutant Consortium, 2012), as it leads to embryonic

lethality in 100% of homozygotes (Buck et al., 2009) (Fig. S2A). Instead, we crossed the *cdc-*

14(he141) null allele into the three-color imaging strain (Saito et al., 2004). *cdc-14* encodes a

dual-specificity phosphatase that positively regulates CKI-1 activity in a number of larval tissues

(Clayton et al., 2008; Roy et al., 2011; Roy et al., 2014; Saito et al., 2004), likely by preventing

its phosphorylation-dependent degradation (Saito et al., 2004). Nonetheless, 100% of *cdc-*

14(he141) animals had a post-mitotic, invasive AC, even in the presence of *cki-1(RNAi)* (Fig.

S2B,C; Table 1), suggesting that CDC-14 is not a regulator of CKI-1 in the AC or that its

function is redundant with that of another phosphatase.

In the following subsection, we introduce viable null or hypomorphic alleles that impact other

negative regulators of the cell cycle (*cki-2*, *lin-35*, *fzr-1*, and *lin-23*) to determine their role in

maintaining the post-mitotic state of the AC.

CKI-1 is critical for maintaining the post-mitotic state of the AC

Only a handful of factors negatively regulate the somatic cell cycle during *C. elegans* larval

development (Kipreos and van den Heuvel, 2019). These include the CDK inhibitors CKI-1 and

CKI-2 (p21^{CIP1}/p27^{KIP1}) (Buck et al., 2009; Hong et al., 1998), the sole Rb family member LIN-35

(pRb/p107/p130) (Boxem and van den Heuvel, 2001; Boxem and van den Heuvel, 2002), the

APC/C co-activator FZR-1 (Cdh1/Hct1) (Fay et al., 2002), and the F-box/WD-repeat protein LIN-

23 (β-TrCP) (Kipreos et al., 1996; Kipreos et al., 2000; Nayak et al., 2002). Although they all

ensure normal cell cycle exit either generally (CKI-1 and LIN-23), redundantly (LIN-35 and FZR-

1), or tissue specifically (CKI-2), their role in maintaining the post-mitotic state is poorly

understood.

To determine their role in the AC, we generated four separate three-color imaging strains. Three

of these strains include putative null alleles, i.e., *cki-2(ok2105)*, *lin-35(n745)*, or *lin-23(e1883)*

(Kipreos et al., 2000; Lu and Horvitz, 1998; The *C. elegans* Deletion Mutant Consortium, 2012).

The fourth strain harbors *fzr-1(ku298)*, a hypomorphic allele, as FZR-1 function is required for

viability (Fay et al., 2002). However, regardless of their genotype, live imaging and subsequent

quantification of CDK activity demonstrated that they all possessed a post-mitotic, invasive AC

(Fig. 2; Table 1).

These results were consistent with what we found after depleting CKI-1 by RNAi in a wild-type

background of the three-color imaging strain (Fig. S1B,C; Table 1). We next explored whether

we could induce AC proliferation by depleting CKI-1 in each mutant background. Following *cki-*

1(RNAi) treatment, we observed ≥2 ACs in 13%, 15%, 35%, and 81% of *cki-2(ok2105)*, *lin-*

35(n745), *fzr-1(ku298)*, and *lin-23(e1883)* animals, respectively (Fig. 2). Compared to mutant

controls, this corresponded to a significant increase in AC-specific CDK activity (0.14 ± 0.03 (n =

30) vs. 0.55 ± 0.50 (n = 39), 0.15 ± 0.04 (n = 31) vs. 0.77 ± 0.52 (n = 54), 0.17 ± 0.05 (n = 32)

vs. 0.64 ± 0.57 (n = 55), and 0.13 ± 0.03 (n = 34) vs. 1.22 ± 0.72 (n = 111), respectively).

Additionally, we observed an increase in the percentage of proliferating ACs that underwent BM

invasion (Table 1). Overall, these findings, together with our *cki-1(RNAi)* results in RNAi-

sensitive backgrounds (Fig. 1D,E; Fig. S1A), highlight the critical role of CKI-1 in maintaining the

post-mitotic state of the AC.

Maintenance of the post-mitotic state is not required for AC invasion

To further investigate the lack of AC invasion defects following robust cell cycle induction (Table

1), we collected time-lapse movies of cycling ACs from pre-invasion to post-invasion using a

variant of the three-color imaging strain (Fig. 3; Fig. S3). Included in this strain is an AC-specific

F-actin probe, *cdh-3p*::mCherry::moesinABD, which we paired with a ubiquitous CDK activity

sensor, DHB::2xmTurquoise2, and an endogenous *lam-2* translational reporter, LAM-

2::mNeonGreen. We also added the *fzr-1(ku298)* hypomorphic allele rather than the *lin-*

23(e1883) null allele, as *fzr-1(ku298)* animals were easier to propagate and had the next

highest penetrance of the proliferative AC phenotype in the presence of *cki-1(RNAi)* (Fig. 2C,G;

Table 1).

Prior to time-lapse imaging, we captured static images of the invasive AC in *fzr-1(ku298);*

control(RNAi) animals (Fig. 3A), which allowed us to obtain a baseline level of CDK activity

values (0.31 ± 0.05 (n = 23)). This baseline allowed us to selectively record the activity of pre-

invasive ACs in *fzr-1(ku298); cki-1(RNAi)* animals that had already started cycling. Remarkably,

we collected five time-lapse datasets in which single or multiple cycling, pre-invasive ACs

showed an enrichment of the F-actin probe at the invasive membrane (Hagedorn et al., 2013)

(1.74 ± 0.36 (n = 7)), followed by successful invasion of the underlying BM (Fig. 3B; Movie 1). In

one of these datasets, we observed the division of a pre-invasive AC, with both daughter cells

re-entering the cell cycle before invading (Fig. S3). Altogether, these observations led us to

conclude that maintenance of the post-mitotic state is not required for AC invasion.

The pro-invasive GRN is functionally intact in proliferating, invasive ACs

After finding that direct disruption of the cell cycle leads to proliferating, invasive ACs, we

investigated whether these ACs maintain a functional pro-invasive gene regulatory network

(GRN) (Deng et al., 2020; Medwig-Kinney et al., 2020). This GRN consists of a cell cycle-

dependent and cell cycle-independent arm. We predicted that gene expression in both arms

would remain unchanged in proliferating, invasive ACs.

To test our hypothesis, we quantified endogenous levels of NHR-67, which is the distal

transcription factor in the cell cycle-dependent arm and is upstream of CKI-1. Additionally, we

quantified transcriptional levels of the matrix metalloproteinase (MMP) gene, *zmp-1*, which is

one of the downstream effectors of the cell cycle-independent arm. We quantified these levels in

two separate *fzr-1(ku298)* strains, each carrying most, if not all, of the original three-color

imaging alleles in the presence of *cki-1(RNAi)*. In one strain, we inserted an mNeonGreen tag

into the C-terminus of the endogenous *nhr-67* locus using CRISPR/Cas9 (Paix et al., 2015). In

the other strain, we exchanged the endogenous *lag-2* transcriptional reporter, LAG-

2::P2A::H2B::mTurquoise2, for a *zmp-1* transcriptional reporter transgene, *zmp-1p*::CFP (Inoue

et al., 2002). Supporting our prediction, NHR-67::mNeonGreen expression decreased by only

17% in proliferating, invasive ACs (Fig. 4A,B), whereas *zmp-1p*::CFP expression decreased by

73% in proliferating, invasive ACs (Fig. 4C,D). However, given that depletion of GRN

transcription factors results in proliferating, non-invasive ACs and a complete loss of *zmp-1*

transgene expression (Hwang et al., 2007; Matus et al., 2015; Medwig-Kinney et al., 2020;

Rimann and Hajnal, 2007; Sherwood et al., 2005), these data indicate that the pro-invasive

GRN is functionally intact in proliferating, invasive ACs (Fig. 4E).

Conclusions

In this study, we establish a critical role for CKI-1 in maintaining the post-mitotic state of the AC.

However, without knowing the null phenotype of *cki-1*, it remains difficult to ascertain the

redundancies, if any, in the AC cell cycle machinery (Matus et al., 2015). In *C. elegans*, there

are two CDK inhibitors: CKI-1 and CKI-2. Recent work has shown that *cki-2* becomes

upregulated in terminally differentiated vulval cells that are devoid of *cki-1* expression (Portegijs,

2019). This same compensatory relationship may exist in the AC. This could in part explain the

enhanced penetrance of the proliferative AC phenotype following loss of CKI-1 in *rrf-3(pk1426)*

and *cki-2(ok2105)* null mutant backgrounds (Fig. 1D,E; Fig. 2A,E). Interestingly, we observed a

synergistic increase in the number of proliferating ACs in *lin-23(e1883); cki-1(RNAi)* animals

(Fig. 2D,H). Perhaps complete loss of CKI-1, CKI-2, and LIN-23 is required to achieve a fully

penetrant ≥2 AC phenotype. We also do not know the null phenotype of *fzr-1*. Therefore, it may

have a more significant role in maintaining the post-mitotic state of the AC. However, use of the

fzr-1(ku298) hypomorphic allele sufficiently triggered cell cycle entry in post-mitotic neurons

(The et al., 2015). Lastly, as LIN-35 functions in both canonical cell cycle regulation and

negative regulation of RNAi pathways (Boxem and van den Heuvel, 2001; Boxem and van den

Heuvel, 2002; Lehner et al., 2006; Wang et al., 2005), we cannot rule out the possibility that the

lin-35(n745) null allele is simply enhancing the efficiency of *cki-1(RNAi)* (Fig. 2B,F). These

ambiguities could be addressed in future studies by creating and/or combining conditional null

alleles using the auxin-inducible degron (AID) system (Ashley et al., 2021; Hills-Muckey et al.,

2021; Martinez et al., 2020; Negishi et al., 2021; Sepers et al., 2022; Xiao et al., 2023; Zhang et

al., 2015) or an analogous approach (Nance and Frøkjær-Jensen, 2019).

Finally, we demonstrate that AC invasion can occur irrespective of cell cycle state (Fig. 3),

which breaks the dichotomy between invasion and proliferation in *C. elegans* (Matus et al.,

2015). We evaluated the functionality of the GRN that controls AC invasion (Deng et al., 2020;

Medwig-Kinney et al., 2020), specifically examining the expression of two components: NHR-67

and *zmp-1* (Fig. 4). These components provide readouts of the cell cycle-dependent and cell

cycle-independent arm, respectively. We observed a considerable reduction in *zmp-1*

expression in proliferating, invasive ACs, but not in NHR-67 expression. The presence of

reduced *zmp-1* expression in these cells suggests that cell cycle progression downregulates the

production of MMPs independently of the GRN components that transcriptionally regulate them

(Medwig-Kinney et al., 2020). However, MMPs are not necessary for AC invasion (Kelley et al.,

2019). In the absence of MMPs, the AC adapts by expanding its invasive F-actin network and

physically forcing its way through the underlying BM in an ATP-dependent manner. Given that
cell invasion and proliferation are both energy-intensive processes, future investigation is
warranted into whether they can occur simultaneously in the absence of MMPs. To our
knowledge, this is the first example in development where an invasively differentiated cell can
perform its specialized function while proliferating. Although evidence suggests that tumor cells
must alternate between proliferative and invasive states (Kohrman and Matus, 2017; Mondal et
al., 2022), additional research targeting the cell cycle in invasive tumor cells is required to
determine the necessity of this phenotypic switch.
281
282 **MATERIALS AND METHODS**
283
284 *Strains*
285
286 All strains were grown and maintained following standard procedures (Brenner, 1974). A
287 complete list of strains used in this study is available in Table S1.
288
289 *Alleles*
290
The following alleles were used in this study: *qyIs102[fos-1ap::rde-1]* (Hagedorn et al., 2009);
LG I *bmd156[rps-27p::DHB::2xmKate2]* (Adikes et al., 2020), *bmd294[rps-*
27p::DHB::2xmTurquoise2] (this study), and *lin-35(n745)* (Lu and Horvitz, 1998); **LG II**
bmd168[rps-27p::DHB::2xmKate2] (Adikes et al., 2020), *lin-23(e1883)* (Kipreos et al., 2000),
cki-1(gk132) (The *C. elegans* Deletion Mutant Consortium, 2012), *cdc-14(he141)* (Saito et al.,
2004), *fzr-1(ku298)* (Fay et al., 2002), *cki-2(ok2105)* (The *C. elegans* Deletion Mutant
Consortium, 2012), and *rrf-3(pk1426)* (Simmer et al., 2002); **LG IV** *qyIs10[lam-1p::lam-1::GFP]*
(Ziel et al., 2009), *qyIs225[cdh-3p::mCherry::moesinABD]* (Matus et al., 2015), and *nhr-*
67(wy1787[nhr-67::mNeonGreen]) (this study); **LG V** *lag-2(bmd202[lag-*
2::P2A::H2B::mTurquoise2]) (Medwig-Kinney et al., 2022), *rde-1(ne219)* (Tabara et al., 1999),
and *syIs67[zmp-1p::CFP]* (Inoue et al., 2002); **LG X** *lam-2(qy20[lam-2::mNeonGreen])* (Jayadev
et al., 2019) and *qyIs24[cdh-3p::mCherry::PLCδPH]* (Ziel et al., 2009).
303
304 *Generation of the transgenic bmd294 allele*
305

Both pWZ186 (Addgene plasmid #163641) and a plasmid containing 2xmTurquoise2 were

double digested with Bsu36I and NgoMIV to excise 2xmKate2 and 2xmTurquoise2,

respectively. pMAM038 (rps-27p::DHB::2xmTurquoise2) was cloned by T4 ligation using the

backbone from pWZ186 and 2xmTurquoise2 as an insert. After sequence confirmation,

pMAM038 was used as a repair template for insertion into the genome at a safe harbor site on

chromosome I corresponding to the MosSCI insertion site, ttTi4348 (Frøkjær-Jensen et al.,

2012). pAP082 was used as the sgRNA plasmid for chromosome I insertion by CRISPR/Cas9

(Pani and Goldstein, 2018). Young adults were transformed using standard microinjection

techniques, and integrants were identified through the SEC method (Dickinson et al., 2015).

Generation of the endogenous wy1787 allele

A repair template containing 30xlinker-mNeonGreen, with homology at the 5' and 3' ends to the

C-terminus of the *nhr-67* locus, was PCR amplified from pJW2171 (Addgene plasmid #163095)

and concentrated using a PCR purification kit (Qiagen). 3 µl of 10 µM stock tracRNA (IDT) was

incubated with 0.5 µl of a 100 µM crRNA (IDT) for 5 min at 95°C followed by 5 min at 25°C.

Immediately following incubation, the tracRNA:cRNA mixture was incubated with 0.5 µl of Alt-R

Cas9 protein (IDT) for 10 min at 37°C. The repair template and co-injection marker (pRF6) were

combined with the mixture to a final concentration of 200 ng/µl and 50 ng/µl, respectively, for a

final injection mix volume of 10 µl. Adult worms possessing no more than a single row of eggs

were transformed using standard microinjection techniques. Roller F1 progeny were singled out,

and F2 progeny were genotyped for insertions (Paix et al., 2015). The sequence of the PCR

primers and guide are available in Tables S2 and S3, respectively.

RNAi

All RNAi experiments were performed by first synchronizing animals at the L1 larval stage using

the bleaching technique (Porta-de-la-Riva et al., 2012). These animals were then fed bacteria

expressing the T444T-based RNAi construct for *cki-1* until the mid-L3 larval stage (P6.p four-cell

stage), when AC invasion normally occurs (Sherwood and Sternberg, 2003). Control animals

were fed bacteria carrying the empty T444T vector.

Generation of a T444T-based RNAi construct for cki-1

A T444T-based RNAi construct targeting *cki-1* was generated by inserting a 552 bp synthetic

DNA fragment, based on the *cki-1* cDNA sequence from WormBase (Davis et al., 2022), into

the T444T vector (Addgene plasmid #113081) using the BglII and XhoI restriction sites (Sturm

et al., 2018). The synthetic DNA was purchased from IDT as a gBlock with homology arms to

T444T for Gibson assembly. The sequence of the synthetic DNA is available in Table S4.

Image acquisition

Images were acquired using a spinning disk confocal microscope supported by Nobska

Imaging. This confocal system consists of a Hamamatsu ORCA EM-CCD camera mounted on

an upright Zeiss Axio Imager.A2, equipped with a Borealis-modified Yokogawa CSU-10

spinning disk scanning unit with 6 solid state (405, 440, 488, 514, 561, and 640 nm) lasers and

a Zeiss Plan-Apochromat 100x/1.4 oil DIC objective.

For static imaging, animals were anesthetized by placing them into a drop of M9 on a 5%

agarose pad containing 7 mM sodium azide and securing them with a coverslip. Time-lapse

imaging was performed using a modified version of a previously published protocol (Kelley et

al., 2017), as described in Adikes et al. (2020).

Imaging processing and analysis

The acquired images were processed using ImageJ/Fiji (Schneider et al., 2012). DHB ratios

were quantified as previously described (Adikes et al., 2020). NHR-67::mNeonGreen and *zmp-*

1::CFP fluorescence were quantified as previously described (Medwig-Kinney et al., 2020). AC

invasion was defined as the complete loss of fluorescent BM signal underneath the AC. Plots

were generated using Prism software, and figures, including cartoons, were created using a

combination of Adobe Photoshop and Illustrator.

Statistics

A power analysis was performed to determine the number of animals (N) needed per

experiment (Pollard et al., 2019). The figure legends specify the measures of central tendency,

error bars, numeric *P*-values, and statistical tests used.

Acknowledgements

We thank Nicholas Palmisano, Sam Stettnisch, and Abigail Nishimura for their comments and

suggestions.

Data availability

All relevant data can be found within the article and its supplementary information.

Competing interests

D.Q.M. is a paid employee of Arcadia Science.

Author contributions

Conceptualization: M.A.Q.M., D.Q.M.; Methodology: M.A.Q.M., D.Q.M.; Validation: M.A.Q.M.,

C.Z.Z.; Formal analysis: M.A.Q.M.; Investigation: M.A.Q.M., C.Z.Z., F.E.Q.M.; Resources:

F.E.Q.M., C.Y., W.Z.; Writing – original draft: M.A.Q.M.; Writing – review and editing: M.A.Q.M.,

D.Q.M.; Visualization: M.A.Q.M., C.Z.Z., D.Q.M.; Supervision: B.L.M., D.Q.M.; Project

administration: M.A.Q.M.; Funding acquisition: K.S., B.L.M., D.Q.M.

Funding

M.A.Q.M. is supported by the National Cancer Institute (F30CA257383), F.E.Q.M. is supported

by the National Institute of General Medical Sciences Diversity Supplement Program

(R01GM121597), C.Y. is supported by the Human Frontiers Science Program (LT000127/2016-

L), and K.S. is a Howard Hughes Medical Institute Investigator. Additionally, both B.L.M. and

D.Q.M. are supported by the Damon Runyon Cancer Research Foundation (DRR4714), and

D.Q.M. is also supported by the National Institute of General Medical Sciences

(R01GM121597).

REFERENCES

Adikes, R. C., Kohrman, A. Q., Martinez, M. A. Q., Palmisano, N. J., Smith, J. J., Medwig-
Kinney, T. N., Min, M., Sallee, M. D., Ahmed, O. B., Kim, N., et al. (2020). Visualizing the
metazoan proliferation-quiescence decision in vivo. *eLife* 9, e63265.

Ajioka, I., Martins, R. A. P., Bayazitov, I. T., Donovan, S., Johnson, D. A., Frase, S., Cicero, S.
A., Boyd, K., Zakharenko, S. S. and Dyer, M. A. (2007). Differentiated Horizontal
Interneurons Clonally Expand to Form Metastatic Retinoblastoma in Mice. *Cell* 131,
378–390.

Ashley, G. E., Duong, T., Levenson, M. T., Martinez, M. A. Q., Johnson, L. C., Hibshman, J. D.,
Saeger, H. N., Palmisano, N. J., Doonan, R., Martinez-Mendez, R., et al. (2021). An
expanded auxin-inducible degron toolkit for *Caenorhabditis elegans*. *Genetics* 217,
iyab006.

Boxem, M. and van den Heuvel, S. (2001). *lin-35* Rb and *cki-1* Cip/Kip cooperate in
developmental regulation of G1 progression in *C. elegans*. *Development* 128, 4349–
4359.

Boxem, M. and van den Heuvel, S. (2002). *C. elegans* Class B Synthetic Multivulva Genes Act
in G1 Regulation. *Current Biology* 12, 906–911.

Brenner, S. (1974). The genetics of *Caenorhabditis elegans*. *Genetics* 77, 71–94.

Buck, S. H., Chiu, D. and Saito, R. M. (2009). The cyclin-dependent kinase inhibitors, *cki-1* and
cki-2, act in overlapping but distinct pathways to control cell-cycle quiescence during *C.*
elegans development. *Cell Cycle* 8, 2613–2620.

Buttitta, L. A. and Edgar, B. A. (2007). Mechanisms controlling cell cycle exit upon terminal
differentiation. *Current Opinion in Cell Biology* 19, 697–704.

Clayton, J. E., van den Heuvel, S. J. L. and Saito, R. M. (2008). Transcriptional control of cell-
cycle quiescence during *C. elegans* development. *Developmental Biology* 313, 603–613.

Davis, P., Zarowiecki, M., Arnaboldi, V., Becerra, A., Cain, S., Chan, J., Chen, W. J., Cho, J., da
Veiga Beltrame, E., Diamantakis, S., et al. (2022). WormBase in 2022—data, processes,
and tools for analyzing *Caenorhabditis elegans*. *Genetics* 220, iyac003.

Deng, T., Stempor, P., Appert, A., Daube, M., Ahringer, J., Hajnal, A. and Lattmann, E. (2020).
The *Caenorhabditis elegans* homolog of the Evi1 proto-oncogene, *egl-43*, coordinates
G1 cell cycle arrest with pro-invasive gene expression during anchor cell invasion. *PLoS*
Genet 16, e1008470.

Dickinson, D. J., Pani, A. M., Heppert, J. K., Higgins, C. D. and Goldstein, B. (2015).
Streamlined Genome Engineering with a Self-Excising Drug Selection Cassette.
Genetics 200, 1035–1049.

Fay, D. S., Keenan, S. and Han, M. (2002). *fzr-1* and *lin-35* /Rb function redundantly to control
cell proliferation in *C. elegans* as revealed by a nonbiased synthetic screen. *Genes Dev.*
16, 503–517.

Feng, H., Zhong, W., Punkosdy, G., Gu, S., Zhou, L., Seabolt, E. K. and Kipreos, E. T. (1999).
CUL-2 is required for the G1-to-S-phase transition and mitotic chromosome
condensation in *Caenorhabditis elegans*. *Nature Cell Biology* 1, 486–492.

Frøkjær-Jensen, C., Davis, M. W., Ailion, M. and Jorgensen, E. M. (2012). Improved Mos1-
mediated transgenesis in *C. elegans*. *Nat Methods* 9, 117–118.

Fukuyama, M., Gendreau, S. B., Derry, W. B. and Rothman, J. H. (2003). Essential embryonic
roles of the CKI-1 cyclin-dependent kinase inhibitor in cell-cycle exit and morphogenesis
in *C. elegans*. *Developmental Biology* 260, 273–286.

Haass, N. K., Beaumont, K. A., Hill, D. S., Anfosso, A., Mrass, P., Munoz, M. A., Kinjyo, I. and
Weninger, W. (2014). Real-time cell cycle imaging during melanoma growth, invasion,
and drug response. *Pigment Cell & Melanoma Research* 27, 764–776.

Hagedorn, E. J., Yashiro, H., Ziel, J. W., Ihara, S., Wang, Z. and Sherwood, D. R. (2009).
Integrin Acts Upstream of Netrin Signaling to Regulate Formation of the Anchor Cell's
Invasive Membrane in *C. elegans*. *Developmental Cell* 17, 187–198.

Hagedorn, E. J., Ziel, J. W., Morrissey, M. A., Linden, L. M., Wang, Z., Chi, Q., Johnson, S. A.
and Sherwood, D. R. (2013). The netrin receptor DCC focuses invadopodia-driven
basement membrane transmigration in vivo. *Journal of Cell Biology* 201, 903–913.

Hills-Muckey, K., Martinez, M. A. Q., Stec, N., Hebbar, S., Saldanha, J., Medwig-Kinney, T. N.,
Moore, F. E. Q., Ivanova, M., Morao, A., Ward, J. D., et al. (2021). An engineered,
orthogonal auxin analog/AtTIR1(F79G) pairing improves both specificity and efficacy of
the auxin degradation system in *Caenorhabditis elegans*. *Genetics* 220, iyab174.

Hong, Y., Roy, R. and Ambros, V. (1998). Developmental regulation of a cyclin-dependent
kinase inhibitor controls postembryonic cell cycle progression in *Caenorhabditis elegans*.
Development 125, 3585–3597.

Hwang, B. J., Meruelo, A. D. and Sternberg, P. W. (2007). *C. elegans* EVI1 proto-oncogene,
EGL-43, is necessary for Notch-mediated cell fate specification and regulates cell
invasion. *Development* 134, 669–679.

Inoue, T., Sherwood, D. R., Aspöck, G., Butler, J. A., Gupta, B. P., Kirouac, M., Wang, M., Lee,
P.-Y., Kramer, J. M., Hope, I., et al. (2002). Gene expression markers for *Caenorhabditis*
elegans vulval cells. *Mechanisms of Development* 119, S203–S209.

Jayadev, R., Chi, Q., Keeley, D. P., Hastie, E. L., Kelley, L. C. and Sherwood, D. R. (2019). α-
Integrins dictate distinct modes of type IV collagen recruitment to basement membranes.
Journal of Cell Biology 218, 3098–3116.

Kelley, L. C., Wang, Z., Hagedorn, E. J., Wang, L., Shen, W., Lei, S., Johnson, S. A. and
Sherwood, D. R. (2017). Live-cell confocal microscopy and quantitative 4D image
analysis of anchor-cell invasion through the basement membrane in *Caenorhabditis*
elegans. *Nature Protocols* 12, 2081–2096.

Kelley, L. C., Chi, Q., Cáceres, R., Hastie, E., Schindler, A. J., Jiang, Y., Matus, D. Q., Plastino,
J. and Sherwood, D. R. (2019). Adaptive F-Actin Polymerization and Localized ATP
Production Drive Basement Membrane Invasion in the Absence of MMPs.
Developmental Cell 48, 313-328.e8.

Kipreos, E. T. and van den Heuvel, S. (2019). Developmental Control of the Cell Cycle: Insights
from *Caenorhabditis elegans*. *Genetics* 211, 797–829.

Kipreos, E. T., Lander, L. E., Wing, J. P., He, W. W. and Hedgecock, E. M. (1996). *cul-1* Is
Required for Cell Cycle Exit in C. elegans and Identifies a Novel Gene Family. *Cell* 85,
829–839.

Kipreos, E. T., Gohel, S. P. and Hedgecock, E. M. (2000). The *C. elegans* F-box/WD-repeat
protein LIN-23 functions to limit cell division during development. *Development* 127,
5071–5082.

Kohrman, A. Q. and Matus, D. Q. (2017). Divide or Conquer: Cell Cycle Regulation of Invasive
Behavior. *Trends in Cell Biology* 27, 12–25.

Lehner, B., Calixto, A., Crombie, C., Tischler, J., Fortunato, A., Chalfie, M. and Fraser, A. G.
(2006). Loss of LIN-35, the *Caenorhabditis elegans* ortholog of the tumor suppressor
p105Rb, results in enhanced RNA interference. *Genome Biology* 7, R4.

Lu, X. and Horvitz, H. R. (1998). *lin-35* and *lin-53*, Two Genes that Antagonize a *C. elegans* Ras
Pathway, Encode Proteins Similar to Rb and Its Binding Protein RbAp48. *Cell* 95, 981–
991.

Martinez, M. A. Q. and Matus, D. Q. (2022). CDK activity sensors: genetically encoded
ratiometric biosensors for live analysis of the cell cycle. *Biochemical Society*
Transactions 50, 1081–1090.

Martinez, M. A. Q., Kinney, B. A., Medwig-Kinney, T. N., Ashley, G., Ragle, J. M., Johnson, L.,
Aguilera, J., Hammell, C. M., Ward, J. D. and Matus, D. Q. (2020). Rapid Degradation of
Caenorhabditis elegans Proteins at Single-Cell Resolution with a Synthetic Auxin. *G3*
Genes|Genomes|Genetics 10, 267–280.

Matus, D. Q., Li, X.-Y., Durbin, S., Agarwal, D., Chi, Q., Weiss, S. J. and Sherwood, D. R.
(2010). In Vivo Identification of Regulators of Cell Invasion Across Basement
Membranes. *Sci. Signal.* 3, ra35.

Matus, D. Q., Lohmer, L. L., Kelley, L. C., Schindler, A. J., Kohrman, A. Q., Barkoulas, M.,
Zhang, W., Chi, Q. and Sherwood, D. R. (2015). Invasive Cell Fate Requires G1 Cell-
Cycle Arrest and Histone Deacetylase-Mediated Changes in Gene Expression.
Developmental Cell 35, 162–174.

Medwig-Kinney, T. N., Smith, J. J., Palmisano, N. J., Tank, S., Zhang, W. and Matus, D. Q.

(2020). A developmental gene regulatory network for *C. elegans* anchor cell invasion.

Development 147, dev.185850.

Medwig-Kinney, T. N., Sirota, S. S., Gibney, T. V., Pani, A. M. and Matus, D. Q. (2022). An *in*

vivo toolkit to visualize endogenous LAG-2/Delta and LIN-12/Notch signaling in *C.*

elegans. *MicroPublication Biology* 2022, 10.17912/micropub.biology.000602.

Mondal, C., Gacha-Garay, M. J., Larkin, K. A., Adikes, R. C., Di Martino, J. S., Chien, C.-C.,

Fraser, M., Eni-aganga, I., Agullo-Pascual, E., Cialowicz, K., et al. (2022). A proliferative

to invasive switch is mediated by srGAP1 downregulation through the activation of TGF-

β2 signaling. *Cell Reports* 40,.

Nance, J. and Frøkjær-Jensen, C. (2019). The *Caenorhabditis elegans* Transgenic Toolbox.

Genetics 212, 959–990.

Nayak, S., Santiago, F. E., Jin, H., Lin, D., Schedl, T. and Kipreos, E. T. (2002). The

Caenorhabditis elegans Skp1-Related Gene Family: Diverse Functions in Cell

Proliferation, Morphogenesis, and Meiosis. *Current Biology* 12, 277–287.

Negishi, T., Kitagawa, S., Horii, N., Tanaka, Y., Haruta, N., Sugimoto, A., Sawa, H., Hayashi, K.,

Harata, M. and Kanemaki, M. T. (2021). The auxin-inducible degron 2 (AID2) system

enables controlled protein knockdown during embryogenesis and development in

Caenorhabditis elegans. *Genetics* 220, iyab218.

Paix, A., Folkmann, A., Rasoloson, D. and Seydoux, G. (2015). High Efficiency, Homology-

Directed Genome Editing in *Caenorhabditis elegans* Using CRISPR-Cas9

Ribonucleoprotein Complexes. *Genetics* 201, 47–54.

Pani, A. M. and Goldstein, B. (2018). Direct visualization of a native Wnt in vivo reveals that a

long-range Wnt gradient forms by extracellular dispersal. *eLife* 7, e38325.

Pollard, D. A., Pollard, T. D. and Pollard, K. S. (2019). Empowering statistical methods for

cellular and molecular biologists. *MBoC* 30, 1359–1368.

Porta-de-la-Riva, M., Fontrodona, L., Villanueva, A. and Cerón, J. (2012). Basic *Caenorhabditis*

elegans Methods: Synchronization and Observation. *JoVE* e4019.

Portegijs, V. (2019). Developmental control of the *C. elegans* cell cycle.

Rimann, I. and Hajnal, A. (2007). Regulation of anchor cell invasion and uterine cell fates by the
egl-43 Evi-1 proto-oncogene in *Caenorhabditis elegans*. *Developmental Biology* 308,
187–195.

Roy, S. H., Clayton, J. E., Holmen, J., Beltz, E. and Saito, R. M. (2011). Control of Cdc14
activity coordinates cell cycle and development in *Caenorhabditis elegans*. *Mechanisms*
of Development 128, 317–326.

Roy, S. H., Tobin, D. V., Memar, N., Beltz, E., Holmen, J., Clayton, J. E., Chiu, D. J., Young, L.
D., Green, T. H., Lubin, I., et al. (2014). A Complex Regulatory Network Coordinating
Cell Cycles During *C. elegans* Development Is Revealed by a Genome-Wide RNAi
Screen. *G3 Genes|Genomes|Genetics* 4, 795–804.

Ruijtenberg, S. and van den Heuvel, S. (2016). Coordinating cell proliferation and differentiation:
Antagonism between cell cycle regulators and cell type-specific gene expression. *Cell*
Cycle 15, 196–212.

Sage, C., Huang, M., Karimi, K., Gutierrez, G., Vollrath, M. A., Zhang, D.-S., García-Añoveros,
J., Hinds, P. W., Corwin, J. T., Corey, D. P., et al. (2005). Proliferation of Functional Hair
Cells in Vivo in the Absence of the Retinoblastoma Protein. *Science* 307, 1114–1118.

Saito, R. M., Perreault, A., Peach, B., Satterlee, J. S. and van den Heuvel, S. (2004). The CDC-
14 phosphatase controls developmental cell-cycle arrest in *C. elegans*. *Nat Cell Biol* 6,
777–783.

Schneider, C. A., Rasband, W. S. and Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of
image analysis. *Nat Methods* 9, 671–675.

Sepers, J. J., Verstappen, N. H. M., Vo, A. A., Ragle, J. M., Ruijtenberg, S., Ward, J. D. and
Boxem, M. (2022). The mIAA7 degron improves auxin-mediated degradation in *C.*
elegans. *G3 Genes|Genomes|Genetics* 12, jkac222.

Sherwood, D. R. and Sternberg, P. W. (2003). Anchor Cell Invasion into the Vulval Epithelium in
C. elegans. *Developmental Cell* 5, 21–31.

Sherwood, D. R., Butler, J. A., Kramer, J. M. and Sternberg, P. W. (2005). FOS-1 Promotes
Basement-Membrane Removal during Anchor-Cell Invasion in *C. elegans*. *Cell* 121,
951–962.

Simmer, F., Tijsterman, M., Parrish, S., Koushika, S. P., Nonet, M. L., Fire, A., Ahringer, J. and
Plasterk, R. H. A. (2002). Loss of the Putative RNA-Directed RNA Polymerase RRF-3
Makes *C. elegans* Hypersensitive to RNAi. *Current Biology* 12, 1317–1319.

Smith, J. J., Xiao, Y., Parsan, N., Medwig-Kinney, T. N., Martinez, M. A. Q., Moore, F. E. Q.,
Palmisano, N. J., Kohrman, A. Q., Chandhok Delos Reyes, M., Adikes, R. C., et al.
(2022). The SWI/SNF chromatin remodeling assemblies BAF and PBAF differentially
regulate cell cycle exit and cellular invasion *in vivo*. *PLoS Genet* 18, e1009981.

Soufi, A. and Dalton, S. (2016). Cycling through developmental decisions: how cell cycle
dynamics control pluripotency, differentiation and reprogramming. *Development* 143,
4301–4311.

Spencer, S. L., Cappell, S. D., Tsai, F.-C., Overton, K. W., Wang, C. L. and Meyer, T. (2013).
The Proliferation-Quiescence Decision Is Controlled by a Bifurcation in CDK2 Activity at
Mitotic Exit. *Cell* 155, 369–383.

Sturm, Á., Saskői, É., Tibor, K., Weinhardt, N. and Vellai, T. (2018). Highly efficient RNAi and
Cas9-based auto-cloning systems for *C. elegans* research. *Nucleic Acids Research* 46,
e105.

Tabara, H., Sarkissian, M., Kelly, W. G., Fleenor, J., Grishok, A., Timmons, L., Fire, A. and
Mello, C. C. (1999). The *rde-1* Gene, RNA Interference, and Transposon Silencing in *C.
elegans*. *Cell* 99, 123–132.

The, I., Ruijtenberg, S., Bouchet, B. P., Cristobal, A., Prinsen, M. B. W., van Mourik, T., Koreth,
J., Xu, H., Heck, A. J. R., Akhmanova, A., et al. (2015). Rb and FZR1/Cdh1 determine
CDK4/6-cyclin D requirement in *C. elegans* and human cancer cells. *Nature
Communications* 6, 5906.

The *C. elegans* Deletion Mutant Consortium (2012). Large-Scale Screening for Targeted
Knockouts in the *Caenorhabditis elegans* Genome. *G3 Genes|Genomes|Genetics* 2,
1415–1425.

Timmons, L. and Fire, A. (1998). Specific interference by ingested dsRNA. *Nature* 395, 854–
854.

van Rijnberk, L. M., van der Horst, S. E. M., van den Heuvel, S. and Ruijtenberg, S. (2017). A
dual transcriptional reporter and CDK-activity sensor marks cell cycle entry and
progression in *C. elegans. PLoS ONE* 12, e0171600.

Vittadello, S. T., McCue, S. W., Gunasingh, G., Haass, N. K. and Simpson, M. J. (2020).
Examining Go-or-Grow Using Fluorescent Cell-Cycle Indicators and Cell-Cycle-Inhibiting
Drugs. *Biophysical Journal* 118, 1243–1247.

Wang, D., Kennedy, S., Conte, D., Kim, J. K., Gabel, H. W., Kamath, R. S., Mello, C. C. and
Ruvkun, G. (2005). Somatic misexpression of germline P granules and enhanced RNA
interference in retinoblastoma pathway mutants. *Nature* 436, 593–597.

Xiao, Y., Yee, C., Zhao, C. Z., Martinez, M. A. Q., Zhang, W., Shen, K., Matus, D. Q. and
Hammell, C. (2023). An expandable FLP-ON::TIR1 system for precise spatiotemporal
protein degradation in *Caenorhabditis elegans. Genetics* iyad013.

Zhang, L., Ward, J. D., Cheng, Z. and Dernburg, A. F. (2015). The auxin-inducible degradation
(AID) system enables versatile conditional protein depletion in *C. elegans. Development*
142, 4374–4384.

Ziel, J. W., Hagedorn, E. J., Audhya, A. and Sherwood, D. R. (2009). UNC-6 (netrin) orients the
invasive membrane of the anchor cell in *C. elegans. Nat Cell Biol* 11, 183–189.

A   *C. elegans* invasion/proliferation dichotomy

breached BM   wild-type AC   proliferating AC   intact BM

VPCs   VPCs

B   CDK activity sensor

DHB $\xrightarrow{CDK}$ DHB-P

nucleus   cytoplasm

cell cycle progression

$$CDK\ activity = \frac{cytoplasmic\ DHB}{nuclear\ DHB}$$

G0 → G1 → S → G2

C   *wild type*

DIC   ▢ LAM-2::mNG   ▢ LAG-2::P2A::H2B::mT2   ▢ DHB::2xmK2

P6.p 1-cell

P6.p 2-cell

P6.p 4-cell

D   *rrf-3(pk1426)*

▢ LAM-2::mNG   ▢ LAG-2::P2A::H2B::mT2   ▢ DHB::2xmK2

control(RNAi)   1 AC (30/30)

cki-1(RNAi)   ≥2 AC (8/34)

E   *rrf-3(pk1426)*

DHB cytoplasmic:nuclear ratio

***

control(RNAi)   *cki-1(RNAi)*

Figure 1. A moderate increase in AC proliferation is observed upon systemic depletion of

CKI-1. (A) Schematic of the *C. elegans* invasion/proliferation dichotomy. (B) Schematic of the

CDK activity sensor, DHB, which is phosphorylated by CDK and translocates from the nucleus

to the cytoplasm in response to CDK activity during cell cycle progression. The equation used to

quantify CDK activity is shown. (C) DIC (left) and confocal images merging LAM-

2::mNeonGreen with LAG-2::P2A::H2B::mTurquoise2 (middle) and DHB::2xmKate2 (right) from

the P6.p 1-cell stage to the P6.p 4-cell stage are shown. Arrowheads and brackets indicate the

position of the AC and P6.p cells, respectively. The scale bar represents 5 µm. (D) Confocal

images merging LAM-2::mNeonGreen with LAG-2::P2A::H2B::mTurquoise2 (left) and

DHB::2xmKate2 (right) from the P6.p 4-cell stage are shown. Arrowheads indicate the position

of the AC(s) in *rrf-3(pk1426)* animals following *control(RNAi)* and *cki-1(RNAi)* treatment. The

scale bar represents 5 µm. (E) Plot displays median and IQR of DHB::2xmKate2 ratios in the

AC(s) of *rrf-3(pk1426)* animals following *control(RNAi)* and *cki-1(RNAi)* treatment (N ≥ 30

animals per treatment). Statistical significance was determined by a Mann-Whitney test (P <

0.0004). See also Figure S1, S2.

Figure 2. CKI-1 is critical for maintaining the post-mitotic state of the AC. (A-D) Confocal

images merging LAM-2::mNeonGreen with LAG-2::P2A::H2B::mTurquoise2 (left) and

DHB::2xmKate2 (right) from the P6.p 4-cell stage are shown. Arrowheads indicate the position

of the AC(s) in *cki-2(ok2105)* (A), *lin-35(n745)* (B), *fzr-1(ku298)* (C), and *lin-23(e1883)* (D)

animals following *control(RNAi)* and *cki-1(RNAi)* treatment. Scale bars represent 5 µm. (E-H)

Plots display median and IQR of DHB::2xmKate2 ratios in the AC(s) of *cki-2(ok2105)* (E), *lin-*

35(n745) (F), *fzr-1(ku298)* (G), and *lin-23(e1883)* (H) animals following *control(RNAi)* and *cki-*

1(RNAi) treatment (N ≥ 30 animals per treatment). Statistical significance was determined by a

Mann-Whitney test (P < 0.0001).

700 **Table 1. Genetic analysis of negative cell cycle regulators during AC invasion.**

Genotype	Treatment	% invasion		% no invasion		
		1 AC	≥2 AC	1 AC	≥2 AC	N
wild type	*control(RNAi)*	100	0	0	0	27
rrf-3(pk1426)		97	0	3	0	30
cdc-14(he141)		100	0	0	0	35
cki-2(ok2105)		100	0	0	0	30
lin-35(n745)		100	0	0	0	31
fzr-1(ku298)		100	0	0	0	32
lin-23(e1883)		100	0	0	0	34
wild type	*cki-1(RNAi)*	100	0	0	0	29
rrf-3(pk1426)		76	9	0	15	34
cdc-14(he141)		100	0	0	0	35
cki-2(ok2105)		87	13	0	0	31
lin-35(n745)		85	12	0	3	33
fzr-1(ku298)		65	29	0	6	34
lin-23(e1883)		19	74	0	7	31

701 AC invasion was scored at the P6.p 4-cell stage of vulval development, the stage when AC

702 invasion normally occurs, using AC/VPC, BM, and cell cycle fluorescent reporters.

Figure 3. Maintenance of the post-mitotic state is not required for AC invasion. (A) Static

images of *cdh-3p*::mCherry::moesinABD (left), LAM-2::mNeonGreen (middle), and

DHB::2xmTurquoise2 (right) from the P6.p 4-cell stage are shown. Arrowheads indicate the

position of the AC in a *fzr-1(ku298); control(RNAi)* animal. The scale bar represents 5 µm. (B)

Time-lapse images of *cdh-3p*::mCherry::moesinABD (left), LAM-2::mNeonGreen (middle), and

DHB::2xmTurquoise2 (right) from pre-AC invasion to post-AC invasion are shown. Arrowheads

indicate the position of multiple invading ACs in a *fzr-1(ku298); cki-1(RNAi)* animal. The scale

bar represents 5 µm. See also Movie 1 and Figure S3.

Movie 1. Time-lapse imaging confirms that cycling ACs can invade. A timelapse of two

cycling ACs from pre-invasion to post-invasion is shown. The AC cell membrane, BM, and CDK

activity are visualized using an F-actin probe (*cdh-3p*::mCherry::moesinABD), an endogenous

lam-2 translational reporter (LAM-2::mNeonGreen), and a ubiquitous CDK activity sensor

(DHB::2xmTurquoise2). The time points were acquired every 5 min for 350 min with a step size

of 1 μm. The scale bar represents 5 μm.

Figure 4. Proliferating, invasive ACs retain a functionally intact pro-invasive GRN. (A)

Confocal images of LAG-2::P2A::H2B::mTurquoise2 (left), NHR-67::mNeonGreen (middle),

LAM-2::mNeonGreen (middle), and DHB::2xmKate2 (right) from the P6.p 4-cell stage are

shown. Arrowheads indicate the position of the AC(s) in *fzr-1(ku298)* animals following

control(RNAi) and *cki-1(RNAi)* treatment. The scale bar represents 5 µm. (B) Plot displays mean

and SD of normalized NHR-67::mNeonGreen intensity in the AC(s) of *fzr-1(ku298)* animals

following *control(RNAi)* and *cki-1(RNAi)* treatment (N ≥ 30 animals per treatment). (C) Confocal

images of *zmp-1p*::CFP (left), LAM-2::mNeonGreen (middle), and DHB::2xmKate2 (right) from

the P6.p 4-cell stage are shown. Arrowheads indicate the position of the AC(s) in *fzr-1(ku298)*

animals following *control(RNAi)* and *cki-1(RNAi)* treatment. The scale bar represents 5 µm. (D)

Plot displays mean and SD of normalized *zmp-1p*::CFP intensity in the AC(s) of *fzr-1(ku298)*

animals following *control(RNAi)* and *cki-1(RNAi)* treatment (N ≥ 31 animals per treatment). (E)

Whereas depletion of pro-invasive GRN transcription factors (TFs) results in proliferating, non-

invasive ACs (left), direct perturbation of the AC cell cycle results in proliferating, invasive ACs

that maintain pro-invasive gene expression (right).