1

Massively integrated coexpression

analysis reveals transcriptional

regulation, evolution and cellular

implications of the noncanonical

translatome

April Rich[+,1,2,3], Omer Acar[+,1,2,3], Anne-Ruxandra Carvunis*[2,3]

[1]*Joint Carnegie Mellon University-University of Pittsburgh Computational Biology PhD Program,*

University of Pittsburgh, Pittsburgh, PA, USA; [2]*Department of Computational and Systems*

Biology, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA; [3]*Pittsburgh Center for*

Evolutionary Biology and Medicine (CEBaM), University of Pittsburgh, Pittsburgh, PA, USA

[+] co-first authors

* corresponding author

Abstract

Background:

Recent studies uncovered pervasive transcription and translation of thousands of noncanonical

open reading frames (nORFs) outside of annotated genes. The contribution of nORFs to cellular

phenotypes is difficult to infer using conventional approaches because nORFs tend to be short,

of recent *de novo* origins, and lowly expressed. Here we develop a dedicated coexpression

analysis framework that accounts for low expression to investigate the transcriptional regulation,

evolution, and potential cellular roles of nORFs in *Saccharomyces cerevisiae*.

Results:

Our results reveal that nORFs tend to be preferentially coexpressed with genes involved in

cellular transport or homeostasis but rarely with genes involved in RNA processing.

Mechanistically, we discover that young *de novo* nORFs located downstream of conserved

genes tend to leverage their neighbors' promoters through transcription readthrough, resulting in

high coexpression and high expression levels. Transcriptional piggybacking also influences the

coexpression profiles of young *de novo* nORFs located upstream of genes, but to a lesser

extent and without detectable impact on expression levels. Transcriptional piggybacking

influences, but does not determine, the transcription profiles of *de novo* nORFs emerging

nearby genes. About 40% of nORFs are not strongly coexpressed with any gene but are

transcriptionally regulated nonetheless and tend to form entirely new transcription modules. We

offer a web browser interface (https://carvunislab.csb.pitt.edu/shiny/coexpression/) to efficiently

query, visualize and download our coexpression inferences.

Conclusions:

Our results suggest that nORF transcription is highly regulated. Our coexpression dataset

serves as an unprecedented resource for unraveling how nORFs integrate into cellular

networks, contribute to cellular phenotypes, and evolve.

Keywords:

<u>Coexpression networks, de novo gene birth, noncanonical ORFs, translatome, smORFs,</u>

<u>transcriptional regulation</u>

43   Background

Eukaryotic genomes encompass thousands of open reading frames (ORFs). The vast majority

are so-called "noncanonical" ORFs (nORFs) excluded from genome annotations because of

their short length, lack of evolutionary conservation, and perceived irrelevance to cellular

physiology [1–3]. The development of RNA sequencing (RNA-seq) [4] and ribosome profiling

[5,6] has revealed genome-wide transcription and translation of nORFs across species ranging

from yeast to humans [6–14]. Recent studies have characterized individual nORFs that form

stable peptides and impact phenotypes, including cell growth [10,13,15], cell cycle regulation

[16], muscle physiology [17–19], and immunity [20–22]. Unraveling the cellular, physiological

and evolutionary implications of nORFs has become an active area of research [14,23].

Many nORFs have evolved *de novo* from previously noncoding regions [24–26]. Thus, the study

of nORFs and *de novo* gene birth as evolutionary innovation carries a synergistic overlap where

findings in one area could improve our understanding of the other. For instance, Sandmann et

al. measured physical protein interactions for hundreds of peptides translated from nORFs and

proposed that short linear motifs present in young *de novo* nORFs could mediate how nORFs

impact essential cellular processes [26]. Other studies observed a gradual integration of

evolutionary young ORFs into cellular networks and showed they could gain essential roles [27–

29]. These studies support an evolutionary model whereby pervasive expression of nORFs

generates the raw material for *de novo* gene birth [24,25].

The biological interpretation of nORF expression is complex. Some studies suggest that the

transcription or translation of nORFs could be attributed to expression noise [30–32], whereby

non-specific binding of RNA polymerases and ribosomes to DNA and RNA might cause

promiscuous transcription or translation, respectively. How do nORFs become expressed in the

first place? There are multiple hypotheses on how *de novo* ORFs gain the ability to become

transcriptionally regulated [33]. One possibility is the emergence of novel regulatory regions

along with or following the emergence of an ORF (ORF-first), as was shown for specific *de novo*

ORFs in *Drosophila melanogaster* [34], codfish [35], human [36,37] and chimpanzee [36].

Alternatively, ORFs may emerge on actively transcribed loci such as near enhancers [38] or on

long noncoding RNAs [39], as was shown for *de novo* ORFs in primates [40] and for *de novo*

ORFs upstream or downstream of transcripts containing genes [37] (transcription-first) [41–43].

Transcription has a ripple effect causing coordinated activation of nearby genes [44,45]. Thus,

de novo ORFs that emerge near established genes or regulatory regions may acquire

transcriptional regulation by 'piggybacking' [45] on the pre-existing regulatory context [41,46].

This piggybacking could predispose *de novo* ORFs to be involved in similar cellular processes

as their neighbors, which in turn would help with characterization. To date, the fraction of

nORFs that are transcriptionally regulated and contribute to cellular phenotypes is unknown for

any species.

An obstacle to studying nORF expression at scale is their detection, as nORF expression levels

are typically low and reliant on specific conditions [24,36]. Recent studies demonstrated that the

integration of omics data [14,47–49] could effectively address detection issues. For example,

Wacholder et al. [14] recently discovered around 19,000 translated nORFs in *Saccharomyces*

cerevisiae by massive integration of ribosome profiling data. This figure is three times larger

than the number of canonical ORFs (cORFs) annotated in the yeast genome. These translated

nORFs have the potential to generate peptides that affect cellular phenotypes but are almost

entirely uncharacterized.

Coexpression is a well-established approach for studying transcriptional regulation through the

massive integration of RNA-seq data. Coexpression refers to the similarity between

transcriptional profiles of ORF pairs across numerous samples. Coexpression has been used

successfully to identify new gene functions [50,51], disease-related genes [22,52,53] and for

studying the conservation of the regulatory machinery [51,54] or gene modules [55] between

species. Based on the assumption that genes involved in similar pathways have correlated

expression patterns, coexpression can reveal relationships between genes and other

transcribed genetic elements [56,57]. Most coexpression studies have focused on cORFs, but

the abundance of publicly available RNA-seq data represents a tractable avenue to interrogate

the transcriptional regulation of thousands of nORFs at once using coexpression approaches

[47,58–61]. Indeed, RNA-seq is probe-agnostic and annotation-agnostic, thereby enabling the

reuse of existing data to explore these novel ORFs. However, low expression levels can distort

coexpression inferences due to statistical biases [62,63]. A coexpression analysis of translated

nORFs that addresses the statistical issues arising from low expression is still lacking for any

species.

Here, we developed a dedicated statistical approach that accounts for low expression levels

when inferring coexpression relationships between ORFs. We applied this approach to the

recently identified 19,000 translated nORFs in *S. cerevisiae* [14] and built the first high-quality

coexpression network spanning the canonical and noncanonical translatome of any species.

Coexpression relationships suggest that the majority of nORFs are transcriptionally regulated.

While many nORFs form entirely new noncanonical transcription modules, approximately half

are transcriptionally associated with genes involved in cellular homeostasis and transport. We

show that *de novo* ORFs that piggyback onto their neighbors' transcription tend to have higher

expression and tend to be highly coexpressed with their neighbors. We provide a web

application to allow researchers to easily access this dataset to investigate the coexpression

relationships and potential cellular roles for thousands of ORFs.

Results

High-quality coexpression inferences show transcriptional and regulatory relationships between nORFs and cORFs

Figure 1: Overview of coexpression inference framework and properties of the dataset

A) Workflow: 3,916 samples were analyzed to create an expression matrix for 11,630 ORFs,

including 5,803 cORFs and 5,827 nORFs; center log ratio transformed (clr) expression values

were used to calculate the coexpression matrix using proportionality metric, ρ, followed by

normalization to correct for expression bias. The coexpression matrix was thresholded using ρ >

0.888 to create a coexpression network (top 0.2% of all pairs). B) Distribution of the number of

ORFs binned based on their median expression values (transcript per million - TPM) and the

number of samples the ORFs were detected in with at least 5 raw counts. C) Coexpressed

cORF pairs (ρ > 0.888) are more likely to encode proteins that form complexes than non-

coexpressed cORF pairs (Fisher's exact test p < 2.2e-16; error bars: standard error of the

proportion); using annotated protein complexes from ref. [64]. D) Coexpressed ORF pairs (ρ >

0.888) are more likely to have their promoters bound by a common transcription factor (TF) than

non-coexpressed ORF pairs (Fisher's exact test p < 2.2e-16; error bars: standard error of the

proportion); genome-wide TF binding profiles from ref. [65] and transcription start sites (TSS)

from ref. [66] were analyzed to define promoter binding (see Methods). E) Hierarchical

clustering of the coexpression matrix reveals functional enrichments for most clusters that

contain at least 5 cORFs; functional enrichments estimated by gene ontology (GO) enrichment

analysis at false discovery rate (FDR) < 0.05 using Fisher's exact test.

To infer coexpression at the translatome scale in *S. cerevisiae*, we considered all cORFs

annotated as "verified", "uncharacterized", or "transposable element" in the *Saccharomyces*

Genome Database (SGD) [67], as well as all nORFs, ORFs that were either unannotated or

annotated as "dubious" and "pseudogene", with evidence of translation according to Wacholder

et al. [14]. To maximize detection of transcripts containing nORFs, we curated and integrated

3,916 publicly available RNA-seq samples from 174 studies (Figure 1A, Supplementary Data 1).

Many nORFs were not detected in most of the samples we collected, creating a very sparse

dataset (Figure 1B). The issue of sparsity has been widely studied in the context of single cell

RNA-seq (scRNA-seq). A recent study looking at multiple measures of association for

constructing coexpression networks from scRNA-seq showed that proportionality methods

coupled with center log ratio (clr) transformation consistently outperformed other measures of

coexpression in a variety of tasks including identification of disease-related genes and protein-

protein network overlap analysis [68]. Thus, we used clr to transform the raw read counts and

quantified coexpression relationships using the proportionality metric, ρ [69].

We further addressed the issue of sparsity with two sample thresholding approaches. First, any

observation with a raw count below five was discarded, such that when calculating ρ only the

samples expressing both ORFs with at least five counts were considered. Second, we

empirically determined that a minimum of 400 samples were required to obtain reliable

coexpression values by assessing the effect of sample counts on the stability of ρ values

(Supplementary Figure 1). These steps resulted in an 11,630 by 11,630 coexpression matrix

encompassing 5,803 cORFs and 5,827 nORFs (ORF list in Supplementary Data 2).

The combined use of clr, ρ, and sample thresholding accounted for statistical issues in

estimating coexpression deriving from sparsity, but the large difference in RNA expression

levels between cORFs and nORFs posed yet another challenge. Indeed, Wang et al. showed

that the distribution of coexpression values is biased by expression level due to statistical

artifacts [62]. We observed this artifactual bias in our dataset (Supplementary Figure 2A) and

corrected for it using spatial quantile normalization (SpQN) as recommended by Wang et al. [62]

(Supplementary Figure 2B). This resulted in a normalized coexpression matrix (Supplementary

Data 3) with ρ values centered around 0.476.

We then created a network representation of the coexpression matrix by considering only the

top 0.2% of ρ values between all ORF pairs (ρ > 0.888). This threshold was chosen to include

90% of cORFs (Supplementary Figure 3). Altogether, our dedicated analysis framework (Figure

1A) inferred 124,382 strong (ρ > 0.888) coexpression relationships between 9,303 ORFs,

encompassing 4,112 nORFs and 5,191 cORFs.

To assess whether our coexpression network captures meaningful biological and regulatory

relationships, we examined its overlap with orthogonal datasets. Using a curated [64] protein

complex dataset for cORFs, we found that coexpressed cORF pairs are significantly more likely

to encode proteins that form a protein complex together compared to non-coexpressed pairs

(Odds ratio = 10.8 Fisher's exact test $p < 2.2e-16$; Figure 1C). Using a previously published [65]

genome-wide chromatin immunoprecipitation with exonuclease digestion (ChIP-exo) dataset

containing DNA-binding information for 73 sequence-specific transcription factors (TFs) and

using transcript isoform sequencing (TIF-seq) [66] data to determine transcription start sites

(TSSs) and promoter regions, we observed that coexpressed ORF pairs were more likely to

have their promoters bound by a common TF than non-coexpressed ORF pairs, whether the

pairs consist of nORFs or cORFs (*canonical-canonical pairs*: Odds ratio = 3.84, *canonical-*

noncanonical pairs: Odds ratio = 2.55, *noncanonical-noncanonical pairs*: Odds ratio = 3.22,

Fisher's exact test $p < 2.2e-16$ for all three comparisons; Figure 1D). Enrichments were robust

to different coexpression cutoffs (Supplementary Figure 4-5). Using the WGCNA [70] method to

cluster the coexpression matrix, we found that more than half of the clusters identified contained

functionally related ORFs (gene ontology (GO) biological process enrichments at Benjamini-

Hochberg (BH) adjusted false discovery rate (FDR) < 0.05; Figure 1E; Supplementary Figure 6).

These analyses demonstrate the high quality of our coexpression network and confirm that it

captures meaningful biological and regulatory relationships for both cORFs and nORFs.

Conventional approaches for coexpression analysis include using transcript per million (TPM) or

reads per kilobase per million (RPKM) normalization, batch correction by removing top principal

components, and Pearson's correlation as the similarity metric [71,56,72]. Compared to these

approaches, our framework increased the proportion of coexpressed ORF pairs whose

promoters are bound by a common TF specifically for pairs containing nORFs (Supplementary

Figure 7), and yielded coexpression networks encompassing the largest number of nORFs at

most thresholds (Supplementary Figure 8). Hence our dedicated analysis framework therefore

outperforms conventional coexpression approaches for the study of nORFs. We offer an R

Shiny [73] interface (https://carvunislab.csb.pitt.edu/shiny/coexpression/) to efficiently query,

visualize and download the coexpression data we generated. To our knowledge, this is the most

comprehensive coexpression dataset focusing on empirically translated elements, both

annotated and unannotated, for any species to date.

nORFs tend to be located at the periphery of the coexpression

network and form new noncanonical transcription modules

Figure 2 Topological properties of the coexpression network

A) Visualization for canonical-only and full coexpression networks using spring embedded graph

layout [74]. The full network contains more cORFs than the canonical-only network since

addition of nORFs also results in addition of many cORFs that are only connected to an nORF.

B) nORFs have fewer coexpression partners (degree in full network) than cORFs (Mann-

Whitney U-test p < 2.2e-16). C) Most cORFs are coexpressed with at least one nORF. D) Only

59% of nORFs are coexpressed with at least one cORFs and this is less than expected by

chance, on average, 89% of nORFs are coexpressed with a cORF across 1,000 randomized

networks generated in a degree-preserving fashion by swapping edges of noncanonical nodes

(Fisher's exact test p < 2.2e-16; error bar: standard error of the mean proportion across

randomized networks). E) Addition of nORFs to the canonical-only network results in the full

network being less compact, whereas the opposite is expected by chance, shown by the

decrease in diameters for the 1,000 randomized networks. F) Addition of nORFs to the

canonical-only network decreases local clustering in the full network, however this is to a lesser

extent than expected by chance as shown by the distribution for the 1,000 randomized

networks. G) Most clusters in the coexpression matrix encompass either primarily nORFs or

primarily cORFs (n= 69 clusters, *green* represents nORF majority clusters, *purple* represents

cORF majority clusters).

Conventional analyses of coexpression networks have been restricted to cORFs. Our full

coexpression network contains twice the number of ORFs and three times the number of strong

($\rho > 0.888$) coexpression relationships compared to the canonical-only network (Figure 2A). We

sought to compare the network properties of the canonical-only and full networks. On average,

nORFs have fewer coexpressed partners (degree) than cORFs, suggesting that nORFs have

distinct transcriptional profiles (Cliff's Delta d = -0.29, Mann-Whitney U-test p < 2.2e-16; Figure

2B). We found that 91% of cORFs are coexpressed with at least one nORF (n = 4,726; Figure

2C), whereas only 59% of nORFs are coexpressed with at least one cORF. In contrast, we

would have expected an average of 89% of nORFs to be coexpressed with a cORF according

to degree preserving simulations of 1,000 randomized networks where edges from nORFs were

shuffled (Odds ratio = 0.174, Fisher's exact test p < 2.2e-16; Figure 2D, Supplementary Figure

9). This suggests that, while most nORFs are integrated in the full coexpression network, they

also have distinct expression profiles that differ markedly from those of all cORFs and are more

similar to those of other nORFs.

To investigate how these seemingly conflicting attributes impact the organization of the

coexpression network, we analyzed two global network properties: diameter, which is the

longest shortest path between any two ORFs; and transitivity, which is the tendency for ORFs

that are coexpressed with a common neighbor to also be coexpressed with each other. The

incorporation of nORFs in the full network led to a larger diameter relative to the canonical-only

network (Figure 2E). This is in sharp contrast with the null expectation, set by 1,000 degree-

preserving simulations, whereby random incorporation of nORFs decreases network diameter.

The full coexpression network is thus much less compact than expected by chance, suggesting

that nORFs tend to be located at the periphery of the network. Network transitivity decreased

with the incorporation of nORFs compared to the canonical-only network, but to a lesser extent

than expected by chance (Figure 2F). This suggests that despite their low degree and

peripheral locations, the connections formed by nORFs are structured and may form

noncanonical clusters.

To investigate this hypothesis, we inspected the ratio of nORFs and cORFs among the cluster

assignments from WGCNA hierarchical clustering of the full coexpression matrix

(Supplementary Figure 6). Strikingly, we observed a bimodal distribution of clusters, with

approximately half of the clusters consisting mostly of nORFs and the other half containing

mostly cORFs (Figure 2G). We conclude that nORFs exhibit a unique and non-random

organization within the coexpression network, simultaneously connecting to all cORFs while

also forming entirely new noncanonical transcription modules.

Coexpression profiles reveal most nORFs are transcriptionally

associated with genes involved in cellular transport and

homeostasis

Figure 3 Biological processes associated with nORF transcriptional regulation

A-B) Biological processes that are more (A) (Odds ratio > 2, n = 16 terms) or less (B) (Odds

ratio < 0.5, n = 23 terms) transcriptionally associated with nORFs than cORFs (y-axis ordered

by nORF enrichment proportion from highest to lowest, BH adjusted FDR < 0.001 for all terms,

Fisher's exact test, GO term enrichments were detected using gene set enrichment analyses

(GSEA), error bars: standard error of the proportion). C) nORFs that are highly coexpressed

with genes involved in transport are more likely to have predicted transmembrane (TM) domains

as determined by TMHMM [75] compared to nORFs that are not (Odds ratio = 1.6, Fisher's

exact test p = 1.3e-4; error bars: standard error of the proportion). D) nORFs and cORFs that

are Sfp1 or Hsf1 targets are more likely to be downregulated when Sfp1 or Hsf1 are deleted

compared to ORFs that are not targets (*Sfp1*: cORFs: p < 2.2e-16; nORFs: p = 2.8e-9; *Hsf1*:

cORFs: p <2.2e-16; nORFs: p = 9.9e-13; Fisher's exact test, error bars: 95% confidence interval

of the odds ratio; *dashed* line shows odds ratio of 1; RNA abundance data from SRA accession

SRP159150 and SRP437124 [76] respectively). E) nORFs that are regulated by TFs are more

likely to be coexpressed with genes involved in processes related to known functions of that TF.

To determine whether nORFs are transcriptionally associated with specific cellular processes,

we performed gene set enrichment analyses [77] (GSEA) on their coexpression partners. GSEA

takes an ordered list of genes, in this case sorted by coexpression level, and seeks to find if the

higher ranked genes are preferentially annotated with specific GO terms. For each cORF and

nORF, we ran GSEA to detect if their highly coexpressed partners were preferentially

associated with any GO terms (Supplementary Figure 10). Almost all ORFs (99.9%), whether

cORF or nORF, had at least one significant GO term associated with their coexpression

partners at BH adjusted FDR < 0.01, suggesting that nORFs are engaged in coherent

transcriptional programs. We then calculated, for each GO term, the number of cORFs and

nORFs with GSEA enrichments in this term (Supplementary Data 4). These analyses identified

specific GO terms that were significantly more (16 terms, BH adjusted FDR < 0.001, Odds ratio

> 2, Fisher's exact test; Figure 3A, Supplementary Data 5) or less (23 terms, BH adjusted FDR

< 0.001, Odds ratio < 2, Fisher's exact test; Figure 3B, Supplementary Data 5) prevalent among

the coexpression partners of nORFs relative to those of cORFs. Most of the GO terms that were

significantly enriched among the coexpression partners of nORFs were related to cellular

homeostasis and transport (Figure 3A) while most of the GO terms significantly depleted among

the coexpression partners of nORFs were related to DNA, RNA, and protein processing (Figure

3B). Running the same GSEA pipeline with Kyoto Encyclopedia of Genes and Genomes

(KEGG) [78] annotations yielded consistent results (Supplementary Figure 11, Supplementary

Data 6-7). Half of nORFs were coexpressed with genes involved in homeostasis (GO:0042592,

53%), monoatomic ion transport (GO:0006811, 49%) and transmembrane transport

(GO:0055085, 47%). The nORFs transcriptionally associated with the parent term 'transport' (n

= 2,718, GO:0006810, GSEA BH adjusted FDR < 0.01) were 1.6 times more likely to contain a

predicted transmembrane domain than other nORFs (p = 1.3e-4, Fisher's exact test; Figure 3C),

in line with potential transport-related activities. These findings reveal a strong and previously

unsuspected transcriptional association between nORFs, and cellular processes related to

homeostasis and transport.

Hsf1 and Sfp1 nORF targets are part of protein folding and

ribosome biogenesis transcriptional programs, respectively

Overall, our analyses relating coexpression to TF binding (Figure 1D) and functional

enrichments (Figure 3A-B) suggest that nORF expression is regulated rather than simply the

consequence of transcriptional noise. To further investigate this hypothesis, we sought to

identify regulatory relationships between specific TFs and nORFs. We reasoned that if nORFs

are regulated by TFs in similar ways as cORFs, then genetic knockout of the TFs that regulate

them should impact their expression levels as it does for cORFs [79]. We focused on two

transcriptional activators for which both ChIP-exo [65] and knockout RNA-seq data [76] were

publicly available: Sfp1, which regulates ribosome biogenesis [80] and Hsf1, which regulates

heat shock and protein folding responses [81].

For both cORFs and nORFs, knockout of Sfp1 or Hsf1 was more likely to trigger a significant

decrease in expression when the ORF's promoter was bound by the respective TF according to

ChIP-exo evidence (Figure 3D). The statistical association between TF binding and knockout-

induced downregulation was as strong for nORFs as it was for cORFs, consistent with nORFs

having similar mechanisms of transcriptional activation (*Sfp1*: cORFs Odds ratio = 11.1, p <

2.2e-16; nORFs Odds ratio = 21.8, p = 2.8e-9, Fisher's exact test; *Hsf1*: cORFs Odds ratio =

12.7, p < 2.2e-16; nORFs Odds ratio = 12.1, p = 9.9e-13, Fisher's exact test). Therefore, the

nORFs whose promoters are bound by these TFs, and whose expression levels decrease upon

deletion of these TFs, are likely genuine regulatory targets of these TFs. By this stringent

definition, our analyses identified 9 nORF targets of Sfp1 (and 34 cORF targets) and 19 nORF

targets of Hsf1 (and 39 cORF targets). The coexpression profiles of these Sfp1 and Hsf1 nORF

targets were preferentially associated with genes involved in processes directly related to the

known functions of Sfp1 and Hsf1 (Supplementary Data 8). For example, the coexpression

profiles of 9 Sfp1 nORF targets revealed preferential associations with genes involved in

'ribosomal large subunit biogenesis' and 7 Sfp1 nORF targets involved in 'regulation of

translation' according to our GSEA pipeline (Fisher's exact test, BH adjusted p-value < 6.7e-4

for both terms). Similarly, 13 Hsf1 nORF targets were preferentially associated with genes

involved in 'Protein Folding' (Fisher's exact test, BH adjusted p-value = 5.7e-9). These results

show that nORF expression can be actively regulated by TFs as part of coherent transcriptional

programs (Figure 3E).

de novo ORF expression and regulation are shaped by genomic

location

352 **Figure 4 Expression, coexpression and biological processes similarity of *de novo* ORFs**

353 **with respect to genomic orientations**

354 A) Pipeline used to reclassify ORFs as conserved or *de novo*. cORFs were considered for both

355 conserved and *de novo* classification while nORFs were only considered for *de novo*

356 classification. Conserved ORFs were determined by either detection of homology outside of

357 *Saccharomyces* or reading frame conservation within *Saccharomyces* (*top*). *De novo* ORFs

358 were determined by evidence of translation, lack of homology outside of *Saccharomyces* as well

359 as lack of a homologous ORF in the two most distant *Saccharomyces* branches (*bottom*). B)

360 Counts of cORFs and nORFs that emerged *de novo*. C) Genomic orientations of *de novo* ORFs

361 that cannot transcriptionally piggyback off neighboring conserved ORF (cannot share promoter

362 with neighbor, *pink shading*) or can transcriptionally piggyback off neighboring conserved ORF

363 (possible to share promoter with neighbor, *green shading*). D) Counts of *de novo* ORFs that are

364 within 500 bp of a conserved ORF in different genomic orientations; ORFs further than 500bp

365 are classified as independent. E) *De novo* ORFs in orientations that can piggyback have higher

366 RNA expression levels than *de novo* ORFs in orientations that cannot piggyback (Cliff's Delta d

367 = 0.4). Only *de novo* ORFs in a single orientation are considered (dashed box in panel *D*).

368 Dashed line represents the median expression of independent *de novo* ORFs. F) *de novo* ORFs

369 in orientations that can piggyback have higher coexpression with neighboring conserved ORFs

370 compared to *de novo* ORFs in orientations that cannot piggyback (Cliff's Delta d = 0.43).

371 Dashed line represents median coexpression of *de novo*-conserved ORF pairs on separate

372 chromosomes. G) *de novo* ORFs in orientations that can piggyback are more likely to be

373 transcriptionally associated with genes involved in the same biological processes as their

374 neighboring conserved ORFs than *de novo* ORFs in orientations that cannot piggyback (Cliff's

375 Delta d = 0.31). Dashed line represents median functional enrichment similarities of *de novo*-

376 conserved ORF pairs on separate chromosomes. (For panels E-F-G: Mann-Whitney U-test, ****:

377 p < 2.2e-16).

Previous literature has shown that many nORFs arise *de novo* from previously noncoding

regions [24,26]. We wanted to investigate how these evolutionarily novel ORFs acquire

expression and whether their locus of emergence influences this acquisition. To define which

ORFs were of recent *de novo* evolutionary origins, we developed a multistep pipeline combining

sequence similarity searches and syntenic alignments (Figure 4A). cORFs were considered

conserved if they had homologues detectable by sequence similarity searches with BLAST in

budding yeasts outside of the *Saccharomyces* genus or if their open reading frames were

maintained within the *Saccharomyce*s genus [14]. cORFs and nORFs were considered *de novo*

if they lacked homologues detectable by sequence similarity outside of the *Saccharomyces*

genus and if less than 60% of syntenic orthologous nucleotides in the two most distant

Saccharomyces branches were in the same reading frame as in *S. cerevisiae*. These criteria

aimed to identify the youngest *de novo* ORFs. Overall, we identified 5,624 conserved cORFs

and 2,756 *de novo* ORFs including 77 *de novo* cORFs and 2,679 *de novo* nORFs (Figure 4B).

In general, the coexpression patterns of *de novo* ORFs (Supplementary Figure 12) were similar

to those of nORFs (Figure 3A-B).

We hypothesized that the locus where *de novo* ORFs arise may influence their expression

profiles through "piggybacking" off their neighboring conserved ORFs' pre-existing regulatory

environment. To investigate this hypothesis, we categorized *de novo* ORFs based on their

positioning relative to neighboring conserved ORFs. The *de novo* ORFs further than 500 bp

from all conserved ORFs were classified as independent. The remaining *de novo* ORFs were

classified as either upstream or downstream on the same strand (up same or down same),

upstream or downstream on the opposite strand (up opposite or down opposite), or as

overlapping on the opposite strand (anti-sense overlap) based on their orientation to the nearest

conserved ORF (Figure 4C-D). We categorized the orientations as being able to piggyback or

unable to piggyback based on their potential of sharing a promoter with neighboring conserved

ORFs, with down opposite and antisense overlap as orientations that cannot piggyback and up

opposite, up same, and down same as orientations that can piggyback (Figure 4C). The

piggybacking hypothesis predicts that *de novo* ORFs that arise in orientations that can

piggyback would be positively influenced by the regulatory environment provided by the

promoters of neighboring conserved ORFs, resulting in similar transcription profiles as their

neighbors and increased expression relative to *de novo* ORFs that do not benefit from a pre-

existing regulatory environment.

We considered three metrics to assess piggybacking: RNA expression level, measured as

median TPM over all the samples analyzed, coexpression with neighboring conserved ORF and

biological process similarity with neighboring conserved ORF. To calculate biological process

similarity between two ORFs, we used significant GO terms at FDR < 0.01 determined by

coexpression GSEA for each ORF (Supplementary Figure 10) and calculated the similarity

between these two sets of GO terms using the relevance method [82]. If two ORFs are enriched

in the same specialized terms, their relevance metric would be higher than if they are enriched

in different terms or in the same generic terms. We found that *de novo* ORFs in orientations that

can piggyback tend to have higher expression (focusing only on ORFs that could be assigned a

single orientation, dashed box in Figure 4D, Cliff's Delta d = 0.4; Figure 4E), higher

coexpression with their neighbor (Cliff's Delta d = 0.43; Figure 4F), and higher biological

process similarity (Cliff's Delta d = 0.31; Figure 4G), compared to ORFs in orientations that

cannot piggyback (p < 2.2e-16 Mann-Whitney U-test for all). Thus, all three metrics supported

the piggybacking hypothesis.

Closer examination revealed a more complex situation. First, the immediate neighbors of *de*

novo ORFs in orientations that can piggyback were rarely among their strongest coexpression

partners (only found in the top 10 coexpressed partners for 15% of down same, 4.5% of up

same, 3% of up opposite ORFs). Therefore, emergence nearby a conserved ORF in a

piggybacking orientation influences, but does not fully determine, the transcription profiles of *de*

novo ORFs. Transcriptional regulation beyond that provided by the pre-existing regulatory

environment may exist. Second, while ORFs in all three orientations that can piggyback

displayed increased coexpression and biological process similarity with their neighbors relative

to background expectations (Supplementary Figure 13A-B), only down same *de novo* ORFs

displayed increased RNA expression levels (Supplementary Figure 13C). The expression levels

of up same *de novo* ORFs were statistically indistinguishable from independent *de novo* ORFs,

while those of up opposite *de novo* ORFs were significantly lower than those of independent *de*

novo ORFs (Supplementary Figure 13C). Down same *de novo* ORFs also showed stronger

coexpression and biological process similarity with their conserved neighbors than up same and

up opposite *de novo* ORFs (Supplementary figure 13A-B). Therefore, the transcription of down

same *de novo* ORFs appeared most susceptible to piggybacking.

Figure 5 Effects of promoter sharing on expression, coexpression and biological process similarities of *de novo* ORFs

A) *De novo* ORFs that share a promoter with neighboring conserved ORFs, as determined by TIF-seq transcript boundaries, have significantly higher expression levels than *de novo* ORFs that do not. Considering only ORFs in a single orientation. Dashed line represents the median expression of independent *de novo* ORFs. B) *De novo* ORFs that share a promoter with neighboring conserved ORFs have higher coexpression with their neighbors than *de novo* ORFs that do not share a promoter. Dashed line represents median coexpression of *de novo*-conserved ORF pairs on separate chromosomes. C) *De novo* ORFs that share a promoter have more similar functional enrichments with neighboring conserved ORFs than *de novo* ORFs that do not share a promoter. Dashed line represents median functional enrichment similarities of the background distribution of *de novo*-conserved ORF pairs on separate chromosomes. D) Down same *de novo* ORFs share a promoter with neighboring conserved ORFs significantly more often than up same ORFs. E) Conserved ORFs with downstream *de novo* ORFs have a significant increase in expression compared to conserved ORFs with upstream *de novo* ORFs. F) Existence of transcription termination factors (Pcf11 or Nrd1) in between conserved ORFs and nearby downstream *de novo* ORFs leads to less shared transcripts. G) Transcript isoforms (*gray*) at an example locus where there are no transcription termination factors present between conserved ORF YBL015W (*pink*) and downstream *de novo* ORF chr2:195794-195847(+) (*blue*). H) Transcript isoforms (*gray*) at an example locus where there is Pcf11 transcription terminator present (*red line*) between conserved ORF YPR034W (*pink*) and downstream *de novo* ORF chr16:641385-641534(+) (*blue*). All detected transcript isoforms on these loci are plotted for G and F. (For all panels: ****: $p \leq 0.0001$, ***: $p \leq 0.001$, **: $p \leq 0.01$, *: $p \leq 0.05$, ns: not-significant; Mann-Whitney U-test)

To understand the molecular mechanisms leading to the differences in expression,

coexpression and biological process similarity between the orientations that can piggyback,

which all have the potential to share a promoter with neighboring conserved ORF, we

investigated which actually do by analyzing transcript architecture. Using a publicly available

TIF-seq dataset [66], we defined down same or up same ORFs as sharing a promoter with their

neighbor if they mapped to the same transcript at least once. We defined up opposite ORFs as

sharing a promoter with their neighbor if their respective transcripts did not have overlapping

TSSs, as would be expected for divergent promoters [83]. According to these criteria, 84% of

down same (n = 174), 64% of up same (n = 368), and 66% of up opposite (n = 185) *de novo*

ORFs share a promoter with their neighboring conserved ORFs (Supplementary Figure 14).

Among all *de novo* ORFs that arose in orientations that can piggyback, those that share

promoters with neighboring conserved ORFs displayed higher expression levels than those that

do not (*down same*: d = 0.75, p = 1.06e-8; *up same*: d = 0.38, p = 1.23e-7; *up opposite*: d = 0.3,

p = 2.9e-3 Mann-Whitney U-test, d: Cliff's Delta; Figure 5A). We also observed a significant

increase in coexpression and biological process similarity between *de novo* ORFs and their

neighboring conserved ORFs when their promoters are shared compared to when they are not

(coexpression: *down same*: d = 0.28, p = 2.99e-9; *up same*: d = 0.31, p < 2.2e-16; *up opposite*:

d = 0.27, p = 2.1e-7; biological process similarity: *down same*: d = 0.24, p = 5.5e-7; *up same*: d

= 0.108, p = 3.78e-3; *up opposite*: d = 0.24, p = 6.1e-6, d: Cliff's Delta, Mann-Whitney U-test;

Figures 5B and 5C, respectively). Hence, sharing a promoter led to increases in the three

piggybacking metrics for the three orientations.

Further supporting the notion that down same ORFs are particularly prone to piggybacking, the

down same *de novo* ORFs that share a promoter with their conserved neighbors displayed

much higher expression levels, and higher coexpression and biological process similarity with

their conserved neighbor, than up same or up opposite ORFs that also share a promoter with

their conserved neighbors (expression: *down same vs up same*: d = 0.58; *down same vs up*

opposite: d = 0.55; coexpression: *down same vs up same:* d = 0.29, *down same vs up opposite:*

d = 0.38; biological process similarity: *down same vs up same*: d = 0.37, *down same vs up*

opposite: d = 0.45; d: Cliff's Delta, p < 2.2e-16 for all comparisons, Mann-Whitney U-test). This

could be due to down same ORF's tendency to share promoters more often than up same

ORFs, as a larger proportion of transcripts containing down same ORFs also contain a

conserved ORF (*down same vs up same*: Cliff's Delta d = 0.26, Mann-Whitney U-test p < 2.2e-

16; Figure 5D), or higher expression levels of conserved ORFs that have down same ORFs on

their transcripts compared to conserved ORFs with up same or up opposite piggybacking ORFs

(*down same vs up same*: d = 0.2, p = 5.4e-3; *down same vs up opposite*: d = 0.34, p = 6.5e-4,

Mann-Whitney U-test, d: Cliff's Delta; Figure 5E).

Based on these results, we reasoned that transcriptional readthrough could be the molecular

mechanism underlying the efficient transcriptional piggybacking of down same *de novo* ORFs.

To investigate this hypothesis, we examined the impact of transcription terminators Pcf11 or

Nrd1 on the frequency of transcript sharing between a conserved ORF and its downstream *de*

novo ORF. Analyzing publicly available ChIP-exo data [65], we found that the presence of

terminators between conserved ORFs and their downstream *de novo* ORF pairs resulted in a

notably lower percentage of shared transcripts (Cliff's Delta d = -0.39, p = 1.59e-10, Mann-

Whitney U-test; Figure 5F). As an illustration, consider the genomic region on chromosome II

from bases 194,000 to 196,000, containing the conserved ORF YBL015W and a downstream

de novo ORF (positions 195,794 to 195,847). No terminator factor is bound to the intervening

DNA between these two ORFs. This pair has high coexpression, with ρ = 0.96 and we observed

that nearly all transcripts in this region containing the *de novo* ORF also include YBL015W

(Figure 5G). In contrast, the genomic region on chromosome XVI from 639,000 to 641,800,

containing the conserved ORF YPR034W and downstream *de novo* ORF (positions 641,385 to

641,534), does have a Pcf11 terminator factor between the pair, and as expected, none of the

transcripts in this region contain both YPR034W and the *de novo* ORF, which have poor

coexpression as a result (ρ = 0.1; Figure 5H). We conclude that sharing a transcript via

transcriptional readthrough is the major transcriptional piggybacking mechanism for down same

de novo ORFs.

Discussion

We explored the transcription of nORFs from multiple angles including network topology,

associations with cellular processes, TF regulation, and influence of the locus of emergence on

de novo ORF expression. Delving into network topology, we find that nORFs have distinct

expression profiles that are strongly correlated with only a few other ORFs. Nearly all cORFs

are coexpressed with at least one nORF, but the converse is not true. Numerous nORFs form

new structured transcriptional modules, possibly involved in both known and unknown cellular

processes. The addition of nORFs to the cellular network resulted in a more clustered network

than expected by chance, highlighting the previously unsuspected influence of nORFs in

shaping the coexpression landscape.

Our study is the first to show a large-scale association between the expression of nORFs and

cellular homeostasis and transport processes. We anticipate that future studies will follow up to

test these associations experimentally. We also found nORFs to be preferentially associated

with cellular processes related to metabolism, transposition and cell adhesion, but rarely with

the core processes of the central dogma, DNA, RNA or protein processing. Genes involved in

transport, metabolism, and stress tend to have more variable expression compared to genes in

other pathways [84]. Pathways with more variable expression could be more likely to

incorporate novel ORFs, possibly as a form of adaptive transcriptional response. There are

several consistent observations in the literature [47,85,86]. For instance, Li et al. [47] showed

that many *de novo* ORFs are upregulated in heat shock. Wilson and Masel [87] found higher

translation of *de novo* ORFs under starvation conditions. Carvunis et al. [24] found *de novo*

cORFs are enriched for the GO term 'response to stress'. Other studies showed examples of

how specific *de novo* ORFs could be involved in stress response [35,88] or homeostasis

[88,89]. For instance the *de novo* antifreeze glycoprotein AFGP allows Arctic codfish to live in

colder environments [35] or *MDF1* in yeast [88,90] was found in a screen to provide resistance

to certain toxins and mediates ion homeostasis [91]. Our results, combined with these previous

investigations, argue that a large fraction of nORFs provide adaptation to stresses and help

maintain homeostasis, perhaps through modulation of transport processes.

Recent research in yeast has revealed an enrichment of transmembrane domains [15,24,92,93]

within *de novo* ORFs. Previous studies identified small nORFs and *de novo* ORFs that localize

to diverse cellular membranes, such as those of the endoplasmic reticulum, Golgi, or

mitochondria in different species [10,15,94–97]. These findings are consistent with the notion

that *de novo* ORFs could play a role in a range of transport processes, such as ion, amino acid,

or protein transport across cellular membranes. By establishing a connection between predicted

transmembrane domains and increased coexpression with transport-related genes, our findings

set the stage for future experimental investigations into the precise molecular mechanisms and

functional roles of nORFs in diverse transport systems.

Lastly, we explored how the preexisting regulatory context influences the transcriptional profiles

of *de novo* ORFs. We found that *de novo* ORFs that piggyback off their neighboring conserved

ORFs' promoters had increases in expression, coexpression and biological process similarity

with their neighboring conserved ORFs. Strikingly, ORFs that emerge *de novo* downstream of

conserved ORFs have the largest increases in expression, coexpression and biological process

similarities with their neighbors compared to other orientations, largely due to transcriptional

readthrough leading to transcript sharing. Previous studies have shown that the transcription of

regions downstream of genes is functional and regulated [98]. A study in humans showed that

readthrough transcription downstream of some genes is responsible for roughly 15%–30% of

intergenic transcription and is induced by osmotic and heat stress creating extended transcripts

that play a role in maintaining nuclear stability during stress [99]. Another study in humans and

zebrafish showed that the translation of small ORFs located in the 3' UTR of mRNAs (dORFs)

increased the translation rate of the upstream gene [100]. Lastly, a study in yeast found that

genes which are preferentially expressed as bicistronic transcripts tend to contain evolutionarily

younger genes compared to adjacent genes that do not share transcripts, suggesting that

transcript sharing could provide a route for novel ORFs to become established genes [101].

These findings together with our results suggest that genomic regions downstream of genes

may provide the most favorable environment for the transcription of *de novo* ORFs.

Our analyses show that the likelihood of a *de novo* ORF being expressed or repressed under

the same conditions as the neighboring conserved ORF is influenced by the extent to which it

piggybacks on the neighboring ORF's regulatory context. Therefore, in addition to the

evolutionary pressure acting on the sequence of emerging ORFs, our results suggest that

transcriptional regulation and genomic context also influence their functional potential. However,

this influence is not entirely deterministic, and much weaker when *de novo* ORFs emerge

upstream than downstream of genes. Future studies are needed to map regulatory networks

controlling nORF expression and reconstruct their evolutionary histories.

There are several limitations to our study. First, while SpQN enhances the coexpression signal

of lowly expressed ORFs, it comes at the cost of reducing signals in highly expressed ORFs

[62]. Given our objective of studying lowly-expressed nORFs this tradeoff is deemed worthwhile.

Second, our study provides evidence of associations between nORFs and cellular processes

such as homeostasis and transport, but these findings are based on transcription profile

similarities which do not necessarily imply cotranslation or correlated protein abundances [102].

Furthermore, our analyses were performed in the yeast *S. cerevisiae* and the generalizability of

our findings to other species requires further investigation.

603 Conclusions

In conclusion, our study represents a significant step forward towards the characterization of

nORFs. We employed advanced statistical methods to account for low expression levels and

generate a high-quality coexpression network. Despite being lowly expressed, nORFs are

coexpressed with almost every cORF. We find that numerous nORFs form structured,

noncanonical-only transcriptional modules which could be involved in regulating novel cellular

processes. We find that many nORFs are coexpressed with genes involved in homeostasis and

transport related processes, suggesting that these pathways are most likely to incorporate novel

ORFs. Additionally, our investigation into the influence of genomic orientation on the expression

and coexpression of *de novo* ORFs showed that ORFs located downstream of conserved ORFs

are most influenced by the pre-existing regulatory environment at their locus of emergence. Our

findings provide a foundation for future research to further elucidate the roles of nORFs and *de*

novo ORFs in cellular processes and their broader implications in adaptation and evolution.

Methods

Creating ORF list

To create our initial ORF list, we utilized two sources. First, we took annotated ORFs in the *S. cerevisiae* genome R64-2-1 downloaded from SGD [103], which included 6,600 ORFs. Second, we utilized the translated ORF list from Wacholder et al. [14] reported in their *Supplementary Table 3.* We filtered to include cORFs (Verified, Uncharacterized or Transposable element genes) as well as any nORFs with evidence of translation at q value < 0.05 (Dubious, Pseudogenes and unannotated ORFs). We removed ORFs with lengths shorter than the alignment index kmer size of 25nt used for RNA-seq alignment. In situations where ORFs overlapped on the same strand with greater than 75% overlap of either ORF, we removed the shorter ORF using bedtools [104]. We removed ORFs that were exact sequence duplicates of another ORF. This left 5,878 cORFs and 18,636 nORFs, for a total of 24,514 ORFs used for RNA-seq alignment.

RNA-seq data preprocessing

Strand specific RNA-seq samples were obtained from the Sequencing Read Archive (SRA) using the search query *(saccharomyces cerevisiae[Organism]) AND rna sequencing*. Each study was manually inspected and only studies that had an accompanying paper or detailed methods on Gene Expression Omnibus (GEO) were included. Samples were quality controlled (nucleotides with Phred score < 20 at end of reads were trimmed) and adapters were removed using TrimGalore version 0.6.4 [105]. Samples were aligned to the transcriptome GTF file containing the ORFs defined above and quantified using Salmon [106] version 0.12.0 with an index kmer size of 25. Samples with less than 1 million reads mapped or unstranded samples

were removed, resulting in an expression dataset of 3,916 samples from 174 studies

(Supplementary Data 1). ORFs were removed to limit sparsity and increase the number of

observations in the subsequent pairwise coexpression analysis. Only ORFs that had at least

400 samples with a raw count > 5 were included for downstream coexpression analysis, n =

11,630 ORFs (5,803 canonical and 5,827 noncanonical, Supplementary Data 2).

Coexpression calculations

The raw counts were transformed using clr. Pairwise proportionality was calculated using ρ [69]

for each ORF pair. Spatial quantile normalization (SpQN) [62] of the coexpression network was

performed using the mean clr expression value for each ORF as confounders to correct for

mean expression bias, which resulted in similar distributions of coexpression values across

varying expression levels (Supplementary Figure 2). Only ORF pairs that had at least 400

samples expressing both ORFs (at raw >5) were included. This threshold was determined

empirically as detailed below.

Since zero values cannot be used with log ratio transformations, all zeros must be removed

from the dataset. Proposed solutions in the literature on how to remove zeros, all of which have

their pros and cons, include removing all genes that contain any zeros, imputing the zeros, or

adding a pseudo count to all genes [107,108]. Removing all ORFs that contain any zeros is not

possible for this analysis since the ORFs of interest are lowly and conditionally expressed. The

addition of pseudocounts can be problematic when dealing with lowly expressed ORFs, for the

addition of a small count is much more substantial for an ORF with a low read count compared

to an ORF with a high read count [109]. For these reasons, all raw counts below 5 were set to

NA prior to clr transformation. These observations were then excluded when calculating the clr

transformation and in the ρ calculations. We used clr and ρ implementations in R package *Propr*

[69] and implementation of SpQN from Wang et al. [62].

To determine the minimum number of samples needed expressing both ORFs in a pair, we

determined the number of samples needed for coexpression values to converge within ρ ± 0.05

or ρ ± 0.1 for 2,167 nORF-cORF pairs which have a ρ > 99th percentile (before SpQN). All

samples expressing both ORFs in a pair were randomly binned into groups of 10, and ρ was

calculated after each addition of another sample. Fluctuations were calculated as max(ρ)-min(ρ)

within a sample bin. Convergence was determined as the first sample bin with fluctuations ≤

fluctuation threshold, either 0.05 or 0.01 (Supplementary Figure 1).

671 Comparing coexpression inference approaches

To compare our approach with a batch correction approach, we used clr to transform the

expression matrix, followed by removing the top principal component (PC1) of the clr expression

matrix to do batch correction using the function *removePrincipalComponents* from the *WGCNA*

[70] R package. We then calculated ρ values and applied SpQN normalization. Additionally, we

created a coexpression matrix based on TPM as well as RPKM normalized expression values

instead of clr and calculated Pearson's correlation coefficient.

678 Protein Complex enrichments

We retrieved a manually curated list of 408 protein complexes in *S. cerevisiae* from the

CYC2008 database by Pu et al. [64]. The coexpression matrix was filtered to contain only the

1,617 cORFs found in the CYC2008 database prior to creating the contingency table. Fisher's

exact test was used to calculate the significance of the association between coexpression and

protein complex formation. Coexpressed was defined as the 99.8th ρ percentile (ρ > 0.888)

considering all ORF pairs in the coexpression matrix (n = 62,204,406 ORF pairs) for Figure 1C.

685 TF binding enrichments

A ChIP-exo dataset from Rossi et al. [65] containing DNA-binding information for 73 sequence-

specific TFs across the whole genome was used. For each ORF we identified which TFs had

binding within 200 bp upstream of the ORF's TSS. The TSSs for all ORFs in the coexpression

matrix was determined by the median 5' transcript isoform (TIF) start positions using TIF-seq

[66] dataset. Only ORFs found in the TIF-seq dataset were considered (n = 5,334 cORFs and

5,362 nORFs). To calculate the enrichments reported in Figures 1D, Supplementary Figure 5

and Supplementary Figure 7, the coexpression matrix was first filtered to only include ORFs that

have at least 1 TF binding within 200 bp upstream of its TSS (n = 973 cORFs and 936 nORFs).

Fisher's exact test was used to calculate the association between coexpression and having their

promoters bound by a common TF. Coexpressed was defined as the 99.8th ρ percentile (ρ >

0.888) considering all ORF pairs in the coexpression matrix (n = 62,204,406 ORF pairs) for

Figure 1D.

698 Coexpression matrix clustering

We used the weighted gene coexpression network analysis (*WGCNA*) package [70] in R to

cluster our coexpression matrix. To do this, we first transformed our coexpression matrix into a

weighted adjacency matrix by applying a soft thresholding which involved raising the

coexpression matrix to the power of 12. This removed weak coexpression relationships from the

matrix. We then used the topological overlap matrix (TOM) similarity to calculate the distances

between each column and row of the matrix. Using the *hclust* function in R with the *ward*

clustering method, we created a hierarchical clustering dendrogram. We then used the dynamic

tree cutting method within the *WGCNA* package to assign ORFs to coexpression clusters,

resulting in 73 clusters of which 69 were mapped to the full coexpression network. ORFs in the

other four clusters were not included in the network as they did not pass the ρ threshold.

709 GO analysis of clusters

We downloaded GO trees (file: go-basic.obo) and annotations (files: sgd.gaf) from ref. [110]. We

used the Python package, *GOATools* [111], to calculate the number of genes associated with

each GO term in a cluster and the overall population of (all) genes in the coexpression matrix.

We excluded annotations based on the evidence codes ND (no biological data available). We

identified GO term enrichments by calculating the likelihood of the ratio of the cORFs associated

with a GO term within a cluster given the total number of cORFs associated with the same GO

term in the background set of all cORFs in the coexpression matrix. We applied Fisher's exact

test and FDR with BH multiple testing correction [112] to calculate corrected p-values for the

enrichment of GO term in the clusters. FDR < 0.05 was taken as a requirement for significance.

We applied GO enrichment calculations only when there were at least 5 cORFs in the cluster

(n=54).

721 Network randomization and topology analyses

To create random networks while preserving the same degree distribution, we used an edge

swapping method (Supplementary Figure 9). This involved randomly selecting two edges in the

network, which were either cORF-nORF or nORF-nORF edges and swapping them. The swap

was accepted only if it did not disconnect any nodes from the network and the newly generated

edges were not already present in the network. We repeated this process for at least ten times

the number of edges in the network. Network diameter and transitivity were calculated using R

package *igraph* [113] and networks were plotted using spring embedded layout [74] in Python

package *networkx* [114].

Gene set enrichment analysis

Gene set enrichment analysis (GSEA) calculates enrichments of an ordered list of genes given

a biological annotation such as GO or KEGG. For each ORF in our dataset, we used ρ values to

order annotated ORFs and provided this sorted set to *fgsea* [115] . We used the GO slim file

downloaded from SGD [103] for GO annotations. We used *clusterProfiler* [116] R package to

download KEGG annotations using KEGG REST API [78] on 1 April 2023 and then used

fgseaMultilevel function in *fgsea* R package to calculate enrichments for both annotations

individually. To calculate GO or KEGG terms that are enriched or depleted for nORFs compared

to cORFs, we calculated the number of cORFs and nORFs that had GSEA enrichments at BH

adjusted FDR < 0.01. Using these counts we calculated the proportion of nORFs and cORFs

associated with a GO or KEGG term and used Fisher's exact test to assess the significance of

association. P values returned by Fisher's exact test were corrected for multiple hypothesis

testing using BH correction. Odds ratios were calculated by dividing proportion of nORFs to

proportion of cORFs. Proportions for the GO terms with BH adjusted FDR < 0.001 and Odds

ratio greater than 2 or less than 0.5 are plotted in Figures 3A-B and are reported in

Supplementary Data 5 and proportions for KEGG terms are plotted in Supplementary Figure 11

and reported in Supplementary Data 6.

Transmembrane domain enrichment

Transmembrane domains were predicted using TMHMM 2.0 [75] for all nORFs. An ORF was

classified as having a transmembrane domain if it was predicted to have at least one

transmembrane domain. nORFs were classified as "coexpressed with transport-related genes" if

the ORF had a GSEA enrichment at FDR < 0.01 with any of the 15 GO slim transport terms:

transport, ion transport, amino acid transport, lipid transport, carbohydrate transport, regulation

of transport, transmembrane transport, vacuolar transport, vesicle-mediated transport,

endosomal transport, nucleobase-containing compound transport, Golgi vesicle transport,

nucleocytoplasmic transport, nuclear transport, or cytoskeleton-dependent intracellular

transport. Fisher's exact test was used to calculate the significance of association between

transport-related processes and transmembrane domain.

758   Differential expression analysis for TF deletion and

759   overrepresentation tests

For Hsf1 analysis, RNA-seq samples were from Ciccarelli et al. (SRA accession SRP437124)

[76]. Hsf1 deletion strains were compared to wild type (WT) strains when exposed to heat shock

conditions. For Sfp1 analysis, RNA-seq samples were from SRA accession SRP159150. In both

cases, deletion strains were compared to WT strains. Differential expression was calculated

using R package *DESeq2* [117], and ORFs were defined as differentially expressed if the log

fold change (FC) in RNA expression between WT and control strains was greater than or less

than 0.5 i.e. log(FC) > 0.5 or log(FC) < -0.5 and BH adjusted p-value < 0.05. ChIP-exo data for

Hsf1 and Sfp1 binding was taken from Rossi et al. [65] and an ORF was labeled as having Hsf1

or Sfp1 binding if the TF was found within 200 bp upstream of the ORF's TSS. Fisher's exact

test was performed to see if there is an association between an nORF in a GO biological

process and being regulated by the TF. We define an nORF to be "in" a GO term if it has a

GSEA enrichment for that GO term at FDR < 0.01. We defined an nORF as regulated by a TF if

the nORF had evidence of the TF binding within 200 bp of the nORF's TSS in ChIP-exo and has

significantly downregulated expression in the TF deletion RNA-seq samples compared to the

WT samples. BH p-value correction was performed for all GO terms tested. Significant GO

terms and the associated regulated nORFs are reported in Supplementary Data 8.

776 Detection of homologs using BLAST

We obtained the genomes of 332 budding yeasts from Shen et al. [118]. To investigate the

homology of each non overlapping ORF in our dataset, we used TBLASTN and BLASTP [119]

against each genome in the dataset, excluding the *Saccharomyces* genus. Default settings

were used, with an e-value threshold of 0.0001. The BLASTP analysis was run against the list

of protein coding genes used in Shen et al., while the TBLASTN analysis was run against each

entire genome. We also applied BLASTP to annotated ORFs within the *S. cerevisiae* genome to

identify homology that could be caused by whole genome duplication or transposons.

784 Identification of *de* novo and conserved ORFs

To identify *de novo* ORFs, we applied several strict criteria. Firstly, we obtained translation q-

values and reading frame conservation (RFC) data from Wacholder et al. [14]. All cORFs and

only nORFs with a translation q-value less than 0.05 were considered as potential *de novo*

candidates. We excluded ORFs that overlapped with another cORF on the same strand or had

TBLASTN or BLASTP hits outside of the *Saccharomyces* genus at e-value < 0.0001. Moreover,

we eliminated ORFs that had BLASTP hits to another cORF in *S. cerevisiae*. From the

remaining list of candidate *de novo* ORFs, we investigated whether their ancestral sequence

could be noncoding. To do this, we utilized RFC values for each species within *Saccharomyces*

genus. We classified ORFs as *de novo* if the RFC values for the most distant two branches

were less than 0.6, suggesting the absence of a homologous ORF in those two species.

We identified conserved ORFs if a nonoverlapping cORF has an average RFC > 0.8 or has

either TBLASTN or BLASTP hit at e-value < 0.0001 threshold.

To identify conserved cORFs with overlaps we first considered if the cORFs had a BLASTP

outside of *Saccharomyces* genus with e-value < 0.0001. Then for two overlapping ORFs, if one

had RFC > 0.8 and the other had RFC < 0.8, we considered the one with higher RFC as

conserved. For the ORF pairs that were not assigned as conserved using these two criteria, we

applied TBLASTN for the non-overlapping parts of the overlapping pairs. Those with a

TBLASTN hit with e-value < 0.0001 were considered conserved. We found a total of 5,624

conserved ORFs and 2,756 *de novo* ORFs.

Calculation of GO term similarities

GO term similarities were calculated using the Relevance method developed in Schlicker et al.

[82]. This method considers both the information content (IC) of the GO terms that are being

compared and the IC of their most informative ancestor. IC represents the frequency of a GO

term; thus, an ancestral GO term has lower IC than a descendant. We used the *GOSemSim*

[120] package in R that implements these similarity measures.

Termination factor binding analysis

ChIP-exo data for Pcf11 and Nrd1 termination factor binding sites are taken from Rossi et al. [65]. This study reports binding sites at base pair resolution for *S. cerevisiae* for around 400 proteins. We used supplementary bed formatted files for Pcf11 and Nrd1, which are known transcriptional terminators, and used in house R scripts to find binding sites within the regions between the stop codon of conserved ORFs and the start codon of down same *de novo* ORFs. ORF pairs were classified as having terminators present between them if there was either Pcf11 or Nrd1 binding.

Determining shared promoters

To determine whether two ORFs shared a promoter, we reused the TIF-Seq dataset from Pelechano et al. [66]. TIF-Seq is a sequencing method that detects the boundaries of TIFs. We extracted all reported TIFs from the supplementary data file S1 and identified all TIFs that fully cover each ORF in both YPD and galactose. We then used this information to find ORF pairs that mapped to the same TIFs for down same and up same pairs, as well as found TIFs with non-overlapping TSSs for up opposite *de novo*-conserved ORF pairs. ORF pairs where the conserved ORF was not found in the TIF-seq dataset were not included and pairs where the *de novo* ORF was not found were considered to not share a promoter.

Web application

We utilized R language [121] and the shiny framework [73] to develop a web application which allows querying of ORFs in our dataset for information about their coexpression with other ORFs, network visualization, and GSEA enrichments. It can be accessed through a web browser and is available at https://carvunislab.csb.pitt.edu/shiny/coexpression/.

832 Acknowledgments

833 Figures 1A, 4A, 4C, and Supplementary Figure 10 were created with BioRender.com. The

834 authors are grateful to Dr. Aaron Wacholder, Carly Houghton, Nelson Coelho, Dr. Saurin Bipin

835 Parikh, Jiwon Lee, Lin Chou, Alistair Turcan, Dr. Nikolaos Vakirlis, and Dr. Maria Chikina for

836 reviewing the manuscript prior to submission.

837 Author Contributions

838 Conceptualization: A.R., O.A., and A.-R.C.; Methodology: A.R, O.A.; Investigation: A.R, O.A.;

839 Writing-original draft: A.R, O.A.; Writing-review and editing: A.R., O.A., and A.-R.C.;

840 Supervision: A.-R.C. All authors approved the final version of the manuscript.

841 Funding

842 This work was supported by: the National Science Foundation under Grant No. 2144349

843 awarded to A.-R.C and the National Science Foundation Graduate Research Fellowship under

844 Grant No. 2139321 awarded to A.R. The funders had no role in study design, data collection

845 and analysis, decision to publish, or preparation of the manuscript.

846 Source code

847 All source codes for the analyses conducted are accessible online at

848 https://www.github.com/oacar/noncanonical_coexpression_network

Ethics Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

A.-R.C. is a member of the scientific advisory board for ProFound Therapeutics (Flagship Labs 69, Inc).

Supplementary Data

Supplementary data files are available on Figshare https://doi.org/10.6084/m9.figshare.22289614

Supplementary Data 1: RNA-seq studies and samples used in this study. (CSV)

Supplementary Data 2: ORFs included in the coexpression matrix. (CSV)

Supplementary Data 3: Coexpression matrix generated in this study. (CSV)

Supplementary Data 4: GSEA analysis results for each ORF using GO BP annotations. (CSV)

Supplementary Data 5: List of GO BP terms that are more associated with nORFs than cORFs

and statistics. (CSV)

Supplementary Data 6: List of KEGG terms that are more associated with nORFs than cORFs

and statistics. (CSV)

Supplementary Data 7: GSEA analysis results for each ORF using KEGG annotations. (CSV)

Supplementary Data 8: GO BP terms where nORFs are regulated by either Hsf1 or Sfp1 in GO

BP terms are overrepresented. (CSV)

871 Supplementary Figures

872 Supplementary Figure 1

Supplementary Figure 1 To understand the effect of sample size on coexpression values and to

determine how many samples is sufficient for ρ to converge, we recalculated coexpression for a

given ORF pair using n = 2 samples through n = all samples. Fluctuations were calculated as

max(ρ)-min(ρ) within bins of 10 samples. The number of samples needed for ρ to converge was

calculated as the first sample bin where ρ fluctuations ≤ fluctuation threshold, either 0.1 or 0.05.

Histogram showing the minimum number of samples needed for ρ values to converge within ρ ±

0.05 (*left*) and ρ ± 0.1 (*right*) for 2,167 cORF-nORF pairs with ρ > 99th percentile. Red dashed

lines show the median number of samples needed.

Supplementary Figure 2

Supplementary Figure 2 Distribution of coexpression values (ρ) for ORF pairs binned by

expression level, from lowly expressed pairs *top* to highly expressed pairs *bottom*, A) before

spatial quantile normalization (SpQN) and B) after SpQN, which normalizes the coexpression

values so that the distribution within each expression bin is similar.

Supplementary Figure 3

Supplementary Figure 3 Network threshold affects cORFs and nORFs differently. *Left* shows

the proportion of cORFs or nORFs in the network at each quantile threshold and the *right* shows

the number of connections in the network. Dashed line represents 0.9998 quantile which was

chosen for creating the network.

Supplementary Figure 4

Supplementary Figure 4 Coexpressed cORFs pairs are more likely to encode proteins that form

protein complexes than non-coexpressed cORF pairs, and this is consistent across different

coexpression cutoffs. Coexpression was defined using the top 90th, 95th, 99th, and 99.9th

percentile of all ORF pairs in the network (n = 62,204,406 ORF pairs). 90th percentile (ρ >

0.713) Odds ratio = 8.89; 95th percentile (ρ > 0.763) Odds ratio = 9.59; 99th percentile (ρ >

0.836) Odds ratio = 9.23; 99.9th percentile (ρ > 0.906) Odds ratio = 12.1; Fisher's exact test p <

2.2e-16 for all comparisons. Numbers above bars represent the number of ORF pairs in each

category. Error bars represent the standard error of the proportion. A list of 408 protein

complexes were retrieved from Pu et al. CYC2008 database [64]. Enrichments were calculated

using only the 1,617 cORFs found in the CYC2008 database.

Supplementary Figure 5

Supplementary Figure 5 Coexpressed ORF pairs are more likely to have their promoters bound

by a common TF than non-coexpressed ORF pairs, and this is true across different

coexpression cutoffs and for canonical-canonical (cc), canonical-noncanonical (cn) and

noncanonical-noncanonical (nn) ORF pairs. Coexpression was defined using the top 90th, 95th,

99th, and 99.9th percentile of all ORF pairs in the network (n = 62,204,406 ORF pairs). 90th

percentile (ρ > 0.713): cc Odds ratio = 2.08, cn Odds ratio = 1.42, nn Odds ratio = 1.38; 95th

percentile (ρ > 0.763): cc Odds ratio = 2.38, cn Odds ratio = 1.50, nn Odds ratio = 1.45; 99th

percentile (ρ > 0.836): cc Odds ratio = 3.19, cn Odds ratio = 1.85, nn Odds ratio = 1.82; 99.9th

percentile (ρ > 0.906): cc Odds ratio = 4.57, cn Odds ratio = 3.10, nn Odds ratio = 4.29; ****:

Fisher's exact test p < 2.2e-16 for all comparisons. Error bars represent the standard error of

918   the proportion. Using a ChIP-exo dataset from Rossi et al. [65] containing DNA-binding

919   information for 73 sequence-specific TFs, TF binding was defined as a ChIP-exo peak within

920   200 bp upstream of the ORF's TSS. Only ORFs whose promoter was bound by at least one TF

921   were considered. Numbers above bars represent the number of ORF pairs in each category.

922   # Supplementary Figure 6

923   Supple

mentary Figure 6 Clustered matrix heatmap. Coexpression values are first transformed by

taking power of 12 and then WGCNA pipeline [70] is applied. Clusters are determined by cutting

dendrograms (see methods for details). Colors on 'clusters' section represent the different

clusters. Values of 0.3 and above are represented by red to show the structure of the heatmap.

Supplementary Figure 7

Supplementary Figure 7 Using clr normalization, ρ similarity metric and SpQN normalization

leads to the highest odds ratios for nORF-nORF coexpressed pairs to also have their promoters

bound by common TFs. Our method (*pink*) uses clr to transform the expression matrix, uses

proportionality metric ρ to calculate coexpression and SpQN to normalize the coexpression

matrix. Method TPM + pearson (*green*) uses TPM to normalize the expression matrix followed

by Pearson correlation to calculate coexpression. Method clr + batch correction + rho + SpQN

(*blue*) uses clr to transform the expression matrix, followed by removing the top principal

component of the clr expression matrix to do batch correction, followed by calculating

coexpression using proportionality metric ρ and SpQN normalization of the coexpression matrix.

Method RPKM + pearson correlation (*purple*) uses RPKM to normalize the expression matrix

followed by Pearson correlation to calculate coexpression. Coexpression percentiles were

determined using all ORF pairs (n = 62,204,406 ORF pairs). All odds ratios are significant at p <

2.15e-5, Fisher exact test. Batch correction performed by removing the top principal component

on the clr transformed expression matrix. Error bars represent the 95% confidence interval of

the odds ratio. Dashed line shows an odds ratio of 1.

Supplementary Figure 8

Supplementary Figure 8 Proportion of nORFs defined as coexpressed (and therefore included

in the coexpression network) at various coexpression percentile cutoffs using four different

methods. Our method (*pink*) uses clr to transform the expression matrix, uses proportionality

metric ρ to calculate coexpression and SpQN to normalize the coexpression matrix. Method

TPM + Pearson (*green*) uses TPM to normalize the expression matrix followed by Pearson

correlation to calculate coexpression. Method ρ + batch correction (*blue*) uses clr to transform

the expression matrix, followed by removing the top principal component of the clr expression

matrix to do batch correction, followed by calculating coexpression using proportionality metric ρ

and SpQN normalization of the coexpression matrix. Method RPKM + pearson correlation

(*purple*) uses RPKM to normalize the expression matrix followed by Pearson correlation to

calculate coexpression. Coexpression percentiles were determined using all ORF pairs (n =

62,204,406 ORF pairs).

Supplementary Figure 9

cORF

nORF

— coexpressed

Supplementary Figure 9 Strategy for generating randomized networks. Edges between cORF-nORF and nORF-nORF pairs were swapped in a pairwise manner such that the degree of each node stayed the same. Edges between cORF-cORF pairs were not randomized.

Supplementary Figure 10

Supplementary Figure 10 GSEA pipeline using coexpression profiles to find GO terms that are

more likely to incorporate nORFs.

Supplementary Figure 11

Supplementary Figure 11 KEGG pathways that proportionally have more (*left*) (Odds ratio > 2, n

= 37 terms) or less (*right*) (Odds ratio < 0.5, n = 10 terms) GSEA enrichments with nORFs

compared to cORFs (y-axis ordered by nORF enrichment proportion from highest to lowest, BH

adjusted FDR < 0.001 for all terms, Fisher's exact test). Error bars represent the standard error

of the proportion.

Supplementary Figure 12

Supplementary Figure 12 GO terms that proportionally have more (*left*) (Odds ratio > 2, n = 35

terms) or less (*right*) (Odds ratio < 0.5, n = 11 terms) GSEA enrichments with *de novo* ORFs

compared to conserved ORFs (y-axis ordered by *de novo* ORF enrichment proportion from

highest to lowest, BH adjusted FDR < 0.001 for all terms, Fisher's exact test). Error bars

represent the standard error of the proportion.

Supplementary Figure 13

Supplementary Figure 13 A) Coexpression (y-axis) of *de novo* ORFs with neighboring

conserved ORFs per orientation (x-axis). Down same *de novo* ORFs tend to be highly

coexpressed with their neighbors; background: *de novo*-conserved ORF pairs located on

different chromosomes. B) Biological process similarity (y-axis) of *de novo* ORFs with

neighboring conserved ORFs per orientation (x-axis). Similarity measured by calculating

semantic similarity between GSEA enrichments for neighboring *de novo*-conserved ORF pairs

using relevance metric (0 = no similarity, 1 = perfect overlap); background: *de novo*-conserved

ORF pairs located on different chromosomes. C) Median expression of *de novo* ORFs (y-axis)

per orientation (x-axis). *De novo* ORFs located downstream on the same strand as conserved

ORFs have the highest expression among different orientations (considering only ORFs in only

a single orientation, dashed box in panel 4D; independent: *de novo* ORFs located further than

500 bp from all conserved ORFs). For panels A-B-C: Mann-Whitney U-test, ****: $p \leq 0.0001$, ***:

$p \leq 0.001$, **: $p \leq 0.01$, *: $p \leq 0.05$, ns: not-significant, +: small effect size (Cliff's d < 0.33), ++:

medium effect size (Cliff's d < 0.474), +++: large effect size (Cliff's d $\geq$ 0.474); all orientations

are compared to either background pairs (A, B) or to independent ORFs (C).

Supplementary Figure 14

1000

Supplementary Figure 14 Proportion of *de novo* ORFs that share a promoter with their

neighboring conserved ORF. To determine if ORFs shared a promoter with neighbors we used

a publicly available TIF-seq dataset from Pelechano et al [65]. We defined down same or up

same ORFs as sharing a promoter if they mapped to the same transcript at least once, and

defined up opposite ORFs as sharing a promoter if their respective transcripts did not have

overlapping TSSs. We found that 84% of down same (n = 174), 64% of up same (n = 368), and

66% of up opposite (n = 185) *de novo* ORFs share a promoter with their neighboring conserved

ORF. Error bars represent the standard error of the proportion.

1009 References

[1] Dujon B. The yeast genome project: what did we learn? Trends Genet TIG 1996;12:263–

70. https://doi.org/10.1016/0168-9525(96)10027-5.

[2] Fisk DG, Ball CA, Dolinski K, Engel SR, Hong EL, Issel-Tarver L, et al. Saccharomyces

cerevisiae S288C genome annotation: a working hypothesis. Yeast Chichester Engl

2006;23:857–65. https://doi.org/10.1002/yea.1400.

[3] Basrai MA, Hieter P, Boeke JD. Small Open Reading Frames: Beautiful Needles in the

Haystack. Genome Res 1997;7:768–71. https://doi.org/10.1101/gr.7.8.768.

[4] Nagalakshmi U, Wang Z, Waern K, Shou C, Raha D, Gerstein M, et al. The

Transcriptional Landscape of the Yeast Genome Defined by RNA Sequencing. Science

2008;320:1344–9. https://doi.org/10.1126/science.1158441.

[5] Ingolia NT, Ghaemmaghami S, Newman JRS, Weissman JS. Genome-Wide Analysis in

Vivo of Translation with Nucleotide Resolution Using Ribosome Profiling. Science

2009;324:218–23. https://doi.org/10.1126/science.1168978.

[6] Ingolia NT, Brar GA, Stern-Ginossar N, Harris MS, Talhouarne GJS, Jackson SE, et al.

Ribosome Profiling Reveals Pervasive Translation Outside of Annotated Protein-Coding

Genes. Cell Rep 2014;8:1365–79. https://doi.org/10.1016/j.celrep.2014.07.045.

[7] Bazzini AA, Johnstone TG, Christiano R, Mackowiak SD, Obermayer B, Fleming ES, et

al. Identification of small ORFs in vertebrates using ribosome footprinting and

1028   evolutionary conservation. EMBO J 2014;33:981–93.

1029   https://doi.org/10.1002/embj.201488411.

1030 [8] Couso J-P, Patraquim P. Classification and function of small open reading frames. Nat

1031   Rev Mol Cell Biol 2017;18:575–89. https://doi.org/10.1038/nrm.2017.58.

1032 [9] Lu S, Zhang J, Lian X, Sun L, Meng K, Chen Y, et al. A hidden human proteome encoded

1033   by 'non-coding' genes. Nucleic Acids Res 2019;47:8111–25.

1034   https://doi.org/10.1093/nar/gkz646.

1035 [10] Chen J, Brunner A-D, Cogan JZ, Nuñez JK, Fields AP, Adamson B, et al. Pervasive

1036   functional translation of noncanonical human open reading frames. Science

1037   2020;367:1140–6. https://doi.org/10.1126/science.aay0262.

1038 [11] Orr MW, Mao Y, Storz G, Qian S-B. Alternative ORFs and small ORFs: shedding light on

1039   the dark proteome. Nucleic Acids Res 2020;48:1029–42.

1040   https://doi.org/10.1093/nar/gkz734.

1041 [12] Vitorino R, Guedes S, Amado F, Santos M, Akimitsu N. The role of micropeptides in

1042   biology. Cell Mol Life Sci 2021;78:3285–98. https://doi.org/10.1007/s00018-020-03740-3.

1043 [13] Prensner JR, Enache OM, Luria V, Krug K, Clauser KR, Dempster JM, et al.

1044   Noncanonical open reading frames encode functional proteins essential for cancer cell

1045   survival. Nat Biotechnol 2021;39:697–704. https://doi.org/10.1038/s41587-020-00806-2.

1046 [14] Wacholder A, Parikh SB, Coelho NC, Acar O, Houghton C, Chou L, et al. A vast

1047   evolutionarily transient translatome contributes to phenotype and fitness. Cell Syst

1048   2023;14:363-381.e8. https://doi.org/10.1016/j.cels.2023.04.002.

1049 [15] Vakirlis N, Acar O, Hsu B, Castilho Coelho N, Van Oss SB, Wacholder A, et al. De novo

1050   emergence of adaptive membrane proteins from thymine-rich genomic sequences. Nat

1051   Commun 2020;11:781. https://doi.org/10.1038/s41467-020-14500-z.

1052   [16]   Arnoult N, Correia A, Ma J, Merlo A, Garcia-Gomez S, Maric M, et al. Regulation of DNA

1053   repair pathway choice in S and G2 phases by the NHEJ inhibitor CYREN. Nature

1054   2017;549:548–52. https://doi.org/10.1038/nature24023.

1055   [17]   Anderson DM, Anderson KM, Chang C-L, Makarewich CA, Nelson BR, McAnally JR, et

1056   al. A Micropeptide Encoded by a Putative Long Noncoding RNA Regulates Muscle

1057   Performance. Cell 2015;160:595–606. https://doi.org/10.1016/j.cell.2015.01.009.

1058   [18]   Magny EG, Pueyo JI, Pearl FMG, Cespedes MA, Niven JE, Bishop SA, et al. Conserved

1059   Regulation of Cardiac Calcium Uptake by Peptides Encoded in Small Open Reading

1060   Frames. Science 2013;341:1116–20. https://doi.org/10.1126/science.1238802.

1061   [19]   Matsumoto A, Pasut A, Matsumoto M, Yamashita R, Fung J, Monteleone E, et al.

1062   mTORC1 and muscle regeneration are regulated by the LINC00961-encoded SPAR

1063   polypeptide. Nature 2017;541:228–32. https://doi.org/10.1038/nature21034.

1064   [20]   Jackson R, Kroehling L, Khitun A, Bailis W, Jarret A, York AG, et al. The translation of

1065   non-canonical open reading frames controls mucosal immunity. Nature 2018;564:434–8.

1066   https://doi.org/10.1038/s41586-018-0794-7.

1067   [21]   Bhatta A, Atianand M, Jiang Z, Crabtree J, Blin J, Fitzgerald KA. A Mitochondrial

1068   Micropeptide Is Required for Activation of the Nlrp3 Inflammasome. J Immunol

1069   2020;204:428–37. https://doi.org/10.4049/jimmunol.1900791.

1070   [22]   Niu X, Zhang J, Zhang L, Hou Y, Pu S, Chu A, et al. Weighted Gene Co-Expression

1071   Network Analysis Identifies Critical Genes in the Development of Heart Failure After

1072   Acute Myocardial Infarction. Front Genet 2019;10.

1073   https://doi.org/10.3389/fgene.2019.01214.

1074   [23]   Wright BW, Yi Z, Weissman JS, Chen J. The dark proteome: translation from

1075   noncanonical open reading frames. Trends Cell Biol 2021.

1076   https://doi.org/10.1016/j.tcb.2021.10.010.

[24]   Carvunis A-R, Rolland T, Wapinski I, Calderwood MA, Yildirim MA, Simonis N, et al.

Proto-genes and *de novo* gene birth. Nature 2012;487:370–4.

https://doi.org/10.1038/nature11184.

[25]   Van Oss SB, Carvunis A-R. De novo gene birth. PLOS Genet 2019;15:e1008160.

https://doi.org/10.1371/journal.pgen.1008160.

[26]   Sandmann C-L, Schulz JF, Ruiz-Orera J, Kirchner M, Ziehm M, Adami E, et al.

Evolutionary origins and interactomes of human, young microproteins and small peptides

translated from short open reading frames. Mol Cell 2023;83:994-1011.e18.

https://doi.org/10.1016/j.molcel.2023.01.023.

[27]   Zhang W, Landback P, Gschwend AR, Shen B, Long M. New genes drive the evolution of

gene interaction networks in the human and mouse genomes. Genome Biol 2015;16:202.

https://doi.org/10.1186/s13059-015-0772-4.

[28]   Abrusán G. Integration of New Genes into Cellular Networks, and Their Structural

Maturation. Genetics 2013;195:1407–17. https://doi.org/10.1534/genetics.113.152256.

[29]   Capra JA, Pollard KS, Singh M. Novel genes exhibit distinct patterns of function

acquisition and network integration. Genome Biol 2010;11:R127.

https://doi.org/10.1186/gb-2010-11-12-r127.

[30]   Housman G, Ulitsky I. Methods for distinguishing between protein-coding and long

noncoding RNAs and the elusive biological purpose of translation of long noncoding

RNAs. Biochim Biophys Acta BBA - Gene Regul Mech 2016;1859:31–40.

https://doi.org/10.1016/j.bbagrm.2015.07.017.

[31]   Pertea M, Shumate A, Pertea G, Varabyou A, Breitwieser FP, Chang Y-C, et al. CHESS:

a new human gene catalog curated from thousands of large-scale RNA sequencing

experiments reveals extensive transcriptional noise. Genome Biol 2018;19:208.

https://doi.org/10.1186/s13059-018-1590-2.

[32] Xu H, Li C, Xu C, Zhang J. Chance promoter activities illuminate the origins of eukaryotic

intergenic transcriptions. Nat Commun 2023;14:1826. https://doi.org/10.1038/s41467-

023-37610-w.

[33] Schlötterer C. Genes from scratch – the evolutionary fate of de novo genes. Trends

Genet 2015;31:215–9. https://doi.org/10.1016/j.tig.2015.02.007.

[34] Zhao L, Saelao P, Jones CD, Begun DJ. Origin and spread of de novo genes in

Drosophila melanogaster populations. Science 2014;343:769–72.

https://doi.org/10.1126/science.1248286.

[35] Zhuang X, Yang C, Murphy KR, Cheng C-HC. Molecular mechanism and history of non-

sense to sense evolution of antifreeze glycoprotein gene in northern gadids. Proc Natl

Acad Sci 2019;116:4400–5. https://doi.org/10.1073/pnas.1817138116.

[36] Ruiz-Orera J, Hernandez-Rodriguez J, Chiva C, Sabidó E, Kondova I, Bontrop R, et al.

Origins of De Novo Genes in Human and Chimpanzee. PLOS Genet 2015;11:e1005721.

https://doi.org/10.1371/journal.pgen.1005721.

[37] Vakirlis N, Vance Z, Duggan KM, McLysaght A. De novo birth of functional microproteins

in the human lineage. Cell Rep 2022;41:111808.

https://doi.org/10.1016/j.celrep.2022.111808.

[38] Majic P, Payne JL. Enhancers Facilitate the Birth of De Novo Genes and Gene

Integration into Regulatory Networks. Mol Biol Evol 2020;37:1165–78.

https://doi.org/10.1093/molbev/msz300.

[39] Ruiz-Orera J, Villanueva-Cañas JL, Albà MM. Evolution of new proteins from translated

sORFs in long non-coding RNAs. Exp Cell Res 2020;391:111940.

https://doi.org/10.1016/j.yexcr.2020.111940.

[40] Chen J-Y, Shen QS, Zhou W-Z, Peng J, He BZ, Li Y, et al. Emergence, Retention and

Selection: A Trilogy of Origination for Functional De Novo Proteins from Ancestral

LncRNAs in Primates. PLOS Genet 2015;11:e1005391.

https://doi.org/10.1371/journal.pgen.1005391.

[41] Vakirlis N, Hebert AS, Opulente DA, Achaz G, Hittinger CT, Fischer G, et al. A Molecular

Portrait of De Novo Genes in Yeasts. Mol Biol Evol 2018;35:631–45.

https://doi.org/10.1093/molbev/msx315.

[42] Neme R, Tautz D. Fast turnover of genome transcription across evolutionary time

exposes entire non-coding DNA to de novo gene emergence. ELife 2016;5:e09977.

https://doi.org/10.7554/eLife.09977.

[43] Knowles DG, McLysaght A. Recent de novo origin of human protein-coding genes.

Genome Res 2009;19:1752–9. https://doi.org/10.1101/gr.095026.109.

[44] Ebisuya M, Yamamoto T, Nakajima M, Nishida E. Ripples from neighbouring

transcription. Nat Cell Biol 2008;10:1106–13. https://doi.org/10.1038/ncb1771.

[45] Ghanbarian AT, Hurst LD. Neighboring Genes Show Correlated Evolution in Gene

Expression. Mol Biol Evol 2015;32:1748–66. https://doi.org/10.1093/molbev/msv053.

[46] Ji Z, Song R, Regev A, Struhl K. Many lncRNAs, 5'UTRs, and pseudogenes are

translated and some are likely to express functional proteins. ELife 2015;4:e08890.

https://doi.org/10.7554/eLife.08890.

[47] Li J, Singh U, Arendsee Z, Wurtele ES. Landscape of the Dark Transcriptome Revealed

Through Re-mining Massive RNA-Seq Data. Front Genet 2021;12.

[48] O'Meara TR, O'Meara MJ. DeORFanizing Candida albicans Genes using Coexpression.

MSphere 2021;6:e01245-20. https://doi.org/10.1128/mSphere.01245-20.

[49] Chothani SP, Adami E, Widjaja AA, Langley SR, Viswanathan S, Pua CJ, et al. A high-

resolution map of human RNA translation. Mol Cell 2022;82:2885-2899.e8.

https://doi.org/10.1016/j.molcel.2022.06.023.

[50] Kim SK, Lund J, Kiraly M, Duke K, Jiang M, Stuart JM, et al. A Gene Expression Map for

Caenorhabditis elegans. Science 2001;293:2087–92.

https://doi.org/10.1126/science.1061603.

[51] Stuart JM, Segal E, Koller D, Kim SK. A Gene-Coexpression Network for Global

Discovery of Conserved Genetic Modules. Science 2003;302:249–55.

https://doi.org/10.1126/science.1087447.

[52] Yang Y, Han L, Yuan Y, Li J, Hei N, Liang H. Gene co-expression network analysis

reveals common system-level properties of prognostic genes across cancer types. Nat

Commun 2014;5:3231. https://doi.org/10.1038/ncomms4231.

[53] Voineagu I, Wang X, Johnston P, Lowe JK, Tian Y, Horvath S, et al. Transcriptomic

analysis of autistic brain reveals convergent molecular pathology. Nature 2011;474:380–

4. https://doi.org/10.1038/nature10110.

[54] Xue Z, Huang K, Cai C, Cai L, Jiang C, Feng Y, et al. Genetic programs in human and

mouse early embryos revealed by single-cell RNA sequencing. Nature 2013;500:593–7.

https://doi.org/10.1038/nature12364.

[55] Lee J, Shah M, Ballouz S, Crow M, Gillis J. CoCoCoNet: conserved and comparative co-

expression across a diverse set of species. Nucleic Acids Res 2020;48:W566–71.

https://doi.org/10.1093/nar/gkaa348.

[56] van Dam S, Võsa U, van der Graaf A, Franke L, de Magalhães JP. Gene co-expression

analysis for functional classification and gene–disease predictions. Brief Bioinform

2018;19:575–92. https://doi.org/10.1093/bib/bbw139.

[57] Yin W, Mendoza L, Monzon-Sandoval J, Urrutia AO, Gutierrez H. Emergence of co-

expression in gene regulatory networks. PLOS ONE 2021;16:e0247671.

https://doi.org/10.1371/journal.pone.0247671.

[58] Hanada K, Higuchi-Takeuchi M, Okamoto M, Yoshizumi T, Shimizu M, Nakaminami K, et

al. Small open reading frames associated with morphogenesis are hidden in plant

genomes. Proc Natl Acad Sci 2013;110:2395–400.
https://doi.org/10.1073/pnas.1213958110.

[59] Bashir K, Hanada K, Shimizu M, Seki M, Nakanishi H, Nishizawa NK. Transcriptomic analysis of rice in response to iron deficiency and excess. Rice 2014;7:18. https://doi.org/10.1186/s12284-014-0018-1.

[60] Stiens J, Tan YY, Joyce R, Arnvig KB, Kendall SL, Nobeli I. Using a Whole Genome Co-expression Network to Inform the Functional Characterisation of Predicted Genomic Elements from Mycobacterium tuberculosis Transcriptomic Data 2022:2022.06.22.497203. https://doi.org/10.1101/2022.06.22.497203.

[61] Li H, Xiao L, Zhang L, Wu J, Wei B, Sun N, et al. FSPP: A Tool for Genome-Wide Prediction of smORF-Encoded Peptides and Their Functions. Front Genet 2018;9. https://doi.org/10.3389/fgene.2018.00096.

[62] Wang Y, Hicks SC, Hansen KD. Addressing the mean-correlation relationship in co-expression analysis. PLOS Comput Biol 2022;18:e1009954. https://doi.org/10.1371/journal.pcbi.1009954.

[63] Crow M, Paul A, Ballouz S, Huang ZJ, Gillis J. Exploiting single-cell expression to characterize co-expression replicability. Genome Biol 2016;17:101. https://doi.org/10.1186/s13059-016-0964-6.

[64] Pu S, Wong J, Turner B, Cho E, Wodak SJ. Up-to-date catalogues of yeast protein complexes. Nucleic Acids Res 2009;37:825–31. https://doi.org/10.1093/nar/gkn1005.

[65] Rossi MJ, Kuntala PK, Lai WKM, Yamada N, Badjatia N, Mittal C, et al. A high-resolution protein architecture of the budding yeast genome. Nature 2021;592:309–14. https://doi.org/10.1038/s41586-021-03314-8.

[66] Pelechano V, Wei W, Steinmetz LM. Extensive transcriptional heterogeneity revealed by isoform profiling. Nature 2013;497:127–31. https://doi.org/10.1038/nature12121.

1202   [67]   Cherry JM, Hong EL, Amundsen C, Balakrishnan R, Binkley G, Chan ET, et al.

1203   Saccharomyces Genome Database: the genomics resource of budding yeast. Nucleic

1204   Acids Res 2012;40:D700–5. https://doi.org/10.1093/nar/gkr1029.

1205   [68]   Skinnider MA, Squair JW, Foster LJ. Evaluating measures of association for single-cell

1206   transcriptomics. Nat Methods 2019;16:381–6. https://doi.org/10.1038/s41592-019-0372-4.

1207   [69]   Quinn TP, Richardson MF, Lovell D, Crowley TM. propr: An R-package for Identifying

1208   Proportionally Abundant Features Using Compositional Data Analysis. Sci Rep

1209   2017;7:16252. https://doi.org/10.1038/s41598-017-16520-0.

1210   [70]   Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network

1211   analysis. BMC Bioinformatics 2008;9:559. https://doi.org/10.1186/1471-2105-9-559.

1212   [71]   Ballouz S, Verleyen W, Gillis J. Guidance for RNA-seq co-expression network

1213   construction and analysis: safety in numbers. Bioinformatics 2015;31:2123–30.

1214   https://doi.org/10.1093/bioinformatics/btv118.

1215   [72]   Parsana P, Ruberman C, Jaffe AE, Schatz MC, Battle A, Leek JT. Addressing

1216   confounding artifacts in reconstruction of gene co-expression networks. Genome Biol

1217   2019;20:94. https://doi.org/10.1186/s13059-019-1700-9.

1218   [73]   Chang W, Cheng J, Allaire J, Sievert C, Schloerke B, Xie Y, et al. shiny: Web application

1219   framework for R. 2023.

1220   [74]   Fruchterman TMJ, Reingold EM. Graph drawing by force-directed placement. Softw Pract

1221   Exp 1991;21:1129–64. https://doi.org/10.1002/spe.4380211102.

1222   [75]   Krogh A, Larsson B, von Heijne G, Sonnhammer ELL. Predicting transmembrane protein

1223   topology with a hidden markov model: application to complete genomes. J Mol Biol

1224   2001;305:567–80. https://doi.org/10.1006/jmbi.2000.4315.

1225   [76]   Ciccarelli M, Masser AE, Kaimal JM, Planells J, Andréasson C. Genetic inactivation of

1226   essential HSF1 reveals an isolated transcriptional stress response selectively induced by

1227   protein misfolding 2023:2023.05.05.539545. https://doi.org/10.1101/2023.05.05.539545.

[77]   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene

set enrichment analysis: A knowledge-based approach for interpreting genome-wide

expression profiles. Proc Natl Acad Sci 2005;102:15545–50.

https://doi.org/10.1073/pnas.0506580102.

[78]   Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. KEGG as a reference

resource for gene and protein annotation. Nucleic Acids Res 2016;44:D457–62.

https://doi.org/10.1093/nar/gkv1070.

[79]   Hu Z, Killion PJ, Iyer VR. Genetic reconstruction of a functional transcriptional regulatory

network. Nat Genet 2007;39:683–7. https://doi.org/10.1038/ng2012.

[80]   Marion RM, Regev A, Segal E, Barash Y, Koller D, Friedman N, et al. Sfp1 is a stress-

and nutrient-sensitive regulator of ribosomal protein gene expression. Proc Natl Acad Sci

2004;101:14315–22. https://doi.org/10.1073/pnas.0405353101.

[81]   Masser AE, Kang W, Roy J, Mohanakrishnan Kaimal J, Quintana-Cordero J, Friedländer

MR, et al. Cytoplasmic protein misfolding titrates Hsp70 to activate nuclear Hsf1. ELife

2019;8:e47791. https://doi.org/10.7554/eLife.47791.

[82]   Schlicker A, Domingues FS, Rahnenführer J, Lengauer T. A new measure for functional

similarity of gene products based on Gene Ontology. BMC Bioinformatics 2006;7:302.

https://doi.org/10.1186/1471-2105-7-302.

[83]   Wei W, Pelechano V, Järvelin AI, Steinmetz LM. Functional consequences of bidirectional

promoters. Trends Genet 2011;27:267–76. https://doi.org/10.1016/j.tig.2011.04.002.

[84]   Zrimec J, Börlin CS, Buric F, Muhammad AS, Chen R, Siewers V, et al. Deep learning

suggests that gene expression is encoded in all parts of a co-evolving interacting gene

regulatory structure. Nat Commun 2020;11:6141. https://doi.org/10.1038/s41467-020-

19921-4.

[85] Blevins WR, Ruiz-Orera J, Messeguer X, Blasco-Moreno B, Villanueva-Cañas JL, Espinar L, et al. Uncovering de novo gene birth in yeast using deep transcriptomics. Nat Commun 2021;12:604. https://doi.org/10.1038/s41467-021-20911-3.

[86] Khitun A, Ness TJ, Slavoff SA. Small open reading frames and cellular stress responses. Mol Omics 2019;15:108–16. https://doi.org/10.1039/C8MO00283E.

[87] Wilson BA, Masel J. Putatively Noncoding Transcripts Show Extensive Association with Ribosomes. Genome Biol Evol 2011;3:1245–52. https://doi.org/10.1093/gbe/evr099.

[88] Li D, Yan Z, Lu L, Jiang H, Wang W. Pleiotropy of the de novo-originated gene MDF1. Sci Rep 2014;4. https://doi.org/10.1038/srep07280.

[89] Frumkin I, Laub MT. Selection of a de novo gene that can promote survival of E. coli by modulating protein homeostasis pathways 2023:2023.02.07.527531. https://doi.org/10.1101/2023.02.07.527531.

[90] Li D, Dong Y, Jiang Y, Jiang H, Cai J, Wang W. A de novo originated gene depresses budding yeast mating pathway and is repressed by the protein encoded by its antisense strand. Cell Res 2010;20:408–20. https://doi.org/10.1038/cr.2010.31.

[91] Pagé N, Gérard-Vincent M, Ménard P, Beaulieu M, Azuma M, Dijkgraaf GJP, et al. A Saccharomyces cerevisiae Genome-Wide Mutant Screen for Altered Sensitivity to K1 Killer Toxin. Genetics 2003;163:875–94. https://doi.org/10.1093/genetics/163.3.875.

[92] Tassios E, Nikolaou C, Vakirlis N. Intergenic Regions of Saccharomycotina Yeasts are Enriched in Potential to Encode Transmembrane Domains. Mol Biol Evol 2023;40:msad059. https://doi.org/10.1093/molbev/msad059.

[93] Peng J, Zhao L. The origin and structural evolution of de novo genes in Drosophila 2023:2023.03.13.532420. https://doi.org/10.1101/2023.03.13.532420.

[94] Kesner JS, Chen Z, Aparicio AA, Wu X. A unified model for the surveillance of translation in diverse noncoding sequences 2022:2022.07.20.500724. https://doi.org/10.1101/2022.07.20.500724.

[95]   Slavoff SA, Mitchell AJ, Schwaid AG, Cabili MN, Ma J, Levin JZ, et al. Peptidomic discovery of short open reading frame–encoded peptides in human cells. Nat Chem Biol 2013;9:59–64. https://doi.org/10.1038/nchembio.1120.

[96]   Zhang S, Reljić B, Liang C, Kerouanton B, Francisco JC, Peh JH, et al. Mitochondrial peptide BRAWNIN is essential for vertebrate respiratory complex III assembly. Nat Commun 2020;11:1312. https://doi.org/10.1038/s41467-020-14999-2.

[97]   Leong AZ-X, Lee PY, Mohtar MA, Syafruddin SE, Pung Y-F, Low TY. Short open reading frames (sORFs) and microproteins: an update on their identification and validation measures. J Biomed Sci 2022;29:19. https://doi.org/10.1186/s12929-022-00802-5.

[98]   Mayr C. What Are 3′ UTRs Doing? Cold Spring Harb Perspect Biol 2019;11:a034728. https://doi.org/10.1101/cshperspect.a034728.

[99]   Vilborg A, Passarelli MC, Yario TA, Tycowski KT, Steitz JA. Widespread Inducible Transcription Downstream of Human Genes. Mol Cell 2015;59:449–61. https://doi.org/10.1016/j.molcel.2015.06.016.

[100]   Wu Q, Wright M, Gogol MM, Bradford WD, Zhang N, Bazzini AA. Translation of small downstream ORFs enhances translation of canonical main open reading frames. EMBO J 2020;39:e104763. https://doi.org/10.15252/embj.2020104763.

[101]   Wu B, Cox MP. Characterization of Bicistronic Transcription in Budding Yeast. MSystems 2021;6:e01002-20. https://doi.org/10.1128/mSystems.01002-20.

[102]   Kustatscher G, Grabowski P, Rappsilber J. Pervasive coexpression of spatially proximal genes is buffered at the protein level. Mol Syst Biol 2017;13:937. https://doi.org/10.15252/msb.20177548.

[103]   Saccharomyces Genome Database | SGD n.d. https://www.yeastgenome.org/ (accessed January 20, 2021).

[104]   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics 2010;26:841–2. https://doi.org/10.1093/bioinformatics/btq033.

[105] Krueger F, James F, Ewels P, Afyounian E, Weinstein M, Schuster-Boeckler B, et al.

FelixKrueger/TrimGalore 2023. https://doi.org/10.5281/zenodo.7598955.

[106] Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon: fast and bias-aware

quantification of transcript expression using dual-phase inference. Nat Methods

2017;14:417–9. https://doi.org/10.1038/nmeth.4197.

[107] Lin P, Troup M, Ho JWK. CIDR: Ultrafast and accurate clustering through imputation for

single-cell RNA-seq data. Genome Biol 2017;18:59. https://doi.org/10.1186/s13059-017-

1188-0.

[108] L. Lun AT, Bach K, Marioni JC. Pooling across cells to normalize single-cell RNA

sequencing data with many zero counts. Genome Biol 2016;17:75.

https://doi.org/10.1186/s13059-016-0947-7.

[109] Lovell DR, Chua X-Y, McGrath A. Counts: an outstanding challenge for log-ratio analysis

of compositional data in the molecular biosciences. NAR Genomics Bioinforma

2020;2:lqaa040. https://doi.org/10.1093/nargab/lqaa040.

[110] Gene Ontology Resource. Gene Ontol Resour n.d. http://geneontology.org/ (accessed

March 10, 2022).

[111] Klopfenstein DV, Zhang L, Pedersen BS, Ramírez F, Warwick Vesztrocy A, Naldi A, et al.

GOATOOLS: A Python library for Gene Ontology analyses. Sci Rep 2018;8:1–17.

https://doi.org/10.1038/s41598-018-28948-z.

[112] Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. J R Stat Soc Ser B Methodol 1995;57:289–300.

[113] Csardi G, Nepusz T. The Igraph Software Package for Complex Network Research.

InterJournal 2005;Complex Systems:1695.

[114] Hagberg AA, Schult DA, Swart PJ. Exploring network structure, dynamics, and function

using NetworkX. In: Varoquaux G, Vaught T, Millman J, editors. Proc. 7th Python Sci.

Conf., Pasadena, CA USA: 2008, p. 11–5.

[115] Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov MN, Sergushichev A. Fast gene

set enrichment analysis 2021:060012. https://doi.org/10.1101/060012.

[116] Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal

enrichment tool for interpreting omics data. The Innovation 2021;2:100141.

https://doi.org/10.1016/j.xinn.2021.100141.

[117] Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol 2014;15:550. https://doi.org/10.1186/s13059-

014-0550-8.

[118] Shen X-X, Opulente DA, Kominek J, Zhou X, Steenwyk JL, Buh KV, et al. Tempo and

Mode of Genome Evolution in the Budding Yeast Subphylum. Cell 2018;175:1533-

1545.e20. https://doi.org/10.1016/j.cell.2018.10.023.

[119] Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:

architecture and applications. BMC Bioinformatics 2009;10:421.

https://doi.org/10.1186/1471-2105-10-421.

[120] Yu G, Li F, Qin Y, Bo X, Wu Y, Wang S. GOSemSim: an R package for measuring

semantic similarity among GO terms and gene products. Bioinformatics 2010;26:976–8.

https://doi.org/10.1093/bioinformatics/btq064.

[121] R Core Team. R: A Language and Environment for Statistical Computing. Vienna,

Austria: R Foundation for Statistical Computing; 2017.

1349