Genomic insights into greater yam tuber quality traits

Ana Zotta Mota[1,2,3†], Komivi Dossa[1,4†], Mathieu Lechaudel[5,6], Denis Cornet[1,2], Pierre Mournet[1,2], David Lopez[1,2], Hana Chaïr[1,2]

[1]CIRAD, UMR AGAP, F34398-Montpellier, France,
[2] AGAP, Univ Montpellier, CIRAD, INRA, Montpellier SupAgro,Montpellier, France,
[3] Institut Sophia Agrobiotech, Sophia Antipolis, France
[4]CIRAD, UMR AGAP Institut, 97170 Petit Bourg, Guadeloupe, France.
[5]UMR Qualisud, CIRAD, F97130-Capesterre-Belle-Eau, Guadeloupe, France
[6]QualiSud, Université Montpellier, Institut Agro, CIRAD, Avignon Université, Université de la Réunion, 34398 Montpellier, France

†These authors contributed equally to this work
Corresponding author: hana.chair@cirad.fr

ABSTRACT

Yams are important tubers widely consumed in developing countries in various forms, mainly boiled, pounded and fried. Tuber quality is a key criterion for acceptance by the various actors in the value chain. However, the genetics of yam tuber quality has not yet been comprehensively investigated. Given this lack of knowledge, we used population genomics and candidate gene association approaches to unravel the genetic basis of the three main quality attributes of boiled yam, namely texture, starch content and colour indices. We re-sequenced the whole genome of 127 yam genotypes with different quality attributes, and performed an enrichment of the already available functional genome annotation using comparative genomics. Population genomics analysis highlighted three main genetic groups and 1,053 genes under selection. We focused this work on three known quality trait-related pathways: pectin, starch content and flavonoid biosynthesis, and inventoried in the genome all the structural genes related to those pathways by comparative genomics. We improved the functional annotation of the three pathways with 48% more genes. A total of 18 candidate genes containing genetic variants significantly associated with the target traits were detected, including eight genes that were also found under selection. The texture-related candidate genes were distributed between the pectin and starch pathways. Overall,

the use of comparative genomics has allowed the generation of an unprecedented genomic
resource. The improved functional annotation of the yam genome is a promising predictive tool
for identifying other core genes associated with any trait of interest to meet the growing need and
diversified demands for yams.

Keywords: comparative genomics, texture, pectin, starch, flavonoids, *Dioscorea alata*

INTRODUCTION

Releasing varieties with enhanced quality is a target in the breeding process, and failure to achieve
it can lead to the rejection of improved varieties. However, food quality is a complex attribute that
reflects the preferences of all actors in the value chain: producers, processors, retailers and
consumers. In recent years, there has been an increasing amount of studies on the identification of
genomic regions associated with quality traits (Liao et al., 2021; Xia et al., 2020), with a focus on
economically important crops such as rice, tomato, citrus, etc. (Xiao et al., 2021; Dominguez et
al., 2020; Butelli et al., 2019). Underutilised crops such as tropical root and tuber crops have been
largely neglected. However, root and tuber crops are among the most important crops for
subsistence and commercial purposes after cereals.

Yams, of the Dioscoreaceae family, are important edible tuber crops, mainly in developing
countries. The most widespread species is greater yam (*Dioscorea alata* L.), cultivated mainly for
its starchy tubers in tropical regions (Wu et al., 2014; Siqueira et al., 2014; Girma et al., 2014
Lebot et al., 1998). Greater yam displays a wide tolerance to different environments (Orkwor and
Asadu, 1998). It is a dioecious and autopolyploid species ($2n = 2x = 40$, $3x = 60$ and $4x = 80$)
(Arnau et al., 2009). The recent publication of the genetic map of this species (Cormier et al., 2021)
followed by the release of its reference genome (Bredeson et al., 2022) have paved the way to
work on the genetic architectures of the traits of interest. Genomic regions associated to
anthracnose resistance, tuber oxidative browning, tuber morphological attributes, and dry matter
have been revealed (Bredeson et al., 2022; Gatarira et al., 2020; Ehounou et al., 2022). Although
it is well known that tuber quality is important for varieties adoption in root and tuber crops

(Dufour et al. 2021), the genetics of yam tuber quality has not yet been comprehensively
investigated.

Greater yam is consumed in various forms, mainly boiled, pounded, fried or baked (Honfozo et
al., 2021). The quality of boiled yam, as characterised by sensory testing with consumers, is related
to white colour, crumbly, sticky to the fingers, non-fibrous and easy to chew texture, sweet taste
and good smell (Honfozo et al., 2021). The choice of colour and the consistency of the tubers after
cooking are essential characteristics that have been previously selected in other crops, such as
cassava and potato (Alves-Pereira et al., 2022; Hardigan et al., 2017). Texture, which determines
the ability of the raw material to soften after cooking while maintaining its firmness, is a complex
trait. The main factors influencing texture are starch content and its distribution in the tuber, cell
wall structure and composition, and degradation of the middle lamella of the cell wall (reviewed
in Taylor et al., 2008; Ross et al., 2011). The cell wall is composed of 90-95% polysaccharides,
divided into cellulose, hemicellulose and pectin, and only 5-10% protein. Cell wall
polysaccharides and starch content are mainly associated with two metabolic pathways: pentose
and glucuronate interconversions, and starch and sucrose metabolism. Colour, one of the most
important varietal rejection factors, is partly determined by genes in the flavonoid biosynthetic
pathway (He et al., 2021). Despite their importance for the main quality attributes of boiled yam,
these pathways have been very little studied.

Over the last decade, hundreds of genomes have been released. Likewise, much efforts have been
put on improving the genome annotation in order to extract biological knowledge from genomic
sequences (Marks et al., 2021). However, annotating a genome is time and resources consuming.
Automatic pipelines can produce inaccurate genome annotation and their results often require
manual curation (Bolger et al., 2017). In non-model species, this can be especially true, since
several genes lack a precise functional annotation, relying on public databases and gene similarity
only. The use of comparative analysis allows to find homologous and orthologous genes from
different species and within the species of interest (Mota et al., 2020), and predict more precisely
their functions. Using the annotation produced with non-manual curated methods (similarity-
based), together with comparative genomics methods can increase the number of genes with a
functional annotation, and also help to better assign the functional annotation of other genes.

In this study, we re-sequenced the whole genome of 127 yam genotypes with different quality traits from distinct geographical locations. We focused our work on the three metabolic pathways involved in texture and in starch content (pentose and glucuronate interconversions, starch and sucrose metabolism), and in colour (flavonoid biosynthesis). Through the use of comparative genomics, we were able to improve the annotation of a number of genes, including those in the three major metabolic pathways targeted, thereby enriching the annotation of the *D. alata* genome. We conducted a population genomics study to identify genes under selection. To complement our results, we performed a candidate gene association analysis using phenotypic data on texture, colour and starch content. We identified candidate genes associated with key quality traits for boiled yam and assessed their expression profiles using transcriptomic analysis. Overall, we generated a consistent and valuable amount of genomic resources for future research on important agronomic traits in yam.

RESULTS

Pattern of genome-wide variation, population structure and signatures of selection

We generated the whole-genome sequences of 127 genotypes of *Dioscorea* spp. from diverse origins (Africa, Caribbean, Pacific and Asia) (Figure 1A) using short-read sequencing with a mean coverage of 37X (Supplemental Figure 1, Supplemental Table 1). The polymorphism analyses in *D. alata* genotypes yielded a total of 63 M SNPs after filtering. The type of SNPs was divided as 2.8% of variants present in exons, 1.3% of synonymous variants, and 48% intergenic variants (Figure 1B, Supplemental Figure 2). For diversity purposes, only diploid genotypes (107) were used (Supplemental Table 1), since the diversity of the diploid genotypes represents the overall diversity of the species as demonstrated previously (Sharif et al., 2020). Three major groups were separated phylogenetic tree (Supplemental Figure 3A) and by principal component analysis (PCA). A first group consisting mainly of Asian genotypes, a second group composed of Pacific genotypes, and a final group of African and Caribbean genotypes (Figure 1C), which agrees with our previous study, performed by the genotype-by-sequencing analysis on a larger *D. alata* population (Sharif et al., 2020). The genetic structure separated the genotypes into four clusters

(Figure 1D). However, as the fourth group had only six genotypes, a robust statistical analysis
could not be conducted. Consequently, we conducted our further analysis with $K=3$. The cluster
A composed mainly of Pacific representatives and, to a lesser extent, African and Caribbean
genotypes, while the cluster B was composed mostly by representatives from Africa and
Caribbean, and finally the cluster C had mostly representatives from Asia (26 out of 30) (Figure
1D, Supplemental Figure 3B), which is in accordance with the PCA results. The nucleotide
diversity (π) was low in all gene pools $0.407e^{-4}$. The π values were significantly different between
the clusters, with the highest values obtained for the cluster A and C ($\pi = 0.399e^{-4}$ and $0.396e^{-4}$,
respectively) and the lowest for the Cluster B ($\pi = 0.33e^{-4}$).
To evaluate the genetic variance among the three clusters, we calculated the fixation index values
(*Fst*) for each region in a window of 50kb and a step-window size of 10kb. We found in total 2,141
outlier regions *i.e.,* with significant values of *Fst* (varying from $7.85e^{-06}$ to 0.64) (Figure 2A, B and
C). The comparison of Cluster B *vs* C, showed the highest values of *Fst* and the highest number
of outlier regions (881), followed by the comparison of A *vs* C (789) then A *vs* B (471) (Figure
2D). These outlier regions harboured a total of 1,053 unique genes (Figure 2D). We found 11
outlier regions in common among the three genetic clusters, which could be the result of an early
purifying selection in *D. alata*. Only four genes were present in these 11 common outlier regions:
two closely located genes on chromosome 1 and two closely located genes on chromosome 4. The
genes found on chromosome 1 were both annotated as non-specific serine/threonine kinase
(EC:2.7.11.1). Similarly, both genes on chromosome 4 are annotated as proteasome endopeptidase
complexes (EC:3.4.25.1). Functional analysis of the 1,053 genes based on Gene Ontology (GO)
terms showed that various functional pathways were enriched. The most enriched GO terms were
tripeptide transmembrane transporter, transmembrane receptor protein binding, stamen filament
development, hormone-mediated signalling pathway and lignin biosynthetic process (Figure 2E).
148
149 **Search of genes involved in pathways associated with tuber quality using combined**
150 **strategies**
151
152 To better understand the genetic basis of *D. alata* tuber quality, we focused on the genes described
153 to be involved in one of the three pathways targeted (pentose and glucuronate interconversions,
154 starch and sucrose metabolism, and flavonoid biosynthesis). To obtain the most complete set of

genes from these pathways, we used two complementary strategies: a keyword search and a comparative genomics approach. For the first strategy, genes annotated on the *D. alata* reference genome (Bredeson et al., 2022) with the enzyme commission numbers (EC number) in one of the three pathways were selected. In total, we could retrieve 322 genes from these pathways encompassing 177 EC numbers (Figure 3A; Supplemental Table 2). The pentose and glucuronate interconversions pathway include most genes (150) and EC numbers (73), followed by starch and sucrose metabolism (136 genes, 76 EC numbers) and flavonoid biosynthesis metabolism (83 genes, 28 EC numbers). To complement the set of annotated genes, a comparative genomics approach was employed using a comparison of 45 diverse plant genomes (Mota et al., 2020) (Supplemental Table 3). Groups of orthologous predicted proteins from different plants (OG), were used to infer the function of proteins that were not annotated by other methods. Since the clustering of proteins in OG does not depend on their previous function annotation but rather on their protein sequence, this method can uncover missing genes from a keyword search. The orthology analysis resulted in a total of 58,916 OG from 1,706,645 proteins (weblink dataverse Cirad). Out of the total proteins analysed, 93.7% could be assigned to an OG. We identified 3,982 OG that are commonly found in all 45 species, despite their significant evolutionary distance. As expected, species that are closely related either phylogenetically or by genus have a greater number of OG in common than distantly ones (Supplemental Figure 4A, B). When we narrowed our comparison to monocotyledons (which comprised 18 species), we found a total of 5,544 OG that are commonly shared among the species. Only three OG were not present in *D. alata* (Figure 3B). The low number of missing groups in *D. alata* species was also observed by Bredeson et al. (2022), when comparing it with species exclusively from the Dioscoreales family.

Among the five Dioscoreales species, we found 8,029 OG in common. *Trichopus zeylanicus* showed the highest number of exclusive OG (1,951) compared to the other *Dioscorea* species (*D. dumetorum*, *D. rotundata*, *D. zingiberensis*, and *D. alata*) and *D. alata* the lowest one (336) (Figure 3C). Additionally, *D. alata* and *D. rotundata* had a high similarity, and likely underwent the same duplication events (Bredeson et al., 2022). We identified 14 OG that comprise 163 exclusive proteins of Dioscoreales (Supplemental Table 4), with most of these proteins (39) belonging to *D. alata*. Based on their EC numbers, most of these OG are hydrolases. They are not exclusive to *Dioscorea* species but their sequence conservation among other species is low, resulting in exclusive OG (*i.e.,* species-specific).

Once we identified all the OG that included representatives from *D. alata*, we narrowed these

groups down to the three pathways targeted in this study. For most of the OG found in the three

pathways, at least one gene from each of the 45 species was present, or the majority of them

(weblink dataverse Cirad; Supplemental Figure 4B). These OG were not exclusive to any

particular phylogenetic clade, genus, or species, indicating the importance of these proteins in

biological processes for all plant species. Although present in all species, the number of proteins

for each OG was uneven, which could be the result of both bias originating from assembly and

annotation discrepancies, as well as duplication events that occurred after the differentiation of

certain species (in-paralogs) (Pont et al., 2011).

The comparative analysis was not used to increase the number of EC numbers for these pathways

but to complete the set of previously annotated genes (Bredeson et al., 2022). We could retrieve

new protein coding genes from the comparative genomics analysis which were not found from the

keyword search. Thus, we found additional 137 genes for the pentose and glucuronate

interconversions pathway, 12 genes for starch and sucrose metabolism pathway and 46 genes for

flavonoid biosynthesis metabolism (Figure 3A; Supplemental Figure 5, 6, 7). We were able to

associate one orthologous group to each EC number, which contains one or more genes, except

for the pentose and glucuronate interconversions pathway, for which the 73 EC numbers were

collapsed in only 57 orthogroups. For the other two pathways, each EC number corresponds to an

orthologous group. After combining these two approaches, we could retrieve a total of 564 genes

from 161 orthogroups for the three pathways, representing 48% more genes than using only one

strategy. These genes were distributed along all the chromosomes of *D. alata* with chromosomes

2, 5, 11, and 17 having the highest number of genes and chromosome 16 the lowest (Figure 1B).

To determine the polymorphism within the genes found in the three pathways, we selected the

open read frame regions since we are only interested in the genic polymorphisms in this study. A

total of 406,325 SNPs on 564 candidate genes, from the three pathways was selected. By keeping

only SNPs annotated as high impact (*i.e.* non-synonymous, stop codon gain, frame shift)

responsible for the most significant differences on the protein sequences, a final number of 4,287

SNPs was used for further analysis, with the pathway of pentose and glucuronate interconversions

having the highest number (2,912), followed by starch and sucrose pathway (765) and flavonoid

biosynthesis (610).

Identification of genes under selection in the three targeted pathways

Once we have determined the complete set of *D. alata* genes belonging to the three pathways, we searched for regions under selective pressure, using *Fst* analysis. When comparing the 564 genes from the three pathways to the 1,053 genes from the *Fs*t analysis, we found 27 genes in common including nine genes found exclusively through the comparative genomic approach (Figure 3D, Supplemental table 5). The majority of the 158 SNPs on these 27 candidate genes were on the pentose and glucuronate interconversions pathway (119 polymorphisms in 10 genes) only found between Cluster B (mostly African genotypes) and C (Asian genotypes), followed by starch and sucrose (23 polymorphisms in 7 genes) and flavonoids (16 polymorphisms in 10 genes).

On the pentose and glucuronate interconversions pathway, we found three pectate lyase genes (*Dioal.01G025700.1*, *Dioal.01G025800.1*, *Dioal.01G035100.1*), which together with the polygalactunorase gene (PG) (*Dioal.06G005800.1*) and pectin methylesterase gene (PME) (*Dioal.08G105700.1*) are known as pectin-degrading enzymes involved in the demethylesterification of homogalacturonans (Leng et al., 2017). PME family acts prior to hydrolysis by PG enzyme that degrades the homogalacturonate backbone (polygalacturonate). Pectate lyase gene family is responsible for the cleavage of pectate, which is the product of pectin degradation. From the same pathway, we also found four malectin-like receptor-like kinase genes (MLD) under selection (*Dioal.20G055000.1*, *Dioal.20G055100.1*, *Dioal.20G055200.1*, *Dioal.20G055300.1*). The MLD genes, frequently erroneously described as Receptor-Like Kinases, can bind to pectin, with a preference for more highly methyl esterified pectin, causing a loosening in the cell-wall (Li et al., 2016).

On the starch and sucrose pathway, seven genes were under selection. Among them, two cellulose synthase genes (*Dioal.08G106000* and *Dioal.08G130500*), were under selection between clusters B and C. Additionally, we identified a starch synthase gene (*Dioal.09G032200*) and a nudix-hydrolase (*Dioal.11G000200*), under selection among A and C and among A and B, respectively, a 1,3-beta-glucan synthase (*Dioal.10G014800*), among A and B and a trehalose-6-phosphate phosphatase (*Dioal.04G184200*) under selective pressure among A and C genetic clusters. Those genes are responsible for starch synthesis in plants (Kolbe et al., 2005; Muñoz et al., 2006; Keeling and Myers, 2010). Trehalose phosphate phosphatase is involved in the synthesis of Trehalose consisting of two 1,1-linked glucose molecules (Kolbe et al., 2005). Lastly, we found a beta-

amylase gene (*Dioal.04G168200*) under selection. The beta-amylase gene family was previously described to increase the firmness of cooked sweet potatoes when its enzyme activity is low (Banda et al., 2021).

On the flavonoids pathway were found under selection 2-oxoglutarate-dependent dioxygenase (*Dioal.01G016100* and *Dioal.01G016200*) and chalcone isomerase (*Dioal.08G131200.1*). The involvement of these genes was mostly associated with plant-pathogen interaction, and their relationship with quality traits remains to be investigated (Fang et al., 2022).

Candidate gene association studies for quality traits in *D. alata*

To get a deeper insight in the involvement of the genes annotated from the three metabolic pathways on the tuber quality, we conducted a candidate gene association study. We used phenotypic data collected from two-years field trials on 45 genotypes among the 125 *D. alata* genotypes. They were selected to cover the genetic diversity and to minimize the genetic structure. A set of 10 traits related to starch content, colour indices (Brown Index, the Hue for purple and the Browning over time) and texture were measured. Texture was measured with two different types of probes. A conical probe was used to measure the total area (N sec), which reflects the total force required to penetrate the sample at a constant speed. A flat plate probe was used during a double compression cycle to measure hardness, gumminess, springiness, and cohesiveness. Descriptive statistics of the various tuber quality traits evaluated in this study are shown in Supplemental Table 6. The colour indices BI, HI exhibited wide range of variation (12.57-143 and -11.99-99.92, respectively), showing the presence of varying coloured yam tubers in the association panel. The tuber SC was on average 77.93±2.10 with the minimum (69.46 %) and maximum (81.26 %) observed in CRB417 and Roujol9, respectively. The textural traits also varied extensively within the panel, indicating varying boiled yam qualities of the accessions (Supplemental Table 6). Taking together, our results indicate a wide variation in the tuber quality traits within the 45 genotypes and pinpoint that these traits are largely quantitative in nature.

Based on the general linear model, we obtained significant associations between SNPs and the phenotypic data. We detected a total of 22 quantitative trait nucleotides (QTNs) distributed on 10 chromosomes, for seven out of the ten traits evaluated in this study (Table 1). Hereafter, we present in more details five key genes associated with textural traits, starch content and colour. The

expression profiles of the candidate genes were assessed in six phenotypically diverse genotypes
using transcriptomic analysis.

Out of the 22 QTNs, 17 were detected for texture traits with fourteen and three within pentose and
glucuronate interconversions, and sucrose and starch pathways, respectively (Table 1 and
Supplemental Figure 8). Within pentose and glucuronate interconversions pathway, we found
pectin esterase (*Dioal.19G187400.1*) and xylose isomerase genes (*Dioal.13G053400*) associated
to springiness and gumminess. Leucine-rich repeat (*Dioal.10G024400)*, protein kinases
(*Dioal.02G062700*, *Dioal.02G062800*), a malectin-like encoding gene (*Dioal.20G055400*) were
associated to springiness and had a stop gain caused by a transition and by a transversion. Two
other malectin-like genes found under selection were associated to hardness. Within starch and
sucrose pathway we found three candidate genes associated to gumminess, total area and hardness,
α-amylase (*Dioal.07G051400.1*), 4-α-glucanotransferase *(Dioal.07G067100)* and beta-
glucosidase (*Dioal.13G093500*), respectively.

Pectin methylesterase (*Dioal.04G130600.1*), was associated to gumminess and total area and
found under selection. A tuber pectin methyl esterase activity has been identified as a potential
factor impacting cooked potato textural properties (Ross et al. 2011). The SNP S4_20715506 (C/T)
located in the only exon of *Dioal.04G130600.1* (Figure 4A), results in an amino acid change (H to
W), with a potential impact on the enzyme activity. We observed that the C-allele at S4_20715506
decreases gumminess and total area (Figure 4B and C). None of the accessions used for
transcriptome analysis had the T-allele, hence we were unable to evaluate the expression pattern
of this gene between two haplotypes. Nonetheless, our results suggest that allelic variation in
Dioal.04G130600.1 probably affects the enzyme activity, leading to varying textural properties of
boiled yam.

Beta-glucosidase, associated to hardness, is a cellulase enzyme playing a part in metabolism of
cell wall polysaccharides (Sampedro et al., 2017). The alternative C-allele at the qH13.2 had a
strong effect on hardness (62%) by modifying the splice donor on position 837 in the gene
Dioal.13G093500 (Table 1). The genotypes CRB47, Roujol49, and Roujol75 which are
homozygous for the G-allele showed an overall higher expression than the genotypes with the C-
allele (CRB96, Roujol62, and Roujol9).

The variation of the starch content in the association panel was associated with three QTNs (Table
1). They were annotated as 1,3-beta-glucan-synthase (*Dioal.10G014800*), beta-amylase

(*Dioal.15G119500*) and glucose-1-phosphate adenyl transferase (*Dioal.17G112400*). The 1,3-
beta-glucan-synthase had the highest effect on the starch content and was also found in an outlier
region by our *Fst* analysis (Supplemental Table 5). A recent study demonstrated that manipulation
of this gene induces different starch and polysaccharide profiles in barley grain (Garcia-Gimenez
et al. 2020). The genotypes harbouring the G-allele had a premature stop codon instead of a
glycine. They had 14% less starch content compared to the genotypes with the T-allele. We
speculated that the gain of a stop codon results in a non-functional or altered protein which affects
the SC.

Lastly, within the flavonoid biosynthesis pathway, we identified two QTNs qHI4.1 and qBI19.1
for Brown index (BI) and HUE index (HI), respectively (Table 1). The gene *Dioal.19G083800.1*
encodes for flavanone 3-hydroxylase (F3H). It catalyses the 3-beta-hydroxylation of 2S-
flavanones to 2R,3R-dihydroflavonols which are intermediates in the biosynthesis of major
pigments (flavonols, anthocyanidins, and proanthocyanidins) (Falcone Ferreyra et al., 2012).
Dioal.19G083800.1 contains SNP S19_17029453 (G/C) in the last exon, with the G-allele leading
to a premature stop codon, a truncated protein and likely a non-functional enzyme (Figure 4A).
S19_17029453 was significantly associated with BI. The genotypes with the G-allele tend to have
brown colour in contrast to the genotypes with the C-allele displaying mostly whitish tuber flesh
(Figure 4D). Thus, this mutation may block the formation of flavonoid compounds and redirect
the phenylpropanoid pathway to the formation of other brownish compounds like oxidized
phenolic acids. Expression pattern of *Dioal.19G083800* between two contrasting accessions with
G/C alleles showed that it was expressed nearly 20 times higher in heterozygous (Roujol75; C/G)
than in homozygous (Roujol62; G/G) genotypes (Figure 4E). Overall, these results suggest that a
functional mutation in *Dioal.19G083800* partly modulates colour formation in yam tuber.

Another key gene detected for colour variation (purple and non-purple yam varieties) was
Dioal.04G073400, encoding for UDP-glucuronosyltransferase (UGT). Anthocyanins are unstable
water-soluble pigments. UGTs are key enzymes that stabilizes anthocyanidin by attaching sugar
moieties to the anthocyanin aglycone (Li et al. 2016). In this study, the T-allele at the SNP
S4_16031453 leads to a stop codon in the middle of *Dioal.04G073400* (Figure 4A), which will
probably result in a non-functional enzyme. Intriguingly, all accessions with the T-allele displayed
purple tuber (very low HI score) (Figure 4F). This could be explained by the fact that non-

functionality of an enzyme producing uncoloured flavanones could redirect the pathway to

anthocyanins biosynthesis resulting in the purple colour of the tubers.

Altogether, the candidate gene association analysis showed that variants in several genes affect the

functions and potentially the expression of key structural genes involved in the regulation of yam

tuber quality traits. Interestingly, only 38% of the genes found in this analysis were retrieved from

the keyword annotation, while the remaining were retrieved by comparative genomics.

DISCUSSION

In this study, the use of comparative genomics and genome sequence analysis of 127 *Dioscorea*

alata accessions enabled the identification of candidate genes in the main metabolic pathways

involved in yam tuber quality. The functional annotation of the *D. alata* genome remains

suboptimal since there are few genomic data available for this crop (Chaïr et al., 2022). Only

recently a complete *D. alata* genome reference was published (Bredeson et al., 2022). Therefore,

the comparative genomics combined with keyword search used in this study, allowed the

identification of novel genes associated with the three metabolic pathways targeted, pentose and

glucuronate interconversions, starch and sucrose metabolism, and flavonoid biosynthesis

pathways. We found more paralogs for each metabolic pathway, even if the number of ECs

remained the same. We annotated orthologous groups using the conserved domains, which

facilitated the correction of incomplete or outdated gene assignments, such as for Malectin-like

genes. In addition, based on the orthology analysis we found gene structure errors, such as for

OG0082990 and OG0083049 (weblink CIRAD dataverse), which calls for manual curation of

important gene families to further improve the functional annotation of *D. alata* genome.

We found that various gene ontology categories were enriched from the genes under selection

between the three identified genetic groups. The most enriched pathways under selection were

transmembrane transport, protein phosphorylation, response to wounding, hormone signalling,

lignin biosynthetic process, known to play important physiological roles in defence response

during plant-stress interactions (Zhang and Sun, 2021). This indicates most probably a selection

for adaptation to biotic and abiotic conditions. The downsizing to the three pathways revealed

under selection cellulose synthase, required for primary and secondary cell wall cellulose synthesis

(Richmond and Somerville, 2000), pectate lyase and PG known as pectin degrading enzymes.

Additionally, we also identified genes involved in starch synthesis, nudix hydrolase, soluble starch

synthase 3 and 1-3 beta-glucan synthase (Miao et al., 2017), confirming the importance of starch

content in yam during farmer selection and most probably during domestication (Meyer and

Purugganan, 2013). Genes for the pentose and glucuronate interconversion pathway were found

only between African and Asian genetic groups. This result deserves further investigation to

understand this exclusivity. Five genes from the pectin and starch-sugar pathways were found in

common between the *Fst* analysis and candidate gene association study, showing that the quality

of yam tubers was also a factor of differentiation and selection during yam evolution.

The candidate gene association analysis enabled the identification of fourteen pectin and three

starch candidate genes associated with texture, confirming the involvement of the cell wall and

starch content in boiled yam texture (Kouadio et al., 2013). Among the seventeen candidate genes,

three have high effect and are involved directly in the cell wall composition. Beta-glucosidase

explains more than 60% of the hardness confirming it as a major QTN in boiled yam texture. Its

role in cell wall thickness has been demonstrated in different crops such as potato (Ross et al.,

2011) or barley in which it participates in endosperm cell wall degradation during germination

(Leah et al., 1995). In cotton (*Gossypium hirsutum*), the overexpression of the *GhBG1A* a gene

encoding beta-glucosidase repressed fiber length but promoted cellulose biosynthesis resulting in

thicker fiber cell wall (Watcharatpong et al., 2020). The four remaining predicted genes with high

effect on texture parameters were involved in pectin pathway. Pectin methylesterase, found also

under selection, encodes for an enzyme which plays an important role in both pectin remodelling

and disassembly and consequently in firming and softening of cell wall (Atmodjo et al., 2013). In

our study, it has a negative effect on both total area and gumminess indicating its involvement in

pectin degradation during yam boiling, such as in cooked potato (Ross et al., 2011). In cassava, its

involvement in the root softening process during cassava retting was demonstrated in addition to

pectin/pectate lyase and polygalacturonase genes (Ngolong Ngea et al., 2016). These two last

genes were found under selection but not associated with any trait investigated. Finally, Xylose

isomerase seems to play an important role in boiled yam texture. It was associated with gumminess

and springiness most probably by catalysing the reversible isomerisation of pentoses such as
arabinose, one of the components of pectin side chain (Mu et al., 2018).

Such as for cereals and other tuber crops, the total starch content of yams is a primary determinant
of tuber quality. Preferred mealy cooking yam cultivars had significantly higher starch content
(Kouadio et al., 2013). Among the three genes associated with starch content, the 1,3-beta-glucan
synthase had the highest effect. It is often considered to be a cellulase family member, plays an
important role in cellulose structure. In this study this gene family was found under selection and
predicted to be associated with starch content. The (1,3;1,4)-beta-D-glucans are most abundant in
walls of the cereals, specifically in the starchy endosperm of grain, where they can contribute up
to 70% by weight of the walls in barley, rye, and oats (Burton and Fincher, 2012). Whether it is
related to starch content on yam or other tuberous starchy crops remains unclear.

The colour is a determinant key trait in yam varieties adoption. While in West Africa, white colour
is preferred, in the Pacific or Caribbean colourful plates are more appreciated. In our study the
flavanone 3-hydroxylase (F3H) was found involved in brown index with a high effect. This gene
is involved in the accumulation of catechins in tea plants (*Camelia sinensis* L.) (Singh et al., 2008).
In carnation flower (*Dianthus caryophyllus* L.) it has been found involved in colour and fragrance
(Zuker et al., 2002). RNAi-Mediated silencing of the F3H confirmed that this gene is one of the
key enzymes required for the biosynthesis of flavonoids in strawberry fruit (Jiang et al., 2013).
The UDP-glucosyltransferase was found associated with purple colour confirming its role in
colour formation in yam tuber. None of these genes were found by Wu et al. (2015) through
transcriptome analysis of two contrasting *D. alata* genotypes for tuber purple colour, or by a
genome-wide association analysis (Dossa et al., 2023). These results indicate that more genes
could be involved in this trait. Moreover, we mainly focused our study on SNPs located in coding
regions, and only structural genes involved in three targeted pathways were investigated. This
approach therefore excludes the regulatory genes such as transcription factors as well as other
candidate genes and genome components that control agronomic traits. Similarly, we did not
observe significant gene expression changes between contrasting genotypes for the all identified
candidate genes, probably because coding SNPs mainly affect protein functions but at a less extent
transcript levels.

In conclusion, quality is one of the main criteria selected during evolution and adaptation, which
is reflected in the genes identified in this study. We found twenty-two candidate genes, in pentose
and glucuronate, sucrose and starch and flavonoid biosynthesis pathways, associated to the three
main attributes of boiled yam: texture, starch content and tuber colour. This is the first work that
considers the different cell wall constituents and their effect on texture. In addition, we were able
to confirm the expression of five candidate genes in these pathways. Further validation of these
results on a larger panel of associations by performing a genome-wide association analysis
combined with metabolomic and transcriptomic analyses will confirm the robustness of this
approach. Validation of the identified genes will pave the way for favourable allele pyramiding in
breeding programmes. Moreover, the use of comparative genomics and the keyword search
approach to retrieve genes from the three pathways has proved effective. We have been able to
retrieve new protein-coding genes that were not found in the keyword search. This approach,
applied to orphan crops for which few genomic resources are available, will improve the
annotation of their genomes.

Material and Methods

Plant material

This study used 125 genotypes from *Dioscorea alata* and two outgroup genotypes, belonging to
D. rotundata and *D. trifida* species, collected in 19 different countries. Genotypes were selected
to cover a maximum of worldwide diversity (Figure 1; Supplemental Table 1) based on previous
studies (Sharif et al., 2020). These genotypes were used to find genomic polymorphism, compared
to the *D. alata* reference genome. From the whole set used in this study, a total of 45 *D. alata*
accessions harbouring a high genetic diversity were planted from 2018 to 2020 in May each year
and harvested at full maturity in the following year between January and March at three different
locations in Guadeloupe (Godet, Roujol and Duclos). The three locations have contrasting pedo-
climatic conditions (Cormier et al., 2021). The phenotyping was performed on freshly harvested
tubers. Finally, a set of six genotypes was selected from the list, based on their previously described
phenotypic traits, to be used for transcriptome sequencing (Supplemental Table 1).

Phenotyping for quality traits

Colour was analyzed by computerized image analysis techniques (Detailed method is described in

Supplemental Note 1). Images were collected on three tubers per genotype (45) and location (3)

during the 2018 cropping season. To test the repeatability and accuracy of the system, a custom

colour chart was placed on every picture. Acquired images were analyzed using the Rvision library

(Garnier and Muschelli, 2022) with R programming language (R Core Team, 2021). Different

colour indexes were tested to characterize purple yam. Finally, the Hue was found to be

discriminant within the studied diversity panel. Brown (BI) index was also calculated according

to (Buera et al., 1986).

Starch content was predicted using near infrared spectroscopy (NIRS). Two replicates of yam flour

samples were scanned with a FOSS-NIRSystems model 6500 scanning monochromator (FOSS-

NIRSystems, Silver Spring, MD, USA) equipped with an autocup. The spectroscopic procedures

and data recording were conducted with ISIscan (TM) software (FOSS, Hillerød, Denmark). The

model was calibrated using 2016 and 2017 data and validated on an external independent 2018

dataset. (Supplemental Note 2).

The texture was measured on steam-cooked yam by penetrometry using the TAX-TPlus texture

analyzer (Stable Micro Systems, Ltd., Surrey, UK). Yam tubers harvested at Godet 2019 were

used for this analysis. Three tubers per variety were sampled and divided into three equal sections

(proximal, distal, central) and prepared as described in Supplemental Note 3. Each part was used

to produce three cubes at the central section. Each cube was steam-cooked up to 15 min, followed

by a cooling time of 7 min, corresponding to a cube temperature of 45°C. Total area was calculated

from the puncture test, while four texture profile analyzer parameters were computed from the

force-time curve: hardness (N), cohesiveness, gumminess, and springiness (Supplemental Note 3).

Whole-genome sequencing

The genomic DNA from 127 genotypes of *Dioscorea spp*. was extracted from leaves according to

a protocol using mixed alkyl trimethylammonium bromide (MATAB) buffer and NucleoMag Plant

Kit (Macherey–Nagel, Germany) already described by (Cormier et al., 2019). Sequencing libraries

were prepared as described in Dossa et al. (2023). Paired-end high throughput sequencing (2 × 150

bp) was performed on an Illumina NovaSeq 6000 instrument on GeT-PlaGe platform, (Toulouse,

France) and Genewiz company (Leipzig, Germany).

The reads quality and mapping and Variant discovery and filtering were performed as described

in Dossa et al. in press. For the filtering, the following parameters were used: --max-missing 0.5 -

-minQ 30 --minDP 10 --maxDP 200. For gene candidate purposes, we selected only the

polymorphisms annotated with a high impact on the amino acid sequences.

Genetic population analyses

Among all the 127 genotypes of *Dioscorea* spp., 107 diploid genotypes (Supplemental Table 1)

were selected for further population genetics analyses. We calculated the population structure

using the original filters described above, in addition to: --maf 0.01 --max-alleles 2 --min-alleles

2 --remove-indels --max-missing 0.5 --thin 10 --ld-window 50 --min-r2 0.1. The final VCF file

obtained from this analysis was used for population structure analysis using ADMIXTURE version

1.23 (Alexander and Lange, 2011). After testing the replicates of K = 1 to 15, we limited the

ancestry threshold to 75%, where genotypes with a value lower than this were considered in the

admixture group. After Bayesian clustering analysis, populations were redefined according to the

results obtained.

To identify genes under selection, *Fst* analysis was conducted using vcftools with an *Fst*-window-

size of 50kb and a *Fst*-window-step of 10kb. A cutoff of the 5% top values and a Weibull

distribution analysis on regions with extreme *Fst* between the different genetic clusters identified,

were used to select candidates. Subsequently, we searched for the genes on these regions using

BEDTools intersect version 2.29 (Quinlan and Hall, 2010).

The gene ontology enrichment analysis of the genes from outlier regions was performed with

KOBAS-i webserver (Bu et al., 2021) in order to find out their biological significance.

Comparative genomics and orthology analyses

For comparative genomics analyses 45 species from diverse plant lineages were selected

(Supplemental Table 2). The species used in this analysis were chosen based on the completeness

of their genomes, phylogenetic distribution along the tree of viridiplantae and in the use of plant-

models. These plants belong to the monocotyledons and dicotyledons, encompassing 17 different

clades. Five genomes belonging to Dioscoreales were added (*Dioscorea alata, D. rotundata, D.*

dumetorum, D. zingiberensis, Trichopus zeylanicus). The completeness score of each plant

proteome was verified by BUSCO (Simão et al., 2015) using the Viridiplantae dataset of BUSCO.

We chose a threshold of 70% of completeness to include public proteomes. For the inference of

orthologous groups, OrthoFinder software v2.4.0 (Emms and Kelly, 2015) was used with the

default parameters. Gene presence and absence of species in orthogroups was represented by

UpSetR (Lex et al., 2014), using a binary matrix as input.

Metabolic pathways selection and gene retrieval from D. alata

We downloaded the complete list of enzyme codes (EC) from the pathway of pentose and

glucuronate interconversions (map00040), starch and sucrose metabolism (map00500) and

flavonoid biosynthesis (map00941) from KEGG database (https://www.genome.jp). These EC

numbers were used to retrieve the genes annotated in the GFF file of *D. alata* (https://phytozome-

next.jgi.doe.gov/info/Dalata_v2_1). The genes annotated with these EC numbers were further

searched in the orthogroups produced by our comparative genomics study, and all the homologous

genes from *D. alata* were retrieved. The VCF file was reduced to the middle coding sequences of

the annotated genes of interest.

Candidate gene association analysis

Three VCF files, one for each target metabolic pathway genes, containing non-synonymous SNPs

were extracted from the whole VCF file and further filtered for minor allele frequency ≥ 0.05,

missing rate $< 20\%$ using TASSEL5.0 (Bradbury et al., 2007). The associations between the

polymorphisms from each metabolic pathway candidate genes and the corresponding phenotypic

traits were tested using TASSEL5.0 based on the General Linear Model. Only significant variants

with $P \leq 0.001$ were retained. The association analysis was conducted with phenotypic data from

each location independently. The effect of alleles at significant SNPs was assessed by comparing

phenotyping data for haplotype groups. A Student's t-test was used to compare the groups of

haplotypes ($P<0.05$) in the R4.0.23 software with the "ggpubr" packages and "rstatix"

Transcriptome analysis of six D. alata genotypes

RNA was extracted from six of the most *D. alata* diverse genotypes, based on phenotypic traits.

Three biological replicates for each genotype were used. For each genotype, a 50 mg sample cut

from the middle of freshly harvested tuber was ground in liquid nitrogen and total cellular RNA

was extracted using a Sigma-Aldrich (St. Louis, MO) Spectrum™ Plant Total RNA kit with a

DNAse treatment. This was followed by quantification by the Invitrogen (Carlsbad, CA) Quant-

iT™ RiboGreen® RNA Reagent based on the manufacturer's protocol and verification of RNA

quality by 5200 Fragment Analyzer™ System (Agilent, Santa Clara, CA) profile.

Synthesis of cDNA and construction of libraries were done with Illumina, Inc. TruSeq RNA

Sample Preparation v2 Kit (San Diego, CA). Fragment size of selected cDNA were between 200pb

and 400pb. The 18 libraries were indexed, mixed and sequenced using one lane of an Illumina

HiSeq 3000 sequencer with the 2 x 150 cycles, paired-end, indexed protocol (Genewiz facility,

Liepzig, Germany).

The quality of the raw reads from the 18 libraries were assessed by FastQC version 0.11.7

(Andrews, 2010). The low-quality sequences and the Illumina adapters were trimmed by

Trimmomatic version 0.39 (Bolger et al., 2017). Trimmed reads were quantified using Kallisto

version 0.46.1 (Bray et al., 2016) using the reference genome of *D. alata* (Bredeson et al., 2022).

To conduct the expression analysis of the genes, we used the log10 of the average value of

transcripts per million over the three biological replicates. A functional annotation was performed

using the conserved domains of PFAM version 35 and a gene ontology annotation using Pfam2go

(http://current.geneontology.org/ontology/external2go/pfam2go).

All the bioinformatics analyses were performed with the support of MESO@LR-Platform at

University of Montpellier, CIRAD UMR-AGAP HPC (South Green Platform) and IFB core

cluster.

DATA AVAILABILITY

570 The Illumina HiSeq 3000 sequencing raw data are available in the NCBI SRA (Sequence Read

571 Archive), and transcriptome data are under the BioProject number: PRJNA918625. The

572 phenotypic datasets are available from the corresponding author upon request.

FUNDING

575 We acknowledge the support from Breeding RTB Products for End User Preferences (RTBfoods)
576 Project (Grant OPP1178942) through funds received from the Bill and Melinda Gates Foundation.
577 The production of the genomic resources was supported by a fund from the CGIAR Research
578 Program on Roots, Tubers and Bananas (CRP-RTB).

ACKNOWLEDGMENTS

We are grateful to Christian Mestres for his valuable advice on yam quality and its attributes. We
gratefully acknowledge Nancy Terrier for her critical reading of the manuscript. Our greatest debts
go to CRP-RTB, IITA and PRC genebanks and Juliane Kaoh, Mamy Tiana Rajaonah, Senanayake
Ravinda Lakshan, and Babil Pachakkil for providing us with plant leaves material. We are also
thankful to Elie Nudol, Marie-Claire Gravillon, Christophe Perrot and Erick Maledon for their
invaluable help with field work and phenotyping, and Sandrine Causse and Hélène Vignes for
laboratory assistance in preparing *D. alata* libraries. Finally, special thanks are due to Sylvain
Santoni and Muriel Latreille for their assistance in transcriptome preparation.

REFERENCES

Alexander, D. H., and Lange, K. (2011). Enhancements to the ADMIXTURE algorithm for individual ancestry estimation. *BMC Bioinformatics* **12**:246.

Alves-Pereira, A., Zucchi, M. I., Clement, C. R., Viana, J. P. G., Pinheiro, J. B., Veasey, E. A., and de Souza, A. P. (2022). Selective signatures and high genome-wide diversity in traditional Brazilian manioc (Manihot esculenta Crantz) varieties. *Sci. Rep.* **12**:1268.

Andrews, S. and others (2010). FastQC: a quality control tool for high throughput sequence data Advance Access published 2010.

Arnau, G., Nemorin, A., Maledon, E., and Abraham, K. (2009). Revision of ploidy status of Dioscorea alata L. (Dioscoreaceae) by cytogenetic and microsatellite segregation analysis. *Theor. Appl. Genet.* **118**:1239–1249.

Atmodjo, M. A., Hao, Z., and Mohnen, D. (2013). Evolving Views of Pectin Biosynthesis. *Annu. Rev. Plant Biol.* **64**:747–779.

Banda, L., Kyallo, M., Domelevo Entfellner, J.-B., Moyo, M., Swanckaert, J., Mwanga, R. O. M., Onyango, A., Magiri, E., Gemenet, D. C., Yao, N., et al. (2021). Analysis of β-amylase gene (Amyβ) variation reveals allele association with low enzyme activity and increased firmness in cooked sweetpotato (Ipomoea batatas) from East Africa. *J. Agric. Food Res.* **4**:100121.

Beleia, A., Butarelo, S. S., and Silva, R. S. F. (2006). Modeling of starch gelatinization during cooking of cassava (Manihot esculenta Crantz). *LWT - Food Sci. Technol.* **39**:400–405.

Bhosale, S. H., Rao, M. B., and Deshpande, V. V. (1996). Molecular and industrial aspects of glucose isomerase. *MICROBIOL REV* **60**.

Bolger, M. E., Arsova, B., and Usadel, B. (2017). Plant genome and transcriptome annotations: from misconceptions to simple solutions. *Brief. Bioinform.* Advance Access published January 5, 2017, doi:10.1093/bib/bbw135.

Bradbury, P. J., Zhang, Z., Kroon, D. E., Casstevens, T. M., Ramdoss, Y., and Buckler, E.

S. (2007). TASSEL: software for association mapping of complex traits in diverse

samples. *Bioinformatics* **23**:2633–2635.

Bray, N. L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic

RNA-seq quantification. *Nat. Biotechnol.* **34**:525–527.

Bredeson, J. V., Lyons, J. B., Oniyinde, I. O., Okereke, N. R., Kolade, O., Nnabue, I.,

Nwadili, C. O., Hřibová, E., Parker, M., Nwogha, J., et al. (2022). Chromosome

evolution and the genetic basis of agronomically important traits in greater yam. *Nat.*

Commun. **13**:2001.

Bu, D., Luo, H., Huo, P., Wang, Z., Zhang, S., He, Z., Wu, Y., Zhao, L., Liu, J., Guo, J., Fang,

S., Cao, W., Yi, L., Zhao, Y., & Kong, L. (2021). KOBAS-i: intelligent prioritization and

exploratory visualization of biological functions for gene enrichment analysis. *Nucleic*

Acids Research **49**:W317–W325.

Buera, M. P., Lozano, R., and Petriella, C. (1986). Definition of colour in the non enzymatic

browning process. *Farbe* **32**:318–322.

Burton, R. A., and Fincher, G. B. (2012). Current challenges in cell wall biology in the cereals

and grasses. *Front. Plant Sci.* **3**.

Butelli, E., Licciardello, C., Ramadugu, C., Durand-Hulak, M., Celant, A., Reforgiato

Recupero, G., Froelicher, Y., and Martin, C. (2019). Noemi Controls Production of

Flavonoid Pigments and Fruit Acidity and Illustrates the Domestication Routes of

Modern Citrus Varieties. *Curr. Biol.* **29**:158-164.e2.

Chair, H., Arnau, G., and Mota, A. P. Z. Yam Genomics. In *Underutilised Crop Genomes*, pp.

373–398. Springer International Publishing.

Cormier, F., Lawac, F., Maledon, E., Gravillon, M.-C., Nudol, E., Mournet, P., Vignes, H.,

Chaïr, H., and Arnau, G. (2019). A reference high-density genetic map of greater yam

(Dioscorea alata L.). *TAG Theor. Appl. Genet. Theor. Angew. Genet.* **132**:1733.

Cormier, F., Martin, G., Vignes, H., Lachman, L., Cornet, D., Faure, Y., Maledon, E.,
Mournet, P., Arnau, G., and Chaïr, H. (2021). Genetic control of flowering in greater
yam (Dioscorea alata L.). *BMC Plant Biol.* **21**:163.

Critchley, J. H., Zeeman, S. C., Takaha, T., Smith, A. M., and Smith, S. M. (2001). A critical
role for disproportionating enzyme in starch breakdown is revealed by a knock-out
mutation in Arabidopsis: Function of D-enzyme in starch breakdown. *Plant J.* **26**:89–100.

Das, M., Haberer, G., Panda, A., Das Laha, S., Ghosh, T. C., and Schäffner, A. R. (2016).
Expression Pattern Similarities Support the Prediction of Orthologs Retaining Common
Functions after Gene Duplication Events. *Plant Physiol.* **171**:2343–2357.

do Nascimento, J. R. O., Júnior, A. V., Bassinello, P. Z., Cordenunsi, B. R., Mainardi, J. A.,
Purgatto, E., and Lajolo, F. M. (2006). Beta-amylase expression and starch degradation
during banana ripening. *Postharvest Biol. Technol.* **40**:41–47.

Domínguez, M., Dugas, E., Benchouaia, M., Leduque, B., Jiménez-Gómez, J. M., Colot, V.,
and Quadrana, L. (2020). The impact of transposable elements on tomato diversity. *Nat.*
Commun. **11**:4058.

Dong, M.-Y., Fan, X.-W., and Li, Y.-Z. (2019). Cassava AGPase genes and their encoded
proteins are different from those of other plants. *Planta* **250**:1621–1635.

Dossa, K., Morel, A., Houngbo, E.M., Mota, A.Z., Maledon, E. et al. (2023). Genome-wide
association studies reveal novel loci controlling tuber flesh colour and oxidative browning
in *Dioscorea alata. bioRxiv* 2023.03.12.532275.

Dufour, D., Hershey, C., Hamaker, B. R., and Lorenzen, J. (2021). Integrating end-user
preferences into breeding programmes for roots, tubers and bananas. *Int. J. Food Sci.*
Technol. **56**:1071–1075.

Ehounou, A. E., Cormier, F., Maledon, E., Nudol, E., Vignes, H., Gravillon, M. C.,
N'guetta, A. S. P., Mournet, P., Chaïr, H., Kouakou, A. M., et al. (2022).
Identification and validation of QTLs for tuber quality related traits in greater yam
(*Dioscorea alata* L.). *Sci. Rep.* **12**:8423.

Emms, D. M., and Kelly, S. (2015). OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. *Genome Biol.*
16:157.

Falcone Ferreyra, M. L., Rius, S. P., and Casati, P. (2012). Flavonoids: biosynthesis,
biological functions, and biotechnological applications. *Front. Plant Sci.* **3**.

Fang, H., Zhang, F., Zhang, C., Wang, D., Shen, S., He, F., Tao, H., Wang, R., Wang, M.,
Wang, D., et al. (2022). Function of hydroxycinnamoyl transferases for the biosynthesis
of phenolamides in rice resistance to Magnaporthe oryzae. *J. Genet. Genomics* **49**:776–
786.

Garcia-Gimenez, G., Barakate, A., Smith, P., Stephens, J., Khor, S.F., Doblin, M.S., Hao,
P., Bacic, A., Fincher, G.B., Burton, R.A., Waugh, R., Tucker, M.R. and Houston,
K. (2020), Targeted mutation of barley (1,3;1,4)-β-glucan synthases reveals complex
relationships between the storage and cell wall polysaccharide content. Plant J., 104:
1009-1022.

Garnier, S., and Muschelli, J. (2022). *Rvision - A computer vision library for R*.

Gatarira, C., Agre, P., Matsumoto, R., Edemodu, A., Adetimirin, V., Bhattacharjee, R.,
Asiedu, R., and Asfaw, A. (2020). Genome-Wide Association Analysis for Tuber Dry
Matter and Oxidative Browning in Water Yam (Dioscorea alata L.). *Plants* **9**:969.

Girma, G., Hyma, K. E., Asiedu, R., Mitchell, S. E., Gedil, M., and Spillane, C. (2014).
Next-generation sequencing based genotyping, cytometry and phenotyping for
understanding diversity and evolution of guinea yams. *Theor. Appl. Genet.* **127**:1783–
1794.

Hardigan, M. A., Laimbeer, F. P. E., Newton, L., Crisovan, E., Hamilton, J. P.,
Vaillancourt, B., Wiegert-Rininger, K., Wood, J. C., Douches, D. S., Farré, E. M., et
al. (2017). Genome diversity of tuber-bearing *Solanum* uncovers complex evolutionary
history and targets of domestication in the cultivated potato. *Proc. Natl. Acad. Sci.* **114**.

He, Y., Zhu, D., Sun, Y., Wang, Q., Zhu, L., and Zeng, H. (2021). Metabonomic Profiling
695   Analyses Reveal ANS Upregulation to Enhance the Flavonoid Pathway of Purple-
696   Fleshed Sweet Potato Storage Root in Response to Deep Shading. *Agronomy* **11**:737.

697   **Hendriks, J. H. M., Kolbe, A., Gibon, Y., Stitt, M., and Geigenberger, P.** (2003). ADP-
698   Glucose Pyrophosphorylase Is Activated by Posttranslational Redox-Modification in
699   Response to Light and to Sugars in Leaves of Arabidopsis and Other Plant Species. *Plant*
700   *Physiol.* **133**:838–849.

701   **Honfozo, L., Adinsi, L., Bouniol, A., Adetonah, S., Forsythe, L., Kleih, U., Hounhouigan, J.
702   D., Fliedel, G., and Akissoe, N. H.** (2021). Boiled yam end-user preferences and
703   implications for trait evaluation. *Int. J. Food Sci. Technol.* **56**:1447–1457.

704   **Hou, J., Liu, T., Reid, S., Zhang, H., Peng, X., Sun, K., Du, J., Sonnewald, U., and Song, B.**
705   (2019). Silencing of α-amylase StAmy23 in potato tuber leads to delayed sprouting. *Plant*
706   *Physiol. Biochem.* **139**:411–418.

707   **Janeček, Š., Svensson, B., and MacGregor, E. A.** (2014). α-Amylase: an enzyme specificity
708   found in various families of glycoside hydrolases. *Cell. Mol. Life Sci.* **71**:1149–1170.

709   **Jiang, F., Wang, J.-Y., Jia, H.-F., Jia, W.-S., Wang, H.-Q., and Xiao, M.** (2013). RNAi-
710   Mediated Silencing of the Flavanone 3-Hydroxylase Gene and Its Effect on Flavonoid
711   Biosynthesis in Strawberry Fruit. *J. Plant Growth Regul.* **32**:182–190.

712   **Keeling, P. L., and Myers, A. M.** (2010). Biochemistry and Genetics of Starch Synthesis. *Annu.*
713   *Rev. Food Sci. Technol.* **1**:271–303.

714   **Kim, Y.-L., Mun, S., Park, K.-H., Shim, J.-Y., and Kim, Y.-R.** (2013). Physicochemical
715   functionality of 4-α-glucanotransferase-treated rice flour in food application. *Int. J. Biol.*
716   *Macromol.* **60**:422–426.

717   **Kolbe, A., Tiessen, A., Schluepmann, H., Paul, M., Ulrich, S., and Geigenberger, P.** (2005).
718   Trehalose 6-phosphate regulates starch synthesis via posttranslational redox activation of
719   ADP-glucose pyrophosphorylase. *Proc. Natl. Acad. Sci.* **102**:11118–11123.

Kouadio, O. K., N'dri, D. Y., Nindjin, C., Marti, A., Casiraghi, M. C., Faoro, F., Erba, D.,

Bonfoh, B., and Amani, N. G. (2013). Effect of resistant starch on the cooking quality of

yam (*Dioscorea* spp.) and cassava (*Manihot esculenta*) based paste products. *Int. J.*

Food Sci. Nutr. **64**:484–493.

Leah, R., Kigel, J., Svendsen, I., and Mundy, J. (1995). Biochemical and Molecular

Characterization of a Barley Seed β-Glucosidase. *J. Biol. Chem.* **270**:15789–15797.

Lebot, V., Trilles, B., Noyer, J. L., and Modesto, J. Genetic relationships between Dioscorea

alata L. cultivars.

Leng, Y., Yang, Y., Ren, D., Huang, L., Dai, L., Wang, Y., Chen, L., Tu, Z., Gao, Y., Li, X.,

et al. (2017). A Rice *PECTATE LYASE-LIKE* Gene Is Required for Plant Growth and

Leaf Senescence. *Plant Physiol.* **174**:1151–1166.

Lex, A., Gehlenborg, N., Strobelt, H., Vuillemot, R., and Pfister, H. (2014). UpSet:

Visualization of Intersecting Sets. *IEEE Trans. Vis. Comput. Graph.* **20**:1983–1992.

Li, X., Shin, S., Heinen, S., Dill-Macky, R., Berthiller, F., Nersesian, N., Clemente, T.,

McCormick, S., and Muehlbauer, G. J. (2015). Transgenic Wheat Expressing a Barley

UDP-Glucosyltransferase Detoxifies Deoxynivalenol and Provides High Levels of

Resistance to *Fusarium graminearum*. *Mol. Plant-Microbe Interactions®* **28**:1237–1246.

Li, C., Wu, H.-M., and Cheung, A. Y. (2016). FERONIA and Her Pals: Functions and

Mechanisms. *Plant Physiol.* **171**:2379–2392.

Li, X.J., Zhang, J.Q., Wu, Z.C., Lai, B., Huang, X.M., Qin, Y.H., Wang, H.C., Hu, G.B.

(2016). Functional characterization of a glucosyltransferase gene, LcUFGT1, involved in

the formation of cyanidin glucoside in the pericarp of *Litchi chinensis*. *Physiol. Plant.*

156:139-149.

Liao, L., Zhang, W., Zhang, B., Fang, T., Wang, X.-F., Cai, Y., Ogutu, C., Gao, L., Chen,

G., Nie, X., et al. (2021). Unraveling a genetic roadmap for improved taste in the

domesticated apple. *Mol. Plant* **14**:1454–1471.

Liu, C.; Yu, Q.; Li, Z.; Jin, X.; Xing, W. (2020). Metabolic and transcriptomic analysis related
to flavonoid biosynthesis during the colour formation of *Michelia crassipes* tepal. *Plant*
Physiology and Biochemistry **155**: 938-951.

Lou, H., Tucker, M. R., Shirley, N. J., Lahnstein, J., Yang, X., Ma, C., Schwerdt, J., Fusi,
R., Burton, R. A., Band, L. R., et al. (2022). The *cellulose synthase-like F3* (*CslF3*)
gene mediates cell wall polysaccharide synthesis and affects root growth and
differentiation in barley. *Plant J.* **110**:1681–1699.

Marks, R. A., Hotaling, S., Frandsen, P. B., and VanBuren, R. (2021). Representation and
participation across 20 years of plant genome sequencing. *Nat. Plants* **7**:1571–1578.

Meyer, R. S., and Purugganan, M. D. (2013). Evolution of crop species: genetics of
domestication and diversification. *Nat. Rev. Genet.* **14**:840–852.

Miao, H., Sun, P., Liu, Q., Jia, C., Liu, J., Hu, W., Jin, Z., and Xu, B. (2017). Soluble Starch
Synthase III-1 in Amylopectin Metabolism of Banana Fruit: Characterization,
Expression, Enzyme Activity, and Functional Analyses. *Front. Plant Sci.* **8**.

Mota, A. P. Z., Fernandez, D., Arraes, F. B. M., Petitot, A.-S., de Melo, B. P., de Sa, M. E.
L., Grynberg, P., Saraiva, M. A. P., Guimaraes, P. M., Brasileiro, A. C. M., et al.
(2020). Evolutionarily conserved plant genes responsive to root-knot nematodes
identified by comparative genomics. *Mol. Genet. Genomics* **295**:1063–1078.

Mu, W., Hassanin, H. A. M., Zhou, L., and Jiang, B. (2018). Chemistry Behind Rare Sugars
and Bioprocessing. *J. Agric. Food Chem.* **66**:13343–13345.

Muñoz, F. J., Baroja-Fernández, E., Morán-Zorzano, M. T., Alonso-Casajús, N., and
Pozueta-Romero, J. (2006). Cloning, Expression and Characterization of a Nudix
Hydrolase that Catalyzes the Hydrolytic Breakdown of ADP-glucose Linked to Starch
Biosynthesis in Arabidopsis thaliana. *Plant Cell Physiol.* **47**:926–934.

Ngolong Ngea, G. L., Guillon, F., Essia Ngang, J. J., Bonnin, E., Bouchet, B., and Saulnier,
L. (2016). Modification of cell wall polysaccharides during retting of cassava roots. *Food*
Chem. **213**:402–409.

Orkwor, G. C., Asiedu, R., and Ekanayake, I. (1998). *Food yams: Advances in research.*

Pont, C., Murat, F., Confolent, C., Balzergue, S., and Salse, J. (2011). RNA-seq in grain
unveils fate of neo- and paleopolyploidization events in bread wheat (Triticum aestivum
L.). *Genome Biol.* **12**:R119.

Prüfer, K., Muetzel, B., Do, H.-H., Weiss, G., Khaitovich, P., Rahm, E., Pääbo, S.,
Lachmann, M., and Enard, W. (2007). FUNC: a package for detecting significant
associations between gene sets and ontological annotations. *BMC Bioinformatics* **8**:41.

Quinlan, A. R., and Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* **26**:841–842.

Richmond, T. A., and Somerville, C. R. (2000). The Cellulose Synthase Superfamily. *Plant*
Physiol. **124**:495–498.

Ross, H. A., Morris, W. L., Ducreux, L. J. M., Hancock, R. D., Verrall, S. R., Morris, J. A.,
Tucker, G. A., Stewart, D., Hedley, P. E., McDougall, G. J., et al. (2011). Pectin
engineering to modify product quality in potato: Pectin engineering in potato. *Plant*
Biotechnol. J. **9**:848–856.

Sampedro, J., Valdivia, E. R., Fraga, P., Iglesias, N., Revilla, G., and Zarra, I. (2017).
Soluble and Membrane-Bound β-Glucosidases Are Involved in Trimming the Xyloglucan
Backbone. *Plant Physiol.* **173**:1017–1030.

Sharif, B. M., Burgarella, C., Cormier, F., Mournet, P., Causse, S., Van, K. N., Kaoh, J.,
Rajaonah, M. T., Lakshan, S. R., Waki, J., et al. (2020). Genome-wide genotyping
elucidates the geographical diversification and dispersal of the polyploid and clonally
propagated yam (*Dioscorea alata*). *Ann. Bot.* **126**:1029–1038.

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., and Zdobnov, E. M.
(2015). BUSCO: assessing genome assembly and annotation completeness with single-
copy orthologs. *Bioinformatics* **31**:3210–3212.

Singh, K., Rani, A., Kumar, S., Sood, P., Mahajan, M., Yadav, S. K., Singh, B., and Ahuja,
P. S. (2008). An early gene of the flavonoid pathway, flavanone 3-hydroxylase, exhibits a
positive relationship with the concentration of catechins in tea (*Camellia sinensis*). *Tree*
Physiol. **28**:1349–1356.

Siqueira, M. V., Dequigiovanni, G., Corazon-Guivin, M. A., Feltran, J. C., and Veasey, E.
A. (2012). DNA fingerprinting of water yam (*Dioscorea alata*) cultivars in Brazil based
on microsatellite markers. *Hortic. Bras.* **30**:653–659.

Soliman, A., Ayele, B. T., and Daayf, F. (2014). Biochemical and Molecular Characterization
of Barley Plastidial ADP-Glucose Transporter (HvBT1). *PLoS ONE* **9**:e98524.

Taylor, N. G. (2008). Cellulose biosynthesis and deposition in higher plants. *New Phytol.*
178:239–252.

Tetlow, I. J. (2011). Starch biosynthesis in developing seeds. *Seed Sci. Res.* **21**:5–32.

Van Harsselaar, J. K., Lorenz, J., Senning, M., Sonnewald, U., and Sonnewald, S. (2017).
Genome-wide analysis of starch metabolism genes in potato (*Solanum tuberosum* L.).
BMC Genomics **18**:37.

Watcharatpong, P., Kaga, A., Chen, X., and Somta, P. (2020) Narrowing Down a Major QTL
Region Conferring Pod Fiber Contents in Yardlong Bean (*Vigna unguiculata*), a
Vegetable Cowpea. *Genes*, **11**, 363.

Wiberley-Bradford, A. E., Busse, J. S., Jiang, J., and Bethke, P. C. (2014). Sugar
metabolism, chip colour, invertase activity, and gene expression during long-term cold
storage of potato (*Solanum tuberosum*) tubers from wild-type and vacuolar invertase
silencing lines of Katahdin. *BMC Res. Notes* **7**:801.

Wu, Z. G., Li, X. X., Lin, X. C., Jiang, W., Tao, Z. M., Mantri, N., Fan, C. Y., and Bao, X.
Q. (2014). Genetic diversity analysis of yams (Dioscorea spp.) cultivated in China using
ISSR and SRAP markers. *Genet. Resour. Crop Evol.* **61**:639–650.

Wu, Z.-G., Jiang, W., Mantri, N., Bao, X.-Q., Chen, S.-L., and Tao, Z.-M. (2015).
Transciptome analysis reveals flavonoid biosynthesis regulation and simple sequence
repeats in yam (*Dioscorea alata* L.) tubers. *BMC Genomics* **16**:346.

Xia, E., Tong, W., Hou, Y., An, Y., Chen, L., Wu, Q., Liu, Y., Yu, J., Li, F., Li, R., et al.
(2020). The Reference Genome of Tea Plant and Resequencing of 81 Diverse Accessions
Provide Insights into Its Genome Evolution and Adaptation. *Mol. Plant* **13**:1013–1026.

Xiao, N., Pan, C., Li, Y., Wu, Y., Cai, Y., Lu, Y., Wang, R., Yu, L., Shi, W., Kang, H., et al.
(2021). Genomic insight into balancing high yield, good quality, and blast resistance of
japonica rice. *Genome Biol.* **22**:283.

Zhang, J., and Sun, X. (2021). Recent advances in polyphenol oxidase-mediated plant stress
responses. *Phytochemistry* **181**:112588.

Zhou, Y.-X., Chen, Y.-X., Tao, X., Cheng, X.-J., and Wang, H.-Y. (2016). Isolation and
characterization of cDNAs and genomic DNAs encoding ADP-glucose
pyrophosphorylase large and small subunits from sweet potato. *Mol. Genet. Genomics*
291:609–620.

Zuker, A., Tzfira, T., Ben-Meir, H., Ovadis, M., Shklarman, E., Forkmann, G., Martens, S.,
Neta-Sharir, I., Weiss, D., and Vainstein, A. (2002) Modification of flower colour and
fragrance by antisense suppression of the flavanone 3-hydroxylase gene. *Molecular*
Breeding **9** :33–41

Figure Legends

Figure 1: Population structure of *Dioscorea alata*. A: Geographical origin of the 125 *Dioscorea alata* accessions. B: Circos plot of the 20 chromosomes of *D. alata*, the circles represent from outer to the inner circle: chromosomes, gene density, localization of genes from pentose and glucuronate interconversions, genes from starch and sucrose metabolism and genes from flavonoid biosynthesis. C: Principal component analysis depicting the 107 diploid genotypes of *D. alata*. Accessions are coloured according to their assignment to the three genetic clusters after Admixture analysis. The threshold to assign a genotype to a cluster is set at 75 %. D: Admixture barplot showing the distribution of the $K = 2$ to $K = 4$ genetic clusters.

Figure 2: Selection signature found from the comparison of clusters. A: Manhattan plot of Fst analysis of cluster A *versus* B, each colour represents one chromosome. B: *Fst* analysis of clusters A *versus* C. C: *Fst* analysis of clusters B *versus* C. D: Venn diagram of the common genes found among the three clusters, and the number of common outlier regions. E: The gene ontology enrichment of genes found in outlier regions against the whole set of genes.

Figure 3: Orthology analysis of 45 plant species. A: Number of genes found in the keyword analysis per pathway and the EC number which they were associated with (light grey). Number of genes found in the orthology analysis, number of EC numbers and number of orthologous groups found (dark grey). B: UpSetR of the comparison of all the monocotyledon plants used in orthology analysis. Each dot represents a connection, and the bar represents the number of orthologous groups found for each association. The red dots show the species from the Dioscoreales family, and the red numbers represent the exclusive associations of Dioscoreales. C: UpSetR of the five Dioscoreales species. D: Intersection between the genes found in the *Fst* analysis and in the pathways.

Figure 4: Candidate genes associations and gene expression profile. Selected genes and their expression profile, and allelic effect on colour and texture traits. A: gene structures (*Dioal.19G083800, Dioal.04G073400* and *Dioal.04G130600*) with positions of the significant SNPs, their alleles and resulting amino acid changes; S4_20715506 (C/T) effect on B: gumminess; C: total area; S10_17029453 (G/C) effect on D: brown index (BI); E: Transcriptome based expression of *Dioal.19G083800* between two genotypes with G and C alleles (FPKM= fragments per kilobase of transcript per million fragments mapped); FPKM values was extracted from three biological replicates. S4_16031453 (C/T) effect on F: Hue index leading purple colour change displayed by the two representative tubers. Means were separated by two-tailed *t*-test at 0.05 probability.

Trait	SNP_ID	Chr	Pos (bp)	Allele Major	Allele Minor	Consequence Type	Genomic Location	Effect (%)	Gene Model	Predicted Annotation	Pathway	QTN
Springiness	S1_10223458	1	10223458	C	T	Stop gain	p.Trp287*	0.38	Dioal.01G024400.1	Leucine-rich repeat	Pectin	qSp1.1
	S1_10224508	1	10224508	G	A	Stop gain	p.Gln104*	0.36	Dioal.01G024400.1	Leucine-rich repeat	Pectin	qSp1.2
	S2_19082482	2	19082482	C	T	Stop gain	p.Gln1032*	-0.05	Dioal.02G062700.1	Protein kinase	Pectin	qSp2.1
	S2_19095245	2	19095245	C	G	Stop gain	p.Tyr578*	-0.04	Dioal.02G062800.1	Protein kinase	Pectin	qSp2.2
	S13_11781422	13	11781422	A	G	splice_acceptor_variant&intron_variant	c.63-2A>G	-0.37	Dioal.13G053400.1	Xylose isomerase	Pectin	qSp13.1
	S20_14416887	20	14416887	G	T	Stop gain	p.Gly381*	-0.46	Dioal.20G055400.1	Malectin-like	Pectin	qS20.1
Gumminess	S4_20715506	4	20715506	C	T	Start Loss	p.Met1?	-3.97	Dioal.04G130600.1	Pectin methylesterase	Pectin	qG4.1
	S7_6378890	7	6378890	G	A	splice_donor_variant&intron_variant	c.-109+1G>A	-9.23	Dioal.07G051400.1	Alpha amylase	Starch	qG7.1
	S13_11783621	13	11783621	G	T	Stop gain	p.Glu208*	-2.61	Dioal.13G053400.1	Xylose isomerase	Pectin	qG13.1
	S13_11784207	13	11784207	A	G	splice_acceptor_variant&intron_variant	c.723-1G>A	23.60	Dioal.13G053400.1	Xylose isomerase	Pectin	qG13.2
	S19_24800556	19	24800556	G	A	splice_donor_variant&intron_variant	c.537+2C>T	4.87	Dioal.19G187400.1	Pectinesterase	Pectin	qG19.2
Total area	S2_18814720	2	18814720	T	A	Stop gain	p.Tyr289*	21.88	Dioal.02G061200.1	Protein kinase	Pectin	qTa2.1
	S4_20715506	4	20715506	C	T	Start Loss	p.Met1?	-27.90	Dioal.04G130600.1	Pectin methylesterase	Pectin	qTa4.1
	S7_16450864	7	16450864	T	T	Stop gain	p.Tyr228*	-29.42	Dioal.07G067100.1	4-Alpha-glucanotransferase	Starch	qTa7.1
Hardness	S13_28669451	13	28669451	C	G	splice_donor_variant&intron_variant	c.-837+1G>C	62.22	Dioal.13G093500.1	Beta-glucosidase	Starch	qH13.2
	S20_14334067	20	14334067	G	T	Stop gain	p.Gly6*	-12.80	Dioal.20G055000.1	Malectin-like	Pectin	qH20.1
	S20_14364743	20	14364743	A	C	Stop loss	p.Ter592Serext*?	10.75	Dioal.20G055100.1	Malectin-like	Pectin	qH20.2
Starch Content	S10_2314181	10	2314181	G	T	Stop gain	p.Gly35*	-14.75	Dioal.10G014800.1	1,3-beta-Glucan synthase	Starch	qSC10.1
	S15_23348104	15	23348104	T	C	splice_donor_variant&intron_variant	c.1029+2C>T	4.99	Dioal.15G119500.1	beta-Amylase	Starch	qSC15.2
	S17_19982533	17	19982533	T	C	splice_acceptor_variant&intron_variant	c.811-2A>G	-9.70	Dioal.17G112400.1	Glucose-1-phosphate adenylyltransferase	Starch	qSC17.1
Brown Index	S19_17029453	19	17029453	G	C	Stop gain	p.Tyr339*	34.69	Dioal.19G083800.1	Flavanone 3-dioxygenase	Flavonoid	qYI19.1
HUE Index	S4_16031453	4	16031453	C	T	Stop gain	p.Trp272*	50.28	Dioal.04G073400.1	UDP-glucosyltransferase protein	Flavonoid	qHI4.1