Overcoming donor variability and risks associated with fecal microbiota transplants through bacteriophage-mediated treatments

Torben Sølbeck Rasmussen[1*], Sarah Forster[1], Sabina Birgitte Larsen[1], Alexandra Von Münchow[2], Kaare Dyekær Tranæs[1], Anders Brunse[3], Josue Leonardo Castro Mejia[1], Signe Adamberg[4], Axel Kornerup Hansen[2], Kaarel Adamberg[4,5], Camilla Hartmann Friis Hansen[2], Dennis Sandris Nielsen[1*]

[1] Section of Microbiology and Fermentation, Dept. of Food Science, University of Copenhagen, Frederiksberg, Denmark

[2] Section of Experimental Animal Models, Dept. of Veterinary and Animal Sciences, University of Copenhagen, Frederiksberg, Denmark

[3] Section of Comparative Pediatrics and Nutrition, Department of Veterinary and Animal Sciences, University of Copenhagen, Frederiksberg, Denmark

[4] Department of Chemistry and Biotechnology, Tallinn University of Technology, Tallinn, Estonia

[5] Center of Food and Fermentation Technologies, Tallinn, Estonia

*Corresponding authors. Email: dn@food.ku.dk / torben@food.ku.dk

Summary: Fecal viromes depleted of enveloped viruses efficiently treat *Clostridioides difficile*-associated diarrhea in a murine model.

ABSTRACT

Fecal microbiota transplantation (FMT) and fecal virome transplantation (FVT, sterile filtrated donor feces) have proven effective in treating recurrent *Clostridioides difficile* infection (rCDI). However, variability in donor materials and costly screening, coupled with concerns (motivated by case reports) over pathogen transfer with FMT or FVT, hinders their wider clinical application in treating less acute diseases. Although FVT presents safety advantages over FMT by transferring no bacteria, the presence of eukaryotic viruses poses a risk of harmful infections. To overcome these challenges, we developed methods to broaden FVT's clinical application while maintaining efficacy and increasing safety. Namely: 1) chemostat-fermentation to reproduce the bacteriophage FVT component and remove eukaryotic viruses (FVT-ChP); 2) solvent-detergent treatment to inactivate enveloped viruses (FVT-SDT); and 3) pyronin-Y treatment to block RNA-virus replication (FVT-PyT). We evaluated the efficacy of these processed FVTs using a CDI mouse model and compared them with untreated FVT (FVT-UnT), FMT, and saline control. FVT-UnT, FVT-SDT, and FVT-ChP significantly reduced *C. difficile* colonization, limited CDI symptoms and reduced the incidence of mice reaching humane endpoints compared with the saline control. This proof-of-concept opens the possibility of developing safer and more effective phage-mediated therapies for a broad range of gut-related diseases and addresses the reproducibility issues associated with fecal donor material.

INTRODUCTION

During the past decade, it has become evident that various diseases are associated with gut microbiome (GM) dysbiosis, including recurrent *Clostridioides difficile* infection (rCDI)[1–3]. Fecal microbiota transplantation (FMT) from a healthy donor to rCDI patients has proven highly effective in curing the disease, with a success rate exceeding 90%[4]. However, FMT faces challenges such as expensive and labor-intensive donor screening[5,6], donor variability in term of treatment efficacy and reproducibility[6–8], and safety concerns since no screening methods can definitively exclude the transfer of agents that could cause adverse effects from the donor[3]. The importance of the latter was highlighted in June 2019 when two patients in the United States experienced severe bacterial infections after FMT, resulting in one fatality[9]. Subsequent safety alerts from the U.S. Food & Drug Administration have warned against potential severe adverse effects associated with the transfer of pathogenic microorganisms during FMT[10,11].

Interestingly, Ott et al.[3] and Kao et al.[12] successfully treated rCDI patients using sterile-filtered donor feces (containing viruses but no intact bacteria), a method often referred to as fecal virome transplantation (FVT). The efficacy of FVT was comparable to other clinical studies using FMT (containing bacteria, etc.), indicating that the gut virome plays a significant role in treating rCDI with FMT[3,12]. The gut virome is predominantly composed of bacteriophages (in short phages), which are host-specific bacterial viruses, but also include eukaryotic and archaeal viruses[13]. The concept of applying FVT from a healthy phenotype to a symptomatic phenotype has been further investigated in preclinical studies. For instance, Draper et al. demonstrated that FVT influenced the composition of the murine GM following initial perturbation with antibiotics[14]. Additionally, FVT treatment from lean donors alleviated symptoms of metabolic syndrome in two independent diet-induced obesity murine models[15,16], and FVT using term piglet donors prevented development of necrotizing enterocolitis in a preterm piglet model[17]. FVT has the advantage over FMT in that it transfers little to no bacteria. However, the sterile filtration process commonly used in FVT (typically using 0.45 µm filters) does not eliminate the risk of transferring eukaryotic viruses. While it is possible to screen donor feces for known pathogenic viruses, recent studies have revealed that the human gastrointestinal tract harbors hundreds of eukaryotic viruses with unknown functions[13,18,19]. Most of these viruses are likely harmless to the human host. Nevertheless, it cannot be ruled out that they may contribute to later disease development, as exemplified by the human papillomavirus, which turned out as a significant risk factor for cervical cancer years after

infection[20]. Therefore, there are good reasons to minimize the transfer of eukaryotic viruses when

applying FVT to alleviate conditions associated with gut microbiome dysbiosis, particularly when

treating individuals with compromised immune systems. In contrast, phages do not actively infect

and replicate in eukaryotic cells and are believed to be key players in the successful treatment of

gut-related diseases through FMT/FVT[3,15,17,21], and have recently been suggested to harbor immune

modulating effects[22,23]. Phage therapy using defined phage cocktails has been applied for almost a

century[24] and is generally considered safe since severe adverse effects are rarely reported[25–28].

FMT and FVT hold the potential to revolutionize treatments for numerous gut-related diseases,

but their widespread use is unlikely due to safety concerns and donor variability. Therefore, our

objective was to develop methodologies that mitigate the risk of viral infections associated with

FVT by removing or inactivating the eukaryotic viral component in fecal donor material.

Additionally, we aimed to address the challenge of donor variability by ensuring reproducible

production of enteric donor phage communities[29]. The efficacy of these processed fecal viromes

was evaluated using a murine CDI model. This proof-of-concept study represents an important

first step towards developing safer and more consistent therapeutic approaches that can effectively

target a wide range of gut-related diseases that are less acute than rCDI, and potentially replace

FMT with phage-mediated therapies.

RESULTS

To generate fecal viromes free of eukaryotic viruses, we developed methodologies that exploit the

fundamental differences in characteristics between eukaryotic viruses and phages. The majority of

eukaryotic viruses are enveloped RNA viruses[30–32] and rely on eukaryotic hosts for replication. In

contrast, the majority of phages are non-enveloped DNA viruses[32,33] that require bacterial hosts for

replication. We employed a solvent/detergent method to inactivate enveloped viruses (FVT-SDT),

utilized pyronin Y to inhibit the replication of RNA viruses (FVT-PyT), and created a chemostat-

propagated virome (FVT-ChP) to dilute and remove the majority of eukaryotic viruses[29]. These

differently processed fecal viromes were evaluated in a murine CDI model and compared with a

saline solution (sham treatment), fecal microbiota transplantation (FMT, previously shown to be

efficient in treating CDI in the used preclinical model[34,35]), and untreated donor-filtrated feces

(FVT-UnT). The content of all treatments originated from the same intestinal donor material.

Effective inactivation of enveloped phages through solvent/detergent treatment and replication inhibition of RNA phages using pyronin Y

The applicability of solvent/detergent and pyronin Y treatments to inactivate eukaryotic viruses while preserving phage activity was assessed using phages with distinct characteristics. These characteristics included enveloped (phi6) versus non-enveloped (phiX174, T4, and C2) phage structures and ss/dsDNA (phiX174, T4, C2) versus ss/dsRNA (MS2, phi6) genomes (Fig. 1A & 1B). The solvent/detergent treatment completely inactivated the enveloped phage phi6, as indicated by the plaque-forming units (PFU)/mL dropping to below detection limit, while the non-enveloped phages phiX174 and T4 remained unaffected. Phage C2, on the other hand, experienced a minor decrease of 1 $\log_{10}$ in PFU/mL (Fig. 1A). Pyronin Y was used to inhibit the replication of viruses carrying RNA genomes. After exploring various combinations of pyronin Y concentrations, temperatures, and incubation times, an overnight incubation at 40°C with 100 µM pyronin Y was chosen. This treatment resulted in a 5 $\log_{10}$ reduction in PFU/mL for the ssRNA phage MS2 and complete inactivation of the dsRNA phage phi6. Phi6 demonstrated temperature sensitivity as incubation at 40°C alone was sufficient to inactivate this enveloped phage, while incubation temperature at 20°C along with 100 µM pyronin Y led to a reduction of 4 $\log_{10}$ PFU/mL of phage phi6. Unfortunately, the pyronin Y treatment at 40°C also affected the plaque-forming ability of phages C2 (dsDNA), T4 (dsDNA), and phiX174 (ssDNA), resulting in reductions of 1, 2.5, and 5 $\log_{10}$ PFU/mL, respectively (Fig. 1B).

By utilizing chemostat propagation, eukaryotic viruses were depleted from fecal viromes through dilution, leading to a reduction from an average of 20 to 1 eukaryotic virus that was taxonomically classified[29].

Sustained treatment efficacy of fecal viromes following inactivation of enveloped viruses

The treatment efficacy, specifically preventing the mice from reaching the humane endpoint, was assessed in a CDI model using C57BL/6J mice. The survival probability rate was evaluated using a Kaplan-Meier estimate (Fig. 2A) and compared to the mice treated with saline (2/7 mice survived). The analysis revealed a significantly improved survival rate (8/8 mice, p = 0.004) for mice treated with FVT-SDT. Additionally, FVT-UnT showed a tendency towards an improved survival rate (5/7 mice, p = 0.067) compared to the saline control. On the other hand, the FVT-PyT treated mice, and unexpectedly, the FMT treatment, showed no improvement in survival rate (1/8 and 3/8 mice survived, respectively). The effect of FVT-ChP on the survival rate showed

numerical, but non-significant, improvement (5/8 mice, p = 0.235). The pathological score of the cecum tissue supported the observed survival rate, as FVT-UnT, FVT-ChP, and FVT-SDT treated mice showed a significant decrease (p < 0.05) in pathological score compared to the saline control (Fig. 2B, Fig. S1). The concentrations of 10 pro- and anti-inflammatory cytokines were also measured in the cecum tissue to further validate the cecum histology score and the qualitative health evaluation (Fig. S2). The cytokine profiles generally reflected whether the mice survived (with low or no inflammatory response when measured at study termination) or were euthanized (with a high inflammatory response) due to *C. difficile* infection (Fig. 2C-L). Overall, the FVT-SDT treatment, which involved the inactivation of enveloped viruses, effectively prevented severe *C. difficile* infection, and increased the chances of survival. The average pathological score at 6.7 of the saline treatment was in line with the original published CDI mouse model by Chen et al. that reported a pathological score at 7.0 for *C. difficile* infected mice, compared to mice not infected with *C. difficile* that showed a score at 1.3[36]. This supported that the pathological scores of 2.5 - 3.0 observed in the FVT-UnT, FVT-ChP, and FVT-SDT treated mice corresponded to limited symptoms of *C. difficile* infection.

Successful treatments impede *C. difficile* colonization and subsequent disease development
C. difficile abundance in feces was quantified using qPCR (Fig. 3A-3C). No *C. difficile* was detected before inoculation with *C. difficile* (Fig. 3A). The FVT-SDT treated mice exhibited an average of 2 $\log_{10}$ lower *C. difficile* abundance (gene copies per gram feces) compared to the saline treated mice before the 2nd treatment, and the FVT-UnT and -ChP treatments resulted in a 1.5 $\log_{10}$ lower abundance. This suggested that these three treatments effectively impeded *C. difficile* colonization in the gut. In contrast, the FMT and FVT-PyT treated mice had similar *C. difficile* abundance as the saline treated group. A possible clearance of *C. difficile* colonization was observed at termination, as 7/8 FVT-SDT treated mice tested negative for *C. difficile*, while all other treatments showed persistent *C. difficile* presence (Fig. 3C).
The levels of the *C. difficile*-associated toxin A/B were measured using an ELISA-based assay, which showed similar patterns as the qPCR data (Fig. 3D-3F). The toxin A/B levels in the FVT-SDT treated mice were significantly lower than those in the saline group (p < 0.05), and only 2 mice in the FVT-SDT group exhibited detectable toxin A/B levels before the 2nd treatment. At termination, toxin A/B could not be detected in any of the FVT-SDT treated mice (Fig. 3F). The decrease in *C. difficile* abundance (Fig. 3A-3C) and the low levels of toxin A/B (Fig. 3D-3F) in

the FVT-SDT, -UnT, and -ChP treated mice support the phenotypic data of survival rate, histology,

and cytokine profile (Fig. 2).

Fecal virome treated with solvent/detergent supports recovery of the bacterial community in

a dysbiotic GM

There were no initial differences in bacterial and viral GM profiles at both baseline (before

antibiotics) and before *C. difficile* infection between the mice that were later either euthanized or

survived the *C. difficile* infection (Fig. S3, S4, S5, & S6). The antibiotic intake through the drinking

water was similar across the cages (Table S1). The bacterial taxonomic profile of the mice that

were euthanized due to reaching the humane endpoint was consistently dominated by

Escherichia/Shigella, *Enterococcus*, *Clostridioides*, *Bacteroides*, and *Parabacteroides* (Fig. 4C,

4D, & Fig. S7), and the bacterial composition (Bray-Curtis dissimilarity) and diversity (Shannon

diversity index) were significantly different (q < 0.003) from the mice that had survived the

infection (Fig. 4A & 4B). In the FMT treated mice that survived the infection, bacterial taxa were

observed that overlapped with those present in the FMT inoculum, suggesting potential

colonization of these bacteria in the gut of the FMT treated mice (Fig. 4C & 4D). Two mice treated

with FVT-PyT were colonized with 5-20% relative abundance of *Salmonella* spp., which may have

contributed to increased disease severity for these mice. The bacterial taxonomic profile of the

mice that had survived the infection was dominated by lactobacilli, *Bacteroides*, *Lachnospiraceae*,

Bifidobacterium, *Akkermansia*, *Porphyromonadaceaea*, *Bacteroidales*, and *Turibacter*, which

were also the dominant taxa in the mice at baseline (Fig. 4C, 4D, & Fig. S7A), suggesting partial

restoration of the GM profile in mice that survived the *C. difficile* infection.

The FVT-SDT treatment improved the chances of restoring the GM composition after antibiotic

treatment, thereby impacting the mice's ability to fight off the *C. difficile* infection. The viral

composition and diversity also differed significantly (p < 0.005) between the euthanized and

surviving mice (Fig. 5A & 5B). Euthanized mice showed a significant (p < 0.05) increase in the

relative abundance of the viral order *Petitvirales* and phages predicted to have *Sutterella* as host.

In contrast, mice that survived the infection until termination were characterized by a significant

increase (p < 0.05) in the viral order *Caudovirales*, as well as phages predicted to target the genera

Parabacteroides, *Bacteroides*, *Prevotella*, *Odorifactor*, and *Escherichia* (Fig. S7B & S7C). The

eukaryotic viral profile was similar among the different FVT inoculates (Fig. S8) and constituted

0.5-2.5% of the total relative abundance (Fig. S9). However, it is important to note that the
taxonomic resolution of eukaryotic viruses was insufficient to differentiate between treatments or
outcomes in relation to the relative abundance of eukaryotic viruses (Fig. S8). It should also be
emphasized that inactivation by dissolving the viral envelope or inhibiting replication does not
exclude the detection of viruses through sequencing.

DISCUSSION

Here, we have developed methodologies to address both the risks associated with transferring
pathogenic microorganisms and the variability of donors in FMT/FVT for gut-related diseases.
The preservation of treatment efficacy of the modified FVTs against *C. difficile* infections (CDI)[3,12]
was assessed using a CDI mouse model. The inactivation of enveloped viruses through
solvent/detergent treatment emerged as the superior method to modify fecal viromes while
preserving treatment efficacy against CDI, as all FVT-SDT treated mice survived without reaching
the humane endpoint.

The systemic inflammatory response observed during CDI is driven by *C. difficile* toxins that
increase gut tissue permeability[37,38], making the gut more susceptible to other microbial
infections[39,40]. Therefore, transferring untreated fecal donor viromes (including intact viruses) may
result in additional inflammation due to microbes translocating through the damaged intestinal
tissue. Enveloped viruses, including those belonging to the *Herpesviridae* family, which includes
major mammalian pathogens[41] were detected in both the FVT-UnT and -SDT inoculum (Fig. S8).
Enveloped *Herpesviridae* were, however, expected to be inactivated by the solvent/detergent
which could explain the mild symptoms and high survival rate associated with the FVT-SDT-
treated mice compared to the untreated equivalent of FVT-UnT. Treating fecal viromes with
solvent/detergent prior to transfer to patients may be particularly relevant for gut-associated
diseases due to increased gut tissue permeability[39,40].

FMT is generally linked to a treatment efficacy above 90% in preclinical[34,35] and clinical CDI
studies[3]. Surprisingly, the survival rate of FMT-treated mice was similar to that of the saline control
group. Analysis of the inoculum by 16S rRNA gene amplicon sequencing revealed that the FMT
inoculum contained approximately 20% of *Clostridium sensu stricto* spp., which has been
associated with *C. difficile*-positive calves[42] and to diarrhea in pigs[43]. The relatively high abundance
of *Clostridium sensu stricto* spp. in the FMT inoculum may have counteracted the curative effects
typically associated with FMT[3,34,35]. While the FMT and FVT inocula originated from the same

donor material, the removal of bacteria during FVT processing may explain the higher survival
rates observed with FVT-UnT, -ChP, and -SDT compared to FMT. This also suggests that even
unsuitable fecal donor material for FMT could potentially be suitable for FVT. The lowest survival
rate was observed with the RNA-targeting compound pyronin Y. During the initial evaluation of
pyronin Y's ability to inactivate RNA phages, it became evident that DNA phages were also
affected. The pyronin Y treatment may have caused a reduction in phage activity, which could
have impacted the efficacy of the treatment. This finding aligns with other studies that emphasize
the potential role of phages in restoring GM balance following FMT or FVT[3,14,15,17,21,44–46].
The ability of different FVT treatments to reduce the abundance of the infectious *C. difficile* strain
was associated with the survival rate, histology, and cytokine profiles. Mice treated with FVT-
UnT, -ChP, and -SDT showed a significant decrease in *C. difficile* abundance compared to those
treated with saline, FMT, and FVT-PyT. This suggests that when fewer *C. difficile* cells are present
to produce toxins associated with CDI etiology[37,38], fewer cells will reach the stationary growth
phase where toxin production is highest[47]. We believe that the phages transferred along with the
FVT play a role in allowing commensal bacteria associated with a healthy state to compete with
the infectious *C. difficile* strain, as well as commensal bacteria that can act as opportunistic
pathogens. This is supported by a significant ($p < 0.05$) increase in the relative abundance of
viruses predicted to infect *Escherichia*/*Shigella*, *Parabacteroides*, and *Bacteroides*, while these
same bacterial genera constituted a significantly smaller ($p < 0.05$) fraction of the bacterial relative
abundance when comparing the surviving mice with the euthanized mice.
243
Metavirome sequencing of fecal samples only revealed ssRNA viruses (phages) belonging to the
Leviviridae family, while the positive controls of both the dsRNA Phi6 and ssRNA MS2 phages
(titer ~10^6 PFU/mL) were detected in both mock communities and spiked fecal samples (Fig. S10).
This suggests that the metavirome library preparation was able to detect RNA viruses at
concentrations of at least ~10^6 viral particles/mL. The limited detection of enteric RNA viruses in
our study is consistent with other studies using similar techniques[48] and may be due to insufficient
viral databases[49], the "too clean" nature of specific pathogen-free (SPF) mice[50,51], or the
concentration of these viruses being below the detection limit (~10^6 viral particles/mL).
A major challenge of CDI is the risk of recurrent infections[52–54]. It is therefore interesting to note
that 7 out of 8 FVT-SDT treated mice showed a depletion of *C. difficile* at termination, indicating

a decrease in the risk of recurrent infections when treated with a solvent/detergent modified fecal
virome. Thus, the solvent/detergent treated fecal virome demonstrated superiority over other
treatments against CDI in our study. Interestingly, the treatment efficacy and decrease in CDI-
associated symptoms were also pronounced in mice treated with the chemostat propagated enteric
virome, that in a previous study showed high reproducibility of the viral component[29]. Also
Santiago-Rodriguez et al. have previously demonstrated how propagation of fecal inoculum in a
chemostat-fermentation holds the potential for reproducing the enteric viral component[55].

The inherent challenges of variability and reproducibility in fecal donor material exist for both
FMT and FVT[6–8]. When optimized, the chemostat fermentation-based approach has the potential
for higher reproducibility, standardization, large-scale production, increased removal of eukaryotic
viruses through further dilution, and relies on only a few effective donors. Therefore, it could be
argued that the solvent/detergent methodology, already approved by WHO as a safe procedure for
treating blood plasma[56], holds the potential to complement FMT in the treatment of CDI in the
short-term perspective. In the long-term perspective, a cost-effective, standardized, and
reproducible chemostat propagated enteric phageome for CDI treatment may also have tremendous
potential for treating other diseases associated with gut dysbiosis.

METHODS

Study design - *Clostridioides difficile* infection model

The *C. difficile* infection model was based on Chen et al. 2008[36]. Forty-eight female C57BL/6J (JAX) mice, 8 weeks old, were obtained from Charles River Laboratories (European distributor of JAX mice) and housed at the animal facilities of the Department of Experimental Medicine (AEM, University of Copenhagen, Denmark) in Innovive disposable IVC cages. The cages provided water, food (Altromin 1324 chow diet), bedding, cardboard housing, nesting material, felt pad, and biting stem. Upon arrival, the mice were ear tagged, randomly assigned to cages with 4 mice each, and acclimatized for one week (Fig. 6). An antibiotic mixture (kanamycin 0.4 mg/mL, gentamicin 0.035 mg/mL, colistin 850 U/mL, metronidazole 0.215 mg/mL, and vancomycin 0.045 mg/mL) was prepared in the drinking water and provided to the mice through the IVCs for 3 days, and the water was replaced with clean antibiotic-free drinking water for 2 days. Subsequently, the mice received an intraperitoneal injection of clindamycin (2 mg/mL) diluted in sterile 0.9% (w/v) NaCl water (based on the average body weight of the mice, around 20 g). Twenty-four hours later, the mice were orally inoculated with 1.21 x 10^4 CFU of *C. difficile* VPI 10463 (CCUG 19126) via gavage. The mice were then divided into six different treatment groups (n = 8): saline (control), FMT, FVT-UnT (untreated FVT), FVT-ChP (chemostat propagated virome), FVT-SDT (solvent/detergent treated FVT for inactivating enveloped viruses), FVT-PyT (FVT treated with pyronin Y for inactivation of RNA viruses). The respective treatments were administered orally in two doses of 0.15 mL each, at 18 hours and 72 hours after *C. difficile* inoculation (Fig. 6). One mouse from the saline treated group was euthanized immediately after oral gavage of *C. difficile*, as the culture was accidentally administered via the trachea, reducing this group to n = 7. Unfortunately, one mouse from the FVT-UnT group was euthanized 1 week after the 2nd FVT treatment due to malocclusions that had led to malnutrition, reducing this group to n = 7. Frequent (up to every 3rd hour – day and night) blinded monitoring of the animal health status was performed after *C. difficile* infection to determine if the mice had reached the humane endpoint, leading to immediate euthanization when necessary. The monitoring was supervised by the blinded veterinarian associated with the study (co-author AVM). The following qualitative parameters were used: physical activity level (i.e., decreased spontaneous or provoked activity), consistency of feces (watery or normal), body posture (hunching or normal), and whether their fur was kept clean or not. The mice were scored on a scale of 0-2: 0 (healthy), 1 (mild symptoms), and 2 (clear

symptoms). Mice with a score of 2 that showed no improvement in the above parameters during the subsequent checkup were euthanized. Fecal pellets were sampled whenever possible at different time points until the mice were euthanized (Fig. 6). At the time of euthanization, samples of the intestinal content from the cecum and colon were taken. A portion of the cecum tissue (55) was fixed in 10% neutral-buffered formalin (Sarstedt) for histological analysis and stored at room temperature. Another part of the cecum tissue, along with the intestinal content, was preserved for cytokine analysis and stored at -80°C. All procedures involving the handling of these animals were conducted in accordance with Directive 2010/63/EU and the Danish Animal Experimentation Act, under license ID: 2021-15-0201-00836.

Preparation of the *Clostridioides difficile* inoculum

Clostridioides difficile VPI 10463 (CCUG 19126), originating from a human infection, was used as infectious agent in the mouse CDI described by Chen et al.[36]. The bacteria were cultured in brain-heart-infusion supplement (BHIS) medium[57] with 0.02% (w/v) 1,4-dithiothreitol (Fisher Scientific) and 0.05% (w/v) L-cysteine (Fisher Scientific), grown at 37°C in Hungate tubes (SciQuip), and handled anaerobically as previously described[58]. For solid media 1.5% (w/v) agar (Fischer Scientific) was added. Optical density (OD_{600nm}) of bacterial cultures was measured with a Genesys30 Visible spectrophotometer (Fischer Scientific). *C. difficile* primarily synthesize its toxins during the stationary phase[47], thus the *C. difficile* culture in its early exponential phase was used as inoculum to minimize transfer of toxins. The CDI inoculum was prepared as follows: a single colony of *C. difficile* was transferred to a Hungate tube containing 10 mL BHIS medium and incubated overnight at 37°C. Then 150 µL of the *C. difficile* overnight culture was transferred to a new Hungate tube containing 10 mL BHIS medium, incubated for 3.5 hours and its OD_{600nm} was measured. A bacterial calibration curve (OD_{600nm} vs. CFU/mL) of *C. difficile* VPI 10463 was used to dilute the culture to the desired concentration (1×10^5 CFU/mL). This constituted the *C. difficile* inoculum. The cell concentration of the CDI inoculum was evaluated with CFU counts on BHIS agar plates (Fig. S11).

Propagation of host-phage pairs

The bacteria were grown in media and at a temperature suitable for each strain (Table S2). Five phages representing different characteristics (genome, size, and structure), along with their bacterial hosts, were included to assess the influence of pyronin Y and solvent/detergent treatment on phage activity; *Lactococcus* phage c2 (host: *Lactococcus lactic* MG1363), coliphage T4 (host:

Escherichia coli Luria)*,* coliphage phiX174 (host: *E. coli* PC 0886), coliphage MS2, (host: *E. coli*

W1485), and *Pseudomonas* phage phi6 (host: *Pseudomonas* sp. HER1102). Solid media were

supplemented with 1.5% (w/v) agar (Thermo Fisher Scientific) for plates and 0.7% (w/v) for soft

agar. An end concentration of 10 mM $MgCl_2$ and $CaCl_2$ was supplemented to media during phage

propagation. Plaque activity of phages was evaluated by spot testing where 100 µL bacterial

culture was mixed with 4 mL soft agar (temperature at 49°C), poured to an agar plate, and 10 µL

of a phage suspension (dilution series) was deposited on the surface of the solidified soft agar,

followed by incubation according to the specific bacterial strain (Table S2).

The animal origin and preparation of donor viromes and FMT

A total of 54 male C57BL/6N mice were purchased for the purpose of harvesting intestinal content

for downstream FVT/FMT applications. Upon arrival, the mice were five weeks old and obtained

from three vendors: 18 C57BL/6NTac mice from Taconic (Denmark), 18 C57BL/6NRj mice from

Janvier (France), and 18 C57BL/6NCrl mice from Charles River (Germany). The mice were

earmarked upon arrival and housed at the Section of Experimental Animal Models, University of

Copenhagen, Denmark, following previously described conditions[59]. They were provided with ad

libitum access to a low-fat diet (LF, Research Diets D12450J) for a period of 13 weeks until they

reached 18 weeks of age, which was the planned termination point. Unfortunately, malocclusions

resulted in malnutrition for two C57BL/6NRj mice, and they were euthanized before the intended

termination date. All mice were euthanized by cervical dislocation, and samples of intestinal

content from the cecum and colon were collected and suspended in 500 µL of autoclaved anoxic

PBS buffer (NaCl 137 mM, KCl 2.7 mM, Na_2HPO_4 10 mM, KH_2PO_4 1.8 mM). Subsequently, all

samples were stored at -80°C. In order to preserve the viability of strict anaerobic bacteria, 6 mice

from each vendor (a total of 18 mice) were sacrificed and immediately transferred to an anaerobic

chamber (Coy Laboratory) containing an atmosphere of approximately 93% N_2, 2% H_2, and 5%

CO_2, maintained at room temperature. The samples collected from these mice within the anaerobic

chamber were used for FMT and anaerobic chemostat cultivation to produce the FVT-ChP. The

intestinal content from the remaining 34 mice was sampled under aerobic conditions and used to

generate the fecal virome for downstream processing of the FVT-UnT, FVT-SDT, and FVT-PyT

treatments. A flow diagram illustrating the aforementioned processes is provided in Figure S12.

All procedures involving the handling of donor animals were conducted in accordance with

Directive 2010/63/EU and the Danish Animal Experimentation Act, under license ID: 2012-15-
2934-00256.

Untreated fecal virome (FVT-UnT)

For processing FVT solutions[15], thawed intestinal content from the cecum and colon was
suspended in 29 mL autoclaved SM buffer (100 mM NaCl, 8 mM $MgSO_4 \cdot 7H_2O$, 50 mM Tris-HCl
with pH 7.5), followed by homogenization in BagPage+ 100 mL filter bags (Interscience) with a
laboratory blender (Seward) at medium speed for 120 sec. The filtered and homogenized
suspension was subsequently centrifuged using an Allegra 25R centrifuge (Beckman Coulter) at
5,000 x g for 30 min at 4°C. The fecal supernatant was sampled for further processing of FVT
solutions, while the pellet was resuspended in PBS buffer for bacterial DNA extraction. The fecal
supernatant was filtered through a 0.45 µm Minisart High Flow PES syringe filter (Sartorius) to
remove bacteria and other larger particles. This filtrate was concentrated using Centriprep Ultracel
YM-30K units (Millipore) that by its design of an inner and outer tube removes metabolites below
the size of 30 kDa. The permeate in the inner tube was discarded several times during
centrifugation at 1,500 x g at 20°C until approximately 0.5 mL was left in the outer tube, which at
this point was considered as a fecal virome. The 30 kDa filter from the Centriprep Ultracel YM-
30K units was removed with a sterile scalpel and added to the fecal virome to allow viral particles
to diffuse. In order to trace back the origin of specific bacterial or viral taxa, the fecal viromes
were mixed based on cages, taking into account the coprophagic behavior of mice[60,61]. These fecal
viromes were mixed into one final mixture from mice of all three vendors representing the
"untreated fecal virome", FVT-UnT, which was immediately stored at -80°C. The remaining fecal
viromes were stored at 4°C prior to downstream processing to inactivate the eukaryotic viruses in
the fecal viromes by either dissolving the lipid membrane of enveloped viruses with
solvent/detergent treatment or inhibit replication of RNA viruses with pyronin Y.

Solvent/detergent treated fecal virome (FVT-SDT)

The solvent/detergent treatment is commonly used to inactivate enveloped viruses, as most
eukaryotic viruses possess an envelope, while non-enveloped viruses, including phages, are not
affected by this treatment[62–64]. Following the guidelines set by the World Health Organization
(WHO)[56] and Horowitz et al.[62] for the clinical use of solvent/detergent-treated plasma, the fecal
viromes were subjected to incubation in a solution containing 1% (w/v) tri(n-butyl) phosphate
(TnBP) and 1% (w/v) Triton X-100 at 30°C for 4 hours. It is important to note that the majority of

inactivation typically occurs within the first 30-60 minutes of the solvent/detergent treatment[56].
The removal of TnBP and Triton X-100 was performed according to the method described by
Treščec et al.[65]. The removal of the solvent/detergent agents from the fecal viromes yielded approx.
100 mL viral-flow-through from the column which was concentrated to 0.5 mL using Centriprep
Ultracel YM-30K units as described in the previous section. The final product constituted the FVT-
SDT treatment. The effect of the solvent/detergent treatment on phage activity was assessed with
traditional plaque assay of both non-enveloped phages (C2, T4, and phiX174) and one enveloped
phage (phi6).

Pyronin Y treated fecal virome (FVT-PYT)
Pyronin Y (Merck) is a strong red-colored fluorescent compound. It has been reported to exhibit
efficient binding to single-stranded and double-stranded RNA (ss/dsRNA), while its binding to
single-stranded and double-stranded DNA (ss/dsDNA) is less effective[66,67]. Initial screening was
conducted to determine the optimal conditions for inactivation using various concentrations of
pyronin Y, different incubation times, and temperatures for RNA and DNA phages. The fecal
filtrate was treated with 100 µM pyronin Y and incubated at 40°C overnight to inactivate viral
particles containing RNA genomes. To remove the pyronin Y molecules that were not bound, the
pyronin Y-treated fecal filtrate suspensions were diluted in 50 mL SM buffer and subsequently
concentrated to 0.5 mL using Centriprep Ultracel YM-30K units. This process was repeated three
times, resulting in a transparent appearance of the pyronin Y-treated fecal filtrate, which
constituted the FVT-PyT treatment.

Fecal microbiota transplantation (FMT)
The mouse intestinal content that was sampled anaerobically (Fig. S12) was diluted 1:20 in an
anoxic cryoprotectant consisting of PBS buffer and 20% (v/v) glycerol and stored at -80°C until
administration.

Chemostat propagated fecal virome (FVT-ChP)
The preparation of the chemostat propagated virome was performed as described previously[29].
Briefly, anaerobic-handled mouse cecum content was utilized for chemostat propagation. The
culture medium was formulated to resemble the low-fat (LF) diet (Research Diets D12450J)
provided to the donor mice as their feed (Table S3), and growth conditions such as temperature
and pH were set to simulate the environmental conditions present in the mouse cecum. The end
cultures, which underwent fermentation with a slow dilution rate (0.05 volumes per hour),

exhibited a microbial composition that closely resembled the initial microbial composition profile

of the donor[29]. These batches were combined to form the FVT-ChP treatment.

Fluorescence microscopy

427   Virus-like particle (VLP) counts were evaluated of all fecal viromes (FVT-UnT, FVT-SDT, FVT-

428   ChP, and FVT-PyT, Fig. S13) by epifluorescence microscopy using SYBR Gold staining (Thermo

429   Scientific) as described online dx.doi.org/10.17504/protocols.io.bx6cpraw.

Cytokine analysis

Pre-weighted cecum tissue was homogenized in 400μl lysis buffer (stock solution: 10ml Tris lysis

buffer, 100μl phosphatase inhibitor 1, 100μl phosphatase inhibitor 2, and 200μl protease inhibitor

(MSD inhibitor pack, Meso Scale Discovery, Rockville, MD) using a FastPrep Bead Beater

Homogenizer (MP Biomedicals, Irvine, CA), and centrifuged (8000g; 4°C; 5 min). Samples were

diluted 1:2 and analyzed for IFN-γ, GM-CSF, IL-15, IL-6, IL-10, KC/GRO, MIP-2, TNF-α IL-

17A/F, and IL-22 in a customized metabolic group 1 U-PLEX (MSD) according to manufacturer's

instructions. The plates were read and analyzed on a MESO QuickPlex SQ 120MM instrument

using MSD's own analysis software Discovery Workbench.

Histology and cytotoxicity assay

Formalin-fixed, paraffin-embedded cecum tissue sections were stained with hematoxylin and

eosin for histopathological evaluation by a blinded investigator (co-author AB). A composite score

was assigned, taking into account the following pathological features: 1) immune cell infiltration,

2) submucosal edema or hemorrhage, 3) epithelial injury, each with a range of severity/extent as

follows: 0: none, 1: mild, 2: moderate, 3: severe) for a cumulative pathology grade between 0 and

9 [15]. The RIDASCREEN *C. difficile* Toxin A/B ELISA kit (r-biopharm) was used to measure the

toxin concentrations in the mice feces by following the instructions of the manufacturer. The

OD_{450nm} was measured with a Varioskan Flash plate reader (Thermo Fisher Scientific).

Quantitative real-time PCR measuring *C. difficile* abundance

C. difficile in the fecal samples was enumerated using quantitative real-time polymerase chain

reaction (qPCR) with specie specific primers (C.Diff_ToxA_Fwd: 5'-TCT ACC ACT GAA GCA

TTA C-3', C.Diff_ToxA_Rev: 5'-TAG GTA CTG TAG GTT TAT TG-3'[68]) purchased from

Integrated DNA Technologies. Standard curves (Table S4) were based on a dilution series of total

DNA extracted from monocultures *C. difficile* VPI 10463. The qPCR results were obtained using

the CFX96 Touch Real-Time PCR Detections System (Bio-Rad Laboratories) and the reagent

RealQ plus 2x Master Mix Green low Rox (Amplicon) as previously described[69].

Pre-processing of fecal samples for separation of viruses and bacteria

Fecal samples from three different time points were included to investigate gut microbiome

changes over time: baseline (before antibiotic treatment), before *C. difficile* infection, and at

termination. This represented in total 142 fecal samples. Separation of the viruses and bacteria

from the fecal samples generated a fecal pellet and fecal supernatant by centrifugation and 0.45

µm filtering as described previously[59], except the volume of fecal homogenate was adjusted to 5

mL using SM buffer.

Bacterial DNA extraction, sequencing and pre-processing of raw data

The DNeasy PowerSoil Pro Kit (Qiagen) was used to extract bacterial DNA from the fecal pellet

by following the instructions of the manufacturer. The final purified DNA was stored at -80ºC and

the DNA concentration was determined using Qubit HS Assay Kit (Invitrogen) on the Qubit 4

Fluorometric Quantification device (Invitrogen). The bacterial community composition was

determined by Illumina NextSeq-based high-throughput sequencing of the 16S rRNA gene V3-

region, as previously described [59]. Quality control of reads, de-replicating, purging from chimeric

reads and constructing zOTU was conducted with the UNOISE pipeline[70] and taxonomy assigned

with Sintax [71] using the EZtaxon for 16S rRNA gene database[72]. Code describing this pipeline can

be accessed in github.com/jcame/Fastq_2_zOTUtable. The average sequencing depth after quality

control (Accession: PRJEB58777, available at ENA) for the fecal 16S rRNA gene amplicons was

60,719 reads (min. 11,961 reads and max. 198,197 reads).

Viral RNA/DNA extraction, sequencing and pre-processing of raw data

The sterile filtered fecal supernatant was concentrated using Centrisart centrifugal filters with a

filter cut-off at 100 kDA (Sartorius) by centrifugation at 1,500 x g at 4ºC

(dx.doi.org/10.17504/protocols.io.b2qaqdse). The viral DNA/RNA was extracted from the fecal

supernatants using the Viral RNA mini kit (Qiagen) as previously described [59]. Reverse

transcription was executed with SuperScript IV VILO Master mix by following the instructions of

the manufacturer and subsequently cleaned with DNeasy blood and tissue kit (Qiagen) by only

following step 3-8 in the instructions from the manufacturer. In brief, the DNA/cDNA samples

were mixed with ethanol, bound to the silica filter, washed two times, and eluted with 40 µL elution

buffer. Multiple displacement amplification (MDA, to include ssDNA viruses) using GenomiPhi

V3 DNA amplification kit (Cytiva) and sequencing library preparation using the Nextera XT kit

(Illumina) was performed at previously described[59], and sequenced using the Illumina NovaSeq

platform at the sequencing facilities of Novogene (Cambridge, UK). The average sequencing depth

of raw reads (Accession: PRJEB58777, available at ENA) for the fecal viral metagenome was

17,384,372 reads (min. 53,960 reads and max. 81,642,750 reads). Pre-processing of raw

sequencing data and viral contig assembly were performed as previously described[73]. Code

describing this pipeline can be accessed in github.com/jcame/virome_analysis-FOOD.

Bioinformatic analysis of bacterial and viral sequences

Initially the dataset was purged for zOTU's/viral contigs, which were detected in less than 5% of

the samples, but the resulting dataset still maintained >99.8% of the total reads. Cumulative sum

scaling (CSS)[74] was applied for the analysis of β-diversity. CSS normalization was performed

using the R software using the metagenomeSeq package[75]. R version 4.2.2 was used for subsequent

analysis and presentation of data. The main packages used were phyloseq[76], vegan[77], DESeq2[78],

ampvis2[79], ggpubr[80], and ggplot2[80].

Statistical analysis

α-diversity analysis was based on raw read counts for bacteriome analysis and normalized (RPKM)

read counts for virome analysis, while statistics were based on ANOVA. β-diversity was

represented by Bray-Curtis dissimilarity and statistics were based on PERMANOVA. Cytokine

levels, toxin levels, *C. difficile* abundance, and histology data were analyzed in R using linear

models with saline as control group, while log rank test was used to compare the survival

distributions.

List of Supplementary Materials

This PDF file includes:

Fig. S1 to S13

Table S1 to S4

Acknowledgement

We would like to express our gratitude to the staff of veterinarians and animal caretakers at the

Department of Experimental Medicine (AEM, University of Copenhagen, Denmark) for their

cooperation in housing, handling, and monitoring the mice. We would also like to extend our

thanks to PhD Casper Normann Nurup for his assistance in the initial setup of the column

chromatography, and to lab trainee Mariam Al-Batool Samir S Bagi for performing the ELISA

assay. Lastly, we would like to acknowledge the Food & Health Open Innovation project

(FOODHAY), granted by the Danish Ministry of Education and Research, for funding the qPCR

equipment (Bio-Rad Laboratories CFX96) used in this study.

Funding

520 Funding was provided by:

521 The Lundbeck Foundation with grant ID: R324-2019-1880 under the acronym "SafeVir".

522 The Novo Nordisk Foundation with grant ID: NNF-20OC0063874 under the acronym "PrePhage".

Author contributions

Conceptualization: TSR, DSN

Methodology: TSR, SF, SBL, KDT, AVM (veterinarian, supervising the health status monitoring),

AB (scored histology images), JLCM, SA, KA, CHFH

Investigation: TSR, SF, AB, AVM, CHFH, AKH, DSN

Visualization: TSR, AB

Funding acquisition: TSR, DSN

Project administration: TSR, DSN

Supervision: DSN, AKH

Writing – original draft: TSR

Writing – review & editing: All authors critically revised and approved the final version of the

manuscript.

Competing interests

536 All authors declare no financial or personal conflicts of interest.

Data and materials availability

538 All data associated with this study are present in the paper or the Supplementary Materials. All

539 sequencing datasets are available in the ENA database under accession number PRJEB58777.

References

1 Millan Braden, Park Heekuk, Hotte Naomi, Mathieu Olivier, Burguiere Pierre, Tompkins Thomas A, et al.
Fecal Microbial Transplants Reduce Antibiotic-resistant Genes in Patients With Recurrent *Clostridium*
difficile Infection. *Clin Infect Dis* 2016;**62**(12):1479–86. Doi: 10.1093/cid/ciw185.

2 Maruvada Padma, Leone Vanessa, Kaplan Lee M, Chang Eugene B. The Human Microbiome and Obesity:
Moving beyond Associations. *Cell Host Microbe* 2017;**22**(5):589–99. Doi: 10.1016/j.chom.2017.10.005.

3 Ott Stephan J, Waetzig Georg H, Rehman Ateequr, Moltzau-Anderson Jacqueline, Bharti Richa, Grasis Juris
A, et al. Efficacy of Sterile Fecal Filtrate Transfer for Treating Patients With *Clostridium difficile* Infection.
Gastroenterology 2017;**152**(4):799-811.e7. Doi: 10.1053/j.gastro.2016.11.010.

4 Gupta Shaan, Allen-Vercoe Emma, Petrof Elaine O. Fecal microbiota transplantation: in perspective. *Therap*
Adv Gastroenterol 2016;**9**(2):229–39. Doi: 10.1177/1756283X15607414.

5 Kassam Zain, Dubois Nancy, Ramakrishna Bharat, Ling Kelly, Qazi Taha, Smith Mark, et al. Donor
Screening for Fecal Microbiota Transplantation. *New England Journal of Medicine* 2019;**381**(21):2070–2.
Doi: 10.1056/NEJMc1913670.

6 Hota Susy S., Surangiwala Salman, Paterson Aimee S., Coburn Bryan, Poutanen Susan M. Regional
variability in fecal microbiota transplantation practices: a survey of the Southern Ontario Fecal Microbiota
Transplantation Movement. *CMAJ Open* 2018;**6**(2):E184–90. Doi: 10.9778/cmajo.20170109.

7 Olesen Scott W. Power calculations for detecting differences in efficacy of fecal microbiota donors.
Contemp Clin Trials Commun 2020;**20**. Doi: 10.1016/j.conctc.2020.100674.

8 Wilson Brooke C., Vatanen Tommi, Cutfield Wayne S., O'Sullivan Justin M. The super-donor phenomenon
in fecal microbiota transplantation. *Front Cell Infect Microbiol* 2019;**9**(January):1–11. Doi:
10.3389/fcimb.2019.00002.

9 U.S. Food & Drug Administration. Important Safety Alert Regarding Use of Fecal Microbiota for
Transplantation and Risk of Serious Adverse Reactions Due to Transmission of Multi-Drug Resistant
Organisms. Available at https://www.fda.gov/vaccines-blood-biologics/safety-availability-
biologics/important-safety-alert-regarding-use-fecal-microbiota-transplantation-and-risk-serious-adverse.
Accessed July 4, 2019, 2019.

10 U.S. Food & Drug Administration. Safety Alert Regarding Use of Fecal Microbiota for Transplantation and
Additional Safety Protections Pertaining to Monkeypox Virus. 2022.

11 U.S. Food & Drug Administration. Safety Alert Regarding Use of Fecal Microbiota for Transplantation and
Risk of Serious Adverse Events Likely Due to Transmission of Pathogenic Organisms. Available at
https://www.fda.gov/vaccines-blood-biologics/safety-availability-biologics/safety-alert-regarding-use-fecal-
microbiota-transplantation-and-risk-serious-adverse-events-likely. Accessed May 23, 2023, 2020.

12 Kao D H, Roach B, Walter J, Lobenberg R, Wong K. Effect of lyophilized sterile fecal filtrate vs lyophilized
donor stool on recurrent *Clostridium difficile* infection (rCDI): Preliminary results from a randomized,
double-blind pilot study. *J Can Assoc Gastroenterol* 2019;**2**(Supplement_2):101–2. Doi:
10.1093/jcag/gwz006.050.

13   Cao Zhirui, Sugimura Naoki, Burgermeister Elke, Ebert Matthias P., Zuo Tao, Lan Ping. The gut virome: A
578   new microbiome component in health and disease. *EBioMedicine* 2022;**81**:104113. Doi:
579   10.1016/j.ebiom.2022.104113.

580   14   Draper Lorraine A, Ryan Feargal J, Dalmasso Marion, Casey Pat G, McCann Angela, Velayudhan
581   Vimalkumar, et al. Autochthonous faecal viral transfer (FVT) impacts the murine microbiome after
582   antibiotic perturbation. *BMC Biol* 2020;**18**(1):173. Doi: 10.1186/s12915-020-00906-0.

583   15   Rasmussen Torben Sølbeck, Mentzel Caroline Märta Junker, Kot Witold, Castro-Mejía Josué Leonardo,
584   Zuffa Simone, Swann Jonathan Richard, et al. Faecal virome transplantation decreases symptoms of type 2
585   diabetes and obesity in a murine model. *Gut* 2020;**69**(12):2122–30. Doi: 10.1136/gutjnl-2019-320005.

586   16   Borin Joshua M., Liu Roland, Wang Yanhan, Wu Tsung-Chin, Chopyk Jessica, Huang Lina, et al. Fecal
587   virome transplantation is sufficient to alter fecal microbiota and drive lean and obese body phenotypes in
588   mice. *BioRxiv* 2023. Doi: 10.1101/2023.02.03.527064.

589   17   Brunse Anders, Deng Ling, Pan Xiaoyu, Hui Yan, Castro-Mejía Josué L, Kot Witold, et al. Fecal filtrate
590   transplantation protects against necrotizing enterocolitis. *ISME J* 2022;**16**(3):686–94. Doi: 10.1038/s41396-
591   021-01107-5.

592   18   Lim Efrem S., Zhou Yanjiao, Zhao Guoyan, Bauer Irma K., Droit Lindsay, Ndao I. Malick, et al. Early life
593   dynamics of the human gut virome and bacterial microbiome in infants. *Nat Med* 2015;**21**(10):1228–34.
594   Doi: 10.1038/nm.3950.

595   19   Shah Shiraz A., Deng Ling, Thorsen Jonathan, Pedersen Anders G., Dion Moïra B., Castro-Mejía Josué L.,
596   et al. Expanding known viral diversity in the healthy infant gut. *Nat Microbiol* 2023. Doi: 10.1038/s41564-
597   023-01345-7.

598   20   Doorbar John, Egawa Nagayasu, Griffin Heather, Kranjec Christian, Murakami Isao. Human papillomavirus
599   molecular biology and disease association. *Rev Med Virol* 2015;**25**(S1):2–23. Doi: 10.1002/rmv.1822.

600   21   Rasmussen Torben Sølbeck, Koefoed Anna Kirstine, Jakobsen Rasmus Riemer, Deng Ling, Castro-Mejía
601   Josué L, Brunse Anders, et al. Bacteriophage-mediated manipulation of the gut microbiome - promises and
602   presents limitations. *FEMS Microbiol Rev* 2020;**44**(4):507–21. Doi: 10.1093/femsre/fuaa020.

603   22   Champagne-Jorgensen Kevin, Luong Tiffany, Darby Taylor, Roach Dwayne R. Immunogenicity of
604   bacteriophages. *Trends Microbiol* 2023. Doi: 10.1016/j.tim.2023.04.008.

605   23   Bichet Marion C., Chin Wai Hoe, Richards William, Lin Yu-Wei, Avellaneda-Franco Laura, Hernandez
606   Catherine A., et al. Bacteriophage uptake by mammalian cell layers represents a potential sink that may
607   impact phage therapy. *IScience* 2021;**24**(4):102287. Doi: 10.1016/j.isci.2021.102287.

608   24   D'Herelle Felix. On an invisible microbe antagonistic toward dysenteric bacilli: brief note by Mr. F.
609   D'Herelle, presented by Mr. Roux. 1917. *Res Microbiol* 1917;**158**(7):553–4. Doi:
610   10.1016/j.resmic.2007.07.005.

611   25   Górski Andrzej, Miedzybrodzki Ryszard, Weber-Dabrowska Beata, Fortuna Wojciech, Letkiewicz
612   Sławomir, Rogóz Paweł, et al. Phage therapy: Combating infections with potential for evolving from merely

a treatment for complications to targeting diseases. *Front Microbiol* 2016;**7**(SEP):1–9. Doi:
10.3389/fmicb.2016.01515.

26   Al-Ishaq Raghad Khalid, Skariah Sini, Büsselberg Dietrich. Bacteriophage Treatment: Critical Evaluation of
Its Application on World Health Organization Priority Pathogens. *Viruses* 2020;**13**(1). Doi:
10.3390/v13010051.

27   Loc-Carrillo Catherine, Abedon Stephen T. Pros and cons of phage therapy. *Bacteriophage* 2011;**1**(2):111–
4. Doi: 10.4161/bact.1.2.14590.

28   Liu Dan, van Belleghem Jonas D, de Vries Christiaan R, Burgener Elizabeth, Chen Qingquan, Manasherob
Robert, et al. The Safety and Toxicity of Phage Therapy: A Review of Animal and Clinical Studies. *Viruses*
2021;**13**(7):1–20. Doi: 10.3390/v13071268.

29   Adamberg Signe, Rasmussen Torben Sølbeck, Larsen Sabina Birgitte, Nielsen Dennis Sandris, Adamberg
Kaarel. Reproducible chemostat cultures to eliminate eukaryotic viruses from fecal transplant material.
BioRxiv 2023. Doi: 10.1101/2023.03.15.529189.

30   Rey Felix A., Lok Shee Mei. Common Features of Enveloped Viruses and Implications for Immunogen
Design for Next-Generation Vaccines. *Cell* 2018:1319–34. Doi: 10.1016/j.cell.2018.02.054.

31   Bell Philip J L. Evidence supporting a viral origin of the eukaryotic nucleus. *Virus Res* 2020;**289**:198168.
Doi: 10.1016/j.virusres.2020.198168.

32   Koonin Eugene v., Dolja Valerian v., Krupovic Mart. Origins and evolution of viruses of eukaryotes: The
ultimate modularity. *Virology* 2015:2–25. Doi: 10.1016/j.virol.2015.02.039.

33   Sausset R., Petit M. A., Gaboriau-Routhiau V., De Paepe M. New insights into intestinal phages. *Mucosal*
Immunol 2020;**13**(2):205–15. Doi: 10.1038/s41385-019-0250-5.

34   Xu Qiaomai, Zhang Shumeng, Quan Jiazheng, Wu Zhengjie, Gu Silan, Chen Yunbo, et al. The evaluation of
fecal microbiota transplantation vs vancomycin in a *Clostridioides difficile* infection model. *Appl Microbiol*
Biotechnol 2022;**106**(19–20):6689–700. Doi: 10.1007/s00253-022-12154-z.

35   Seekatz Anna M, Theriot Casey M, Molloy Caitlyn T, Wozniak Katherine L, Bergin Ingrid L, Young
Vincent B. Fecal Microbiota Transplantation Eliminates *Clostridium difficile* in a Murine Model of
Relapsing Disease. *Infect Immun* 2015;**83**(10):3838–46. Doi: 10.1128/IAI.00459-15.

36   Chen Xinhua, Katchar Kianoosh, Goldsmith Jeffrey D, Nanthakumar Nanda, Cheknis Adam, Gerding Dale
N, et al. A mouse model of *Clostridium difficile*-associated disease. *Gastroenterology* 2008;**135**(6):1984–92.
Doi: 10.1053/j.gastro.2008.09.002.

37   Moore R, Pothoulakis C, LaMont J T, Carlson S, Madara J L. *C. difficile* toxin A increases intestinal
permeability and induces Cl- secretion. *Am J Physiol* 1990;**259**(2 Pt 1):G165-72. Doi:
10.1152/ajpgi.1990.259.2.G165.

38   Rao Krishna, Erb-Downward John R., Walk Seth T., Micic Dejan, Falkowski Nicole, Santhosh Kavitha, et
al. The Systemic Inflammatory Response to *Clostridium difficile* Infection. *PLoS One* 2014;**9**(3):e92578.
Doi: 10.1371/journal.pone.0092578.

39 Bischoff Stephan C., Barbara Giovanni, Buurman Wim, Ockhuizen Theo, Schulzke Jörg Dieter, Serino
Matteo, et al. Intestinal permeability - a new target for disease prevention and therapy. *BMC Gastroenterol*
2014:189. Doi: 10.1186/s12876-014-0189-7.

40 Camilleri Michael. Leaky gut: mechanisms, measurement and clinical implications in humans. *Gut*
2019:1516–26. Doi: 10.1136/gutjnl-2019-318427.

41 Sharma Vikas, Mobeen Fauzul, Prakash Tulika. Comparative Genomics of *Herpesviridae* Family to Look
for Potential Signatures of Human Infecting Strains. *Int J Genomics* 2016;**2016**:1–10. Doi:
10.1155/2016/9543274.

42 Redding Laurel E, Berry Alexander S, Indugu Nagaraju, Huang Elizabeth, Beiting Daniel P, Pitta Dipti. Gut
microbiota features associated with *Clostridioides difficile* colonization in dairy calves. *PLoS One*
2021;**16**(12):e0251999. Doi: 10.1371/journal.pone.0251999.

43 Zhu Jia-Jia, Gao Ming-Xing, Song Xiao-Jun, Zhao Li, Li Yu-Wen, Hao Zhi-Hui. Changes in bacterial
diversity and composition in the faeces and colon of weaned piglets after feeding fermented soybean meal. *J*
Med Microbiol 2018;**67**(8):1181–90. Doi: 10.1099/jmm.0.000766.

44 Zuo Tao, Wong Sunny H, Lam Kelvin, Lui Rashid, Cheung Kitty, Tang Whitney, et al. Bacteriophage
transfer during faecal microbiota transplantation in *Clostridium difficile* infection is associated with
treatment outcome. *Gut* 2018;**67**(4):634–43. Doi: 10.1136/gutjnl-2017-313952.

45 Fujimoto Kosuke, Kimura Yasumasa, Allegretti Jessica R., Yamamoto Mako, Zhang Yao-zhong, Katayama
Kotoe, et al. Functional Restoration of Bacteriomes and Viromes by Fecal Microbiota Transplantation.
Gastroenterology 2021;**160**(6):2089-2102.e12. Doi: 10.1053/j.gastro.2021.02.013.

46 Mao Xiaotian, Larsen Sabina Birgitte, Zachariassen Line Sidsel Fisker, Brunse Anders, Adamberg Signe,
Castro-Meija Josue Leonardo, et al. A reproducible enteric phage community improves blood glucose
regulation in an obesity mouse model. *BioRxiv* 2023. Doi: 10.1101/2023.03.20.532903.

47 Darkoh Charles, DuPont Herbert L, Norris Steven J, Kaplan Heidi B. Toxin synthesis by *Clostridium*
difficile is regulated through quorum signaling. *MBio* 2015;**6**(2):e02569. Doi: 10.1128/mBio.02569-14.

48 Callanan Julie, Stockdale Stephen R., Shkoporov Andrey, Draper Lorraine A., Ross R. Paul, Hill Colin.
Biases in viral metagenomics-based detection, cataloguing and quantification of bacteriophage genomes in
human faeces, a review. *Microorganisms* 2021:1–15. Doi: 10.3390/microorganisms9030524.

49 Goodacre Norman, Aljanahi Aisha, Nandakumar Subhiksha, Mikailov Mike, Khan Arifa S. A Reference
Viral Database (RVDB) To Enhance Bioinformatics Analysis of High-Throughput Sequencing for Novel
Virus Detection. *MSphere* 2018;**3**(2):1–18. Doi: 10.1128/mSphereDirect.00069-18.

50 Rasmussen Torben Sølbeck, Jakobsen Rasmus Riemer, Castro-Mejía Josué L, Kot Witold, Thomsen Allan
Randrup, Vogensen Finn Kvist, et al. Inter-vendor variance of enteric eukaryotic DNA viruses in specific
pathogen free C57BL/6N mice. *Res Vet Sci* 2021;**136**(January):1–5. Doi: 10.1016/j.rvsc.2021.01.022.

51 Fay Elizabeth J, Balla Keir M, Roach Shanley N, Shepherd Frances K, Putri Dira S, Wiggen Talia D, et al.
Natural rodent model of viral transmission reveals biological features of virus population dynamics. *J Exp*
Med 2022;**219**(2). Doi: 10.1084/jem.20211220.

52   Mullish Benjamin H, Quraishi Mohammed Nabil, Segal Jonathan P, McCune Victoria L, Baxter Melissa,
Marsden Gemma L, et al. The use of faecal microbiota transplant as treatment for recurrent or refractory
Clostridium difficile infection and other potential indications: joint British Society of Gastroenterology
(BSG) and Healthcare Infection Society (HIS) guidelines. *Gut* 2018;**67**(11):1920–41. Doi: 10.1136/gutjnl-
2018-316818.

53   Hui Wenjia, Li Ting, Liu Weidong, Zhou Chunyan, Gao Feng. Fecal microbiota transplantation for
treatment of recurrent *C. difficile* infection: An updated randomized controlled trial meta-analysis. *PLoS*
One 2019;**14**(1):e0210016. Doi: 10.1371/journal.pone.0210016.

54   Martinez-Gili Laura, McDonald Julie A.K. K, Liu Zhigang, Kao Dina, Allegretti Jessica R., Monaghan
Tanya M., et al. Understanding the mechanisms of efficacy of fecal microbiota transplant in treating
recurrent *Clostridioides difficile* infection and beyond: the contribution of gut microbial-derived metabolites.
Gut Microbes 2020;**12**(1):1810531. Doi: 10.1080/19490976.2020.1810531.

55   Santiago-Rodriguez Tasha M., Ly Melissa, Daigneault Michelle C., Brown Ian H.L., McDonald Julie A.K.,
Bonilla Natasha, et al. Chemostat culture systems support diverse bacteriophage communities from human
feces. *Microbiome* 2015;**3**:58. Doi: 10.1186/s40168-015-0124-3.

56   WHO Health Product Policy and Standards Team. Guidelines on viral inactivation and removal procedures
intended to assure the viral safety of human blood plasma products. Available at
https://www.who.int/publications/m/item/WHO-TRS924-Annex4. Accessed January 2, 2023, 2004.

57   Edwards Adrianne N, McBride Shonna M. Isolating and Purifying *Clostridium difficile* Spores. *Methods*
Mol Biol 2016;**1476**:117–28. Doi: 10.1007/978-1-4939-6361-4_9.

58   Rasmussen Torben Sølbeck, Streidl Theresa, Hitch Thomas C. A., Wortmann Esther, Deptula Paulina,
Kofoed Michael V. W., et al. *Sporofaciens musculi* gen. nov., sp. nov., a novel bacterium isolated from the
caecum of an obese mouse. *Int J Syst Evol Microbiol* 2021;**71**(3). Doi: 10.1099/ijsem.0.004673.

59   Rasmussen Torben Sølbeck, de Vries Liv, Kot Witold, Hansen Lars Hestbjerg, Castro-Mejía Josué L,
Vogensen Finn Kvist, et al. Mouse Vendor Influence on the Bacterial and Viral Gut Composition Exceeds
the Effect of Diet. *Viruses* 2019;**11**(5). Doi: 10.3390/v11050435.

60   Bibiloni Rodrigo. Rodent models to study the relationships between mammals and their bacterial
inhabitants. *Gut Microbes* 2012;**3**(6):536–43. Doi: 10.4161/gmic.21905.

61   Torres Pedro J, Ho Bryan S, Arroyo Pablo, Sau Lillian, Chen Annie, Kelley Scott T, et al. Exposure to a
Healthy Gut Microbiome Protects Against Reproductive and Metabolic Dysregulation in a PCOS Mouse
Model. *Endocrinology* 2019;**160**(5):1193–204. Doi: 10.1210/en.2019-00050.

62   Horowitz B, Bonomo R, Prince A M, Chin S N, Brotman B, Shulman R W. Solvent/detergent-treated
plasma: a virus-inactivated substitute for fresh frozen plasma. *Blood* 1992;**79**(3):826–31.

63   Prince A M, Horowitz B, Brotman B. Sterilisation of hepatitis and HTLV-III viruses by exposure to tri(n-
butyl)phosphate and sodium cholate. *Lancet* 1986;**1**(8483):706–10. Doi: 10.1016/s0140-6736(86)91101-3.

64   Remy M M, Alfter M, Chiem M-N, Barbani M T, Engler O B, Suter-Riniker F. Effective chemical virus

inactivation of patient serum compatible with accurate serodiagnosis of infections. *Clin Microbiol Infect*

2019;**25**(7):907.e7-907.e12. Doi: 10.1016/j.cmi.2018.10.016.

65   Trescec A, Simić M, Branović K, Gebauer B, Benko B. Removal of detergent and solvent from solvent-

detergent-treated immunoglobulins. *J Chromatogr A* 1999;**852**(1):87–91. Doi: 10.1016/s0021-

9673(99)00178-8.

66   Darzynkiewicz Z, Kapuscinski J, Carter S P, Schmid F A, Melamed M R. Cytostatic and cytotoxic

properties of pyronin Y: relation to mitochondrial localization of the dye and its interaction with RNA.

Cancer Res 1986;**46**(11):5760–6.

67   Kapuscinski J, Darzynkiewicz Z. Interactions of pyronin Y(G) with nucleic acids. *Cytometry*

1987;**8**(2):129–37. Doi: 10.1002/cyto.990080205.

68   Gillers Sara, Atkinson Christopher D, Bartoo Aaron C, Mahalanabis Madhumita, Boylan Michael O,

Schwartz John H, et al. Microscale sample preparation for PCR of *C. difficile* infected stool. *J Microbiol*

Methods 2009;**78**(2):203–7. Doi: 10.1016/j.mimet.2009.05.020.

69   Ellekilde M, Krych L, Hansen C H F, Hufeldt M R, Dahl K, Hansen L H, et al. Characterization of the gut

microbiota in leptin deficient obese mice - Correlation to inflammatory and diabetic parameters. *Res Vet Sci*

2014;**96**(2):241–50. Doi: 10.1016/j.rvsc.2014.01.007.

70   Edgar Robert C. Updating the 97% identity threshold for 16S ribosomal RNA OTUs. *Bioinformatics*

2018;**34**(14):2371–5. Doi: 10.1093/bioinformatics/bty113.

71   Edgar Robert. SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS sequences. *BioRxiv*

2016:074161. Doi: 10.1101/074161.

72   Kim Ok-Sun, Cho Yong-Joon, Lee Kihyun, Yoon Seok-Hwan, Kim Mincheol, Na Hyunsoo, et al.

Introducing EzTaxon-e: a prokaryotic 16S rRNA gene sequence database with phylotypes that represent

uncultured species. *Int J Syst Evol Microbiol* 2012;**62**(Pt_3):716–21. Doi: 10.1099/ijs.0.038075-0.

73   Rasmussen Torben Sølbeck, Mentzel Caroline M Junker, Refslund Malene, Jakobsen Rasmus Riemer,

Zachariassen Line Sidsel Fisker, Castro-Mejia Leonardo, et al. Fecal virome transfer improves proliferation

of commensal gut Akkermansia muciniphila and unexpectedly enhances the fertility rate in laboratory mice.

BioRxiv 2022. Doi: 10.1101/2022.06.30.498243.

74   Paulson Joseph N, Stine O Colin, Bravo Héctor Corrada, Pop Mihai. Differential abundance analysis for

microbial marker-gene surveys. *Nat Methods* 2013;**10**(12):1200–2. Doi: 10.1038/nmeth.2658.

75   Paulson Jn. metagenomeSeq: Statistical analysis for sparse high-throughput sequencing. *BioconductorJp*

2014:1–20.

76   McMurdie Paul J., Holmes Susan. phyloseq: an R package for reproducible interactive analysis and graphics

of microbiome census data. *PLoS One* 2013;**8**(4):e61217. Doi: 10.1371/journal.pone.0061217.

77   Dixon Philip. VEGAN, a package of R functions for community ecology. *Journal of Vegetation Science*

2003;**14**(6):927–30. Doi: 10.1111/j.1654-1103.2003.tb02228.x.

78   Love Michael I, Huber Wolfgang, Anders Simon. Moderated estimation of fold change and dispersion for

758   RNA-seq data with DESeq2. *Genome Biol* 2014;**15**(12):550. Doi: 10.1186/s13059-014-0550-8.

759   79   Andersen Kasper S., Kirkegaard Rasmus H., Karst Søren M., Albertsen Mads. ampvis2: an R package to

760   analyse and visualise 16S rRNA amplicon data. *BioRxiv* 2018. Doi: 10.1101/299537.

761   80   Wickham Hadley. ggplot2. *Wiley Interdiscip Rev Comput Stat* 2011;**3**(2):180–5. Doi: 10.1002/wics.147.

765 Fig. 1: Evaluation of inactivation of phage activity with solvent/detergent (S/D) or pyronin Y treatment. A) Three non-enveloped

766 phages (phiX174, C2, and T4) and one enveloped phage (phi6) were treated with solvent/detergent, and their plaque activity (plaque

767 forming units per mL) was evaluated on their respective bacterial hosts. B) Phages representing ssDNA (phiX174), dsDNA (C2

768 and T4), ssRNA (MS2), and dsRNA (phi6) were treated with pyronin Y, and their plaque activity (PFU/mL) at different incubation

769 conditions was evaluated on their respective bacterial hosts. Dashed lines mark the detection limit of the applied assay.

Fig. 2: Overview of mouse phenotypic characteristics. A) Kaplan-Meier curve showing the survival probability of the mice that

was associated to the different treatments. B) Pathological score of cecum tissue evaluating the effect of the treatments' ability to

prevent *C. difficile* associated damage of the cecum tissue. C) to L) Showing the overall cytokine profile in the mouse cecum tissue

of the different treatments. The euthanized (cross) mice are differentiated from the mice that survived (circle) the *C. difficile*

infection. "**" = p < 0.005, "*" = p < 0.05, "." = p < 0.1.

Clostridioides difficile - Abundance

Clostridioides difficile - Toxin A/B levels

Fig. 3: Evaluation of *C. difficile* abundance A) to C) by qPCR targeting the *toxA* gene and D) to F) the associated toxin A/B levels measured with ELISA on feces samples from three different time points: before *C. difficile* inoculum, before 2nd treatment, and at study termination. The fraction below the bar plots highlights the number of mice that were detected positive of either *C. difficile* or toxin A/B. Dashed line marks the detection limit of the applied assay. * = p < 0.05, ** = p < 0.01, *** = p < 0.005.

Fig.4: Bacteriome analysis based on 16S rRNA gene amplicon sequencing. A) The general bacterial Shannon diversity index (α-diversity) and B) Bray-Curtis dissimilarity based PCoA plot (β-diversity) at baseline (before antibiotic treatment), that was compared with the mice that had survived *C. difficile* infection regardless of the treatment and the euthanized mice. C) Barplot and D) heatmap illustrating the bacterial relative abundance of the dominating bacterial taxa that was associated to the different mice that either survived the infection or were euthanized. The "n" above the bar plots highlights the number of mice of which the taxonomical average was based on. *** = p < 0.005. **** = p < 0.0005.

Fig. 5: Metavirome analysis based on whole-genome sequencing. A) The viral Shannon diversity index (α-diversity) and B) Bray-Curtis dissimilarity based PCoA plot (β-diversity) at baseline (before antibiotic treatment), that was compared with the mice that had survived *C. difficile* infection regardless of the treatment and the euthanized mice. C) Barplot showing the relative abundance of the viral taxonomy, and D) heatmap illustrating the relative abundance of the bacterial hosts that are predicted on the basis of the viral sequences. The "n" above the bar plots highlights the number of mice of which the taxonomical average was based on. ** = p < 0.01, *** = p < 0.005.

Fig. 6: Overview of the animal model. The mice were initially treated with an antibiotic mixture in their drinking water, intraperitoneal (I.P.) injection of clindamycin, and then inoculated with *C. difficile* (~10^4 CFU). Eighteen hours after the mice were treated with either saline (as control), FMT (fecal microbiota transplantation), FVT-UnT (fecal virome transplantation – Untreated, e.i. sterile filtered donor feces), FVT-ChP (FVT-chemostat propagated fecal donor virome to remove eukaryotic viruses by dilution), FVT-SDT (FVT-solvent/detergent treated to inactivate enveloped viruses), and FVT-PyT (FVT-pyronin Y treated to inactivate RNA viruses). Crosses marks time points of feces sampling.