Title

Drosophila model to clarify the pathological significance of OPA1 in autosomal dominant optic

atrophy

Authors

Yohei Nitta[1,†], Jiro Osaka[1,2,†], Ryuto Maki[2], Satoko Hakeda-Suzuki[2,3], Emiko Suzuki[4,5], Satoshi Ueki[6],

Takashi Suzuki[2] and Atsushi Sugie[1*]

[1] Brain Research Institute, Niigata University, Niigata, 951-8585, Japan

[2] School of Life Science and Technology, Tokyo Institute of Technology, Yokohama, 226-8501, Japan

[3] Research Initiatives and Promotion Organization, Yokohama National University, Yokohama, 240-

8501, Japan

[4] Department of Biological Sciences, Graduate School of Science, Tokyo Metropolitan University,

Hachioji 192-0397, Japan

[5] Department of Gene Function and Phenomics, National Institute of Genetics, Mishima, 411-8540,

Japan

[6] Division of Ophthalmology and Visual Science, Graduate School of Medical and Dental Sciences,

Niigata University, Niigata, 951-8585, Japan

[†] Equal contribution

[*] Corresponding author: atsushi.sugie@bri.niigata-u.ac.jp (A.S.) Tel (+81) 25-227-0616.

Abstract

Autosomal dominant optic atrophy (DOA) is a progressive form of blindness caused by degeneration

of retinal ganglion cells and their axons, mainly caused by mutations in the OPA1 mitochondrial

dynamin *like* GTPase (*OPA1*) gene. *OPA1* encodes a dynamin-like GTPase present in the

mitochondrial inner membrane. When associated with OPA1 mutations, DOA can present not only

ocular symptoms but also multi-organ symptoms (DOA plus). DOA plus often results from point

mutations in the GTPase domain, which are assumed to have dominant negative effects. However, the

presence of mutations in the GTPase domain does not always result in DOA plus. Therefore, an

experimental system to distinguish between DOA and DOA plus is needed. In this study, we found

that loss-of-function mutations of the *dOPA1* gene in *Drosophila* can imitate the pathology of optic

nerve degeneration observed in DOA. We successfully rescued this degeneration by expressing the

human *OPA1* (*hOPA1*) gene, indicating that *hOPA1* is functionally interchangeable with *dOPA1* in the

fly system. However, mutations previously identified did not ameliorate the *dOPA1* deficiency

phenotype. By expressing both WT and DOA plus mutant *hOPA1* forms in the optic nerve of *dOPA1*

mutants, we observed that DOA plus mutations suppressed the rescue, facilitating the distinction

between loss-of-function and dominant negative mutations in *hOPA1*. This fly model aids in

distinguishing DOA from DOA plus and guides initial *hOPA1* mutation treatment strategies.

Introduction

Autosomal dominant optic atrophy (DOA) is a progressive form of blindness characterized by

selective degeneration of retinal ganglion cells (RGCs) and the axons that form the optic nerve. Despite

being a rare disease with a frequency of 1/12,000 to 1/50,000, DOA is the most commonly diagnosed

form of hereditary optic neuropathy. In 2000, *OPA1 mitochondrial dynamin like GTPase (OPA1)* was

identified as the causative gene for DOA (Alexander et al., 2000; Delettre et al., 2000).

OPA1 encodes a dynamin-like GTPase located in the inner mitochondrial membrane (Olichon et

al., 2002). It has various functions, including mitochondrial fusion, mtDNA maintenance, control of

cell death, and resistance to reactive oxygen species (ROS) (Lenaers et al., 2021). The most common

pathogenic mutation in *OPA1* is the c.2708-2711 delTTAG deletion (Toomes et al., 2001). Five

pathogenic mutations, including the one above, result in a substantial decrease in OPA1 protein in the

fibroblasts from affected patients (Zanna et al., 2008), supporting the theory that DOA is caused by

haploinsufficiency.

Optic atrophy is a characteristic feature of *OPA1*-associated DOA; however, multisystem

symptoms have been reported in up to 20% of *OPA1* mutation carriers (Yu-Wai-Man et al., 2010a).

This is called DOA plus. DOA plus can include sensorineural deafness (Amati-Bonneau et al., 2008),

multiple sclerosis-like illness (Verny et al., 2008; Yu-Wai-Man et al., 2016), parkinsonism and

dementia (Carelli et al., 2015; Lynch et al., 2017), and cardiomyopathy (Spiegel et al., 2016). The

R445H mutation is a well-known mutation associated with DOA plus (Amati-Bonneau et al., 2008,

2003; Shimizu et al., 2003). The proportion of mutated OPA1 locations differs between patients with

DOA and DOA plus. Missense mutations in the dynamin-featured structure, especially the GTPase

domain of *OPA1*, are more likely to cause severe symptoms compared with loss-of-function (LOF)

mutations such as deletions or splice site mutations (Yu-Wai-Man et al., 2010a). This is probably due

to the dominant negative effect (DN) of a DOA plus mutation on the normal *OPA1* allele, since OPA1

functions as a homo-oligomer (Frezza et al., 2006; Olichon et al., 2006); in fact, its yeast ortholog,

Mgm1, forms oligomers, increasing the GTPase activity (Meglei and McQuibban, 2009; Rujiviphat

et al., 2009). Particularly, the dynamin-featured structure accounts for 70% of the 233 pathogenic

variants of DOA and DOA plus described in the locus-specific database (Ferré et al., 2015). However,

mutations in these domains do not determine DOA plus, as it affects only 20% of all *OPA1* mutation

carriers. Thus, experimental models to determine whether a gene mutation is LOF or DN are required

to distinguish between DOA and DOA plus. Although currently there is no effective treatment for

DOA or DOA plus, distinguishing between them allows earlier planning of interventions such as

hearing aids, rather than relying solely on visual aids, necessary only for those with DOA plus. In the

quest to differentiate between LOF and DN effects within the context of genetic mutations,

precedents exist in simpler systems such as yeast and human fibroblasts. These models have

provided valuable insights into the conserved functions of OPA1 across species, as evidenced by

studies in yeast models (Del Dotto et al., 2017) and fibroblasts derived from patients harboring

OPA1 mutations (Kane et al., 2017). However, the ability to distinguish between LOF and DN

effects in an in vivo model organism, particularly at the structural level of retinal axon

degeneration, has remained elusive. This gap underscores the necessity for a more complex model

that not only facilitates molecular analysis but also enables the examination of structural changes

in axons and mitochondria, akin to those observed in the actual disease state.

As *OPA1* is conserved in various species, DOA models have been reported in vertebrates such as

mice and zebrafish, simple organisms such as nematodes and fruit flies, and *in vitro* yeast and cultured

cell models (Del Dotto and Carelli, 2021). In mouse models, focal RGC axons decrease in number

with mitochondrial abnormalities at the electron microscope level, but the phenotype appears slowly,

not allowing for analysis in a short time frame (Alavi et al., 2007; Davies et al., 2007; Nguyen et al.,

2011; Sarzi et al., 2012). Zebrafish models have displayed developmental delays in eyes and heads,

short length, and body axis abnormalities (Eijkenboom et al., 2019; Rahn et al., 2013) but have not

been used to investigate optic nerve degeneration.

In *C. elegans*, most studies focused on muscle cells using mutants of *eat-3*, an *OPA1* homolog

(Kanazawa et al., 2008; Rolland et al., 2009). However, the mitochondria in the posterior lateral

microtubule neurons did not show any difference in size from its wild-type (WT) counterparts (Byrne

et al., 2019). Thus, LOF of *eat-3* may have a limited impact on phenotypic outcomes in the nervous

system. Nevertheless, a DOA model in which $hOPA1^{K301A}$ is expressed in GABAergic motor neurons,

showed a decreased number of mitochondria in those cells (Zaninello et al., 2020). In a fly model, the

dOPA1 somatic clone in the eye exhibited a rough-eye phenotype, eye structural abnormalities, and

increased MitoSOX fluorescence, which was partially rescued by vitamin E or SOD1 expression

(Yarosh et al., 2008). Heterozygous *dOPA1* mutants showed shortened lifespan, elevated ROS levels,

and irregular muscle tissue mitochondrial structures (Tang et al., 2009). Moreover, the

Electroretinogram (ERG) pattern showed age-dependent decreases in the on transient response, heart

rate, negative geotaxis response, and increased heart arrhythmia due to heat shock stress (Shahrestani

et al., 2009). However, except for mice, the evidence of optic nerve axonal abnormalities is limited in

model organisms such as zebrafish, worms, and flies.

The structure of the *Drosophila* visual system is similar to that of mammals (Sanes and Zipursky,

2010), with conserved mechanisms of synaptogenesis and neural circuit formation (Sanes and

Zipursky, 2020). *Drosophila* photoreceptors type R7/8 extend their retinal axons from the retina

directly into the brain, forming synapses in the second optic ganglion medulla via the chiasma (Figure

1A), having potential application as a model for mammalian photoreceptors and RGCs. In this study,

we defined the retinal axons of *Drosophila* as analogous to the human optic nerve. Based on this, we

aimed to develop an experimental model for observing and quantifying degeneration in the *Drosophila*

retinal axon (Nitta et al., 2023; Richard et al., 2022).

In this study, a *dOPA1* LOF could mimic the human DOA, in which mitochondrial fragmentation,

increased ROS levels, and neurodegeneration occur in retinal axons. The axonal degeneration induced

by the *dOPA1* mutant was rescued by the expression of human *OPA1* (*hOPA1*). This demonstrated that

the function of *hOPA1* is the same as in the fly. Previously reported mutations failed to rescue the

dOPA1 deficiency phenotype. To distinguish between the effect of LOF and DN mutations, we

expressed the *hOPA1* gene of both WT and mutation in a genetic background in which *dOPA1* was

deleted in retinal axons. The DOA plus mutations D438V and R445H inhibited the rescue by the WT

of *hOPA1*. Taken together, we generated a new *Drosophila* DOA model which served to isolate the

LOF or DN effect of *hOPA1* mutations.

Results

122   ***dOPA1* depletion caused mitochondrial fragmentation in axon terminals**

123   The OPA1 protein is located in the inner mitochondrial membrane. To confirm whether dOPA1 is also

124   localized in the mitochondria of *Drosophila* retinal axons, we used a Gal4/UAS system (Brand and

125   Perrimon, 1993) and expressed HA-tagged dOPA1 proteins using the eye-specific *GMR-Gal4* driver

126   (Newsome et al., 2000a, 2000b). The outer membrane of the mitochondria was visualized with

127   mCherry-mito, which tagged the transmembrane domain of Miro with mCherry (Vagnoni and Bullock,

128   2016). Confocal microscopy revealed that the HA signals colocalized with mCherry-mito in the axon

129   terminals of photoreceptor cells (Figure 1B). This indicated that dOPA1 colocalizes with mitochondria

130   in retinal axons.

131   To examine the impact of *dOPA1* on mitochondrial structure and density, we analyzed

132   mitochondria in *dOPA1* RNAi flies. The intensity of the mitochondrial signal was significantly

133   reduced in the retinal axons of these flies (Figure 1C and D; quantification in 1E). Previously, impaired

134   mitochondrial transport was reported in the LOF of *marf* (MFN 1/2 homolog), which is necessary for

135   the fusion of the outer membrane of mitochondria (Sandoval et al., 2014). Thus, we investigated if the

136   LOF of *dOPA1* may cause transport defects, which may be responsible for the reduced mitochondrial

137   density in retinal axons in *dOPA1* RNAi flies. To this end, we observed the photoreceptor neurons

138   after puparium formation at 24 h, which allowed us to simultaneously visualize the cell bodies and

139   axon terminals of the photoreceptors (Figure S1A). Both *dOPA1* knockdown and control flies showed

140   mitochondrial signals in the axon terminals (Figure S1B and C). However, no mitochondrial signals

141   were observed in the axon terminals of flies with a knockdown of *milton*, which is thought to act as

142   an adaptor between kinesin to mitochondria, promoting anterograde transport (Figure S1D). These

143   findings suggest that trafficking defects are not the main cause of the mitochondrial mislocalization

144   observed in the retinal axons of the *dOPA1* knockdown.

145   To further analyze the mitochondrial morphology of retinal axons in the LOF of *dOPA1*, we

146   conducted an electron microscopy analysis (Figure 1F and G). Although *Drosophila* has eight types

of photoreceptors, the axons of R1–6, which project to the lamina of the first optic lobe, can be

identified without markers; thus, we chose them for this experiment. The *dOPA1* knockdown led to a

significant decrease in mitochondria size in R1–6 axons (Figure 1F–H). In addition, there was

significant heterogeneity in healthy mitochondria, ranging from dense-matrix (Class A) to swollen

mitochondria with a hypodense matrix (Class B) (Figure 1I). Knocking down *dOpa1* in R1–6 axon

terminals resulted in severely abnormal mitochondrial ultrastructure (22.8%; Figure 1J). These results

suggest that *dOPA1* is involved in maintaining proper mitochondrial morphology by promoting

mitochondrial fusion in retinal axons.

To verify whether ROS is increased in the retinal axons in *dOPA1* knockdown photoreceptors, we

measured the ROS levels in the knockdown retinal axons using MitoSOX, a mitochondrial superoxide

indicator (Yarosh et al., 2008). The results showed significantly elevated ROS levels in *dOPA1*

knockdown retinal axons compared to control (Figure 2A and B; quantification in 2C).

Given that an increase in mitophagy activity has been reported in mouse RGCs and nematode

ADOA models (Zaninello et al., 2022, 2020), the mitoQC marker, an established indicator of

mitophagy activity, was expressed in the photoreceptors of *Drosophila*. The mito-QC reporter consists

of a tandem mCherry-GFP tag that localizes to the outer membrane of mitochondria (Lee et al., 2018).

This construct allows the measurement of mitophagy by detecting an increase in the red-only mCherry

signal when the GFP is degraded after mitochondria are transported to lysosomes.

Post *dOPA1* knockdown, we observed a significant elevation in mCherry positive and GFP negative

puncta signals at one week, demonstrating an activation of mitophagy as a consequence

of *dOPA1* knockdown (Figure 2D–H).

The *dOPA1* LOF leads to progressive distal degeneration of *Drosophila* photoreceptors

We tested whether knocking down *dOPA1* causes optic nerve degeneration in *Drosophila*. To this end,

we performed an RNAi experiment using the *GMR-Gal4* driver and evaluated the number of retinal

axons projecting to the second optic ganglion medulla (Figure 3A). To compare the number of retinal

axon terminals–R7 axons–between genotypes, we used a previously developed automated method,

MeDUsA (method for the quantification of degeneration using fly axons) (Nitta et al., 2023). We have

assessed the extent of their reduction from the total axonal terminal count, thereby determining the

degree of axonal terminal degeneration (Nitta et al., 2023; Richard et al., 2022). Our results showed

that the *dOPA1* knockdown caused axonal degeneration one day after eclosion, further decreasing one

week later (Figure 3B–E, quantification in 3F). We obtained similar results with an independent

alternative RNAi line (Figure S2A and B). To determine whether the degeneration was limited to the

axons, we counted the number of rhabdomeres in the cell bodies. Each compound eye comprises 700-

800 ommatidia, with each ommatidium containing eight types of photoreceptors, designated as R1–

R8. The distal region of an ommatidium reveals the R1–R7 photoreceptors with a stereotypical

arrangement of rhabdomeres (arrowheads in Figure 3G–J). Since it is easy to evaluate retinal

degeneration by counting the number of rhabdomeres, a decrease in their number indicates

degeneration of the photoreceptor cell body. The *dOPA1* knockdown caused a significant decrease in

the number of rhabdomeres per ommatidium in 1-week-old adults (Figure 3G–J, quantification in 3K).

Interestingly, although the retinal axons had already decreased on the day of eclosion, the ommatidia

remained intact (Figure 3F and K). Note that the ommatidia in R7 remained intact after one week

(Figure 3J). Our results indicate that the *dOPA1* knockdown causes degeneration in both axonal

terminals and cell bodies, with the neuronal degeneration starting from the axons and continuing in

the cell bodies. A previous report indicated that the compound eyes of homozygous mutations of

dOPA1 displayed a glossy eye phenotype (Yarosh et al., 2008). Upon knocking down *dOPA1* using the

GMR-Gal4 driver, we also observed a glossy eye-like rough eye phenotype in the compound eyes

(Figure S3). The *GMR-Gal4* driver does not exclusively target Gal4 expression to photoreceptor cells.

Consequently, the observed retinal axonal degeneration could potentially be secondary to

abnormalities in support cells external to the photoreceptors.

To test whether our RNAi results reflect *dOPA1* downregulation, we performed mutant analysis of

the *dOPA1* gene in retinal axons. For this, we used the *dOPA1^{s3475}* hypomorphic mutant allele, which

has a P-element insertion in exon 2 (Yarosh et al., 2008). Somatic mosaic clones were generated in the

retinal axons using the FLP/FRT system (Golic and Lindquist, 1989) because homozygous mutant

alleles are embryonic lethal (Sandoval et al., 2014; Yarosh et al., 2008). Note that the mutant clone

analysis was conducted in a context where mutant and heterozygous cells coexist as a mosaic, and it

was not possible to distinguish between them. Using MeDUsA, we found a significantly lower number

of *dOPA1* mutant retinal axons than in controls one day after eclosion (Figure 4A and B, quantification

in 4D). To determine if *dOPA1* is responsible for axon neurodegeneration, we quantify the number of

the axons in the *dOPA1* eye clone fly with the expression of *dOPA1* at one day after eclosion and found

that *dOPA1* expression partially rescued the axonal degeneration (Figure 4C, quantification in 4D).

Our results indicate that *dOPA1* is necessary for maintaining the number of retinal axons in the

Drosophila visual system. In conclusion, the neurodegeneration observed in the LOF of *dOPA1* is due

to the progressive loss of retinal axons, as well as the mammal optic nerve.

Investigating disease mutations in OPA1 using a fly DOA model to confirm their pathological

significance

Our observations in *dOPA1* LOF flies confirmed mitochondria fragmentation, increased ROS levels,

and degeneration of retinal axons, as reported in mammals. These data indicated that the function of

OPA1 may be conserved across *Drosophila* and humans. Thus, we tested whether *hOPA1* could

effectively replace *dOPA1* in flies by generating a transgenic organism to express *hOPA1* in

Drosophila. Following the UAS sequence, that the yeast-derived transcription factor Gal4 binds, the

construct included an HA tag, followed by the *hOPA1* gene, and a myc tag (Figure 5A). This construct

was expressed in *Drosophila* and protein expression was confirmed by Western blotting. The HA tag,

attached to the N-terminus, was immunoblotted resulting in a band of the expected full-length size

(Figure 5B). For the myc tag at the C-terminus, a strong band and two weak bands were observed at

the upper and lower positions (Figure 5B). Comparison of the expression levels across the variants

revealed no significant differences in protein expression (Figure 5C). We also employed a human anti-

OPA1 antibody to verify the expression of hOPA1 in vivo in *Drosophila*, detecting both the long and

short forms of OPA1, band 2 and band 4 respectively (Figure 5D).

Using this *UAS-hOPA1*, we performed rescue experiments. *hOPA1* expression in the retinal axons

of the *dOPA1* mutant, which generated somatic cell clones, partially rescued the number of axons

(Figure 5E). These results suggest that *hOPA1* and *dOPA1* are interchangeable. We also generated

and expressed the *hOPA1* mutation 2708-2711del, which is known to cause DOA, the I382M mutation,

located in the GTPase domain and associated with DOA, as well as the D438V and R445H mutations,

also located in the GTPase domain and associated with DOA plus. The expression of these mutations

in retinal axons failed to rescue the *dOPA1* deficiency to the same extent as the WT *hOPA1* (Figure

5E). Importantly, unlike the D438V and R445H mutations, the 2708-2711del and I382M mutations

could be weakly rescued. This suggests that these mutations result in LOF, but may retain some

residual activity. These findings are also in line with the observed severity of the associated diseases,

DOA and DOA plus. As the D438V and R445H mutations were not significantly rescued, unlike the

2708-2711del and I382M mutations. Our methodology distinctively facilitates the quantitative

evaluation of LOF severity by comparing the rescue capabilities of various mutations. Notably, the

2708-2711del and I382M mutations demonstrated only partial rescue, indicative of a hypomorphic

effect with residual activity. In contrast, the D438V and R445H mutations failed to show significant

rescue, suggesting a more profound LOF. The correlation between the partial rescue by the 2708-

2711del and I382M mutations and their classification as hypomorphic is significant. Moreover, the

observed differences in rescue efficacy correspond to the clinical severities associated with these

mutations, namely in DOA and DOA plus disorders. Thus, our results substantiate the model's ability

to quantitatively discriminate among mutations based on their impact on protein functionality,

providing an insightful measure of LOF magnitude.

Furthermore, we assessed the potential for rescuing ROS signals. Similar to its effect on axonal

degeneration, wild-type hOPA1 effectively mitigated the phenotype, whereas the 2708-2711del,

D438V, and R445H mutants did not (Figure 5F). Importantly, the I382M variant also reduced ROS

levels comparably to the wild type. These findings demonstrate that both axonal degeneration and the

increase in ROS caused by dOPA1 downregulation can be effectively counteracted by hOPA1.

Although I382M retains partial functionality, it acts as a relatively weak hypomorph in this

experimental setup.

We also conducted Western blot analyses using anti-COXII and anti-Atg8a antibodies to assess

changes in mitochondrial quantity and autophagy activity following the knockdown of *dOPA1*.

Mitochondrial protein levels, indicated by COXII quantification, were evaluated to verify

mitochondrial content, and the ratio of Atg8a-1 to Atg8a-2 was used to measure autophagy activation.

For these experiments, *Tub-Gal4* was employed to systemically knockdown *dOPA1*. Considering the

lethality of a whole-body *dOPA1* knockdown, *Tub-Gal80^{TS}* was utilized to repress the knockdown

until eclosion by maintaining the flies at 20°C. After eclosion, we increased the temperature to 29°C

for two weeks to induce the knockdown or expression of *hOPA1* variants. The results revealed no

significant differences across the genotypes tested (Figure S4A–D).

Distinction between LOF and DN mutations in *hOPA1* linked to DOA or DOA plus

Currently, the *hOPA1* mutations that contribute to DOA plus have been primarily located in the

GTPase domain of *hOPA1*. The role of *hOPA1* as a GTPase is facilitated by its ability to form a

polymer via the GED domain and Middle domain (Li et al., 2019). This led to the hypothesis that

mutations in the GTPase domain can interact with WT hOPA1 but cannot activate the GTPase activity

to show a DN effect. However, it has been challenging to clarify whether these mutations are DN or

LOF, as there is a significant number of LOF mutations in the GTPase domain. Given that *dOPA1* can

be substituted with *hOPA1*, we suspected that a DN effect could also be achieved by expressing the

D438V or R445H mutations in the WT background. However, the axons did not degenerate despite

expression and monitoring for two weeks in adult flies (Fig. 6A). The results presented in Figure 5C

indicate that there are no significant differences in the expression levels among the variants, suggesting

that variations in expression levels do not influence the outcomes. The amino acid sequences of hOPA1

and dOPA1 are highly conserved (72% identity), particularly in the GTPase domain. The Middle and

GED domains are also relatively well-preserved, with 54% and 64% agreement, respectively (Figure

S5). These results imply that, although *dOPA1* and *hOPA1* are functionally interchangeable, they may

not interact with each other. Consequently, we expressed *hOPA1* in the photoreceptor depleted of

dOPA1 by somatic clone; under these conditions, we further expressed *hOPA1* WT, 2708-2711del,

and mutations in the GTPase domains, including R445H, I382M, and D438V. This allowed *hOPA1*

to interact with itself and to verify the DN effect. As a result, the expression of the WT, 2708del, and

I382M did not result in any changes when rescued by *hOPA1* substitution (Figure 6B). However, the

rescue was significantly suppressed for D438V and R445H, which are known to cause DN; thus, the

DN effect could be replicated (Figure 6B). In conclusion, we established an experimental model that

can separate LOF and DN, the pathological significance of *hOPA1* mutations. We investigated the

impacts of dominant negative mutations on mitochondrial oxidation levels, mitochondrial quantity,

and autophagy activation levels; however, none of these parameters showed statistical significance

(Figure S6).

Discussion

In this study, we found that LOF of *dOPA1* in *Drosophila* can imitate human DOA, in which the optic

nerve degenerates. We were successful in reversing this degeneration by expressing *hOPA1* (Figure

6C). However, we could not rescue any previously reported mutations known to cause either DOA or

DOA plus except for I382M. In the context of rescuing the retinal axons of the *dOPA1* mutant by

expressing the WT of *hOPA1*, it was observed that only the DOA plus mutations suppressed the rescue

(Figure 6C). This allowed us to distinguish between LOF and DN mutations in *hOPA1*. The fly model

developed in this study will allow investigating new *hOPA1* mutations for DOA or DOA plus. We have

previously utilized MeDUsA to quantify axonal degeneration, applying this methodology extensively

to various neurological disorders. The robust adaptability of this experimental system is demonstrated

by its application in exploring a wide spectrum of genetic mutations associated with neurological

conditions, highlighting its broad utility in neurogenetic research. We identified a novel de novo

variant in Spliceosome Associated Factor 1, Recruiter of U4/U6.U5 Tri-SnRNP (*SART1*). The patient,

born at 37 weeks with a birth weight of 2934g, exhibited significant developmental delays, including

an inability to support head movement at 7 months, reliance on tube feeding, unresponsiveness to

visual stimuli, and development of infantile spasms with hypsarrhythmia, as evidenced by EEG

findings. Profound hearing loss and brain atrophy were confirmed through MRI imaging. To assess

the functional impact of this novel human gene variant, we engineered transgenic Drosophila lines

expressing both wild type and mutant *SART1* under the control of a UAS promoter. Our MeDUsA

analysis suggested that the variant may confer a gain-of-toxic-function (Nitta et al., 2023). Moreover,

we identified heterozygous loss-of-function mutations in *DHX9* as potentially causative for a newly

characterized neurodevelopmental disorder. We further investigated the pathogenic potential of a novel

heterozygous de novo missense mutation in *DHX9* in a patient presenting with short stature,

intellectual disability, and myocardial compaction. Our findings indicated a loss of function in the

G414R and R1052Q variants of *DHX9* (Yamada et al., 2023). This experimental framework has been

instrumental in elucidating the impact of gene mutations, enhancing our ability to diagnose how novel

variants influence gene function. Our research established that *dOPA1* knockdown precipitates

axonal degeneration and elevates ROS signals in retinal axons. Expression of human *OPA1* within

this context effectively mitigated both phenomena; it partially reversed axonal degeneration and

nearly completely normalized ROS levels. These results imply that factors other than increased

ROS may drive the axonal degeneration observed post-knockdown. Furthermore, while

differences between the impacts of DN mutations and loss-of-function mutations were evident in

axonal degeneration, they were less apparent when using ROS as a biomarker. The extensive use

of transgenes in our experiments might have mitigated the knockdown effects. In a systemic

dOPA1 knockdown, assessments of mitochondrial quantity and autophagy activity revealed no

significant changes, suggesting that the cellular consequences of reduced *OPA1* expression might

vary across different cell types.

In our study, expressing *dOPA1* in the retinal axons of *dOPA1* mutants resulted in significant

rescue, but it did not return to control levels. There are three possible explanations for this result. The

first concerns gene expression levels. The Gal4-line used for the rescue experiments may not replicate

the expression levels or timing of endogenous *dOPA1*. Considering that the optimal functionality

of *dOPA1* may be contingent upon specific gene expression levels, attaining a wild-type-like state

necessitates the precise regulation of these expression levels. The second is a non-autonomous issue.

Although *dOPA1* gene expression was induced in the retinal axons for the rescue experiments, many

retinal axons were homozygous mutants, while other cell types were heterozygous for

the *dOPA1* mutation. If there is a non-autonomous effect of *dOPA1* in cells other than retinal axons, it

might not be possible to restore the wild-type-like state fully. The third potential issue is that only one

isoform of *dOPA1* was expressed. In mouse OPA1, to completely restore mitochondrial network shape,

an appropriate balance of at least two different isoforms, l-OPA1 and s-OPA1, is required (Del Dotto

et al., 2017). This requirement implies that multiple isoforms of *dOPA1* are essential for the dynamic

activities of mitochondria.

Our established fly model is the first simple organism to allow observation of degeneration of the

retinal axons. The mitochondria in the axons showed fragmentation of mitochondria. Former studies

have observed mitochondrial fragmentation in S2 cells (McQuibban et al., 2006), muscle tissue (Deng

et al., 2008), segmental nerves (Trevisan et al., 2018), and ommatidia (Yarosh et al., 2008) due to the

LOF of *dOPA1*. Our study presents compelling evidence that *dOPA1* knockdown instigates neuronal

degeneration, characterized by a sequential deterioration at the axonal terminals and extending to the

cell bodies. This degenerative pattern, commencing from the distal axons and progressing proximally

towards the cell soma, aligns with the paradigm of 'dying-back' neuropathy, a phenomenon extensively

documented in various neurodegenerative disorders (Wang et al., 2012). Previously, the function of *OPA1* in mice (*mOPA1*) was substituted by *hOPA1* (Del Dotto et al., 2017; Sarzi et al., 2018). Here we demonstrated that the function of *dOPA1* can also be substituted by *hOPA1*. Thus, the function of *OPA1* is conserved across a wide range of species but our fly model has the advantage of allowing rapid observation of phenotypes. In the mouse model of DOA with the heterozygous deletion of the *mOPA1* gene at positions 329–355 (Alavi et al., 2007) or delTTAG mutant (Sarzi et al., 2012), a decrease in the number of RGC was observed after 17 months or 16 months of age, respectively. To save time, using a fly model for quick analysis and direct observation of retinal axons would be useful. Previous models using nematodes, flies, and zebrafish did not observe axonal degeneration directly, but our new model has made this possible. As a result, our model has the potential to be used for screening for modifiers of unclear molecular pathologies and drug screening.

Our model could also demonstrate the pathological significance of *hOPA1* mutations and be used for genetic diagnosis to differentiate between LOF and DN mutations. To elucidate the pathophysiological implications of mutations in the *OPA1* gene, we engineered and expressed several human *OPA1* variants, including the 2708-2711del mutation, associated with DOA, and the I382M mutation, located in the GTPase domain and linked to DOA. We also investigated the D438V and R445H mutations in the GTPase domain and correlated with the more severe DOA plus phenotype. The 2708-2711del mutation exhibited limited detectability via HA-tag probing. Still, it was undetectable with a myc tag, likely due to a frameshift event leading to the mutation's characteristic truncated protein product, as delineated in prior studies (Zanna et al., 2008). Contrastingly, the I382M, D438V, and R445H mutations demonstrated expression levels comparable to the WT *hOPA1*.

However, the expression of these mutants in retinal axons did not restore the *dOPA1* deficiency to the same extent as the WT hOPA1, as evidenced in Figure 5E. This finding indicates a functional impairment imparted by these mutations, aligning with established understanding (Zanna et al., 2008). Notably, while the 2708-2711del and I382M mutations exhibited limited functional rescue, the D438V and R445H mutations did not show significant rescue activity. This differential rescue efficiency

suggests that the former mutations, particularly the I382M, categorized as a hypomorph (Del Dotto et

al., 2018), may retain partial functional capacity, indicative of a LOF effect but with residual activity.

The I382M missense mutation within the GTPase domain of OPA1 has been described as a mild

hypomorph or a disease modifier. Intriguingly, this mutation alone does not induce significant clinical

outcomes, as evidenced by multiple studies (Bonifert et al., 2014; Bonneau et al., 2014; Carelli et al.,

2015; Schaaf et al., 2011). A significant reduction in protein levels has been observed in fibroblasts

originating from patients harboring the I382M mutation. However, mitochondrial volume remains

unaffected, and the fusion activity of mitochondria is only minimally influenced (Del Dotto et al.,

2018; Kane et al., 2017). This observation is consistent with findings reported by de la Barca et al.

(Chao De La Barca et al., 2020), where a targeted metabolomics approach classified I382M as a mild

hypomorph. In our current study, the I382M mutation preserves more OPA1 function compared to DN

mutations, as depicted in Figures 5E and F. Considering the results from our *Drosophila* model and

previous research, we hypothesize that the I382M mutation may constitute a mild hypomorphic variant.

This might explain its failure to manifest a phenotype on its own, yet its contribution to increased

severity when it occurs in compound heterozygosity.

Our results for LOF or DN segmentation were consistent with previously reported yeast models

(Del Dotto et al., 2018) and fibroblasts obtained from patients with OPA1 mutations (Kane et al., 2017),

and confirmed the conserved function of *OPA1* across species. A major advantage of the *Drosophila*

model over yeast or fibroblasts, is that we can observe a phenotype that mimics the actual disease state

of axonal degeneration in retinal axons and analyze it not only at the molecular level, but also the

structural level of axons and mitochondria. The transgenic *Drosophila* of disease-associated *hOpa1*

mutations created in this study can also serve to understand the pathophysiology of these mutations.

Advances in genome analysis allowed the identification of many variants and genetic mutations

associated with DOA using next-generation sequencing. This could confirm diagnoses and identify

new forms of DOA. The widespread use of these sequencers has also created new challenges, such as

the growing number of variants of unknown significance (VUS) in publicly available databases such

as CliniVar for DOA. These variants are categorized as pathogenic, likely pathogenic, VUS, likely

benign, or benign, and the number of VUS is increasing as more variants are registered. The ClinVar

database had classified 110 out of 319 *Opa1* variants as VUS at the end of 2019. By the end of 2022,

this number had increased to 357 out of 840, indicating an increase from 34% to 42% over the last

three years (Henrie et al., 2018). This trend suggests that the number of VUS in *Opa1* may continue

to increase in the future. Analyzing VUS requires testing the effects of a mutation *in vitro* or *in vivo*;

this can be expensive and time-consuming without prior confirmation or high probability of

pathological significance. However, our fly model can be used to easily and inexpensively determine

the pathological effects of disease-causing gene mutations among the growing number of rare

mutations identified in DOA.

In this study, we tested the effect of expressing the D438V and R445H mutations in the *dOPA1*

background and found that retinal axons did not degenerate. Regarding the interactions among OPA1

proteins, in yeast Mgm1, K854 in the region near the C-terminus is important for aggregation, and it

becomes an oligomer to increase the activity of GTPase (Rujiviphat et al., 2009). Although hOPA1

normally has low GTPase activity, its interaction with negatively charged phospholipids such as

cardiolipin causes hOPA1 to aggregate, increasing GTP hydrolysis activity (Ban et al., 2010). In the

thermophilic fungus *Chaetomium thermophilum*, Mgm1 forms a dimer and then a tetramer at the C-

terminal Stalk domain of Mgm1 (Faelber et al., 2019). Structural analysis also predicted that the

Middle and GED domains are necessary for dimer and oligomer formation in hOPA1 (Li et al., 2019).

In addition, peptide-binding assays have revealed that the coiled-coil domain, which is part of the GED

domain, is required for hOPA1 interactions (Akepati et al., 2008). Comparing the amino acid

sequences of hOPA1 and dOPA1, the Middle and GED domains, which are necessary for interactions

between OPA1 molecules, exhibit concordance rates of 54.3% and 63.6%, respectively. These

percentages are low compared to the 71.8% identity observed in the GTPase domains. These suggest

that the OPA1 homologs of each species form oligomers with their unique OPA1. Note that the

expression of *dOPA1^{K273A}*, a presumably GTPase-negative form of *dOPA1* (Tsuyama et al., 2017),

degenerated retinal axons (Figure S7). These results imply that, while dOPA1 and hOPA1 are

interchangeable, they may not interact with each other. This could pave the way for the development

of a chimeric hOPA1, which would retain its functional properties while avoiding the interaction with

endogenous hOPA1. Such chimeric hOPA1 could potentially be used as gene therapy since induced

pluripotent stem cells have been generated from a patient with R445H (Sladen et al., 2021).

OPA1 has been implicated in various other diseases as well, including normal tension glaucoma

(Powell et al., 2003; Yu-Wai-Man et al., 2010b), multiple sclerosis-like illness (Verny et al., 2008;

Yu-Wai-Man et al., 2016), Parkinson's disease and dementia (Carelli et al., 2015; Lynch et al., 2017),

and cardiomyopathy (Spiegel et al., 2016). Although the connection between these diseases and OPA1

is not fully understood, further research on DOA may provide a deeper insight into this relationship.

In the future, pathological analysis using the present model could have implications for understanding

the mechanisms underlying these diseases.

DOA has also been reported to involve mitochondrial DNA (mtDNA) depletion (Amati-Bonneau

et al., 2008), but our model, while useful for nuclear gene analysis, is limited for mtDNA analysis.

The gene content in the *Drosophila* mtDNA genome is similar as that in vertebrates, but the gene order

and distribution on both DNA strands differ. mOPA1 interacts with mtDNA nucleoids through exon

4b binding to mtDNA D-loops, independent of mitochondrial fusion (Yang et al., 2020). However,

the homologous region of mOPA1 or hOPA1 to exon4b is not found in *Drosophila*, and the D-loop of

mtDNA has not been found in cultured cells of flies or in mtDNA of embryos using electron

microscopy (Rubenstein et al., 1977). In addition, while the D-loop is involved in mtDNA replication

in humans (Fish et al., 2004), the A + T region is the origin of replication in mtDNA replication in

Drosophila (Goddard and Wolstenholme, 1978). Thus, there may be differences in the mechanism of

mtDNA replication between humans and flies (Garesse and Kaguni, 2005). Therefore, the association

of dOPA1 with mtDNA may differ from that in mammals and the regulatory mechanism of mtDNA

homeostasis in which dOPA1 is involved requires further investigation.

454 Another limitation of our model is that the molecular mechanisms underlying the formation of

455 long-form OPA1 (L-OPA1) and short-form OPA1 (S-OPA1), which are well analyzed in mammals

456 (MacVicar and Langer, 2016), have not been elucidated in *Drosophila*. In hOPA1, S1 or S2 sites are

457 processed by the i-AAA protease OMA1 and Yme1L respectively to produce S-hOPA1 (Anand et al.,

458 2014). Despite the fact that Yme1L is conserved in *Drosophila*, it has been involved in the degradation

459 of the dOPA1 protein rather than its cleavage by dYme1L (Liu et al., 2020). Moreover, the cleavage

460 site for i-AAA does not exist in dOPA1 (Olichon et al., 2007). Regarding other proteases, yeast Mgm1

461 is processed by the rhomboid protease Pcp1 (Esser et al., 2002; Herlan et al., 2003). However, the

462 Rhomboid-7, a Pcp1 ortholog of *Drosophila*, was not required for dOPA1 processing (Whitworth et

463 al., 2008). Although hOPA1 expression in *Drosophila* has identified three different sizes of hOPA1 in

464 our Western blotting result, it is unclear how they are cleaved. From a size perspective, the upper band

465 was inferred to represent the full-length hOPA1 including the mitochondria import sequence (MIS).

466 Since the Middle and the lowest bands were not present in HA-tagged samples, it is possible that the

467 bands at the Middle size represent the long-form hOPA1 (L-hOPA1) in which probably MIS was

468 processed. The band detected at the lowest position may represent a short form of hOPA1 (S-hOPA1).

469 However, *Drosophila* does not have OMA1, an i-AAA protease that cleaves the S1 site. Although

470 another i-AAA protease that cleaves the S2 site, YME1L, is conserved, our expressed hOPA1 is

471 isoform 1 and lacks the S2 site (Fig. 5A). Therefore, it is not clear how S-hOPA1 is processed in

472 *Drosophila*. The anti-dOPA1 antiserum detects three bands in the overexpression of a FLAG-tagged

473 dOPA1 construct (Poole et al., 2010), suggesting a molecular mechanism by which dOPA1, like yeast

474 and mammals, is cleaved after translation. However, the proteases involved in the mechanism are not

475 yet clarified. Whether *Drosophila's* endogenous mechanisms can be used to study how L-OPA1 and

476 S-OPA1 are involved in DOA remains unclear.

477 In this study, we established a new model of DOA in *Drosophila*, in which we discovered the

478 following: 1. We could replicate the human disease of optic nerve degeneration, allowing rapid genetic

479 disease analysis. 2. We could distinguish the LOF and DN effects of *hOPA1* by observation of axonal

480 degeneration phenotype but not with ROS signal, autophagy activity, and amount of the mitochondria

in this model. These findings can reveal the pathological significance of *de novo* mutations and are

useful in diagnosing DOA or DOA plus. Additionally, while *hOPA1* is the major gene involved in

DOA, other genes involved in DOA and interacting proteins have not been investigated yet. Our model

can be used for pathological modifier screening, exploring modifiers, and contributing to drug

development. In the future, we hope to provide insights that can be applied to the fundamental

treatment of DOA.

Materials and Methods

Fly strains

Flies were maintained at 25°C on standard fly food. For knockdown experiments (Figures 1C–E,

1F–J, 2A–H, 3B–K, 5F, S1, S2, S6A, and S7), flies were kept at 29°C in darkness. Female flies

were used in all experiments to ensure consistency in the number of retinal axons. The following fly

strains were obtained from the Bloomington Drosophila Stock Center (BDSC): *GMR-Gal4* (BDSC

1104), *Tub-Gal4* (BDSC 5138), *UAS-dicer2(III)* (BDSC 24651), *UAS-mito-HA-GFP* (BDSC 8443),

UAS-mCherry-mito (BDSC 66532), *UAS-mitoQC* (BDSC 91641), *UAS-dOPA1 RNAi* (BDSC 32358),

UAS-Milton RNAi (BDSC 44477), *FRT42D, GMR-hid, l(2)cl-R11[1]/CyO; ey-Gal4, UAS-flp* (BDSC

5251), *ey3.5-flp* (BDSC 35542), *Tub-Gal80^{TS}* (BDSC 7017), and *UAS-luciferase RNAi* (BDSC 31603).

The following fly strains were obtained from the Vienna Drosophila Resource Center (VDRC): The

40D-UAS (VDRC 60101) and *UAS-dOPA1 RNAi* (VDRC 106290). The following fly strains were

obtained from the Kyoto Drosophila Stock Center (Kyoto): *FRT19* (Kyoto 101231), *FRT42D* (Kyoto

101878), and *dOPA1^{s3475}* (Kyoto 111438). The *UAS-dOPA1-HA* and *UAS-dOPA1-HA^{K273A}* were

generously provided by T. Uemura (Kyoto University, Japan). The *dOPA1* clone analysis was

performed by inducing flippase expression in the eyes using either *ey-Gal4* with *UAS-flp* or *ey3.5-flp*,

followed by recombination at the chromosomal location *FRT42D* to generate a mosaic of cells

homozygous for *dOPA1^{s3475}*.

Generation of transgenic flies

The cDNA for WT, I382M, D438V, R445H, and 2708-2711del forms of *hOPA1* [NM_015560.3] were

generated by Vectorbuilder Inc. (Yokohama, Japan). Then, the HA-hOPA1-myc sequence was

amplified and inserted into pJFRC81-10XUAS-IVS-Syn21-GFP-p10 (ID36432, Addgene, USA)

digested with *NotI* and *XbaI* using primers with the N-terminal containing the Kozak and HA

sequences (TTACTTCAGGCGGCCGCGGCCAAAATGTACCCATACGATGTTCCAGATTAC) and

the   C-terminal   containing   the   myc   sequence

(TTAAAAACGATTCATTCTAGTTACAGATCCTCTTCTGAGATGAGTTTTTGTTCTTTCTCCT

GATGAAGAGCTTCAATG). To confirm production of the complete coding sequence, it was

bidirectionally sequenced using the Sanger method. These plasmids were injected into embryos and

integrated into the attP2 landing site via ΦC31 recombination (WellGenetics, Taiwan).

Immunohistochemistry and imaging

Immunohistochemistry was performed as described previously (Sugie et al., 2017).   Flies were

anesthetized on a CO_2 pad, and specimens of the correct genotype were selected. Specimens were

briefly washed in ethanol for a few seconds before being transferred to a Petri dish containing

phosphate-buffered saline (PBS). Using forceps, the proboscis was secured, and the right and left eyes

were dissected out with another set of forceps. The proboscis and attached trachea were also removed

from the brain. The brains were then carefully transferred with forceps into a 1.5 mL tube containing

150 μL of 0.3% PBT (PBS with Triton-X100) at room temperature.

To fix the tissues, 50 μL of 16% formaldehyde was added to each tube, followed by gentle mixing.

The samples were incubated in the fixative for 50 minutes at room temperature. Subsequent washing

involved three rinses with 200 μL of 0.3% PBT, using a micropipette for fluid exchange. After the

final wash, the PBT was removed, and the samples were incubated in the primary antibody solution

overnight (O/N) at 4 °C. This was followed by three washes with 0.3% PBT, after which the secondary

antibody solution (diluted in 0.3% PBT) was added, and samples were incubated overnight in the dark

at 4 °C. After another three washes in 0.3% PBT, the samples were prepared for imaging. The

following antibodies were used: rat anti-HA (3F10, 1:400; Roche, Switzer land), mouse anti-Chaoptin

(24B10, 1:25; Developmental Studies Hybridoma Bank, USA), rat anti-CadN (DN-Ex#8, 1:50;

Developmental Studies Hybridoma Bank), anti-rat Alexa Fluor 488 (1:400; Thermo Fisher Scientific,

USA), anti-mouse Alexa Fluor 488 (1:400; Thermo Fisher Scientific), anti-mouse Alexa Fluor 568

(1:400; Thermo Fisher Scientific), and anti-rat Alexa Fluor 633 (1:400; Thermo Fisher Scientific).

Specimens were mounted using a Vectashield mounting medium (Vector Laboratories, USA). Images

were captured using an FV3000 confocal microscope (Olympus, Japan) or a C2 (Nikon, Japan)

confocal microscope for Figure S1 and processed using the IMARIS software (Oxford Instruments,

UK) or Fiji software, an open-source image analysis software (Schindelin et al., 2012).

Electron microscopy

Fly heads were fixed overnight with 2.5% glutaraldehyde and 2% paraformaldehyde in 0.1 M sodium

cacodylate buffer. Heads were then post-fixed with 1% osmium tetroxide, dehydrated in ethanol, and

embedded in Epon. Ultra-thin tangential sections of the laminas (70 nm) were stained with uranyl

acetate and lead citrate. Random retinas or lamina sections were imaged at a magnification of ×100 k

with a VELETA CCD Camera (Olympus) mounted on a JEM 1010 transmission electron microscope

(JEOL, Japan). In the R1–6 axons of the lamina, the areas corresponding to mitochondria were

quantified using the freehand line tool in NIS Elements software (Nikon). Concurrently, mitochondria

were manually categorized based on their structural characteristics: those with a densely packed matrix

were classified as Class A, and those with a collapsed structure as Class B.

Quantification of Mito-GFP intensity in retinal axons and retina

To measure the relative intensity of Mito-GFP in the photoreceptor R7/8 axons of the medulla, the imaging analysis software IMARIS was used. First, the M1–6 layers of the medulla, including the R7/8 axons, were manually covered using the surface function. Then, the surface area was masked, the surface of the left axons was stained with anti-Chaoptin, and the Mito-GFP signal was generated automatically using the surface function. Finally, the voxel number of the Mito-GFP surface was divided by the voxel number of the R axon surface. These quantifications were performed by experimenters blind to the genotype.

Mitochondrial superoxide detection and quantification

The ROS level was determined using the MitoSOX Red Mitochondrial Superoxide Indicator (Thermo Fisher Scientific). The procedure involved dissecting adult brains in phosphate-buffered saline (PBS), transferring them to 5 µM MitoSOX solution, and incubating them for 10 min in the dark at room temperature. The brains were then washed with PBS containing 20% Vectashield (Vector Laboratories), followed by sequential washes with PBS containing 40% and 60% Vectashield. The samples were mounted in Vectashield and scanned using an FV3000 confocal microscope (Olympus). To quantify the ROS level in photoreceptor R7/8 axons, the IMARIS software (Bitplane) was used for imaging analysis. The software generated automatic surfaces of the MitoSOX and Mito-GFP signals. Then, the ROS level was calculated by dividing the voxel number of the MitoSOX surface on the Mito-GFP surface by the voxel number of the Mito-GFP surface. These quantifications were performed by experimenters blind to the genotype.

Measurement of mitophagy levels in retinal axons

The mitophagy levels in the photoreceptor R7/8 axons of the medulla were analyzed using the mitoQC

reporter. Mitolysosomes were identified by referencing the red-only mCherry puncta of mitoQC. The

number of mitolysomes of each voxel image was manually quantified with IMARIS software

(Bitplane). These quantifications were performed by experimenters blind to the genotype.

Western blot analysis of the whole Drosophila body

We regulated protein expression temporally across the whole body using the *Tub-Gal4* and *Tub-*

GAL80^{TS} system. Flies harboring each *hOPA1* variant were maintained at a permissive

temperature of 20°C, and upon emergence, females were transferred to a restrictive temperature

of 29°C for subsequent experiments. The flies were collected in plastic tubes and sonicated (2 x 30

s) using a Q55 sonicator (Qsonica, Newtown, CT, USA) in lysis buffer (10 mM Tris–HCl, pH 7.5, 150

mM NaCl, 1 mM EDTA, 2% DDM: 10μl/mg flies) supplemented with 1:1000 (v/v) Protease Inhibitor

Cocktail Set III (Calbiochem, USA). Rat anti-HRP-conjugated-HA (3F10, 1: 3000; proteintech,

Germany), mouse anti-myc (9B11, 1: 10000; Cell Signaling Technology, USA), rabbit anti-OPA1

(ab42364, 1: 1000; Abcam, UK), rabbit anti-COXII (1: 100; Murari et al. *JCB* 2020), rabbit anti-Atg8a

(ab109364, 1: 8000; Abcam), mouse anti-α-Tubulin (T9026, 1:100,000; Sigma‑Aldrich, USA) ,

goat HRP-conjugated anti-mouse IgG (SA00001-1, 1: 10,000; proteintech), and goat HRP-conjugated

anti-rabbit IgG (7074S, 1: 5,000; Cell Signaling Technology) antibodies were used for Western

blotting.

Pairwise alignment of hOPA1 and dOPA1

The amino acid sequences for hOPA1 and dOPA1 were referenced from NM_015560.3 and

600 NM_166040.2, respectively, and aligned using the EMBOSS Needle (Rice et al., 2000). The GTPase,

601 Middle, and GED domains in dOPA1 were identified by cross-referencing the previously described

602 amino acid positions of these domains in hOPA1 (Liesa et al., 2009). The degree of identity between

603 the aligned amino acid sequences was subsequently determined.

605 ***Experimental design and statistical analyses***

606 Experimental analyses were performed using Prism 9 (GraphPad Software, USA). For Figures 1E, 1H,

607 2C, S2B and S7B, data were analyzed using t tests (and nonparametric tests) and Mann-Whitney tests.

608 For Figure 1J, Fisher's exact test was used. For Figures 2H, 3F, 3K, 4D, 5C, 5E, 5F, 6A, 6B, S3B, S4B,

609 S4D, S6C and S6E, differences among multiple groups were examined using the Kruskal−Wallis test

610 followed by the two-stage linear step-up procedure of Benjamini, Krieger, and Yekutieli tests. The null

611 hypothesis was rejected at a 0.05 level of significance. Sample sizes are indicated in each figure. The

612 number of retinal axons and rhabdomeres, the intensity of Mito-GFP and MitoSOX, the mitochondrial

613 area and the classification, and the corrected P values are described in the Results section. All data are

614 expressed as the mean ± SEM.

616 ***Acknowledgements***

617 We would like to thank Dr. Uemura for providing us with the transgenic fly strain and Dr. Owusu-

618 Ansah for gifting us the antibody. The Ministry of Education, Culture, Sports, Science, and Technology

619 of Japan (#18K14835, #18J00367 and #21K15619 to Y.N., #21K06184 to S.H.S., #16H06457,

620 #21H02483 and 21H05682 to T.S., #17H04983, #19K22592 and #21H02837 to A.S.), the Japan

621 Agency for Medical Research and Development (AMED) (#JP22ek019484s) to A.S., NIG–JOINT to

622 E.S., Takeda Science Foundation Takeda Visionary Research Grant to T.S., Takeda Science Foundation

623 Life Science Research Grant to A.S., and GSK Japan Research Grant 2022 [AS2021A000166849] to

A.S..

References

Akepati VR, Müller EC, Otto A, Strauss HM, Portwich M, Alexander C. 2008. Characterization of OPA1

isoforms isolated from mouse tissues. *J Neurochem* **106**:372–383. doi:10.1111/j.1471-

4159.2008.05401.x

Alavi M V., Bette S, Schimpf S, Schuettauf F, Schraermeyer U, Wehrl HF, Ruttiger L, Beck SC, Tonagel

F, Pichler BJ, Knipper M, Peters T, Laufs J, Wissinger B. 2007. A splice site mutation in the murine

Opa1 gene features pathology of autosomal dominant optic atrophy. *Brain* **130**:1029–1042.

doi:10.1093/brain/awm005

Alexander C, Votruba M, Pesch UEA, Thiselton DL, Mayer S, Moore A, Rodriguez M, Kellner U, Leo-

Kottler B, Auburger G, Bhattacharya SS, Wissinger B. 2000. OPA1, encoding a dynamin-related

GTPase, is mutated in autosomal dominant optic atrophy linked to chromosome 3q28. *Nat Genet*

26:211–215. doi:10.1038/79944

Amati-Bonneau P, Odent S, Derrien C, Pasquier L, Malthiéry Y, Reynier P, Bonneau D. 2003. The

association of autosomal dominant optic atrophy and moderate deafness may be due to the R445H

mutation in the OPA1 gene. *Am J Ophthalmol* **136**:1170–1171. doi:10.1016/S0002-9394(03)00665-

2

Amati-Bonneau P, Valentino ML, Reynier P, Gallardo ME, Bornstein B, Boissière A, Campos Y, Rivera

H, De La Aleja JG, Carroccia R, Iommarini L, Labauge P, Figarella-Branger D, Marcorelles P, Furby

A, Beauvais K, Letournel F, Liguori R, La Morgia C, Montagna P, Liguori M, Zanna C, Rugolo M,

Cossarizza A, Wissinger B, Verny C, Schwarzenbacher R, Martín MÁ, Arenas J, Ayuso C, Garesse

R, Lenaers G, Bonneau D, Carelli V. 2008. OPA1 mutations induce mitochondrial DNA instability

and optic atrophy "plus" phenotypes. *Brain* **131**:338–351. doi:10.1093/brain/awm298

Anand R, Wai T, Baker MJ, Kladt N, Schauss AC, Rugarli E, Langer T. 2014. The i-AAA protease YME1L

and OMA1 cleave OPA1 to balance mitochondrial fusion and fission. *J Cell Biol* **204**:919–929.

doi:10.1083/jcb.201308006

Ban T, Heymann JAW, Song Z, Hinshaw JE, Chan DC. 2010. OPA1 disease alleles causing dominant optic

atrophy have defects in cardiolipin-stimulated GTP hydrolysis and membrane tubulation. *Hum Mol*

Genet **19**:2113–2122. doi:10.1093/hmg/ddq088

Bonifert T, Karle KN, Tonagel F, Batra M, Wilhelm C, Theurer Y, Schoenfeld C, Kluba T, Kamenisch Y,

Carelli V, Wolf J, Gonzalez MA, Speziani F, Schüle R, Züchner S, Schöls L, Wissinger B, Synofzik

M. 2014. Pure and syndromic optic atrophy explained by deep intronic OPA1 mutations and an

intralocus modifier. *Brain* **137**:2164–2177. doi:10.1093/brain/awu165

Bonneau D, Colin E, Oca F, Ferre M, Chevrollier A, Guéguen N, Desquiret-Dumas V, N'Guyen S, Barth

M, Zanlonghi X, Rio M, Desguerre I, Barnerias C, Momtchilova M, Rodriguez D, Slama A, Lenaers

G, Procaccio V, Amati-Bonneau P, Reynier P. 2014. Letter to the editor: Early-onset Behr syndrome

due to compound heterozygous mutations in OPA1. *Brain* **137**. doi:10.1093/brain/awu184

Brand AH, Perrimon N. 1993. Targeted gene expression as a means of altering cell fates and generating

dominant phenotypes. *Development* **118**:401–415. doi:10.1242/dev.118.2.401

Byrne JJ, Soh MS, Chandhok G, Vijayaraghavan T, Teoh JS, Crawford S, Cobham AE, Yapa NMB, Mirth

CK, Neumann B. 2019. Disruption of mitochondrial dynamics affects behaviour and lifespan in

Caenorhabditis elegans. *Cell Mol Life Sci* **76**:1967–1985. doi:10.1007/s00018-019-03024-5

Carelli V, Musumeci O, Caporali L, Zanna C, La Morgia C, Del Dotto V, Porcelli AM, Rugolo M,

Valentino ML, Iommarini L, Maresca A, Barboni P, Carbonelli M, Trombetta C, Valente EM,

Patergnani S, Giorgi C, Pinton P, Rizzo G, Tonon C, Lodi R, Avoni P, Liguori R, Baruzzi A, Toscano

A, Zeviani M. 2015. Syndromic parkinsonism and dementia associated with OPA1 missense

mutations. *Ann Neurol* **78**:21–38. doi:10.1002/ana.24410

Chao De La Barca JM, Fogazza M, Rugolo M, Chupin S, Del Dotto V, Ghelli AM, Carelli V, Simard G,

Procaccio V, Bonneau D, Lenaers G, Reynier P, Zanna C. 2020. Metabolomics hallmarks OPA1

variants correlating with their in vitro phenotype and predicting clinical severity. *Hum Mol Genet*

29:1319–1329. doi:10.1093/hmg/ddaa047

Davies VJ, Hollins AJ, Piechota MJ, Yip W, Davies JR, White KE, Nicols PP, Boulton ME, Votruba M.

2007. Opa1 deficiency in a mouse model of autosomal dominant optic atrophy impairs mitochondrial
morphology, optic nerve structure and visual function. *Hum Mol Genet* **16**:1307–1318.
doi:10.1093/hmg/ddm079

Del Dotto V, Carelli V. 2021. Dominant Optic Atrophy (DOA): Modeling the Kaleidoscopic Roles of
OPA1 in Mitochondrial Homeostasis. *Front Neurol*. doi:10.3389/fneur.2021.681326

Del Dotto V, Fogazza M, Musiani F, Maresca A, Aleo SJ, Caporali L, La Morgia C, Nolli C, Lodi T,
Goffrini P, Chan D, Carelli V, Rugolo M, Baruffini E, Zanna C. 2018. Deciphering OPA1 mutations
pathogenicity by combined analysis of human, mouse and yeast cell models. *Biochim Biophys Acta -*
Mol Basis Dis **1864**:3496–3514. doi:10.1016/j.bbadis.2018.08.004

Del Dotto V, Mishra P, Vidoni S, Fogazza M, Maresca A, Caporali L, McCaffery JM, Cappelletti M,
Baruffini E, Lenaers G, Chan D, Rugolo M, Carelli V, Zanna C. 2017. OPA1 Isoforms in the
Hierarchical Organization of Mitochondrial Functions. *Cell Rep* **19**:2557–2571.
doi:10.1016/j.celrep.2017.05.073

Delettre C, Lenaers G, Griffoin JM, Gigarel N, Lorenzo C, Belenguer P, Pelloquin L, Grosgeorge J, Turc-
Carel C, Perret E, Astarie-Dequeker C, Lasquellec L, Arnaud B, Ducommun B, Kaplan J, Hamel CP.
2000. Nuclear gene OPA1, encoding a mitochondrial dynamin-related protein, is mutated in dominant
optic atrophy. *Nat Genet* **26**:207–210. doi:10.1038/79936

Deng H, Dodson MW, Huang H, Guo M. 2008. The Parkinson's disease genes pink1 and parkin promote
mitochondrial fission and/or inhibit fusion in Drosophila. *Proc Natl Acad Sci U S A* **105**:14503–
14508. doi:10.1073/pnas.0803998105

Eijkenboom I, Vanoevelen JM, Hoeijmakers JGJ, Wijnen I, Gerards M, Faber CG, Smeets HJM. 2019. A
zebrafish model to study small-fiber neuropathy reveals a potential role for GDAP1. *Mitochondrion*
47:273–281. doi:10.1016/j.mito.2019.01.002

Esser K, Tursun B, Ingenhoven M, Michaelis G, Pratje E. 2002. A novel two-step mechanism for removal
of a mitochondrial signal sequence involves the mAAA complex and the putative rhomboid protease
Pcp1. *J Mol Biol* **323**:835–843. doi:10.1016/S0022-2836(02)01000-8

Faelber K, Dietrich L, Noel JK, Wollweber F, Pfitzner AK, Mühleip A, Sánchez R, Kudryashev M,

Chiaruttini N, Lilie H, Schlegel J, Rosenbaum E, Hessenberger M, Matthaeus C, Kunz S, von der

Malsburg A, Noé F, Roux A, van der Laan M, Kühlbrandt W, Daumke O. 2019. Structure and

assembly of the mitochondrial membrane remodelling GTPase Mgm1. *Nature* **571**:429–433.

doi:10.1038/s41586-019-1372-3

Ferré M, Caignard A, Milea D, Leruez S, Cassereau J, Chevrollier A, Amati-Bonneau P, Verny C, Bonneau

D, Procaccio V, Reynier P. 2015. Improved locus-specific database for OPA1 mutations allows

inclusion of advanced clinical data. *Hum Mutat* **36**:20–25. doi:10.1002/humu.22703

Fish J, Raule N, Attardi G. 2004. Discovery of a major D-loop replication origin reveals two modes of

human mtDNA synthesis. *Science (80-)* **306**:2098–2101. doi:10.1126/science.1102077

Frezza C, Cipolat S, Martins de Brito O, Micaroni M, Beznoussenko G V., Rudka T, Bartoli D, Polishuck

RS, Danial NN, De Strooper B, Scorrano L. 2006. OPA1 Controls Apoptotic Cristae Remodeling

Independently from Mitochondrial Fusion. *Cell* **126**:177–189. doi:10.1016/j.cell.2006.06.025

Garesse R, Kaguni LS. 2005. A Drosophila model of mitochondrial DNA replication: Proteins, genes and

regulation. *IUBMB Life*. doi:10.1080/15216540500215572

Goddard JM, Wolstenholme DR. 1978. Origin and direction of replication in mitochondrial DNA molecules

from Drosophila melanogaster. *Proc Natl Acad Sci U S A* **75**:3886–3890. doi:10.1073/pnas.75.8.3886

Golic KG, Lindquist S. 1989. The FLP recombinase of yeast catalyzes site-specific recombination in the

drosophila genome. *Cell* **59**:499–509. doi:10.1016/0092-8674(89)90033-0

Henrie A, Hemphill SE, Ruiz-Schultz N, Cushman B, DiStefano MT, Azzariti D, Harrison SM, Rehm HL,

Eilbeck K. 2018. ClinVar Miner: Demonstrating utility of a Web-based tool for viewing and filtering

ClinVar data. *Hum Mutat* **39**:1051–1060. doi:10.1002/humu.23555

Herlan M, Vogel F, Bornhövd C, Neupert W, Reichert AS. 2003. Processing of Mgm1 by the rhomboid-

type protease Pcp1 is required for maintenance of mitochondrial morphology and of mitochondrial

DNA. *J Biol Chem* **278**:27781–27788. doi:10.1074/jbc.M211311200

Kanazawa T, Zappaterra MD, Hasegawa A, Wright AP, Newman-Smith ED, Buttle KF, McDonald K,

Mannella CA, Van Der Bliek AM. 2008. The C. elegans Opa1 homologue EAT-3 is essential for

resistance to free radicals. *PLoS Genet* **4**. doi:10.1371/journal.pgen.1000022

Kane MS, Alban J, Desquiret-Dumas V, Gueguen N, Ishak L, Ferre M, Amati-Bonneau P, Procaccio V,

Bonneau D, Lenaers G, Reynier P, Chevrollier A. 2017. Autophagy controls the pathogenicity of

OPA1 mutations in dominant optic atrophy. *J Cell Mol Med* **21**:2284–2297. doi:10.1111/jcmm.13149

Lee JJ, Sanchez-Martinez A, Zarate AM, Benincá C, Mayor U, Clague MJ, Whitworth AJ. 2018. Basal

mitophagy is widespread in Drosophila but minimally affected by loss of Pink1 or parkin. *J Cell Biol*

217:1613–1622. doi:10.1083/jcb.201801044

Lenaers G, Neutzner A, Le Dantec Y, Jüschke C, Xiao T, Decembrini S, Swirski S, Kieninger S, Agca C,

Kim US, Reynier P, Yu-Wai-Man P, Neidhardt J, Wissinger B. 2021. Dominant optic atrophy:

Culprit mitochondria in the optic nerve. *Prog Retin Eye Res*. doi:10.1016/j.preteyeres.2020.100935

Li D, Wang J, Jin Z, Zhang Z. 2019. Structural and evolutionary characteristics of dynamin-related GTPase

OPA1. *PeerJ* **2019**. doi:10.7717/peerj.7285

Liesa M, Palacín M, Zorzano A. 2009. Mitochondrial dynamics in mammalian health and disease. *Physiol*

Rev. doi:10.1152/physrev.00030.2008

Liu W, Duan X, Xu L, Shang W, Zhao J, Wang L, Li JC, Chen CH, Liu JP, Tong C. 2020. Chchd2 regulates

mitochondrial morphology by modulating the levels of Opa1. *Cell Death Differ* **27**:2014–2029.

doi:10.1038/s41418-019-0482-7

Lynch DS, Loh SHY, Harley J, Noyce AJ, Martins LM, Wood NW, Houlden H, Plun-Favreau H. 2017.

Clinical/Scientific Notes: Nonsyndromic Parkinson disease in a family with autosomal dominant

optic atrophy due to opa1 mutations. *Neurol Genet* **3**. doi:10.1212/NXG.0000000000000188

MacVicar T, Langer T. 2016. OPA1 processing in cell death and disease - the long and short of it. *J Cell*

Sci. doi:10.1242/jcs.159186

McQuibban GA, Lee JR, Zheng L, Juusola M, Freeman M. 2006. Normal Mitochondrial Dynamics

Requires Rhomboid-7 and Affects Drosophila Lifespan and Neuronal Function. *Curr Biol* **16**:982–

989. doi:10.1016/j.cub.2006.03.062

Meglei G, McQuibban GA. 2009. The dynamin-related protein Mgm1p assembles into oligomers and

hydrolyzes GTP to function in mitochondrial membrane fusion. *Biochemistry* **48**:1774–1784.

doi:10.1021/bi801723d

Newsome TP, Åsling B, Dickson BJ. 2000a. Analysis of Drosophila photoreceptor axon guidance in eye-
specific mosaics. *Development* **127**:851–860. doi:10.1242/dev.127.4.851

Newsome TP, Schmidt S, Dietzl G, Keleman K, Åsling B, Debant A, Dickson BJ. 2000b. Trio combines
with dock to regulate Pak activity during photoreceptor axon pathfinding in Drosophila. *Cell*
101:283–294. doi:10.1016/S0092-8674(00)80838-7

Nguyen D, Alavi M V., Kim KY, Kang T, Scott RT, Noh YH, Lindsey JD, Wissinger B, Ellisman MH,
Weinreb RN, Perkins GA, Ju WK. 2011. A new vicious cycle involving glutamate excitotoxicity,
oxidative stress and mitochondrial dynamics. *Cell Death Dis* **2**. doi:10.1038/cddis.2011.117

Nitta Y, Kawai H, Maki R, Osaka J, Hakeda-Suzuki S, Nagai Y, Doubková K, Uehara T, Watanabe K,
Kosaki K, Suzuki T, Tavosanis G, Sugie A. 2023.   Direct evaluation of neuroaxonal degeneration
with the causative genes of neurodegenerative diseases in Drosophila using the automated axon
quantification system, MeDUsA . *Hum Mol Genet*. doi:10.1093/hmg/ddac307

Olichon A, ElAchouri G, Baricault L, Delettre C, Belenguer P, Lenaers G. 2007. OPA1 alternate splicing
uncouples an evolutionary conserved function in mitochondrial fusion from a vertebrate restricted
function in apoptosis. *Cell Death Differ* **14**:682–692. doi:10.1038/sj.cdd.4402048

Olichon A, Emorine LJ, Descoins E, Pelloquin L, Brichese L, Gas N, Guillou E, Delettre C, Valette A,
Hamel CP, Ducommun B, Lenaers G, Belenguer P. 2002. The human dynamin-related protein OPA1
is anchored to the mitochondrial inner membrane facing the inter-membrane space. *FEBS Lett*
523:171–176. doi:10.1016/S0014-5793(02)02985-X

Olichon A, Guillou E, Delettre C, Landes T, Arnauné-Pelloquin L, Emorine LJ, Mils V, Daloyau M, Hamel
C, Amati-Bonneau P, Bonneau D, Reynier P, Lenaers G, Belenguer P. 2006. Mitochondrial dynamics
and disease, OPA1. *Biochim Biophys Acta - Mol Cell Res*. doi:10.1016/j.bbamcr.2006.04.003

Poole AC, Thomas RE, Yu S, Vincow ES, Pallanck L. 2010. The mitochondrial fusion-promoting factor
mitofusin   is   a   substrate   of   the   PINK1/parkin   pathway.   *PLoS   One*   **5**.
doi:10.1371/journal.pone.0010054

Powell BL, Toomes C, Scott S, Yeung A, Marchbank NJ, Spry PGD, Lumb R, Inglehearn CF, Churchill
AJ. 2003. Polymorphisms in OPA1 are associated with normal tension glaucoma. *Mol Vis* **9**:460–464.

Rahn JJ, Stackley KD, Chan SSL. 2013. Opa1 Is Required for Proper Mitochondrial Metabolism in Early

Development. *PLoS One* **8**. doi:10.1371/journal.pone.0059218

Rice P, Longden L, Bleasby A. 2000. EMBOSS: The European Molecular Biology Open Software Suite.

Trends Genet **16**:276–277. doi:10.1016/S0168-9525(00)02024-2

Richard M, Doubková K, Nitta Y, Kawai H, Sugie A, Tavosanis G. 2022. A Quantitative Model of Sporadic

Axonal Degeneration in the Drosophila Visual System. *J Neurosci* **42**:4937–4952.

doi:10.1523/JNEUROSCI.2115-21.2022

Rolland SG, Lu Y, David CN, Conradt B. 2009. The BCL-2-like protein CED-9 of C. elegans promotes

FZO-1/Mfn1,2- and EAT-3/Opa1-dependent mitochondrial fusion. *J Cell Biol* **186**:525–540.

doi:10.1083/jcb.200905070

Rubenstein JLR, Brutlag D, Clayton DA. 1977. The mitochondrial DNA of drosophila melanogaster exists

in two distinct and stable superhelical forms. *Cell* **12**:471–482. doi:10.1016/0092-8674(77)90123-4

Rujiviphat J, Meglei G, Rubinstein JL, McQuibban GA. 2009. Phospholipid association is essential for

dynamin-related protein Mgm1 to function in mitochondrial membrane fusion. *J Biol Chem*

284:28682–28686. doi:10.1074/jbc.M109.044933

Sandoval H, Yao CK, Chen K, Jaiswal M, Donti T, Lin YQ, Bayat V, Xiong B, Zhang K, David G, Charng

WL, Yamamoto S, Duraine L, Graham BH, Bellen HJ. 2014. Mitochondrial fusion but not fission

regulates larval growth and synaptic development through steroid hormone production. *Elife* **3**:1–23.

doi:10.7554/eLife.03558

Sanes JR, Zipursky SL. 2020. Synaptic Specificity, Recognition Molecules, and Assembly of Neural

Circuits. *Cell*. doi:10.1016/j.cell.2020.04.008

Sanes JR, Zipursky SL. 2010. Design Principles of Insect and Vertebrate Visual Systems. *Neuron*.

doi:10.1016/j.neuron.2010.01.018

Sarzi E, Angebault C, Seveno M, Gueguen N, Chaix B, Bielicki G, Boddaert N, Mausset-Bonnefont AL,

Cazevieille C, Rigau V, Renou JP, Wang J, Delettre C, Brabet P, Puel JL, Hamel CP, Reynier P,

Lenaers G. 2012. The human OPA1delTTAG mutation induces premature age-related systemic

neurodegeneration in mouse. *Brain* **135**:3599–3613. doi:10.1093/brain/aws303

Sarzi E, Seveno M, Piro-Mégy C, Elzière L, Quilès M, Péquignot M, Müller A, Hamel CP, Lenaers G,

Delettre C. 2018. OPA1 gene therapy prevents retinal ganglion cell loss in a Dominant Optic Atrophy

mouse model. *Sci Rep* **8**. doi:10.1038/s41598-018-20838-8

Schaaf CP, Blazo M, Lewis RA, Tonini RE, Takei H, Wang J, Wong LJ, Scaglia F. 2011. Early-onset

severe neuromuscular phenotype associated with compound heterozygosity for OPA1 mutations. *Mol*

Genet Metab **103**:383–387. doi:10.1016/j.ymgme.2011.04.018

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C,

Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A.

2012. Fiji: An open-source platform for biological-image analysis. *Nat Methods*.

doi:10.1038/nmeth.2019

Shahrestani P, Leung HT, Le PK, Pak WL, Tse S, Ocorr K, Huang T. 2009. Heterozygous mutation of

Drosophila Opa1 causes the development of multiple organ abnormalities in an age-dependent and

organ-specific manner. *PLoS One* **4**. doi:10.1371/journal.pone.0006867

Shimizu S, Mori N, Kishi M, Sugata H, Tsuda A, Kubota N. 2003. A novel mutation in the OPA1 gene in

a Japanese patient with optic atrophy. *Am J Ophthalmol* **135**:256–257. doi:10.1016/S0002-

9394(02)01929-3

Sladen PE, Perdigão PRL, Salsbury G, Novoselova T, van der Spuy J, Chapple JP, Yu-Wai-Man P,

Cheetham ME. 2021. CRISPR-Cas9 correction of OPA1 c.1334G>A: p.R445H restores

mitochondrial homeostasis in dominant optic atrophy patient-derived iPSCs. *Mol Ther - Nucleic*

Acids **26**:432–443. doi:10.1016/j.omtn.2021.08.015

Spiegel R, Saada A, Flannery PJ, Burté F, Soiferman D, Khayat M, Eisner V, Vladovski E, Taylor RW,

Bindoff LA, Shaag A, Mandel H, Schuler-Furman O, Shalev SA, Elpeleg O, Yu-Wai-Man P. 2016.

Fatal infantile mitochondrial encephalomyopathy, hypertrophic cardiomyopathy and optic atrophy

associated with a homozygous OPA1 mutation. *J Med Genet* **53**:127–131. doi:10.1136/jmedgenet-

2015-103361

Sugie A, Möhl C, Hakeda-Suzuki S, Matsui H, Suzuki T, Tavosanis G. 2017. Analyzing synaptic

modulation of drosophila melanogaster photoreceptors after exposure to prolonged light. *J Vis Exp*

2017. doi:10.3791/55176

Tang S, Le PK, Tse S, Wallace DC, Huang T. 2009. Heterozygous mutation of Opa1 in Drosophila shortens
lifespan mediated through increased reactive oxygen species production. *PLoS One* **4**.
doi:10.1371/journal.pone.0004492

Toomes C, Marchbank NJ, Mackey DA, Craig JE, Newbury-Ecob RA, Bennett CP, Vize CJ, Desai SP,
Black GCM, Patel N, Teimory M, Markham AF, Inglehearn CF, Churchill AJ. 2001. Spectrum,
frequency and penetrance of OPA1 mutations in dominant optic atrophy. *Hum Mol Genet* **10**:1369–
1378. doi:10.1093/hmg/10.13.1369

Trevisan T, Pendin D, Montagna A, Bova S, Ghelli AM, Daga A. 2018. Manipulation of Mitochondria
Dynamics Reveals Separate Roles for Form and Function in Mitochondria Distribution. *Cell Rep*
23:1742–1753. doi:10.1016/j.celrep.2018.04.017

Tsuyama T, Tsubouchi A, Usui T, Imamura H, Uemura T. 2017. Mitochondrial dysfunction induces
dendritic loss via eIF2α phosphorylation. *J Cell Biol* **216**:815–834. doi:10.1083/jcb.201604065

Vagnoni A, Bullock SL. 2016. A simple method for imaging axonal transport in aging neurons using the
adult Drosophila wing. *Nat Protoc* **11**:1711–1723. doi:10.1038/nprot.2016.112

Verny C, Loiseau D, Scherer C, Lejeune P, Chevrollier A, Gueguen N, Guillet V, Dubas F, Reynier P,
Amati-Bonneau P, Bonneau D. 2008. Multiple sclerosis-like disorder in OPA1-related autosomal
dominant optic atrophy. *Neurology* **70**:1152–1153. doi:10.1212/01.wnl.0000289194.89359.a1

Wang JT, Medress ZA, Barres BA. 2012. Axon degeneration: Molecular mechanisms of a self-destruction
pathway. *J Cell Biol*. doi:10.1083/jcb.201108111

Whitworth AJ, Lee JR, Ho VMW, Flick R, Chowdhury R, McQuibban GA. 2008. Rhomboid-7 and
HtrA2/Omi act in a common pathway with the Parkinson's disease factors Pink1 and Parkin. *DMM*
Dis Model Mech **1**:168–174. doi:10.1242/dmm.000109

Yamada M, Nitta Y, Uehara T, Suzuki H, Miya F, Takenouchi T, Tamura M, Ayabe S, Yoshiki A, Maeno
A, Saga Y, Furuse T, Yamada I, Okamoto N, Kosaki K, Sugie A. 2023. Heterozygous loss-of-
function DHX9 variants are associated with neurodevelopmental disorders: Human genetic and
experimental evidences. *Eur J Med Genet* **66**:104804. doi:10.1016/j.ejmg.2023.104804

Yang L, Tang H, Lin X, Wu Y, Zeng S, Pan Y, Li Y, Xiang G, Lin YF, Zhuang SM, Song Z, Jiang Y, Liu
X. 2020. OPA1-Exon4b Binds to mtDNA D-Loop for Transcriptional and Metabolic Modulation,
Independent of Mitochondrial Fusion. *Front Cell Dev Biol* **8**. doi:10.3389/fcell.2020.00180

Yarosh W, Monserrate J, Tong JJ, Tse S, Le PK, Nguyen K, Brachmann CB, Wallace DC, Huang T. 2008.
The molecular mechanisms of OPA1-mediated optic atrophy in Drosophila model and prospects for
antioxidant treatment. *PLoS Genet* **4**:0062–0071. doi:10.1371/journal.pgen.0040006

Yu-Wai-Man P, Griffiths PG, Gorman GS, Lourenco CM, Wright AF, Auer-Grumbach M, Toscano A,
Musumeci O, Valentino ML, Caporali L, Lamperti C, Tallaksen CM, Duffey P, Miller J, Whittaker
RG, Baker MR, Jackson MJ, Clarke MP, Dhillon B, Czermin B, Stewart JD, Hudson G, Reynier P,
Bonneau D, Marques W, Lenaers G, McFarland R, Taylor RW, Turnbull DM, Votruba M, Zeviani
M, Carelli V, Bindoff LA, Horvath R, Amati-Bonneau P, Chinnery PF. 2010a. Multi-system
neurological disease is common in patients with OPA1 mutations. *Brain* **133**:771–786.
doi:10.1093/brain/awq007

Yu-Wai-Man P, Spyropoulos A, Duncan HJ, Guadagno J V., Chinnery PF. 2016. A multiple sclerosis-like
disorder in patients with OPA1 mutations. *Ann Clin Transl Neurol* **3**:723–729. doi:10.1002/acn3.323

Yu-Wai-Man P, Stewart JD, Hudson G, Andrews RM, Griffiths PG, Birch MK, Chinnery PF. 2010b. OPA1
increases the risk of normal but not high tension glaucoma. *J Med Genet* **47**:120–125.
doi:10.1136/jmg.2009.067512

Zaninello M, Palikaras K, Naon D, Iwata K, Herkenne S, Quintana-Cabrera R, Semenzato M, Grespi F,
Ross-Cisneros FN, Carelli V, Sadun AA, Tavernarakis N, Scorrano L. 2020. Inhibition of autophagy
curtails visual loss in a model of autosomal dominant optic atrophy. *Nat Commun* **11**.
doi:10.1038/s41467-020-17821-1

Zaninello M, Palikaras K, Sotiriou A, Tavernarakis N, Scorrano L. 2022. Sustained intracellular calcium
rise mediates neuronal mitophagy in models of autosomal dominant optic atrophy. *Cell Death Differ*
29:167–177. doi:10.1038/s41418-021-00847-3

Zanna C, Ghelli A, Porcelli AM, Karbowski M, Youle RJ, Schimpf S, Wissinger B, Pinti M, Cossarizza A,
Vidoni S, Valentino ML, Rugolo M, Carelli V. 2008. OPA1 mutations associated with dominant

optic atrophy impair oxidative phosphorylation and mitochondrial fusion. *Brain* **131**:352–367.

doi:10.1093/brain/awm335

Figures with the corresponding legend

Figure 1: Effect of *dOPA1* knockdown on mitochondrial density and size in retinal axons.

(A) Schematic illustration of the *Drosophila* visual system from a dorsal view, highlighting the

arrangement of photoreceptor neurons. The rectangular area indicated in B represents a single R7/8

photoreceptor axon, as illustrated in the figure for B. The rectangular areas highlighted in C and D

encompass the entirety of R7/8 photoreceptor axons, as shown in the figures for C and D. The lines

marked in F and G denote cross-sections of R1–6 photoreceptor axons, as depicted in the figures for

F and G. (B) Visualization of mitochondria and dOPA1 in a set of R7/8 axons. Mitochondria were

visualized using the expression of *mCherry-mito* driven by *GMR-Gal4* (magenta) while HA-tagged

dOPA1 was immunostained using the anti-HA antibody (green). Scale bar: 3 μm. (C, D)

Representations of mitochondria visualized using the expression of *Mito-GFP* driven by *GMR-Gal4*

(green), one day after eclosion. Retinal axons were labeled using the anti-Chaoptin antibody (magenta).

Scale bar: 30 μm. (E) Quantification of Mito-GFP intensity. Mito-GFP levels were calculated by

dividing the Mito-GFP fluorescence intensity by the anti-Chaoptin signal intensity. Control (n = 28

optic lobes) and *dOPA1* RNAi (n = 28 optic lobes). The data are presented as mean ± SEM. (F, G)

Electron micrographs of cross sections of the R1–6 retinal axons in the lamina of the Control (F) and

d*OPA1* knockdown (G) on the day of eclosion. Yellow circles and white dotted lines indicate

mitochondria and a lamina column, respectively. Scale bar: 2 μm. (H) Quantification of the

mitochondrial area. Control (n = 96 mitochondria) and *dOPA1* RNAi (n = 92 mitochondria). Data are

expressed as mean ± SEM. (I) Representative EM images showing mitochondria with a densely

packed matrix structure (classified as class A) and collapsed mitochondria (classified as class B). Scale

bar: 500 nm. (J) Quantification of the mitochondria classified into class A and class B. Control (n =

96 mitochondria) and *dOPA1* RNAi (n = 92 mitochondria). See the Table1 for the genotypes of the

Drosophila used in the experiments.

Figure S1. Analysis of trafficking defects in axons of *dOPA1* knockdown flies.

Schematic illustration of the *Drosophila* visual system around 24 h after puparium formation. The cell

bodies of photoreceptors (magenta) are found in the retina; type 1–6 photoreceptors project their axons

to the lamina of the primary optic ganglion. The axons of type 7/8 photoreceptors project into the

medulla of the secondary optic ganglion. Mitochondria are abundantly located in their cell bodies and

axon terminals (green). Higher-order visual centers include the lobula and lobula plate. (B–D)

Neuropil and photoreceptor cells were labeled using an anti-CadN antibody (cyan) and anti-Chaoptin

antibody (magenta), respectively. Mitochondria were visualized using the expression of Mito-GFP

driven by *GMR-Gal4* (green). (B, B') and (C, C') show whole photoreceptors in the RNAi Control and

dOPA1 knockdown, respectively. The regions indicated by dashed and solid lines in (B', C', D') indicate

the cell body and axonal areas, respectively. (D, D') shows whole photoreceptors in *milton* knockdown.

Scale bar: 50 μm. See the Table1 for the genotypes of the *Drosophila* used in the experiments.

Figure 2. elevated reactive oxygen species and mitophagy activity levels in the *dOPA1* knockdown.

(A, B) Mitochondrial reactive oxygen species (ROS) levels in a set of the R7/8 retinal axon visualized using Mito-GFP (green) driven by *GMR-Gal4* and MitoSOX (magenta) to specifically detect superoxide, an indicator of oxidative stress. Representative images of a retinal axon in the Control (A, A') and *dOPA1* knockdown (B, B'). (C) Quantification of ROS levels for each genotype. ROS levels were determined by dividing the fluorescence intensity of MitoSOX by that of Mito-GFP, highlighting the significant increase in ROS levels in *dOPA1* knockdown flies compared to controls, suggesting

enhanced mitochondrial stress and dysfunction upon *dOPA1* depletion. Control (n = 20 optic lobes)

and *dOPA1* RNAi (n = 18 optic lobes). Scale bar: 3 μm. Data are presented as mean ± SEM. (D, E)

Mitophagy activity in a set of the R7/8 retinal axon visualized by a genetic marker, mito-QC, driven

by *GMR-Gal4*. Representative images of a retinal axon in the Control 1 day after eclosion (D), *dOPA1*

knockdown 1 day after eclosion (E), the Control 1 week after eclosion (F), *dOPA1* knockdown 1 week

after eclosion (G). The arrowheads indicate a mitolysosome that is positive for mCherry and negative

for GFP. (H) Quantification of the mitophagy activity each genotype. The number of the

mitolysosomes were counted in each genotype. The significant increase in the number of the

mitolysosomes in *dOPA1* knockdown flies compared to controls after 1 week, suggesting enhanced

mitophagy upon *dOPA1* depletion. Control 1d (n = 18 optic lobes), Control 1week (n = 19 optic lobe),

dOPA1 RNAi 1d (n = 24 optic lobes), and *dOPA1* RNAi 1 week (n = 11 optic lobes). Scale bar: 3 μm.

Data are presented as mean ± SEM. See the Table1 for the genotypes of the *Drosophila* used in the

experiments.

Figure 3. Effect of *dOPA1* knockdown on photoreceptor neurodegeneration in *Drosophila*.

Schematic of the visual system in *Drosophila*. A set of R7/8 photoreceptors project their axons to the

second optic ganglion medulla. The rectangular prism indicated in B–E represents the area of the

whole medulla, as shown in the figure for B–E. The squire marked in G–J denote cross-sections of the

cell bodies of the photoreceptors, as depicted in the figures for G–J. (B–E) All R7 and R8 axon

terminals project to the medulla. *dOPA1* in the photoreceptor was knocked down by *GMR-Gal4*. *40D-*

UAS was used as a Control to match the number of *UAS* sequences recognized by Gal4. One day after

eclosion in Control (B) and *dOPA1* knockdown (C), and one week after in Control (D) and *dOPA1*

knockdown (E). The retinal axons were stained with anti-Chaoptin, a photoreceptor-specific antibody.

Scale bar = 50 μm. (F) Number of axons at each time point in R7 neurons for each situation quantified

by MeDUsA. At 1 day of Control (n = 34 optic lobes) and *dOPA1* RNAi (n = 23 optic lobes), and at 1

week of Control (n = 28 optic lobes) and *dOPA1* RNAi (n = 26 optic lobes). Data are expressed as

mean ± SEM. (G–J) Electron micrographs of cross sections of the photoreceptor cell bodies in the

ommatidia of the retina. Rhabdomeres are shown as yellow arrowheads. The day of eclosion in Control

(G) and *dOPA1* knockdown (H), and 1-week-old adults of Control (I) and *dOPA1* knockdown (J).

Scale bar = 2 μm. **(K)** Quantification of the number of rhabdomeres for each genotype and time point.

At 0 day in Control (n = 50 ommatidia) and *dOPA1* RNAi (n = 127 ommatidia), and at 1 week in

Control (n = 50 ommatidia) and *dOPA1* RNAi (n = 117 ommatidia). Data are expressed as mean ±

SEM. See the Table1 for the genotypes of the *Drosophila* used in the experiments.

Figure S2. The impact on the retinal axons of another RNAi line in *Drosophila OPA1.*

(A) The images show the representations of retinal axons labeled with an anti-Chaoptin antibody (gray). Control and *dOPA1* RNAi (VDRC 106290) from VDRC two weeks after eclosion. Scale bar: 50 μm. (B) Quantification of the number of axons for each genotype. Control (n = 14 optic lobes) and *dOPA1* RNAi (VDRC 106290) (n = 22 optic lobes). The data are presented as mean ± SEM. See the Table1 for the genotypes of the *Drosophila* used in the experiments.

Figure S3. Evaluation of the impact on *Drosophila* compound eyes due to knockdown of *dOPA1*.

(A) Bright-field microscopic images of representative fly eyes expressing 40D-UAS (control), RNAi #1 (BDSC 32358), and RNAi #2 (VDRC 106290) using the *GMR-Gal4* driver. (B) Graph representing the P score of each genotype: female *control* (n = 19), *female RNAi #1* (n = 19), *female RNAi #2* (n = 20), male control (n = 21), *male RNAi #1* (n = 20), and *male RNAi #2* (n = 20). See the Table1 for the genotypes of the *Drosophila* used in the experiments.

1005

Figure 4. Role of *dOPA1* in the retinal axon according to mutant analysis.

(A–C) Representations of retinal axons labeled with an anti-Chaoptin antibody (gray), one day after

eclosion. The images show Control (A), *dOPA1^{s3475}* somatic mosaic flies (B), and *dOPA1^{s3475}* somatic

mosaic flies expressing the eye-specific full-length *dOPA1* (C). Scale bar: 50 μm. (D) Quantification

of the number of axons for each genotype. Control (n = 41 optic lobes), mutant (n = 45 optic lobes),

and rescue (n = 41 optic lobes). The data are presented as mean ± SEM. See the Table1 for the

genotypes of the *Drosophila* used in the experiments.

1013

1014

Figure 5. Verification of the pathological significance of disease mutations in *hOPA1*.

(A) Schematic illustration of the *hOPA1* gene construct with HA and myc tags in the UAS-based vector.

hOPA1 includes a mitochondrial import sequence (MIS) cleaved by mitochondrial processing

peptidase (MPP), a transmembrane region (TM), a coiled-coil region (CC1), a GTPase domain, a

Middle domain, and a GTPase effector domain (GED) containing a coiled-coil region (CC2). The sites

of variants (I382M, D438V, R445H, and a deletion from 2708 to 2711) are shown in red. S1 is the site

cleaved by OMA1. (B) Western blot analysis to confirm the expression of hOPA1 variants.

hOPA1_WT, hOPA1_D438V, hOPA1_I382M, hOPA1_R445H, and hOPA1_2708del were expressed

in whole *Drosophila* bodies and detected using anti-HA and anti-myc antibodies. α-Tubulin was used

as a loading Control. (C) Quantification of the expression levels of the OPA1 protein for each variant.

hOPA1_WT (n = 3), D438V (n = 3), I382M (n = 3), R445H (n = 3), and 2708del (n = 3). The data are

presented as mean ± SEM. (D) Detection of hOPA1 expression in vivo in *Drosophila* using anti-OPA1.

Band 1 represents the full-length OPA1, which includes the MIS, while Band 2 corresponds to the

long-form OPA1 (L-OPA1), cleaved at the MPP. Bands 3 and 5 potentially detect the endogenous

dOPA1. Band 4 is thought to represent the short-form OPA1 (S-OPA1). (E) Rescue experiments were

conducted to assess the expression of each *hOPA1* variant, including *dOPA1*, in the retina axons of

dOPA1 mutant somatic clones. The sample size is indicated (n). Data are presented as mean ± SEM.

(F) Quantification of mitochondrial ROS levels upon expression of hOPA1 variants in the context of

dOPA1 knockdown. Control (n = 22 optic lobes), *dOPA1* RNAi (n = 16 optic lobes), *hOPA1_WT* with

dOPA1 RNAi (n = 18 optic lobes) , *hOPA1_2708-2711del* with *dOPA1* RNAi (n = 16 optic lobes),

hOPA1_I382M with *dOPA1* RNAi (n = 14 optic lobes), *hOPA1_D438V* with *dOPA1* RNAi (n = 16

optic lobes), and *hOPA1_R445H* with *dOPA1* RNAi (n = 18 optic lobes). Data are presented as mean

± SEM. See the Table1 for the genotypes of the *Drosophila* used in the experiments.

Figure S4. Effects of *hOPA1* variants on COXII expression and Autophagy in the context of

***dOPA1* knockdown.**

(A) Western blot analysis to show the expression of COXII upon expression of *hOPA1* variants in the

context of *dOPA1* knockdown. (B) Quantification of COXII protein levels. Control (n = 3), *luciferase*

RNAi with *dOPA1* RNAi (n = 3), *hOPA1_WT* with *dOPA1* RNAi (n = 3) , *hOPA1_2708-2711del* with

dOPA1 RNAi (n = 3), *hOPA1_I382M* with *dOPA1* RNAi (n = 3), *hOPA1_D438V* with *dOPA1* RNAi

(n = 3), and *hOPA1_R445H* with *dOPA1* RNAi (n = 3). Data are presented as mean ± SEM. (C) Western

blot analysis to show the expression of Atg8a-1 and Atg8a-2 upon expression of *hOPA1* variants in the

context of *dOPA1* knockdown. (D) Quantification of the ratio of Atg8a-2 and Atg8a-1. Control (n =

3), luciferase RNAi with *dOPA1* RNAi (n = 3), *hOPA1_WT* with *dOPA1* RNAi (n = 3) , *hOPA1_2708-*

2711del with *dOPA1* RNAi (n = 3), *hOPA1_I382M* with *dOPA1* RNAi (n = 3), hOPA1_D438V with

dOPA1 RNAi (n = 3), and *hOPA1_R445H* with *dOPA1* RNAi (n = 3). Data are presented as mean ±

SEM.

1053

1054

1055

1056

1057

1058

1059

Figure 6. LOF or DN effects of disease mutations in *hOPA1*.

(A) Impact of each human OPA1 variant on the axon number in the optic nerve of *Drosophila* as

quantified using MeDUsA. (B) Expression of both *hOPA1* WT and its respective variants was analyzed

in photoreceptors lacking *dOPA1*, and the number of axons was quantified using the MeDUsA. The

sample size is indicated (n). Data are expressed as mean ± SEM. (C) Schematic representation of

1065   interspecies differences in OPA1 interactions and the interchangeability of OPA1 between human and

1066   fly. See the Table1 for the genotypes of the *Drosophila* used in the experiments.

1067

1068

1069

1070

1071

1072

1073

A   GTPase domain   173/241 (71.8%)

```
hOPA1 [NM_015560.3] 280 ASYNTQDHLPRVVVVGDQSAGKTSVLEMIAQARIFPRGSGEMMTRSPVKVTLSEGPHHVALFKDSSREFDLTKEEDLAALRHEIELRMRKNVKEGCTVSP 379
                        .|...||||||||||||||||:||||||||.||:|||||||||||||||||:||||:|||||:||.|:|||.|||.||..||.::||.::.|.|||.
dOPA1 [NM_166040.2] 291 TGYTMADHLPRVVVVGDQSSGKTSVLESIAKARIFPRGSGEMMTRAPVKVTLAEGPYHVAQFRDSDREYDLTKESDLQDLRRDVEFRMKASVRGGKTVSN 390

                    380 ETISLNVKGPGLQRMVLVDLPGVINTVTSGMAPDTKETIFSISKAYMQNPNAIILCIQDGSVDAERSIVTDLVSQMDPHGRRTIFVLTKVDLAEKNVASP 479
                        .|.||||||||||||||||||||.||:.||||:||||.||.|.||..|.||||||||||||||||||.|||||.||.||..|||||||||||||.|:.|
                    391 EVIAMTVKGPGLQRMVLVDLPGIISTMTVDMASDTKDSIHQMTKHYMSNPNAIILCIQDGSVDAERSNVTDLVMQCDPLGRRTIFVLTKVDLAEE-LADP 489

                    480 SRIQQIIEGKLFPMKALGYFAVVTGKGNSSESIEAIREYEE 520
                        .||:|:.||||||||||||.||||.|.|||:..:|:||||||
                    490 DRIRKILSGKLFPMKALGYYAVVTGRGRKDDSIDAIRQYEE 530
```

B   Middle domain   204/376 (54.3%)

```
hOPA1 [NM_015560.3] 521 EFFQNSKLL-KTSMLKAHQVTTRNLSLAVSDCFWKMVRESVEQQADSFKATRFNLETEWKNNYPRLRELDRNELFEKAKNEILDEVISLSQVTPKHWEEI 619
                        :||:|||.  .:..:..||||:||||||||.|||||:||||||:|||:||||||||||||||:|||:||||:.:||||||.|:|||||:.|||||::.|:
dOPA1 [NM_166040.2] 531 DFFKNSKLFHRRGVIMPHQVTSRNLSLAVSDRFWKMVRETIEQQADAFKATRFNLETEWKNNFPRLRESGRDELFDKAKGEILDEVVTLSQISAKKWDDA 630

                    620 LQQSLWERVSTHVIENIYLPAAQTMNSGTFNTTVDIKLKQWTDKQLPNKAVEVAWETLQEEFSRFM-TEPKGKEHDDIFDKLKEAVKEESIKRHKWNDFA 718
                        |...||:|:.:.|.|:|||||:.:.:.|||.|:|||||||.|||||.|||::||||||||:|:||.  :..|...|::||:|:|||.||.|:.:||.|.|
                    631 LSTKLWEKLSNYVFESIYLPAAQSGSQNSFNTMVDIKLRQWAEQALPAKSVEAGWEALQQEFISLMERSKKAQDHDGIFDQLKSAVVDEAIRRHSWEDKA 730

                    719 EDSLRVIQHNALEDRSISDKQQWDAAIYFMEEALQARLKDTENAIENMVGPDWKKRWLYWKNRTQEQCVHNETKNELEKMLKCNEEHPAYLASDEITTVR 818
                        .|.|||:|.||:..|:.:|||.|:|..|:.|:.|:|:.|.:.|.::||.|..||.:.:||..|.:||:.|.|||..:.|:.||||.|||:.|.|.:||||
                    731 IDMLRVIQLNTLEDRFVHDKQEWDSAVKFLESSVNAKLVQTEETLAQMFGPGQMRRITHWQYLTQDQQKRRSVKNELDKILKNDTKHLPTLTHDELTTVR 830

                    819 KNLESRGVEVDPSLIKDTWHQVYRRHFLKTALNHCNLCRRGFYYYQRHFVDSELECNDVVLFWRIQRMLAITANTL 894
                        |||:...|:||...|:..||.|||||||.||....||:.:|.|.:..||.||||||||||||||:::.||.|.|
                    831 KNLQRDNVDVDTDYIRQTWFPVYRKHFLQQALQRAKDCRKAYYLYTQQGAECEISCSDVVLFWRIQQVIKITGNAL 906
```

C   GED domain   42/66 (63.6%)

```
hOPA1 [NM_015560.3] 895 RQQLTNTEVRRLEKNVKEVLEDFAEDGEKKIKLLTGKRVQLAEDLKKVREIQEKLDAFIEALHQEK 960
                        ||:.|.|.||||.:|:.|||:.|.:|||||||||||||.||||:|||||.||:|||.||.:|:||
dOPA1 [NM_166040.2] 907 RQQVINREARRLDKEIKAVLDEFSDDEEKKGYLLTGKRVLLAEELIKVRQIQEKLEEFINSLNQEK 972
```

Figure S5. Pairwise Alignment of GTPase, Middle, and GED Domains of hOPA1 and dOPA1.

(A–C) The amino acid sequences of hOPA1 ([NM_015560.3]) and dOPA1 ([NM_166040.2]) were

aligned using EMBOSS Needle. GTPase domain (A), Middle domain (B), and GED domain (C). Each

percentage value represents the degree of identity within the amino acid sequence. GTPase domain

(280–520), Middle domain (521–894), and GED domain (895–907). The beginning and end of each

domain were reported previously (Liesa et al., 2009).

1082

Figure S6. Comparative analysis of mitochondrial ROS levels, COXII expression, and

Autophagy between *hOPA1* wild-type and the variants expression under *dOPA1* knockdown

conditions.

(A) Impact on mitochondrial ROS levels upon expression of wild-type *hOPA1* and its variants under

conditions of *dOPA1* knockdown. Control (n = 16 optic lobes), *hOPA1_WT* (n = 20 optic lobes) ,

hOPA1_2708-2711del (n = 18 optic lobes), *hOPA1_I382M* (n = 20 optic lobes), *hOPA1_D438V* (n =

18 optic lobes), and *hOPA1_R445H* (n = 15 optic lobes). Data are presented as mean ± SEM. (B)

Western blot analysis to show the expression of COXII upon expression of wild-type *hOPA1* and its

variants under condition of *dOPA1* knockdown. (C) Quantification of COXII protein levels. Control

(n = 3), *hOPA1_WT* (n = 3) , *hOPA1_2708-2711del* (n = 3), *hOPA1_I382M* (n = 3), *hOPA1_D438V* (n

= 3), and *hOPA1_R445H* (n = 3). Data are presented as mean ± SEM. (D) Western blot analysis to

show the expression of Atg8a-1 and Atg8a-2 upon expression of wild-type *hOPA1* and its variants

under condition of *dOPA1* knockdown. (E) Quantification of the ratio of Atg8a-2 and Atg8a-1. Control

(n = 3), *hOPA1_WT* (n = 3) , *hOPA1_2708-2711del* (n = 3), *hOPA1_I382M* (n = 3), *hOPA1_D438V* (n

= 3), and *hOPA1_R445H* (n = 3). Data are presented as mean ± SEM. See the Table1 for the genotypes

of the *Drosophila* used in the experiments.

1099

1100

1101

1102

Figure S7. The effect of a dominant-negative mutation in *dOPA1*.

(A) The images show the representations of retinal axons labeled with an anti-Chaoptin antibody

(gray). Control and the expression of a presumably GTPase-negative form of *dOPA1^{K273A}* 1 week after

eclosion. Scale bar: 50 μm. (B) Quantification of the number of axons for each genotype. Control (n

= 26 optic lobes) and *dOPA1^{K273A}* (n = 22 optic lobes). The data are presented as mean ± SEM. See

the Table1 for the genotypes of the *Drosophila* used in the experiments.

1109