Title

Drosophila model to clarify the pathological significance of OPA1 in autosomal dominant optic

atrophy

Authors

Yohei Nitta[1,7], Jiro Osaka[2,7], Ryuto Maki[2], Satoko Hakeda-Suzuki[2,3], Emiko Suzuki[4,5], Satoshi Ueki[6],

Takashi Suzuki[2] and Atsushi Sugie[1] [*]

[1] Brain Research Institute, Niigata University, Niigata, 951-8585, Japan

[2] School of Life Science and Technology, Tokyo Institute of Technology, Yokohama, 226-8501, Japan

[3] Research Initiatives and Promotion Organization, Yokohama National University, Yokohama, 240-

8501, Japan

[4] Department of Biological Sciences, Graduate School of Science, Tokyo Metropolitan University,

Hachioji 192-0397, Japan

[5] Department of Gene Function and Phenomics, National Institute of Genetics, Mishima, 411-8540,

Japan

[6] Division of Ophthalmology and Visual Science, Graduate School of Medical and Dental Sciences,

Niigata University, Niigata, 951-8585, Japan

[7] Equal contribution

* Corresponding author: atsushi.sugie@bri.niigata-u.ac.jp (A.S.) Tel (+81) 25-227-0616.

Abstract

Autosomal dominant optic atrophy (DOA) is a progressive form of blindness caused by degeneration

of retinal ganglion cells and their axons, mainly caused by mutations in the OPA1 mitochondrial

dynamin *like* GTPase (*OPA1*) gene. *OPA1* encodes a dynamin-like GTPase present in the

mitochondrial inner membrane. When associated with OPA1 mutations, DOA can present not only

ocular symptoms but also multi-organ symptoms (DOA plus). DOA plus often results from point

mutations in the GTPase domain, which are assumed to have dominant negative effects. However, the

presence of mutations in the GTPase domain does not always result in DOA plus. Therefore, an

experimental system to distinguish between DOA and DOA plus is needed. In this study, we found

that loss-of-function mutations of the *dOPA1* gene in *Drosophila* can imitate the pathology of optic

nerve degeneration observed in DOA. We successfully rescued this degeneration by expressing the

human *OPA1* (*hOPA1*) gene, indicating that *hOPA1* is functionally interchangeable with *dOPA1* in the

fly system. However, we could not rescue any previously reported mutations known to cause either

DOA or DOA plus. By expressing both WT and DOA plus mutant *hOPA1* forms in the optic nerve of

dOPA1 mutants, we observed that DOA plus mutations suppressed the rescue, facilitating the

distinction between loss-of-function and dominant negative mutations in *hOPA1*. The fly model

developed in this study can assist in the differential diagnosis between DOA and DOA plus and inform

early treatment decisions in patients with mutations in *hOPA1*.

Introduction

Autosomal dominant optic atrophy (DOA) is a progressive form of blindness characterized by

selective degeneration of retinal ganglion cells (RGCs) and the axons that form the optic nerve.

Despite being a rare disease with a frequency of 1/12,000 to 1/50,000, DOA is the most commonly

diagnosed form of hereditary optic neuropathy. In 2000, *OPA1 mitochondrial dynamin like GTPase*

(OPA1) was identified as the causative gene for DOA (Alexander et al., 2000; Delettre et al., 2000).

OPA1 encodes a dynamin-like GTPase located in the inner mitochondrial membrane

(Olichon et al., 2002). It has various functions, including mitochondrial fusion, mtDNA maintenance,

control of cell death, and resistance to reactive oxygen species (ROS) (Lenaers et al., 2021). The most

common pathogenic mutation in *OPA1* is the c.2708_2711 delTTAG deletion (Toomes et al., 2001).

Five pathogenic mutations, including the one above, result in a substantial decrease in OPA1 protein

in the fibroblasts from affected patients (Zanna et al., 2008), supporting the theory that DOA is caused

by haploinsufficiency.

Optic atrophy is a characteristic feature of *OPA1*-associated DOA; however, multisystem

symptoms have been reported in up to 20% of *OPA1* mutation carriers (Yu-Wai-Man et al., 2010a).

This is called DOA plus. DOA plus can include sensorineural deafness (Amati-Bonneau et al., 2008),

multiple sclerosis (Verny et al., 2008; Yu-Wai-Man et al., 2016), parkinsonism and dementia (Carelli

et al., 2015; Lynch et al., 2017), and cardiomyopathy (Spiegel et al., 2016). The R445H mutation is a

well-known mutation associated with DOA plus (Amati-Bonneau et al., 2008, 2003; Shimizu et al.,

2003). The proportion of mutated OPA1 locations differs between patients with DOA and DOA plus.

Missense mutations in the dynamin-featured structure, especially the GTPase domain of *OPA1*, are

more likely to cause severe symptoms compared with loss-of-function (LOF) mutations such as

deletions or splice site mutations (Yu-Wai-Man et al., 2010a). This is probably due to the dominant

negative effect (DN) of a DOA plus mutation on the normal *OPA1* allele, since OPA1 functions as a

homo-oligomer (Frezza et al., 2006; Olichon et al., 2006); in fact, its yeast ortholog, Mgm1, forms

oligomers, increasing the GTPase activity (Meglei and McQuibban, 2009; Rujiviphat et al., 2009).

Particularly, the dynamin-featured structure accounts for 70% of the 233 pathogenic variants of DOA

and DOA plus described in the locus-specific database (Ferré et al., 2015). However, mutations in

these domains do not determine DOA plus, as it affects only 20% of all *OPA1* mutation carriers. Thus,

experimental models to determine whether a gene mutation is LOF or DN are required to distinguish

between DOA and DOA plus. Although currently there is no effective treatment for DOA or DOA

plus, distinguishing between them allows earlier planning of interventions such as hearing aids, rather

than relying solely on visual aids, necessary only for those with DOA plus.

As *OPA1* is conserved in various species, DOA models have been reported in vertebrates

such as mice and zebrafish, simple organisms such as nematodes and fruit flies, and *in vitro* yeast and

cultured cell models (Del Dotto and Carelli, 2021). In mouse models, focal RGC axons decrease in

number with mitochondrial abnormalities at the electron microscope level, but the phenotype appears

slowly, not allowing for analysis in a short time frame (Alavi et al., 2007; Davies et al., 2007; Nguyen

et al., 2011; Sarzi et al., 2012). Zebrafish models have displayed developmental delays in eyes and

heads, short length, and body axis abnormalities (Eijkenboom et al., 2019; Rahn et al., 2013) but have

not been used to investigate optic nerve degeneration.

In *C. elegans*, most studies focused on muscle cells using mutants of *eat-3*, an *OPA1*

homolog (Kanazawa et al., 2008; Rolland et al., 2009). However, the mitochondria in the posterior

lateral microtubule neurons did not show any difference in size from its wild-type (WT) counterparts

(Byrne et al., 2019). Thus, LOF of *eat-3* may have a limited impact on phenotypic outcomes in the

nervous system. Nevertheless, a DOA model in which $hOPA1^{K301A}$ is expressed in GABAergic motor

neurons, showed a decreased number of mitochondria in those cells (Zaninello et al., 2020). In a fly

model, the *dOPA1* somatic clone in the eye exhibited a rough-eye phenotype, eye structural

abnormalities, and increased MitoSOX fluorescence, which was partially rescued by vitamin E or

SOD1 expression (Yarosh et al., 2008). Heterozygous *dOPA1* mutants showed shortened lifespan,

elevated ROS levels, and irregular muscle tissue mitochondrial structures (Tang et al., 2009).

Moreover, the Electroretinogram (ERG) pattern showed age-dependent decreases in the on transient

response, heart rate, negative geotaxis response, and increased heart arrhythmia due to heat shock

stress (Shahrestani et al., 2009). However, except for mice, the evidence of optic nerve axonal

abnormalities is limited in model organisms such as zebrafish, worms, and flies.

The structure of the *Drosophila* visual system is similar to that of mammals (Sanes and

Zipursky, 2010), with conserved mechanisms of synaptogenesis and neural circuit formation (Sanes

and Zipursky, 2020). *Drosophila* photoreceptors type R7/8 extend their retinal axons from the retina

directly into the brain, forming synapses in the second optic ganglion medulla via the chiasma (Figure

1A), having potential application as a model for mammalian photoreceptors and RGCs. Based on this,

we aimed to develop an experimental model for observing and quantifying degeneration in the

Drosophila retinal axon (Nitta et al., 2023; Richard et al., 2022).

In this study, a *dOPA1* LOF could mimic the human DOA, in which mitochondrial

fragmentation, increased ROS levels, and neurodegeneration occur in retinal axons. The axonal

degeneration induced by the *dOPA1* mutant was rescued by the expression of human *OPA1* (*hOPA1*).

This demonstrated that the function of *hOPA1* is the same as in the fly. As none of the previously

reported mutations known to cause DOA or DOA plus were rescued, their function seems to be

impaired. To distinguish between the effect of LOF and DN mutations, we expressed the *hOPA1* gene

of both WT and mutation in a genetic background in which *dOPA1* was deleted in retinal axons. The

DOA plus mutations D438V and R445H inhibited the rescue by the WT of *hOPA1*. Taken together,

we generated a new *Drosophila* DOA model which served to isolate the LOF or DN effect of *hOPA1*

mutations.

Results

***dOPA1* depletion caused mitochondrial fragmentation in axon terminals**

The OPA1 protein is located in the inner mitochondrial membrane. To confirm whether dOPA1 is also

localized in the mitochondria of *Drosophila* retinal axons, we used a Gal4/UAS system (Brand and

Perrimon, 1993) and expressed HA-tagged dOPA1 proteins using the eye-specific *GMR-Gal4* driver

(Newsome et al., 2000a, 2000b). The outer membrane of the mitochondria was visualized with

mCherry-mito, which tagged the transmembrane domain of Miro with mCherry (Vagnoni and Bullock,

2016). Confocal microscopy revealed that the HA signals colocalized with mCherry-mito in the axon

terminals of photoreceptor cells (Figure 1B). This indicated that dOPA1 colocalizes with mitochondria

in retinal axons.

To examine the impact of *dOPA1* on mitochondrial structure and density, we analyzed

mitochondria in *dOPA1* RNAi flies. The intensity of the mitochondrial signal was significantly

reduced in the retinal axons of these flies (Figure 1C and D; quantification in 1E), suggesting a

decreased number of mitochondria in retinal axons. However, whether this decrease is due to

degradation resulting from a decline in mitochondrial quality or axonal transport failure remains

unclear. Previously, impaired mitochondrial transport was reported in the LOF of *marf* (MFN 1/2

homolog), which is necessary for the fusion of the outer membrane of mitochondria (Sandoval et al.,

2014). Thus, we investigated if the LOF of *dOPA1* may cause transport defects, which may be

responsible for the reduced mitochondrial density in retinal axons in *dOPA1* RNAi flies. To this end,

we observed the photoreceptor neurons after puparium formation at 24 h, which allowed us to

simultaneously visualize the cell bodies and axon terminals of the photoreceptors (Figure S1A). Both

dOPA1 knockdown and control flies showed mitochondrial signals in the axon terminals (Figure S1B

and C). However, no mitochondrial signals were observed in the axon terminals of flies with a

knockdown of *milton*, which is thought to act as an adaptor between kinesin to mitochondria,

promoting anterograde transport (Figure S1D). These findings suggest that trafficking defects are not

the main cause of the mitochondrial mislocalization observed in the retinal axons of the *dOPA1*

knockdown.

Nevertheless, the degradation resulting from mitochondrial fragmentation may have

decreased the mitochondrial signal. To further analyze the mitochondrial morphology of retinal axons

in the LOF of *dOPA1*, we conducted an electron microscopy analysis (Figure 1F and G). Although

Drosophila has eight types of photoreceptors, the axons of R1–6, which project to the lamina of the

first optic lobe, can be identified without markers; thus, we chose them for this experiment. The *dOPA1*

knockdown led to a significant decrease in mitochondria size in R1–6 axons (Figure 1F–H). In addition,

there was significant heterogeneity in healthy mitochondria, ranging from dense-matrix (Class A) to

swollen mitochondria with a hypodense matrix (Class B) (Figures 1I). Knocking down *dOpa1* in R1–

6 axon terminals resulted in severely abnormal mitochondrial ultrastructure (22.8%; Figure 1J). These

results suggest that *dOPA1* is involved in maintaining proper mitochondrial morphology by promoting

mitochondrial fusion in retinal axons. Previous studies have observed mitochondrial fragmentation in

S2 cells (McQuibban et al., 2006), muscle tissue (Deng et al., 2008), segmental nerves (Trevisan et

al., 2018), and ommatidia (Yarosh et al., 2008) due to LOF of *dOPA1*, which is consistent with the

results observed in the retinal axons.

To verify whether ROS is increased in the retinal axons in *dOPA1* knockdown

photoreceptors, we measured the ROS levels in the knockdown retinal axons using MitoSOX, a

mitochondrial superoxide indicator (Yarosh et al., 2008). The results showed significantly elevated

ROS levels in *dOPA1* knockdown retinal axons compared to control (Figure 2A and B; quantification

in 2C).

The *dOPA1* LOF leads to progressive distal degeneration of *Drosophila* photoreceptors

We tested whether knocking down *dOPA1* causes optic nerve degeneration in *Drosophila*. To this end,

we performed an RNAi experiment using the *GMR-Gal4* driver and evaluated the number of retinal

axons projecting to the second optic ganglion medulla (Figure 3A). To compare the number of retinal

axon terminals–R7 axons–between genotypes, we used a previously developed automated method,

MeDUsA (method for the quantification of degeneration using fly axons) (Nitta et al., 2023). Our

results showed that the *dOPA1* knockdown caused axonal degeneration one day after eclosion, further

decreasing one week later (Figure 3B–E, quantification in 3F). To determine whether the degeneration

was limited to the axons, we counted the number of rhabdomeres in the cell bodies. Each compound

eye comprises 700–800 ommatidia, with each ommatidium containing eight types of photoreceptors,

designated as R1–R8. The distal region of an ommatidium reveals the R1–R7 photoreceptors with a

stereotypical arrangement of rhabdomeres (arrowheads in Figure 3G–J). Since it is easy to evaluate

retinal degeneration by counting the number of rhabdomeres, a decrease in their number indicates

degeneration of the photoreceptor cell body. The *dOPA1* knockdown caused a significant decrease in

the number of rhabdomeres per ommatidium in 1-week-old adults (Figure 3G–J, quantification in 3K).

Interestingly, although the retinal axons had already decreased on the day of eclosion, the ommatidia

remained intact (Figure 3F and 3K). Note that the ommatidia in R7 remained intact after one week

(Figure 3J). Our results indicate that the *dOPA1* knockdown causes degeneration in both axonal

terminals and cell bodies, with the neuronal degeneration starting from the axons and continuing in

the cell bodies. This suggests that cell body loss precedes neurite retrograde degeneration, as described

in many neurodegenerative diseases (Wang et al., 2012).

To test whether our RNAi results reflect *dOPA1* downregulation, we performed mutant

analysis of the *dOPA1* gene in retinal axons. For this, we used the *dOPA1^{s3475}* hypomorphic mutant

allele, which has a P-element insertion in exon 2 (Yarosh et al., 2008). Somatic mosaic clones were

generated in the retinal axons using the FLP/FRT system (Golic and Lindquist, 1989) because

homozygous mutant alleles are embryonic lethal (Sandoval et al., 2014; Yarosh et al., 2008). Using

MeDUsA, we found a significantly lower number of *dOPA1* mutant retinal axons than in controls one

day after eclosion (Figure 4A and B, quantification in 4D). To determine if *dOPA1* is responsible for

axon neurodegeneration, we observed the *dOPA1* mutant axons by expressing full-length versions of

dOPA1 in the photoreceptors at one day after eclosion and found that dOPA1 expression significantly

rescued the axonal degeneration (Figure 4C, quantification in Figure 4D). Our results indicate that

dOPA1 is necessary for maintaining the number of retinal axons in the *Drosophila* visual system. In

conclusion, the neurodegeneration observed in the LOF of *dOPA1* is due to the progressive loss of

retinal axons, as well as the mammal optic nerve.

Investigating disease mutations in OPA1 using a fly DOA model to confirm their pathological

significance

Our observations in *dOPA1* LOF flies confirmed mitochondria fragmentation, increased ROS levels,

and degeneration of retinal axons, as reported in mammals. These data indicated that the function of

OPA1 may be conserved across *Drosophila* and humans. Thus, we tested whether *hOPA1* could

effectively replace *dOPA1* in flies by generating a transgenic organism to express *hOPA1* in

Drosophila. Following the UAS sequence, that the yeast-derived transcription factor Gal4 binds, the

construct included an HA tag, followed by the *hOPA1* gene, and a myc tag (Figure 5A). This construct

was expressed in *Drosophila* and protein expression was confirmed by Western blotting. The HA tag,

attached to the N-terminus, was immunoblotted resulting in a band of the expected full-length size

(Figure 5B). For the myc tag at the C-terminus, a strong band and two weak bands were observed at

the upper and lower positions (Figure 5B).

Using this UAS-hOpa1, we performed rescue experiments. *hOPA1* expression in the retinal

axons of the *dOPA1* mutant, which generated somatic cell clones, significantly rescued the number of

axons (Figure 5C). These results suggest that *hOPA1* and *dOPA1* are interchangeable. We also

generated and expressed the *hOPA1* mutation 2708_2711del, which is known to cause DOA, the

I382M mutation, located in the GTPase domain and associated with DOA, as well as the D438V and

R445H mutations, also located in the GTPase domain and associated with DOA plus. The 2708-

2711del mutation was weakly detectable by HA but not detectable by myc due to the frameshift caused

by the deletion (Fig. 5B). This appears to be a result from the mutation to generate a truncated protein,

as previously reported (Zanna et al., 2008). The I382M, D438V, and R445H mutations had similar

expression levels to the WT *hOPA1* (Figure 5B). However, the expression of these mutations in retinal

axons failed to rescue the *dOPA1* deficiency to the same extent as the WT *hOPA1* (Figure 5C). These

results suggest that these mutations result in functional impairment, consistent with the established

understanding (Zanna et al., 2008). Importantly, unlike the D438V and R445H mutations, the 2708-

2711del and I382M mutations could be weakly rescued. This suggests that these mutations result in

LOF, but may retain some residual activity. This is consistent with the characterization of I382M as a

hypomorph (Del Dotto et al., 2018). These findings are also in line with the observed severity of the

associated diseases, DOA and DOA plus. As the D438V and R445H mutations were not significantly

rescued, unlike the 2708-2711del and I382M mutations. Our results demonstrate that our model can

quantitatively measure the degree of LOF.

Distinction between LOF and DN mutations in *hOPA1* linked to DOA or DOA plus

Currently, the *hOPA1* mutations that contribute to DOA plus have been primarily located in the

GTPase domain of *hOPA1*. The role of *hOPA1* as a GTPase is facilitated by its ability to form a

polymer via the GED domain and Middle domain (Li et al., 2019). This led to the hypothesis that

mutations in the GTPase domain can interact with WT hOPA1 but cannot activate the GTPase activity

to show a DN effect. However, it has been challenging to clarify whether these mutations are DN or

LOF, as there is a significant number of LOF mutations in the GTPase domain. Given that *dOPA1* can

be substituted with *hOPA1*, we suspected that a DN effect could also be achieved by expressing the

D438V or R445H mutations in the WT background. However, the axons did not degenerate despite

expression and monitoring for two weeks in adult flies (Fig. 6A). The amino acid sequences of hOPA1

and dOPA1 are highly conserved (72% identity), particularly in the GTPase domain. The Middle and

GED domains are also relatively well-preserved, with 54% and 64% agreement, respectively (Figure

S2). These results imply that, although *dOPA1* and *hOPA1* are functionally interchangeable, they may

not interact with each other. Consequently, we expressed *hOPA1* in the photoreceptor depleted of

dOPA1 by somatic clone; under these conditions, we further expressed *hOPA1* WT, 2708-2711del,

and mutations in the GTPase domains, including R445H, I382M, and D438V. This allowed *hOPA1*

to interact with itself and to verify the DN effect. As a result, the expression of the WT, 2708del, and

I382M did not result in any changes when rescued by *hOPA1* substitution (Figure 6B). However, the

rescue was significantly suppressed for D438V and R445H, which are known to cause DN; thus, the

DN effect could be replicated (Figure 6B). In conclusion, we established an experimental model that

can separate LOF and DN, the pathological significance of *hOPA1* mutations.

Discussion

In this study, we found that LOF of *dOPA1* in *Drosophila* can imitate human DOA, in which the optic

nerve degenerates. We were successful in reversing this degeneration by expressing *hOPA1* (Figure

6C). However, we could not rescue any previously reported mutations known to cause either DOA or

DOA plus. In the context of rescuing the retinal axons of the *dOPA1* mutant by expressing the WT of

hOPA1, it was observed that only the DOA plus mutations suppressed the rescue (Figure 6C). This

allowed us to distinguish between LOF and DN mutations in *hOPA1*. The fly model developed in this

study will allow investigating new *hOPA1* mutations for DOA or DOA plus.

Our established fly model is the first simple organism allowing observation of degeneration

of axons in the optic nerve and fragmentation of mitochondria. Previously, the function of *OPA1* in

mice (*mOPA1*) was substituted by *hOPA1* (Del Dotto et al., 2017; Sarzi et al., 2018). Here we

demonstrated that the function of *dOPA1* can also be substituted by *hOPA1*. Thus, the function of

OPA1 is conserved across a wide range of species but our fly model has the advantage of allowing

rapid observation of phenotypes. In the mouse model of DOA with the heterozygous deletion of the

mOPA1 gene at positions 329–355 (Alavi et al., 2007) or delTTAG mutant (Sarzi et al., 2012), a

decrease in the number of RGC was observed after 17 months or 16 months of age, respectively. To

save time, using a fly model for quick analysis and direct observation of retinal axons would be useful.

Previous models using nematodes, flies, and zebrafish did not observe axonal degeneration directly,

but our new model has made this possible. As a result, our model has the potential to be used for

screening for modifiers of unclear molecular pathologies and drug screening.

Our model could also demonstrate the pathological significance of *hOPA1* mutations and be

used for genetic diagnosis to differentiate between LOF and DN mutations. Our results for LOF or

DN segmentation were consistent with previously reported yeast models (Del Dotto et al., 2018) and

fibroblasts obtained from patients with OPA1 mutations (Kane et al., 2017), and confirmed the

conserved function of *OPA1* across species. A major advantage of the *Drosophila* model over yeast

or fibroblasts, is that we can observe a phenotype that mimics the actual disease state of axonal

degeneration in retinal axons and analyze it not only at the molecular level, but also the structural level

of axons and mitochondria. The transgenic *Drosophila* of disease-associated *hOpa1* mutations created

in this study can also serve to understand the pathophysiology of these mutations.

Advances in genome analysis allowed the identification of many variants and genetic

mutations associated with DOA using next-generation sequencing. This could confirm diagnoses and

identify new forms of DOA. The widespread use of these sequencers has also created new challenges,

such as the growing number of variants of unknown significance (VUS) in publicly available databases

such as CliniVar for DOA. These variants are categorized as pathogenic, likely pathogenic, VUS,

likely benign, or benign, and the number of VUS is increasing as more variants are registered. The

ClinVar database had classified 110 out of 319 *Opa1* variants as VUS at the end of 2019. By the end

of 2022, this number had increased to 357 out of 840, indicating an increase from 34% to 42% over

the last three years (Henrie et al., 2018). This trend suggests that the number of VUS in *Opa1* may

continue to increase in the future. Analyzing VUS requires testing the effects of a mutation *in vitro* or

in vivo; this can be expensive and time-consuming without prior confirmation or high probability of

pathological significance. However, our fly model can be used to easily and inexpensively determine

the pathological effects of disease-causing gene mutations among the growing number of rare

mutations identified in DOA.

In this study, we tested the effect of expressing the D438V and R445H mutations in the

dOPA1 background and found that retinal axons did not degenerate. Regarding the interactions among

OPA1 proteins, in yeast Mgm1, K854 in the region near the C-terminus is important for aggregation,

and it becomes an oligomer to increase the activity of GTPase (Rujiviphat et al., 2009). Although

hOPA1 normally has low GTPase activity, its interaction with negatively charged phospholipids such

as cardiolipin causes hOPA1 to aggregate, increasing GTP hydrolysis activity (Ban et al., 2010). In

the thermophilic fungus *Chaetomium thermophilum*, Mgm1 forms a dimer and then a tetramer at the

C-terminal Stalk domain of Mgm1 (Faelber et al., 2019). Structural analysis also predicted that the

Middle and GED domains are necessary for dimer and oligomer formation in hOPA1 (Li et al., 2019).

In addition, peptide-binding assays have revealed that the coiled-coil domain, which is part of the

GED domain, is required for hOPA1 interactions (Akepati et al., 2008). Comparing the amino acid

sequences of hOPA1 and dOPA1, the Middle and GED domains, which are necessary for interactions

between OPA1 molecules, exhibit concordance rates of 54.3% and 63.6%, respectively. These

percentages are low compared to the 71.8% identity observed in the GTPase domains. These suggest

that the OPA1 homologs of each species form oligomers with their unique OPA1. Note that the

expression of *dOPA1^{K273A}*, a presumably GTPase-negative form of *dOPA1* (Tsuyama et al., 2017),

degenerated retinal axons (data not shown). These results imply that, while dOPA1 and hOPA1 are

interchangeable, they may not interact with each other. This could pave the way for the development

of a chimeric hOPA1, which would retain its functional properties while avoiding the interaction with

endogenous hOPA1. Such chimeric hOPA1 could potentially be used as gene therapy since induced

pluripotent stem cells have been generated from a patient with R445H (Sladen et al., 2021).

OPA1 has been implicated in various other diseases as well, including normal tension

glaucoma (Powell et al., 2003; Yu-Wai-Man et al., 2010b), multiple sclerosis (Verny et al., 2008; Yu-

Wai-Man et al., 2016), Parkinson's disease and dementia (Carelli et al., 2015; Lynch et al., 2017), and

cardiomyopathy (Spiegel et al., 2016). Although the connection between these diseases and OPA1 is

not fully understood, further research on DOA may provide a deeper insight into this relationship. In

the future, pathological analysis using the present model could have implications for understanding

the mechanisms underlying these diseases.

DOA has also been reported to involve mitochondrial DNA (mtDNA) depletion (Amati-

Bonneau et al., 2008), but our model, while useful for nuclear gene analysis, is limited for mtDNA

analysis. The gene content in the *Drosophila* mtDNA genome is similar as that in vertebrates, but the

gene order and distribution on both DNA strands differ. mOPA1 interacts with mtDNA nucleoids

through exon 4b binding to mtDNA D-loops, independent of mitochondrial fusion (Yang et al., 2020).

However, the homologous region of mOPA1 or hOPA1 to exon4b is not found in *Drosophila*, and the

D-loop of mtDNA has not been found in cultured cells of flies or in mtDNA of embryos using electron

microscopy (Rubenstein et al., 1977). In addition, while the D-loop is involved in mtDNA replication

in humans (Fish et al., 2004), the A + T region is the origin of replication in mtDNA replication in

Drosophila (Goddard and Wolstenholme, 1978). Thus, there may be differences in the mechanism of

mtDNA replication between humans and flies (Garesse and Kaguni, 2005). Therefore, the association

of dOPA1 with mtDNA may differ from that in mammals and the regulatory mechanism of mtDNA

homeostasis in which dOPA1 is involved requires further investigation.

334   Another limitation of our model is that the molecular mechanisms underlying the formation

of long-form OPA1 (L-OPA1) and short-form OPA1 (S-OPA1), which are well analyzed in mammals

(MacVicar and Langer, 2016), have not been elucidated in *Drosophila*. In hOPA1, S1 or S2 sites are

processed by the i-AAA protease OMA1 and Yme1L respectively to produce S-hOPA1 (Anand et al.,

2014). Despite the fact that Yme1L is conserved in *Drosophila*, it has been involved in the degradation

of the dOPA1 protein rather than its cleavage by dYme1L (Liu et al., 2020). Moreover, the cleavage

site for i-AAA does not exist in dOPA1 (Olichon et al., 2007). Regarding other proteases, yeast Mgm1

is processed by the rhomboid protease Pcp1 (Esser et al., 2002; Herlan et al., 2003). However, the

Rhomboid-7, a Pcp1 ortholog of *Drosophila*, was not required for dOPA1 processing (Whitworth et

al., 2008). Although hOPA1 expression in *Drosophila* has identified three different sizes of hOPA1

in our Western blotting result, it is unclear how they are cleaved. The upper band was expected as the

full-length hOPA1 including the mitochondria import sequence (MIS). Since the Middle and the

lowest bands were not present in HA-tagged samples, it is possible that the bands at the Middle size

represent the long-form hOPA1 (L-hOPA1) in which probably MIS was processed. The band detected

at the lowest position may represent a short form of hOPA1 (S-hOPA1). However, *Drosophila* does

not have OMA1, an i-AAA protease that cleaves the S1 site. Although another i-AAA protease that

cleaves the S2 site, YME1L, is conserved, our expressed hOPA1 is isoform 1 and lacks the S2 site

(Fig. 5a). Therefore, it is not clear how S-hOPA1 is processed in *Drosophila*. The anti-dOPA1

antiserum detects three bands in the overexpression of a FLAG-tagged dOPA1 construct (Poole et al.,

2010), suggesting a molecular mechanism by which dOPA1, like yeast and mammals, is cleaved after

translation. However, the proteases involved in the mechanism are not yet clarified. Whether

Drosophila's endogenous mechanisms can be used to study how L-OPA1 and S-OPA1 are involved

in DOA remains unclear.

357   In this study, we established a new model of DOA in *Drosophila*, in which we discovered

the following: 1. We could replicate the human disease of optic nerve degeneration, allowing rapid

genetic disease analysis. 2. We could distinguish the LOF and DN effects of *hOPA1*. These findings

can reveal the pathological significance of *de novo* mutations and are useful in diagnosing DOA or

DOA plus. Additionally, while *hOPA1* is the major gene involved in DOA, other genes involved in

DOA and interacting proteins have not been investigated yet. Our model can be used for pathological

modifier screening, exploring modifiers, and contributing to drug development. In the future, we hope

to provide insights that can be applied to the fundamental treatment of DOA.

Materials and Methods

Fly strains

Flies were maintained at a temperature of 25°C on standard fly food. For the knockdown experiments

(Figure 1C–E, 1F–H, 2A–C, 3B–K), flies were maintained at 29°C. Female flies were used in all

experiments to ensure consistency in the number of retinal axons. The following fly strains were

obtained from the Bloomington Drosophila Stock Center: *GMR-Gal4* (#1104), *UAS-dicer2(III)*

(#24651), *UAS-mito-HA-GFP* (#8443), *UAS-mCherry-mito* (#66532), *UAS-dOPA1 RNAi* (#32358),

UAS-Milton RNAi (#44477), *FRT42D, GMR-hid, l(2)cl-R11[1]/CyO; ey-Gal4, UAS-flp* (#5251), *ey-*

FLP; FRT42D, l(2)cl-R11[1]/CyO (#5617), and *Tub-Gal80^{TS}* (#7017). The *40D-UAS* (VDRC 60101)

fly strain was obtained from the Vienna Drosophila Resource Center. *FRT19* (Kyoto 101231),

FRT42D (Kyoto 101878), and *dOPA1^{s3475}* (Kyoto 111438) were obtained from the Kyoto Drosophila

Stock Center. The *UAS-dOPA1-HA* was generously provided by T. Uemura (Kyoto University, Japan).

Generation of transgenic flies

The cDNA for WT, I382M, D438V, R445H, and 2708-2711del forms of *hOPA1* [NM_015560.3] were

generated by Vectorbuilder Inc. (Yokohama, Japan). Then, the HA-hOPA1-myc sequence was

amplified and inserted into pJFRC81-10XUAS-IVS-Syn21-GFP-p10 (ID36432, Addgene,

Massachusetts, USA) using primers with the N-terminal containing the Kozak and HA sequences

(TTACTTCAGGCGGCCGCGGCCAAAATGTACCCATACGATGTTCCAGATTAC) and the C-

terminal   containing   the   myc   sequence

(TTAAAAACGATTCATTCTAGTTACAGATCCTCTTCTGAGATGAGTTTTTGTTCTTTCTCCT

GATGAAGAGCTTCAATG). To confirm production of the complete coding sequence, it was

bidirectionally sequenced using the Sanger method. These plasmids were injected into embryos and

integrated into the aTTP2 landing site (WellGenetics, Taipei, Taiwan).

Immunohistochemistry and imaging

Immunohistochemistry was performed as described previously (Sugie et al., 2017). The following

antibodies were used: rat anti-HA (3F10, 1:400; Roche, Switzer land), mouse anti-Chaoptin (24B10,

1:25; Developmental Studies Hybridoma Bank, USA), rat anti-CadN (DN-Ex#8, 1:50; Developmental

Studies Hybridoma Bank), anti-rat Alexa Fluor 488 (1:400; Thermo Fisher Scientific, USA), anti-

mouse Alexa Fluor 488 (1:400; Thermo Fisher Scientific), anti-mouse Alexa Fluor 568 (1:400;

Thermo Fisher Scientific), and anti-rat Alexa Fluor 633 (1:400; Thermo Fisher Scientific). Specimens

were mounted using a Vectashield mounting medium (Vector Laboratories, USA). Images were

captured using an FV3000 confocal microscope (Olympus, Japan) or a C2 (Nikon, Japan) confocal

microscope for Figure S1 and processed using the IMARIS software (Oxford Instruments, UK) or Fiji

software, an open-source image analysis software (Schindelin et al., 2012).

Electron microscopy

Fly heads were fixed overnight with 2.5% glutaraldehyde and 2% paraformaldehyde in 0.1 M sodium

cacodylate buffer. Heads were then post-fixed with 1% osmium tetroxide, dehydrated in ethanol, and

embedded in Epon. Ultra-thin tangential sections of the laminas (70 nm) were stained with uranyl

acetate and lead citrate. Random retinas or lamina sections were imaged at a magnification of ×100 k

with a VELETA CCD Camera (Olympus Soft Imaging Solutions) mounted on a JEM 1010

transmission electron microscope (JEOL). The area corresponding to the mitochondria in the R1–6

axons of the lamina was measured using the freehand line tool in NIS elements (Nikon).

Quantification of mito-GFP intensity in retinal axons and retina

To measure the relative intensity of mito-GFP in the photoreceptor R7/8 axons of the medulla, the

imaging analysis software IMARIS was used. First, the M1–6 layers of the medulla, including the

R7/8 axons, were manually covered using the surface function. Then, the surface area was masked,

the surface of the left axons was stained with anti-Chaoptin, and the mito-GFP signal was generated

automatically using the surface function. Finally, the voxel number of the mito-GFP surface was

divided by the voxel number of the R axon surface. These quantifications were performed by

experimenters blind to the genotype.

Mitochondrial superoxide detection and quantification

The ROS level was determined using the MitoSox Red Mitochondrial Superoxide Indicator

(Invitrogen). The procedure involved dissecting adult brains in phosphate-buffered saline (PBS),

transferring them to 5 µM MitoSox solution, and incubating them for 10 min in the dark at room

temperature. The brains were then washed with PBS containing 20% Vectashield (Vector Laboratories),

followed by sequential washes with PBS containing 40% and 60% Vectashield. The samples were

mounted in Vectashield and scanned using an FV3000 confocal microscope (Olympus). To quantify

the ROS level in photoreceptor R7/8 axons, the IMARIS software (Bitplane) was used for imaging

analysis. The software generated automatic surfaces of the MitoSox and mito-GFP signals. Then, the

ROS level was calculated by dividing the voxel number of the MitoSox surface on the mito-GFP

surface by the voxel number of the mito-GFP surface. These quantifications were performed by

experimenters blind to the genotype.

Western blot analysis of the whole Drosophila body

Protein expression was temporarily controlled in the whole body by combining *Tub-Gal4* and *Tub-*

GAL80^{TS}. The flies crossed with each hOPA1 variant were raised at a permissive temperature (20°C),

and newly eclosed females were held at the restrictive temperature (29°C). After two days, 10 whole

bodies were sonicated (2 x 30 s) using a Q55 sonicator (Qsonica, Newtown, CT, USA) in lysis buffer

(10 mM Tris–HCl, pH 7.5, 150 mM NaCl, 1 mM EDTA, 2% DDM) supplemented with 1:1000 (v/v)

Protease Inhibitor Cocktail Set III (Calbiochem, La Jolla, CA, USA). Rat anti-HA (3F10, 1:5000;

Roche, Mannheim, Germany), mouse anti-myc (9B11, 1: 10000; Cell Signaling Technology, Danvers,

MA, USA), and mouse anti-α-Tubulin (T9026, 1:100,000; Sigma‑Aldrich, St. Louis, MO, USA)

antibodies were used for Western blotting.

Pairwise alignment of hOPA1 and dOPA1

The amino acid sequences for hOPA1 and dOPA1 were referenced from NM_015560.3 and

NM_166040.2, respectively, and aligned using the EMBOSS Needle. The GTPase, Middle, and GED

domains in dOPA1 were identified by cross-referencing the previously described amino acid positions

of these domains in hOPA1 (Liesa et al., 2009). The degree of identity between the aligned amino acid

sequences was subsequently determined.

Experimental design and statistical analyses

Experimental analyses were performed using Prism 9 (GraphPad Software, San Diego, CA, USA).

For Figures 3–5, data were analyzed using t tests (and nonparametric tests) and Mann-Whitney tests.

For Figures 1, 2, 6, and 7, differences among multiple groups were examined using the Kruskal−Wallis

test followed by the two-stage linear step-up procedure of Benjamini, Krieger, and Yekutieli tests. The

null hypothesis was rejected at a 0.05 level of significance. Sample sizes are indicated in each figure.

The number of retinal axons and rhabdomeres, the intensity of mito-GFP and MitoSox, the

mitochondrial area and the classification, and the corrected P values are described in the Results

section. All data are expressed as the mean ± SEM.

Acknowledgements

We would like to thank Dr Uemura for providing us with the transgenic fly strain. The Ministry of

Education, Culture, Sports, Science, and Technology of Japan (#18K14835, #18J00367 and

#21K15619 to Y.N., #21K06184 to S.H.S., #16H06457, #21H02483 and 21H05682 to T.S.,

#17H04983, #19K22592 and #21H02837 to A.S.), the Japan Agency for Medical Research and

Development (AMED) (#JP22ek019484s) to A.S., NIG–JOINT to E.S., Takeda Science Foundation

Takeda Visionary Research Grant to T.S., Takeda Science Foundation Life Science Research Grant to

A.S., and GSK Japan Research Grant 2022 [AS2021A000166849] to A.S..

References

Akepati VR, Müller EC, Otto A, Strauss HM, Portwich M, Alexander C. 2008. Characterization of OPA1

isoforms isolated from mouse tissues. *J Neurochem* **106**:372–383. doi:10.1111/j.1471-

4159.2008.05401.x

Alavi M V., Bette S, Schimpf S, Schuettauf F, Schraermeyer U, Wehrl HF, Ruttiger L, Beck SC, Tonagel

F, Pichler BJ, Knipper M, Peters T, Laufs J, Wissinger B. 2007. A splice site mutation in the

murine Opa1 gene features pathology of autosomal dominant optic atrophy. *Brain* **130**:1029–1042.

doi:10.1093/brain/awm005

Alexander C, Votruba M, Pesch UEA, Thiselton DL, Mayer S, Moore A, Rodriguez M, Kellner U, Leo-

Kottler B, Auburger G, Bhattacharya SS, Wissinger B. 2000. OPA1, encoding a dynamin-related

GTPase, is mutated in autosomal dominant optic atrophy linked to chromosome 3q28. *Nat Genet*

26:211–215. doi:10.1038/79944

Amati-Bonneau P, Odent S, Derrien C, Pasquier L, Malthiéry Y, Reynier P, Bonneau D. 2003. The

association of autosomal dominant optic atrophy and moderate deafness may be due to the R445H

mutation in the OPA1 gene. *Am J Ophthalmol* **136**:1170–1171. doi:10.1016/S0002-9394(03)00665-

2

Amati-Bonneau P, Valentino ML, Reynier P, Gallardo ME, Bornstein B, Boissière A, Campos Y, Rivera

H, De La Aleja JG, Carroccia R, Iommarini L, Labauge P, Figarella-Branger D, Marcorelles P,

Furby A, Beauvais K, Letournel F, Liguori R, La Morgia C, Montagna P, Liguori M, Zanna C,

Rugolo M, Cossarizza A, Wissinger B, Verny C, Schwarzenbacher R, Martín MÁ, Arenas J, Ayuso

C, Garesse R, Lenaers G, Bonneau D, Carelli V. 2008. OPA1 mutations induce mitochondrial DNA

instability and optic atrophy "plus" phenotypes. *Brain* **131**:338–351. doi:10.1093/brain/awm298

Anand R, Wai T, Baker MJ, Kladt N, Schauss AC, Rugarli E, Langer T. 2014. The i-AAA protease

YME1L and OMA1 cleave OPA1 to balance mitochondrial fusion and fission. *J Cell Biol* **204**:919–

929. doi:10.1083/jcb.201308006

Ban T, Heymann JAW, Song Z, Hinshaw JE, Chan DC. 2010. OPA1 disease alleles causing dominant

optic atrophy have defects in cardiolipin-stimulated GTP hydrolysis and membrane tubulation.

Hum Mol Genet **19**:2113–2122. doi:10.1093/hmg/ddq088

Brand AH, Perrimon N. 1993. Targeted gene expression as a means of altering cell fates and generating

dominant phenotypes. *Development* **118**:401–415. doi:10.1242/dev.118.2.401

Byrne JJ, Soh MS, Chandhok G, Vijayaraghavan T, Teoh JS, Crawford S, Cobham AE, Yapa NMB,

Mirth CK, Neumann B. 2019. Disruption of mitochondrial dynamics affects behaviour and lifespan

in Caenorhabditis elegans. *Cell Mol Life Sci* **76**:1967–1985. doi:10.1007/s00018-019-03024-5

Carelli V, Musumeci O, Caporali L, Zanna C, La Morgia C, Del Dotto V, Porcelli AM, Rugolo M,

Valentino ML, Iommarini L, Maresca A, Barboni P, Carbonelli M, Trombetta C, Valente EM,

Patergnani S, Giorgi C, Pinton P, Rizzo G, Tonon C, Lodi R, Avoni P, Liguori R, Baruzzi A,

Toscano A, Zeviani M. 2015. Syndromic parkinsonism and dementia associated with OPA1

missense mutations. *Ann Neurol* **78**:21–38. doi:10.1002/ana.24410

Davies VJ, Hollins AJ, Piechota MJ, Yip W, Davies JR, White KE, Nicols PP, Boulton ME, Votruba M.

2007. Opa1 deficiency in a mouse model of autosomal dominant optic atrophy impairs

mitochondrial morphology, optic nerve structure and visual function. *Hum Mol Genet* **16**:1307–

1318. doi:10.1093/hmg/ddm079

Del Dotto V, Carelli V. 2021. Dominant Optic Atrophy (DOA): Modeling the Kaleidoscopic Roles of

OPA1 in Mitochondrial Homeostasis. *Front Neurol*. doi:10.3389/fneur.2021.681326

Del Dotto V, Fogazza M, Musiani F, Maresca A, Aleo SJ, Caporali L, La Morgia C, Nolli C, Lodi T,

Goffrini P, Chan D, Carelli V, Rugolo M, Baruffini E, Zanna C. 2018. Deciphering OPA1

mutations pathogenicity by combined analysis of human, mouse and yeast cell models. *Biochim*

Biophys Acta - Mol Basis Dis **1864**:3496–3514. doi:10.1016/j.bbadis.2018.08.004

Del Dotto V, Mishra P, Vidoni S, Fogazza M, Maresca A, Caporali L, McCaffery JM, Cappelletti M,

Baruffini E, Lenaers G, Chan D, Rugolo M, Carelli V, Zanna C. 2017. OPA1 Isoforms in the

Hierarchical Organization of Mitochondrial Functions. *Cell Rep* **19**:2557–2571.

doi:10.1016/j.celrep.2017.05.073

Delettre C, Lenaers G, Griffoin JM, Gigarel N, Lorenzo C, Belenguer P, Pelloquin L, Grosgeorge J, Turc-

Carel C, Perret E, Astarie-Dequeker C, Lasquellec L, Arnaud B, Ducommun B, Kaplan J, Hamel

CP. 2000. Nuclear gene OPA1, encoding a mitochondrial dynamin-related protein, is mutated in

dominant optic atrophy. *Nat Genet* **26**:207–210. doi:10.1038/79936

Deng H, Dodson MW, Huang H, Guo M. 2008. The Parkinson's disease genes pink1 and parkin promote

mitochondrial fission and/or inhibit fusion in Drosophila. *Proc Natl Acad Sci U S A* **105**:14503–

14508. doi:10.1073/pnas.0803998105

Eijkenboom I, Vanoevelen JM, Hoeijmakers JGJ, Wijnen I, Gerards M, Faber CG, Smeets HJM. 2019. A

zebrafish model to study small-fiber neuropathy reveals a potential role for GDAP1. *Mitochondrion*

47:273–281. doi:10.1016/j.mito.2019.01.002

Esser K, Tursun B, Ingenhoven M, Michaelis G, Pratje E. 2002. A novel two-step mechanism for

removal of a mitochondrial signal sequence involves the mAAA complex and the putative

rhomboid protease Pcp1. *J Mol Biol* **323**:835–843. doi:10.1016/S0022-2836(02)01000-8

Faelber K, Dietrich L, Noel JK, Wollweber F, Pfitzner AK, Mühleip A, Sánchez R, Kudryashev M,

Chiaruttini N, Lilie H, Schlegel J, Rosenbaum E, Hessenberger M, Matthaeus C, Kunz S, von der

538   Malsburg A, Noé F, Roux A, van der Laan M, Kühlbrandt W, Daumke O. 2019. Structure and

539   assembly of the mitochondrial membrane remodelling GTPase Mgm1. *Nature* **571**:429–433.

540   doi:10.1038/s41586-019-1372-3

541 Ferré M, Caignard A, Milea D, Leruez S, Cassereau J, Chevrollier A, Amati-Bonneau P, Verny C,

542   Bonneau D, Procaccio V, Reynier P. 2015. Improved locus-specific database for OPA1 mutations

543   allows inclusion of advanced clinical data. *Hum Mutat* **36**:20–25. doi:10.1002/humu.22703

544 Fish J, Raule N, Attardi G. 2004. Discovery of a major D-loop replication origin reveals two modes of

545   human mtDNA synthesis. *Science (80-)* **306**:2098–2101. doi:10.1126/science.1102077

546 Frezza C, Cipolat S, Martins de Brito O, Micaroni M, Beznoussenko G V., Rudka T, Bartoli D, Polishuck

547   RS, Danial NN, De Strooper B, Scorrano L. 2006. OPA1 Controls Apoptotic Cristae Remodeling

548   Independently from Mitochondrial Fusion. *Cell* **126**:177–189. doi:10.1016/j.cell.2006.06.025

549 Garesse R, Kaguni LS. 2005. A Drosophila model of mitochondrial DNA replication: Proteins, genes and

550   regulation. *IUBMB Life*. doi:10.1080/15216540500215572

551 Goddard JM, Wolstenholme DR. 1978. Origin and direction of replication in mitochondrial DNA

552   molecules from Drosophila melanogaster. *Proc Natl Acad Sci U S A* **75**:3886–3890.

553   doi:10.1073/pnas.75.8.3886

554 Golic KG, Lindquist S. 1989. The FLP recombinase of yeast catalyzes site-specific recombination in the

555   drosophila genome. *Cell* **59**:499–509. doi:10.1016/0092-8674(89)90033-0

Henrie A, Hemphill SE, Ruiz-Schultz N, Cushman B, DiStefano MT, Azzariti D, Harrison SM, Rehm

HL, Eilbeck K. 2018. ClinVar Miner: Demonstrating utility of a Web-based tool for viewing and

filtering ClinVar data. *Hum Mutat* **39**:1051–1060. doi:10.1002/humu.23555

Herlan M, Vogel F, Bornhövd C, Neupert W, Reichert AS. 2003. Processing of Mgm1 by the rhomboid-

type protease Pcp1 is required for maintenance of mitochondrial morphology and of mitochondrial

DNA. *J Biol Chem* **278**:27781–27788. doi:10.1074/jbc.M211311200

Kanazawa T, Zappaterra MD, Hasegawa A, Wright AP, Newman-Smith ED, Buttle KF, McDonald K,

Mannella CA, Van Der Bliek AM. 2008. The C. elegans Opa1 homologue EAT-3 is essential for

resistance to free radicals. *PLoS Genet* **4**. doi:10.1371/journal.pgen.1000022

Kane MS, Alban J, Desquiret-Dumas V, Gueguen N, Ishak L, Ferre M, Amati-Bonneau P, Procaccio V,

Bonneau D, Lenaers G, Reynier P, Chevrollier A. 2017. Autophagy controls the pathogenicity of

OPA1 mutations in dominant optic atrophy. *J Cell Mol Med* **21**:2284–2297.

doi:10.1111/jcmm.13149

Lenaers G, Neutzner A, Le Dantec Y, Jüschke C, Xiao T, Decembrini S, Swirski S, Kieninger S, Agca C,

Kim US, Reynier P, Yu-Wai-Man P, Neidhardt J, Wissinger B. 2021. Dominant optic atrophy:

Culprit mitochondria in the optic nerve. *Prog Retin Eye Res*. doi:10.1016/j.preteyeres.2020.100935

Li D, Wang J, Jin Z, Zhang Z. 2019. Structural and evolutionary characteristics of dynamin-related

GTPase OPA1. *PeerJ* **2019**. doi:10.7717/peerj.7285

Liesa M, Palacín M, Zorzano A. 2009. Mitochondrial dynamics in mammalian health and disease.

Physiol Rev. doi:10.1152/physrev.00030.2008

Liu W, Duan X, Xu L, Shang W, Zhao J, Wang L, Li JC, Chen CH, Liu JP, Tong C. 2020. Chchd2

regulates mitochondrial morphology by modulating the levels of Opa1. *Cell Death Differ* **27**:2014–

2029. doi:10.1038/s41418-019-0482-7

Lynch DS, Loh SHY, Harley J, Noyce AJ, Martins LM, Wood NW, Houlden H, Plun-Favreau H. 2017.

Clinical/Scientific Notes: Nonsyndromic Parkinson disease in a family with autosomal dominant

optic atrophy due to opa1 mutations. *Neurol Genet* **3**. doi:10.1212/NXG.0000000000000188

MacVicar T, Langer T. 2016. OPA1 processing in cell death and disease - the long and short of it. *J Cell

Sci*. doi:10.1242/jcs.159186

McQuibban GA, Lee JR, Zheng L, Juusola M, Freeman M. 2006. Normal Mitochondrial Dynamics

Requires Rhomboid-7 and Affects Drosophila Lifespan and Neuronal Function. *Curr Biol* **16**:982–

989. doi:10.1016/j.cub.2006.03.062

Meglei G, McQuibban GA. 2009. The dynamin-related protein Mgm1p assembles into oligomers and

hydrolyzes GTP to function in mitochondrial membrane fusion. *Biochemistry* **48**:1774–1784.

doi:10.1021/bi801723d

Newsome TP, Åsling B, Dickson BJ. 2000a. Analysis of Drosophila photoreceptor axon guidance in eye-

specific mosaics. *Development* **127**:851–860. doi:10.1242/dev.127.4.851

Newsome TP, Schmidt S, Dietzl G, Keleman K, Åsling B, Debant A, Dickson BJ. 2000b. Trio combines

with dock to regulate Pak activity during photoreceptor axon pathfinding in Drosophila. *Cell*

101:283–294. doi:10.1016/S0092-8674(00)80838-7

Nguyen D, Alavi M V., Kim KY, Kang T, Scott RT, Noh YH, Lindsey JD, Wissinger B, Ellisman MH,

Weinreb RN, Perkins GA, Ju WK. 2011. A new vicious cycle involving glutamate excitotoxicity,

oxidative stress and mitochondrial dynamics. *Cell Death Dis* **2**. doi:10.1038/cddis.2011.117

Nitta Y, Kawai H, Maki R, Osaka J, Hakeda-Suzuki S, Nagai Y, Doubková K, Uehara T, Watanabe K,

Kosaki K, Suzuki T, Tavosanis G, Sugie A. 2023.   Direct evaluation of neuroaxonal degeneration

with the causative genes of neurodegenerative diseases in Drosophila using the automated axon

quantification system, MeDUsA . *Hum Mol Genet*. doi:10.1093/hmg/ddac307

Olichon A, ElAchouri G, Baricault L, Delettre C, Belenguer P, Lenaers G. 2007. OPA1 alternate splicing

uncouples an evolutionary conserved function in mitochondrial fusion from a vertebrate restricted

function in apoptosis. *Cell Death Differ* **14**:682–692. doi:10.1038/sj.cdd.4402048

Olichon A, Emorine LJ, Descoins E, Pelloquin L, Brichese L, Gas N, Guillou E, Delettre C, Valette A,

Hamel CP, Ducommun B, Lenaers G, Belenguer P. 2002. The human dynamin-related protein

OPA1 is anchored to the mitochondrial inner membrane facing the inter-membrane space. *FEBS*

Lett **523**:171–176. doi:10.1016/S0014-5793(02)02985-X

Olichon A, Guillou E, Delettre C, Landes T, Arnauné-Pelloquin L, Emorine LJ, Mils V, Daloyau M,

Hamel C, Amati-Bonneau P, Bonneau D, Reynier P, Lenaers G, Belenguer P. 2006. Mitochondrial

dynamics and disease, OPA1. *Biochim Biophys Acta - Mol Cell Res*.

doi:10.1016/j.bbamcr.2006.04.003

Poole AC, Thomas RE, Yu S, Vincow ES, Pallanck L. 2010. The mitochondrial fusion-promoting factor

mitofusin is a substrate of the PINK1/parkin pathway. *PLoS One* **5**.

doi:10.1371/journal.pone.0010054

Powell BL, Toomes C, Scott S, Yeung A, Marchbank NJ, Spry PGD, Lumb R, Inglehearn CF, Churchill

AJ. 2003. Polymorphisms in OPA1 are associated with normal tension glaucoma. *Mol Vis* **9**:460–

464.

Rahn JJ, Stackley KD, Chan SSL. 2013. Opa1 Is Required for Proper Mitochondrial Metabolism in Early

Development. *PLoS One* **8**. doi:10.1371/journal.pone.0059218

Richard M, Doubková K, Nitta Y, Kawai H, Sugie A, Tavosanis G. 2022. A Quantitative Model of

Sporadic Axonal Degeneration in the Drosophila Visual System. *J Neurosci* **42**:4937–4952.

doi:10.1523/JNEUROSCI.2115-21.2022

Rolland SG, Lu Y, David CN, Conradt B. 2009. The BCL-2-like protein CED-9 of C. elegans promotes

FZO-1/Mfn1,2- and EAT-3/Opa1-dependent mitochondrial fusion. *J Cell Biol* **186**:525–540.

doi:10.1083/jcb.200905070

Rubenstein JLR, Brutlag D, Clayton DA. 1977. The mitochondrial DNA of drosophila melanogaster

exists in two distinct and stable superhelical forms. *Cell* **12**:471–482. doi:10.1016/0092-

8674(77)90123-4

Rujiviphat J, Meglei G, Rubinstein JL, McQuibban GA. 2009. Phospholipid association is essential for

dynamin-related protein Mgm1 to function in mitochondrial membrane fusion. *J Biol Chem*

284:28682–28686. doi:10.1074/jbc.M109.044933

Sandoval H, Yao CK, Chen K, Jaiswal M, Donti T, Lin YQ, Bayat V, Xiong B, Zhang K, David G,

Charng WL, Yamamoto S, Duraine L, Graham BH, Bellen HJ. 2014. Mitochondrial fusion but not

fission regulates larval growth and synaptic development through steroid hormone production. *Elife*

3:1–23. doi:10.7554/eLife.03558

Sanes JR, Zipursky SL. 2020. Synaptic Specificity, Recognition Molecules, and Assembly of Neural

Circuits. *Cell*. doi:10.1016/j.cell.2020.04.008

Sanes JR, Zipursky SL. 2010. Design Principles of Insect and Vertebrate Visual Systems. *Neuron*.

doi:10.1016/j.neuron.2010.01.018

Sarzi E, Angebault C, Seveno M, Gueguen N, Chaix B, Bielicki G, Boddaert N, Mausset-Bonnefont AL,

Cazevieille C, Rigau V, Renou JP, Wang J, Delettre C, Brabet P, Puel JL, Hamel CP, Reynier P,

Lenaers G. 2012. The human OPA1delTTAG mutation induces premature age-related systemic

neurodegeneration in mouse. *Brain* **135**:3599–3613. doi:10.1093/brain/aws303

Sarzi E, Seveno M, Piro-Mégy C, Elzière L, Quilès M, Péquignot M, Müller A, Hamel CP, Lenaers G,

Delettre C. 2018. OPA1 gene therapy prevents retinal ganglion cell loss in a Dominant Optic

Atrophy mouse model. *Sci Rep* **8**. doi:10.1038/s41598-018-20838-8

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C,

Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A.

2012. Fiji: An open-source platform for biological-image analysis. *Nat Methods*.

doi:10.1038/nmeth.2019

Shahrestani P, Leung HT, Le PK, Pak WL, Tse S, Ocorr K, Huang T. 2009. Heterozygous mutation of

Drosophila Opa1 causes the development of multiple organ abnormalities in an age-dependent and

organ-specific manner. *PLoS One* **4**. doi:10.1371/journal.pone.0006867

Shimizu S, Mori N, Kishi M, Sugata H, Tsuda A, Kubota N. 2003. A novel mutation in the OPA1 gene in

a Japanese patient with optic atrophy. *Am J Ophthalmol* **135**:256–257. doi:10.1016/S0002-

9394(02)01929-3

Sladen PE, Perdigão PRL, Salsbury G, Novoselova T, van der Spuy J, Chapple JP, Yu-Wai-Man P,

Cheetham ME. 2021. CRISPR-Cas9 correction of OPA1 c.1334G>A: p.R445H restores

mitochondrial homeostasis in dominant optic atrophy patient-derived iPSCs. *Mol Ther - Nucleic*

Acids **26**:432–443. doi:10.1016/j.omtn.2021.08.015

Spiegel R, Saada A, Flannery PJ, Burté F, Soiferman D, Khayat M, Eisner V, Vladovski E, Taylor RW,

Bindoff LA, Shaag A, Mandel H, Schuler-Furman O, Shalev SA, Elpeleg O, Yu-Wai-Man P. 2016.

664 Fatal infantile mitochondrial encephalomyopathy, hypertrophic cardiomyopathy and optic atrophy

665 associated with a homozygous OPA1 mutation. *J Med Genet* **53**:127–131. doi:10.1136/jmedgenet-

666 2015-103361

667 Sugie A, Möhl C, Hakeda-Suzuki S, Matsui H, Suzuki T, Tavosanis G. 2017. Analyzing synaptic

668 modulation of drosophila melanogaster photoreceptors after exposure to prolonged light. *J Vis Exp*

669 **2017**. doi:10.3791/55176

670 Tang S, Le PK, Tse S, Wallace DC, Huang T. 2009. Heterozygous mutation of Opa1 in Drosophila

671 shortens lifespan mediated through increased reactive oxygen species production. *PLoS One* **4**.

672 doi:10.1371/journal.pone.0004492

673 Toomes C, Marchbank NJ, Mackey DA, Craig JE, Newbury-Ecob RA, Bennett CP, Vize CJ, Desai SP,

674 Black GCM, Patel N, Teimory M, Markham AF, Inglehearn CF, Churchill AJ. 2001. Spectrum,

675 frequency and penetrance of OPA1 mutations in dominant optic atrophy. *Hum Mol Genet* **10**:1369–

676 1378. doi:10.1093/hmg/10.13.1369

677 Trevisan T, Pendin D, Montagna A, Bova S, Ghelli AM, Daga A. 2018. Manipulation of Mitochondria

678 Dynamics Reveals Separate Roles for Form and Function in Mitochondria Distribution. *Cell Rep*

679 **23**:1742–1753. doi:10.1016/j.celrep.2018.04.017

680 Tsuyama T, Tsubouchi A, Usui T, Imamura H, Uemura T. 2017. Mitochondrial dysfunction induces

681 dendritic loss via eIF2α phosphorylation. *J Cell Biol* **216**:815–834. doi:10.1083/jcb.201604065

Vagnoni A, Bullock SL. 2016. A simple method for imaging axonal transport in aging neurons using the

adult Drosophila wing. *Nat Protoc* **11**:1711–1723. doi:10.1038/nprot.2016.112

Verny C, Loiseau D, Scherer C, Lejeune P, Chevrollier A, Gueguen N, Guillet V, Dubas F, Reynier P,

Amati-Bonneau P, Bonneau D. 2008. Multiple sclerosis-like disorder in OPA1-related autosomal

dominant optic atrophy. *Neurology* **70**:1152–1153. doi:10.1212/01.wnl.0000289194.89359.a1

Wang JT, Medress ZA, Barres BA. 2012. Axon degeneration: Molecular mechanisms of a self-

destruction pathway. *J Cell Biol*. doi:10.1083/jcb.201108111

Whitworth AJ, Lee JR, Ho VMW, Flick R, Chowdhury R, McQuibban GA. 2008. Rhomboid-7 and

HtrA2/Omi act in a common pathway with the Parkinson's disease factors Pink1 and Parkin. *DMM*

Dis Model Mech **1**:168–174. doi:10.1242/dmm.000109

Yang L, Tang H, Lin X, Wu Y, Zeng S, Pan Y, Li Y, Xiang G, Lin YF, Zhuang SM, Song Z, Jiang Y,

Liu X. 2020. OPA1-Exon4b Binds to mtDNA D-Loop for Transcriptional and Metabolic

Modulation, Independent of Mitochondrial Fusion. *Front Cell Dev Biol* **8**.

doi:10.3389/fcell.2020.00180

Yarosh W, Monserrate J, Tong JJ, Tse S, Le PK, Nguyen K, Brachmann CB, Wallace DC, Huang T.

2008. The molecular mechanisms of OPA1-mediated optic atrophy in Drosophila model and

prospects for antioxidant treatment. *PLoS Genet* **4**:0062–0071. doi:10.1371/journal.pgen.0040006

Yu-Wai-Man P, Griffiths PG, Gorman GS, Lourenco CM, Wright AF, Auer-Grumbach M, Toscano A,

Musumeci O, Valentino ML, Caporali L, Lamperti C, Tallaksen CM, Duffey P, Miller J, Whittaker

RG, Baker MR, Jackson MJ, Clarke MP, Dhillon B, Czermin B, Stewart JD, Hudson G, Reynier P,

Bonneau D, Marques W, Lenaers G, McFarland R, Taylor RW, Turnbull DM, Votruba M, Zeviani

M, Carelli V, Bindoff LA, Horvath R, Amati-Bonneau P, Chinnery PF. 2010a. Multi-system

neurological disease is common in patients with OPA1 mutations. *Brain* **133**:771–786.

doi:10.1093/brain/awq007

Yu-Wai-Man P, Spyropoulos A, Duncan HJ, Guadagno J V., Chinnery PF. 2016. A multiple sclerosis-

like disorder in patients with OPA1 mutations. *Ann Clin Transl Neurol* **3**:723–729.

doi:10.1002/acn3.323

Yu-Wai-Man P, Stewart JD, Hudson G, Andrews RM, Griffiths PG, Birch MK, Chinnery PF. 2010b.

OPA1 increases the risk of normal but not high tension glaucoma. *J Med Genet* **47**:120–125.

doi:10.1136/jmg.2009.067512

Zaninello M, Palikaras K, Naon D, Iwata K, Herkenne S, Quintana-Cabrera R, Semenzato M, Grespi F,

Ross-Cisneros FN, Carelli V, Sadun AA, Tavernarakis N, Scorrano L. 2020. Inhibition of

autophagy curtails visual loss in a model of autosomal dominant optic atrophy. *Nat Commun* **11**.

doi:10.1038/s41467-020-17821-1

Zanna C, Ghelli A, Porcelli AM, Karbowski M, Youle RJ, Schimpf S, Wissinger B, Pinti M, Cossarizza

A, Vidoni S, Valentino ML, Rugolo M, Carelli V. 2008. OPA1 mutations associated with dominant

optic atrophy impair oxidative phosphorylation and mitochondrial fusion. *Brain* **131**:352–367.

doi:10.1093/brain/awm335

 Figures with the corresponding legend

Figure 1: Effect of *dOPA1* knockdown on mitochondrial density and size in retinal axons.

(A) Schematic illustration of the *Drosophila* visual system from a dorsal view. (B) Visualization of

mitochondria and dOPA1 in a set of R7/8 axons. Mitochondria were visualized using the expression

of *mCherry-mito* driven by *GMR-Gal4* (magenta) while HA-tagged dOPA1 was immunostained using

the anti-HA antibody (green). Scale bar: 3 μm. (C, D) Representations of mitochondria visualized

using the expression of *Mito::GFP* driven by *GMR-Gal4* (green), one day after eclosion. Retinal axons

were labeled using the anti-Chaoptin antibody (magenta). Scale bar: 30 μm. (E) Quantification of

Mito::GFP intensity. Mito::GFP levels were calculated by dividing the Mito::GFP fluorescence

intensity by the anti-Chaoptin signal intensity. Control (n = 28 optic lobes) and *dOPA1* RNAi (n = 28

optic lobes). The data are presented as mean ± SEM. (F, G) Electron micrographs of cross sections of

the R1–6 retinal axons in the lamina of the Control (F) and d*OPA1* knockdown (G) on the day of

eclosion. Yellow circles and white dotted lines indicate mitochondria and a lamina column,

respectively. Scale bar: 2 μm. (H) Quantification of the mitochondrial area. Control (n = 96

mitochondria) and *dOPA1* RNAi (n = 92 mitochondria). Data are expressed as mean ± SEM. (I)

Representative EM images showing mitochondria with a densely packed matrix structure (classified

as class A) and collapsed mitochondria (classified as class B). Scale bar: 500 nm. (H) Quantification

of the mitochondria classified into class A and class B. Control (n = 96 mitochondria) and *dOPA1*

RNAi (n = 92 mitochondria). Data are analyzed by Chi-square test.

Figure 2. elevated reactive oxygen species levels in the *dOPA1* knockdown.

(A, B) Mitochondrial reactive oxygen species (ROS) levels in a retinal axon visualized using mito-

GFP (green) driven by *GMR-Gal4* and MitoSOX (magenta). Representative images of a retinal axon

in the Control (A, A') and *dOPA1* knockdown (B, B'). (C) Quantification of ROS levels for each

genotype. ROS levels were determined by dividing the fluorescence intensity of MitoSOX by that of

mito-GFP. Control (n = 20 optic lobes) and *dOPA1* RNAi (n = 18 optic lobes). Scale bar: 3 μm. Data

are presented as mean ± SEM.

Figure 3. Effect of *dOPA1* knockdown on photoreceptor neurodegeneration in *Drosophila*.

Schematic of the visual system in *Drosophila*. A set of R7/8 photoreceptors project their axons to the

second optic ganglion medulla. (B–E) All R7 and R8 axon terminals project to the medulla. *dOPA1* in

the photoreceptor was knocked down by *GMR-Gal4*. *40D-UAS* was used as a Control to match the

number of *UAS* sequences recognized by Gal4. One day after eclosion in Control (B) and *dOPA1*

knockdown (C), and one week after in Control (D) and *dOPA1* knockdown €. The retinal axons were

stained with anti-Chaoptin, a photoreceptor-specific antibody. Scale bar = 50 μm. (F) Number of axons

at each time point in R7 neurons for each situation quantified by MeDUsA. At 1 day of Control (n =

34 optic lobes) and *dOPA1* RNAi (n = 23 optic lobes), and at 1 week of Control (n = 28 optic lobes)

and *dOPA1* RNAi (n = 26 optic lobes). Data are expressed as mean ± SEM. (G–J) Electron

micrographs of cross sections of the photoreceptor cell bodies in the ommatidia of the retina.

Rhabdomeres are shown as yellow arrowheads. The day of eclosion in Control **(G)** and *dOPA1*

knockdown (H), and 1-week-old adults of Control (I) and *dOPA1* knockdown (J). Scale bar = 2 μm.

(K) Quantification of the number of rhabdomeres for each genotype and time point. At 0 day in Control

(n = 50 ommatidia) and *dOPA1* RNAi (n = 127 ommatidia), and at 1 week in Control (n = 50

ommatidia) and *dOPA1* RNAi (n = 117 ommatidia). Data are expressed as mean ± SEM.

Figure 4. Role of *dOPA1* in the retinal axon according to mutant analysis.

(A–C) Representations of retinal axons labeled with an anti-Chaoptin antibody (gray), one day after

eclosion. The images show Control (A), *dOPA1ˢ³⁴⁷⁵* somatic mosaic flies (B), and *dOPA1ˢ³⁴⁷⁵* somatic

mosaic flies expressing the eye-specific full-length *dOPA1* (C). Scale bar: 50 μm. (D) Quantification

of the number of axons for each genotype. Control (n = 41 optic lobes), mutant (n = 45 optic lobes),

and rescue (n = 41 optic lobes). The data are presented as mean ± SEM.

Figure 5. Verification of the pathological significance of disease mutations in *hOPA1*.

(A) Schematic illustration of the *hOPA1* gene construct with HA and myc tags in the UAS-based vector.

hOPA1 includes a mitochondrial import sequence (MIS) cleaved by mitochondrial processing

peptidase (MPP), a transmembrane region (TM), a coiled-coil region (CC1), a GTPase domain, a

Middle domain, and a GTPase effector domain (GED) containing a coiled-coil region (CC2). The sites

of variants (I382M, D438V, R445H, and a deletion from 2708 to 2711) are shown in red. S1 is the site

cleaved by OMA1. (B) Western blot analysis to confirm the expression of hOPA1 variants.

hOPA1_WT, hOPA1_D438V, hOPA1_I382M, hOPA1_R445H, and hOPA1_2708del were expressed

802   in whole *Drosophila* bodies and detected using anti-HA and anti-myc antibodies. α-Tubulin was used

803   as a loading Control. (C) Rescue experiments were conducted to assess the expression of each *hOPA1*

804   variant, including *dOPA1*, in the retina axons of *dOPA1* mutant somatic clones. The sample size is

805   indicated (n). Data are presented as mean ± SEM.

Figure 6. LOF or DN effects of disease mutations in *hOPA1*.

(A) Impact of each human OPA1 variant on the axon number in the optic nerve of *Drosophila* as

quantified using MeDUsA. (B) Expression of both *hOPA1* WT and its respective variants was analyzed

in photoreceptors lacking *dOPA1*, and the number of axons was quantified using the MeDUsA. The

sample size is indicated (n). Data are expressed as mean ± SEM. (C) Schematic representation of

814 interspecies differences in OPA1 interactions and the interchangeability of OPA1 between human and

815 fly.