A primer-independent DNA polymerase-based method for competent whole-genome amplification of intermediate to high GC sequences

Carlos D. Ordóñez[1], Carmen Mayoral-Campos[2], Conceição Egas[3,4] and Modesto Redrejo-Rodríguez[2*]

[1]Centro de Biología Molecular Severo Ochoa, CSIC-UAM, Madrid, Spain
[2]Departamento de Bioquímica, Universidad Autónoma de Madrid (UAM) and Instituto de Investigaciones Biomédicas Alberto Sols (CSIC-UAM), Madrid, Spain
[3]Center for Neuroscience and Cell Biology, University of Coimbra, Coimbra, Portugal
[4]Biocant, Transfer Technology Association, Cantanhede, Portugal

Present address: Carlos D. Ordóñez, CIC bioGUNE, Bizkaia Science and Technology Park, building 800, Derio (Bizkaia), Spain

*Correspondence to Modesto Redrejo Rodríguez
Departamento de Bioquímica
School of Medicine (Universidad Autónoma de Madrid)
Arzobispo Morcillo, 4
28029 – Madrid (SPAIN)
Telephone: +34 91 497 5963
E-mail: modesto.redrejo@uam.es

36 **Graphical abstract**
37

38
39
Schematic representation of methods based on multiple displacement
amplification (MDA) for whole genome amplification. The diagrams above
represent protocols initiated by random primers (RP-MDA) or a DNA primase-
generated short DNA primers (PrimPol-MDA) and continued by Φ29DNAP,
whereas the schematics below show piPolB-mediated MDA (left) and the
piMDA protocol (right), in which piPolB synthesizes DNA strands that are further
extended by Φ29DNAP.
47

48
49 **Abstract**
50
51
Multiple displacement amplification (MDA) has proven to be a useful technique
for obtaining large amounts of DNA from tiny samples in genomics and
metagenomics. However, MDA has limitations, such as amplification artifacts
and biases that can interfere with subsequent quantitative analysis. To
overcome these challenges, alternative methods and engineered DNA
polymerase variants have been developed. Here, we present new MDA
protocols based on the primer-independent DNA polymerase (piPolB), a
replicative-like DNA polymerase endowed with DNA priming and proofreading
capacities. These new methods were tested on a genomes mixture containing
diverse sequences with high-GC content, followed by deep sequencing.
Protocols relying on piPolB as a single enzyme cannot achieve competent
amplification due to its limited processivity and the presence of *ab initio* DNA
synthesis. However, an alternative method called piMDA, which combines
piPolB with Φ29 DNA polymerases, allows proficient and faithful amplification of
the genomes. In addition, the prior denaturation step commonly performed in
MDA protocols is dispensable, resulting in a more straightforward protocol. In
summary, piMDA outperforms commercial methods in the amplification of
metagenomes containing high GC sequences and exhibits similar profiling,
error rate, and variant determination as the non-amplified samples.
71
72
73
74 **Keywords**
75
76 WGA, MDA, primer-independent DNA polymerase, piPolB, metagenomics,
77 genomics
78

Introduction

Genomics and metagenomics have undergone a revolution in the last two decades. This is due to advances in whole-genome amplification methods and the development of massively parallel sequencing methods that have opened up previously unimagined applications in biotechnology and biomedicine (1). This has led to an increasing demand for more efficient, sensitive, and unbiased protocols aimed at developing high-throughput analysis of biodiversity and personalized medicine, among other applications.

Current technologies allow direct sequencing of DNA samples without a prior amplification step, which can reduce bias and increase data reliability. However, this is not possible when the quantity or quality of DNA is limited. Therefore, sequencing of the original raw sample is still experimental or limited to targeted testing in some scenarios (2–4). In addition, sample origin and characteristics, genome isolation procedures, and sequencing platforms can also be sources of error, bias, and underrepresentation of genomic sequences that are present in lower quantities (5–8).

To ensure sample availability for DNA sequencing, whole (meta)genome amplification (WGA), which includes PCR-based approaches, multiple displacement amplification (MDA), and a number of variations and derived methods (9–11), enables efficient amplification of minute nucleic acid samples. MDA provides faithful and reliable amplification of (meta)genomic DNA without the need for prior knowledge of the target sequence. It has been used for the amplification of complex (meta)genomic samples as well as for whole genome amplification at the single-cell level (scWGA) for almost 20 years (12–14). In addition, it is based on an isothermal protocol that simplifies its use at the point of care and promotes the development of various applications beyond genomic analysis (15–17). Most MDA protocols are based on the use of hexamer random primers (RP) and the highly processive DNA polymerase from *Bacillus virus Φ29* (Φ29DNAP) (15, 18, 19), which can generate very long amplicons, that allow high coverage and detection of single nucleotide polymorphisms (SNP) (20, 21). However, MDA also brings some disadvantages, such as the generation of primer-related artifacts, chimeric DNA sequences, or biased poorer amplification of sequences with extreme GC content (22–27), which may affect uniform genome coverage and sensitivity for the detection of minor alleles or underrepresented sequences (23, 28, 29). In addition, the extreme processivity of Φ29DNAP leads to the overamplification of circular DNA molecules, which compromises its use for some metagenomic applications, such as metaviromic analysis (30, 31). Accordingly, several new tools and methods have been developed to increase the accuracy of WGA. These include both engineered versions of DNA polymerases with improved MDA performance, thermostability, or tolerance to chemicals (32–34), and alternatives that combine both PCR and MDA, such as MALBAC or PicoFLEX/GenomeFlex. The latter methods have been reported to have limited coverage compared with Φ29DNAP-based MDA but offer higher single nucleotide variants (SNV) detection rates, with MALBAC having higher uniformity and a lower allelic dropout (ADO) rate in the portion of the genome covered (29, 35). However, the specialized polymerase required for this reaction lacks proofreading capacity, resulting in increased error rates (36). Another promising modification of MDA-based WGA

is TruePrime, which is based on the combined use of a *Thermus thermophilus*
primase-polymerase enzyme (TthPrimPol) to generate short DNA primers that
are subsequently extended by Φ29DNAP (37). This protocol is expected to
reduce primer-related artifacts and provide a better representation of
sequences with high-GC content (38).
In this work, we have developed and assessed new MDA methods based on
recombinant *Escherichia coli* primer-independent PolB (piPolB), previously
characterized as having faithful DNA polymerase activity, along with
proofreading, translesion synthesis, and DNA primase capacities (39). Our
results have shown that the use of piPolB in the absence of DNA primers or
accessory factors can lead to successful DNA amplification, although the
amplified DNA product also contains a high proportion of duplicated and low
complexity sequences, corresponding to spurious *ab initio* DNA synthesis (40).
In contrast, the combination of piPolB with a highly processive enzyme such as
Φ29DNAP, a protocol hereafter referred to as primer-independent MDA
(piMDA), results in the amplification of genomic samples at a level similar to
commercially available kits, which is further increased when a prior alkaline
denaturation step is added (piMDA+D protocol).
Deep sequencing of MDA products from a mixture of bacterial genomes
containing high-GC sequences obtained with piMDA and piMDA+D protocols
shows that both alternatives achieve competent amplification and an improved
assembly compared to commercial random primers and PrimPol-based MDA.
Overall, our results suggest that the piPolB polymerase, primase, and
proofreading capacities result in high-fidelity DNA molecules that can
subsequently be extended by the highly processive Φ29DNAP for competent
and accurate amplification of DNA samples. This work paves the way for the
use of novel tailored MDA methods for environmental and biomedical
applications.
157

MATERIALS AND METHODS

DNA amplification substrates

The single- and double-stranded M13 DNA was obtained from New England Biolabs (NEB). The genomes mixture was generated from the combination of four different bacterial genomes. *Escherichia coli JM83, Micrococcus luteus* (later corrected to *Kocuria rhizophila*) and *Pseudomonas aeruginosa* strains were taken from the teaching collection of the Molecular Biology Department at Universidad Autónoma de Madrid. *Bacillus subtilis 110NA* (41) was taken from Margarita Salas laboratory stock. Genomic DNAs were purified from 3 mL overnight cultures grew at optimum temperature (30 ºC or 37 ºC) by Dneasy Blood & Tissue Kits (QIAGEN). Additionally, 1 mg/mL metapolyzyme mix from a stock diluted in PBS pH 7.5 was included in the lysis step to enhance digestion. The mock metagenome sample was then prepared by mixing an equal mass of purified genomic DNAs.

Genomic reference sequences were selected from the top hits of BLASTN queries on NCBI Nucleotide Database, using a subset of the largest contigs from the control assemblies (Table S1).

Multiple displacement amplification assays

The DNA polymerase from bacteriophage Φ29 from Margarita Salas laboratory was purified as described (42). The piPolB was purified as untagged recombinant protein as reported previously (39).

Unless otherwise stated, piPolB isothermal MDA reactions were performed employing 250 nM piPolB in the presence of 20 ng of either M13 ssDNA (m13mp18, NEB), dsDNA (m13mp18 RF I, NEB) or genomes mixture DNA. The piPolB MDA reactions contained 50 mM Tris-HCl pH 7.5, 1 mM DTT, 4% (v/v) glycerol, 0.1 mg/mL BSA, 0.05% (v/v) Tween 20, 20 mM $MgCl_2$, 500 μM dNTPs, 10 mM ammonium sulfate, in a final volume of 10 μL for 30 ºC for 16 h. Characterization of piPolB amplification product was performed by digestion of the indicated volume of amplification product with diverse nucleases. Samples treated with 0.2 U/μL Micrococcal Nuclease (Worthington) were incubated at 37 ºC for 2 h in 50 mM Tris-HCl pH 7.5, 1 mM DTT, 4% (v/v) glycerol, 0.1 mg/mL BSA, 0.05% (v/v) Tween 20, 25 mM $CaCl_2$. 1 U/μL or the indicated concentration of EcoRI-HF, HhaI, or MnlI was at 37 ºC for 2 h (EcoRI-HF) or 3 h (HhaI or MnlI) in rCutSmartTM Buffer (NEB). Nuclease S1 (Boehringer Mannheim) was tempered at 37 ºC for 3 min before the incubation with the sample at 37 ºC for 1 min in Reaction Buffer for S1 nuclease (Fermentas). After the digestion, enzymes were inactivated according to manufacturer's instructions.

The so-called primer-independent DNA amplification protocol (piMDA) (see below) reactions were performed in 35 mM Tris-HCl pH 7.5, 1 mM DTT, 0.1 mg/mL BSA, 10 mM ammonium sulfate, 50 nM piPolB, 500 nM Φ29DNAP, 20 mM $MgCl_2$, and 500 μM dNTPs. In the piPolB+D and piMDA+D protocols, DNA was subjected to a previous denaturation step with 75 mM NaOH in a volume of 5 μL for 3 min, and subsequently neutralized adding 5 μL of 250 mM Tris-HCl pH 7.5. Amplification reactions were carried out at 30 ºC for 3 h unless other conditions are specified. Reactions were terminated by heating samples at 65 ºC for 10 min. MDA controls were performed employing a random-primers-based MDA (RP-MDA) kit as random-primer-based method: Repli-g Mini Kit

from QIAGEN; and a TruePrime-based-MDA (PrimPol-MDA) kit: TruePrime
Whole Genome Amplification Kit from 4BaseBio. Manufacturer's specifications
were followed in commercial protocols. Furthermore, a final heating step at 65
ºC for 10 min was always conducted for protein inactivation in MDA protocols.
Electrophoresis analyses of amplification product were carried out by digestion
of 2 µL (for amplification of circular DNA) or 5 µL (for samples from
metagenomic DNA) of each reaction with 1 U/µL EcoRI or EcoRI-HF at 37 ºC in
a final volume of 10 µL prior to electrophoresis in 0.7% agarose 1X-TAE and
visualized with ethidium bromide (Sigma-Aldrich) or GreenSafe Premium
(NZYTech).
Absolute amplification measure of DNA input sample and amplification products
was performed by fluorescence quantitation with the AccuBlue High Sensitivity
dsDNA Quantitation Kit (Biotum). Black opaque microplates with 96 wells
(Greiner) were used for accomplishing the fluorescence reactions which were
measured with FLUOstar Omega (BMG Labtech).
223
Amplified metagenomes sequencing and reads processing
Control genomic DNA and amplification products were ethanol-precipitated and
~1 mg of DNA was sheared into ~350-bp fragments using a Covaris M220
focused acoustic shearer (Covaris, Woburn, MA, USA) following the
manufacturer protocol. Sequencing libraries were prepared with the TruSeq
DNA PCR-Free Low Throughput Library Prep kit (Illumina) according to the
manufacturer's protocol. Samples were then sequenced in the Illumina NextSeq
550 with the 2x75bp NextSeq V2.5 High Output kit, at Genoinseq - Next
Generation Sequencing Unit (Biocant, Cantanhede).
Adapters and low-quality reads were filtered with Trimmomatic v.0.35
(parameters: LEADING:3 TRAILING:3 SLIDINGWINDOW:4:20 MINLEN:35).
Quality check (QC) before and after trimming was performed with Seqkit v2.3
(43) and FastQC v0.11.9
(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/), and individual
reports subsequently aggregated with MultiQC 1.11 (44) for comparison.
Trimmomatic filtered reads were used for *de novo* profiling using Metaphlan
v.4.0.0 (45) using the --unclassified_estimation option. Additionally, trimmed
reads were mapped against the reference genomes with Bowtie2 v2.3.5.1 (46)
and further analyzed for coverage depth, breadth, and aligning mismatches with
Weesam v1.6 (https://bioinformatics.cvr.ac.uk/weesam-version-1-5/) and Alfred
v0.1.16 (https://www.gear-genomics.com/alfred).
Fastp (47) was additionally used to obtain a list of overrepresented sequences
in each dataset or raw reads. Identification of low-complexity sequences in
Metaphlan unclassified reads was performed with BBduk
(https://jgi.doe.gov/data-and-tools/software-tools/bbtools/). GC bias was
assessed using Picard tool CollectGcBiasMetrics
(https://broadinstitute.github.io/picard/).
Trimmomatic filtered reads from each sample were assembled with
metaSPAdes (48). Genomic assemblies were analyzed as minimal
metagenomes with Metaquast (49) and also annotated with Bakta (50).
Comparative assessment of k-mers diversity present in the reads and in the
corresponding assembly was carried out with KAT (51) using a K-mer size of
16, according to the size of the amplified metagenome, as suggested previously
(52). Single-nucleotide variants (SNV) detection with Snippy 4.6.0

(https://github.com/tseemann/snippy). Finally, MetaCoAG (53) was used for *de*
novo metagenomes binning (coverage >5% and p-value <10^{-10}) and bins were
assigned to reference genomes using Mash distances (54).
261
262
Data analysis
Data were analyzed using R and RStudio (55) and plots were generated with
ggplot2 package (56). The final figures also contain minor modifications made
with Illustrator (Adobe). As indicated, analysis of inter-groups significant
differences was carried out by pair-wise t-test or Wilcoxon rank tests, when
normality and homogeneous variance test prescribed the use of non-parametric
analyses.
The correlation between normalized coverage and frequency of sequence
windows at each GC content from Picard output was analyzed using the
Corrplot package (57). Outliers that would impair these analyses were removed
as follows. For metagenome correlation analyses GC values below 20% and
over 80% were removed. In the case of correlations for individual reference
genomes (Figure S6), GC contents with a sequence windows frequency below
10 were discarded.
277
278
279   **Data availability**
280   Raw sequencing data has been deposited in the e-cienciaDatos repository with
281   DOI: 10.21950/HCNDGF
282   An annotated markdown report of the analysis containing the R code and plots
283   is available in a Github repository (https://mredrejo.github.io/pimda/).
284

RESULTS
Primer-independent isothermal MDA by recombinant piPolB
The DNA priming capacity of piPolB together with a 3'-5' exonuclease proofreading activity and strand displacement capacity (39) make it a promising candidate for use in DNA synthesis by isothermal multiple displacement amplification (MDA). As shown in Figure 1A, piPolB can amplify both single- and double-stranded DNA starting from 1 and 10 ng of input nucleic acid, respectively. Similar to other MDA protocols, piPolB MDA is affected by high ionic strength and requires magnesium as a cofactor (Figure S1A).

Figure 1. Isothermal multiple displacement amplification by piPolB (piPolB-MDA).
(A) Agarose electrophoresis of EcoRI-digested product from single-stranded and double-stranded DNA amplification by piPolB. (B) Digestion analysis of amplified DNA product. Amplification reactions were performed with 1 ng input ssDNA and subsequently digested with increasing concentrations (0.07, 0.7 and 7 units) of EcoRI (single-site) and two different promiscuous restriction enzymes, HhaI (26 sites) and MnlI (62 sites). As indicated 200 ng of M13 dsDNA was also digested as a control (lanes 13-15). Lane 12 corresponds to undigested M13 dsDNA, thus containing supercoiled and relax circular forms. Migration of linear M13 is indicated with a black arrow (7.25 Kbp). (C) Schematic representation of piPolB MDA DNA product as an hyperbranched DNA structure

because of multiple initiation events. Thus, single-site enzymatic digestion gives rise to a mixture of small fragments, unit-length molecules and branched molecules that move slower than unit length.

As expected, given the limited processivity of piPolB (1-6 kB, see ref. (39)), the amplified DNA product migrated as a smear in agarose electrophoresis, even after digestion with a single-site restriction enzyme that would generate unit length only if long concatemers were produced (see below). Digestion of DNA product synthetized by piPolB with different nucleases (Figure 1B and Figure S1C) indicates that piPolB synthesizes dsDNA by means of repeated initiation events that give rise to hyperbranched dsDNA structures in which fragments longer than unit length (in this case 7.25 kb) would be a minority (Figure 1C).
To obtain larger DNA products we then combined piPolB with Φ29DNAP (Figure 2). As expected, Φ29DNAP can extend DNA fragments generated by piPolB, resulting in long amplicons that in the case of plasmid substrates are made up of concatemers that can be digested into unit-length fragments. This

new protocol also allows the amplification of smaller amounts of DNA, especially for dsDNA (Figure 2B). Amplification of dsDNA (plasmid or metagenome) is impaired with increasing piPolB concentration, likely due to competition between the two DNA polymerases for the free 3'OH ends, which would be more scarce for dsDNA than for ssDNA. However, the amplification yield correlates with Φ29DNAP concentration, suggesting that it is responsible for the main amplification product. This combined protocol of piPolB+Φ29DNAP in a single step will be referred to hereafter as piMDA or piMDA+D, when a denaturation step was performed previously.

Figure 2. Proficient isothermal DNA amplification by piPolB coupled to Φ29DNAP (piMDA) generates long amplicons.
(A) EcoRI-digested DNA products of piMDA product on different DNA substrates reveals the formation of long dsDNA amplicons, larger than the M13 genome unit, increasing the sensitivity for amplification of ssDNA (lanes 1-9) and, especially, dsDNA substrate (lanes 10-17). The ratio of piPolB: Φ29DNAP affects the DNA yield on DNA amplification of plasmid (M13, black arrow). (B) and a mock metagenomic DNA (mgDNA, C).

To evaluate the amplification efficiency of the piPolB MDA and piMDA methods in detail, we prepared a genomic DNA sample (see Methods) consisting of four genomes from Gram-positive and Gram-negative bacteria, of different sizes (2.7-6.5 MB) and moderate to high (45-70%) GC content. The final sample (Table S1) is approximately 18 MB long and has an average GC content of 58%. This sample was used for comparative MDA assays with piPolB-based protocols and two commercially available kits based on Φ29DNAP, namely RepliG (Qiagen) for Random-Primers MDA (RP-MDA) and TruePrime

(4BaseBio) for a primase-based MDA (PrimPol-MDA). The results (Figure 3 and
Figure S2) show that the piPolB MDA protocol is comparable to the available
MDA alternatives in terms of DNA yield, but the yield of the piMDA method,
especially the version with a DNA denaturation step (piMDA+D), outperforms
the classical, random-primers based MDA (RP-MDA) in amplifying of our
genomic sample. Under these conditions, amplification using the RP-MDA
method was significantly less productive than the other methods, likely due to
the abundance of high-GC sequences in the analyzed sample.
403

404
Figure 3. Time-course assays of different DNA amplification protocols.
Quantification of piPolB-based MDA protocols and commercially available random-primers MDA
and PrimPol-MDA DNA products. Amplification assays with commercial kits were performed
following manufacturer's instructions. When indicated (+D), a prior alkaline denaturation step
was performed for piPolB and piMDA protocols. Four independent MDA experiments with 0.4
ng/µl genomic DNA input were quantified after the indicated amplification time at 30 ºC.
Significant p-values from unpaired Wilcoxon rank sum tests for pairwise comparisons of DNA
product/hour are indicated above the graph.
413
High sequencing depth underscores the need for a highly processive DNA
polymerase in MDA
Samples from MDA experiments of the genomes mixture were sequenced using
a PCR-free library preparation and short reads paired-end strategy to minimize
the sequencing bias (58). The amplification products of piMDA, piMDA+D, and
RP-MDA and the nonamplified control samples (NA) were analyzed in
duplicates from independent amplification experiments (Table S2). Deep
sequencing yielded an average total length of more than 280X of the input
metagenome and provided high sequence coverage for all samples.
Adaptor trimming and quality filtering revealed the first differences between
samples, with reads from piPolB and piPolB+D MDA products having lower

average quality scores (Table S2) and a higher proportion of reads discarded at
this stage (Table S3). This suggests that library generation was defective in
these samples, likely due to the hyperbranched structure of the amplified DNA
product generated by piPolB. This could hinder the generation of a
homogeneous DNA shearing and adapter ligation, making detailed evaluation
of the piPolB and piPolB+D MDA methods difficult.
As expected for DNA sequences obtained by MDA, the quality assessment also
showed that all samples contained more over-duplicated reads (duplication
level > 3-4) than the non-amplified control samples (Figure 4A). Importantly, the
proportion of duplicated reads was lower for the piMDA and piMDA+D samples
than for RP-MDA and PrimPol-MDA up to a duplication level ≤10, suggesting
that the latter protocols are more prone to overduplication of some sequences.
However, the piPolB and piPolB+D samples contained a higher proportion of
highly duplicated sequences. This pattern would be consistent with repetitive
initiation events followed by distributive strand elongation, resulting in a biased
amplification product.
441
Sequence coverage bias is highly dependent on GC-content
The GC content of the input genomic mixture is 57.70% (Table S1), which is
very similar to the value obtained in the non-amplified samples, rendering
curves with two peaks, one around 45% GC, which would correspond to the
genomes of *E. coli* and *B. subtilis* and the second up to ~68% GC,
corresponding to the GC content of *P. aeruginosa* and *K. rhizophila* (Figure 4B).
However, the average GC content in RP-MDA (~44%), PrimPol-MDA (~46%),
piPolB+D MDA(~46%), and piPolB MDA (~42%) is reduced because the peak
of high-GC content is absent from the curves, indicating a strong negative GC
bias. On the other hand, the reads from the piMDA and piMDA+D products
show a positive GC bias as their average GC content ranges from 60 to 65%
and shows curves with a main peak at 70% GC and a shoulder around 50%
GC.
We then performed the profiling of the raw data of each sample against a
recently curated microbial database using Metaphlan (45). As shown in Figure
4C, reads from the nonamplified control samples show an overall even
distribution among the four species that make up the mixture, with small
differences between duplicates resulting in a slight underrepresentation of the
larger genome of *P. aeruginosa* and an overrepresentation of the moderate-GC
sequences of the *B. subtilis* genome, consistent with the expected bias and
variability resulting from library preparation and short reads sequencing (58,
59). However, the random-primers- and PrimPol-based MDA promoted a very
strong underrepresentation of high-GC genomes (*K. rhizophila* and *P.*
aeruginosa) and, in turn, an overrepresentation of *E. coli* and especially *Bacillus*
sequences. In contrast, the DNA samples generated with the piMDA and
piMDA+D protocols contain some overrepresentation of genomes with high-GC
content compared to the NA samples, although they are distributed across the
four species and show the highest relatedness to the non-amplified samples
(Figure S3). Strikingly, the samples from piPolB and piPolB+D MDA contain a
high proportion of unknown DNA sequences. The proportion of unknown DNA
sequences decreases with the addition of the previous DNA denaturation step
in piPolB+D MDA, suggesting that amplification specificity increases when
access to the DNA priming substrate is favored by double helix denaturation.

Accordingly, a negligible number of unclassified reads in piMDA reads (around
1%) is absent in the samples obtained piMDA+D protocol. Importantly, the fact
that these unknown sequences have no similarities in the metaphlan database
rules out the presence of contaminated DNA in any of the samples analyzed.

479
Figure 4. Characterization of the reads from each MDA experiment.
Trimmomatic output files containing paired and unpaired reads from forward and reverse raw
reads were analyzed with FastQC and subsequently evaluated for metagenome profiling with
Metaphlan4. (A) The relative level of duplication found in each sequence. The piPolB and
piPolB+D overrepresented sequences were also extracted with Fastp (see Tables S9-S12). (B)
GC content proportion per sample. (C) Profiling of reads across the Metaphlan database at the
species or genus level. Unclassified reads are also indicated. For simplicity, samples in A and B
are colored by MDA method, although paired and unpaired reads for each of the eleven
samples sequenced are represented.

489
The poor quality control and profiling results of the sequencing data of piPolB
and piPolB+D DNA amplification suggest that piPolB may be able to perform *ab*
initio DNA synthesis, independent of primers and template. *Ab initio* DNA
synthesis has been described in various DNA polymerases since the 1960-
1970s, and its contribution *in vivo* and *in vitro* to the bulk of DNA synthesis is
controversial (40). The mechanism of generation of these undirected DNA
products is poorly understood, but the characterization of the *ab initio* activity of
Bst DNA polymerase indicated that it is strongly stimulated by DNA nickases
and it results in low-GC content DNA enriched in non-palindromic repetitive
sequences (40, 60). In the case of piPolB, prolonged incubations of MDA
assays in the absence of input DNA (Figure S4), confirmed the spurious
amplification. Extraction of the overrepresented in forward and reverse reads of
piPolB and piPolB+D amplified samples shows that they consist of similar
repetitive sequences (Tables S9-S12). Furthermore, examination of the
unmapped reads from the Metaphlan database shows that the addition of a
denaturation step not only increases mapping efficiency but also decreases the
proportion of low complexity reads, from 12.6% in piPolB MDA to 0.25% in
piPolB+D MDA, similar to the level in any of the other samples (Table S4).
These results, together with a lower GC content in piPolB than in piPolB+D raw
reads (Figure 4B) but not in the mapped reads (see below and Figure 6E),
points to *ab initio* DNA synthesis by piPolB that would be reduced when access
to single-stranded DNA is facilitated by the denaturation step.
In conclusion, the use of piPolB for MDA as a single enzyme is hampered by
the limited processivity of the enzyme as well as by spurious DNA synthesis,
whereas its combination with Φ29DNAP reduces or eliminates *ab initio* DNA
synthesis and allows proficient whole genome amplification with a sequence
profiling similar to that of the unamplified sample.
517
GC bias affects coverage depth, breadth, and amplification fidelity.
We then analyzed the mapping, coverage, and alignment mismatch statistics of
each sample with the independent reference genomes in our genomic mixture.
As a reference, Figure 5 shows the coverage plots for the *E. coli* (A) and *K.*
rhizophila (B) genomes. The control NA samples show the most consistent
coverage, although they contain spikes of overrepresented sequences that
range up to 10-fold the average depth. Mapping of reads from the RP-MDA and
PrimPol-MDA samples shows a jittered but overall uniform plot in the *E. coli*
genome, but coverage of *K. rhizophila* is scattered with some regions of great
depth coverage. In the case of the piPolB and piPolB+D samples, the plots
indicate an uneven coverage depth, with peaks of strongly overrepresented
sequences and a low average depth, resulting in a high standard deviation of
depth, especially in the piPolB reads (Figure 6C). As for the piMDA and
piMDA+D reads, the mapped reads show a high jitter but an overall similar
profile for all reference genomes with few large peaks of overrepresented
sequences but with high coverage of reference genomes with high-GC.
534

535
Figure 5. Mapping coverage of sequenced samples across the reference genomes of *E*
***coli* (A, 50% GC) and *K. rhizophila* (B, 70% GC).**
Average coverage depth, x0.2, and x1.8 ranks are indicated with a green, orange, and red
dotted line, respectively. Plots were generated using weeSAM (see Methods). Note that a
different Y-axis scale was calculated for each sample, see Figure 8 for a coverage plot with a
constant scale per sample. As indicated, mapping coverage is shown for one sample of each
protocol. A full report with all samples mapping over the four reference genomes is available at
the online Github repository (https://mredrejo.github.io/pimda/weeSam/weesam_full.html).
544

545
The proportion of mapped reads (Figure 6A) agrees with the profiling results
(Figure 4C). We also analyzed the Weesam coefficient of coverage variation,
which indicates the variability of coverage depth and breadth, with values > 1
(Figure 6C) considered indicative of non-uniform coverage. High-GC genomes
of *P. aeruginosa* and *K. rhizophila* are covered by RP-MDA, PrimPol-MDA,
piPolB-MDA, and piPolB+D-MDA with lower depth, with the reads from both
RP-MDA samples with a distinct higher coefficient of variation (Figure 6C). This
is consistent with low depth and breadth (Figure 5 and Figure 6B) of the high-
GC reference genomes and also with previous reports (61). This less proficient
amplification of high-GC sequences is also reflected in a higher mismatch rate
of the RP-MDA and PrimPol-MDA methods for these genomes (Figure 6D).
However, the samples generated by the piPolB and piPolB+D MDA protocols
include a lower amplification fidelity of *B. subtilis* reference sequences that also
exhibits higher depth standard deviation suggesting that overamplification may
be the reason for the lower replication fidelity. In contrast, the piMDA(+D)
samples show a more uniform average coverage depth of the four genomes,
although with a lower depth for the *E. coli* and *B. subtilis* reference genomes
and a similar mismatch rate for all reference genomes as the control samples.
Detailed analysis of GC bias in the genomes mixture (Figure 6E and Figure S5)
highlights the differences in high-GC content samples, with a significant
negative correlation of coverage per GC content not only in the RP-MDA and
PrimPol-MDA samples but also in the NA control samples, indicating a negative
effect of the sequencing method in our sample, likely at the library preparation
step (58, 62). Thus, the normalized coverage in the control NA samples
decreases from ~1.2 to ~0.8-0.9 at GC content above 50-55%. In line with
previous results, the piPolB(+D) amplification gives rise to scattered peaks of
highly overrepresented sequences encompassing 30-50% GC content.
Previous analysis of GC bias induced by PrimPol-MDA method shows some
discrepancies that might be due to the different nature of the amplified sample,
as it was reported to have a minor negative GC bias, similar to that of RP-MDA
in human DNA (37) but it outperformed the classic MDA protocol on metavirome
amplification (38). In our genomes, we found that PrimPol-MDA exhibits strong
negative GC bias, only slightly less pronounced than the RP-MDA samples.
Thus, both sets of samples show a statistically similar (Table S5) normalized
coverage of over 2.5 at low-GC content, dropping to 0.1-0.2 at a high-GC
content of >60%. Contrary to that and consistent with the profiling results, the
piMDA and piMDA+D assays promoted a significant bias toward high-GC
sequences, but with a more balanced normalized coverage, ranging from 0.4-
0.5 at 45% GC content (*B. subtilis*) to 1.4-1.7 at 70% GC (*K. rhizophila*). The
analysis of GC bias per reference genome suggests that bias is higher in
intermediate-GC genomes, and it also reveals certain differences between
duplicates (Figures S5 & S6). Nevertheless, pairwise comparisons of the
normalized coverage of all samples show that piMDA+D samples are not
statistically different from the non-amplified samples (Table S5).
All in all, the piMDA and piMDA+D methods provide reasonable coverage at
any GC value, with higher coverage for sequences with high-GC content.
592

Figure 6. Coverage of reference genomes mixture by different MDA protocols.
Mapping reads rate (A), coverage breadth (B), and average depth ± standard deviation (C) were
analyzed using Weesam v1.6. The small points in panel C represent the coverage variation
coefficient per reference genome (right axis). The dashed horizontal line marks the value 1 for a
homogeneous coefficient of variation. The overall mismatch rate (D) was obtained using Alfred
v.0.1.16. (E) Normalized sequence coverage by sample reads at a given GC content was
determined using Picard v2.25.0. Percentages of GC in the reference sequence are determined
from sequence bins of 100 bases, whose abundance is shown as steel blue bars (right axis).
The horizontal line represents the reference value of homogeneous coverage at all GC content
values. The average GC content of each sample and the correlation of normalized coverages to
sequence abundance at each GC content (95% confidence interval) are also shown in the right
panel. Nonsignificant R coefficients (p>0.05) are shown in grey.

Extraction of high-quality genomes from high-GC bacterial genomes amplified with piMDA

We then attempted to assemble the reads to compare the assembled sequences and considered them as a minimal metagenome and evaluate the extraction of individual genomes. We could obtain *de novo* assemblies with Metaspades, except for reads from the piPolB MDA sample, which could not be assembled due to the abundance of low-complexity sequences. The assembly statistics (Figure 7 and Table S6) and circular visualization (Figure 8) show that the obtained assemblies appear to be different from the reference genome for all samples of the *P. aeruginosa* genome. On the other hand, the obtained *B. subtilis* 110NA assemblies in all samples have a gap corresponding to the expected deletion across the Spo0A gene (41). In contrast, the assemblies of *E. coli* and *K. rhizophila* match the reference genomes very well in most samples.

Figure 7. Assembly of amplified sequences.
Metaspades (v. 3.13.1) *de novo-generated* assemblies were analyzed with MetaQuast v. 5.1.0rc1 using the indicated genomes as references (Table S1). (A) Number of contigs (large points) and the L50 (small points), i.e., the smallest number of contigs whose length sum makes up half of the genome size. (B) Mismatches (big points) and the number of insertions and deletions (small points) per 100 kbp. (C) Total number of misassemblies (big points) and local misassemblies (small points). (D) Number of contigs per reference genome. Non-aligned contigs are also shown. (E) Mismatches per genome and per 100 kbp are represented normalized to the control NA sample.

The number and length of contigs in the final assemblies were similar in NA, PrimPol-MDA, piMDA, and piMDA+D samples (Table S6 and Figure 7). The assembly of piPolB+D sample, and its reads were also assembled in a higher

number of contigs and the L50 was the largest obtained, which also indicates
the presence of several small contigs, mostly from *E. coli* and *B. subtilis*. The
RP-MDA assembly has a very high number of contigs (Figure 7A), suggesting
many small contigs with high-GC content (> 50%, see Figure S7) mapped to the
genomes of *P. aeruginosa* and *K. rhizophila*. Accordingly, this assembly also
had a higher number of misassemblies and mismatches, mainly related to the
K. rhizophila sequences. The assembly of the PrimPol-MDA sample also
contains a large number of small contigs with very high-GC content (<60%), but
the assembly show a better quality. K-mer comparison of reads and assemblies
indicated that, as expected, only the piPolB+D sample contain a high number of
k-mers excluded from the assembly and RP-MDA, RP-MDA2, PrimPol-MDA
and piPolB+D samples contain k-mers in the reads occurring >2 times in the
assembly, in agreement with a higher level of biased overamplification (Figure
S8). Finally, the annotation of all the sequences was very similar, except for
those from RP-MDA and piPolB+D amplified DNA, which span lower coding
density and a higher number of pseudogenes, most likely as a consequence of
poor assembly (Table S7).
For short variant calling, we used the Snippy caller, which was recently
benchmarked as the best variant caller for microbial genomes and showed both
strong and consistent performance across species (63). The capacity of variant
calling varied widely among samples (Table S8). For example, RP-MDA and
piPolB(+D) amplification result in poor variant detection of both single
nucleotide polymorphisms (SNPs) and complex variants. It should be noted that
with the exception of samples NA and NA2, which can call the same SNVs,
there are differences between the duplicates and the second batch of sample
sequencing in the case of RP-MDA, piMDA and especially piMDA+D allow
better variant detection, with piMDA+D2 being identical to the control samples.
Nevertheless, the PrimPol-MDA and piMDA(+D) methods show similar and
good performance in variant detection, with up to 60-80% of variants coinciding
with the non-amplified samples.
We then performed reference-free contig binning of the assemblies to test
whether amplification affected the extraction of the metagenome-assembled
genomes (Figure 9). We used MetaCoAG because it generates contig bins with
abundance, base composition, and connectivity information from assembly
graphs to produce high-quality bins (53), which we then assigned to the
corresponding reference genome. As expected from the high coverage breadth
and the statistics of the assemblies, the *E. coli* genome in all the samples can
be assigned to a single bin covering 95-98% of the reference genome, whereas
the genomes of *B. subtilis* and *P. aeruginosa* were more unevenly covered by
the obtained bins. Consistent with the above results, the contigs of the RP-MDA
assemblies result only in bins covering half of the genomes of *P. aeruginosa*
and *K. rhizophila*. Interestingly, all other assemblies yield a similar binning
profile, even the piPolB+D assembly, although supported by an overall lower
coverage depth and breadth.

Figure 8. Circos plot of assemblies mapped against the reference genomes.
As indicated on the NA control sample, the outer circle represents reference sequences with GC (%) heatmap [from 25% (white) to 79% (black)]. Yellow, orange, cyan, and purple color bars help to distinguish between the reference genomes of *B. subtilis, E. coli, P. aeruginosa, and K. rhizophila*, respectively. Green/red bars in the second circle represent correctly assembled and misassembled contigs, respectively, and black bars indicate mismatches. The inner circle contains a plot of the reads coverage depth, with a uniform scale for each sample in the different references.

Figure 9. Sample binning for each amplification protocol.
Bins were reference-free extracted with MetaCoAG (–mg_threshold 0.4 –bin_mg_threshold 0.2)
and subsequently assigned to a reference genome using the smallest Mash distance. The
proportion (%) of each bin spanned in its reference genome is indicated and it is represented
accumulated in the Y axis up to the full genomes mixture (up to 400%).

DISCUSSION

Performance of piPolB in alternative MDA methods

The combination of 5'-3' DNA polymerase and DNA primase activities, coupled with 3'-5' proofreading, together with the ability for translesion synthesis and strand displacement capacity, suggests that the recombinant *E. coli* piPolB has potential for its application in novel MDA methods (39). Here we developed new MDA protocols consisting of the use of piPolB as a single enzyme or coupled with another faithful, highly processive DNA polymerase, such as Φ29DNAP. The performance of these new methods was analyzed in comparison with commercially available MDA kits for amplification of genomic sequences with moderate (45%) to high-GC content (70%), to evaluate the GC bias of each method, which has been described in detail as a major source of bias in MDA as well as in short-read high-throughput sequencing (26, 28, 38, 59, 62). We performed a detailed comparative analysis of the short reads sequenced samples, either unamplified or after isothermal 30 ºC amplification with different piPolB-based protocols. Sequencing of DNA products synthesized with piPolB MDA has revealed two major drawbacks to using this DNA polymerase as a single synthesizing enzyme. First, the piPolB DNA product appears to have a hyperbranched structure resulting from repeated priming events, which may impair competent WGA and also hinders sequencing library preparation. We also detected a high proportion of unknown DNA sequences. Random primers and the presence of preexisting contaminant DNA have been referred to as the main sources of artifacts in MDA (24, 25, 27, 64), but that spurious DNA has not yet been analyzed in detail by high-throughput sequencing. In the case of the piPolB MDA products, there are no RPs in the reaction and high-throughput sequencing ruled out contamination for the not assigned reads during Metaphlan profiling. Therefore, we hypothesized that piPolB is capable of synthesizing DNA in a template-independent manner, commonly referred to as *ab initio* DNA synthesis. Although neglected or controversial in the literature, *ab initio* DNA synthesis has been reported for several DNA polymerases, particularly those of ancient origin (65–68), as is likely the case for piPolB (39). Consistent with the literature, these spurious DNA products generated by piPolB consist of low-complexity, repetitive, and low-GC DNA sequences (Tables S9-S12). Overall, these results highlight the limitations of piPolB for MDA, but also caution against the presence of *ab initio* DNA synthesis in whole-genome amplification reactions under certain reactions, that may be overlooked in MDA products. We believe that this should be further addressed in the future by developing modified protocols that could control this activity using SSBs or other factors that can prevent the production of spurious DNA (69).

The interplay of piPolB and Φ29DNAP in the new piMDA protocol outperforms available MDA kits in amplifying sequences with high-GC content

As expected (28), GC bias in the amplified samples is the major limitation to homogeneous and reliable coverage and hinders the sequence assembling. Despite the low complexity of the amplified genomic sample, we found large differences in sequence coverage and assembly quality among the different WGA protocols tested. Thus, WGA amplification with the traditional random-primed MDA protocol was particularly poor for the genomes with high-GC content and promoted a very strong bias toward intermediate-GC sequences.

This negative bias was only slightly reduced in the sample generated with
PrimPol-MDA, with a similar pattern of overrepresented sequences. Despite a
bias toward high-GC sequences, the combination of piPolB and Φ29DNAP
achieved a more competent amplification of our sample, obtaining good
assemblies, comparable to those of unamplified control samples.
Previous analyses reported either similar GC bias in RP-MDA and PrimPol-
MDA (37) or better GC representation in PrimPol-MDA (38), using different DNA
samples. Consistent with the latter, the use of T7 gp4 primase-helicase instead
of random primers (pWGA) reduced GC bias and allowed the identification of
high-GC species missing in MDA product, although overrepresentation of
intermediate GC sequences was still similar to classical MDA (26). In this case,
the authors proposed that the helicase activity of gp4 might favor the
amplification of high-GC sequences (26). In the piMDA and piMDA+D methods,
the priming and initial DNA synthesis are performed by the piPolB, whose
strand displacement capacity might play a similar role. However, piPolB shows
a strong bias towards 40-50% GC when used alone for MDA, downplaying that
mechanism. Moreover, the alkaline denaturation step increases the coverage
depth and breadth of all the reference genomes (Figure 6B), suggesting that
high-GC content limits the initiation and also the elongation of new DNA strands
by piPolB, downplaying a relevant contribution of strand displacement capacity
to reduce GC bias.
Remarkably, in RP-MDA, PrimPol-MDA, and piMDA(+D) most of the
amplification is performed by the same enzyme, the Φ29DNAP, suggesting that
the random primers or the onset of DNA synthesis are the key factors
determining the GC bias. The random primers used in MDA are usually 6-mers,
and the primers of PrimPol or gp4 are also very short DNA or RNA
oligonucleotides, respectively (37, 70). However, piPolB can processively
synthesize longer DNA molecules of up to 6 kb (39). Conversely, greater
stability of high-GC sequences would reduce processivity and possibly favor
dissociation of piPolB and distributive DNA synthesis. This would generate a
higher number of initiation products, albeit longer than random primers or
PrimPol/gp4-synthesized primers, which in the case of piMDA(+D) could be
resumed processively by the Φ29DNAP. This is consistent with the fact that the
denaturation step, which would allow a higher rate of priming events by piPolB,
slightly increases the GC bias in piMDA+D but not in piPolB+D (Figure 6).
Furthermore, because piPolB exhibits translesion synthesis capacity (39), it is
reasonable to speculate that piPolB-mediated WGA may offer some
advantages in amplifying damaged DNA, which is often present in metagenomic
DNA samples. However, since piMDA uses Φ29DNAP as the main
amplification enzyme, we do not expect significant differences compared with
related MDA methods. Rather, an improved WGA method based only on piPolB
would be more suitable for damaged DNA samples.
789
Consistency of piMDA performance among duplicates and protocol
modifications
The high number of reads obtained and the use of duplicates for some of the
samples allowed us to analyze in detail the performance and variability of each
of the amplification methods used and also the TruSeq library generation and
Illumina sequencing. The overall results of the duplicate samples are very
similar, although the results of the second sequencing batch resulted in a

slightly lower GC bias and a better assembly that contained fewer and larger
contigs and fewer misassemblies, among other parameters (Table S6-S8).
Population variant detection was also better for samples from the second
sequencing batch, particularly for RP-MDA2 and piMDA+D2 samples, as
compared with RP-MDA and piMDA+D, respectively. Indeed, variant calling in
the piMDA+D2 sample was almost identical to the NA samples.Although
conventional MDA protocols like the commercial kits tested in this work include
a DNA denaturation step, we found that piPolB can perform MDA with dsDNA
templates and decided to analyze the quality of the DNA products obtained with
and without prior denaturation. As expected, the denaturation step increases
the amount of synthesized DNA with shorter reaction times in piMDA+D
compared to the piMDA protocol (Figure 3). In addition, it increases the
alignment rate of reads to reference genomes (Figure 6) and improves
assembly with lower N50 and L50 and fewer misassemblies (Figure 7).
However, as mentioned earlier, the denaturation step can increase GC bias
(Figure 6E) and affects assembly binning (Figure 9). Therefore, unlike other
available methods, piMDA can be successfully used as a simple protocol
without a denaturation step or, if the amount of MDA product for downstream
analysis is limiting, the prior denaturation step can be introduced. We can
conclude that piMDA methods enable proficient WGA of a wide range of
genomes for downstream applications, including those related to the study of
microbiome diversity in different environments, especially in environments
where high-GC microorganisms, such as halophiles or thermophiles, would
predominate. In addition, the results shown here suggest that piMDA has great
potential for application in microbiome studies involving DNA amplification, such
as those using single-cell metagenomics to reconstruct strain-resolved
genomes of microbial communities at once, at the risk of missing poorly
represented sequences with high GC content.

Funding

This work was supported by grants PGC2018-09723-A-I00 and PID2021-
123403NB-I00 funded by MCIN/AEI/10.13039/501100011033 and *ERDF A way
of making Europe* to M.R.R.
C. Egas' laboratory was funded by the European Union´s Horizon 2020
Research and Innovation Program under Grant Agreement No 685474, the
METAFLUIDICS project.
C.D.O. and C.M.C. were holder of Fellowships from the Spanish Ministry of
University (FPU16/02665) and the Spanish Ministry of Science and Innovation
(PRE2019-087304), respectively.

Conflict of interest

MRR is listed as inventor on an international patent application on the use of
piPolB for DNA synthesis. The patent holder or licensee were not involved in
this project in any way.

Acknowledgments

We would like to dedicate this work to the memory of Professor Margarita
Salas, for her long and inspiring support and contributions to the discovery of
piPolB and the early development of this project.

847　We also thank to all members of MRR lab for helpful discussions and critical
848　reading of the manuscript.
849
850

REFERENCES

1. McCombie,W.R., McPherson,J.D. and Mardis,E.R. (2019) Next-Generation Sequencing Technologies. *Cold Spring Harb Perspect Med*, **9**, a036798.

2. van der Meij,K.R.M., Sistermans,E.A., Macville,M.V.E., Stevens,S.J.C., Bax,C.J., Bekker,M.N., Bilardo,C.M., Boon,E.M.J., Boter,M., Diderich,K.E.M., *et al.* (2019) TRIDENT-2: National Implementation of Genome-wide Non-invasive Prenatal Testing as a First-Tier Screening Test in the Netherlands. *The American Journal of Human Genetics*, **105**, 1091–1101.

3. Yamaguchi,I., Watanabe,T., Ohara,O. and Hasegawa,Y. (2019) PCR-free whole exome sequencing: Cost-effective and efficient in detecting rare mutations. *PLOS ONE*, **14**, e0222562.

4. Shi,Y., Wang,G., Lau,H.C.-H. and Yu,J. (2022) Metagenomic Sequencing for Microbial DNA in Human Samples: Emerging Technological Advances. *International Journal of Molecular Sciences*, **23**, 2181.

5. Costea,P.I., Zeller,G., Sunagawa,S., Pelletier,E., Alberti,A., Levenez,F., Tramontano,M., Driessen,M., Hercog,R., Jung,F.-E., *et al.* (2017) Towards standards for human fecal sample processing in metagenomic studies. *Nat Biotechnol*, **35**, 1069–1076.

6. Knudsen,B.E., Bergmark,L., Munk,P., Lukjancenko,O., Priemé,A., Aarestrup,F.M. and Pamp,S.J. (2016) Impact of Sample Type and DNA Isolation Procedure on Genomic Inference of Microbiome Composition. *mSystems*, **1**, e00095-16.

7. Poulsen,C.S., Ekstrøm,C.T., Aarestrup,F.M. and Pamp,S.J. (2022) Library Preparation and Sequencing Platform Introduce Bias in Metagenomic-Based Characterizations of Microbiomes. 10.1101/592154.

8. Stoler,N. and Nekrutenko,A. (2021) Sequencing error profiles of Illumina sequencing instruments. *NAR Genomics and Bioinformatics*, **3**, lqab019.

9. Glökler,J., Lim,T.S., Ida,J. and Frohme,M. (2021) Isothermal amplifications – a comprehensive review on current methods. *Critical Reviews in Biochemistry and Molecular Biology*, **56**, 543–586.

10. Wang,X., Liu,Y., Liu,H., Pan,W., Ren,J., Zheng,X., Tan,Y., Chen,Z., Deng,Y., He,N., *et al.* (2022) Recent advances and application of whole genome amplification in molecular diagnosis and medicine. *MedComm (2020)*, **3**, e116.

11. Czyz,Z.T., Kirsch,S. and Polzer,B. (2015) Principles of Whole-Genome Amplification. *Methods Mol Biol*, **1347**, 1–14.

12. Handyside,A.H., Robinson,M.D., Simpson,R.J., Omar,M.B., Shaw,M.-A., Grudzinskas,J.G. and Rutherford,A. (2004) Isothermal whole genome amplification from single and small numbers of cells: a new era for preimplantation genetic diagnosis of inherited disease. *Mol Hum Reprod*, **10**, 767–772.

13. Hutchison,C.A., Smith,H.O., Pfannkoch,C. and Venter,J.C. (2005) Cell-free cloning using phi29 DNA polymerase. *Proc Natl Acad Sci U S A*, **102**, 17332–6.

14. Lasken,R.S. and Egholm,M. (2003) Whole genome amplification: abundant supplies of DNA from precious samples or clinical specimens. *Trends Biotechnol*, **21**, 531–535.

15. Nelson,J.R. (2014) Random-primed, Phi29 DNA polymerase-based whole genome amplification. *Curr Protoc Mol Biol*, **105**, Unit 15 13.

16. Oliveira,B.B., Veigas,B. and Baptista,P.V. (2021) Isothermal Amplification of

Nucleic Acids: The Race for the Next "Gold Standard". *Frontiers in Sensors*, **2**.

17. Jain,S., S. Dandy,D., J. Geiss,B. and S. Henry,C. (2021) Padlock probe-
based rolling circle amplification lateral flow assay for point-of-need nucleic acid
detection. *Analyst*, **146**, 4340–4347.

18. Salas,M., Holguera,I., Redrejo-Rodriguez,M. and de Vega,M. (2016) DNA-
binding proteins essential for protein-primed bacteriophage Φ29 DNA
replication. *Front Mol Biosci*, **3**, 37.

19. Blanco,L., Bernad,A., Lázaro,J.M., Martín,G., Garmendia,C. and Salas,M.
(1989) Highly efficient DNA synthesis by the phage Φ29 DNA polymerase.
Symmetrical mode of DNA replication. *J Biol Chem*, **264**, 8935–40.

20. Paez,J.G., Lin,M., Beroukhim,R., Lee,J.C., Zhao,X., Richter,D.J., Gabriel,S.,
Herman,P., Sasaki,H., Altshuler,D., *et al.* (2004) Genome coverage and
sequence fidelity of phi29 polymerase-based multiple strand displacement
whole genome amplification. *Nucleic Acids Res*, **32**, e71.

21. Ling,J., Deng,Y., Long,X., Liu,J., Du,H., Cao,B. and Xu,K. (2012) Single-
nucleotide polymorphism array coupled with multiple displacement
amplification: accuracy and spatial resolution for analysis of chromosome copy
numbers in few cells. *Biotechnol Appl Biochem*, **59**, 35–44.

22. Ahsanuddin,S., Afshinnekoo,E., Gandara,J., Hakyemezoğlu,M., Bezdan,D.,
Minot,S., Greenfield,N. and Mason,C.E. (2017) Assessment of REPLI-g Multiple
Displacement Whole Genome Amplification (WGA) Techniques for
Metagenomic Applications. *J Biomol Tech*, **28**, 46–55.

23. Lasken,R.S. and Stockwell,T.B. (2007) Mechanism of chimera formation
during the Multiple Displacement Amplification reaction. *BMC Biotechnol*, **7**, 19.

24. Lage,J.M., Leamon,J.H., Pejovic,T., Hamann,S., Lacey,M., Dillon,D.,
Segraves,R., Vossbrinck,B., González,A., Pinkel,D., *et al.* (2003) Whole
Genome Analysis of Genetic Alterations in Small DNA Samples Using
Hyperbranched Strand Displacement Amplification and Array–CGH. *Genome*
Res., **13**, 294–307.

25. Sabina,J. and Leamon,J.H. (2015) Bias in Whole Genome Amplification:
Causes and Considerations. *Methods Mol Biol*, **1347**, 15–41.

26. Direito,S.O., Zaura,E., Little,M., Ehrenfreund,P. and Roling,W.F. (2014)
Systematic evaluation of bias in microbial community profiles induced by whole
genome amplification. *Environ Microbiol*, **16**, 643–57.

27. Alsmadi,O., Alkayal,F., Monies,D. and Meyer,B.F. (2009) Specific and
complete human genome amplification with improved yield achieved by phi29
DNA polymerase and a novel primer at elevated temperature. *BMC Res Notes*,
2, 48.

28. Jamal,R., Li,X. and Weidhaas,J. (2021) Template length, concentration and
guanidine and cytosine content influence on multiple displacement amplification
efficiency. *J Microbiol Methods*, **181**, 106146.

29. Huang,L., Ma,F., Chapman,A., Lu,S. and Xie,X.S. (2015) Single-Cell Whole-
Genome Amplification and Sequencing: Methodology and Applications. *Annu*
Rev Genomics Hum Genet, **16**, 79–102.

30. López-Labrador,F.X., Brown,J.R., Fischer,N., Harvala,H., Van
Boheemen,S., Cinek,O., Sayiner,A., Madsen,T.V., Auvinen,E., Kufner,V., *et al.*
(2021) Recommendations for the introduction of metagenomic high-throughput
sequencing in clinical virology, part I: Wet lab procedure. *Journal of Clinical*
Virology, **134**, 104691.

31. Regnault,B., Bigot,T., Ma,L., Pérot,P., Temmam,S. and Eloit,M. (2021)

Deep Impact of Random Amplification and Library Construction Methods on
Viral Metagenomics Results. *Viruses*, **13**, 253.
32. de Vega,M., Lázaro,J.M., Mencía,M., Blanco,L. and Salas,M. (2010)
Improvement of Φ29 DNA polymerase amplification performance by fusion of
DNA binding motifs. *Proc Natl Acad Sci USA*, **107**, 16506–11.
33. Ordóñez,C.D., Lechuga,A., Salas,M. and Redrejo-Rodríguez,M. (2020)
Engineered viral DNA polymerase with enhanced DNA amplification capacity: a
proof-of-concept of isothermal amplification of damaged DNA. *Sci Rep*, **10**,
15046.
34. Povilaitis,T., Alzbutas,G., Sukackaite,R., Siurkus,J. and Skirgaila,R. (2016)
In vitro evolution of phi29 DNA polymerase using isothermal compartmentalized
self replication technique. *Protein Eng Des Sel*, **29**, 617–628.
35. Mendez,P., Fang,L.T., Jablons,D.M. and Kim,I.-J. (2017) Systematic
comparison of two whole-genome amplification methods for targeted next-
generation sequencing using frozen and FFPE normal and cancer tissues. *Sci
Rep*, **7**, 4055.
36. Deleye,L., De Coninck,D., Dheedene,A., De Sutter,P., Menten,B.,
Deforce,D. and Van Nieuwerburgh,F. (2016) Performance of a TthPrimPol-
based whole genome amplification kit for copy number alteration detection
using massively parallel sequencing. *Sci Rep*, **6**, 31825.
37. Picher,A.J., Budeus,B., Wafzig,O., Kruger,C., Garcia-Gomez,S., Martinez-
Jimenez,M.I., Diaz-Talavera,A., Weber,D., Blanco,L. and Schneider,A. (2016)
TruePrime is a novel method for whole-genome amplification from single cells
based on TthPrimPol. *Nature communications*, **7**, 13296.
38. Parras-Moltó,M., Rodríguez-Galet,A., Suárez-Rodríguez,P. and López-
Bueno,A. (2018) Evaluation of bias induced by viral enrichment and random
amplification protocols in metagenomic surveys of saliva DNA viruses.
Microbiome, **6**, 119.
39. Redrejo-Rodríguez,M., Ordóñez,C.D., Berjón-Otero,M., Moreno-
González,J., Aparicio-Maldonado,C., Forterre,P., Salas,M. and Krupovic,M.
(2017) Primer-Independent DNA Synthesis by a Family B DNA Polymerase
from Self-Replicating Mobile Genetic Elements. *Cell reports*, **21**, 1574–1587.
40. Zyrina,N.V., Antipova,V.N. and Zheleznaya,L.A. (2014) Ab initio synthesis
by DNA polymerases. *FEMS Microbiology Letters*, **351**, 1–6.
41. Michel, Janine F.,C.,Brigitte (1969) Sélection de mutants de Bacillus subtilis
bloqués au début de la sporulation. *Annales de l'Institut Pasteur*, **116**, 3–18.
42. Lázaro,J.M., Blanco,L. and Salas,M. (1995) Purification of bacteriophage
Φ29 DNA polymerase. *Methods in Enzymology*, **262**, 42–9.
43. Shen,W., Le,S., Li,Y. and Hu,F. (2016) SeqKit: A Cross-Platform and
Ultrafast Toolkit for FASTA/Q File Manipulation. *PLoS One*, **11**, e0163962.
44. Ewels,P., Magnusson,M., Lundin,S. and Käller,M. (2016) MultiQC:
summarize analysis results for multiple tools and samples in a single report.
Bioinformatics, **32**, 3047–3048.
45. Blanco-Miguez,A., Beghini,F., Cumbo,F., McIver,L.J., Thompson,K.N.,
Zolfo,M., Manghi,P., Dubois,L., Huang,K.D., Thomas,A.M., *et al.* (2022)
Extending and improving metagenomic taxonomic profiling with uncharacterized
species with MetaPhlAn 4. 10.1101/2022.08.22.504593.
46. Langmead,B. and Salzberg,S.L. (2012) Fast gapped-read alignment with
Bowtie 2. *Nat Methods*, **9**, 357–359.
47. Chen,S., Zhou,Y., Chen,Y. and Gu,J. (2018) fastp: an ultra-fast all-in-one

FASTQ preprocessor. *Bioinformatics*, **34**, i884–i890.

48. Nurk,S., Meleshko,D., Korobeynikov,A. and Pevzner,P.A. (2017)
metaSPAdes: a new versatile metagenomic assembler. *Genome Res*, **27**, 824–
834.

49. Mikheenko,A., Saveliev,V. and Gurevich,A. (2016) MetaQUAST: evaluation
of metagenome assemblies. *Bioinformatics*, **32**, 1088–1090.

50. Schwengers,O., Jelonek,L., Dieckmann,M.A., Beyvers,S., Blom,J. and
Goesmann,A. (2021) Bakta: rapid and standardized annotation of bacterial
genomes via alignment-free sequence identification. *Microb Genom*, **7**, 000685.

51. Mapleson,D., Garcia Accinelli,G., Kettleborough,G., Wright,J. and
Clavijo,B.J. (2017) KAT: a K-mer analysis toolkit to quality control NGS datasets
and genome assemblies. *Bioinformatics*, **33**, 574–576.

52. Kelley,D.R., Schatz,M.C. and Salzberg,S.L. (2010) Quake: quality-aware
detection and correction of sequencing errors. *Genome Biol*, **11**, R116.

53. Mallawaarachchi,V. and Lin,Y. (2022) Accurate Binning of Metagenomic
Contigs Using Composition, Coverage, and Assembly Graphs. *J Comput Biol*,
29, 1357–1376.

54. Ondov,B.D., Treangen,T.J., Melsted,P., Mallonee,A.B., Bergman,N.H.,
Koren,S. and Phillippy,A.M. (2016) Mash: fast genome and metagenome
distance estimation using MinHash. *Genome Biol*, **17**, 132.

55. RStudio Team (2020) RStudio: Integrated Development Environment for R
RStudio, PBC., Boston, MA.

56. Wickham,H. (2009) ggplot2: Elegant Graphics for Data Analysis rstu.
Springer-Verlag, New York.

57. Wei,T. and Simko,V. (2021) R package 'corrplot': Visualization of a
Correlation Matrix.

58. Sato,M.P., Ogura,Y., Nakamura,K., Nishida,R., Gotoh,Y., Hayashi,M.,
Hisatsune,J., Sugai,M., Takehiko,I. and Hayashi,T. (2019) Comparison of the
sequencing bias of currently available library preparation kits for Illumina
sequencing of bacterial genomes and metagenomes. *DNA Research*, **26**, 391–
398.

59. Benjamini,Y. and Speed,T.P. (2012) Summarizing and correcting the GC
content bias in high-throughput sequencing. *Nucleic Acids Res*, **40**, e72.

60. Zyrina,N.V., Zheleznaya,L.A., Dvoretsky,E.V., Vasiliev,V.D., Chernov,A. and
Matvienko,N.I. (2007) N.BspD6I DNA nickase strongly stimulates template-
independent synthesis of non-palindromic repetitive DNA by Bst DNA
polymerase. *Biol Chem*, **388**, 367–372.

61. Pinard,R., de Winter,A., Sarkis,G.J., Gerstein,M.B., Tartaro,K.R., Plant,R.N.,
Egholm,M., Rothberg,J.M. and Leamon,J.H. (2006) Assessment of whole
genome amplification-induced bias through high-throughput, massively parallel
whole genome sequencing. *BMC Genomics*, **7**, 216.

62. Chen,Y.-C., Liu,T., Yu,C.-H., Chiang,T.-Y. and Hwang,C.-C. (2013) Effects
of GC bias in next-generation-sequencing data on de novo genome assembly.
PLoS One, **8**, e62856.

63. Bush,S.J., Foster,D., Eyre,D.W., Clark,E.L., De Maio,N., Shaw,L.P.,
Stoesser,N., Peto,T.E.A., Crook,D.W. and Walker,A.S. (2020) Genomic
diversity affects the accuracy of bacterial single-nucleotide polymorphism-
calling pipelines. *Gigascience*, **9**, giaa007.

64. Pan,X., Urban,A.E., Palejev,D., Schulz,V., Grubert,F., Hu,Y., Snyder,M. and
Weissman,S.M. (2008) A procedure for highly specific, sensitive, and unbiased

whole-genome amplification. *Proc Natl Acad Sci U S A*, **105**, 15499–15504.

65. Ogata,N. and Miura,T. (1997) Genetic information 'created' by archaebacterial DNA polymerase. *Biochem J*, **324 (Pt 2)**, 667–671.

66. Béguin,P., Gill,S., Charpin,N. and Forterre,P. (2015) Synergistic template-free synthesis of dsDNA by Thermococcus nautili primase PolpTN2, DNA polymerase PolB, and pTN2 helicase. *Extremophiles*, **19**, 69–76.

67. Zyrina,N.V. and Antipova,V.N. (2021) Nonspecific Synthesis in the Reactions of Isothermal Nucleic Acid Amplification. *Biochemistry Moscow*, **86**, 887–897.

68. Liang,X., Jensen,K. and Frank-Kamenetskii,M.D. (2004) Very efficient template/primer-independent DNA synthesis by thermophilic DNA polymerase in the presence of a thermophilic restriction endonuclease. *Biochemistry*, **43**, 13459–13466.

69. Zyrina,N.V., Artyukh,R.I., Svad'bina,I.V., Zheleznaya,L.A. and Matvienko,N.I. (2012) The effect of single-stranded DNA binding proteins on template/primer-independent DNA synthesis in the presence of nicking endonuclease Nt.BspD6I. *Russ J Bioorg Chem*, **38**, 171–176.

70. Tabor,S. and Richardson,C.C. (1981) Template recognition sequence for RNA primer synthesis by gene 4 protein of bacteriophage T7. *Proc Natl Acad Sci U S A*, **78**, 205–209.

A Overall alignment rate

B Coverage Breadth

C Coverage Depth

D Mismatch rate

E

Mapped sample	GC%	R
NA	49.5%	-0.93
NA2	52.4%	-0.53
RP-MDA	47.4%	-0.91
RP-MDA2	48.1%	-0.94
PrimPol-MDA	47.3%	-0.95
piPolB-MDA	33.2%	-0.12
piPolB+D-MDA	33.2%	-0.20
piMDA	51%	0.89
piMDA2	54.9%	0.91
piMDA+D	63%	0.92
piMDA+D2	57.2%	0.91