Cortical and subcortical substrates of minutes and days-long object value memory in humans

Sepideh Farmani[2], Kiomars Sharifi[1], Ali Ghazizadeh[1,2]

[1] Bio-intelligence Research Unit, Sharif Brain Center, Electrical Engineering Department, Sharif University of Technology, Tehran, Iran

[2] School of Cognitive Sciences, Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

Obtaining valuable objects motivates many of our daily decisions. However, the neural underpinnings of object processing based on the value memory in humans are not yet fully understood. Here, we used whole-brain fMRI while participants passively viewed objects before, minutes after, and 1-70 days after value training to examine activations due to value memory. The behavioral performance showed significant value memory for objects, which nevertheless faded over days after training. Strong value discrimination was observed shortly after training in the occipital, ventral temporal, interparietal, and prefrontal areas. Days after training, activation in the prefrontal and parietal regions disappeared, whereas the ventral temporal areas showed sustained activation. In addition, days-long value responses emerged in certain subcortical regions, including the caudate, ventral striatum, and thalamus, with the former two showing a significant correlation with participants' performance in days-long memory. Resting-state analysis revealed that the functional distance of value-coding brain regions from clusters, which included the intraparietal and striatal areas, determined the strength of their minutes- and days-long value memory, respectively. These findings shed light on the neural basis of value memory in humans with implications for object habit formation and cross-species comparisons.

Keywords: object value, long-term memory, fMRI, cortical, subcortical, resting state, human

Introduction

Value-based memory, which relies on past stimulus-response associations, can evoke automatic responses independent of immediate outcomes (Dickinson & Balleine, 2002; Graybiel, 2008; Ashby et al., 2010; Ghazizadeh et al., 2016a). Such automatic behavior arises from stable object-value pairings formed through repeated experiences and can persist over a long period (Kim & Hikosaka, 2015; Anderson & Yantis, 2013). While habitual processes supported by value-based memory lead to efficient behavior on the positive side (Ghazizadeh et al., 2016b), on the negative side, they underlie disruptive attention to distractors (Anderson et al., 2011) and maladaptive behaviors in a number of neuropsychiatric disorders such as Tourette's syndrome (Delorme et al., 2016) and drug addiction (Everitt & Robbins, 2005; Sjoerds et al., 2014).

A substantial body of work in animals has focused on the neural mechanisms of value-based memory. Electrophysiological evidence supports the role of the posterior basal ganglia in long-term value memory in non-human primates (NHPs) (Yasuda et al., 2012; Kim & Hikosaka, 2013; Kim et al., 2015). Furthermore, robust value discrimination of objects in regions of the temporal and prefrontal cortex as well as in the striatum, amygdala, and claustrum, is demonstrated using fMRI (Ghazizadeh et al., 2018a). Notably, these differential responses persist for several months after training (Yasuda et al., 2012; Ghazizadeh et al., 2018a; Ghazizadeh & Hikosaka, 2021). The neural correlates of value memory in the human brain are remarkably less clear. A number of human studies aimed to distinguish neural representations of goal-directed and habitual processing across the whole brain, suggesting that the dorsolateral striatum is involved in habitual processing, whereas the dorsomedial striatum and orbitofrontal cortex (OFC) are implicated in goal-directed processing (Tricomi et al., 2009; de Wit et al., 2012; Wunderlich et al., 2012; Graybiel & Grafton, 2015; Glascher et al., 2010; Liljeholm et al., 2015; Daw & O'Doherty, 2014; McNamee et al., 2015). Moreover, previously reward-associated objects, despite being task-irrelevant, are shown to evoke preferential responses in the caudate tail and extra-striate visual cortex (Anderson et al., 2014). In addition, the ventral striatum (VS) has recently been shown to be involved in retaining long-term value memory of objects (Kang et al., 2021). However, the whole-brain functional circuitry of object value memory spanning minutes to days remains unexplored.

In the current study, participants were trained to associate novel fractal stimuli with monetary reward or no reward. Brain responses to the passive viewing of objects were measured before, minutes after, and days after the final training session using fMRI. Value training was conducted across three sessions to elicit stable value-based memory (Wimmer et al., 2018). We adopted the passive-viewing paradigm previously used to investigate the long-term value memory of objects in NHPs (Ghazizadeh et al., 2018a; Ghazizadeh et al., 2018b), which allowed us to explicitly examine the modulations in the visual processing of objects attributed to value memory, from minutes to days. Task and resting state fMRI results revealed a network of prefrontal and parietal cortical regions demonstrating value memory immediately after training, while in the days-long memory, the striatum, thalamus, and ventral temporal areas showed significant activations.

Fig. 1 Experimental timeline and task paradigms. (A) Experimental timeline: Participants learned the value of fractal objects across three sessions. Brain responses were measured using fMRI before, immediately after, and within 1-70 days after the last training session. (B) Stimuli: 40 fractals were randomly assigned into two groups of good and bad objects (the assignment switched across subjects). (C) Value learning task consisted of two types of trials. In force trials, fractal objects were repeatedly associated with a monetary reward (good objects) or no reward (bad objects). In choice trials, participants were instructed to choose the value of a single object shown in the center by pressing the up or down keys without receiving any feedback about the accuracy or monetary outcome. (D) In the novelty test, participants were instructed to indicate the familiarity of a single object shown in the center. The value test was the same as choice trials in the training sessions. (E) fMRI Passive-viewing task consisted of alternating base and probe blocks. In the base blocks, participants looked at the central fixation, and no object was shown. In the probe blocks, a single good or bad object was shown in the center, on the left, or the right hemifield.

Results

Participants learned the value of 40 fractal objects randomly associated with a monetary reward (good objects) or no reward (bad objects) across three sessions (Fig. 1A). The large number of stimuli and arbitrary value assignments resembled real-life experiences involving many objects and ensured that the contrasts between good and bad objects were not confounded by the idiosyncratic physical features of objects (Fig. 1B). During the training, after a central fixation offset, a single fractal was displayed on the left or right side of the screen. Participants had to select that object by pressing the left or right key, and then the associated reward was presented (80% force trials). Interspersed among force trials were unitary choice trials in which single objects were presented, and the participants were instructed to indicate the value of the objects (20% choice trials) (Fig. 1C). No monetary feedback was provided in the unitary choice trials. Following the value training sessions, the neural responses elicited by passive exposure to good and bad objects were measured using fMRI, minutes (post-training fMRI session for minutes-long memory), and within 1-70 days after the third value training session (memory fMRI session for days-long memory) in a block design paradigm consisting of base blocks with only central fixation and probe blocks with a single fractal flashing in the center or periphery (Fig. 1E) (see methods for details). As a control, the same passive-viewing fMRI paradigm was performed prior to the first training session (pre-training fMRI session, Fig. 1A). Participants were instructed to keep the central fixation during the passive-viewing task, and a fixation-break classification indicated no significant difference in the number of fixation breaks between good and bad objects (t_{12}=0.04, p=0.96) (see methods).

A repeated-measures ANOVA performed on the accuracy in choice trials yielded a significant improvement by training ($F_{2,90}$= 6.21, p = 0.003), a significant effect of value ($F_{1,90}$= 15.41, p = 0.0002), and no significant interaction ($F_{2,90}$= 0.62, p = 0.54) (Fig. 2B). Performance significantly increased across the three training sessions (session 1: M = 0.87, SD = 0.01; session 2: M = 0.91, SD = 0.01 and session 3: M = 0.93, SD = 0.01). Furthermore, participants showed higher accuracy for good vs. bad objects in the last two sessions (session 2: t_{15}= 2.54, p = 0.03; session 3: t_{15}= 4.09, p < 0.001). In force trials, performance was high throughout the training sessions (M > 0.99, SD < 0.014) (Fig. S1). In addition, an analogous ANOVA performed on the choice response time showed a significant effect of training with participants becoming faster ($F_{2,90}$ = 15.87, p < 0.001), but no effect of value ($F_{1,90}$= 2.47, p = 0.1) nor interaction ($F_{2,90}$= 0.07, p = 0.9) (Fig 2C). Response time significantly decreased across the three training sessions (session 1: M = 0.1.07, SD = 0.02; session 2: M = 0.96, SD = 0.02 and session 3: M = 0.88, SD = 0.02). Moreover, in the final training session, participants responded faster to good objects compared to bad objects (t_{15} = -2.56, p = 0.02), consistent with previous literature showing faster response to higher value objects (Wimmer et al., 2018; Tankelvitch et al., 2020; Kim et al., 2015). The reduction in response time was also observed in force trials (Fig. S1). The differential responses to good and bad objects, and ~90% choice accuracy for object values, demonstrated that participants successfully learned the value associations across the training sessions.

A days-long value test was performed 1-70 days after the final training session. Despite a significant decrease in average performance for both good and bad objects compared to the initial training session (t_{15} = -3.98, p = 0.001), performance was still significantly above chance for both object types (t_{15} = 2.44, p = 0.03 for bad and t_{15}= 4.48, p < 0.001 for good objects) with performance for good objects remaining higher than bad objects (t_{15} = -3.97, p = 0.001) (Fig 2B). Choice response time for good and bad objects in the value test further decreased

Fig. 2 Humans can readily learn and form days-long value memory of a large number of abstract fractal objects (A) Behavioral results are from choice trials in the training sessions highlighted in light pink. (B) Choice performance across the three training sessions and the memory session. Boxplots show the choice rate of good and bad objects in each session. The central mark within each box indicates the median, and the bottom and top of the box correspond to the 25th and 75th percentiles, respectively. The whiskers extend to the extreme values within 1.5 times the interquartile range. Outliers are plotted as dots. (C) Response time in choice trials across the three training sessions and value test. Same conventions as in (B). (*p < 0.05, **p < 0.01, ***p < 0.005) (D) Choice performance in the value test vs. the duration of the intervening period from the last training session (Pearson's correlation "r" and significance "p" are noted in the plot). (E) Choice performance in the novelty test for good, bad, and novel stimuli. Same conventions as in (B).

compared to the final training session (t_{15} = -7.20, p < 0.001) (Fig 2C). As expected, the decline
in value memory was significantly and negatively correlated with time from the last training
(Fig. 2D).

Reduced performance in session 4 (value test) could have been caused by forgetting the object
values or the objects themselves. In the first scenario, participants should have remembered
that they had seen an object before but should have been skeptical about its value, while in the
second, the object should have looked novel to them. To address this possibility, some of our
participants took part in a novelty test before the memory scan, in which good and bad objects
were presented along with the same number of novel objects. Participants were instructed to
choose whether the object was novel or familiar. Notably, the average performance for old and
new objects was 97.25% and 95% (p < 0.001), respectively, with no difference between good
and bad objects (t_4 = -0.03, p = 0.4) (Fig. 2E). These results suggest that participants had a clear
idea of which objects were novel and that the errors in value judgments were not due to
forgetting previously seen objects, but most likely to forget their value.

Brain areas involved in object value memory

To evaluate the neural substrates of value-based memory, we employed a passive-viewing task
with no reward delivery. Good and bad objects were presented in the scanner in the center or
in the right or left hemifield, and participants were instructed to simply keep their eyes on the
central fixation point (Fig. S2). Given the large number of objects, their random visual features,
and their arbitrary assignment to good and bad groups, differential responses to good and bad
objects could only arise due to previously learned values. As expected, we found no significant
difference in responses to good and bad objects before training (pre-learning session, p < 0.05,
alpha = 0.01, cluster corrected). In contrast, significant differentiations were observed in
several brain regions minutes after the last training session (post-training fMRI session) in the
intraparietal sulcus (IPS), middle occipital gyrus, ventral lateral prefrontal cortex (vlPFC),
insula, fusiform gyrus (FFG), dorsolateral putamen, and parahippocampal gyrus (PHC) (Fig.
3B, Table 1). These regions were more strongly activated in response to good objects than bad
ones. Limiting the good vs. bad contrast in the post-learning session to objects correctly
identified in the final training session (memorized objects) yielded similar results due to the
small number of errors.

Object value discrimination based on days-long memory was found in a subset of regions
activated in the post-training session, including the fusiform gyrus and inferior occipital gyrus.
(Fig. 3C, Table 1). In contrast, certain areas, including the IPS, vlPFC, insula, and PHC, did
not pass the cluster correction in the memory session. Nevertheless, some of these areas, most
notably the IPS, remained significantly active when we limited the good vs. bad contrast to
memorized objects in the value test (Fig S3, Table S1). Interestingly, robust value coding was
observed in several subcortical areas in the memory fMRI session that were not significantly
activated in the post-training scan. This value coding was highly evident in the caudate,
thalamus, and ventral striatum (VS) (Fig. 3C). Together, these results suggest that activity in
the dorsal visual stream and prefrontal cortex was more pronounced in minutes-long memory.
In contrast, the ventral visual stream (including the bilateral fusiform gyrus) and certain
subcortical regions such as the basal ganglia and thalamic circuits, demonstrated enhanced
activation in days-long memory (Fig 3D-E).

Fig. 3 Whole-brain neural correlates of minutes-long and days-long value memory. (A) Experimental timeline: post-learning and memory fMRI sessions are highlighted in light and dark pink, respectively. (B) Brain regions with significant differential responses to good and bad objects in the post-learning session (minutes-long memory) ($p < 0.05$, $\alpha < 0.01$, cluster corrected). (C) Same as B but for the memory fMRI session (days-long memory) ($p < 0.05$, $\alpha < 0.01$, cluster corrected). (D) Cortical regions with significant good vs. bad discrimination in the post-learning session (minutes-long, yellow), in the memory session (days-long, green), and both sessions (pink). (E) Same as D but for subcortical regions.

Table 1. Summary of whole-brain analysis results

Contrast	Regions	Cluster size	x	y	z
Good vs. bad post-learning	R vlPFC	390	-52.5	-22.5	-4.5
	R insular lobe				
	R dorsolateral putamen				
	R hippocampus				
	R amygdala				
	R temporal pole				
	L fusiform gyrus	224	46.5	76.5	-16.5
	L lingual gyrus				
	L parahippocampal gyrus				
	L cerebellum				
	R middle occipital gyrus	191	-31.5	65.5	31.5
	R intraparietal sulcus				
	R superior occipital gyrus				
	R inferior parietal lobule				
	L inferior temporal cortex	171	52.5	58.5	-22.5
	L inferior occipital cortex				
	L superior parietal lobule	149	25.5	70.5	55.5
	L intraparietal sulcus				
	L inferior parietal lobule				
Good vs. bad memory	L thalamus	479	1.5	7.5	10.5
	R thalamus				
	R dorsal caudate				
	R ventral caudate				
	R ventral striatum				
	R lingual gyrus	368	-16.5	88.5	-16.5
	R cerebellum				
	L inferior occipital cortex	359	40.5	88.5	-16.5
	L fusiform gyrus				
	L cerebellum				
	L lingual gyrus				
	L inferior temporal cortex	202	58.5	61.5	-16.5
	L inferior occipital cortex				
	L dorsal caudate	146	10.5	4.5	16.5
	L ventral caudate				
	L ventral striatum				

Clusters of activity exceeding whole-brain p < 0.05, α < 0.01, cluster-corrected (cluster size = 62 voxels for the post-learning and cluster size = 69 voxels for the memory fMRI session).

Resting-state networks are differentially engaged in maintaining object value memory.

The appearance and disappearance of certain cortical and subcortical areas in post-learning and memory sessions suggested that the observed activations might be organized at the network level. To investigate the network properties of value-based memory, we leveraged the functional connectivity among value-coding regions during rest. Pairwise correlations between the resting-state time course of regions significantly active in either the post-training or memory session were computed and converted into a distance matrix (see methods). A 2D embedding of areal distances was implemented using multidimensional scaling, and areas were clustered using a density-based clustering algorithm (Ghazizadeh et al., 2018a). In this embedding, each point corresponded to one area, and a smaller distance between the two points indicated a stronger resting state correlation (Fig. 4A). The results demonstrated that the value-coding regions could be divided into five clusters based on functional connectivity, namely: 1. Thalamo-striatal cluster, 2. Temporo-occipito-cerebellar cluster, 3. Parieto-occipital cluster, 4. Parahippocampal cluster, and 5. Prefrontal cluster. We compared the average value signal across each cluster in the post-training and memory sessions to gain further insight into the relationship between these clusters and the value signal. The results indicated that in the thalamo-striatal cluster, the value signal was significantly higher in the memory session than in the post-training session, while the opposite effect was observed in the parieto-occipital, parahippocampal and prefrontal clusters (Fig. 4B). Furthermore, the value signal in the memory session (but not in the post-training session) showed a negative relationship with the distance from the center of mass of the thalamo-striatal and temporo-occipito-cerebellar clusters together, whereas the value signal in the post-training session was negatively correlated with distance from the center of the parieto-occipital cluster, showing that areas functionally connected to the thalamo-striatal and temporo-occipito-cerebellar clusters played an essential role in encoding days-long value memory, while areas functionally connected to the parieto-occipital cluster, were more engaged in minutes-long memory after the value training (Fig. 4 C-D).

Fig. 4 Functional connectivity between regions activated by minutes- or days-long value memory. (A) Functional distance of brain regions with significant value discrimination shown in 2D space using low dimensional embedding and clustered using a density-based algorithm. Clusters are plotted with different colors. (B) Good vs. bad beta coefficients for different clusters in post-learning and memory fMRI sessions. (C) Good vs. bad activation in post-learning and memory sessions as a function of the Euclidean distance from the center of mass of the thalamo-striatal and temporo-occipito-cerebellar clusters. (D) Same as C but for the Euclidian distance from the center of mass of the parieto-occipital cluster. (Pearson's correlation "r" is noted in (C), (D), *p < 0.05, **p < 0.01, ***p < 0.005).

Activation in the caudate nucleus correlates with days-long memory performance.

Our results revealed regions with significant value discrimination in the memory session. In particular, the thalamo-striatal cluster was significantly active in the memory session. However, it is not clear which areas in this or other clusters contributed to the participants' behavior in the value test. Remarkably, the magnitude of value-dependent activation in the right and left caudate and in the right VS showed a significant positive correlation with the performance in the value (Fig. 5A, Fig. S4). In our experiment, days-long value memory was tested over a relatively long period of 1-70 days. As shown previously, memory performance was dependent on the time elapsed since the last training session and decreased significantly over time (Fig. 1D). Similar to performance, activation of the caudate nucleus decreased significantly with the duration of the intervening period (Fig. 5B, Fig. S4). Notably, among the regions showing activation in the memory scan, activation in the right caudate nucleus also passed the cluster threshold for the memory > post-learning contrast (Fig. 5C) (Fig. 3-4).

Fig. 5 Activation in the caudate correlates with days-long value memory performance. (A) Good vs. bad beta coefficient averaged over right caudate voxels as a function of participants' performance in the memory session. (B) Same format as (A) but as a function of time elapsed since the last training session (Pearson's correlation "r" and significance "p" are noted in (A) and (B)). (C) Areas activated by the memory minus post-learning value contrast ($p < 0.05$, $\alpha < 0.01$, cluster corrected, voxel size = 68).

Discussion

Building on previous animal research that has demonstrated robust sensitivity to object value memory in multiple cortical and subcortical areas (Ghazizadeh et al., 2018a), we investigated the neural underpinnings of value-based memory in humans. Participants were trained across three sessions to learn the value of novel fractal objects. Neural responses were measured using fMRI before, 15-30 minutes after (minutes-long memory), and days after (days-long memory) the last training session. We employed a passive-viewing task in the absence of reward to assess visual responses modulated by the previously learned values of objects. Minutes after the final training session, we found enhanced responses in the middle and inferior occipital gyrus, FFG, PHC, vlPFC, insula, and IPS (Fig. 3). Despite the time-dependent decline in performance, participants were able to remember the value of many objects significantly above chance days after the last training (Fig. 2). This decline in performance was specific to forgetting the object values, not the objects themselves (Fig. 2E). Imaging results during this time showed persistent activation in some areas, including the FFG, and inferior occipital gyrus. Remarkably, we observed the emergence of value coding in the caudate, thalamus, and VS in the memory session (Fig. 3). Further investigation revealed that the activated regions in either scan formed several functional networks with different behavior in minutes-long and days-long memory (Fig. 4). While the functional distance from the parieto-occipital cluster determined the strength of minutes-long value memory, it was the distance from thalamo-striatal and temporo-occipito-cerebellar clusters that determined the strength of days-long memory. In particular. among the regions with strong days-long value memory, the dorsal caudate showed greater activation in the memory vs. post-learning scans, which was significantly correlated with participants' performance in the days-long value test (Fig. 5).

Minutes after training, regions in the parieto-occipital, prefrontal, and parahippocampal clusters exhibited enhanced value coding. Value discrimination in the temporal and occipital areas is in accordance with previous studies showing value-based modulation of representations in the visual cortical regions (Anderson, 2016; Qi et al., 2013; Hickey and Peelen, 2015; Tankelevitch et al., 2020; Ataei et al., 2022). Nevertheless, activation in several cortical areas decreased significantly across days. One possible explanation could be that some of these areas are more involved in the minutes-long value memory of objects, and as consolidation takes place, memory is stored elsewhere in the brain. Among these fading areas, the prefrontal and parietal cortices may be involved in regulating and forming lasting value memories but could disengage once the memories are ingrained (Cabeza et al., 2012). On the other hand, it is known that repeated pairing of objects with rewards induces habitual processing and leads to the formation of stable value memories (Kim & Hikosaka, 2015; Kim et al., 2015; Yasuda et al., 2012; Ghazizadeh et al., 2016b). Previous studies have provided evidence for the involvement of the dorsolateral striatum in habitual processing (Tricomi et al., 2009; de Wit et al., 2012; Wunderlich et al., 2012; Lee et al., 2013). These findings accord with electrophysiological studies in animals (Yin & Knowlton, 2006; Kim & Hikosaka, 2013). Interestingly, evidence from lesion studies has indicated that after habitual responses were established in the dorsal striatum (DS), the disconnection of the VS and DS disrupted habitual seeking behavior supporting the mediating role of the VS (Belin & Everitt, 2008). Consistent with these results, we found significant activity in the caudate and VS in the memory session, significantly correlated with participants' behavioral days-long memory (Fig. 3,5).

In a previous study, the VS has been shown to be significantly activated in response to high-value objects presented centrally 2-5 days after value training (Kang et al., 2021). This aligns with our

results, demonstrating the emergence of activity in the VS in the days-long memory session.
However, in contrast to our results, in the aforementioned study, no activation was found
beyond the VS in other cortical and subcortical areas, and also no behavioral measure of
participants' performance in the days-long memory session was provided. One possibility for
the discrepancy between the whole-brain results is the differences in task designs. In Kang et
al. (2021), object values were presented via binary choice trials, which could lead to a selection
bias, modulating brain responses irrelevant to reward value (Failing & Theeuwes, 2018). In
our task, selection bias was avoided by training participants using force trials and probing the
value learning using unitary choice trials (Fig. 1). In addition, in our passive-viewing task,
objects were presented in both the periphery and the center, which might have resulted in the
engagement of several cortical and subcortical areas that are known to have lateralized
receptive fields (Hasson et al., 2002; Yu et al., 2014).

Extensive work in human literature has focused on value-driven attentional capture (VDAC),
which has revealed the influence of value learning on attention. This influence persists even
when the previously reward-associated stimuli are currently task-irrelevant (Della Libera &
Chelazzi, 2006; Hickey et al., 2010; Anderson et al., 2011; Anderson & Yantis, 2013;
Anderson, 2016; Failing & Theeuwes, 2018). Neuroimaging studies have shown that the
VDAC leads to greater activity in the visual cortex (Saproo & Serences, 2010; Hopf et al.,
2015; MacLean & Giesbrecht, 2015; Anderson et al., 2014; Hickey & Peelen, 2015; Anderson,
2017), posterior parietal cortex (Anderson et al., 2014; Barbaro et al., 2017; Lee & Shomstein,
2013; Qi et al., 2013; Anderson, 2017), and the caudate tail (Anderson et al., 2014; Anderson,
2016; Moore & Zirnsak, 2017). Brain responses contributing to salience coding (such as
VDAC) and value-based memory are highly entangled. Therefore, some of the activated areas
in the current study may also be involved in other processes. Indeed, it is known that the insula
and thalamus encode stimulus salience irrespective of the valance (Seeley, 2019; Zhou et al.,
2021). In addition, the role of vlPFC in salience coding beyond value has recently been
confirmed in macaques (Ghazizadeh & Hikosaka, 2022). Nevertheless, we emphasize that
regardless of the interpretation and the downstream functionality of the observed activations in
our study, they were all caused by the value memory of objects. Future studies are needed to
elucidate the distinct functionality of neural representations attributed to attentional, emotional,
motivational, or pure valuation domains.

In summary, our results revealed cortical and subcortical networks spanning minutes-long to
days-long memory of previously learned object values. While the parietal, prefrontal, and
parahippocampal networks were activated within minutes after value learning, the visual
ventral stream areas and the thalamo-striatal functional circuitry retained the value memory for
days. Although the habitual mechanisms underlying value-based memory can render efficient
visual processing of objects (Ghazizadeh et al., 2016b) and free up precious cognitive resources
for new experiences, the role of habit formation in maladaptive behaviors cannot be overstated
(Delorme et al., 2016; Everitt & Robbins, 2005; Sjoerds et al., 2014). Causal manipulations in
these areas using electrical, magnetic, chemical, or genetic methods to disrupt such value-based
maladaptive behaviors is a crucial direction for future investigations (Luigjes et al., 2019; Song
et al., 2019; Antonelli et al., 2021).

Materials and Methods

Participants. 20 healthy adults with normal or corrected-to-normal vision participated in the present study. Four participants were excluded due to either incomplete sessions or excessive head motion. Our final sample included 16 participants (6 females, mean age: 22.5 years, range:19–32). All participants provided written informed consent and were compensated for their participation with a flat monetary reward and an additional monetary bonus based on their performance on the value learning task. The experimental protocol was approved by the ethics committee of the Institute for Research in Fundamental Sciences (IPM) (protocol number 98/60.1/3025).

By performing power analysis on data from a pilot experiment (n = 6, mean beta = 0.02, SD = 0.019), we found that the estimated sample size to achieve an 80% power (β=0.2) at α=0.05 for detecting a difference between the good and bad objects in the caudate was 10 (Faul et al., 2009).

Stimuli. 80 fractal images similar to those used to study the neural basis of value-based memory in NHPs were utilized (Ghazizadeh et al., 2018a) (Fig. 1B). The parameters, including size, color, etc., were randomly selected for each fractal. 40 fractals were used in the value learning task. Half of these stimuli were associated with reward (good objects), and the other half received no reward (bad objects) during training. The order of objects was counterbalanced across participants. The remaining 40 fractals were used as novel stimuli in the novelty test.

Experimental Procedure. To examine the neural substrates of value-based memory, participants were trained over three sessions outside the scanner, and their brain responses were measured across three rounds of fMRI scanning. The scans took place before the initial training session (pre-learning session), immediately after the final training session (post-learning session), and one or more days (1- 70) after completing the training (memory session) (Fig. 1A). All tasks were presented using MATLAB (MathWorks) and Psychtoolbox-3 (Kleiner et al., 2007).

Value Learning Task. Participants completed three training sessions, each consisting of five blocks. Each block was performed with a set of 8 fractals (4 good and 4 bad fractals) and comprised 80 trials, 64 force trials, and 16 choice trials. On each force trial, a white fixation dot was presented for 800 ms, followed by a blank screen for 200 ms (Fig. 1C). An object was then presented on the left or right side of the screen for 1000 ms, and participants were required to indicate their choice by pressing the left or right button. After the selection, visual and auditory feedback indicated the outcome for 1000 ms. The successful performance led to a reward (+200 tomans) for the good objects and no reward (0 tomans) for the bad objects. Each trial was followed by an intertrial interval (ITI) of 1000 ms. Choice trials were interspersed among the force trials, yielding training blocks consisting of 4 force trials and 1 choice trial. Each choice trial started with the presentation of a red fixation dot for 800 ms, followed by a blank screen for 200 ms. An object was then presented for 2000 ms in the center, along with the options "good" and "bad" above or below the object (the position of the options was randomized across subjects). Each trial was followed by an intertrial interval (ITI) of 1000 ms. Participants were instructed to indicate the object's value by pressing the up and down arrow keys on the keyboard. No feedback was given on the accuracy of choice.

Novelty Task. 5 participants underwent a novelty test before the memory scan. The task comprised of 80 trials. Each trial started with the presentation of a fixation dot for 800 ms, followed by a 200 ms blank screen (Fig. 1D). An object was then presented in the center for 1000 ms. Participants were required to choose whether they had seen the objects in the training sessions (old) or not (new) by pressing the left and right arrow keys on the keyboard. No feedback was given throughout the task. Each trial was followed by an ITI of 2000 ms. The order of the objects was counterbalanced across participants.

Passive-viewing Task. Neural responses to good and bad objects were measured before the first training session, immediately after, and within 1-70 days of the final training session. Participants engaged in a passive-viewing task across 7 runs. Each run consisted of 36 blocks (18 base and 18 probe) of 10 s duration (Fig. 1E). There were 6 different probe blocks: [good, bad] x [left, center, right]. The order of the blocks was pseudorandomized, and each stimulus was presented three times during each session, once in each of the left, center, or right hemifields. During each probe block, a fractal was flashed (800 ms on, 200 ms off) in the center or one hemifield ($\sim6^{\circ}$ eccentricity across horizontal and 45° oblique directions) while the participant maintained central fixation. The base blocks had the same duration and contingencies as the probe block, except that no object was shown, and participants only had to keep their eyes on the central fixation. The order of objects was pseudorandomized across participants.

Resting-state fMRI. In addition to the passive-viewing task, 12 participants completed resting scans in two to three sessions to examine functional connectivity in the brain. Each resting scan consisted of 36 consecutive base blocks, and participants were instructed to keep their eyes on the central fixation.

Fixation-break Classification. To classify fixation-break vs fixation-hold labels, support vector classification (SVC) was performed using leave-one-session-out cross-validation. The cross-validation procedure involved pre-learning, post-learning, and memory sessions. Participants performed a fixation-break task (2 runs, 90 TRs each) inside the scanner, where they were required to saccade to a new dot point in each TR that represented the break of fixation. The locations of dots matched the locations in which fractals were presented in the passive-viewing task. Data from both the fixation-break and resting-state tasks were used to train and validate the classifier, with an average validation accuracy of 99%. The best classifier was then used to predict fixation breaks in each TR of the fMRI sessions (pre-learning, post-learning, and memory). On average, the classifier predicted that fixation breaks occurred 27.3% of the time during these sessions. Notably, there were no significant differences in the number of fixation breaks observed in the base vs. probe (t_{12}=-0.10, p=0.91), good vs. bad (t_{12}=0.04, p=0.96), and left vs. right blocks (t_{12}=-0.09, p=0.92). These findings suggest that the classifier accurately predicted the occurrence of fixation breaks during the task and that the occurrence of fixation breaks was consistent across different experimental conditions.

fMRI Data Acquisition. Functional and structural imaging was conducted on a 3 T Siemens MAGNETOM Prisma scanner equipped with a 64-channel head coil at National Brain Mapping Laboratory (NBML; Tehran, Iran). For each participant, a whole-brain structural image was acquired using a T1-weighted MPRAGE pulse sequence (TR = 1.8 s, TE = 3.53 ms, voxel size = 1 x 1 x 1 mm^3, flip angle = 7°). Functional images were collected using an EPI sequence (TR = 2 s, TE = 30 ms, voxel size = 3 x 3 x 3 mm^3, flip angle = 90°).

Behavioral Analysis. Behavioral analyses were conducted in MATLAB (MathWorks). To determine whether the learning procedure differentially affected the performance and response time for the good and bad objects, repeated-measures ANOVA was performed, followed by Tukey multiple comparison test. Furthermore, average performance and reaction time across training and memory sessions were compared using t-tests.

fMRI Data Analysis. fMRI data were analyzed using the AFNI/SUMA software package (Cox, 1996; Saad et al., 2004) and MATLAB (MathWorks) scripts. The pre-processing steps included: 1. Detrending and despiking (3dToutcount, 3dDespike); 2. Slice time correction (3dTshift); 3. Alignment to the anatomical T1 image (align_epi_anat.py); 4. Skull stripping and non-linear warping of the T1 image into MNI space (auto_warp.py); 5. Motion correction (3dvolreg) and 6. Non-linear warping of EPI data into MNI space (3dNwarpApply). The resulting EPI time series from all runs were concatenated and converted into percentage change from the mean. General linear model (GLM) analysis is then performed to estimate the contribution of each factor on percentage change from the mean (3dDeconvolve). The model consisted of six regressors ([good, bad] × [left, center, right]) convolved with the canonical hemodynamic response function. Six rigid-body motion parameters (x, y, z dimensions, roll, pitch, and yaw) were included in the GLM as additional regressors to account for the residual effects of participants' motion. These regressors were entered into contrasts to yield 'Good vs. Bad,' 'Left vs. Right,' and 'Center vs. Periphery' responses. After spatial smoothing with a 6 mm FWHM kernel (3dBlurInMask), linear contrasts were taken to a group-level analysis to calculate the mean and test whether the average estimates were significant (3dttest++). Finally, the cluster correction method was applied to find the significant clusters. Estimates of spatial smoothness based on the group residuals were found by simulating noise-only volumes, assuming the autocorrelation function (ACF) was given by a mixed-model and then determining the threshold for the significant clusters (3dClustSim). The minimum cluster size for two-sided $\alpha < 0.05$ and voxelwise significance of $P < 0.05$ was 127 and 134 voxels in the post-learning and memory sessions, respectively.

Functional Connectivity Analysis. Resting-state correlation analysis was carried out using the residuals extracted from each region of interest (ROI). The anatomical parts that were significantly activated in the post-learning or memory session were included in each ROI. The physiological fluctuations of the white matter, the ventricles, and the cerebrospinal fluid (CSF) were regressed out from the residual time series, and bandpass filtering between 0.01 and 0.1 Hz was applied. Then Pearson's correlation between the average signal of each pair of ROIs was calculated. The dissimilarity was measured as 1- ρ, with ρ being the pairwise Pearson's correlation. Next, we applied multidimensional scaling to the dissimilarity matrix to visualize the resting-state distances in 2D (mdscale in MATLAB). Clustering of value coding ROIs based on their resting-state distance was implemented using the density-based clustering algorithm (DBSCAN). It required no prior knowledge of the number of clusters and allowed arbitrarily shaped clusters. The minimum number of data points to define a cluster was set to 2, and the radius of the neighborhood was set to the 75th percentile of all pairwise dissimilarities.

References

Anderson B. A. (2016). The attention habit: how reward learning shapes attentional selection. *Annals of the New York Academy of Sciences*, *1369*(1), 24–39. https://doi.org/10.1111/nyas.12957

Anderson B. A. (2017). Reward processing in the value-driven attention network: reward signals tracking cue identity and location. *Social cognitive and affective neuroscience*, *12*(3), 461–467. https://doi.org/10.1093/scan/nsw141

Anderson, B. A., & Yantis, S. (2013). Persistence of value-driven attentional capture. *Journal of experimental psychology. Human perception and performance*, *39*(1), 6–9. https://doi.org/10.1037/a0030860

Anderson, B. A., Laurent, P. A., & Yantis, S. (2011). Value-driven attentional capture. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(25), 10367–10371. https://doi.org/10.1073/pnas.1104047108

Anderson, B. A., Laurent, P. A., & Yantis, S. (2014). Value-driven attentional priority signals in human basal ganglia and visual cortex. *Brain research*, *1587*, 88–96. https://doi.org/10.1016/j.brainres.2014.08.062

Antonelli, M., Fattore, L., Sestito, L., Di Giuda, D., Diana, M., & Addolorato, G. (2021). Transcranial Magnetic Stimulation: A review about its efficacy in the treatment of alcohol, tobacco and cocaine addiction. *Addictive behaviors*, *114*, 106760. https://doi.org/10.1016/j.addbeh.2020.106760

Ashby, F. G., Turner, B. O., & Horvitz, J. C. (2010). Cortical and basal ganglia contributions to habit learning and automaticity. *Trends in cognitive sciences*, *14*(5), 208–215. https://doi.org/10.1016/j.tics.2010.02.001

Ataei, A., Amini, A., & Ghazizadeh, A. (2022). Gustatory Cortex Is Involved in Evidence Accumulation during Food Choice. *eNeuro*, 9(3), ENEURO.0006-22.2022. https://doi.org/10.1523/ENEURO.0006-22.2022

Barbaro, L., Peelen, M. V., & Hickey, C. (2017). Valence, Not Utility, Underlies Reward-Driven Prioritization in Human Vision. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, *37*(43), 10438–10450. https://doi.org/10.1523/JNEUROSCI.1128-17.2017

Belin, D., & Everitt, B. J. (2008). Cocaine seeking habits depend upon dopamine-dependent serial connectivity linking the ventral with the dorsal striatum. *Neuron*, *57*(3), 432–441. https://doi.org/10.1016/j.neuron.2007.12.019

Cabeza, R., Ciaramelli, E., & Moscovitch, M. (2012). Cognitive contributions of the ventral parietal cortex: an integrative theoretical account. *Trends in cognitive sciences*, *16*(6), 338–352. https://doi.org/10.1016/j.tics.2012.04.008

Cox RW (1996) AFNI: Software for analysis and visualization of functional magnetic resonance neuroimages. *Comput Biomed Res* 29:162–173.

Daw, N.D., & O'Doherty, J.P. (2014). Multiple Systems for Value Learning. https://doi.org/10.1016/B978-0-12-416008-8.00021-8

de Wit, S., Watson, P., Harsay, H. A., Cohen, M. X., van de Vijver, I., & Ridderinkhof, K. R. (2012). Corticostriatal connectivity underlies individual differences in the balance between habitual and goal-directed action control. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, *32*(35), 12066–12075. https://doi.org/10.1523/JNEUROSCI.1088-12.2012

Della Libera, C., & Chelazzi, L. (2006). Visual selective attention and the effects of monetary rewards. *Psychological science*, *17*(3), 222–227. https://doi.org/10.1111/j.1467-9280.2006.01689.x

Delorme, C., Salvador, A., Valabrègue, R., Roze, E., Palminteri, S., Vidailhet, M., de Wit, S., Robbins, T., Hartmann, A., & Worbe, Y. (2016). Enhanced habit formation in Gilles de la Tourette syndrome. *Brain: a journal of neurology*, *139*(Pt 2), 605–615. https://doi.org/10.1093/brain/awv307

Dickinson, Anthony & Balleine, Bernard. (2002). The role of learning in motivation. Stevens' Handbook of Experimental Psychology. Vol. 3.

Everitt, B. J., & Robbins, T. W. (2005). Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nature neuroscience*, *8*(11), 1481–1489. https://doi.org/10.1038/nn1579

Faul, Franz & Erdfelder, Edgar & Buchner, Axel & Lang, Albert-Georg. (2009). Statistical Power Analyses Using G*Power 3.1: Tests for Correlation and Regression Analyses. *Behavior research methods.* 41. 1149-60. 10.3758/BRM.41.4.1149.

Failing, M., & Theeuwes, J. (2018). Selection history: How reward modulates selectivity of visual attention. *Psychonomic bulletin & review*, *25*(2), 514–538. https://doi.org/10.3758/s13423-017-1380-y

Ghazizadeh, A., & Hikosaka, O. (2021). Common coding of expected value and value uncertainty memories in the prefrontal cortex and basal ganglia output. *Science advances*, *7*(20), eabe0693. https://doi.org/10.1126/sciadv.abe0693

Ghazizadeh, A., & Hikosaka, O. (2022). Salience memories formed by value, novelty and aversiveness jointly shape object responses in the prefrontal cortex and basal ganglia. *Nature communications*, *13*(1), 6338. https://doi.org/10.1038/s41467-022-33514-3

Ghazizadeh, A., Griggs, W., & Hikosaka, O. (2016a). Ecological Origins of Object Salience: Reward, Uncertainty, Aversiveness, and Novelty. *Frontiers in neuroscience*, *10*, 378. https://doi.org/10.3389/fnins.2016.00378

Ghazizadeh, A., Griggs, W., Leopold, D. A., & Hikosaka, O. (2018a). Temporal-prefrontal cortical network for discrimination of valuable objects in long-term memory. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(9), E2135–E2144. https://doi.org/10.1073/pnas.1707695115

Ghazizadeh, A., Hong, S., & Hikosaka, O. (2018b). Prefrontal Cortex Represents Long-Term Memory of Object Values for Months. *Current biology: CB*, *28*(14), 2206–2217.e5. https://doi.org/10.1016/j.cub.2018.05.017

Ghazizadeh, Ali & Griggs, Whitney & Hikosaka, Okihide. (2016b). Object-finding skill created by repeated reward experience. *Journal of Vision*. 16. 17. 10.1167/16.10.17. https://doi.org/10.1167/16.10.17

Gläscher, J., Daw, N., Dayan, P., & O'Doherty, J. P. (2010). States versus rewards: dissociable neural prediction error signals underlying model-based and model-free reinforcement learning. *Neuron*, *66*(4), 585–595. https://doi.org/10.1016/j.neuron.2010.04.016

Graybiel A. M. (2008). Habits, rituals, and the evaluative brain. *Annual review of neuroscience*, *31*, 359–387. https://doi.org/10.1146/annurev.neuro.29.051605.112851

Graybiel, A. M., & Grafton, S. T. (2015). The striatum: where skills and habits meet. *Cold Spring Harbor perspectives in biology*, *7*(8), a021691. https://doi.org/10.1101/cshperspect.a021691

Hasson, U., Levy, I., Behrmann, M., Hendler, T., & Malach, R. (2002). Eccentricity bias as an organizing principle for human high-order object areas. *Neuron*, *34*(3), 479–490. https://doi.org/10.1016/s0896-6273(02)00662-1

Hickey, C., & Peelen, M. V. (2015). Neural mechanisms of incentive salience in naturalistic human vision. *Neuron*, *85*(3), 512–518. https://doi.org/10.1016/j.neuron.2014.12.049

Hickey, C., Chelazzi, L., & Theeuwes, J. (2010). Reward changes salience in human vision via the anterior cingulate. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, *30*(33), 11096–11103. https://doi.org/10.1523/JNEUROSCI.1026-10.2010

J.-M. Hopf, M. A. Schoenfeld, A. Buschschulte, A. Rautzenberg, R. M. Krebs & C. N. Boehler (2015) The modulatory impact of reward and attention on global feature selection in human visual cortex, *Visual Cognition*, 23:1-2, 229-248. https://doi.org/10.1080/13506285.2015.1011252

Kang, J., Kim, H., Hwang, S. H., Han, M., Lee, S. H., & Kim, H. F. (2021). Primate ventral striatum maintains neural representations of the value of previously rewarded objects for habitual seeking. *Nature communications*, *12*(1), 2100. https://doi.org/10.1038/s41467-021-22335-5

Kim, H. F., & Hikosaka, O. (2013). Distinct basal ganglia circuits controlling behaviors guided by flexible and stable values. *Neuron*, *79*(5), 1001–1010. https://doi.org/10.1016/j.neuron.2013.06.044

Kim, H. F., & Hikosaka, O. (2015). Parallel basal ganglia circuits for voluntary and automatic behaviour to reach rewards. *Brain: a journal of neurology*, *138*(Pt 7), 1776–1800. https://doi.org/10.1093/brain/awv134

Kim, H. F., Ghazizadeh, A., & Hikosaka, O. (2015). Dopamine Neurons Encoding Long-Term Memory of Object Value for Habitual Behavior. *Cell*, *163*(5), 1165–1175. https://doi.org/10.1016/j.cell.2015.10.063

Kleiner, Mario & Brainard, D.H. & Pelli, Denis & Ingling, A. & Murray, R. & Broussard, C. (2007).
What's new in Psychtoolbox-3. *Perception*. 36. 1-16. 10.1068/v070821.

Lee, J., & Shomstein, S. (2013). The differential effects of reward on space- and object-based
attentional allocation. *The Journal of neuroscience: the official journal of the Society for
Neuroscience*, *33*(26), 10625–10633. https://doi.org/10.1523/JNEUROSCI.5575-12.2013

Liljeholm, M., Dunne, S., & O'Doherty, J. P. (2015). Differentiating neural systems mediating
the acquisition vs. expression of goal-directed and habitual behavioral control. *The European
journal of neuroscience*, *41*(10), 1358–1371. https://doi.org/10.1111/ejn.12897

Luigjes, J., Segrave, R., de Joode, N. *et al.* Efficacy of Invasive and Non-Invasive Brain
Modulation Interventions for Addiction. *Neuropsychol Rev* **29**, 116–138 (2019).
https://doi.org/10.1007/s11065-018-9393-5

MacLean, M. H., & Giesbrecht, B. (2015). Neural evidence reveals the rapid effects of reward
history on selective attention. *Brain research*, *1606*, 86–94.
https://doi.org/10.1016/j.brainres.2015.02.016

McNamee, D., Liljeholm, M., Zika, O., & O'Doherty, J. P. (2015). Characterizing the associative
content of brain structures involved in habitual and goal-directed actions in humans: a
multivariate FMRI study. *The Journal of neuroscience: the official journal of the Society for
Neuroscience*, *35*(9), 3764–3771. https://doi.org/10.1523/JNEUROSCI.4677-14.2015

Moore, T., & Zirnsak, M. (2017). Neural Mechanisms of Selective Visual Attention. *Annual
review of psychology*, *68*, 47–72. https://doi.org/10.1146/annurev-psych-122414-033400

Qi, S., Zeng, Q., Ding, C., & Li, H. (2013). Neural correlates of reward-driven attentional capture
in visual search. *Brain research*, *1532*, 32–43. https://doi.org/10.1016/j.brainres.2013.07.044

Saad ZS, Reynolds RC, Argall B, Japee S, Cox RW (2004) Suma: An interface for surface- based
intra- and inter-subject analysis with AFNI. 2004 *Second IEEE International Symposium on
Biomedical Imaging*: *Macro to Nano* (IEEE, Piscataway, NJ), pp 1510–1513.

Saproo, S., & Serences, J. T. (2010). Spatial attention improves the quality of population codes
in human visual cortex. *Journal of neurophysiology*, *104*(2), 885–895.
https://doi.org/10.1152/jn.00369.2010

Seeley W. W. (2019). The Salience Network: A Neural System for Perceiving and Responding to
Homeostatic Demands. *The Journal of neuroscience: the official journal of the Society for
Neuroscience*, *39*(50), 9878–9882. https://doi.org/10.1523/JNEUROSCI.1138-17.2019

Sjoerds, Z., de Wit, S., van den Brink, W., Robbins, T. W., Beekman, A. T., Penninx, B. W., &
Veltman, D. J. (2013). Behavioral and neuroimaging evidence for overreliance on habit learning
in alcohol-dependent patients. *Translational psychiatry*, *3*(12), e337.
https://doi.org/10.1038/tp.2013.107

Song, S., Zilverstand, A., Gui, W., Li, H. J., & Zhou, X. (2019). Effects of single-session versus
multi-session non-invasive brain stimulation on craving and consumption in individuals with
drug addiction, eating disorders or obesity: A meta-analysis. *Brain stimulation*, *12*(3), 606–618.
https://doi.org/10.1016/j.brs.2018.12.975

Tankelevitch, L., Spaak, E., Rushworth, M. F. S., & Stokes, M. G. (2020). Previously Reward-
Associated Stimuli Capture Spatial Attention in the Absence of Changes in the Corresponding
Sensory Representations as Measured with MEG. *The Journal of neuroscience: the official*
journal   of   the   Society   for   Neuroscience,   *40*(26),   5033–5050.
https://doi.org/10.1523/JNEUROSCI.1172-19.2020

Tricomi, E., Balleine, B. W., & O'Doherty, J. P. (2009). A specific role for posterior dorsolateral
striatum in human habit learning. *The European journal of neuroscience*, *29*(11), 2225–2232.
https://doi.org/10.1111/j.1460-9568.2009.06796.x

Watson, P., Pavri, Y., Le, J., Pearson, D., & Le Pelley, M. E. (2022). Attentional capture by signals
of reward persists following outcome devaluation. *Learning & memory (Cold Spring Harbor,*
N.Y.), *29*(7), 181–191. https://doi.org/10.1101/lm.053569.122

Watson, P., van Wingen, G., & de Wit, S. (2018). Conflicted between Goal-Directed and
Habitual   Control,   an   fMRI   Investigation.   *eNeuro*,   *5*(4),   ENEURO.0240-18.2018.
https://doi.org/10.1523/ENEURO.0240-18.2018

Wimmer, G. E., Li, J. K., Gorgolewski, K. J., & Poldrack, R. A. (2018). Reward Learning over Weeks
Versus Minutes Increases the Neural Representation of Value in the Human Brain. *The Journal*
of neuroscience: the official journal of the Society for Neuroscience, *38*(35), 7649–7666.
https://doi.org/10.1523/JNEUROSCI.0075-18.2018

Wunderlich, K., Dayan, P., & Dolan, R. J. (2012). Mapping value based planning and extensively
trained   choice   in   the   human   brain.   *Nature   neuroscience*,   *15*(5),   786–791.
https://doi.org/10.1038/nn.3068

Yasuda, M., Yamamoto, S., & Hikosaka, O. (2012). Robust representation of stable object values
in the oculomotor Basal Ganglia. *The Journal of neuroscience: the official journal of the Society*
for Neuroscience, *32*(47), 16917–16932. https://doi.org/10.1523/JNEUROSCI.3438-12.2012

Yin, H. H., & Knowlton, B. J. (2006). The role of the basal ganglia in habit formation. *Nature*
reviews. Neuroscience, *7*(6), 464–476. https://doi.org/10.1038/nrn1919

Yu, H. H., Chaplin, T. A., & Rosa, M. G. (2015). Representation of central and peripheral vision
in the primate cerebral cortex: Insights from studies of the marmoset brain. *Neuroscience*
research, *93*, 47–61. https://doi.org/10.1016/j.neures.2014.09.004

Zhou, K., Zhu, L., Hou, G., Chen, X., Chen, B., Yang, C., & Zhu, Y. (2021). The Contribution of
Thalamic Nuclei in Salience Processing. *Frontiers in behavioral neuroscience*, *15*, 634618.
https://doi.org/10.3389/fnbeh.2021.634618

Fig. S1 Behavioral results in force trials. (A) Performance in force trials across the three training sessions. Boxplots show the choice rate of good and bad objects in each session. The central mark within each box indicates the median, and the bottom and top of the box correspond to the 25th and 75th percentiles, respectively. The whiskers extend to the extreme values within 1.5 times the interquartile range. Outliers are plotted as dots. (B) Response time in force trials across the three training sessions. Same conventions as in (A) ($*p < 0.05$).

Fig. S2 Brain regions with significant spatial coding. (A) Brain regions with significant hemifield selectivity. Warm colors show regions more strongly activated in response to the objects displayed on the left hemifield, and cool colors show the opposite (p < 0.05, α < 0.01 cluster corrected). (B) Brain regions with significant central vs. peripheral discrimination. Warm colors show areas more strongly activated in response to the objects displayed in the center, and cool colors show the opposite (p < 0.05, α < 0.01 cluster corrected).

Fig. S3 Regions showing significant activation in response to the memorized value. Brain regions with significantly different responses to the memorized good vs. memorized bad objects in the memory session (p < 0.05, α < 0.01 cluster corrected).

Table S1. Summary of whole-brain analysis results

Contrast	Regions	Cluster size	x	y	z
Memorized good vs.	R thalamus	241	1.5	7.5	10.5
Memorized bad	L thalamus				
	R caudate				
	R ventral striatum				
	R lingual gyrus	211	-10.5	88.5	-16.5
	R cerebellum				
	R intraparietal sulcus	89	-31.5	73.5	49.5
	L intraparietal sulcus	85	28.5	70.5	52.5
	L caudate	78	10.5	4.5	16.5
	L ventral striatum				
	R inferior temporal gyrus	73	-55.5	61.5	-19.5
	R inferior occipital gyrus				
	R fusiform gyrus				
	L fusiform gyrus	67	46.5	76.5	-16.5
	L lingual gyrus				

Clusters of activity exceeding whole-brain p < 0.05, α < 0.01, cluster-corrected (cluster size = 66 voxels).

Fig. S4 Areas showing significant correlation with days-long value test performance. (A-B) Same format as Fig. 5 A-B, but for the left caudate nucleus (C-D) Same format as Fig. 5 A-B, but for the right ventral striatum (Pearson's correlation "r" and significance "p" are noted in the plots).