Genetically diverse populations hold the keys to climatic adaptation: a lesson from a cosmopolitan raptor

Hugo Corval[1,2Δ]*, Tristan Cumer[1,2Δ]*, Alexandros Topaloudis[1,2], Flavien Collart[3], Antoine Guisan[1], Alexandre Roulin[1] and Jérôme Goudet[1,2]*

* corresponding authors

Δ co-first authors – sorted alphabetically

1 Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland

2 Swiss Institute of Bioinformatics, Lausanne, Switzerland

3 Institute of Botany, University of Liège, Liège, Belgium

Contact details for corresponding author: Department of Ecology and Evolution, Biophore Building, University of Lausanne, CH-1015 Lausanne, Switzerland. Email:

hugo.corval@unil.ch; tristan.cumer@unil.ch; jerome.goudet@unil.ch

Orcid

Hugo Corval - 0009-0008-1842-7754

Tristan Cumer - 0000-0002-0276-7462

Alexandros Topaloudis - 0000-0002-0909-8695

23   Flavien Collart - 0000-0002-4342-5848

24   Antoine Guisan - 0000-0002-3998-4815

25   Alexandre Roulin - 0000-0003-1940-6927

26   Jérôme Goudet - 0000-0002-5318-7601

Abstract

Although local adaptation influences species distributions, its role in driving evolutionary resilience under climate change remains unclear. Current predictive models focus on genetic adaptation to present climates, providing limited insight into future adaptive capacity. We hypothesise that historical responses to climatic shifts can reveal future adaptive potential. Combining ecological niche modelling and genomic analyses, we investigated spatiotemporal patterns and mechanisms of local adaptation of the Western Palearctic barn owl (*Tyto alba*). Ecological modelling revealed that barn owls now occupy a broader climatic niche than during the Last Glacial Maximum. Genomic analyses indicated ongoing adaptation, with regions under selection linked to environmental factors across all populations. Our findings demonstrate that local adaptation drives evolutionary changes across populations, enabling colonisation of new habitats and shaping responses to climate change in resident populations. We demonstrate that standing genetic diversity plays a crucial role in adaptation to past, present, and future environmental shifts.

Key words

Local adaptation, Climate change, Genomics, Diversity, Chromosomal inversion

Introduction

Climatic variations affect biodiversity by impacting individual's fitness, by driving population differentiation, and ultimately by shaping species distributions[1–3]. The extent of these impacts, however, depends on the speed and intensity of climatic variations[4]: sudden or extreme shifts can lead to local or global extinction if individuals fail to survive or reproduce[5]. In case of more gradual changes, the persistence of populations and species will depend on individuals' ability to migrate or cope with their new environment[1,6,7]. Through migration, individuals track their suitable conditions in time and space to survive, inducing a shift in species distribution[1]. Individuals can also change physiologically, morphologically, or phenologically to cope with the new local conditions[7]. These changes can occur via phenotypic plasticity, the ability of individual genotypes to produce different phenotypes when exposed to various environmental conditions[8] or through genetic adaptation, favouring different genotypes better adapted to the local ecological conditions[9].

Considering the critical role of local adaptation in determining population persistence[10], understanding whether individuals possess the intrinsic capacity to adapt to new environmental conditions is critical. In recent years, genomic offset has emerged as a key metric for predicting genetic maladaptation to future climates by linking environmental factors to allele frequencies and estimating the genetic changes needed for individuals to survive to new conditions[11–13]. However, while such predictions provide valuable insights into potential risks, these predictions assume populations are adapted to current conditions and cannot evolve further, overlooking their adaptive potential. Therefore, a key challenge remains understanding whether standing genetic variation can fuel adaptation to new climate - a question our study seeks to answer.

At the end of the Last Glacial Maximum (LGM) temperatures rose and ice caps melted,

allowing species to (re-)colonise previously unsuitable lands[2,14,15]. Climatic variations since

the LGM offer an excellent opportunity to study species' adaptation. The interplay between

migration and selection gives rise to several possible scenarios for how local adaptation may

occur: In populations that migrated northward, individuals either tracked their suitable niche

or faced new conditions. In the latter case, selection may have driven local adaptation by

favouring the most suited individuals[9], but repeated migration events eventually caused

founder effects and a loss of adaptive potential along the front of colonisation[16]. In contrast,

populations that remained at the core of the distribution may also have faced a change in

conditions. These populations often harbour a higher genetic diversity[17], possibly enhancing

their ability to adapt to their changing environment. Given these three main scenarios,

where and how local adaptation happens remains elusive and exploring these questions can

enhance our understanding of the adaptive potential of populations.

The barn owl (*Tyto alba*), a nonmigratory raptor distributed all over the Western Palearctic,

faces heterogeneous climates. The species recolonised the Northern part of Europe at the

end of the LGM from two main glacial refugia located in the Iberian Peninsula and the Italian

and Balkan Peninsulas[15]. Presently, its range stretches across the Western Palearctic[15].

Despite a generally low level of genetic differentiation across its range, southern populations

host a higher genetic diversity than northern populations[15,18,19], and exhibit notable

phenotypic variation, such as a cline in plumage coloration between southern and northern

populations[18–21]. A previous study demonstrated that the colour cline cannot be explained

by purely neutral processes and argued that selection for local adaptation was or is still

acting on this phenotype[21]. The well-understood history of the barn owl in Europe since the

Last Glacial Maximum, combined with the low genetic structure at the continental level and

previous evidence of adaptive selection throughout its range, make this species an attractive

model to test and quantify the extent and location of local adaptation.

Here, we evaluated where and how heterogeneous climate induce local adaptation using the

European barn owl as a model organism. We first used ecological information and Species

Distribution Modelling to (i) quantify the climatic heterogeneity faced by the barn owl

nowadays and (ii) measure how these climatic conditions differ from those experienced

during the Last Glacial Maximum. We then looked for an association between climatic

variables and genomic variants from the entire genome of 74 owls from 9 different localities

across Europe. We combined these results with a new approach to scan the genomes for

traces of selection and identified genomic regions and genes potentially involved in the local

adaptation of the different populations. We found a strong and common signal in southern

populations. Overall, our results demonstrate how the most diverse populations, often

located at the core of the distribution, may host the adaptive potential to face climate

change, giving clues on how standing genetic variation can fuel local adaptation.

104 Results

105 Suitable conditions nowadays are more diverse than during the

106 Last Glacial Maximum

The spatial prediction obtained by the species distribution model (SDM)[22] highlights a

striking increase of the area occupied by the barn owl since the Last Glacial Maximum[23]

(LGM, ~ 20,000 years ago), similar to what was observed in trees[24], mammals[25], and birds[26].

During the LGM, suitable habitats were mostly confined to regions around the

Mediterranean — covering northern Africa as well as the Iberian, Italian, and Greek

peninsulas (Supplementary Figure 1). Today, favourable conditions extend well into

Western, Central, and Northern Europe, making most of the continent suitable for barn

owls. This notable northward and inland expansion prompted us to investigate the climatic

variables driving these changes.

To identify the climatic factors defining the climatic niche of the barn owl, we performed a

Principal Component Analysis (PCA) fitting a multidimensional climatic space both from the

LGM and modern periods. The first three principal components explain 91.50% of the

variance (Figure 1b). The first component, explaining 49.66% of the variance, is driven by

temperature-related variables - such as the Mean Temperature of the Coldest Month (bio6),

Temperature Annual Range (bio7), and Mean Temperature of the Wettest Quarter (bio8) -

which differentiate the Mediterranean region from the rest of Europe (Figure 1c). The

second component, explaining 33.6% variance, is mainly influenced by precipitation factors,

including Precipitation Seasonality (bio15), Precipitation of the Driest Quarter (bio17), and

Precipitation of the Coldest Quarter (bio19), effectively distinguishing coastal areas from the

continental interior (Figure 1c). The third component, explaining 8.25% variance, is largely

driven by the Minimum Temperature of the Coldest Month (bio6; Supplementary Figure 2

and 3).

We next explored the overlap between past and present climatic conditions in areas suitable

for barn owls and showed that owls now occupy a wider variety of climatic conditions than

during the LGM[24]. We found that only a small fraction of today's suitable habitats shares

characteristics with those from the LGM. We found that only 25.54% of modern suitable

cells exhibit LGM-like climatic conditions (Figure 1a, dark grey areas). These overlapping

regions are found in northern Africa, the northwestern Iberian Peninsula, western France,

the British Isles, and western Turkey. In contrast, 74.45% of current suitable cells are

characterized by climatic conditions that did not exist during the LGM (Figure 1a, RGB-

coloured areas). This significant shift suggests that most of today's barn owl habitats are

defined by new climatic conditions, potentially driving local adaptation to these emerging

environments.

Species-wide sampling reveals that a substantial portion of the

genome is under climatic selection

To test the hypothesis that local climates may have driven the local adaptation of the barn

owl in the Western Palearctic, we have sampled and sequenced 79 individuals from nine

populations across the climatic and European geographical distribution of the species

(Supplementary Figure 4 and 5, Supplementary Table 1) and identified 12,309,943 Single

Nucleotide Polymorphisms (SNPs). The overall differentiation was low (overall F_{ST} = 0.034)

and in line with previous estimates[15]. The first axis of the genomic PCA (explaining 3.88% of

the total variance) contrasted individuals from the Levant populations (Israel—IS) to all other

individuals (Figure 1d). The second axis of the PCA (explaining 2.26% of the variance)

opposed individuals from the most diverse population (from the Iberic peninsula—PT) to all

others (Figure 1d, Table 1). Overall, southern populations (PT, IT, GR, AE, IS) were more

genetically diverse than northern populations (CH, FR, DK, SB) (Table 1). Next, we employed

two complementary approaches - a genome scan for selection and a Genotype-Environment

Association (GEA) analysis - to pinpoint genomic regions potentially influenced by these

climatic conditions.

In our first approach, we scanned the genomes for signatures of selection using population-

specific FST[27]. This metric, which accounts for population structure, enables the detection of

regions of the genome where individuals have higher genetic similarity than what is

observed along the rest of the genome - a pattern possibly resulting from selective

events[27,28]. Out of 52,429 windows of 100 kbp (with a 20 kbp slide), we identified 10,607

outlier windows: 7,034 were unique to a single population, and 3,573 were shared between

at least two populations. On average, each population had about 1,756 outlier windows

(Table S2), with Israel and France at the extremes (989 and 2,087 outliers, respectively).

To ensure that selection signals were caused by climatic conditions, we directly linked

genetic variation to temperature and precipitation conditions with a Genotype-Environment

Association (GEA). We conducted a Redundancy Analysis (RDA, Supplementary Figure 6)[29]

that we integrated into genomic windows with the Weighted-Z Analysis (WZA) method[30].

We detected 2,181 outlier windows significantly linked to temperature and precipitation

variables (Supplementary Figure 7) and overlaid it with the population-specific FST scans.

This combined analysis yielded a refined list of 1,246 outlier windows (displayed in dark blue

in Figure 2a; Table S2). 59.71% were unique to a single population while the rest was shared

between at least two populations. By merging successive outlier windows, we delineated

270 genomic regions, many forming distinct adaptive peaks along the genome. Particularly

striking associations were observed on Super-Scaffold 14 and 45 (Figure 2b). Additionally, we

detected a high density of outlier windows in the first half of Super-Scaffold 22 (Figure 2b-d);

this signal, shared by owls from France, Italy, and Portugal, corresponded with a marked

increase of population-specific FST in these genomic regions (Supplementary Figure 8),

indicating that individuals within each of these populations are significantly more similar to

one another than to the rest of the samples.

Metabolic pathways associated with climate

To explore the functional implications of selection and climatic adaptation, we extracted a

total of 550 genes from the 270 genomic regions that showed signatures of selection and

variants associated with climatic variables. Among them, 324 genes were unique to a single

population (Table S3). Within the 550 extracted genes, we detected significant enrichments

in pathways related to cellular physiology (details of the Gene Ontology (GO) terms are

provided in Table S4). We observed the same results when we used the population-specific

list of genes from Greece, Israel and Serbia (Table S5). For the other populations, the GO

enrichment analysis did not yield any significant results. Overall, the functional grouping of

the 550 genes included functions such as immunity, locomotion, and anatomy in all the

populations, with different genes in each population (Table S6).

Climatic differences drive differential selection between Southern and Northern populations

Our analysis of the region with the highest signal of association with climate reveals that

differential selection has led to marked genetic differentiation between populations at

opposite ends of the temperature gradient. To explore the signal located on Super-Scaffold

45, we first examined climatic association values within this genomic region (Figure 3). Out

of the 8 outlier windows showing the highest association with climate (red dots in Figure 3a-

b), all were outliers in the population-specific FST scan for the Danish population, and two of

them were also outliers in the Portuguese population, two populations at the opposite ends

of the temperature gradient (first axis of Supplementary Figure 4). We assessed the extent of

genetic similarity in this genomic region between the two populations by computing the

population-pair FST, an estimate of the standardised mean kinship of individuals[31]. With this

statistic, we detected a substantial reduction of the genetic similarity between the

individuals from these two populations within the region harbouring the highest association

with climatic conditions (red rectangle on Figure 3c). Consistently, we observed an increase

of genetic dissimilarity by using a pairwise FST computed on an SNP basis between the two

populations (Figure 3d). We examined the haplotypes within the highest section of the peak

which showed a clear distinction between the Danish and Portuguese populations (Figure

3e).

A putative inversion linked to climatic adaptation in past refugia

We identified a long consecutive signal of climatic association shared between populations

from France, Italy, and Portugal in the first half of Super-Scaffold 22 (Figure 2b-c; Figure 4a).

We used the population-pair FST to assess whether the same genomic variants were shared

in the three populations. We detected a higher genetic similarity between the three pairs of

populations (FR-IT, FR-PT, and IT-PT) within a 14 Mb region at the beginning of the Scaffold

than along the rest of the genome (Figure 4b). This coincides with a climatic convergence in

former glacial refugia, where shifting conditions may have contributed to the shared

adaptive signals (Supplementary Figure 9, 10, 11 and 12).

To further investigate the genomic features of this region, we applied a PCA to the first 14

Mb of the scaffold, encompassing 120,953 SNPs, using data from all 74 individuals (Figure 4c

illustrates the first two axes, explaining respectively 26.91 and 2.73% of the variation).

Individuals were divided along the first axis into three distinct clusters, corresponding to the

expected genomic pattern in case of a chromosomal inversion[32]. The leftmost cluster

included more individuals than the one on the rightmost part of the x-axis. In comparison,

the PCA conducted on the whole genome only separates the Portuguese samples from the

rest along the x-axis and the Israel individuals from the rest along the y-axis (Figure 1d).

227 Discussion

Climatic conditions and their past fluctuations are known to have shaped today's distribution

and genetic diversity of many species[1–3]. While genetic local adaptation is widely recognized

as a key mechanism driving species persistence[6], its role in shaping populations' capacity to

respond to future climatic shifts remains unclear. In this study, we integrated niche-based

species distribution modelling (SDM[22]) with genome-wide analyses to investigate how

climate-driven changes since the Last Glacial Maximum (LGM) have shaped the genetic

composition and adaptive potential of the Western Palearctic barn owl. First, we showed

that suitable conditions nowadays are more climatically diverse than the ones from the

LGM[24–26], exposing the individuals to heterogeneous and new selective pressures. Then, we

identified genomic regions harbouring strong signals of selection and carrying variants

associated with climatic variables. We observed signals of genomic adaptation in all the

population we sampled across the species' range, with a particularly strong signal shared

among southern populations. Overall, our findings challenge the hypothesis that local

adaptation primarily occurs at recolonising margins[33,34], showing instead that genetically

diverse populations such as past glacial refugia are a key source for adaptation[35]. We

highlight that refugial populations, by maintaining standing genetic variation, serve as crucial

reservoirs of adaptive potential, enhancing species' ability to respond to future climatic

changes[35–37].

Species distribution modelling of the European barn owl based on its realized climatic niche

suggests that the latter is broader nowadays than it was during the Last Glacial Maximum

(LGM). We used the Maximum Entropy method[38] for fitting the SDM and project it onto past

climatic conditions to compare the extent of suitable habitat under current versus historical

climates. Our results indicate that some contemporary climatic conditions along the Atlantic

coast, up to the UK, were already present 20,000 years ago. This supports the hypothesis

that, as temperatures rose at the end of the LGM, favourable conditions shifted toward

northwestern Europe, facilitating barn owl range expansion[39]. However, we also identified

newly emerged suitable climatic conditions that were absent during the LGM in recolonised

Central and Eastern Europe as well as in the glacial refugia where conditions changed for

resident populations. All these results suggest that the realised climatic niche of the barn owl

was contracted during the Last Glacial Maximum and extended with the subsequent

warming. We advance that genetic adaptation played a significant role in this process and

further explored this hypothesis.

By combining population-specific FST scans[27,28] to identify genomic regions under selection

with genome-environment association approaches, we found genomic evidence of local

adaptation to humidity and temperature in all populations. Genes among selected regions

showed significant GO term enrichment in several populations, but we found no clear link to

climate adaptation. However, the functional clustering revealed a concentration of genes

related to immunity, with GO terms such as "immune system process" and "immune

response" (Table S6). This finding aligns with previous research showing that pathogens and

infectious diseases exert strong selective pressures on both humans and birds[40,41].

Moreover, temperature and rainfall significantly influence pathogen community

composition[42,43]. Supporting this connection between climate and immune genes, O'Connor

et al. (2020) demonstrated that MHC-I genes in birds vary in diversity according to

humidity[44]. GO terms related to anatomy and growth were also found across population,

encompassing several dozens of genes linked to "anatomical structure morphogenesis",

"regulation of the developmental process" and "growth" (Table S6). This result was

consistent with Bergmann's rule which predicts that body sizes of warm-blooded vertebrates

negatively correlate with temperature, leading to smaller body sizes in warmer climates[45,46].

Overall, our results suggest an implication of these genes in local adaptation to various

climates, but further work is needed to identify and confirm their precise role.

A closer look at the regions of the genome with many variants linked to the environment and

with strong traces of selection revealed a wide range of mechanisms shaping the barn owl's

local adaptation, from opposite directional selection to recurrent evolution.

The region located on the Super-Scaffold 45 showed a clear signal of association with

climatic variables. In this region, population-specific FST highlights the high differentiation

between the Portuguese and Danish populations, which lie at the opposite ends of the

temperature gradient (Supplementary Figure 4). This, along with their distinct haplotypic

structure (Figure 3), suggests that different alleles or haplotypes are under selection in these

populations[47]. A closer look at the genes present in the region did not reveal any genes that

can easily be linked to temperature adaptation and further work should confirm whether

this region plays a role in thermal adaptation.

Our results also revealed a strong and consistent genomic signal of selection linked to

climate across all individuals from three populations, one in a region with a climate similar to

that of the LGM (France) and two in past glacial refugia (the Iberic and Italian peninsulas).

The size of the region involved; the pronounced pattern of genetic differentiation compared

to the rest of the populations as well as the cluster patterns found by our local PCA strongly

suggest the presence of an inversion in this region[32]. Supported by the growing body of

literature linking inversion and adaptation[48–50], we suggest that this inversion is adaptive to

climatic conditions. Briefly, considering the genomic disruption that breakpoints can cause,

as well as the fact that nearly suppressing recombination will prevent purging of deleterious

alleles in the inverted haplotype, it is unlikely for a newly appeared inversion to persist and

spread in a population[51]. However, by bringing adaptive alleles together and averting

maladapted gene flow by blocking recombination, an inversion can promote local adaptation

and thus be favoured by selection[50,52,53].

The history of this inversion remains to be explored. The pattern of selection is shared

between the two peninsulas that hosted the species during the LGM. Considering the

isolation of the two populations at the time and the reduced connectivity nowadays (see

Figure 4 in Cumer et al., 2021 for details)[27] we suppose that this inversion predates the LGM,

and that independent and recurrent selection drove the increase of frequency of this

haplotype independently in these two populations. Further work should formally test this

hypothesis.

A key question is the origin of adaptive alleles and given the pattern of genomic traces of

local adaptation we observed, we propose that standing variation played a crucial role in this

process[54,55]. One possibility for local adaptation to occur is *de novo* mutations bringing new

favourable alleles in the different populations[54,56]. Considering the low mutation rate[57,58] and

the even lower probability of a mutation being both advantageous and occurring in the right

environmental context[59], this mechanism is unlikely to fully explain the extent of genomic

regions involved in local adaptation (Figure 2). A more parsimonious explanation relies on

adaptation to pre-existing genetic variation[55]. In this scenario, allele sorting from standing

genetic variation would have driven adaptation[60]. This occurs when previously neutral or

mildly deleterious alleles become advantageous following shifts in selective pressures, such

as habitat colonisation or climate change[60]. This scenario implied that within glacial refugia,

a pool of segregating alleles was maintained in sufficiently large populations[61]. These alleles

were then sorted through space by local adaptation, with different environments (biotic and

abiotic) acting as a filter on deleterious and/or less advantageous alleles. This whole process

has been reported in species as diverse as insects (*Drosophila melanogaster*[62]), birds

(Darwin's finches[63]) mammals (humans[59,64]) and fish (sticklebacks[65]). We did not explicitly

test whether local adaptation in barn owls resulted from de novo mutations, allele sorting

from standing variation, or a combination of both. Therefore, additional work, focusing on

the date of mutations[66] as well as the timing of selection[67], should confirm our results.

However, the pattern of local adaptation that we identified in populations from past glacial

refugia already provides insight into how standing variation fuels adaptation during climatic

fluctuations.

The climatic variation since the Last Glacial Maximum provides a good framework to gain

knowledge on where and how populations adapt to climate change. A wide body of

literature explored many aspects of what happens in populations that expanded into newly

available habitat, from dispersal limitation[68,69] to the genetic load accumulated during an

expansion[70–74], and how these mechanisms interact with local adaptation[75–78]. However, the

fate of refugial populations remains, to our knowledge, understudied. In this work, we first

demonstrate that the climatic conditions experienced by these populations have changed

over time, creating ongoing pressures for adaptation. Our findings suggest that refugial

populations are not only well adapted to their current environments but are also continuing

to adapt. Based on our results, we propose that adaptation in refugial populations is a

recurrent and dynamic process. As the climate continues to shift, the species' optimal niche

moves northward, and southern populations will be among the first to experience novel

climatic conditions. These populations often harbour the highest levels of genetic diversity[17],

which likely confers them the greatest adaptive potential in the face of climate change. This

underscores the importance of conserving these genetically diverse populations, whose

adaptive capacity may be critical for the long-term resilience of the species. While we

acknowledge that the pace of current climate change is far more rapid compared to shifts

since the Last Glacial Maximum, our study las the groundwork for a deeper of the genomic

foundations of adaptative potential. Future research will be essential to unravel the

mechanisms that enable species to persist and thrive in rapidly changing environments.

Material and methods

Ecological modelling

Species distribution modelling

We first conducted species distribution modelling (SDM[22]) to identify suitable areas for the

species during the Last Glacial Maximum and nowadays. We fitted SDM using the Maximum

Entropy modelling software (MaxEnt[38,79], v.3.4.3), a presence-only based procedure in a

similar approach as the one described in Cumer et al. (2022)[15].

First, we extracted 19 bioclimatic variables at a 5 arc-minute (~9.3 km at the equator)

resolution from the WorldClim 1.4 database[80] using the rbioclim R package[81]. To avoid

redundancy between variables, we removed variables with a correlation equal to or higher

than 0.8[82], leading to a set of 7 uncorrelated climatic variables: Mean Diurnal Range (Bio2),

Min Temperature of Coldest Month (Bio6), Temperature Annual Range (Bio7), Mean

Temperature of Wettest Quarter (Bio8), Precipitation Seasonality (Bio15), Precipitation of

Driest Quarter (Bio17), and Precipitation of Coldest Quarter (Bio19).

We performed the analysis using the dismo R package[83] v.1.3-14 on R v.4.3.2. To determine

which combination of parameters optimised the model without over-complexifying it[84], we

built models with linear, quadratic, and hinge features, using a range of regularisation

multipliers from 1 to 5 (as recommended in Warren & Seifert, 2011[85]). A quadratic feature

with a regularisation multiplier of 1 yielded the lowest AIC and was chosen for further

modelling. We ran a total of 100 quadratic MaxEnt models (with a regularisation multiplier =

1), omitting 25% of the data during training to test the model and using 10'000 background

points randomly sampled across the study area. For each model, we randomly sampled

1,000 presence points from the IUCN distribution map[86]. We evaluated the predictive

performances of the models by assessing the area under the curve (AUC) of the receiver

operating characteristic (ROC) plot of the test data[87]. All models had an AUC between 0.756

and 0.807 (see Supplementary Figure 13), thus classified as fair to good according to Li et al.

(2020)[88].

We projected the 100 models to the present climatic conditions and to the climatic

conditions from the Last Glacial Maximum (about 20'000 years ago), also extracted from the

WorldClim 1.4 database (scenario CCSM from PMIP2[80]) at 5 arc-minute resolution. We used

the "maximum training sensitivity plus specificity" (MaxSSS) threshold, as recommended for

presence-only data[89], to transform the projected output from the models into binary

suitability maps (0 unsuitable, 1 suitable). Finally, we averaged the values among replicates

and retained cells as suitable only if they were so in at least 90% of the models. To avoid

model extrapolation when projecting the models in the past, we used the Multivariate

Environmental Similarity Surface (MESS) approach[90] to identify and discard areas from the

past with climatic conditions absent from those in the calibration data.

Identification of newly suitable conditions absent from the LGM

Then, we determined whether currently suitable areas display climatic conditions different

from those found during the LGM. To do so, we extracted values of the 7 climatic variables

used in the SDM (see Species distribution modelling section for details) at every continental

cell of the studied area for both past and present and performed a Principal Component

Analysis (PCA) to represent the environmental space. We retained the first three principal

components that explained up to 91.50% of the climatic variance. In this environmental

space, we generated a "multidimensional climatic space", representing the entire range of

climatic conditions suitable for the barn owl, by combining past and present suitable pixels

within this PCA space. We then characterised the past suitable conditions for the species by

creating a polygon around the LGM suitable conditions, using the alphashape3d R package

($\alpha = 0.5$, keeping all the data[91]). Next, we classified the suitable conditions at current time as

located inside or outside the polygon, thus respectively present or absent from the range of

LGM suitable conditions.

Estimation of the shift of climate within suitable areas

To quantify shifts in climatic conditions between LGM and now, we identified cells that were

suitable both during the LGM and present time (Supplementary Figure 9) and computed the

Euclidean distance between their past and present positions in the multidimensional climatic

space. We then rescaled the distance values from 0 to 1 (Supplementary Figure 10).

Because we discovered a strong signal of selection shared between the two glacial refugia

populations (see Results section), we investigated how the climate has changed since the

LGM in the two peninsulas to assess whether the climate converged or diverged in these two

locations. We extracted values of the 7 climatic variables used in the ecological modelling for

the LGM and today (see Species distribution modelling section) at the GPS coordinates for

the Italian and Portuguese samples. We projected these conditions inside the

multidimensional climatic space (Supplementary Figure 11). Then, we computed the

pairwise Euclidean distances between all sampling localities of the two peninsulas during the

LGM and the present time (8 (IT) x 8 (PT) pairwise distances at both time points). This way,

we quantified the climatic difference between the Italian and Portuguese sampling localities

at the two different time points (Supplementary Figure 12).

Individual barn owl sampling

Biological samples

420 This study took advantage of the datasets of European barn owls previously published by

421 Cumer et al. (2022), Machado, Cumer, et al. (2021), and Machado, Topaloudis, et al.

422 (2021)[15,27,92]. We retrieved the whole genome sequences from the Sequence Read Archive

423 (SRA - Bioprojects PRJNA700797, PRJNA727915, and PRJNA727977, Table S1).

Genetic data preparation

We performed the read mapping, variant discovery, and variant filtering following Cumer et

al. (2022)[15]. In brief, we mapped raw reads to the reference barn owl genome (GenBank

accession GCA_018691265.1[39]) with BWA-MEM v.0.7.15[93]. We performed Base quality score

recalibration (BQSR) in GATK v.4.1.3 using high-confidence calls described in Cumer et al.

(2022)[15]. We called variants with GATK's HaplotypeCaller and GenotypeGVCF v.4.1.3 from

the recalibrated BAM files. We filtered genotype calls using a hard-filtering approach as

suggested for nonmodel organisms, using GATK and VCFtools[94]. Details of technical filtration

can be retrieved from Cumer et al. (2022)[15].

To prevent some alleles from being over-represented in the dataset (relatedness statistic

based on the method of Manichaikul et al., 2010[95], implemented in VCFtools v0.1.14[94]), we

identified pairs of individuals with a relatedness higher than 0.1. We removed one of the two

individuals for each identified pair, leading to a dataset of 74 individuals (Table S1).

We then excluded genomic regions with uncertain SNP calling by removing regions of the

genome where our ability to confidently map reads is limited (i.e., a "mappability" mask). To

achieve   this,   we   followed   the   procedure   documented   at

lh3lh3.users.sourceforge.net/snpable.shtml. In summary, we divided the reference genome

into reads of 150 base pairs (bp) with a sliding window of 1 bp. These artificial reads were

then mapped back to the reference using BWA-MEM v.0.7.17. Regions of the sequence

where less than 90% of the reads mapped perfectly and uniquely were discarded by

excluding variants using a bed file in VCFtools v0.1.14. We retained only bi-allelic SNPs,

removed loci with more than 5% missing data, and excluded from the analysis 15 scaffolds

(out of 60) showing less than 1000 SNPs and one extra sexual scaffold.

At the end of the filtration process, we retained a set of 12,309,943 SNPs from 39 scaffolds,

genotyped in 74 European barn owls (10 individuals from the Aegean Islands (AE), 10 from

Denmark (DK), 4 from France (FR), 9 from Greece (GR), 9 from Italy (IT), 10 from Israel (IS), 9

from Portugal (PT), 5 from Serbia (SB), and 9 from Switzerland (CH)).

To explore the genome-wide variations of the Western palearctic populations and check its

concordance with previous results, we conducted a PCA with the SNPRelate R package[96]

using this entire set of SNPs and individuals. Additionally, we assessed genetic diversity

among the 9 sampled populations following the procedure of Cumer et al., (2022)[15]. Briefly,

we identified the number of polymorphic sites, private alleles, rare alleles and the whole

genome population-specific F_{ST} for each population independently. To account for

differences in sample sizes (ranging from 4 to 10), we randomly sampled 5 individuals from

each population - except for FR and SB - and calculated these diversity estimates on the

resulting subsets. This resampling process was repeated 10 times, and we reported the

mean and standard deviation of the diversity estimates.

Phasing process and evaluation

We performed the phasing and imputation of the individuals' genotypes in two steps. First,

we conducted a read-based phasing of each individual using WhatsHap v1.0[97]. During this

step, we reconstructed haplotypes based on the mapped sequencing reads covering multiple

variants. Between the two filtering steps, we applied a Minor Allele Frequency (MAF) filter to

ensure it was higher than 5% using VCFtools v.0.1.14[94], thereby removing rare alleles from

the dataset that could influence the second round of phasing, resulting in a dataset of

4'689'284 SNPs. Then, we conducted the complementary round of phasing with ShapeIt4

v4.1.3[98] with default parameters. The latter uses a statistical approach to infer individuals'

haplotypes based on the population genotypes[99] and incorporates the phase information

from the read based phasing.

To evaluate phasing performance, we calculated the switch error rate (SER) of the phasing

generated by ShapeIt4 for each individual[100]. For each individual, we conducted a statistical

phasing using ShapeIt without considering the read-based phasing from WhatsHap for the

focal individual. Subsequently, we compared this phasing to the "true" local phasing,

inferred from the read-based approach (WhatsHap). We estimated the switch error rate

between both sets of phasing using the switchError code (available at

https://github.com/SPG-group/switchError). Among the 74 phased individuals, the mean

error rate was 2.23 * 10^{-4} and none exceeded 0.7% (Supplementary Figure 14).

Detection of traces of selection in each population

We computed a single summary statistic to identify genomic regions potentially under

selection: population-specific FST, using the hierfstat R package[101]. The statistic was

calculated across the genome in overlapping windows of 100 kbp with 20 kbp steps for each

independent population. Only windows containing at least 250 SNPs, corresponding to two

standard deviations below the mean, were included in the analysis.

To identify outlier windows (regions with extreme values of positive population-specific FST),

we first transformed the statistic into Z-scores by subtracting the population mean from

each estimate and dividing it by the standard deviation for each population independently.

We then combined Z-scores from all populations and considered a window as an outlier in

each population if its Z-score for population-specific FST was equal to or higher than 2

standard deviations from the mean of the merged Z-scores. This approach allowed us to

focus on regions exhibiting an excess similarity in the population compared to the rest of the

genome (high population-specific FST). For further details about this method, refer to Cumer

et al. (2022)[27]. We computed the statistic on the set of 12,309,943 SNPs, unfiltered for Minor

Allele Frequency (MAF). To ensure that rare alleles did not influence population-specific FST,

we also calculated this statistic on the filtered variants from MAF (set of 12'309'943 SNPs).

The high consistency between the two estimates (with and without MAF filtering), as

depicted in Supplementary Figure 15, supported our decision to retain the statistic

computed on the unfiltered variants set (12,309,943 SNPs).

Genotype-Environment Association

Redundancy Analysis

We independently conducted a Genotype-Environment Association (GEA) analysis to assess

the relationship between the genotypes of the barn owl and their surrounding environment

for all populations simultaneously.

Based on the GPS coordinates of the 74 samples (Table S1), we extracted values for the

same 7 climatic variables as those used in the species distribution modelling (Mean Diurnal

Range (Bio2), Min Temperature of Coldest Month (Bio6), Temperature Annual Range (Bio7),

Mean Temperature of Wettest Quarter (Bio8), Precipitation Seasonality (Bio15),

Precipitation of Driest Quarter (Bio17), and Precipitation of Coldest Quarter (Bio19); see

Species distribution modelling section for details). We conducted a Principal Component

Analysis (PCA) on these climatic data to assess the level of climatic dissimilarity experienced

by the barn owls sampled in this study nowadays. We retained the first three principal

components, explaining 94.97% of the climatic variance.

We then associated variants with genomic information using Redundancy Analysis (RDA)

with the vegan R package[29,102]. This method relies on a multiple linear regression of the

observed genotypes on a set of abiotic or biotic predictors. The expected genotypes based

on the model (also called fitted values) are then extracted and used as input for a PCA called

RDA space. The projection of the principal axes and components in this RDA space allows the

detection of the SNPs that contribute the most to the RDA axis and whose allelic frequency

might be putatively driven by the explanatory variables[29]. We used the imputed genotype

matrix (phased set of 4'689'284 SNPs) as the response matrix and the bioclimatic variables

extracted at each sampling locality for the multiple linear regression.

To evaluate the significance of the relationship between genotypes and climatic variables,

we performed a permutation test[103]. In brief, we computed a test statistic (F-statistic) from

the regression using the true data. Afterwards, we carried out 999 additional regressions on

permuted rows of the response data (i.e., the genotype matrix), allowing us to establish the

empirical null distribution of the statistics, to which we compared the observed statistic[103].

From this test, we found a significant relationship between the environmental variables and

genetic components (Table S7). To select the number of RDA axes to retain, we also

performed a permutation test for each axis (n = 100) by following the procedure given by

Borcard et al. (2011)[103]. Applying this method, we kept the first five RDA axes, explaining

80.41% of the constrained variance (Table S8). To detect loci that strongly contribute to the

individual's discrimination in the RDA space (outlier loci), we followed the procedure

described in Capblancq et al. (2018)[104]: we computed Mahalanobis distances between each

locus and the centre of the RDA space using the previously retained five axes. P-values were

adjusted for the false discovery rate (FDR) by computing q-values using the qvalue R

package[105]. We considered a SNP as an outlier if its q-value was less than 0.1, following

Capblancq et al. (2018)[104].

Weighted-Z analysis

To improve the robustness of our GEA approach, we decided to transform the SNPs p-values

from the Redundancy Analysis into window-based statistics through the Weighted-Z analysis

(WZA) proposed by Booker et al. (2021)[30]. This method takes as input individual p-values

from any SNP-based GEA approach and calculates a weighted-Z statistic for a given genomic

region. To do so, it transforms the p-values of the focal window into z-scores and computes

the weighted-Z statistic using the equation provided by Booker et al. (2021)[30], which

considers the variation in the number of SNPs among windows along the genome.

We computed the weighted-Z statistics on the same windows as population-specific FST.

Since WZA does not support overlapping windows, we split the window set based into five

sets of non-overlapping windows. We ran 5 separate weighted-Z analyses, one with each

input file, and merged the outputs to obtain the final one. We considered a window as an

outlier when the -log10 of its p-value was equal to or higher than 2 standard deviations from

the mean of all windows (equivalent to a p-value of 0.03 assuming a normal distribution).

Genomic signal of local adaptation to climate

Concordance between the genome scans and landscape genomics

Because the study aimed to detect traces of selection linked with local climatic conditions,

we considered the final list of outlier windows as the overlap between the outlier set from

the genome scans (see Detection of traces of selection in each population section) and the

one from the Genotype-Environment Association analysis (see Weighted-Z Analysis section).

Based on the annotated barn owl genome (GenBank Assembly Accession:

GCA_018691265.1), we considered genes that partially or fully overlapped the final list of

outlier windows as potentially involved in local adaptation to climate and extracted a list of

genes for each population.

Gene Ontology Enrichment

We conducted Gene Ontology Enrichment (GOE) analyses using ShinyGo v.0.8[106] to assess

which biological pathways the genes located in the final list of outlier windows could be

involved in and whether we could link some to local adaptation to abiotic conditions. We

performed GOEs for each set of genes from each population separately and one additional

GOE using all genes detected in at least one population. As a baseline set of genes against

which we compared the observed enrichment signature, also called the background list, we

used all the genes annotated in the barn owl genome that could have been detected with

our 52'429 non-overlapping windows from the genome scans or WZA. For each analysis, we

used the default pathway databases, namely KEGG (Kyoto Encyclopedia of Genes and

Genomes), as well as the GO Biological Process database.

Exploration of signals specific to some populations

Population-pair FST on Super-Scaffold 45

We identified a strong environmental association in Denmark and Portugal, two populations

at the opposite of our environmental gradient, with strong population-specific FST in this

region in each of them. To know whether populations have the same genetic variants or

opposite ones, we assessed the level of genetic (dis)similarity between pairs of populations

on this part of the genome. To do so, we computed a population-pair FST, described in detail

in Goudet & Weir (2023)[31], using the hierfstat R package[101]. In brief, for every pair of

populations, we calculated the average kinship among pairs of individuals and standardised

it by the average kinship between all populations using the same dataset and windows

employed for population-specific FST analysis. A schematic example is provided in

Supplementary Figure 16 for the Denmark - Portugal population pair.

Pairwise FST at the SNP level

To calculate a pairwise FST at the SNP level[31] and confirm the signal obtained through

population-pair FST, we used the *fs.dosage* function of the hierfstat R package[101]. We

computed this pairwise FST on the entire Super-Scaffold 45 between Denmark and Portugal

using the same dataset as for population-specific FST or population-pair FST.

Local PCA on Super-Scaffold 22

As we detected a strong signal of selection related to climate in the first half of Super-

Scaffold 22, we decided to investigate the genetic architecture of this region. To do so, we

conducted a PCA on the first 14 Mb of the scaffold (120,953 SNPs) using all individuals, using

the SNPRelate R package[96].

Acknowledgements

We thank Olivier Delaneau for his advice on the phasing of the individuals and its evaluation.

We thank Marc Robinson Rechavi for his advice on the Gene Ontology Enrichment analysis.

This work was supported by the Swiss National Science Foundation with grants 31003A_179358 to JG.

Data and code accessibility

The data underlying this article are available in the GenBank Sequence Read Archive Database at https://www.ncbi.nlm.nih.gov/sra, and can be accessed within Bioproject PRJNA700797, Bioproject PRJNA727915 and Bioproject PRJNA727977. Code used in this article is available at https://github.com/hugocrvl/TytoAlbaLocalAdaptation. Additional data can be provided by the corresponding authors.

Authors contribution

HC, TC, and JG designed this study; TC called the variants; AT produced the mappability mask; HC conducted the analyses with the help of TC and suggestions of all authors; FC and AG advised on the SDM; HC and TC led the writing of the manuscript with inputs from all authors.

References

1. Huntley, B. & Webb, T. Migration: Species' Response to Climatic Variations Caused by Changes in the Earth's Orbit. *J. Biogeogr.* **16**, 5 (1989).

2. Hewitt, G. The genetic legacy of the Quaternary ice ages. *Nature* **405**, 907–913 (2000).

3. Davis, M. B. & Shaw, R. G. Range Shifts and Adaptive Responses to Quaternary Climate Change. *Science* **292**, 673–679 (2001).

4. Bernatchez, L., Ferchaud, A.-L., Berger, C. S., Venney, C. J. & Xuereb, A. Genomics for monitoring and understanding species responses to global climate change. *Nat. Rev. Genet.* (2023) doi:10.1038/s41576-023-00657-y.

5. Jackson, S. T. & Overpeck, J. T. Responses of plant populations and communities to environmental changes of the late Quaternary. *Paleobiology* **26**, 194–220 (2000).

6. Lafontaine, G., Napier, J. D., Petit, R. J. & Hu, F. S. Invoking adaptation to decipher the genetic legacy of past climate change. *Ecology* **99**, 1530–1546 (2018).

7. Ackerly, D. D. Community Assembly, Niche Conservatism, and Adaptive Evolution in Changing Environments. *Int. J. Plant Sci.* **164**, S165–S184 (2003).

8. Murren, C. J. *et al.* Constraints on the evolution of phenotypic plasticity: limits and costs of phenotype and plasticity. *Heredity* **115**, 293–301 (2015).

9. Kawecki, T. J. & Ebert, D. Conceptual issues in local adaptation. *Ecol. Lett.* **7**, 1225–1241 (2004).

10. DeMarche, M. L., Doak, D. F. & Morris, W. F. Incorporating local adaptation into forecasts of species' distribution and abundance under climate change. *Glob. Change Biol.* **25**, 775–793 (2019).

11. Sang, Y. *et al.* Genomic insights into local adaptation and future climate-induced vulnerability of a keystone forest tree in East Asia. *Nat. Commun.* **13**, 6541 (2022).

12. Gain, C. *et al.* A Quantitative Theory for Genomic Offset Statistics. *Mol. Biol. Evol.* **40**, msad140 (2023).

13. Marková, S. *et al.* Local adaptation and future climate vulnerability in a wild rodent. *Nat. Commun.* **14**, 7840 (2023).

14. Hortal, J. *et al.* Ice age climate, evolutionary constraints and diversity patterns of European dung beetles: Ice age determines European scarab diversity. *Ecol. Lett.* **14**, 741–748 (2011).

15. Cumer, T. *et al.* Landscape and Climatic Variations Shaped Secondary Contacts amid Barn Owls of the Western Palearctic. *Mol. Biol. Evol.* **39**, msab343 (2022).

16. Slatkin, M. & Excoffier, L. Serial Founder Effects During Range Expansion: A Spatial
Analog of Genetic Drift. *Genetics* **191**, 171–181 (2012).

17. Svenning, J.-C., Eiserhardt, W. L., Normand, S., Ordonez, A. & Sandel, B. The Influence
of Paleoclimate on Present-Day Patterns in Biodiversity and Ecosystems. *Annu. Rev.*
Ecol. Evol. Syst. **46**, 551–572 (2015).

18. Antoniazza, S., Burri, R., Fumagalli, L., Goudet, J. & Roulin, A. Local adaptation
maintains clinal variation in melanin-based coloration of European barn owls (*Tyto*
alba). *Evolution* (2010) doi:10.1111/j.1558-5646.2010.00969.x.

19. Burri, R. *et al.* The genetic basis of color-related local adaptation in a ring-like
colonization around the Mediterranean. *Evolution* **70**, 140–153 (2016).

20. Roulin, A. & Ducrest, A.-L. Genetics of colouration in birds. *Semin. Cell Dev. Biol.* **24**,
594–608 (2013).

21. Antoniazza, S. *et al.* Natural selection in a postglacial range expansion: the case of the
colour cline in the European barn owl. *Mol. Ecol.* **23**, 5508–5523 (2014).

22. Guisan, A., Thuiller, W. & Zimmermann, N. E. *Habitat Suitability and Distribution*
Models: With Applications in R. (Cambridge university press, Cambridge, United
Kingdom, 2017).

23. Clark, P. U. *et al.* The Last Glacial Maximum. *Science* **325**, 710–714 (2009).

24. Maiorano, L. *et al.* Building the niche through time: using 13,000 years of data to
predict the effects of climate change on three tree species in Europe. *Glob. Ecol.*
Biogeogr. **22**, 302–317 (2013).

25. Leonardi, M., Boschin, F., Boscato, P. & Manica, A. Following the niche: the
differential impact of the last glacial maximum on four European ungulates.
Commun. Biol. **5**, 1038 (2022).

26. Carrera, L., Pavia, M. & Varela, S. Birds adapted to cold conditions show greater
changes in range size related to past climatic oscillations than temperate birds. *Sci.*
Rep. **12**, 10813 (2022).

27. Cumer, T. *et al.* Genomic basis of insularity and ecological divergence in barn owls
(Tyto alba) of the Canary Islands. *Heredity* **129**, 281–294 (2022).

28. Weir, B. S. & Goudet, J. A Unified Characterization of Population Structure and
Relatedness. *Genetics* **206**, 2085–2103 (2017).

29. Capblancq, T. & Forester, B. R. Redundancy analysis: A Swiss Army Knife for
landscape genomics. *Methods Ecol. Evol.* 2041–210X.13722 (2021) doi:10.1111/2041-
210X.13722.

30. Booker, T. R., Yeaman, S., Whiting, J. R. & Whitlock, M. C. The WZA: A
window-based method for characterizing genotype–environment associations. *Mol.*
Ecol. Resour. 1755–0998.13768 (2023) doi:10.1111/1755-0998.13768.

31. Goudet, J. & Weir, B. S. An allele-sharing, moment-based estimator of global,
population-specific and population-pair FST under a general model of population
structure. *PLOS Genet.* **19**, e1010871 (2023).

32. Li, H. & Ralph, P. Local PCA Shows How the Effect of Population Structure Differs
Along the Genome. *Genetics* **211**, 289–304 (2019).

33. Kawecki, T. J. Adaptation to Marginal Habitats. *Annu. Rev. Ecol. Evol. Syst.* **39**, 321–
342 (2008).

34. Angert, A. L., Bontrager, M. G. & Ågren, J. What Do We Really Know About
Adaptation at Range Edges? *Annu. Rev. Ecol. Evol. Syst.* **51**, 341–361 (2020).

35. Hampe, A. & Petit, R. J. Conserving biodiversity under climate change: the rear edge
matters. *Ecol. Lett.* **8**, 461–467 (2005).

36. Yannic, G. *et al.* Genetic diversity in caribou linked to past and future climate change.
Nat. Clim. Change **4**, 132–137 (2014).

37. Pellissier, L. *et al.* Past climate-driven range shifts and population genetic diversity in
arctic plants. *J. Biogeogr.* **43**, 461–470 (2016).

38. Phillips, S. J., Anderson, R. P. & Schapire, R. E. Maximum entropy modeling of species
geographic distributions. *Ecol. Model.* **190**, 231–259 (2006).

39. Machado, A. P. *et al.* Unexpected post-glacial colonisation route explains the white
colour of barn owls (*Tyto alba*) from the British Isles. *Mol. Ecol.* mec.16250 (2021)
doi:10.1111/mec.16250.

40. Fumagalli, M. *et al.* Signatures of Environmental Genetic Adaptation Pinpoint
Pathogens as the Main Selective Pressure through Human Evolution. *PLoS Genet.* **7**,
e1002355 (2011).

41. Shultz, A. J. & Sackton, T. B. Immune genes are hotspots of shared positive selection
across birds and mammals. *eLife* **8**, e41815 (2019).

42. Clark, N. J. *et al.* Climate, host phylogeny and the connectivity of host communities
govern regional parasite assembly. *Divers. Distrib.* **24**, 13–23 (2018).

43. Harvey, J. A. & Voelker, G. Host associations and climate influence avian
haemosporidian distributions in Benin. *Int. J. Parasitol.* **49**, 27–36 (2019).

44. O'Connor, E. A., Hasselquist, D., Nilsson, J.-Å., Westerdahl, H. & Cornwallis, C. K.
Wetter climates select for higher immune gene diversity in resident, but not
migratory, songbirds. *Proc. R. Soc. B Biol. Sci.* **287**, 20192675 (2020).

45.   Salewski, V. & Watt, C. Bergmann's rule: a biophysiological rule examined in birds.
Oikos **126**, oik.03698 (2017).

46.   Goodman, R. E., Lebuhn, G., Seavy, N. E., Gardali, T. & Bluso-Demers, J. D. Avian body
size changes and climate change: warming or increasing variability? *Glob. Change*
Biol. **18**, 63–73 (2012).

47.   Bonin, A. Explorative Genome Scan to Detect Candidate Loci for Adaptation Along a
Gradient of Altitude in the Common Frog (Rana temporaria). *Mol. Biol. Evol.* **23**, 773–
783 (2006).

48.   Kirkpatrick, M. & Barton, N. Chromosome Inversions, Local Adaptation and
Speciation. *Genetics* **173**, 419–434 (2006).

49.   Wellenreuther, M. & Bernatchez, L. Eco-Evolutionary Genomics of Chromosomal
Inversions. *Trends Ecol. Evol.* **33**, 427–440 (2018).

50.   Berdan, E. L. *et al.* How chromosomal inversions reorient the evolutionary process. *J.*
Evol. Biol. jeb.14242 (2023) doi:10.1111/jeb.14242.

51.   Faria, R., Johannesson, K., Butlin, R. K. & Westram, A. M. Evolving Inversions. *Trends*
Ecol. Evol. **34**, 239–248 (2019).

52.   Tigano, A. & Friesen, V. L. Genomics of local adaptation with gene flow. *Mol. Ecol.* **25**,
2144–2164 (2016).

53.   Durmaz, E., Kerdaffrec, E., Katsianis, G., Kapun, M. & Flatt, T. How Selection Acts on
Chromosomal Inversions. in *Encyclopedia of Life Sciences* 307–315 (Wiley, 2020).
doi:10.1002/9780470015902.a0028745.

54.   Bomblies, K. & Peichel, C. L. Genetics of adaptation. *Proc. Natl. Acad. Sci.* **119**,
e2122152119 (2022).

55.   Barrett, R. & Schluter, D. Adaptation from standing genetic variation. *Trends Ecol.*
Evol. **23**, 38–44 (2008).

56.   Chan, Y. F. *et al.* Adaptive Evolution of Pelvic Reduction in Sticklebacks by Recurrent
Deletion of a *Pitx1* Enhancer. *Science* **327**, 302–305 (2010).

57.   Smeds, L., Qvarnström, A. & Ellegren, H. Direct estimate of the rate of germline
mutation in a bird. *Genome Res.* **26**, 1211–1218 (2016).

58.   Bergeron, L. A. *et al.* Evolution of the germline mutation rate across vertebrates.
Nature **615**, 285–291 (2023).

59.   Rees, J. S., Castellano, S. & Andrés, A. M. The Genomics of Human Local Adaptation.
Trends Genet. **36**, 415–428 (2020).

60. Luqman, H., Wegmann, D., Fior, S. & Widmer, A. Climate-induced range shifts drive
adaptive response via spatio-temporal sieving of alleles. *Nat. Commun.* **14**, 1080
(2023).

61. de Filippo, C. *et al.* Recent Selection Changes in Human Genes under Long-Term
Balancing Selection. *Mol. Biol. Evol.* **33**, 1435–1447 (2016).

62. Shapiro, J. A. *et al.* Adaptive genic evolution in the *Drosophila* genomes. *Proc. Natl.*
Acad. Sci. **104**, 2271–2276 (2007).

63. Enbody, E. D. *et al.* Community-wide genome sequencing reveals 30 years of
Darwin's finch evolution. *Science* **381**, eadf6218 (2023).

64. Haber, M. *et al.* A Rare Deep-Rooting D0 African Y-Chromosomal Haplogroup and Its
Implications for the Expansion of Modern Humans Out of Africa. *Genetics* **212**, 1421–
1428 (2019).

65. Jones, F. C. *et al.* The genomic basis of adaptive evolution in threespine sticklebacks.
Nature **484**, 55–61 (2012).

66. Gandolfo, L. C., Bahlo, M. & Speed, T. P. Dating Rare Mutations from Small Samples
with Dense Marker Data. *Genetics* **197**, 1315–1327 (2014).

67. Albers, P. K. & McVean, G. Dating genomic variants and shared ancestry in
population-scale sequencing data. *PLOS Biol.* **18**, e3000586 (2020).

68. Marsico, T. D. & Hellmann, J. J. Dispersal limitation inferred from an experimental
translocation of *Lomatium* (Apiaceae) species outside their geographic ranges. *Oikos*
118, 1783–1792 (2009).

69. Hargreaves, A. L. & Eckert, C. G. Evolution of dispersal and mating systems along
geographic gradients: implications for shifting ranges. *Funct. Ecol.* **28**, 5–21 (2014).

70. Excoffier, L., Foll, M. & Petit, R. J. Genetic Consequences of Range Expansions. *Annu.*
Rev. Ecol. Evol. Syst. **40**, 481–501 (2009).

71. Hallatschek, O. & Nelson, D. R. LIFE AT THE FRONT OF AN EXPANDING POPULATION.
Evolution **64**, 193–206 (2010).

72. Peischl, S., Dupanloup, I., Kirkpatrick, M. & Excoffier, L. On the accumulation of
deleterious mutations during range expansions. *Mol. Ecol.* **22**, 5972–5982 (2013).

73. Peischl, S., Kirkpatrick, M. & Excoffier, L. Expansion Load and the Evolutionary
Dynamics of a Species Range. *Am. Nat.* **185**, E81–E93 (2015).

74. Peischl, S. & Excoffier, L. Expansion load: recessive mutations and the role of standing
genetic variation. *Mol. Ecol.* **24**, 2084–2094 (2015).

75. Angert, A. L., Bradshaw Jr, H. D. & Schemske, D. W. USING EXPERIMENTAL
EVOLUTION TO INVESTIGATE GEOGRAPHIC RANGE LIMITS IN MONKEYFLOWERS.
Evolution **62**, 2660–2675 (2008).

76. Holt, R. D. & Barfield, M. Theoretical Perspectives on the Statics and Dynamics of
Species' Borders in Patchy Environments. *Am. Nat.* **178**, S6–S25 (2011).

77. Polechová, J. & Barton, N. H. Limits to adaptation along environmental gradients.
Proc. Natl. Acad. Sci. **112**, 6401–6406 (2015).

78. Gilbert, K. J. *et al.* Local Adaptation Interacts with Expansion Load during Range
Expansion: Maladaptation Reduces Expansion Load. *Am. Nat.* **189**, 368–380 (2017).

79. Phillips, S. J., Anderson, R. P., Dudík, M., Schapire, R. E. & Blair, M. E. Opening the
black box: an open-source release of Maxent. *Ecography* **40**, 887–893 (2017).

80. Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G. & Jarvis, A. Very high resolution
interpolated climate surfaces for global land areas. *Int. J. Climatol.* **25**, 1965–1978
(2005).

81. Exposito-Alonso, M. rbioclim: Improved getData function from the raster R package
to interact with past, present and future climate data from worldclim.org.
github.com/MoisesExpositoAlonso/rbioclim (2017).

82. Feng, X., Park, D. S., Liang, Y., Pandey, R. & Papeş, M. Collinearity in ecological niche
modeling: Confusions and challenges. *Ecol. Evol.* **9**, 10365–10376 (2019).

83. Hijmans, R. J., Phillips, S. & Elith, J. L. and J. dismo: Species Distribution Modeling.
(2023).

84. Merow, C. *et al.* What do we gain from simplicity versus complexity in species
distribution models? *Ecography* **37**, 1267–1281 (2014).

85. Warren, D. L. & Seifert, S. N. Ecological niche modeling in Maxent: the importance of
model complexity and the performance of model selection criteria. *Ecol. Appl.* **21**,
335–342 (2011).

86. BirdLife International. IUCN Red List of Threatened Species: Tyto alba. *IUCN Red List*
Threat. Species (2016).

87. Swets, J. A. Measuring the Accuracy of Diagnostic Systems. *Science* **240**, 1285–1293
(1988).

88. Li, Y., Li, M., Li, C. & Liu, Z. Optimized Maxent Model Predictions of Climate Change
Impacts on the Suitable Distribution of Cunninghamia lanceolata in China. *Forests* **11**,
302 (2020).

89. Liu, C., White, M. & Newell, G. Selecting thresholds for the prediction of species
occurrence with presence-only data. *J. Biogeogr.* **40**, 778–789 (2013).

90. Elith, J. *et al.* A statistical explanation of MaxEnt for ecologists: Statistical explanation
of MaxEnt. *Divers. Distrib.* **17**, 43–57 (2011).

91. Lafarge, T. & Pateiro-Lopez, B. alphashape3d: Implementation of the 3D Alpha-Shape
for the Reconstruction of 3D Sets from a Point Cloud. (2020).

92. Machado, A. P. *et al.* Genomic consequences of colonisation, migration and genetic
drift in barn owl insular populations of the eastern Mediterranean. *Mol. Ecol.* **31**,
1375–1388 (2022).

93. Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler
transform. *Bioinforma. Oxf. Engl.* **26**, 589–595 (2010).

94. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158
(2011).

95. Manichaikul, A. *et al.* Robust relationship inference in genome-wide association
studies. *Bioinformatics* **26**, 2867–2873 (2010).

96. Xiuwen   Zheng   [Aut,   C.   SNPRelate.   Bioconductor
https://doi.org/10.18129/B9.BIOC.SNPRELATE (2017).

97. Martin, M. *et al.* *WhatsHap: Fast and Accurate Read-Based Phasing.*
http://biorxiv.org/lookup/doi/10.1101/085050 (2016) doi:10.1101/085050.

98. Delaneau, O., Zagury, J.-F., Robinson, M. R., Marchini, J. L. & Dermitzakis, E. T.
Accurate, scalable and integrative haplotype estimation. *Nat. Commun.* **10**, 5436
(2019).

99. Delaneau, O., Marchini, J. & Zagury, J.-F. A linear complexity phasing method for
thousands of genomes. *Nat. Methods* **9**, 179–181 (2012).

100. Choi, Y., Chan, A. P., Kirkness, E., Telenti, A. & Schork, N. J. Comparison of phasing
strategies for whole human genomes. *PLOS Genet.* **14**, e1007308 (2018).

101. Goudet, J., Jombart, T., Kamvar, Z. N., Archer, E. & Hardy, O. hierfstat: Estimation and
Tests of Hierarchical F-Statistics. (2021).

102. Oksanen, J. *et al.* vegan: Community Ecology Package. (2020).

103. Borcard, D., Gillet, F. & Legendre, P. *Numerical Ecology with R*. (Springer New York,
New York, NY, 2011). doi:10.1007/978-1-4419-7976-6.

104. Capblancq, T., Luu, K., Blum, M. G. B. & Bazin, E. Evaluation of redundancy analysis to
identify signatures of local adaptation. *Mol. Ecol. Resour.* **18**, 1223–1233 (2018).

105. Storey, J. D., Bass, A. J., Dabney, A., Robinson, D. & Warnes, G. qvalue: Q-value
estimation for false discovery rate control. Bioconductor version: Release (3.16)
https://doi.org/10.18129/B9.bioc.qvalue (2022).

106.   Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for animals
852   and plants. *Bioinformatics* **36**, 2628–2629 (2020).

Figures and tables

Figure 1 - Environmental and genetic variation across the suitable range of the European barn owl. (a) Map depicting the climatic heterogeneity across the range of the European barn owl according to the species distribution modelling. Dark grey cells surrounded by a white border have suitable climatic conditions similar to conditions present during the Last glacial Maximum (LGM). Coloured cells outside the white-bordered polygon have climatic conditions not present during the LGM. Colours are based on the multidimensional climatic space of bioclimatic data shown in (b): the scores of the first three principal components (PCs) were converted into values of RGB (PC1: red; PC2: green; PC3: blue) to represent variation in climate. Similar colours represent similar climates. Symbols represent sampling coordinates of individuals from 9 different populations. A jitter has been added for better visualisation (longitude: 0.425; latitude: 0.42). (b) Variance explained by the 7 first principal components of the PCA made on bioclimatic variables from the entire study area pictured on

(a). Climatic variables came both from the LGM and today to picture the overall climatic variability. (**c**) Correlation between climatic variables and the two first axes of the PCA. Dark to white gradients picture the contribution of each axis projected at the European scale - Abbreviations : CT = Coldest ; DT = Driest ; MTH = Month ; Precip = Precipitation ; QR = Quarter ; Temp = Temperature ; WT = Wettest (**d**) PCA based on the whole genome of the 74 European barn owls identified in (a). Symbols legend is the same as for panel (a). Only the two first principal components are represented.

Figure 2 - Genomic signatures of selection linked to climate in the European barn owls. (a) Scatterplot of genomic windows (100kbp each) across nine populations. Vertical and horizontal dashed lines are at 2 standard deviation from the mean of z-scores (population-specific FST) and z-scores (p-values) of WZA respectively. Each colour presents a class of windows. Dark blue dots represent outlier windows in both population-specific FST and WZA scans. Light blue dots are outlier according to WZA and have a population-specific FST higher than 1 standard deviation from the mean; (**b**) Genome-wide distribution of the WZA score. Dark blue dots corresponding to the windows identified in panel (a). A switch between light and dark grey represents a change in the scaffold. The names of all scaffolds are displayed on the upper x-axis. (**c**) Distribution of outlier windows in the different populations. Each row represents a population (DK: Denmark; FR: France; CH: Switzerland; PT: Portugal; IT: Italy; SB: Serbia; GR: Greece; AG: Aegean islands; IS: Israel). Each column is a window along the genome, coloured according to the classification in (a). (**d**) Barplot of outlier windows per population. Dark blue bars picture the outlier windows shared with at least one other population while white dashed bars picture the number of outlier windows unique to each population.

Figure 3 - Divergent selection drives the strong climate-driven signal on Super-Scaffold 45
(a) WZA signal across Super-Scaffold 45 with the main peak of outliers highlighted in grey. Dark blue circles are outlier windows from WZA and population-specific FST. Red circles represent outliers with a -log10(p-values) higher than 8. (b) Zoom on this peak of the Super-Scaffold 45. (c) Population-pair FST in this region between two populations at the extreme of the first climatic axis in Figure 1b (namely Denmark and Portugal). Lower value indicates a higher divergence in this region compared to the rest of the genome. (d) Pairwise FST between Denmark and Portugal, computed on a SNP-basis. (e) Genotypes of each SNP within the highlighted region (red rectangle on panel (B), (C) and (D)), using phased haplotypes from Denmark and Portugal. Beige represents the reference allele, and red represents the alternative allele.

Figure 4 - The shared signal in southern populations point to a putative inversion linked
with local adaptation in the European barn owl (**a**) WZA signal along Super-Scaffold 22 with
the first 14 Mb highlighted in grey. Dark blue circles are outlier windows from WZA and
population-specific FST. (**b**) Population-pair FST along the entire Super-Scaffold 22, higher
value indicates a higher similarity in this region compared to the rest of the genome. As in
(A), the first 14 Mb are highlighted in grey. Pairwise comparisons include France (FR), Italy
(IT) and Portugal (PT). (**c**) PCA made on the first 14 Mb of the Super-Scaffold 22, (120,953

SNPs - DK: Denmark; FR: France; CH: Switzerland; PT: Portugal; IT: Italy; SB: Serbia; GR:
Greece; AG: Aegean islands; IS: Israel).

Population	N	# PolymorphicS	# PrivateA	# RareA	PopSpecificFST
AE	10	4,231,387 (44,618)	193,178 (16,440)	687,423 (28,433)	0.03
CH	9	4,259,867 (19,877)	170,886 (4,123)	642,071 (9,990)	0.038
DK	10	4,102,932 (17,633)	135,566 (3,939)	560,136 (9,451)	0.049
FR	4	3,722,772 (0)	124,430 (1,292)	487,175 (1,859)	0.045
GR	9	4,176,443 (19,842)	144,856 (3,756)	606,164 (8,453)	0.046
IS	9	4,402,055 (12,195)	626,578 (7,595)	1,173,410 (6,991)	0.009
IT	9	4,120,120 (10,620)	214,759 (2,352)	679,772 (5,274)	0.054
PT	9	4,678,275 (25,247)	492,983 (7,476)	1,090,608 (12,569)	-0.018
SB	5	4,005,667 (0)	106,496 (2,004)	514,405 (2,428)	0.06

955   **Table 1 - Genetic diversity estimated for 9 populations of Western Palearctic barn owls.**
956   Standard deviations are found between brackets. See Material and Methods for details on
957   calculation. N, sample size; # PolymorphicS, number of polymorphic sites; # PrivateA,
958   number of private alleles; # RareA, number of rare alleles; PopSpecificFST, population-
959   specific FST computed at the whole genome level.