Secondary bile acid production by gut bacteria promotes Western diet-associated colorectal cancer

Esther Wortmann[*1], Annika Osswald[*2], David Wylensek[1], Stephanie Kuhls[3], Olivia I. Coleman[3], Quinten Ducarmon[4], Wei Liang[5], Nicole Treichel[1], Fabian Schumacher[6], Colin Volet[7], Silke Matysik[8], Karin Kleigrewe[9], Michael Gigl[9], Sascha Rohn[10], Burkhard Kleuser[6], Gerhard Liebisch[8], Angelika Schnieke[5], Rizlan Bernier-Latmani[7], Georg Zeller[4], Dirk Haller[3,11], Krzysztof Flisikowski[5], Soeren Ocvirk[**,2,11], Thomas Clavel[**, ***,1]

[1]Functional Microbiome Research Group, Institute of Medical Microbiology, RWTH University Hospital, Aachen, Germany.

[2]Intestinal Microbiology Research Group, German Institute of Human Nutrition, Potsdam-Rehbruecke, Nuthetal, Germany.

[3] Chair of Nutrition and Immunology, School of Life Sciences, Technical University of Munich, Freising, Germany.

[4]Structural and Computational Biology Unit, European Molecular Biology Laboratory, Heidelberg, Germany.

[5]Chair of Livestock Biotechnology, School of Life Sciences, Technical University of Munich, Freising, Germany.

[6]Institute of Pharmacy, Department of Pharmacology and Toxicology, Freie Universität Berlin, Berlin, Germany.

[7]Environmental Microbiology Laboratory, School of Architecture, Civil and Environmental Engineering, École Polytechnique Fédérale de Lausanne, Lausanne, Switzerland.

[8]Institute of Clinical Chemistry and Laboratory Medicine, University Hospital of Regensburg, Regensburg, Germany.

[9]Bavarian Center for Biomolecular Mass Spectrometry, School of Life Sciences, Technical University of Munich, Freising, Germany.

[10]Institute of Food Technology and Food Chemistry, Technische Universität Berlin, Berlin, Germany.

[11]ZIEL – Institute for Food and Health, Technical University of Munich, Freising, Germany

*: shared first authors, **: shared last authors, ***: corresponding author

Conflict of Interest
The authors declare no potential conflicts of interest.

Abstract

Western diet is an important risk factor for the development of sporadic colorectal cancer (CRC). Dietary fat stimulates bile acid (BA) production by the host and their conversion to secondary BAs by 7α-dehydroxylating (7αDH+) bacteria, but causal proof of their tumor-promoting effects *in vivo* is lacking. To address this, we performed feeding studies in a genetically engineered pig model of CRC combined with multi-omics analyses and gnotobiotic mouse studies. Western diet worsened the disease phenotype in $APC^{1311/+}$ pigs. This was accompanied by microbiota changes, increased levels of the secondary bile acid deoxycholic acid (DCA), and higher colonic epithelial cell proliferation. The latter was counteracted by using the BA-scavenging drug colestyramine. Metagenomic analysis across multiple cohorts revealed higher occurrence of *bai* (BA inducible) operons from *Clostridium scindens* and close relatives in stool of CRC subjects (n = 1,034). Using two gnotobiotic mouse models of CRC, we demonstrate that colonization with 7αDH+ bacteria (*C. scindens* or *Extibacter muris*) increased colonic tumor loads. This work provides clear evidence for the causal role of microbiome-derived DCA production in CRC under detrimental dietary conditions, opening avenues for future preventive strategies.

Introduction

Colorectal cancer (CRC) has one of the highest incidences and mortalities among all cancer types worldwide, with approximately 1.9 million new cases and 0.9 million deaths in 2020 (Sung et al., 2021). Besides genetic factors, including mutations in the tumor-suppressing adenomatous polyposis coli (*APC*) gene, sporadic CRC risk is predominantly influenced by environmental and lifestyle factors, including diet and the gut microbiome (Huxley et al., 2009; Keum & Giovannucci, 2019; Lichtenstein et al., 2000). There is convincing evidence for a positive association between CRC risk and Western-style diets rich in fat and red/processed meat, whilst high-fiber intake is negatively associated with CRC risk (Chen et al., 2013; Garcia-Larsen et al., 2019; Reynolds et al., 2019). In healthy individuals in South Africa, who have a very low CRC risk, a diet switch every two weeks from a traditional high-fiber/low-fat to a Western-style low-fiber/high-fat dietary pattern altered gut microbiota composition and metabolism,

leading to an increase in markers associated with CRC risk such as intestinal epithelial
cell (IEC) proliferation and infiltration of CD3+ and CD68+ immune cells (O'Keefe et al.,
2015). Alaskan Native people, who traditionally consume a diet rich in fat and have the
world's highest risk for CRC, have a gut microbiota characterized by lower diversity and
metabolic adaption to bile acid (BA) metabolism (Ocvirk et al., 2020).

High fat intake stimulates BA synthesis by the host, resulting in increased amounts of
BAs entering the colon, where they are converted to secondary BAs by gut bacteria.
Elevated levels of secondary BAs, in particular deoxycholic acid (DCA), were detected in
feces of CRC patients and healthy cohorts at high CRC risk, and they correlated with
higher occurrence of BA-inducible (*bai*) genes involved in BA 7α-dehydroxylation (7αDH)
by gut bacteria (Hill et al., 1975; Kawano et al., 2010; Ocvirk & O'Keefe, 2021; Ou et al.,
2013; Wirbel et al., 2019). In the murine $Apc^{min/+}$ model, a diet supplemented with the
primary BA cholic acid (CA, 0.4% w/w, 12 weeks), the precursor of DCA, increased the
number and size of intestinal tumors (Wang et al., 2019). This effect was prevented by
antibiotic treatment, suggesting a contribution of the gut microbiota to CA-associated
tumorigenesis in this CRC model. Oral intake of DCA promoted tumor development in
the proximal colon of wildtype mice (0.2% w/w in diet for up to 10 months) (Bernstein et
al., 2011), high-grade dysplasia in AKR/J mice treated with azoxymethane (AOM)
(0.25% w/w in diet for >10 weeks) (Flynn et al., 2007) and higher numbers of intestinal
tumors in $Apc^{min/+}$ mice (0.2% w/v in drinking water for 12 weeks) (Cao et al., 2014).
Fecal microbiota transplantation from DCA-treated $Apc^{min/+}$ donor mice to microbiota-
depleted $Apc^{min/+}$ recipients on a control diet induced increased intestinal tumor numbers
(Cao et al., 2017). However, in these studies, mice were fed with DCA directly, which
does not mirror physiological conditions, or DCA levels were not analyzed. Moreover,
the $Apc^{min/+}$ mouse model develops tumors primarily in the small intestine, which does
not reflect the location of tumorigenesis in humans and contrasts to colon being the
main site of secondary BAs production (A.R Moser et al., 1995; Amy Rapaich Moser et
al., 1990). Hence, clear evidence for a causal role of microbially produced DCA in
relation to diet and underlying mechanisms in colonic tumorigenesis is lacking.

Here we investigated the role of secondary BA production by gut bacteria in CRC, based
on a multi-layer approach using dietary interventions in genetically engineered pigs,

patient stool samples, gnotobiotic mouse models of CRC, and a combination of multi-
omics and cultivation methods. This allowed us to span analyses from microbial
community level to detailed investigations of specific microbiota members involved in the
conversion of BAs, collectively demonstrating a diet-driven, tumor-promoting function of
secondary BA-producing bacteria.

Results

Western diet promotes high levels of secondary BAs and aggravates disease in $APC^{1311/+}$ pigs

Pigs and humans are omnivores and share similarities in their gut microbiota,
gastrointestinal anatomy, physiology, and metabolic processes. This has made the pig
an ideal model to study gastrointestinal diseases (Gonzalez et al., 2015; Rose et al.,
2022). In contrast to transgenic mouse models based on the *Apc* gene, $APC^{1311/+}$ pigs
recapitulate key features of human disease, including the location of tumors in the large
intestine (Flisikowska et al., 2012), disease progression, and differences in severity
within families (Flisikowski et al., 2022).

To investigate the role of a Western diet on the development of CRC under controlled
conditions in a preclinical setting, we performed a feeding intervention in $APC^{1311/+}$ pigs.
The animals were fed (1 kg/d) either a control diet (CTRL) or an experimental diet
enriched in red meat and lard (RL diet) for 3 months (see experimental design in **Fig.
S1a** and diet composition in **Table S1**). Compared with control $APC^{1311/+}$ pigs, both the
number and size of colonic polyps were significantly increased in the RL diet group (**Fig.
1a**). Higher numbers of Ki67+ cells in distal colonic crypts indicated increased
proliferation of the epithelium (**Fig. 1b**).

To assess the effects of experimental feeding on the gut microbial ecosystem, fecal
samples were analyzed by 16S rRNA gene amplicon sequencing before and after
intervention. *Alpha*-diversity analysis revealed a trend towards increased richness and
Shannon effective counts in the CTRL group, which was not observed in the RL group,
including a sharp drop in diversity in the two pigs with most severe disease progression
(**Fig. 1c**). According to *beta*-diversity analysis, the microbiota structure between CTRL
and RL pigs overlapped at the beginning of the feeding trial (4 months) but diverged

after experimental feeding (7 months, **Fig. 1d**). At the level of dominant bacterial families, a decrease in *Prevotellaceae* and *Acidaminococcaceae* over time was observed in both groups, but it was significantly more pronounced in RL pigs (**Fig. 1e**). Whilst *Fibrobacteraceae* increased in the CTRL group, RL pigs were characterized by a significant increase in *Desulfovibrionaceae*.

Besides diet-induced effects on microbiota diversity and composition, changes in fecal metabolites were assessed. Targeted LC-MS/MS measurements showed that primary BAs were overall increased in RL pigs after dietary intervention, primarily due to glycocholic acid (GCA), which was not detected in the CTRL group (**Fig. 1f**). Total concentrations of secondary BAs were slightly higher in the CTRL group, mainly driven by hyodeoxycholic acid (HDCA), shown to suppress epithelial proliferation (M. Song et al., 2020). In contrast, levels of the CRC-associated secondary BAs DCA and 12-keto-litocholic acid (12-KLCA) were substantially higher in the RL group (**Fig. 1g**).

The bile acid sequestrant colestyramine limits Western diet-associated epithelial cell proliferation in the colon of *APC*[1311/+] pigs

Next, we investigated whether the increase in BAs contributed to the RL diet-induced phenotype in *APC*[1311/+] pigs. To this end, a second feeding trial was carried out, including a third group fed the RL diet supplemented with the drug colestyramine (COL; 12g/d), a bile acid scavenger that makes BAs unavailable for bacterial transformation and enterohepatic circulation (Zarras & Vogl, 1999) (see experimental design in **Fig. S1b**). Whilst the number of polyps tended to decrease after intervention in all COL pigs (**Fig. 2a;** $p = 0.062$), their size varied markedly between individuals in this group (**Fig. 2b**). Moreover, the number and size of polyps in the CTRL and RL groups showed inconsistent changes over time. However, immunohistochemistry staining of Ki67+ cells confirmed the proliferative effect of the RL diet in crypts of the distal colon, which was prevented by COL (**Fig. 2c**). Targeted analysis of fecal BAs at the end of the feeding period showed higher total levels of primary and secondary BAs in RL and COL pigs (p.adj. = 0.068 and 0.066, respectively), including higher concentrations of GCA (p.adj. = 0.05), allocholic acid (AlloCA) (p.adj. = 0.07), CA, and ursocholic acid (UCA) (p.adj. = 0.05) (**Fig. 2d**). Fecal concentrations of the secondary BAs DCA and 12-KLCA increased with RL diet feeding, whilst they tended to be lower under colestyramine

treatment (**Fig. 2e**). Targeted measurements of BAs were complemented by a lipidomics

approach. Levels of plant sterols were elevated in fecal samples from CTRL pigs and

colestyramine tended to compensate for this effect in the case of cholestanol and 5α

sitostanol (**Fig. 2f**). In contrast, the levels of cholesterol and derivatives as well as total

fatty acids (FA) were highest in feces of COL pigs (**Fig. 2g**). The differences in FA were

mainly driven by total saturated and branched chain (iso and ante-iso) FA, whereas

unsaturated and long-chain FA (22-24) showed no significant differences.

Taken together, the two feeding trials in genetically engineered pigs provide

interventional evidence for the positive association between Western dietary habits,

elevated levels of secondary BAs, and polyp formation. Western diet-driven epithelial

cell proliferation in the colon was prevented by the drug colestyramine, which stimulated

fecal excretion of lipids, making them likely unavailable for the host and BAs not

accessible for bacterial conversion into secondary BAs. To further demonstrate the

causal implication of secondary BAs in CRC development, we next used gnotobiotic

mouse models.

Specific 7αDH+ bacteria are associated with CRC

To identify the most relevant 7αDH+ bacteria in the context of CRC, we analyzed the

occurrence of *bai* operons from different mOTUs (metagenomic operational taxonomic

units) in fecal metagenomes from CRC patients (n = 1,034) and control individuals (n =

1,108) across multiple cohorts. The 7 mOTUs with a prevalence >2% showed specificity

in terms of both, *bai* operon structure and protein sequence comparison (**Fig. 3a, b**).

The occurrence of mOTUs corresponding to *Clostridium scindens* [r03437], *Clostridium*

hylemonae [r11469], and the yet-unknown species *Clostridiales* sp. [m14015] was

significantly higher in CRC patients (**Fig. 3c**). Cumulative relative abundances of the *bai*

operons from *C. scindens* and *C. hylemonae* confirmed the relevance of these two

species in CRC when analyzing all cohorts together (**Fig. 3d**). An increase was

observed in nearly all individual studies, but significance depended on the cohort

considered, likely due to varying size and environmental parameters (*e.g.,* diet) (**Fig.**

3e).

Due to the metagenomic data aforementioned, we used *C. scindens* in a gnotobiotic

mouse model to study the relevance of DCA production in CRC development. Germfree

wildtype mice on a high-fat diet were colonized with a simplified consortium of human

gut bacteria referred to as the "BA-converting synthetic microbiota" (BACOMI) (**Fig.**

S1d). BACOMI includes 7α-dehydroxylation activity (7αDH+) by *C. scindens* to produce

DCA and can be compared to the condition without DCA when *C. scindens* is excluded

from the consortium (7αDH-). Targeted qPCR analysis revealed a high relative

abundance of 23.0 ± 2.9% (mean ± SD) for *C. scindens* in BACOMI-colonized mice and

a relatively uniform colonization pattern of the other bacteria after 3 months (**Fig. 3f**).

Consistently, the presence of *C. scindens* caused high levels of secondary BAs, in

particular DCA, which was not detected when 7αDH activity was lacking in the microbial

consortium (**Fig. 3g**). DCA concentrations were higher in female than male mice when

colonized by *C. scindens*. Interestingly, 7αDH activity and DCA enhanced the

expression of genes within BA-dependent signaling pathways (**Fig. 3h**) and genes

known to be involved in tumorigenesis in the colonic epithelium (**Fig. 3i**). Furthermore,

7αDH activity was linked to higher numbers of Ki67+ epithelial cells in the colon (**Fig.**

3j).

Microbially produced deoxycholic acid promotes tumorigenesis in the colon

To analyze whether 7αDH-mediated BAs affect colonic tumorigenesis, germfree

wildtype mice on high-fat diet were colonized with BACOMI with or without *C. scindens*

and then treated with AOM and dextran sodium sulphate (DSS) (**Fig. 4a**). To

consolidate these findings independent of the 7αDH+ bacterial species used, we

performed additional experiments in mice colonized with the mouse gut isolate

Extibacter muris (Afrizal et al., 2022; Streidl et al., 2019) instead of *C. scindens*. The *bai*

operon of *E. muris* has a high similarity to the *bai* operon of *C. hylemonae* (Streidl et al.,

2021) and the latter species was also significantly associated with CRC in our

metagenomic analysis (**Fig. 3c**). Moreover, *C. hylemonae* is proposed to be reclassified

within the genus *Extibacter* in GTDB (Parks et al., 2022), highlighting the phylogenetic

relatedness of these two species.

Following AOM/DSS treatment, *C. scindens* was present at higher relative abundance

(50.3 ± 7.1%) (**Fig. 4b**) when compared with untreated BACOMI-colonized mice (**Fig.**

3f), suggesting favorable growth conditions for *C. scindens* under chemically induced

colonic tumorigenesis. DCA was only detected in gnotobiotic mice colonized with

C. scindens, including higher levels in female mice (**Fig. 4d**). Levels of oxo-DCA were

also higher in the caecal content of female mice when *C. scindens* was present. Despite

these sex differences in secondary BA concentrations, bacterial 7αDH activity led to

higher tumor numbers in the colon of nearly all mice (**Fig. 4f**). Stratification according to

tumor size revealed significance for small tumors, which suggests early-stage promotion

of tumorigenesis rather than tumor progression.

Compared with *C. scindens*-colonized mice*,* *E. muris* was less abundant, which was

linked to a more uniform distribution of the remaining bacterial species (**Fig. 4c**). DCA

was detected in caecum when 7αDH activity by *E. muris* was present and sex-specific

differences (higher levels in females) were confirmed (**Fig. 4e**). Several other secondary

BAs were significantly elevated when *E. muris* 7αDH was present (e.g., 12-oxo CDCA,

HCA, TDCA or LCA). Interestingly, levels of the primary BA CA were higher in 7αDH-

mice, indicating enhanced conversion in 7αDH+ mice. Similar to the effects of

C. scindens, 7αDH activity by *E. muris* caused higher numbers of tumors in the colon,

which was again linked primarily to small tumors (**Fig. 4g**).

To confirm the role of microbial DCA production in CRC development, we performed

gnotobiotic experiments in a further colonic tumor model using tissue specific nAtf6

transgenic mice (nAtf6[IEC]) (Coleman et al., 2018) with interleukin-10 knockout (*Il10*[-/-])

and a different synthetic community of cultured gut bacteria. Germfree IEC-specific Atf6

transgenic mice deficient for IL-10 (nAtf6[IEC tg/wt];*Il10*[-/-]) colonized with *E. muris* in addition

to the defined murine bacterial community OMM12 (Brugiroux et al., 2016) (OMM12+E)

were characterized by more and earlier dropouts (**Fig. 5a**), higher fraction of responders

(mice that developed tumors compared with non-responders) (**Fig. 5b**), and significantly

higher numbers of tumors (**Fig. 5c** and **5d**). Interestingly, nAtf6[IEC tg/wt];*Il10*[-/-] mice

colonized only with OMM12 and nAtf6[IEC fl/fl];*Il10*[-/-] colonized with *E. muris* (OMM12+E)

had lower number of tumors, indicating that both, DCA-producing bacteria and gene

susceptibility are required for colonic tumorigenesis.

Discussion

This study demonstrates the role of secondary BAs produced via gut bacterial 7αDH as tumor-promoting metabolites in experimental CRC. In contrast to non-physiological DCA supplementation in previous studies, we provide causal evidence linking the CRC risk-promoting effects of Western diets to functional adaptation of the gut microbiota. The experiments in $APC^{1311/+}$ pigs substantiate the clinical relevance of Western diet-mediated stimulation of microbial BA metabolism and suggest that promoting the excretion of BAs is a therapeutic option to limit their tumorigenic effects in the large intestine.

Bacterial 7αDH activity from the specific DCA-producers *C. scindens* and *E. muris* enhanced colitis-associated CRC in gnotobiotic mice. This validates our own findings about the higher occurrence of *bai* operons from *C. scindens* and *C. hylemonae* in metagenomes from CRC patients and it is an important experimental proof of previous observational data on higher numbers of 7αDH+ bacteria in feces of CRC patients or individuals at high CRC risk (Hill et al., 1975; Kawano et al., 2010; Ocvirk et al., 2020; Ocvirk & O'Keefe, 2021; Ou et al., 2013; Wirbel et al., 2019). It also provides physiologically relevant evidence for the tumor-promoting effects of secondary BAs in murine CRC models compared with previously tested oral DCA supplementation and observed effects in the small intestine (Bernstein et al., 2011; Cao et al., 2014, 2017; Flynn et al., 2007). Of note, 7αDH caused significantly higher numbers of small but not large colonic tumors in our gnotobiotic mouse experiments with AOM/DSS treatment. This may indicate that DCA promotes cellular processes underlying tumor initiation rather than progression in the colonic epithelium of susceptible hosts. Data obtained in the nAtf6$^{IEC\ tg/wt}$;$Il10^{-/-}$ gnotobiotic model of CRC demonstrates the importance of host genetic modification besides gut inflammation for the tumor-promoting effect of 7αDH+ bacteria.

There are multiple possible mechanisms for the tumor-promoting effect of DCA. For instance, it was shown to stimulate cancer stemness in colonic epithelial cells by modulating β-catenin signaling (Farhana et al., 2016; Pai et al., 2004). It also triggered proliferation and DNA damage in $Apc^{Min/+}$ Lgr5+ intestinal stem cells by antagonizing the

BA receptor Fxr (T. Fu et al., 2019). Moreover, oral administration of DCA and high-fat

diet-associated secondary BAs promoted proliferation of Lgr5+ stem cells via the BA

receptor Tgr5 (J.-Y. Li et al., 2022). Consistently, 7αDH-mediated production of DCA did

not induce colonic tumors without AOM/DSS in our study but it increased Ki67+

epithelial cell numbers and enhanced the expression of genes previously shown to be

involved in secondary BA-associated colonic tumorigenesis (T. Fu et al., 2019; Pai et al.,

2004). The tumor-promoting role of secondary BAs in susceptible hosts due to diet is

also supported by our observation of Western diet-driven increase in Ki67+ cells in the

colon of $APC^{1311/+}$ pigs. Furthermore, enhanced fecal excretion of BAs by colestyramine

was associated with reduced proliferation in the colonic epithelium. This finding provides

additional pre-clinical evidence for the potential use of colestyramine for CRC

prevention. Previous data in mice showed that colestyramine decreased the expression

of genes involved in epithelial cell proliferation in the colon and that it could prevent

streptozotocin and high-fat diet-induced hepatocellular carcinoma (J.-Y. Li et al., 2022;

Xie et al., 2016). The use of BA sequestrants in individuals at increased CRC risk

appears to be promising for prevention, since pre-diagnostic high plasma levels of BAs

were linked to CRC outcome in a prospective study (Kühn et al., 2020). In this context of

preventive BA reduction, it is worth noticing that secondary BAs also increased the

expression of genes with tumor-suppressive or essential regulatory functions (e.g.,

Guca2a, *Muc2*, *Pparg*) in our gnotobiotic mice, suggesting that 7αDH-associated

microbial metabolites have a fundamental role in regulating intestinal epithelium

homeostasis and regeneration. Thus, low but not completely abolished DCA levels

seem to be a promising preventive or therapeutic target for CRC risk reduction.

Recently, BAs were shown to be required for TGR5-dependent regeneration of intestinal

epithelial cells and protection from colitis in mice (Sorrentino et al., 2020), further

supporting the importance of physiological BA levels.

In the mouse experiments, we detected a larger caecal BA pool and higher DCA

concentrations in female compared to male animals. It is known that female and male

mice differ in their BA composition and female mice tend to have higher serum and liver

DCA concentrations (Z. D. Fu et al., 2012; Ma et al., 2020; Phelps et al., 2019; Selwyn

et al., 2015). Moreover, sex hormones influence hepatic BA metabolism, including their

production, secretion, and absorption (Z. D. Fu et al., 2012; Phelps et al., 2019).
Therefore, our data is in line with already described sex differences in BAs and
illustrates the importance to investigate both sexes. Besides BAs, sex hormones also
affect CRC risk directly: testosterone increases CRC risk (Amos-Landgraf et al., 2014;
C.-H. Song et al., 2021), whilst estrogen has the opposite effect (Chlebowski et al.,
2004; Murphy et al., 2015; Son et al., 2019). This possibly explains why, in our
gnotobiotic experiments, colonic tumor numbers were increased by secondary BA
production in both male and female mice despite markedly higher DCA concentrations in
the latter group.

Besides the causal role of secondary BAs in experimental CRC as demonstrated here,
Western diet may promote CRC risk via the microbiome in several manners. For
example, a recent study by Yang et al. (2022) (Yang et al., 2022) demonstrated that
experimental CRC was associated with higher levels of *Alistipes* spp. and dysregulation
of gut microbial co-metabolism leading to higher levels of glycerophospholipids.
Moreover, diets high in fat content are usually associated with increased occurrence of
Enterobacteriaceae, potentially including pks-expressing *Escherichia coli* with
tumorigenic effects (Arthur et al., 2012). CRC risk does not exclusively arise from the
increase in tumor-promoting factors but also from the lack of tumor-suppressive dietary
compounds, including complex carbohydrates fermented by gut bacteria to SCFA such
as butyrate (Donohoe et al., 2014; O'Keefe et al., 2015; Ocvirk et al., 2020).
Furthermore, Western or high-fat diets are very diverse in terms of fat content and
quality, with consequences regarding effects on the host. A recent study by Dmitrieva-
Posocco et al. (2022) showed that a ketogenic diet, highly enriched in fat and very low in
carbohydrates, reduced experimental colonic tumorigenesis due to anti-proliferative
effects of the ketone body β-hydroxybutyrate. However, the long-term clinical relevance
of ketogenic diets remains to be demonstrated. Taken together, these findings highlight
the complexity of diet-associated changes in gut microbial metabolism and support the
use of model organisms and experimental diets with detailed composition to unravel the
influence of nutrition on CRC development or prevention.

In conclusion, our studies demonstrate the causal role of gut microbial 7αDH activity in
experimental CRC. Secondary BAs, in particular DCA, are implicated in the tumor-

promoting effects of Western diets, which provides a promising target for CRC
prevention via dietary intervention or modulation of gut microbial BA metabolism.

Methods

Animal work

342 All experiments were carried out according to the German Animal Welfare Act and the
343 European Union Normative for Care and Use of Experimental Animals. Individual
344 approval numbers from the corresponding authorities are provided in the following
345 subsections that describe the specific experiments.

Porcine experiments

347 These experiments were approved by the Federal Government of Bavaria (permit no.
348 ROB55.2-2-2532.Vet_02-18-33). Pigs were provided by the Chair of Livestock
349 Biotechnology at the Technical University of Munich, Germany, and were housed and
350 sampled at the animal facility Thalhausen (Faculty of Life Science Weihenstephan,
351 Freising, Germany).

Dietary Interventions in $APC^{1311/+}$ pigs

The experimental design is depicted in**. Fig. S1**. $APC^{1311/+}$ pigs (German landrace ×
minipig crossbreed) (Flisikowski et al., 2022) underwent colonoscopy at the age of 3
months to enable their separation into two groups with a comparable distribution of
individuals in terms of polyp numbers and size, sex, and littermates. Both groups were
fed a standard diet until the dietary intervention started at the age of 4 months to allow
for recovery after colonoscopy. The control (CTRL) and experimental diet enriched in
red meat and lard (RL) were specifically designed for this experiment and purchased
from ssniff Spezialdiäten GmbH (Soest, Germany; cat. no. S5745-S074 and S5745-
S076, respectively). Main differences between the diets are illustrated in **Fig. S1** and
their detailed compositions are provided in **Table S1.** The diets were fed to the pigs in a
restricted manner (approx. 1 kg/d) for 3 months until the age of 7 months. At the end of
the feeding period, fecal samples were taken, and colonoscopy was performed,
including biopsy collection. Two RL pigs dropped out of the study due to tumor-
independent reasons (meningitis and a foot malformation).

For the second feeding trial testing the effects of colestyramine, the overall experimental
design was as in the first experiment (**Fig**. **S1**). However, $APC^{1311/+}$ pigs were assigned
to three groups instead of two (n = 4-5 pigs each): CTRL, RL, and COL (= RL diet
supplemented with 12 g/kg colestyramine; Ratiopharm GmbH, Ulm, Germany; EAN
4150037520609). The recommended dose by the manufacturer is 4-16 g/day for adult
subjects. Additional components of the drug colestyramine besides the active agent
were added to CTRL and RL diet. Detailed diet compositions are available in **Table S1.**

Colonoscopy

Pigs underwent colonoscopy one month prior to dietary intervention and after the three
months of experimental feeding. To reduce the load of fecal material in the large bowel,
pigs were fasted for 24 h before the procedure. Based on bodyweight, pigs were
narcotized (0.2 mg/kg ketamin and 0.05 mL/kg azaperon) and the large bowel was
flushed with tap water to remove remaining luminal content. Tissue biopsies and images
were taken using a STORZ colonoscopy system (Flexible SILVER SCOPE®).

Collection of fecal samples and sampling

In both pig trials, fresh fecal samples were collected prior to (4 months of age) and
immediately after dietary intervention (7 months of age) before colonoscopy. In brief,
fecal material was taken directly from the rectum of each pig using sterile gloves and
placed into individual, sterile plastic containers. Solely the material from core areas of
the feces was collected using a sterile spatula. Samples were immediately frozen on dry
ice for later sequencing and metabolomics and stored at -80°C. Tissue samples from the
rectum were collected either during colonoscopy (pig trial 1) or after the pigs were culled
(pig trial 2). The resulting samples were formalin-fixed and paraffin embedded tissue
blocks were stored at room temperature.

Mouse experiments

BACOMI consortium

The synthetic microbial community BACOMI (Bile acid-converting microbiota) contains
six bacterial species (all obtained from the Leibniz-Institute DSMZ - German Collection
of Microorganisms and Cell Cultures, Germany) that can transform the host-derived
primary BAs into dehydroxylated secondary BAs (adapted from Ridlon et al. (Ridlon et

al., 2020)) (**Fig. S1d**). *Bacteroides uniformis* (*B. uniformis*) DSM 6597, *Phocaeicola*
vulgatus (*P. vulgatus*) DSM 1447 and *Parabacteroides distasonis* (*P. distasonis*) DSM
20701 express bile salt hydrolases (BSH) that produce free primary BAs from
conjugated counterparts. *Bilophila wadsworthia* (*B. wadsworthia*) DSM 11045 can
further metabolize taurine released by BSH activity into H_2S. *Clostridium scindens* (*C.*
scindens) DSM 5676 is the key-member of the BACOMI consortium, as it is the only
species with 7α-dehydroxylating (7αDH) activity. Finally, *Blautia producta* (*B. producta*)
DSM 2950 was added due to being a dominant human gut bacterium that is H_2-
consuming, acetogenic, able to isomerize bile acids and thus increases variability of the
BA pool. The other 7αDH+ species *Extibacter muris* (*E. muris*) DSM 28560 was isolated
and described by us and available in the in-house collection (Lagkouvardos, Pukall, et
al., 2016; Streidl et al., 2021). All bacteria except for *B. wadsworthia* were cultured in
brain heart infusion (BHI) medium supplemented with resazurin, hemin, yeast extract
and L-cysteine under anaerobic conditions. *B. wadsworthia* was cultured as previously
described (Burkhardt et al., 2021).

OMM12 +/- *E. muris* consortium
OMM12 is a synthetic community of cultured bacteria representing the 5 major phyla in
the mouse gut (Brugiroux et al., 2016). We amended this consortium by the addition of
the 7αDH+ species *Extibacter muris* DSM 28560^T (*E. muris*). For the generation of cryo-
stocks, individual strains were grown in anaerobic Akkermansia medium (Brugiroux et
al., 2016). Strains were mixed equally (based on OD measurements) and mixed with
20% glycerol in a 1:1 ratio (exact mixing volumes and strain designations are given in
Table S7).

Gnotobiotic mouse experiments
The experiments in AOM/DSS treated mice were approved by the Ministry of Social
Affairs, Health, Integration and Consumer Protection of the state Brandenburg (permit
no. 2347-15-2021). Germfree male and female wildtype mice (BALB/cJ) were kept in
positive-pressure isolators at the animal facility of the German Institute for Human
Nutrition Potsdam-Rehbruecke (Nuthetal, Germany) with a 12 h light-dark cycle at 22 ±
2 °C and 55 ± 5 % air humidity. Starting 1 week prior to colonization, the mice were
switched from standard diet to a high-fat diet (HFD, **Table S1)**. Mice (n = 10 per group)

were colonized by gavage (10^8 cells per bacterium in 100 µL medium; 50 µL orally and
50 µL rectally at day 0 and 2) with BACOMI either with (7αDH+) or without (7αDH-) *C.*
scindens. After colonization, mice were either kept for 13 weeks (**Fig. S1 and Fig. 5**) or
subjected to further treatment to induce experimental CRC (**Fig. 6**). In the latter case,
mice received three intraperitoneal injections of azoxymethane (AOM) (5 mg/kg body
weight) in the interval of one week, starting at week 2 after colonization. Five days after
the final AOM injection, mice were treated with 3 cycles of dextran sodium sulphate
(DSS) with 5 days of DSS (1.5 % in drinking water) followed by 16 days of normal
drinking water. Mice were culled two weeks after the end of the AOM/DSS treatment
and colonic tumors were counted using a binocular microscope. The colonization and
AOM/DSS treatment of mice with *E. muris* followed the same protocol, but mice were
kept 2 weeks longer after colonization and prior to AOM/DSS treatment to confirm
colonization status*.* For BACOMI with *C. scindens* colonization, three mice (male n = 1,
female n = 2) dropped out in the 7αDH+ group during the first DSS cycle, whereas for
the 7αDH- group, two mice (male n = 1, female n = 1) dropped out at the end of the third
DSS cycle or after the first AOM injection, respectively. For BACOMI with *E. muris*
colonization, two female mice dropped out in the 7αDH+ group after the first DSS cycle.

Germ-free IEC-specific expression of activated (p50) nAtf6 transgenic mice (nAtf6^{IEC})
(Coleman et al., 2018) were crossed to germ-free *Il10$^{-/-}$* mice. Heterozygous mice with
monoallelic nAtf6 expression (nAtf6$^{fl/wt}$;Vil-Cre$^{tg/wt}$;*Il10$^{-/-}$*) and floxed controls (nAtf6$^{fl/wt}$;Vil-
Cre$^{wt/wt}$;*Il10$^{-/-}$*) were used for gnotobiotic work. These experiments were approved by the
Committee on Animal Health Care and Use of the state of Upper Bavaria (Regierung
von Oberbayern; AZ TVA 55.2-2532.Vet_02-18-121). Four-week-old mice were
inoculated orally (50µL) and rectally (100µL) with cryo-stocks of the OMM12 or OMM12
+ *E. muris* (OMM12+E) strain mixture at day 0 and day 3, and kept colonized for 12
weeks. All mice were euthanized with CO_2 at the end of an experiment (age 16 weeks)
or when abortion criteria were met.

Quantitative real-time PCR for intestinal gene expression

The Dynabeads mRNA purification kit (ThermoFisher Scientific, USA) was used for
mRNA isolation. Lysis buffer (200 µl) and 10 zirconium beads were added to

approximately 5 mg of intestinal tissue stored at -80 °C in RNA later tissue reagent
(Qiagen, Germany). A Tissue lyser (Qiagen, Germany) was then used for
homogenization (2 × 2 min at 50 Hz with a 2 min break in between). The lysate was
pulled through a syringe five times with a 1 mL syringe and centrifuged (14,000 rpm, 2
min, room temperature). The rest of the protocol was as per the manufacturer's
instructions. mRNA concentrations were determined using a Nanodrop (Peqlab VWR,
USA) and 20 ng were used for reverse transcription using the RevertAid H Minus First
Strand cDNA Synthesis kit (ThermoFisher Scientific, USA) according to manufacturer's
recommendations. For qPCR, the mastermix was prepared according to the
manufacturer's recommendations given by the Quantinova SYBR Green PCR Kit
(Qiagen, Germany), and forward and reverse primers added (0.5 μM each) (**Table S5**).
Then, 4 ng template cDNA was used in a total reaction volume of 10 μl. Samples were
amplified using the 7500 Fast Real-Time PCR System (Applied Biosystems, USA). To
determine C_T-values, the Life Technologies 7500 Software (version 2.3) was used.
Normalization was to the housekeeping gene *Rpl13a,* which encodes for the 60S
ribosomal protein L13a. Results were calculated relative to the control group of 7αDH-
mice.

Immunohistochemistry (IHC) and immunofluorescence (IF)

Specimens from the two pig trials were fixed in 4 % formaldehyde solution for 24 h,
embedded, and sectioned (3 μm). The tissue sections were then deparaffinized, antigen
unmasked (citrate buffer, pH = 6) and endogenous peroxidases were inactivated in 3%
H_2O_2 for 10 min. After blocking for 1 h (2% goat serum in PBS), sections were incubated
overnight at 4 °C with the primary antibody for Ki67 (Rabbit, DSC innovative Diagnostik-
System KI681C002, dilution 1:400) followed by a second overnight incubation with the
secondary antibody (Goat Anti-Rabbit IgG, Santa Cruz sc-2780, dilution 1:400).
Peroxidase activity was detected using diaminobenzidine (DAB) substrate or the
VECTASTAIN® Elite® ABC Kit (Vectorlab PK-6100), respectively. Staining of Ki67 was
considered positive and quantified only when detected in nuclei. Ten randomly selected
fields (magnification x40) from each tissue section were analyzed. Values are presented
as the percentage of the stained cells per area.

For the gnotobiotic mouse trial (**Fig. 5**), the dissected tissues were fixed using
Methacarn solution and embedded in paraffin. Tissues were cut into 4 μm-thick sections
and deparaffinized by heating (60 °C, 15 min) followed by two times xylol for 3 min. The
specimens were then rehydrated using a decreasing alcohol series: 100 % for 2 min,
96 % for 2 min, and 70 % for 1 min, followed by water for 1 min. After rinsing twice in
water (5 min each), the specimens were boiled in 10 mM citrate buffer (pH = 6) for 30
min for antigen retrieval, followed by 30 min of cooling. Thereafter, the specimens were
washed 3 times with water for 5 min, once with PBS for 5 min, and then blocked with
PBS containing 5 % goat serum for 1 h at room temperature. Primary antibodies against
Ki67 (ab15580, dilution 1:200) were incubated overnight at 4°C. Specimens were then
washed three times in PBS (5 min each) and incubated with the secondary antibody
(AlexaFluor-488, ThermoFisher, dilution 1:200), and Hoechst (dilution 1:200) for 1-2 h at
room temperature. All antibodies were diluted in PBS containing 1 % bovine serum
albumin and 0.3 % Triton X-100. Finally, the specimens were washed in PBS for 5 min
each and cover with glass slides using mounting medium (Vector Laboratories, United
States). For imaging, a confocal laser microscope was used (LSM 780 microscope,
Zeiss, Germany). Tile scans with 40x magnification were acquired and analysed with the
ZEN (black edition) 2.3 software (Zeiss, Germany).

Microbiota composition analysis

Metagenomic DNA was obtained using a modified version of a previously published
protocol (Godon et al., 1997; Just et al., 2018). Cells were lysed mechanically via bead-
beating and DNA was purified on columns (Macherey-Nagel, Germany). A robotized
platform (Biomek4000, Beckman-Coulter, Germany) was then used for library
construction. The V3-V4 regions of 16S rRNA genes were amplified (25 cycles) in a two-
step PCR (Berry et al., 2011) using primers 341F (5′-cctacgggnggcwgcag) and 785R (5′-
gactachvgggtatctaatcc) (Klindworth et al., 2013) and 24 ng of template DNA. A double
combinatorial indexing strategy was used. Amplicons were purified using the AMPure
XP system (Beckmann Coulter, Germany), pooled in equimolar amounts and sequenced
in paired-end mode using a MiSeq system (Illumina).

Raw data were processed using IMNGS (https://www.imngs.org) (Lagkouvardos,
Joseph, et al., 2016), an online platform based on UPARSE (Edgar, 2013). Sequences

were clustered into operational taxonomic units (OTUs) at 97% sequence similarity and
only those with a relative abundance ≥ 0.25% in at least one sample were kept to
exclude spurious OTUs (Reitmeier et al., 2021). Parameters for data processing were:
min/max sequence length, 300-500nt; barcode mismatch, 1; end-trim length, 15; max.
no. expected errors, 3; trimming q score, 3. The number of sequences per sample after
filtering was 26,286 ± 8,044 for the first pig trial and 12,564 ± 3,954 for the gnotobiotic
mouse experiment with *E. muris*. Downstream diversity and composition analyses were
performed in Rhea (https://github.com/Lagkouvardos/Rhea) (Lagkouvardos et al., 2017).
For the first pig trial, taxonomies were assigned using SILVA release 128 (Pruesse et
al., 2012). Several families showed a significantly different behavior between the feeding
groups over the time. A pseudocount (lowest value × 10) was added to the normalized
relative abundance of each family per sample before calculation to avoid division by
zero. For family comparison over time, a start-adjusted delta-value was calculated as
follows: (end − start) / start. For changes in α-diversity, original values without pseudo-
count were used for the same calculation. Identities of the OTUs for the gnotobiotic
mouse experiment were confirmed using Eztaxon (Yoon et al., 2017). Normalized
relative abundances are plotted in **Fig. 2** and **Fig. 6** and provided in **Table S2** and **Table
S3.**

For the gnotobiotic mouse trials with *C. scindens,* the QIAmp Fast DNA Stool Mini Kit
(Qiagen, Germany) was used to extract and purify DNA from colon content.
Approximately 50-80 mg frozen content was added to a cryo-tube containing ca. 100 mg
of 0.1 mm glass beads, and 1 mL of InhibitEx buffer, followed by bead-beating (Uniprep,
UniEquip, Germany) for 2 × 5 min at 3,000 rpm. All samples were processed according
to manufacturer's instructions except for the elution volume (changed to 70 µL). For
DNA isolation from bacterial cultures, the Genomic DNA from Microorganisms Kit
(Macherey-Nagel, Germany) was used as per the manufacturer's instructions. The
disruption time was 12 min, and the samples were eluted twice using the same elution
buffer.

A standard curve plotting bacterial cell number against C_T-value was established for
each BACOMI species using RT-qPCR and species-specific primers (**Table S5**). The
standard curve was generated from overnight bacterial cultures and cell numbers were

determined using a Thoma cell counting chamber (Fein-Optik, Germany). For
quantification, bacterial DNA was isolated from colon content and bacterial cell numbers
(**Table S4**) were calculated using the generated standard curve.

Analysis of *bai* operon occurrence in metagenomes

To quantify the occurrence of *bai* operons in fecal metagenomes, we first extracted *bai*
sequences from Kim et al., (Kim et al., 2022) and retained the operons harboring at least
7 of the 8 genes found in experimentally validated *bai* operons. The complete contig of
each *bai* sequence was downloaded from GenBank. GECCO (v0.9.7) was trained on
the complete contigs with default parameters using Pfam v35.0 (Mistry et al., 2021) and
bai-specific HMMs (Wirbel et al., 2019) as training features. We then ran GECCO on
ProGenomes3 (PG3) representative genomes (n = 41,171) and selected those clusters
with more than seven unique *bai* genes, since all experimentally verified *bai* operons
had >7 unique *bai* genes with our approach, based on *bai* genes HMM annotation. This
resulted in a total of 18 PG3 genomes with complete *bai* operons and we extracted the
protein sequence of each of the individual operons. We then computed protein
sequence similarity across the *bai* genes between operons from all genomes to
understand how divergent the sequences are, which is important to know prior to
quantifying relative abundances in metagenomes. Protein sequence similarity was
computed by pairwise alignment (global mode, BLOSUM62 substitution matrix) as
implemented in Biopython v1.81 (Cock et al., 2009). Next, we used mmseqs map
(v13.45111) to map protein sequences of the complete operons of all 18 clusters against
our reduced GMGC (Global Microbial Gene Catalog) gut catalogue (n = 13,788,251
ORFs) to identify which genes contained parts of the operons. This approach also
enabled species-resolved *bai* operon quantification. We then collected a total of n =
2,142 samples from case-control metagenomic studies of CRC (n = 1,034 CRC, n =
1,108 CTRL, see **Table S8** for an overview). These metagenomes were processed and
profiled as follows: (i) Raw reads were cleaned using bbduk (v38.93), including low-
quality trimming on either side (qtrim=rl trimq=3), discarding low-quality reads (maq=25),
adapter removal (ktrim=r k=23 mink=11 hdist =1 tpe=true tbo=true; against the bbduk
adapter library), and length filtering (ml=45); (ii) Reads were screened for host
contamination using kraken2 (v2.1.2) against the human hg38 reference genome with

ribosomal sequences masked (Silva_v_138); (iii) Cleaned reads were then mapped onto
our reduced human gut gene catalogue using BWA-MEM (v0.7.17) with default
parameters and alignments were filtered to >45bp alignment length and >97% sequence
identity. Reads aligning to multiple genes contributed fractional counts towards each hit
gene. Alignment counts for a gene were normalized against its length, then scaled up
according to the strategy employed by NGLess
(https://ngless.embl.de/Functions.html#count) and propagated to the functional features
with which the gene is annotated. Extensive use of gffquant (v.2.10.0,
https://github.com/cschu/gff_quantifier) was made throughout the functional profiling
workflow. As information about which individual GMGC gene corresponded to a part of
the *bai* operon of a given bacterial species was available, we obtained species-resolved
relative abundances of *bai* operons. Values for all GMGC genes belonging to a specific
species' *bai* operon were summed to compute total relative abundance for the *bai*
operon of the given bacterial species. We only included *bai*-encoding species if they
occurrence at a prevalence of at least 2% in the entire dataset. To calculate prevalence
of the bacterial species, we performed taxonomic profiling using mOTUs (v3.1) with
default parameters after having run the bbduk quality filtering as described above.
Differential abundance analysis was performed using a linear mixed model with the
study as a random effect, through the SIAMCAT package. A pseudocount of 1e-8 was
used for plotting zero values on a logarithmic scale.

BA measurements

Pig stool samples from trial 1 and 2 were analyzed as described previously by Wegner
et al. (2017) and Reiter et al. (2021), respectively (Reiter et al., 2021; Wegner et al.,
2017).

For the BACOMI mouse experiments with *C. scindens,* lyophilized caecal content (125
mg) were mixed with 450 μL acetonitrile, followed by 15 min vortexing and then
centrifugation (13,000 × g, 60 min, 21 °C). Supernatant (300 μL) was mixed with the
internal standards (ISTDs) d_4-CDCA and d_4-LCA (10 μM each) and dried in a SpeedVac
system (Jouan RC10.22, ThermoFisher Scientific, USA). The residue was dissolved with
methanol and water [1:1]. Chromatographic separation of 5 μL sample was achieved on

a 1290 Infinity II HPLC (Agilent Technologies, Germany) equipped with a Poroshell EC-
C18 column (Agilent Technologies) (3.0 × 150 mm, 2.7 µm) connected to a guard
column (3.0 × 5 mm, 2.7 µm) of the same material. The column was tempered to 30 °C.
A mobile phase system consisting of 10 mM ammonium acetate/acetonitrile (80:20 v:v;
solvent A) and water/acetonitrile (20:80 v:v; solvent B), was applied at a flow rate of
0.2 mL/min. The initial solvent composition of 85% solvent A was maintained for 6 min.
Then, solvent A was decreased stepwise to 70, 40 and 20% within 20, 30 and 40 min of
total run-time, respectively. After 45 min, the mobile phase reached initial conditions
again. MS/MS analysis was carried out using a 6495C triple-quadrupole mass
spectrometer (Agilent Technologies) operating in the negative electrospray ionization
mode (ESI-). The ion source parameters were: sheath gas temperature, 400 °C; sheath
gas flow, 12 L/min of nitrogen; nebuliser pressure, 40 psi; drying gas temperature,
120 °C; drying gas flow, 15 L/min of nitrogen; capillary voltage, 4500 V; nozzle
voltage, 0 V; iFunnel high pressure RF voltage, 90 V and iFunnel low pressure RF
voltage, 60 V. A total of 20 compounds (18 bile acids and 2 ISTDs) were simultaneously
analyzed by single reaction monitoring (SRM) or, if feasible, by multiple reaction
monitoring (MRM). HPLC-MS/MS parameters are given in **Table S6.** Quantification was
performed using MassHunter Workstation Quantitative Analysis for QQQ (Version 10.1,
Agilent Technologies). Analytes were externally calibrated in the concentration range of
0.09 to 90 µM (ISTDs constantly 10 µM). BA concentrations were normalized to dry
weight of the caecal content subjected to extraction.

For the gnotobiotic mouse experiment with *E. muris,* samples were lyophilized overnight
at –60 °C. The dried caecal contents were weighed, and 6 ceramic beads (2.5 mm)
were added to each tube. Proportionally to the weight of each sample, between 500 µL
and 1500 µL of MeOH/H_2O (2/1) + 0.1% formic acid was used as extraction solvent.
Samples were homogenized in a Precellys 24 Tissue Homogenizer (Bertin Instruments,
France) at 6500 rpm 2 x 20" beat and 20" rest. The homogenized caecal samples were
centrifuged at 21,000 rcf, for 15 min, at 4 °C. 100uL from each supernatant or calibration
standard were transferred into individual wells of 2 mL 96-well plate. 50 µL of an ISTD
solution (CA-d_4, CDCA-d_4, TCA-d_4, TUDCA-d_4, DCA-d_4 and LCA-d_4, each at 2 µM in
methanol) was pipetted in each well. Immediately after the addition of ISTD, 600 µL of

0.2% formic acid in H_2O was added to each sample or calibration standard level. The
96-well plate was shaken with an orbital shaker at 300 rpm and centrifuged at 3500 rpm,
5 min, 4 °C. The contents of the 96-well plate were extracted by solid phase extraction
with an Oasis HLB 96-well uElution plate (Waters, USA). The extracted samples were
dried in a Biotage® SPE Dry 96 (Biotage, Sweden) at 20 °C and reconstituted with 100
μL of MeOH/H_2O (50/50). The plate was shaken with an orbital shaker at 300 rpm, 5 min
and centrifuged at 3500 rpm, 5 min, 4 °C. The samples were injected on the LC-HRMS
system.

The quantitative method was performed on an Agilent ultrahigh-performance liquid
chromatography 1290 series coupled in tandem to an Agilent 6530 Accurate-Mass Q-
TOF mass spectrometer (Vico-Oton et al., 2022). The separation was done on a Zorbax
Eclipse Plus C18 column (2.1 × 100 mm, 1.8 μm) and a guard column Zorbax Eclipse
Plus C18 (2.1 × 5 mm, 1.8 μm), both provided by Agilent technologies (USA). The
column compartment was kept heated at 50 °C. Two different solutions were used as
eluents: ammonium acetate [5 mM] in water as mobile phase A and pure acetonitrile as
mobile phase B. A constant flow of 0.4 mL/min was maintained over 26 min of run time
with the following gradient (expressed in eluent B percentage): 0-5.5 min, constant
21.5% B; 5.5-6 min, 21.5-24.5% B; 6-10 min, 24.5-25% B; 10-10.5 min, 25-29% B; 10.5-
14.5 min, isocratic 29% B; 14.5-15 min, 29-40% B; 15-18 min, 40-45% B; 18-20.5 min,
45-95% B; 20.5-23 min, constant 95% B; 23-23.1 min, 95-21.5% B; 23.10-26 min,
isocratic 21.50% B. The system equilibration was implemented at the end of the gradient
for 3 min in initial conditions. The autosampler temperature was maintained at 10 °C and
the injection volume was 5 μL. The ionization mode was operated in negative mode for
the detection using the Dual AJS Jet stream ESI Assembly. The QTOF acquisition
settings were configured in 4 GHz high-resolution mode (resolution 17000 FWHM at m/z
1000), data storage in profile mode and the high-resolution full MS chromatograms were
acquired over the range of m/z 100-1700 at a rate of 3 spectra/s. The mass
spectrometer was calibrated in negative mode using ESI-L solution from Agilent
technologies every 6 h to maintain the best possible mass accuracy. Source parameters
were setup as follows: drying gas flow, 8 L/min; gas temperature, 300 °C; nebuliser
pressure, 35 psi; capillary voltage, 3500 V; nozzle voltage, 1000 V. Data were

processed afterwards using the MassHunter Quantitative software and MassHunter
Qualitative software (Agilent technologies, USA) to control the mass accuracy for each
run. In the quantitative method, 43 bile acids were quantified by calibration curves. The
quantification was corrected by addition of internal standards in all samples and
calibration levels. Extracted ion chromatograms were generated using a retention time
window of ± 1.5 min and a mass extraction window of ± 60ppm around the theoretical
mass of the targeted bile acid.

For all experiments, BAs that were not detected by the respective methods were
assumed to absent, set to 0 μmol/g and only those BAs detected in at least 50 % of the
animals in at least one group were considered for downstream analyses. Statistics were
done only on dominant BA species (the concentration thresholds are shown in the
figures; all individual values for all BAs are available in **Table S2-4**). UDCA is a primary
BA in rodents, but a secondary BA in humans (Honda et al., 2020; J. Li & Dawson,
2019). Due to their similarity to humans, we classified UDCA as a secondary BA in pigs.

Lipidomic analyses

Fecal homogenates for analysis of fatty acids, sterols and stanols were prepared in
isopropanol as described before (Schött et al., 2018). Subsequently, the homogenates
were used for both fatty acid and sterol/stanol analysis. Concentrations of total fecal
fatty acids were determined by gas chromatography coupled to mass spectrometry (GC-
MS) as described previously (Ecker et al., 2012) with some modifications. Samples were
derivatized to fatty acid methyl ester (FAME). The initial column temperature of 50 °C
was held for 0.75 min, increased with 40 °C/min to 110 °C, with 6 °C/min to 210 °C, with
15 °C/min to 250 °C and held for 2 min. Iso- and anteiso-FAMEs standards were applied
to identify branched chain fatty acids and to calibrate the instrument response. Samples
for analysis of sterols and stanols were analyzed as described previously (Kunz &
Matysik, 2019).

Data visualization and statistics

Statistical analyses were done using R (Version 4.2.1) (R Core Team, 2022) in RStudio
(Posit team, 2022) using the rstatix package. For microbiota analyses, detailed
information is provided in Rhea (https://github.com/Lagkouvardos/Rhea) (Lagkouvardos

et al., 2017). Comparisons between two groups were done by Wilcoxon tests and
Benjamini-Hochberg adjustment. Comparisons between three groups were done by
Kruskal Wallis tests with Dunn's multiple comparisons and Benjamini-Hochberg
adjustments. Adjusted p-values are with stars as defined in the corresponding legends;
non-adjusted p-values are given as numbers directly in the graphs whenever
appropriate. All graphs were created in R using tidyverse (ggplot2; RRID:SCR_014601)
(Wickham H, 2016), ggpubr, cowplot, and ggbreak (Xu et al., 2021).

Data availability

711 All data used to generate the figures is available in the supplementary tables (**Table S2-**
712 **4**). Raw data of the 16S rRNA gene amplicon sequencing were submitted to ENA and
713 are available under project PRJEB60380.

Acknowledgements

We are grateful to Viola and Steffen Loebnitz, Tatiana Flisikowska, and Thomas
Winogrodzky from the Chair of Livestock Biotechnology at the Technical University of
Munich, Germany, and Dieter Saur from the Klinikum rechts der Isar, Technical
University of Munich, Germany, for their support with pig experiments. We also thank
Ines Grüner, Anika Sander, Sabine Schmidt and Anke Gühler from the German Institute
of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany, for their help with the
germfree mouse colony and colonization experiments, and Tobias Goris and Annett
Braune for their critical review of data.

Funding

TC received funding from the German Research Foundation (DFG): project
no. 460129525, NFDI4Microbiota; project no. 395357507 - SFB1371 "Microbiome
Signatures"; and project no. CL481/8-1. TC, GZ, DH, and KF received funding from the
German Ministry for Research and Education (BMBF): consortium project Mi-EOCRC
(KZ. 01KD2102D). SO and DH received funding from the DFG: project no. 338582098
and no. 395357507 - SFB1371 "Microbiome Signatures", respectively.

References

Afrizal, A., Jennings, S. A., Hitch, T. C., Riedel, T., Basic, M., Panyot, A., Treichel, N., Hager, F. T., Wong, E. O.-Y., Wolter, B., Viehof, A., von Strempel, A., Eberl, C., Buhl, E. M., Abt, B., Bleich, A., Tolba, R., Blank, L. M., Navarre, W. W., … Clavel, T. (2022). Enhanced cultured diversity of the mouse gut microbiota enables custom-made synthetic communities. *Cell Host & Microbe*, *30*(11), 1630-1645.e25. https://doi.org/1

Amos-Landgraf, J. M., Heijmans, J., Wielenga, M. C. B., Dunkin, E., Krentz, K. J., Clipson, L., Ederveen, A. G., Groothuis, P. G., Mosselman, S., Muncan, V., Hommes, D. W., Shedlovsky, A., Dove, W. F., & Van Den Brink, G. R. (2014). Sex disparity in colonic adenomagenesis involves promotion by male hormones, not protection by female hormones. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(46), 16514–16519. https://doi.org/10.1073/pnas.1323064111

Arthur, J. C., Perez-Chanona, E., Mühlbauer, M., Tomkovich, S., Uronis, J. M., Fan, T.-J., Campbell, B. J., Abujamel, T., Dogan, B., Rogers, A. B., Rhodes, J. M., Stintzi, A., Simpson, K. W., Hansen, J. J., Keku, T. O., Fodor, A. A., & Jobin, C. (2012). Intestinal Inflammation Targets Cancer-Inducing Activity of the Microbiota. *Science*, *338*(6103), 120–123. https://doi.org/10.1126/science.1224820

Bernstein, C., Holubec, H., Bhattacharyya, A. K., Nguyen, H., Payne, C. M., Zaitlin, B., & Bernstein, H. (2011). Carcinogenicity of deoxycholate, a secondary bile acid. *Archives of Toxicology*, *85*(8), 863–871. https://doi.org/10.1007/s00204-011-0648-7

Berry, D., Mahfoudh, K. Ben, Wagner, M., & Loy, A. (2011). Barcoded primers used in multiplex amplicon pyrosequencing bias amplification. *Applied and Environmental Microbiology*, *77*(21), 7846–7849. https://doi.org/10.1128/AEM.05220-11

Brugiroux, S., Beutler, M., Pfann, C., Garzetti, D., Ruscheweyh, H. J., Ring, D., Diehl, M., Herp, S., Lötscher, Y., Hussain, S., Bunk, B., Pukall, R., Huson, D. H., Münch, P. C., McHardy, A. C., McCoy, K. D., MacPherson, A. J., Loy, A., Clavel, T., … Stecher, B. (2016). Genome-guided design of a defined mouse microbiota that

confers colonization resistance against Salmonella enterica serovar Typhimurium.

Nature Microbiology, *2*(16215), 1–12. https://doi.org/10.1038/nmicrobiol.2016.215

Burkhardt, W., Rausch, T., Klopfleisch, R., Blaut, M., & Braune, A. (2021). Impact of

dietary sulfolipid-derived sulfoquinovose on gut microbiota composition and

inflammatory status of colitis-prone interleukin-10-deficient mice. *International*

Journal of Medical Microbiology, *311*(3), 151494.

https://doi.org/10.1016/j.ijmm.2021.151494

Cao, H., Luo, S., Xu, M., Zhang, Y., Song, S., Wang, S., Kong, X., He, N., Cao, X., Yan,

F., & Wang, B. (2014). The secondary bile acid, deoxycholate accelerates intestinal

adenoma–adenocarcinoma sequence in $Apc^{min/+}$ mice through enhancing Wnt

signaling. *Familial Cancer*, *13*(4), 563–571. https://doi.org/10.1007/s10689-014-

9742-3

Cao, H., Xu, M., Dong, W., Deng, B., Wang, S., Zhang, Y., Wang, S., Luo, S., Wang, W.,

Qi, Y., Gao, J., Cao, X., Yan, F., & Wang, B. (2017). Secondary bile acid-induced

dysbiosis promotes intestinal carcinogenesis. *International Journal of Cancer*,

140(11), 2545–2556. https://doi.org/10.1002/ijc.30643

Chen, H. M., Yu, Y. N., Wang, J. L., Lin, Y. W., Kong, X., Yang, C. Q., Yang, L., Liu, Z.

J., Yuan, Y. Z., Liu, F., Wu, J. X., Zhong, L., Fang, D. C., Zou, W., & Fang, J. Y.

(2013). Decreased dietary fiber intake and structural alteration of gut microbiota in

patients with advanced colorectal adenoma. *American Journal of Clinical Nutrition*,

97(5), 1044–1052. https://doi.org/10.3945/ajcn.112.046607

Chlebowski, R. T., Wactawski-Wende, J., Ritenbaugh, C., Hubbell, F. A., Ascensao, J.,

Rodabough, R. J., Rosenberg, C. A., Taylor, V. M., Harris, R., Chen, C., Adams-

Campbell, L. L., & White, E. (2004). Estrogen plus Progestin and Colorectal Cancer

in Postmenopausal Women. *New England Journal of Medicine*, *350*(10), 991–1004.

https://doi.org/10.1056/nejmoa032071

Cock, P. J. A., Antao, T., Chang, J. T., Chapman, B. A., Cox, C. J., Dalke, A., Friedberg,

I., Hamelryck, T., Kauff, F., Wilczynski, B., & De Hoon, M. J. L. (2009). Biopython:

Freely available Python tools for computational molecular biology and

bioinformatics. *Bioinformatics*, *25*(11), 1422–1423.

https://doi.org/10.1093/bioinformatics/btp163

Coleman, O. I., Lobner, E. M., Bierwirth, S., Sorbie, A., Waldschmitt, N., Rath, E.,

Berger, E., Lagkouvardos, I., Clavel, T., McCoy, K. D., Weber, A., Heikenwalder,

M., Janssen, K. P., & Haller, D. (2018). Activated ATF6 Induces Intestinal Dysbiosis

and Innate Immune Response to Promote Colorectal Tumorigenesis.

Gastroenterology, *155*(5), 1539-1552.e12.

https://doi.org/10.1053/j.gastro.2018.07.028

Dmitrieva-Posocco, O., Wong, A. C., Lundgren, P., Golos, A. M., Descamps, H. C.,

Dohnalová, L., Cramer, Z., Tian, Y., Yueh, B., Eskiocak, O., Egervari, G., Lan, Y.,

Liu, J., Fan, J., Kim, J., Madhu, B., Schneider, K. M., Khoziainova, S., Andreeva, N.,

… Levy, M. (2022). β-Hydroxybutyrate suppresses colorectal cancer. *Nature*, *605*,

160–165. https://doi.org/10.1038/s41586-022-04649-6

Donohoe, D. R., Holley, D., Collins, L. B., Montgomery, S. A., Whitmore, A. C.,

Hillhouse, A., Curry, K. P., Renner, S. W., Greenwalt, A., Ryan, E. P., Godfrey, V.,

Heise, M. T., Threadgill, D. S., Han, A., Swenberg, J. A., Threadgill, D. W., &

Bultman, S. J. (2014). A gnotobiotic mouse model demonstrates that dietary fiber

protects against colorectal tumorigenesis in a microbiota- and butyrate-dependent

manner. *Cancer Discovery*, *4*(12), 1387–1397. https://doi.org/10.1158/2159-

8290.CD-14-0501

Ecker, J., Scherer, M., Schmitz, G., & Liebisch, G. (2012). A rapid GC-MS method for

quantification of positional and geometric isomers of fatty acid methyl esters.

Journal of Chromatography B: Analytical Technologies in the Biomedical and Life

Sciences, *897*, 98–104. https://doi.org/10.1016/j.jchromb.2012.04.015

Edgar, R. C. (2013). UPARSE: Highly accurate OTU sequences from microbial amplicon

reads. *Nature Methods*, *10*(10), 996–998. https://doi.org/10.1038/nmeth.2604

Farhana, L., Nangia-Makker, P., Arbit, E., Shango, K., Sarkar, S., Mahmud, H., Hadden,

T., Yu, Y., & Majumdar, A. P. N. (2016). Bile acid: a potential inducer of colon

cancer stem cells. *Stem Cell Research & Therapy*, *7*(1), 181.

817   https://doi.org/10.1186/s13287-016-0439-4

818 Flisikowska, T., Merkl, C., Landmann, M., Eser, S., Rezaei, N., Cui, X., Kurome, M.,
819   Zakhartchenko, V., Kessler, B., Wieland, H., Rottmann, O., Schmid, R. M.,
820   Schneider, G., Kind, A., Wolf, E., Saur, D., & Schnieke, A. (2012). A porcine model
821   of familial adenomatous polyposis. *Gastroenterology*, *143*(5), 1173-1175.e7.
822   https://doi.org/10.1053/j.gastro.2012.07.110

823 Flisikowski, K., Perleberg, C., Niu, G., Winogrodzki, T., Bak, A., Liang, W., Grodziecki,
824   A., Zhang, Y., Pausch, H., Flisikowska, T., Klinger, B., Perkowska, A., Kind, A.,
825   Switonski, M., Janssen, K. P., Saur, D., & Schnieke, A. (2022). Wild-type APC
826   Influences the Severity of Familial Adenomatous Polyposis. *Cmgh*, *13*(2), 669-
827   671.e3. https://doi.org/10.1016/j.jcmgh.2021.11.002

828 Flynn, C., Montrose, D. C., Swank, D. L., Nakanishi, M., Ilsley, J. N. M., & Rosenberg, D.
829   W. (2007). Deoxycholic acid promotes the growth of colonic aberrant crypt foci.
830   *Molecular Carcinogenesis*, *46*(1), 60–70. https://doi.org/10.1002/mc.20253

831 Fu, T., Coulter, S., Yoshihara, E., Oh, T. G., Fang, S., Cayabyab, F., Zhu, Q., Zhang, T.,
832   Leblanc, M., Liu, S., He, M., Waizenegger, W., Gasser, E., Schnabl, B., Atkins, A.
833   R., Yu, R. T., Knight, R., Liddle, C., Downes, M., & Evans, R. M. (2019). FXR
834   Regulates Intestinal Cancer Stem Cell Proliferation. *Cell*, *176*(5), 1098-1112.e18.
835   https://doi.org/10.1016/j.cell.2019.01.036

836 Fu, Z. D., Csanaky, I. L., & Klaassen, C. D. (2012). Gender-divergent profile of bile acid
837   homeostasis during aging of mice. *PLoS ONE*, *7*(3), 1–13.
838   https://doi.org/10.1371/journal.pone.0032551

839 Garcia-Larsen, V., Morton, V., Norat, T., Moreira, A., Potts, J. F., Reeves, T., & Bakolis,
840   I. (2019). Dietary patterns derived from principal component analysis (PCA) and
841   risk of colorectal cancer: a systematic review and meta-analysis. *European Journal*
842   *of Clinical Nutrition*, *73*(3), 366–386. https://doi.org/10.1038/s41430-018-0234-7

843 Godon, J. J., Zumstein, E., Dabert, P., Habouzit, F., & Moletta, R. (1997). Molecular
844   microbial diversity of an anaerobic digestor as determined by small-subunit rDNA
845   sequence analysis. *Applied and Environmental Microbiology*, *63*(7), 2802–2813.

https://doi.org/10.1128/aem.63.7.2802-2813.1997

Gonzalez, L. M., Moeser, A. J., & Blikslager, A. T. (2015). Porcine models of digestive
disease: The future of large animal translational research. *Translational Research*,
166(1), 12–27. https://doi.org/10.1016/j.trsl.2015.01.004

Hill, M. J., Drasar, B. S., Williams, R. E. O., Meade, T. W., Cox, A. G., Simpson, J. E. P.,
& Morson, B. C. (1975). Faecal Bile-Acids and Clostridia in Patients With Cancer of
the Large Bowel. *The Lancet*, *305*(7906), 535–539. https://doi.org/10.1016/S0140-
6736(75)91556-1

Honda, A., Miyazaki, T., Iwamoto, J., Hirayama, T., Morishita, Y., Monma, T., Ueda, H.,
Mizuno, S., Sugiyama, F., Takahashi, S., & Ikegami, T. (2020). Regulation of bile
acid metabolism in mouse models with hydrophobic bile acid composition. *Journal*
of Lipid Research, *61*(1), 54–69. https://doi.org/10.1194/jlr.RA119000395

Huxley, R. R., Ansary-Moghaddam, A., Clifton, P., Czernichow, S., Parr, C. L., &
Woodward, M. (2009). The impact of dietary and lifestyle risk factors on risk of
colorectal cancer: A quantitative overview of the epidemiological evidence.
International Journal of Cancer, *125*(1), 171–180. https://doi.org/10.1002/ijc.24343

Just, S., Mondot, S., Ecker, J., Wegner, K., Rath, E., Gau, L., Streidl, T., Hery-Arnaud,
G., Schmidt, S., Lesker, T. R., Bieth, V., Dunkel, A., Strowig, T., Hofmann, T.,
Haller, D., Liebisch, G., Gérard, P., Rohn, S., Lepage, P., & Clavel, T. (2018). The
gut microbiota drives the impact of bile acids and fat source in diet on mouse
metabolism. *Microbiome*, *6*(1), 1–18. https://doi.org/10.1186/s40168-018-0510-8

Kawano, A., Ishikawa, H., Kamano, T., Kanoh, M., Sakamoto, K., Nakamura, T., Otani,
T., Sakai, T., & Kono, K. (2010). Significance of fecal deoxycholic acid
concentration for colorectal tumor enlargement. *Asian Pacific Journal of Cancer*
Prevention, *11*(6), 1541–1546.

Keum, N. N., & Giovannucci, E. (2019). Global burden of colorectal cancer: emerging
trends, risk factors and prevention strategies. *Nature Reviews Gastroenterology*
and Hepatology, *16*(12), 713–732. https://doi.org/10.1038/s41575-019-0189-8

Kim, K. H., Park, D., Jia, B., Baek, J. H., Hahn, Y., & Jeon, C. O. (2022). Identification

and Characterization of Major Bile Acid 7α-Dehydroxylating Bacteria in the Human

Gut. *MSystems*, *7*(4). https://doi.org/10.1128/msystems.00455-22

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., & Glöckner, F.

O. (2013). Evaluation of general 16S ribosomal RNA gene PCR primers for

classical and next-generation sequencing-based diversity studies. *Nucleic Acids*

Research, *41*(1), 1–11. https://doi.org/10.1093/nar/gks808

Kühn, T., Stepien, M., López-Nogueroles, M., Damms-Machado, A., Sookthai, D.,

Johnson, T., Roca, M., Hüsing, A., Maldonado, S. G., Cross, A. J., Murphy, N.,

Freisling, H., Rinaldi, S., Scalbert, A., Fedirko, V., Severi, G., Boutron-Ruault, M.-C.,

Mancini, F. R., Sowah, S. A., … Kaaks, R. (2020). Prediagnostic Plasma Bile Acid

Levels and Colon Cancer Risk: A Prospective Study. *JNCI: Journal of the National*

Cancer Institute, *112*(5), 516–524. https://doi.org/10.1093/jnci/djz166

Kunz, S., & Matysik, S. (2019). A comprehensive method to determine sterol species in

human faeces by GC-triple quadrupole MS. *The Journal of Steroid Biochemistry*

and Molecular Biology, *190*, 99–103. https://doi.org/10.1016/J.JSBMB.2019.03.014

Lagkouvardos, I., Fischer, S., Kumar, N., & Clavel, T. (2017). Rhea: A transparent and

modular R pipeline for microbial profiling based on 16S rRNA gene amplicons.

PeerJ, *5:e2836*(1). https://doi.org/10.7717/peerj.2836

Lagkouvardos, I., Joseph, D., Kapfhammer, M., Giritli, S., Horn, M., Haller, D., & Clavel,

T. (2016). IMNGS: A comprehensive open resource of processed 16S rRNA

microbial profiles for ecology and diversity studies. *Scientific Reports*, *6*(April), 1–9.

https://doi.org/10.1038/srep33721

Lagkouvardos, I., Pukall, R., Abt, B., Foesel, B. U., Meier-Kolthoff, J. P., Kumar, N.,

Bresciani, A., Martínez, I., Just, S., Ziegler, C., Brugiroux, S., Garzetti, D., Wenning,

M., Bui, T. P. N. N., Wang, J., Hugenholtz, F., Plugge, C. M., Peterson, D. A.,

Hornef, M. W., … Clavel, T. (2016). The Mouse Intestinal Bacterial Collection

(miBC) provides host-specific insight into cultured diversity and functional potential

of the gut microbiota. *Nature Microbiology*, *1*(10), 1.

https://doi.org/10.1038/nmicrobiol.2016.131

Li, J.-Y., Gillilland, M., Lee, A. A., Wu, X., Zhou, S.-Y., & Owyang, C. (2022). Secondary
bile acids mediate high-fat diet-induced upregulation of R-spondin 3 and intestinal
epithelial proliferation. *JCI Insight*, *7*(19), 1–17.
https://doi.org/10.1172/jci.insight.148309

Li, J., & Dawson, P. A. (2019). Animal models to study bile acid metabolism. *Biochimica*
et Biophysica Acta - Molecular Basis of Disease, *1865*(5), 895–911.
https://doi.org/10.1016/j.bbadis.2018.05.011

Lichtenstein, P., Holm, N. V., Verkasalo, P. K., Iliadou, A., Kaprio, J., Koskenvuo, M.,
Pukkala, E., Skytthe, A., & Hemminki, K. (2000). Environmental and Heritable
Factors in the Causation of Cancer — Analyses of Cohorts of Twins from Sweden,
Denmark, and Finland. *The New England Journal of Medicine*, *343*(2), 78–85.
https://doi.org/10.1056/NEJM200007133430201

Ma, J., Hong, Y., Zheng, N., Xie, G., Lyu, Y., Gu, Y., Xi, C., Chen, L., Wu, G., Li, Y., Tao,
X., Zhong, J., Huang, Z., Wu, W., Yuan, L., Lin, M., Lu, X., Zhang, W., Jia, W., … Li,
H. (2020). Gut microbiota remodeling reverses aging-associated inflammation and
dysregulation of systemic bile acid homeostasis in mice sex-specifically. *Gut*
Microbes, *11*(5), 1450–1474. https://doi.org/10.1080/19490976.2020.1763770

Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G. A., Sonnhammer, E.
L. L., Tosatto, S. C. E., Paladin, L., Raj, S., Richardson, L. J., Finn, R. D., &
Bateman, A. (2021). Pfam: The protein families database in 2021. *Nucleic Acids*
Research, *49*(D1), D412–D419. https://doi.org/10.1093/nar/gkaa913

Moser, A.R, Luongo, C., Gould, K. ., McNeley, M. ., Shoemaker, A. ., & Dove, W. .
(1995). Apc^{Min}: A mouse model for intestinal and mammary tumorigenesis.
European Journal of Cancer, *31*(7–8), 1061–1064. https://doi.org/10.1016/0959-
8049(95)00181-H

Moser, Amy Rapaich, Pitot, H. C., & Dove, W. F. (1990). A Dominant Mutation That
Predisposes to Multiple Intestinal Neoplasia in the Mouse. *Science*, *247*(4940),
322–324. https://doi.org/10.1126/science.2296722

Murphy, N., Strickler, H. D., Stanczyk, F. Z., Xue, X., Wassertheil-Smoller, S., Rohan, T.
E., Ho, G. Y. F., Anderson, G. L., Potter, J. D., & Gunter, M. J. (2015). A
Prospective Evaluation of Endogenous Sex Hormone Levels and Colorectal Cancer
Risk in Postmenopausal Women. *Journal of the National Cancer Institute*, *107*(10),
1–10. https://doi.org/10.1093/jnci/djv210

O'Keefe, S. J. D., Li, J. V., Lahti, L., Ou, J., Carbonero, F., Mohammed, K., Posma, J.
M., Kinross, J., Wahl, E., Ruder, E., Vipperla, K., Naidoo, V., Mtshali, L., Tims, S.,
Puylaert, P. G. B., Delany, J., Krasinskas, A., Benefiel, A. C., Kaseb, H. O., …
Zoetendal, E. G. (2015). Fat, fibre and cancer risk in African Americans and rural
Africans. *Nature Communications*, *6*(6342), 1–29.
https://doi.org/10.1038/ncomms7342

Ocvirk, S., & O'Keefe, S. J. D. (2021). Dietary fat, bile acid metabolism and colorectal
cancer. *Seminars in Cancer Biology*, *73*, 347–355.
https://doi.org/10.1016/j.semcancer.2020.10.003

Ocvirk, S., Wilson, A. S., Posma, J. M., Li, J. V., Koller, K. R., Day, G. M., Flanagan, C.
A., Otto, J. E., Sacco, P. E., Sacco, F. D., Sapp, F. R., Wilson, A. S., Newton, K.,
Brouard, F., Delany, J. P., Behnning, M., Appolonia, C. N., Soni, D., Bhatti, F., …
O'Keefe, S. J. D. (2020). A prospective cohort analysis of gut microbial co-
metabolism in Alaska Native and rural African people at high and low risk of
colorectal cancer. *American Journal of Clinical Nutrition*, *111*(2), 406–419.
https://doi.org/10.1093/ajcn/nqz301

Ou, J., Carbonero, F., Zoetendal, E. G., DeLany, J. P., Wang, M., Newton, K., Gaskins,
H. R., & O'Keefe, S. J. (2013). Diet, microbiota, and microbial metabolites in colon
cancer risk in rural Africans and African Americans. *The American Journal of*
Clinical Nutrition, *98*(1), 111–120. https://doi.org/10.3945/ajcn.112.056689

Pai, R., Tarnawski, A. S., & Tran, T. (2004). Deoxycholic Acid Activates β-Catenin
Signaling Pathway and Increases Colon Cell Cancer Growth and Invasiveness.
Molecular Biology of the Cell, *15*(5), 2156–2163. https://doi.org/10.1091/mbc.E03-
12-0894

Parks, D. H., Chuvochina, M., Rinke, C., Mussig, A. J., Chaumeil, P. A., & Hugenholtz, P. (2022). GTDB: An ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank normalized and complete genome-based taxonomy. *Nucleic Acids Research*, *50*(D1), D785–D794. https://doi.org/10.1093/nar/gkab776

Phelps, T., Snyder, E., Rodriguez, E., Child, H., & Harvey, P. (2019). The influence of biological sex and sex hormones on bile acid synthesis and cholesterol homeostasis. *Biology of Sex Differences*, *10*(1), 1–12. https://doi.org/10.1186/s13293-019-0265-3

Posit team. (2022). *RStudio: Integrated Development for R*. Posit Software, PBC. http://www.posit.co/

Pruesse, E., Peplies, J., & Glöckner, F. O. (2012). SINA: Accurate high-throughput multiple sequence alignment of ribosomal RNA genes. *Bioinformatics*, *28*(14), 1823–1829. https://doi.org/10.1093/bioinformatics/bts252

R Core Team. (2022). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing. https://www.r-project.org/

Reiter, S., Dunkel, A., Metwaly, A., Panes, J., Salas, A., Haller, D., & Hofmann, T. (2021). Development of a Highly Sensitive Ultra-High-Performance Liquid Chromatography Coupled to Electrospray Ionization Tandem Mass Spectrometry Quantitation Method for Fecal Bile Acids and Application on Crohn's Disease Studies. *Journal of Agricultural and Food Chemistry*, *69*, 5238–5251. https://doi.org/10.1021/acs.jafc.1c00769

Reitmeier, S., Hitch, T. C. A., Treichel, N., Fikas, N., Hausmann, B., Ramer-Tait, A. E., Neuhaus, K., Berry, D., Haller, D., Lagkouvardos, I., & Clavel, T. (2021). Handling of spurious sequences affects the outcome of high-throughput 16S rRNA gene amplicon profiling. *ISME Communications*, *1*(1). https://doi.org/10.1038/s43705-021-00033-z

Reynolds, A., Mann, J., Cummings, J., Winter, N., Mete, E., & Te Morenga, L. (2019). Carbohydrate quality and human health: a series of systematic reviews and meta-

analyses. *The Lancet*, *393*(10170), 434–445. https://doi.org/10.1016/S0140-
6736(18)31809-9

Ridlon, J. M., Devendran, S., Alves, J. M., Doden, H., Wolf, P. G., Pereira, G. V., Ly, L.,
Volland, A., Takei, H., Nittono, H., Murai, T., Kurosawa, T., Chlipala, G. E., Green,
S. J., Hernandez, A. G., Fields, C. J., Wright, C. L., Kakiyama, G., Cann, I., …
Gaskins, H. R. (2020). The '*in vivo* lifestyle' of bile acid 7α-dehydroxylating bacteria:
comparative genomics, metatranscriptomic, and bile acid metabolomics analysis of
a defined microbial community in gnotobiotic mice. *Gut Microbes*, *11*(3), 381–404.
https://doi.org/10.1080/19490976.2019.1618173

Rose, E. C., Blikslager, A. T., & Ziegler, A. L. (2022). Porcine Models of the Intestinal
Microbiota: The Translational Key to Understanding How Gut Commensals
Contribute to Gastrointestinal Disease. *Frontiers in Veterinary Science*, *9*(March),
1–8. https://doi.org/10.3389/fvets.2022.834598

Schött, H. F., Krautbauer, S., Höring, M., Liebisch, G., & Matysik, S. (2018). A Validated,
Fast Method for Quantification of Sterols and Gut Microbiome Derived 5α/β-Stanols
in Human Feces by Isotope Dilution LC-High-Resolution MS. *Analytical Chemistry*,
90(14), 8487–8494. https://doi.org/10.1021/acs.analchem.8b01278

Selwyn, F. P., Csanaky, I. L., Zhang, Y., & Klaassen, C. D. (2015). Special section on
drug metabolism and the microbiome: Importance of large intestine in regulating
bile acids and glucagon-like peptide-1 in germ-free mice. *Drug Metabolism and*
Disposition, *43*(10), 1544–1556. https://doi.org/10.1124/dmd.115.065276

Son, H. J., Sohn, S. H., Kim, N., Lee, H. N., Lee, S. M., Nam, R. H., Park, J. H., Song,
C. H., Shin, E., Na, H. Y., Kim, J. S., Lee, D. H., & Surh, Y. J. (2019). Effect of
estradiol in an azoxymethane/dextran sulfate sodium-treated mouse model of
colorectal cancer: Implication for sex difference in colorectal cancer development.
Cancer Research and Treatment, *51*(2), 632–648.
https://doi.org/10.4143/crt.2018.060

Song, C.-H., Kim, N., Nam, R. H., Choi, S. I., Yu, J. E., Nho, H., Shin, E., Lee, H.-N., &
Surh, Y.-J. (2021). Testosterone strongly enhances azoxymethane/dextran sulfate

sodium-induced colorectal cancer development in C57BL/6 mice. *Am J Cancer*
Res, *11*(6), 3145–3162. www.ajcr.us/ISSN:2156-
6976/ajcr0132476OriginalArticleTestosteronestronglyenhancesazoxymethane/dextr
ansulfatesodium-inducedcolorectalcancer

Song, M., Yang, Q., Zhang, F., Chen, L., Su, H., Yang, X., He, H., Liu, F., Zheng, J.,
Ling, M., Lai, X., Zhu, X., Wang, L., Gao, P., Shu, G., Jiang, Q., & Wang, S. (2020).
Hyodeoxycholic acid (HDCA) suppresses intestinal epithelial cell proliferation
through FXR-PI3K/AKT pathway, accompanied by alteration of bile acids
metabolism profiles induced by gut bacteria. *FASEB Journal, 34*(5), 7103–7117.
https://doi.org/10.1096/fj.201903244R

Sorrentino, G., Perino, A., Yildiz, E., El Alam, G., Bou Sleiman, M., Gioiello, A.,
Pellicciari, R., & Schoonjans, K. (2020). Bile Acids Signal via TGR5 to Activate
Intestinal Stem Cells and Epithelial Regeneration. *Gastroenterology*, *159*(3), 956-
968.e8. https://doi.org/10.1053/j.gastro.2020.05.067

Streidl, T., Karkossa, I., Segura Muñoz, R. R., Eberl, C., Zaufel, A., Plagge, J.,
Schmaltz, R., Schubert, K., Basic, M., Schneider, K. M., Afify, M., Trautwein, C.,
Tolba, R., Stecher, B., Doden, H. L., Ridlon, J. M., Ecker, J., Moustafa, T., von
Bergen, M., … Clavel, T. (2021). The gut bacterium Extibacter muris produces
secondary bile acids and influences liver physiology in gnotobiotic mice. *Gut*
Microbes, *13*(1), 1–21.

Streidl, T., Kumar, N., Suarez, L. N., Rohn, S., & Clavel, T. (2019). *Extibacter*. In
Bergey's Manual of Systematics of Archaea and Bacteria (pp. 1–7). Wiley.
https://onlinelibrary.wiley.com/doi/10.1002/9781118960608.gbm01749

Sung, H., Ferlay, J., Siegel, R. L., Laversanne, M., Soerjomataram, I., Jemal, A., & Bray,
F. (2021). Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and
Mortality Worldwide for 36 Cancers in 185 Countries. *CA: A Cancer Journal for*
Clinicians, *71*(3), 209–249. https://doi.org/10.3322/caac.21660

Vico-Oton, E., Volet, C., Jacquemin, N., Lederballe Meibom, K., & Bernier-Latmani, R.
(2022). Strain-dependent induction of primary bile acid 7-dehydroxylation by cholic

acid. *BioRxiv*. https://doi.org/10.1101/2022.02.15.480494

Wang, S., Dong, W., Liu, L., Xu, M., Wang, Y., Liu, T., Zhang, Y., Wang, B., & Cao, H.
(2019). Interplay between bile acids and the gut microbiota promotes intestinal
carcinogenesis. *Molecular Carcinogenesis*, *58*(7), 1155–1167.
https://doi.org/10.1002/mc.22999

Wegner, K., Just, S., Gau, L., Mueller, H., Gérard, P., Lepage, P., Clavel, T., & Rohn, S.
(2017). Rapid analysis of bile acids in different biological matrices using LC-ESI-
MS/MS for the investigation of bile acid transformation by mammalian gut bacteria.
Analytical and Bioanalytical Chemistry, *409*(5), 1231–1245.
https://doi.org/10.1007/s00216-016-0048-1

Wickham H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New
York. https://ggplot2.tidyverse.org

Wirbel, J., Pyl, P. T., Kartal, E., Zych, K., Kashani, A., Milanese, A., Fleck, J. S., Voigt,
A. Y., Palleja, A., Ponnudurai, R., Sunagawa, S., Coelho, L. P., Schrotz-King, P.,
Vogtmann, E., Habermann, N., Niméus, E., Thomas, A. M., Manghi, P., Gandini, S.,
… Zeller, G. (2019). Meta-analysis of fecal metagenomes reveals global microbial
signatures that are specific for colorectal cancer. *Nature Medicine*, *25*(4), 679–689.
https://doi.org/10.1038/s41591-019-0406-6

Xie, G., Wang, X., Huang, F., Zhao, A., Chen, W., Yan, J., Zhang, Y., Lei, S., Ge, K.,
Zheng, X., Liu, J., Su, M., Liu, P., & Jia, W. (2016). Dysregulated hepatic bile acids
collaboratively promote liver carcinogenesis. *International Journal of Cancer*,
139(8), 1764–1775. https://doi.org/10.1002/ijc.30219

Xu, S., Chen, M., Feng, T., Zhan, L., Zhou, L., & Yu, G. (2021). Use ggbreak to
Effectively Utilize Plotting Space to Deal With Large Datasets and Outliers.
Frontiers in Genetics, *12*(November), 1–7.
https://doi.org/10.3389/fgene.2021.774846

Yang, J., Wei, H., Zhou, Y., Szeto, C.-H., Li, C., Lin, Y., Coker, O. O., Lau, H. C. H.,
Chan, A. W. H., Sung, J. J. Y., & Yu, J. (2022). High-Fat Diet Promotes Colorectal
Tumorigenesis Through Modulating Gut Microbiota and Metabolites.

Gastroenterology, *162*(1), 135-149.e2. https://doi.org/10.1053/j.gastro.2021.08.041

Yoon, S. H., Ha, S. M., Kwon, S., Lim, J., Kim, Y., Seo, H., & Chun, J. (2017).

Introducing EzBioCloud: A taxonomically united database of 16S rRNA gene

sequences and whole-genome assemblies. *International Journal of Systematic and*

Evolutionary Microbiology, *67*(5), 1613–1617.

https://doi.org/10.1099/ijsem.0.001755

Zarras, P., & Vogl, O. (1999). Polycationic salts as bile acid sequestering agents.

Progress in Polymer Science, *24*(4), 485–516. https://doi.org/10.1016/S0079-

6700(99)00002-7

1086

1087

Figure Legends

Fig 1. Effects of Western diet on disease phenotype, fecal microbiota (16S rRNA amplicon sequencing) and bile acids (LC-MS/MS): a, number and size of polyps in the distal colon of $APC^{1311/+}$ pigs aged 3 months (3m, 1 month prior to the feeding intervention) or 7 months (7m, end of feeding intervention) and fed either the red meat and lard diet (RL, n = 8; red symbols) or the control diet (CTRL, n = 9; blue). Litters are indicated by different symbols. Right panel: Exemplary endoscopic pictures (last 40 cm of the colon). **b,** Percentage of Ki67+ cells in crypts of the distal colon from CTRL and RL pigs (n = 3 each) after the feeding intervention; for each pig, 10 randomly selected fields were counted (small colored dots). Mean values and standard deviations per pig are shown in grey. Right panel: Exemplary Ki67 staining of distal colon samples. Black bars on the pictures represent 200 µm. **c,** Richness and Shannon effective counts over time; p = 0.0206 and 0.1390, respectively (comparison of delta values). 4m (4 months) is the start of the dietary intervention, 7m (7 months) the end. **d,** *Beta*-diversity shown as metaNMDS plot of generalized Unifrac distances (p = 0.001, PERMANOVA; pairwise comparisons with Benjamini-Hochberg adjustment: start, p.adj. = 0.399, end, p.adj. = 0.0015). **e,** Relative abundances of dominant bacterial families with significantly different changes over time (end-start/start) between feeding groups (Wilcoxon rank sum test, p ≤ 0.05); **f,** Primary (grey stack bars) and secondary (brown) BA levels in faeces at the end of the feeding period; mean values per feeding group are shown (n = 9 for CTRL, n = 8 for RL). All values are available in **Table S2**. GCA = Glycocholic acid, CDCA = chenodeoxycholic acid, GCDCA= Glycochenodeoxycholic acid, βMCA = β -muricholic acid, CA = cholic acid, HCA = hyocholic acid, TCDCA = tauro-chenodeoxycholic acid, DCA = deoxycholic acid, DHCA = dehydro-lithocholic acid, HDCA= hyo-deoxycolic acid, ωMCA = ω-muricholic acid, 12-KLCA = 12-keto-litocholic acid, 3-DHCA = 3-dehydro cholic acid. **g,** Dot plots of the secondary BAs DCA and 12-KLCA in faeces. Mean values and standard deviations across all pigs per group are shown in grey. Statistics: **a,** Paired Wilcoxon signed rank test; **b,** unpaired Wilcoxon rank sum test; **c** and **e,** Wilcoxon rank sum Benjamini-Hochberg adjustment (* p.adj. ≤ 0.05, ** p.adj. ≤ 0.01).

Fig. 2 Effect of colestyramine treatment in transgenic $APC^{1311/+}$ pigs: The COL diet corresponded to RL diet supplemented with 12 g/kg colestyramine (see detailed composition in **Table S1**). Polyp number (**a**) and size (**b**) in the distal colon (last 40 cm, 3m = 3 months, one month prior to intervention; 7m = 7 months, end of feeding period, n = 5 for CTRL and COL, n = 4 for RL). Litters are indicated by different symbols. **c,** Percentage of Ki67+ cells in crypts of the distal colon. For each pig, 10 randomly selected fields were counted (colored dots). Mean values and standard deviations per pig are shown in grey. Exemplary microscopic pictures of Ki67 staining (bars = 200 µm) are displayed next to the graph. **d**, Mean values of primary (grey) and secondary (brown) BAs at the end of the feeding period (7 months). All values are given in **Table S3**. **e**, Individual values for DCA and 12-KLCA. Mean values and standard deviations are show in grey. **f**, Sterol/stanol profiles. **g**, FA profiles: total saturated FA (sat. FA), monounsaturated FA, di-unsaturated FA, >2 unsaturated FA, FAs separated by chain length, branched chain FA (iso- and ante-iso), and total FA. Individual values are provided in **Table S3**. Values in **f** and **g** are shown as individual values (black dots) and mean (bars) + standard deviation (whiskers). CTRL, control diet (n = 5); RL, red meat and lard diet (n = 4); COL, colestyramine diet (n = 5). Statistics: **a** and **b**, Paired Wilcoxon signed rank test; **c-g** Kruskal-Wallis test with Dunn's multiple comparisons and Benjamini-Hochberg adjustment (* p.adj. ≤ 0.05).

Fig. 3 Identification of specific DCA-producing bacteria associated with CRC and functional evaluation in a gnotobiotic mouse model: a Genomic architecture of the *bai* operon in mOTUs (metagenomic operational taxonomic units) detected at >2% prevalence (n = 2,142). **b** Protein sequence identities between the different genomic architectures of the *bai* operon shown in **a**. **c** Quantile plots of relative abundances of mOTU-specific *bai* operons over all CRC (n = 1,034) vs. control (n = 1,108) samples. Due to high protein sequence similarity between *Clostridiales* sp. [m12265] and *Clostridiales* sp. [m12309] (last two bacteria listed), relative abundances are almost identical. **d** Cumulative relative abundance of the *bai* operons from *C. scindens* and *C. hylemonae* in samples from all cohorts. Only the samples with a detected *bai* operon from either of the two species are plotted. The corresponding number of positive samples is indicated above the x-axis. The total number of samples is indicated below the x-axis. **e** Same as in d, but per study. Studies that included <10 samples with detected *bai* operons in either of the groups (CTRL or CRC) were excluded. Adjusted p-values < 0.5 are shown (Wilcoxon rank-sum test with Benjamini-Hochberg adjustment, excluding samples without *bai* operon detected). **f,** Relative abundances of bacterial species in colonic content of mice colonized with the BACOMI consortium with

(7αDH+, n = 10, 5 female (F) and 5 male (M) mice) or without (7αDH-, n = 10, 5F/5M) *C. scindens* as
measured by targeted qPCR. **g,** Primary (grey) and secondary (brown) BA profile in caecal content
quantified using LC-MS/MS (female and male mice are shown separately; the stacked bars indicate mean
values; all individual concentrations are available in **Table S4**). **h/i,** Expression of genes related to (**h**) BA
metabolism and (**i**) epithelial cell proliferation and differentiation in the colonic mucosa. **j,** Exemplary
images of immunofluorescence staining for Ki67+ cells in colonic crypts of BACOMI(7αDH-) or
BACOMI(7αDH+) mice and corresponding quantification shown as bar plots (n = 4 mice each). **f, h, i, j**
show mean values + standard deviation. Statistics: Wilcoxon rank sum test with Benjamini-Hochberg
adjustment (* p.adj. ≤ 0.05; ** p.adj. ≤ 0.01, *** p.adj. ≤ 0.001, **** p.adj. ≤ 0.0001).

Fig. 4 Effects of bacterial 7αDH on colonic tumorigenesis in gnotobiotic mice treated with
AOM/DSS. a, Setup of AOM/DSS experiments using BACOMI(7αDH-) mice deprived of 7αDH activity and
their BACOMI(7αDH+) counterparts colonised with either *C. scindens* (**b, d, f**, 7αDH-, n = 8-7; 7αDH+, n =
6) or *E. muris* (**c, e, g,** 7αDH-, n =10; 7αDH+, n = 8). Mice were culled after either 16 weeks (*C. scindens*)
or 18 weeks (*E. muris*). **b, c,** Relative abundances of the bacterial species in colonic content as analysed
by qPCR (**b**) or 16S rRNA gene amplicon analysis (**c**). **d, e,** Primary (grey) and secondary (brown) BA
profile in caecal content as quantified using LC-MS/MS (female and male mice shown separately). Data
are shown as stacked bar plots of mean values. Individual concentrations are available in **Table S4**. **f, g,**
Number of colonic tumors; total or stratified according to their size: small (< 1.2 mm^2), medium (1.2 - 2.5
mm^2), large (> 2.5 mm^2). Exemplary macroscopic pictures of the colon are shown below the bar plots;
single tumors or areas with many tumors are indicated with red arrowheads or circles. In **b**, **c**, **f**, **g**, the sex
of mice is indicated by different symbols (triangle = male; dots = female). Statistical analysis (**d-g**):
Wilcoxon rank sum test with Benjamini-Hochberg adjustment (* p.adj. ≤ 0.05; ** p.adj. ≤ 0.01, *** p.adj. ≤
0.001).

Fig. 5 Effects of bacterial 7αDH activity on colonic tumor development in gnotobiotic model of
colitis-associated cancer. Germfree nAtf6$^{IEC\ fl/fl}$;*Il10*$^{-/-}$ (fl/fl;-/-) and nAtf6$^{IEC\ tg/wt}$;*Il10*$^{-/-}$ (tg/wt;-/-) mice were
colonized either with the synthetic community OMM12 (n=6 per genotype) or OMM12+E (n=5 per
genotype). **a** Survival curves of mice during the experiment. **b** Percentage of responders (mice with
tumors) or non-responders (without tumors), including the respective fraction of early dropouts. **c** Number
of colon tumors in responder mice. **d** Representative macroscopic images of the entire colon cut open
longitudinally. Pictures of only tg/wt,-/- mice are shown; left, colonized with OMM12; right, colonized with
OMM12+E. Tumors are indicated with black arrows. Statistics: Wilcoxon rank sum tests with Benjamini-
Hochberg correction.

a

colonisation

BACOMI (7αDH−)

BACOMI (7αDH+)

BALB/cJ

high fat diet

treatment   AOM AOM AOM   DSS   H2O   DSS   H2O   DSS   H2O

−1   1   2   3   4   5   6   10   16/18   weeks

b   7αDH: *C. scindens*

rel. abundance [%]

P. vulgatus   B. uniformis   P. distasonis   B. producta   B. wexlerae   C. scindens   n.d.

c   7αDH: *E. muris*

norm. rel. abundance [%]

P. vulgatus   B. uniformis   B. distasonis   B. producta   B. wexlerae   E. muris   n.d.

d

| 7αDH− | 7αDH+ |

μmol/g dry weight

F   M   F   M

primary BAs
- CDCA
- αMCA
- βMCA
- CA
- TβMCA
- TCA
- other
- (< 0.001 μmol/g)

secondary BAs
- DCA**
- oxo-DCA*
- iso-DCA
- LCA
- other
- (< 0.0005 μmol/g)

e

| 7αDH− | 7αDH+ |

μmol/g dry weight

F   M   F   M

primary BAs
- CDCA
- TαMCA
- αMCA
- βMCA
- CA*
- CA7S
- TβMCA
- other
- (< 0.01 μmol/g)

secondary BAs
- DCA**
- 12-oxo CDCA*
- HCA**
- TDCA
- LCA**
- UDCA
- 12*-KLCA*
- 7-oxo DCA
- 7,12-dioxo CA
- 7,12-dioxo LCA
- iso-LCA
- TUDCA
- other
- (< 0.005 μmol/g)

f

Nr of colonic tumors

| total number | small | medium | large |

F   M

7αDH−   7αDH+ (*C. scindens*)

g

Nr of colonic tumors

| total number | small | medium | large |

F   M

7αDH−   7αDH+ (*E. muris*)

sex ● F ▲ M