Spontaneous helping in pigs is mediated by helper's social attention and distress signals of individuals in need

Liza R. Moscovice[1*], Anja Eggert[2], Christian Manteuffel[1] and Jean-Loup Rault[3]

**Affiliations:**

[1] Institute of Behavioural Physiology, Research Institute for Farm Animal Biology (FBN), Dummerstorf, Germany

[2] Institute of Genetics and Biometry, Research Institute for Farm Animal Biology (FBN), Dummerstorf, Germany

[3] Institute of Animal Welfare Science, University of Veterinary Medicine, Vienna, Austria

**\* Correspondence:**
Liza R. Moscovice
moscovice@fbn-dummerstorf.de

**Word count**:

Abstract: **200**
Full text + references + table and figure text = **9735**
Figures: 3
Tables: 3

**Running title**: Spontaneous helping in pigs

**Abstract**

Helping behaviour is of special interest for prosociality because it appears to be

motivated by the needs of others. We developed a novel paradigm to investigate

helping in pigs (*Sus scrofa domesticus*) and tested 75 individuals in eight groups in their

home pens. Two identical compartments were attached to the pen, equipped with a

window, and a door that could be opened from the outside by lifting a handle. Pigs in all

groups spontaneously opened doors during a five-day familiarization. During testing,

each pig was isolated once from its group and placed in one of the two compartments,

in a counter-balanced order. In 85% of cases, pigs released a trapped group member

from the test compartment within 20 minutes (median latency = 2.2 minutes). Pigs were

more likely and quicker to open a door to free the trapped pig than to open a door to an

empty compartment. Pigs who spent more time looking at the window of the test

compartment were more likely to help. Distress signals by the trapped pig increased its

probability of being helped. Responses are consistent with several criteria for identifying

targeted helping, but results can also be explained by selfish motivations.

## 1. Background

Helping behaviour, in which an individual assists another to reach an otherwise
unachievable goal, is of special interest for the study of prosociality. Helping can be
costly (1), and often does not provide immediate benefits to the helper, but rather
appears to be motivated by an understanding of the needs of others, also referred to as
targeted helping (2). Examples of helping are wide-spread across animal taxa in the
wild, and include intervening on behalf of group members against predators (ants: (3);
primates: (4); whales: (5)), releasing individuals caught in snares, traps or sticky seeds
(ants: (6); wild boars (7); birds (1)) and defending group members from conspecific
attacks (elephants: (8); primates: (9)).

Animal helping is also the subject of experimental research to identify common
biological mechanisms underlying empathy, broadly defined as the ability to recognize
the emotional states of others and generate appropriate responses (2,10). Animal
helping and human empathic responses share several trait features (10), including that
both are mediated by the processing of social affective cues in similar brain regions
(rodents: (11), humans: (12)). Long-standing criticisms maintain that the available
evidence is not sufficient to support empathy-like motivations underlying animal helping,
and that more parsimonious explanations are possible (13–16). One major point of
contention is about whether helping is truly goal-oriented, which would require showing
not only that animals help in response to individuals in need, but also that observers
modify helping appropriately when the target's needs change or when they no longer
require help (17). Other criticisms highlight issues with the design of helping
experiments themselves, which can make it difficult to distinguish between prosocial
and selfish motivations for helping, for example when helping provides access to
desirable resources such as social partners (e.g. social reinforcement, 14,18), or novel
locations (19-20), in an otherwise barren test environment. Moreover, some studies
have assumed that their paradigms elicit distress without measuring responses of target
individuals (e.g. 19), leading others to question whether these individuals are even in

need of help (14,16,18). Interestingly, there is also evidence that high levels of distress

by individuals in need may sometimes hinder helping (21-22). More studies are thus

needed to disentangle the different motivations for helping behaviour (16).

We developed a novel helping paradigm that gives potential helpers and individuals in

need more behavioural flexibility to express a range of responses, in an attempt to

address some of these criticisms. Our paradigm is inspired by the work of Ben-Ami

Bartal and colleagues (19), in which rats could open a door to release a partner trapped

inside a restraint tube. However, our approach differs in several important aspects.

Firstly, subjects are tested in social groups in their home environment (e.g. (23)), which

improves ecological validity by giving animals the choice about whether, when and

whom to help, while allowing for competing motivations (e.g. exploring, feeding, resting,

social interactions with other group members). Secondly, subjects can simultaneously

perform the same behaviour in a range of helping as well as non-helping contexts,

which allows us to assess different motivations for door-opening. Finally, the trapped

individual is able to move around and explore and can also maintain visual contact and

interact with potential helpers through a mesh window, allowing us to investigate how

different behavioural responses of trapped individuals influence helping (24).

We tested the paradigm in pigs, which are a promising model species for studying the

mechanisms mediating helping behaviour. Pigs show evidence of emotional contagion

when observing group members in stressful situations (25), but without opportunities to

assist them. In addition, a case of spontaneous rescue behaviour was recently

documented in wild boars (*Sus Scrofa*), the ancestors of domestic pigs (7). However,

helping has not been systematically studied in pigs.

Our objectives were to determine whether pigs would spontaneously open a door to free

a trapped group member, to identify the factors that influence door-opening behaviour,

and to test whether pigs' responses were consistent with targeted helping or selfish

motivations. We followed the criteria suggested by Pérez-Manrique and Gomila (26) for

demonstrating targeted helping in animals: i) Helpers should show an other-oriented

reaction, including cognitive appraisal of the situation and a moderate level of arousal,

consistent with emotional regulation; ii) Helpers should show a flexible helping response

that is appropriate to the situation and iii) Helping should lead to an improvement in the

situation for the individual in need.

We predicted that pigs would selectively and consistently open doors to release trapped

group members, but not to access empty compartments. We further predicted that

potential helpers would be more likely and quicker to help individuals: i) if they were

more proficient in the novel door-opening behaviour, ii) if they were more attentive to the

trapped individual and iii) if the trapped individual expressed greater need to be

released, either by producing distress signals or by attempting to make social contact.

We measured changes in salivary cortisol (sCORT) in trapped pigs, to test whether

specific behavioural responses to being trapped were associated with greater activation

of the hypothalamic-pituitary-adrenal axis, consistent with a physical or psychological

stress response (reviewed in (27). Whenever the identity of helpers could be quickly

confirmed without requiring video analysis, we also measured their sCORT

concentrations soon after helping, to test whether helpers showed evidence for

increases in cortisol, consistent with emotional contagion of distress from trapped pigs,

or for more moderate levels of arousal, consistent with emotional regulation that can

facilitate targeted helping (21,26).

**2. Methods**

2.1 Animals and husbandry

All research procedures were approved under the German Animal Welfare Act (German

Animal Protection Law, §8 TierSchG) by the Committee for Animal Use and Care of the

Agricultural Department of Mecklenburg-Western Pomerania, Germany (permit LALLF

7221.3-1-036/20).

Subjects were German Landrace pigs born and housed at the Experimental Pig Facility

at the Research Institute for Farm Animal Biology (FBN) in Dummerstorf, Germany.

Pigs were tested in four temporally-separated cohorts, based on the five-week interval

between the births of new litters. Each cohort consisted of two social groups, for a total

of n = 78 subjects. At 28 days of age, subjects were weaned and mixed into new social

groups, by randomly selecting 3-5 pigs from each of two to three different litters to form

groups of 9-10 individuals. Groups were sex-balanced and contained equal numbers of

littermates from each contributing litter, such that each pig had either two (22% of group

members) or four (44% of group members) relatives in their group. Groups within a

cohort were housed in identical, neighbouring 2.8 × 2.8 m pens in the same room, with

partial slatted flooring and a sleeping mat. Throughout the study, pigs in neighbouring

pens did not have direct visual or physical contact but could exchange auditory and

olfactory cues. Pigs were regularly given back numbers using livestock spray paint to

facilitate individual identification. The room was equipped with multiple video cameras in

order to cover the entire home pen and two additional compartments that were attached

to the home pen for the experiment (see section 2.2). Pigs were tested between 46 – 55

days of age. Prior to testing, pigs were habituated to saliva sampling and had

participated in one social play experiment within their social group (28). Of the 78

subjects, one died of unknown causes, and two subjects from the same group were

removed from the study to be treated for illnesses before testing began. We report

results for the remaining n = 75 subjects who completed the study.

2.2 Familiarization, separation and testing

Two weeks after weaning, pigs went through familiarization, with the goal to expose

pigs to novel compartments and a novel door-opening mechanism. Each group within a

cohort had a one-hour familiarization session per day, with groups tested consecutively

in a counter-balanced order across five days. On each familiarization day, pigs were

presented as a group in their home pen with two identical compartments (90 × 90 × 80

cm in height) that were temporarily attached to opposite sides of the home pen (**Fig.**

**S1**). Compartments were open on top to facilitate placement of trapped pigs and video

recording during testing. Each compartment had a mesh window and a door with a

metal handle, placed at 20 cm height (**Fig. S2**). Opening the door required the use of

1.3 kg of force to lift the handle with the snout high enough to release an inner latch,

causing the door of the compartment to swing open into the home pen. This design is

suitable for pigs, who can exert strong force with their snout to dig or lift (29). During

each familiarization session, a group was presented simultaneously with the

compartments, with both doors closed. If a pig spontaneously opened a door, the door

was left open for two minutes so that pigs could freely explore the inside of the

compartment. At that point, any pigs still in the compartment were gently led out and a

researcher closed the door in a standardized way from outside the pen, using a hook to

insert the door handle back into its slot. If no pig had opened the door of a compartment

for five minutes, a researcher opened that door using the hook and left the door open

for two minutes, before removing any pigs inside the compartment and closing the door.

This approach gave all pigs within the group multiple opportunities to open doors within

and across familiarization days.

Each group then experienced the testing phase, with groups again tested consecutively

in a counter-balanced order across five days. The testing phase was initially similar to

the familiarization phase in that on each testing day the two compartments were

simultaneously attached to the home pen of one group, and pigs could freely explore

and open the doors ('pre-testing). The goal of pre-testing was to wait until pigs lost

interest in the doors and compartments, since re-attaching the compartments each day

drew attention to them. Researchers monitored the pigs from a different room using a

live video stream. Every ten minutes a researcher entered the room and if either of the

doors were opened, they led out any pigs inside the compartments and closed the

doors. Once the pigs had not touched or opened any door for ten minutes (this typically

occurred within 30 min of initial exposure on each test day), a researcher entered the

home pen and removed the designated trapped pig, using a pre-determined pseudo-

random order, so that each pig was trapped once over the course of group testing. This

began the 'separation' condition, during which the designated trapped pig was isolated

in a separate room in a 2.8 $m^2$ empty arena. The goal of the separation was to induce

moderate distress in the designated trapped pig, to ensure that trapped pigs would be

similarly motivated to be reunited with their group. Brief social separations can occur as

part of routine husbandry on farm, but such separations are known to be stressful for

pigs (28). During the separation, researchers monitored the test group remotely. If no

doors were opened during the first five minutes, the designated trapped pig was carried

back into the room by an experimenter and placed directly into the pre-designated test

compartment from above, thus having no contact with pigs in the home pen prior to

being placed in the test compartment. In the event that a pig opened any doors during

the initial 5-minute separation, a researcher entered the room, closed the open door(s)

and left the room. We then waited an additional five minutes before returning the

designated trapped pig and placing it in the pre-designated test compartment. As a

result, the total isolation time for each pig ranged from 5–11 minutes (median isolation

time = 5.8 minutes).

The test trial began once the trapped pig was placed in the test compartment ('testing').

The researcher left the room and monitored the pigs remotely using live video. During

the trial, the trapped pig maintained olfactory and auditory contact with their group

members at all times. In addition, the trapped pig could maintain visual and limited

physical contact with group members through the mesh window located on one side of

the compartment. Each test trial continued until either: i) A pig opened the door of the

test compartment, releasing the trapped pig, or ii) Twenty minutes elapsed without

helping, at which point a researcher entered the room and opened the test compartment

door to release the trapped pig. At any point during testing, pigs could also open the

door of the empty (control) compartment, but this had no influence on trial duration. If

the trapped pig showed signs of distress (screaming, squeals and/or escape attempts)

continuously for three minutes and was not let out by a pig, a researcher opened the

door, thus ending the trial for ethical reasons. This happened only once. The pigs in the

home pen were also monitored continuously via live video during each trial, and none

showed evidence of distress during testing. Once the door of the test compartment was

opened (either by a pig or by a researcher), the pigs were left undisturbed for an

additional 15 minutes and then hormone samples were collected (see section 2.4). The

compartments were then cleaned and approximately 1.5 hours after the first trial, the

same group experienced a second trial, with a second pig this time trapped in the

previously empty compartment. After the first group completed two trials, the second

group in the cohort went through the same procedure and we alternated group testing

order across test days. Within and across test days, we alternated the location of the

first test compartment for each group (at the front or side of the pen), so that each group

experienced both locations as test compartments with similar frequencies. At the

completion of the testing phase, each pig had been trapped once, and potential helpers

had between 6-9 opportunities (depending on their group size) to release different

members of their social group from the test compartment.

2.3 Behavioural measures

Behavioural data were coded from video recordings by two trained assistants using the

Observer software (version 15, Noldus, The Netherlands), with an inter-observer

reliability of 91.2%. During the familiarization phase, the daily latency for each door to

first be opened within each group was recorded as a measure of group task

performance across familiarization days. We also recorded each pig's daily frequency of

door-opening, as a measure of task proficiency.

During the separation we recorded the identity of any pigs who opened doors, which

door was opened (the designated empty or test compartment) and each pig's frequency

of door-opening. During testing, we recorded the amount of time that each potential

helper (referring to all other pigs in the group besides the trapped pig), spent looking in

the direction of the window of either compartment, defined as being within 10 cm of the

window with their head oriented towards the window. If a pig opened the door of either

compartment within 20-min, we recorded the identity of the door-opener, and their

latency to open the door, based on the time at which the latch of the door was

successfully lifted and the door swung open. When a potential helper released the

trapped pig from the test compartment within 20 minutes, they were designated as the

helper for that trial. We continued to record any door-opening of the empty compartment

up to 20 minutes, to directly compare pigs' motivation to open both compartments. We

used video cameras mounted directly above the compartment to record all occurrences

of squeals or screams by the trapped pig, which were identified as prolonged (> 2 sec)

vocalizations with high-frequency components and amplitude modulation (30). Squeals

or screams are commonly associated with negative affect in pigs (30,31). As further

validation that we were recording vocalizations related to negative affect, and to test

whether parts of the data analysis can be automated in future studies, we compared our

manual coding of screams and squeals with analysis of vocalizations using the

Stremodo software, which was developed for the automated detection of distress in pigs

((32), see also Supplementary materials). We also recorded each trapped pig's rate of

escape attempts (attempting to climb or jump against the side of the compartment) and

their investigations of either the window or the door, with investigations counted

whenever the snout of the trapped pig was within 5 cm of the object. For two pigs, the

compartment was not entirely visible in the video recording while they were trapped, so

behavioural data were analysed for the remaining n = 73 trapped pigs.

2.4. Hormonal measures

We assessed the physiological responses of trapped pigs by measuring changes in

their concentration of sCORT, which increases in response to physical or psychosocial

challenges in pigs (28). We collected saliva samples from the designated trapped pig in

their home pen before they were removed from the group (pre-test), and 15 minutes

after the door of the test compartment was open and the trapped pig was released

(post-test). During the post-test sampling, we also attempted to collect saliva samples

from the pig that helped when this could be quickly confirmed without additional video

analyses. Due to the time delay between exposure to a stimulus and peak cortisol

responses in pig saliva (33), the post-test samples should reflect free cortisol

concentrations around the time that pigs were released. Samples were collected

voluntarily from pigs using SalivaBio® Infant Swabs (Salimetrics, CA, USA). Cortisol

analyses were conducted at the FBN, using an ELISA kit for salivary cortisol

(Demeditec Diagnostics®, GmbH, Kiel, Germany) that our lab has previously validated

in pigs (28, see also supplementary materials). The inter-assay CVs of low- and high-

concentration pooled controls were 9.3% and 8.3% respectively (n = 16 samples). Intra-

assay CVs of low- and high-concentration pooled samples were 10.7% and 8.6%

respectively.

2.5 Statistical analyses

Data were analysed using the R statistical software (version 4.2.1, (34), see also

supplementary materials). Results are presented as medians and ranges unless

otherwise specified. When test predictors in models were significant, we conducted

pairwise post-hoc comparisons on marginal means, and calculated Tukey's HSD-

adjusted *p*-values (35). To measure door-opening performance during familiarization,

we fit a linear mixed model (LMM) testing whether the latency for each door to first be

opened was influenced by an interaction between familiarization day and location of the

compartment (front or side). We included daily counter-balanced test order of each

group (first or second) as a control predictor and group identity as a random effect. We

also determined the proportion of pigs that successfully opened a door on each

familiarization day and cumulatively across that day and previous days. To test whether

pigs had a side preference to open a specific compartment during familiarization, we

tested whether the proportional door-opening of each compartment was equal at the

group and individual pig level using one-sample binomial tests.

To compare door-opening in helping and non-helping contexts, we fit a generalized

linear mixed model (GLMM) to test for the n = 74 full trials, whether the likelihood to

open either door (response) was influenced by an interaction between condition

(separation or test trial) and the identity of the door (designated empty or test

compartment). We included the location of each compartment (front or side), the test

day (1-5), and the daily test order within a group (first or second) as control predictors,

and group identity as a random effect. Then, for the subset of separations (n = 35, 47%)

and test trials (n = 68, 92%) where at least one door was opened, we fit an LMM to test

whether the latency to first open a door (response) was influenced by the same test and

control predictors.

To test whether helping was influenced by potential helpers' previous experience and

their appraisal of the situation, we fit a GLMM with a zero-inflated negative binomial

error distribution. For each potential helper (n = 7-9 pigs) in each of the 75 trials, we

indicated whether that group member helped the trapped pig or not (response). We then

tested how each pig's likelihood of helping was influenced by their mean daily

proficiency in door-opening during familiarization and their proportion of time spent near

the window of the test compartment with their head oriented towards the window while a

pig was trapped. We considered this to be a measure of social attentiveness, since from

this position potential helpers should have a clear view of the trapped pig. We included

control predictors for the proportion of trial time that each potential helper spent looking

at the window of the empty compartment, their rate of opening either compartment

during the preceding separation, and whether they opened the empty compartment

during the test trial. We also included test day, daily test order within the group and the

potential helpers' relatedness to the trapped pig (sibling or non-sibling). We included

group identity as a random effect.

We tested for correlations between different behavioural responses of trapped pigs

using Pearsons correlation coefficients with Bonferroni corrections for multiple testing.

Rates of distress vocalizations (screams or squeals) and escape attempts were

correlated (see results), so we summed these two measures to obtain a combined rate

of distress signals. We fit a LMM to test whether changes in sCORT concentrations in

trapped pigs from pre- to post-test samples were related to their behavioural responses

while trapped, for n = 70 pigs for which we had complete data on their physiological and

behavioural responses to being trapped. The response was each trapped pig's salivary

cortisol concentrations. To model changes in sCORT in trapped pigs, we tested for two-

way interactions between sampling context (pre- or post-trapped, representing relative

changes in sCORT), and the following test predictors: i) each trapped pig's rate of

investigations to the window, where they could potentially make contact with group

members, ii) their rate of distress signals (the sum of their screams, squeals and escape

attempts), iii) the duration that each pig was separated from the group before being

trapped and iv) the duration that each pig was trapped. As control predictors, we

included each pig's sex, the test day and daily group testing order and the time that

each sample was taken. We included group and subject as random effects. We used

paired t-tests and Pearson correlation coefficients to compare sCORT concentrations of

trapped pigs and their helpers shortly after pigs had been released.

To test whether helpers' responses were sensitive to the expressions of need by the

trapped pig, we fit a Cox proportional hazards regression model using the Survival

package (v 3.2-7; (36)). In survival analyses, latencies and likelihoods of an event of

interest (in this case being released from the test compartment by a pig) are combined

into a hazard ratio (HR), with an HR above one indicating a covariate that is associated

with a greater likelihood and shorter latency for the event to occur. Trials in which pigs

were released by humans were right-censored, to indicate that the event of interest did

not occur during that pig's observation period. To compare behavioural data that varied

over time while ensuring that we did not violate the proportional hazards assumptions,

we stratified each trapped pig's behavioural data at one-minute intervals and removed

any intervals of less than 60-sec at the beginning of each pig's trapped period. We also

excluded from analysis each pig's behaviour during the last 10-secs before they were

released, to avoid committing "immortal biases" (37) by using information from the

future to predict the outcome. After removing n = 15 trapped pigs who were released in

under one minute, and the n = 2 trapped pigs mentioned above whose behavioural data

could not be analysed due to limited test compartment visibility, we ran the survival

analysis on n = 58 trapped pigs. We included each trapped pig's number of

investigations to the mesh window per minute, which could reflect motivation to acquire

information about the social group or to make social contact. The number of distress

signals (screams, squeals or escape attempts) per minute could not be transformed to

approximate normality. Therefore, we included the presence or absence of distress

signals at one-minute intervals as a binomial predictor. We included as control

predictors group identity, sex, the counter-balanced location of the test compartment (at

the front or the side of the pen), the group testing order and the time when the trapped

pig was first placed in the compartment. Diagnostic tools indicated that the proportional

hazards assumptions were not violated (global hazard ratio, $\chi 2$ = 22.07, df = 21, p =

0.40). The Concordance Index (CI) was 0.69, indicating good predictive discrimination

of the covariates. All data and code related to this study are available on-line at:

https://doi.org/10.5281/zenodo.8096753.

**3. Results**

3.1 Door-opening during familiarization

Despite having no exposure or training in the task beforehand a median of 81 (range =

77–93)% of pigs in each group opened doors on each familiarization day and by the end

of familiarization all but one of the 75 pigs had opened a door at least once (see **Fig.**

**S3**). The interaction between location of compartment and familiarization day tended to

influence door-opening latency (LMM, $F_{4,72}$ = 2.50, p = 0.051, see **Fig. S4 and Table**

**S1**). On day 3, pigs were faster to open the side compartment than the front

compartment (0.10 (0.01–0.48) min vs 1.35 (0.04–3.48) min, p = 0.004). Door-opening

latencies on other days were not affected by the location of the compartment (p values

> 0.19). Groups were also faster to open the side compartment on day 3 compared to

day 1 (0.70 (0.35–7.26) min) and day 2 (0.75 (0.18–1.45) min, p-values < 0.027, see

**Fig. S4**). There were no differences in latencies to open the front compartment across

familiarization days (p-values > 0.17). Pigs varied in their door-opening proficiency with

2 (0–11) daily door openings per pig, but only four (5%) showed evidence for a location

preference for door-opening, with three opening the side door more frequently (p-values

< 0.02), and one opening the front door more frequently (p = 0.035, see **Table S2**). At

the group level, there was no preference to open one of the two doors more frequently

during familiarization (binomial tests, p-values > 0.18, see **Table S2**).

3.2 Door-opening in non-helping and helping contexts

During the separation and testing conditions, potential helpers had opportunities to

perform the same door-opening behaviour in non-helping contexts, by opening either

compartment during the separation, before the trapped pig was placed in the

designated test compartment, or by opening the empty compartment during the test

trial. During the separation, 37 pigs (49% of potential helpers) opened doors, during 36

trials (48% of trials), with a median of 0 (range = 0- 5) door-openings per separation.

This accounted for the variable duration of isolation time (from 5 to 11 min) required to

meet the criteria that no door had been opened for five minutes before a test trial began.

During the test trials, 85% (n = 64) of the trapped pigs were helped by a group member,

with a latency of 2.2 (0.2–17) min to be released (See **Fig. S5**). Of the additional 11

pigs, ten were released by a researcher after the 20-minute trial duration ended and one

trial was aborted and the pig released for ethical reasons after exhibiting distress

signals for three minutes without being helped. In total, n = 33 pigs (44% of potential

helpers) helped a trapped pig, by being the quickest to first open the door of the test

compartment, and each helper helped a median of two trapped pigs (range = 1–4 pigs,

or 11–57% of their group members). During 68% (n = 50) of the full n = 74 test trials, n =

28 pigs (37% of potential helpers), opened the door of the empty compartment within

the 20-min maximum trial duration, with a latency of 3.9 (0.8–18.8) min to open the

empty compartment door.

The likelihood of door opening was influenced by an interaction between the condition

and the compartment (est ± SE = 1.50 ± 0.58, p = 0.008, see **Table S3** and **Fig. S6**).

Pigs opened the test compartment during 85 (63–100)% of test trials within their group,

while opening the empty compartment during 68 (40–100)% of test trials, p = 0.006). In

contrast, there were no differences in the likelihood of opening the two compartments

during the separation (21 (0–43)% vs. 30 (10–57)% of separations, p = 0.450). The

latency to first open a door was also influenced by an interaction between condition and

compartment (est ± SE = -0.97 ± 0.36, p = 0.012, see **Table 1** and **Fig. 1**). Pigs were

quicker to open the test compartment than the empty compartment during test trials (p<

0.001), but did not differ in their latencies to open either of the compartments during the

preceding separation period (p = 0.699, **see Fig. 1**). There was no effect of test day,

test order within a group, or location of each compartment (at the front or side of the

pen), on the likelihood or latency to open compartments (see **Table 1** and **Table S3**).

3.3. Behavioural and physiological responses of helpers and influences on helping

While pigs were trapped, potential helpers spent longer looking in the direction of the

test compartment window (7.7 (0.1–59.6) sec per trial min) than they did looking in the

direction of the empty compartment window (0.2 (0–4.4) sec per trial min, Wilcoxon

signed-rank test, n = 75, Z = 8.89, p < 0.001). Potential helpers who spent a greater

proportion of time looking in the window of the test compartment while a pig was

trapped inside were more likely to help the trapped pig (GLMM, est ± SE = 0.51 ± 0.08,

p < 0.001, see **Table 2** and **Fig. 2**). There was no relationship between proportion of

time spent looking in the window of the empty compartment and helping (see Table 2).

Pigs who were more proficient in door-opening across familiarization days were also

more likely to help their group members (est ± SE= 0.34 ± 0.13, p= 0.008). However, a

potential helper's rate of door opening during the preceding separation, and their

likelihood of opening the empty compartment during the test period, did not predict

helping (see Table 2). The potential helper's relatedness to the trapped pig, test day

and daily test order within a group also did not influence the likelihood of helping a

trapped group member.

In 67% of the helping trials (n = 43 out of 64), the helper initiated nose-to-nose or nose-

to-body contact with the previously trapped pig within one minute after opening the test

compartment door. In 81% of helping trials (n = 52), the helper also entered the test

compartment within one minute of helping. In 54% (n = 28) of these trials, the previously

trapped pig had already left the compartment, so entering the compartment did not lead

to social contact with the trapped pig.

For n = 50 trials where we were able to quickly identify the helper without requiring

additional video analyses, we obtained saliva samples from the helper as well as the

released pig fifteen minutes after helping. Helpers had lower sCORT concentrations

than their released group mate (Mean ± SD = 3.9 (± 1.9) ng/mL vs 7.3 (± 2.4) ng/mL,

$t$(49) = 8.14, p < 0.001, see **Fig. S7**), and there was no correlation between sCORT

concentrations of released pigs and their helpers shortly after release (Pearson, r(48)=

= 0.07, p = 0.63).

3.4 Behavioural and physiological responses of trapped pigs and influences on being

helped

Of the 73 trapped pigs for whom behavioural data could be analysed from video

recordings, 72 investigated the window and 71 investigated the door. Trapped pigs

investigated the window 4.3 (0 - 19) times per minute trapped, and they investigated the

door 1.6 (0- 11.6) times per minute. Rates of investigations to the window and door

were positively correlated (r(71) = 0.80, p < 0.001). The majority of trapped pigs (74%, n

= 54) attempted to escape and 44% of trapped pigs (n = 32) emitted screams or

squeals while trapped. Trapped pigs' rates of distress vocalizations and escape

attempts were positively correlated with each other (r(71) = 0.34, p = 0.02), but not with

either of the trapped pigs' investigatory behaviours (r(71) < 0.22, p values > 0.36). We

therefore summed rates of distress vocalizations and escape attempts to create a

combined measure of distress signals.

For n = 72 pigs with sufficient saliva volumes to measure sCORT before and after being

trapped, 87.5% (n = 63) exhibited increases in sCORT in their post-trapped samples

(Mean ± SD = 7.3 (± 2.7) ng/mL), compared to their samples before being trapped

(Mean ± SD = 4.7 (± 2.0) ng/mL, t(71) = 7.97, p < 0.001). Pigs with greater rates of

distress signals while trapped tended to have greater increases in sCORT (interaction:

est ± SE = -0.124 ± 0.061, p = 0.045, **Table S4** and **Fig. S8**). Changes in sCORT

concentrations while trapped were not related to a pig's duration of separation, their

duration of time trapped or other control predictors (see Table S4).

The Cox proportional hazards model indicated that trapped pigs who exhibited more

distress signals were 2.4 times more likely and quicker to be subsequently released,

compared with pigs who exhibited fewer distress signals ($\chi^2$ = 4.65, df = 1, p = 0.031,

**Table 3**, **Fig. 3 and Supplementary videos**). The rate of window investigations by

trapped pigs did not influence helping outcomes (see Table 3). There were also

significant differences among groups in the likelihood that trapped pigs would be helped

($\chi^2$ = 16.26, df = 7, p = 0.023, see Table 3). The other control predictors did not

influence the probability of being helped.

**4. Discussion**

Pigs spontaneously opened doors without training, and subsequently preferred to open

doors to free a trapped group member than to open doors to empty compartments. Pigs

who directed more attention towards the window of the compartment where a pig was

trapped were quicker to help, and distress signals from trapped pigs increased their

chances and decreased their latency to be helped. Our results support anecdotal

evidence for spontaneous rescue behaviour in wild boars (*Sus scrofa*), the ancestor of

domestic pigs (7). We avoid using the term 'rescue' here, since not all trapped pigs

exhibited signs of distress, and helping did not necessarily require a great cost or risk,

which are two of the criteria suggested by Hollis and Nowbahari (38) to define rescue

behaviour.

There was considerable variation in the likelihood that pigs helped trapped group

members, and in the latency for trapped pigs to be released, which enabled us to

investigate the factors mediating helping behaviour. Roughly half (44%) of the pigs in

each group helped others during the test phase. However, given that only one pig could

be a helper per trial, our group-testing paradigm may also underestimate the propensity

for helping behaviour at the population level. Pigs who opened doors at greater rates

during familiarization were more likely to help others, suggesting that their ability or

willingness to solve this novel task played a role in their helping behaviour.

Throughout the testing phase, pigs also opened doors in non-helping contexts, by

opening either door during the separation (49% of potential helpers), or by opening the

empty door during a test trial (37% of potential helpers). However, few of these

individuals (24% or 9 of 37 pigs who opened doors during the separation phase and

32% or 9 of 28 pigs who opened the empty door during a test trial) also opened the door

of the test compartment to help the trapped pig. As a result, pigs' door-opening in non-

helping contexts during the separation or test trial did not influence their likelihood of

opening the test compartment to help the trapped pig. Across the majority of

familiarization days and during separations, pigs had no preference to open one of the

two compartments. In contrast, during test trials pigs were more likely and quicker to

open the compartment containing a trapped group member than the empty

compartment, suggesting that door-opening during test trials was goal-oriented.

While a pig was trapped, potential helpers spent more time looking at the window of the

test compartment than they spent looking at the window of the empty compartment, and

the proportion of time spent looking at the test compartment window was the strongest

predictor of helping. Pigs who spent more time oriented towards the test compartment

window at close range may have been visually assessing the situation of the trapped

pig, which would be consistent with an other-oriented response (26). However, the

majority of trapped pigs produced vocal and locomotor distress signals that should have

been readily detectable without looking through the window, and the helpers' behaviour

is also consistent with more selfish explanations. As an alternative explanation, it is

likely that the movements and vocalizations of the trapped pig acted as salient features

to attract attention to the test compartment and window, and distressed pigs likely

produced more salient signals that were more effective in attracting attention. Potential

helpers may have then looked in the window to gather information about the situation

for themselves (e.g. 13,16). As a result, potential helpers would also be closer to the

door and handle, increasing the chances of door-opening through an effect of local

enhancement. But if local enhancement explained helping behaviour, we would expect

the majority of potential helpers to be similarly interested in and attentive to highly

salient cues from trapped pigs in distress, which was not the case (see Fig. 2 and

supplementary videos).

Distress signals given by the trapped pig, in the form of screams, squeals and escape

attempts, increased the pig's subsequent chances of being released by a 2.4 odds ratio.

This evidence is consistent with the interpretation of helping behaviour as a flexible

response that is appropriate to the situation of the trapped pig, which is another criteria

for targeted helping (26,39). It is important to note that distress signals were neither

necessary nor sufficient for helping to occur. Of the n = 15 pigs who were released in

under one minute and thus could not be included in the survival analysis, 67% (n = 10)

did not produce measurable distress signals before being released. That helpers

sometimes quickly freed these pigs counters arguments that spontaneous helping may

merely be motivated by selfish needs to alleviate the source of aversive distress signals

(24). However, it also raises the question of whether all trapped pigs were in need of

help (16). In addition, of the eleven pigs who were released by humans, nine (82%)

gave distress signals at some point while trapped, although (with the exception of the

aborted trial), these pigs gave distress signals at lower rates compared with pigs who

were released by conspecifics, as indicated by the survival analysis.

A primary concern of animal helping paradigms is that appropriate controls are often not

provided to test whether helping behaviour is motivated by self-interest. One of the main

examples is when helping provides potentially rewarding social interactions for helpers.

Indeed there is ample evidence that helping decreases when post-helping social

interactions are prevented (14,18,22), although such designs may also make it more

difficult for potential helpers to understand the causal effects of their actions. One of the

main advantages of testing animals in stable social groups in their home environment is

that this reduces the likelihood that helping is due to the desire for social reinforcement

more generally. Moreover, pigs were equally likely to help their siblings whom they had

known since birth as they were to help unrelated group members whom they had known

for a shorter time, which argues against helping motivated by social preferences. We

cannot rule out the possibility that the helper may have had an interest in interacting

with the specific trapped individual, which is supported by the fact that the majority of

helpers initiated social contact with the previously trapped pig within a minute of helping.

More work is thus needed to determine whether dyadic social preferences or

dominance relationships influence helping behaviour.

In the majority of cases (81%), pigs entered the test compartment immediately after

helping, although in most of these cases the trapped pig had already left the

compartment. In some rodent helping paradigms, similar proportions of helpers entered

a tube where conspecifics had previously been restrained (19,40). In those cases, it

was hypothesized that rodents may be motivated to enter the restraint tube because it is

novel (40), and may even prefer it to a barren testing arena (41). Such explanations are

unlikely in our paradigm, since pigs were tested in their home environment, had been

familiarized to the compartments prior to testing, and had equal opportunities to access

the empty compartment, but were less likely and slower to do so. It is more likely that

helpers entered the test compartment to gather relevant information for themselves after

witnessing the trapped pig, which is consistent with selfish motivations.

Wild boars reside in stable, cohesive social groups and even brief separations from their

social group can induce behavioural and physiological signs of stress in domestic pigs

(28,33). Either the separation alone, or the separation in combination with being in the

test compartment are likely to have been stressful for pigs. The majority of pigs had

increases in sCORT concentrations while trapped, and trapped pigs with greater

increases in sCORT exhibited more behavioural responses consistent with distress.

However, helpers did not show corresponding increases in sCORT concentrations, as

expected if they experienced emotional contagion of distress. In contrast, their sCORT

around the time of helping was similar to baseline cortisol concentrations of pigs prior to

being trapped. Although emotional contagion has been shown in pigs in non-helping

contexts (25), it does not appear to explain helping behaviour in this case. In fact, in

rodent studies, potential helpers who showed the greatest stress-reactivity to trapped

conspecifics were less likely to help than those who showed more moderate arousal

(21). Indeed, a moderate level of emotional arousal, consistent with emotional control

regulation, is one of the criteria for identifying empathic targeted helping in animals (26).

The cortisol responses of helpers in this study were more consistent with low rather

than moderate arousal, suggesting that helpers were not stressed by witnessing

trapped pigs. This is consistent with a previous study in which juvenile pigs did not show

emotional contagion of fear after hearing playbacks of distress vocalizations from

unfamiliar pigs (42). Further research should compare the physiological responses of

helpers and non-helpers to better assess how arousal relates to helping.

Given the promising evidence that potential helpers who attended more closely to the

test compartment window were more likely to help, follow-up research should test

whether a broader range of communicative signals by trapped pigs may mediate

helping. Two likely candidates are low-frequency grunt vocalizations and nose-to-nose

contacts, both of which are common socio-communicative behaviours in pigs (43).

Grunts and nose-to-nose contacts were observed during helping trials but could not be

systematically investigated with our recording equipment. Helping paradigms in which

individuals in need can express a broader range of behavioural responses are important

for clarifying the role of social communication in influencing helping and for further

distinguishing between self- and other-oriented explanations for helping behaviour in

animals.

**5. Conclusion**

This group-testing paradigm offers a novel experimental approach to systematically

study the proximate factors mediating spontaneous helping in a broad range of social

species, including farm animals. Applying the framework suggested by Pérez-Manrique

& Gomila (26) for evaluating comparative evidence for empathy, our results provide

partial support for targeted helping in that: i) helpers exhibited visual assessment of the

situation, ii) trapped pigs who signalled more distress were more likely and quicker to be

helped and iii) helping improved the situation for trapped pigs by reuniting them with

their social group. However, the lack of evidence for physiological arousal in helpers

argues against an other-oriented response (26). We also point out alternative

explanations based on local enhancement and selfish motivations to enter the test

compartment that can more readily explain why the majority of helpers entered the

compartment after helping, and why a small percentage of trapped pigs were helped

without showing obvious signs of distress. Thus, follow-up studies are needed to clarify

the motivations underlying spontaneous helping in pigs.

**Acknowledgements**

We thank V. Hörsch, B. Sobczak and E. Normann for their assistance with running the

experiments and collecting saliva samples. We thank P. Müntzel for conducting the

laboratory analyses. We are grateful to V. Hörsch and M. Wagenknecht for their help

with behavioural coding of videos. We thank M. Zenk for her help with the coordination

of this project and D. Ameling, H. Asmus, F. Hintze and H. Schumann for their daily

care of the pigs at the Experimental Pig Facility of the FBN. We are especially grateful

to T. Therneua, for his repeated advice that helped us to improve our survival analysis,

and for helpful feedback from two anonymous reviewers that inspired us to conduct

several additional analyses.

**Funding**

This research was funded in part by a Universities Federation of Animal Welfare grant

(44-19/20) to L.R.M.

**References**

1.    Hammers M, Brouwer L. Rescue behaviour in a social bird: Removal of sticky

"bird-catcher tree" seeds by group members. Behaviour [Internet]. 2017 Jan 1

[cited 2022 Dec 19];154(4):403–11. Available from:

https://brill.com/view/journals/beh/154/4/article-p403_2.xml

2.    De Waal FBM, Preston SD. Mammalian empathy: Behavioural manifestations and

neural basis [Internet]. Vol. 18, Nature Reviews Neuroscience. Nature Publishing

Group; 2017 [cited 2022 Dec 19]. p. 498–509. Available from:

https://www.nature.com/articles/nrn.2017.72

3.   Hollis KL, Nowbahari E. A comparative analysis of precision rescue behaviour in

sand-dwelling ants. Anim Behav. 2013 Mar;85(3):537–44.

4.   Boesch C. The effects of leopard predation on grouping patterns in forest

chimpanzees. Behaviour [Internet]. 1991 Jan 1 [cited 2022 Dec 19];117(3–

4):220–42. Available from: https://brill.com/view/journals/beh/117/3-4/article-

p220_5.xml

5.   Pitman RL, Deecke VB, Gabriele CM, Srinivasan M, Black N, Denkinger J, et al.

Humpback whales interfering when mammal-eating killer whales attack other

species: Mobbing behaviour and interspecific altruism? Mar Mammal Sci

[Internet]. 2017 Jan 1 [cited 2022 Dec 19];33(1):7–58. Available from:

https://onlinelibrary.wiley.com/doi/10.1111/mms.12343

6.   Nowbahari E, Scohier A, Durand JL, Hollis KL. Ants, Cataglyphis cursor, use

precisely directed rescue behaviour to free entrapped relatives. PLoS One. 2009

Aug 12;4(8).

7.   Masilkova M, Ježek M, Silovský V, Faltusová M, Rohla J, Kušta T, et al.

Observation of rescue behaviour in wild boar (Sus scrofa). Sci Rep [Internet].

2021 Dec 1 [cited 2022 Jan 26];11(1):16217. Available from:

https://doi.org/10.1038/s41598-021-95682-4

8.   Bates LA, Lee PC, Njiraini N, Poole JH, Sayialel K, Sayialel S, et al. Do elephants

show empathy? J Conscious Stud [Internet]. 2008 [cited 2023 Feb 2];15(10–

11):204–25. Available from: https://philpapers.org/rec/RNEDES

9.   Vogel ER, Fuentes-Jiménez A. Rescue behaviour in white-faced capuchin

monkeys during an intergroup attack: Support for the infanticide avoidance

hypothesis. Am J Primatol. 2006 Oct;68(10):1012–6.

10.   Decety J, Bartal IBA, Uzefovsky F, Knafo-Noam A. Empathy as a driver of

prosocial behaviour: Highly conserved neurobehavioural mechanisms across

species. Philos Trans R Soc B Biol Sci [Internet]. 2016 Jan 19 [cited 2020 Apr

28];371(1686). Available from: http://dx.doi.org/10.1098/rstb.2015.0077

11.   Meyza KZ, Bartal IBA, Monfils MH, Panksepp JB, Knapska E. The roots of

empathy: Through the lens of rodent models. Vol. 76, Neuroscience and

Biobehavioral Reviews. Elsevier Ltd; 2017. p. 216–34.

12.    Wu YE, Hong W. Neural basis of prosocial behaviour. Vol. 45, Trends in
712           Neurosciences. Elsevier Ltd; 2022. p. 749–62.

713    13.    Schwartz LP, Silberberg A, Casey AH, Kearns DN, Slotnick B. Does a rat release
714           a soaked conspecific due to empathy? Anim Cogn [Internet]. 2017 Mar 1 [cited
715           2020 Nov 3];20(2):299–308. Available from:
716           https://link.springer.com/content/pdf/10.1007/s10071-016-1052-8.pdf

717    14.    Silberberg A, Allouch C, Sandfort S, Kearns D, Karpel H, Slotnick B. Desire for
718           social contact, not empathy, may explain "rescue" behaviour in rats. Anim Cogn
719           [Internet]. 2014 [cited 2020 Nov 3];17(3):609–18. Available from:
720           https://link.springer.com/content/pdf/10.1007/s10071-013-0692-1.pdf

721    15.    Vasconcelos M, Hollis K, Nowbahari E, Kacelnik A. Pro-sociality without empathy.
722           Biol Letters. Vol. 8, 2012. p. 910–12.

723    16.    Blystad MH. An opinion on the interpretation of social release in rats [Internet].
724           Vol. 17, Biology Letters. Royal Society Publishing; 2021 [cited 2023 Jun 26].
725           Available from: https://royalsocietypublishing.org/doi/10.1098/rsbl.2021.0355

726    17.    De Wit S, Dickinson A. Associative theories of goal-directed behaviour: A case for
727           animal-human translational models. Psychol Res [Internet]. 2009 Jul 7 [cited 2022
728           Dec 19];73(4):463–76. Available from:
729           https://link.springer.com/article/10.1007/s00426-009-0230-6

730    18.    Hiura LC, Tan L, Hackenberg TD. To free, or not to free: Social reinforcement
731           effects in the social release paradigm with rats. Behav Processes. 2018 Jul
732           1;152:37–46.

733    19.    Ben-Ami Bartal I, Decety J, Mason P. Empathy and pro-social behaviour in rats.
734           Science (80- ) [Internet]. 2011 Dec 9 [cited 2021 Feb 10];334(6061):1427–30.
735           Available from: https://science.sciencemag.org/content/334/6061/1427

736    20.    Sato N, Tan L, Tate K, Okada M. Rats demonstrate helping behaviour toward a
737           soaked conspecific. Anim Cogn [Internet]. 2015 Sep 22 [cited 2020 Nov
738           3];18(5):1039–47. Available from:
739           https://link.springer.com/article/10.1007/s10071-015-0872-2

740    21.    Ben-Ami Bartal I, Shan H, Molasky NMR, Murray TM, Williams JZ, Decety J, et al.
741           Anxiolytic treatment impairs helping behaviour in rats. Front Psychol [Internet].

2016 Jun 8 [cited 2021 Jan 19];7(JUN):850. Available from:

http://journal.frontiersin.org/Article/10.3389/fpsyg.2016.00850/abstract

22.  Kalamari A, Kentrop J, Hinna Danesi C, Graat EAM, van IJzendoorn MH,

Bakermans-Kranenburg MJ, et al. Complex Housing, but Not Maternal

Deprivation Affects Motivation to Liberate a Trapped Cage-Mate in an Operant

Rat Task. Front Behav Neurosci [Internet]. 2021 Aug 26 [cited 2023 Jun

26];15:698501. Available from:

https://www.frontiersin.org/articles/10.3389/fnbeh.2021.698501/full

23.  Burkart JM, Allon O, Amici F, Fichtel C, Finkenwirth C, Heschl A, et al. The

evolutionary origin of human hyper-cooperation. Nat Commun [Internet]. 2014

Aug 27 [cited 2020 Nov 3];5(1):1–9. Available from:

www.nature.com/naturecommunications

24.  Carvalheiro J, Seara-Cardoso A, Mesquita AR, de Sousa L, Oliveira P,

Summavielle T, et al. Helping Behavior in Rats (Rattus norvegicus) When an

Escape Alternative Is Present. J Comp Psychol [Internet]. 2019 [cited 2022 Dec

19]; Available from: /record/2019-15237-001

25.  Goumon S, Špinka M. Emotional contagion of distress in young pigs is

potentiated by previous exposure to the same stressor. Anim Cogn [Internet].

2016 May 11 [cited 2019 Sep 9];19(3):501–11. Available from:

http://link.springer.com/10.1007/s10071-015-0950-5

26.  Pérez-Manrique A, Gomila A. The comparative study of empathy: sympathetic

concern and empathic perspective-taking in non-human animals. Biol Rev

[Internet]. 2018 Feb 1 [cited 2023 Feb 2];93(1):248–69. Available from:

https://onlinelibrary.wiley.com/doi/10.1111/brv.12342

27.  Sapolsky RM, Romero LM, Munck AU. How do glucocorticoids influence stress

responses? Integrating permissive, suppressive, stimulatory, and preparative

actions [Internet]. Vol. 21, Endocrine Reviews. 2000 [cited 2020 Jan 29]. p. 55–

89. Available from: https://academic.oup.com/edrv/article-

abstract/21/1/55/2423840

28.  Moscovice LR, Gimsa U, Otten W, Eggert A. Salivary Cortisol, but Not Oxytocin,

Varies With Social Challenges in Domestic Pigs: Implications for Measuring

Emotions. Front Behav Neurosci [Internet]. 2022 May 23 [cited 2022 Dec

19];16:899397. Available from: /pmc/articles/PMC9169876/

29.  D'Eath RB, Turner SP. The Natural Behaviour of the Pig. In: Marchant-Forde JN,

editor. The Welfare of Pigs [Internet]. Springer Netherlands; 2009. p. 13–45.

Available from: https://link.springer.com/chapter/10.1007/978-1-4020-8909-1_2

30.  Marx G, Horn T, Thielebein J, Knubel B, Von Borell E. Analysis of pain-related

vocalization in young pigs. J Sound Vib. 2003 Sep 18;266(3):687–98.

31.  Briefer EF, Sypherd CCR, Linhart P, Leliveld LMC, Padilla de la Torre M, Read

ER, et al. Classification of pig calls produced from birth to slaughter according to

their emotional valence and context of production. Sci Rep [Internet]. 2022 Dec 1

[cited 2023 Feb 2];12(1):3409. Available from: https://doi.org/10.1038/s41598-

022-07174-8

32.  Schön PC, Puppe B, Manteuffel G. Automated recording of stress vocalisations

as a tool to document impaired welfare in pigs. Anim Welf [Internet]. 2004 May 11

[cited 2023 Feb 2];13(2):105–10. Available from:

https://www.cambridge.org/core/product/identifier/S096272860002683X/type/jour

nal_article

33.  Ruis MAW, Te Brake JHA, Engel B, Buist WG, Blokhuis HJ, Koolhaas JM.

Adaptation to social isolation acute and long-term stress responses of growing

gilts with different coping characteristics. Physiol Behav. 2001 Jul 1;73(4):541–51.

34.  R Core Team. R: A language and environment for statistical computing [Internet].

R Foundation for Statistical Computing. Vienna, Austria; 2018. Available from:

https://www.r-project.org/

35.  Lenth R- emmeans: Estimated Marginal Means, aka Least-Squares Means. 2021

*https://CRAN.R-project.org/package=emmeans*.

36.  Therneau T. A Package for Survival Analysis in R. https://CRANR-

project.org/package=survival. 2023;

37.  Therneau T, Crowson C, Atkinson E. Using Time Dependent Covariates and Time

Dependent Coefficients in the Cox Model. R Proj https//cran.r-project.org ›

vignettes › timedep [Internet]. 2022 [cited 2023 Feb 6];1–8. Available from:

https://www.researchgate.net/publication/265041278_Using_Time_Dependent_C

ovariates_and_Time_Dependent_Coefficients_in_the_Cox_Model

38.    Hollis KL, Nowbahari E. Toward a behavioral ecology of rescue behaviour. Evol

Psychol [Internet]. 2013 [cited 2023 Feb 9];11(3):647–64. Available from:

www.epjournal.net

39.    Adriaense JEC, Koski SE, Huber L, Lamm C. Challenges in the comparative

study of empathy and related phenomena in animals. Vol. 112, Neuroscience and

Biobehavioral Reviews. Elsevier Ltd; 2020. p. 62–82.

40.    Silva PRR, Silva RH, Lima RH, Meurer YS, Ceppi B, Yamamoto ME. Are There

Multiple Motivators for Helping Behavior in Rats? Front Psychol. 2020 Jul 29;11.

41.    Ueno H, Suemitsu S, Murakami S, Kitamura N, Wani K, Takahashi Y, et al.

Rescue-like Behaviour in Mice is Mediated by Their Interest in the Restraint Tool.

Sci Rep [Internet]. 2019 [cited 2023 Jun 26];9(1):10648. Available from:

https://doi.org/10.1038/s41598-019-46128-5

42.    Düpjan S, Tuchscherer A, Langbein J, Schön PC, Manteuffel G, Puppe B.

Behavioural and cardiac responses towards conspecific distress calls in domestic

pigs (Sus scrofa). Physiol Behav. 2011 Jul 6;103(5):445–52.

43.    Camerlink I, Turner SP. The pig's nose and its role in dominance relationships

and harmful behaviour. Appl Anim Behav Sci. 2013 May 1;145(3–4):84–91.

835    **Tables and Figures.**

838    Table 1. Results of a LMM testing for an influence of condition (separation vs test trial)

839    and identity of compartment (designated empty or test compartment) on the latency for

840    pigs to first open a door. Significant predictors are indicated in bold.

| Predictor | Estimate | Std. Error | lower CI | upper CI | t or F statistic | df | *P* |
|---|---|---|---|---|---|---|---|
| (Intercept) | 0.342 | 0.297 | -0.248 | 0.943 | | | (†) |
| Condition_Test | 1.083 | 0.268 | 0.552 | 1.617 | | | (†) |
| Compartment_Test | 0.113 | 0.277 | -0.435 | 0.661 | | | (†) |
| Location compartment_Side§ | -0.116 | 0.161 | -0.435 | 0.202 | -0.72 | 1, 87.41 | 0.497 |
| Test day_2¥ | 0.089 | 0.24 | -0.386 | 0.566 | 5.362 | 4, 86.88 | 0.252 |
| Test day_3¥ | -0.073 | 0.25 | -0.567 | 0.423 | | | (Þ) |
| Test day_4¥ | -0.283 | 0.242 | -0.762 | 0.199 | | | (Þ) |
| Test day_5¥ | -0.454 | 0.268 | -0.987 | 0.082 | | | (Þ) |
| Daily test order in group_2nd# | 0.316 | 0.159 | 0.002 | 0.631 | 1.993 | 1, 86.49 | 0.061 |
| **Condition_Test : Compartment_Test** | -0.968 | 0.358 | -1.678 | -0.259 | **-2.701** | **1, 88.07** | **0.012** |

† Not indicated due to limited interpretability (of intercept, or when there is a significant
interaction)
Þ Refer to F statistic and p-value for 'Test day_2' for overall effect of predictor
§,¥,# Estimates refer to comparison with reference categories:
§ Location compartment_Front
¥ Test day 1
# Daily test order in group- 1st

853 **Table 2.** Results of a zero-inflated negative binomial GLMM predicting influences on

854 helping responses. Significant predictors are indicated in bold.

| Predictor | Estimate | Std. Error | lower CI | upper CI | $\chi^2$ | df | *P* |
|---|---|---|---|---|---|---|---|
| (Intercept) | -2.47 | 0.339 | -3.135 | -1.805 | | | (†) |
| **Proficiency during Familiarization‡** | 0.337 | 0.128 | 0.087 | 0.587 | **2.639** | **1** | **0.008** |
| Rate door-opening during Separation | 0.023 | 0.079 | -0.133 | 0.178 | 0.284 | 1 | 0.776 |
| **Proportion looking window-test compartment** | 0.521 | 0.074 | 0.377 | 0.665 | **7.078** | **1** | **< 0.001** |
| Proportion looking window-empty compartment | 0.015 | 0.138 | -0.256 | 0.285 | 0.106 | 1 | 0.916 |
| Opens empty compartment§ | 0.486 | 0.349 | -0.199 | 1.17 | 1.391 | 1 | 0.164 |
| Test day_2¥ | -0.031 | 0.405 | -0.825 | 0.764 | 0.430 | 4 | 0.980 |
| Test day_3¥ | 0.071 | 0.403 | -0.719 | 0.86 | | | Þ |
| Test day_4¥ | 0.157 | 0.388 | -0.604 | 0.919 | | | Þ |
| Test day_5¥ | -0.077 | 0.429 | -0.918 | 0.763 | | | Þ |
| Group daily test order_2nd# | -0.076 | 0.261 | -0.588 | 0.436 | -0.29 | 1 | 0.771 |
| Related to trapped pig¤ | -0.488 | 0.279 | -1.035 | 0.059 | -1.75 | 1 | 0.08 |

† Not indicated due to limited interpretability

‡ Data were square-root transformed

Þ Refer to chi-sq statistic and p value for 'Test day_2' for overall effect of predictor

§,¥,#,¤ Estimates refer to comparison with reference categories:

§ Did not open empty compartment during test trial

¥ Test day 1

# Daily test order in group- 1st

¤ Not related to trapped pig

872 **Table 3.** Results of a survival analysis predicting the probability for trapped pigs to be

873 helped. Hazard ratios are indicated by the exponentiated coefficients. Significant

874 predictors are indicated in bold.

| Predictor | coef | exp(coef) | se(coef) | robust se | $\chi^2$ | df | *P* |
|---|---|---|---|---|---|---|---|
| **Signals distress_Yes**[†] | 0.86 | 2.37 | 0.41 | 0.40 | **4.65** | **1** | **0.031** |
| Rate window investigations[‡] | -0.18 | 0.84 | 0.22 | 0.20 | 0.81 | 1 | 0.368 |
| Trapped pig sex_Female[§] | 0.05 | 0.95 | 0.34 | 0.33 | 0.03 | 1 | 0.868 |
| Time in compartment | 0.03 | 1.03 | 0.16 | 0.14 | 0.04 | 1 | 0.836 |
| Location compartment_Side[¥] | 0.39 | 1.48 | 0.34 | 0.34 | 1.37 | 1 | 0.242 |
| Test order in group | 0.15 | 1.16 | 0.16 | 0.16 | 0.91 | 1 | 0.341 |
| **Group identity_1B**[#] | 1.46 | 4.31 | 0.92 | 0.80 | **16.26** | **7** | **0.023** |
| **Group identity_2A**[#] | -0.14 | 0.87 | 0.67 | 0.72 | | | (Þ) |
| **Group identity_2B**[#] | 1.00 | 2.72 | 0.62 | 0.65 | | | (Þ) |
| **Group identity_3A**[#] | 0.51 | 1.66 | 0.64 | 0.69 | | | (Þ) |
| **Group identity_3B**[#] | 0.73 | 2.07 | 0.61 | 0.62 | | | (Þ) |
| **Group identity_4A**[#] | -0.04 | 0.97 | 0.62 | 0.67 | | | (Þ) |
| **Group identity_4B**[#] | 1.33 | 3.78 | 0.65 | 0.60 | | | (Þ) |

[†,§,¥,#] Estimates refer to comparison with reference categories:

[†] Trapped pig did not signal distress during the preceding interval

[§] Trapped pig sex_Male

[¥] Location compartment_Front

[#] Group Identity_1A

[‡] Data were square-root transformed

[Þ] Refer to chi-square statistic and p value under "Group Identity_1B" for the over-all

effect of the predictor

**Figure 1**. Latencies to open compartments during test trials, when a pig was trapped in

the test compartment, and during the separation, when both compartments were empty.

Data points are from separations and trials when at least one door was opened. Box

plots indicate medians and 25-75% interquartile ranges. Estimated marginal means and

SEs are also indicated, with results of pair-wise post-hoc comparisons (Tukey's HSD)

visualized using a compact letter display. Different letters indicate significant

differences. For full model results see Table 1.

**Figure 2.** Relationship between potential helpers' proportional time spent looking at the

window of the test compartment while a pig was trapped and their likelihood of helping,

averaged across their 6–9 exposures to trapped group members. Colours indicate

different social groups. The dashed lines depict the estimate and 95% CI from the fitted

model, with control predictors centred around 0. For full model results see Table

**Figure 3.** Survival curves showing that pigs who gave distress signals at any given

minute interval while trapped were more likely to be released than pigs who did not

(Hazard ratio = 2.4, p = 0.03). For full results, see Table 3.

905
906