Single-Soma Deep RNA Sequencing of Human Dorsal Root Ganglion Neurons Reveals

Novel Molecular and Cellular Mechanisms Underlying Somatosensation

Huasheng Yu[1], Dmitry Usoskin[2*], Saad S. Nagi[3*], Yizhou Hu[2], Jussi Kupari[2], Otmane Bouchatta[3], Suna Li Cranfill[1], Mayank Gautam[1], Yijing Su[1], You Lu[1], James Wymer[4], Max Glanz[4], Phillip Albrecht[5], Hongjun Song[1], Guo-Li Ming[1], Stephen Prouty[6], John Seykora[6], Hao Wu[7], Minghong Ma[1], Frank L Rice[5], Håkan Olausson[3#], Patrik Ernfors[2#], Wenqin Luo[1#]

1. Department of Neuroscience, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA.

2. Department of Medical Biochemistry and Biophysics, Division of Molecular Neurobiology, Karolinska Institute, Stockholm 17177, Sweden.

3. Department of Biomedical and Clinical Sciences, Linköping University, Linköping 58185, Sweden.

4. Department of Biochemistry and Molecular Biology, College of Medicine, University of Florida, Gainesville, FL 32603, USA.

5. Neuroscience & Pain Research Group, Integrated Tissue Dynamics, LLC, Rensselaer, NY 12144, USA.

6. Department of Dermatology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA.

7. Departments of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

* These authors contributed equally to this work

Correspondence:

hakan.olausson@liu.se (Håkan Olausson)

24   patrik.ernfors@ki.se (Patrik Ernfors)

25   luow@pennmedicine.upenn.edu (Wenqin Luo)

Abstract:

The versatility of somatosensation arises from heterogeneous dorsal root ganglion (DRG) neurons. However, soma transcriptomes of individual human DRG (hDRG) neurons – critical information to decipher their functions – are lacking due to technical difficulties. Here, we developed a novel approach to isolate individual hDRG neuron somas for deep RNA sequencing (RNA-seq). On average, >9,000 unique genes per neuron were detected, and 16 neuronal types were identified. Cross-species analyses revealed remarkable divergence among pain-sensing neurons and the existence of human-specific nociceptor types. Our deep RNA-seq dataset was especially powerful for providing insight into the molecular mechanisms underlying human somatosensation and identifying high potential novel drug targets. Our dataset also guided the selection of molecular markers to visualize different types of human afferents and the discovery of novel functional properties using single-cell *in vivo* electrophysiological recordings. In summary, by employing a novel soma sequencing method, we generated an unprecedented hDRG neuron atlas, providing new insights into human somatosensation, establishing a critical foundation for translational work, and clarifying human species-specific properties.

Introduction:

The somatosensory system conveys senses, such as temperature, touch, vibration, and body position[1]. Primary somatosensory neurons, which convert stimuli to electrical signals, are located in the dorsal root ganglia (DRG) and trigeminal ganglia (TG)[2]. They are greatly heterogeneous, composed of many different molecularly and functionally distinct populations[3]. Somatosensation is fundamental to our daily lives, but becomes a devastating human health problem when malfunctioning, such as during chronic pain and itch. Safe and effective drug options for chronic pain and itch are still limited[4-6], and the development of novel treatment strategies is greatly needed.

Most of our current knowledge about the mammalian somatosensory system has been obtained from model organisms, mainly rodents. However, the success rate of translating treatment strategies working in model organisms such as rodents to humans has been low[7,8]. There may be multiple reasons for the lack of success, but a noticeable one is species differences of the genetic, molecular and potentially even cellular makeup of somatosensory neurons between rodents and humans. Some significant differences between rodent and human DRG (hDRG) neurons have been noticed in previous studies, with neuropeptides, ion channels, and other markers not always matching between the species[9,10]. For an example, MRGPRX4, a bile acid receptor for human cholestatic itch, does not have a molecular ortholog in mice[11]. In contrast, TGR5, a receptor identified in mouse DRG neurons for bile acid-induced itch[12], is not expressed in hDRG neurons but instead in the surrounding satellite glial cells[11]. Thus, it is critical to elucidate molecular profiles and cell types of hDRG neurons for understanding human somatosensory mechanisms as well as for translational purposes.

Single-cell RNA sequencing (RNA-seq) is a powerful tool to qualitatively and quantitatively

95   study transcripts of individual cells (soma and/or nuclei)[13]. Based on the transcriptome, including

both transcript identities and expression levels, heterogenous cells can be classified into different

97   types[14]. This approach has been successfully used in varying degrees to study DRG and TG neu-

rons in mouse, macaque and other model organisms[15-18], providing comprehensive molecular and

cellular atlases for understanding somatosensory neurons and their differentiation. However,

there are some unique technical challenges for conducting single-cell RNA-seq of human DRG

or TG neurons: 1) Human tissues are more difficult to obtain compared to model organisms, and

the quality of human tissues (RNA integrity) is much more variable; 2) In human DRG/TG tis-

sues, non-neuronal cells, including satellite glial cells, fibroblasts, and other cell types, greatly

104   outnumber the neuronal cells[9,19]; 3) Satellite glial cells tightly wrap around neuronal somas[9,19],

making their physical separation difficult; 4) Human DRG/TG neuronal somas are much larger

than those of mouse DRG/TG neuronal somas, so they are prone to damage by enzyme digestion

and mechanical forces during single-cell isolation and not compatible with many current se-

quencing platforms. In addition, transcriptome changes caused by enzymatic and mechanical

109   dissociation may also affect cell clustering and cell type identification[20]. Due to these difficulties,

single-nucleus RNA-seq and 10x Visium spatial transcriptomics have been employed for human

111   DRG/TG neurons[21-24]. Despite novel insights from these studies, the quantity of transcripts in the

nucleus is much lower than those in the soma, and the nuclear transcripts may not represent the

113   full transcriptome profile of the whole cell[25], while the 10x Visium spatial transcriptomics lack

single-neuron resolution (Fig. 1A). Thus, it is necessary to develop a new method that enables

soma RNA-seq of hDRG neurons while preserving the single-cell resolution.

To meet this challenge, we developed a novel approach by combining laser capture

microdissection (LCM) for isolating individual neuronal somas and the Smart-seq2[18] for generat-

ing full length RNA-seq libraries (Fig. 1A). We sequenced 1066 hDRG neurons with minimum

satellite glial cell contamination from six lower thoracic and lumbar DRGs of three donors, de-

tecting an average of >9,000 unique genes per neuron (~ 3-5 times or more than the previous

single-nucleus RNA-seq results) and identifying 16 molecularly distinctive neuron types. Cross-

species analysis revealed both similarities but considerable differences among human, macaque,

and mouse DRG neurons. In addition, we uncovered a set of novel marker genes that can help to

distinguish different types of sensory neurons and afferents in hDRG and skin tissues. Based on

the molecular profiles, we also predicted novel response properties of human sensory afferents,

which were tested and confirmed by single-cell *in vivo* recordings. These results support a close

relationship between the molecular profiles uncovered by single-soma RNA-seq, histology, and

functional properties of human sensory afferents, highlighting the precision and unique utility of

this dataset in guiding functional studies of human somatosensory neurons. In short, we have

established a novel approach to conduct single-soma deep RNA-seq of hDRG neurons, which

revealed previously unknown neuronal types and functional properties. Given the high number

of unique transcripts recovered from each neuron, our dataset is especially powerful for molecu-

lar discovery, such as identifying potential high value novel drug targets. We believe that this

high-fidelity single-soma RNA-seq dataset will serve as a ground reference for homogenizing

RNA-seq data of human DRG/TG neurons using different approaches and for translating animal

studies into therapeutic applications.

Results

Development of an LCM-based approach for conducting single-soma deep RNA-seq of human DRG neurons

Adult human DRG neurons present considerable technical challenges for high-quality single-cell RNA-seq study. To overcome these hurdles, we developed a novel method that utilizes the laser capture microdissection to isolate individual neuronal soma and combine with Smart-seq2 full length RNA-seq library generation method for single-soma high-depth transcriptomic analysis (Fig. 1A). Six hDRGs at the low thoracic (T11-T12) and lumbar (L2-L5) levels from three post-mortem donors (donor information and screening criteria are summarized in the supplementary tables 1 & 2) were procured through NDRI (National Disease Research Interchange). DRGs were extracted and immediately frozen in OCT (within ~ 6 hours after death) at the NDRI pro-curement sites to minimize RNA degradation. Fresh frozen DRGs were cryo-sectioned, mounted onto LCM slides, and briefly stained with a HistoGene[TM] dye for cell visualization (Fig. S1A-C). Individual neuronal somas were discernable under microscope and dissected by a laser (Fig. 1A). Each detached soma dropped into a tube cap containing cell lysis buffer for library preparation (Fig. 1A). Dissected neuronal somas exhibited a similar size distribution as the whole DRG neu-ron population (Fig. S1D-E), suggesting no obvious sampling bias. Sequencing libraries were generated following the Smart-seq2 protocol[26]. In total, 1136 neuronal somas were dissected and passed through the final quality control for sequencing. During preliminary bioinformatic analy-sis, 70 samples with obvious glial cell contamination (dominant expression of APEO, FABP7, and other glia cell markers) were removed, and the remaining 1066 neurons were used for analy-sis in this study. 16 transcriptomic clusters of hDRG neurons were identified by Seurat[27] (Fig. 1B), with an average of 9486 genes detected per cell (Fig. 1C). No obvious batch effects or do-

nor differences were observed in the clusters (Fig. S2A-E). All these cells expressed high levels

of known peripheral sensory neuronal markers, *SLC17A6* (*VGLUT2*), *SYP* (Synaptophysin), and

UCHL1 (*PGP9.5*) (Fig. 1D, S2F-G).

Anatomical and molecular features of hDRG neuron clusters

Mammalian DRG neurons have some well-known physiological, anatomical, and molecular fea-

tures. DRG somatosensory afferents can be identified as A- or C- afferent fibers according to

axon conduction velocities[28]. A-fiber DRG neurons usually have large-diameter somas and

myelinated axons, while C-fiber DRG neurons have small-diameter somas and non-myelinated

axons. The A- and C- afferents can be further divided into peptidergic and non-peptidergic types,

based on the expression of one neuropeptide, calcitonin related polypeptide alpha

(*CALCA(*CGRP*))*[29]. To determine the broad types of the 16 hDRG neuron clusters, we analyzed

their soma sizes and expression of some key molecular markers (Fig. 1E). Clusters 1-16 were

arranged from small to large soma sizes. In addition, expression of neurofilament intermediate

filament (*INA*), which was enriched in small-diameter DRG neurons, and heavy chain (*NEFH*),

which was highly expressed in large-diameter DRG neurons, showed a clear complementary

pattern. The combined morphological and molecular features suggested that clusters 1-7 were

likely to be unmyelinated, small diameter C-fiber DRG neurons, whereas clusters 8-16 were

myelinated, large diameter A-fiber DRG neurons (Fig. 1E). The two groups could be further di-

vided based on the expression of *CALCA* (clusters 5-12). Moreover, the PR domain zinc finger

protein 12 (*PRDM12*), a transcriptional regulator critical for development of human pain-sensing

afferents (nociceptors) and C fibers[30], was expressed in clusters 1-12, further distinguishing be-

tween C- and A-fiber nociceptors versus A-fiber low-threshold mechanoreceptors (Fig. 1E).

Taken all information into consideration, clusters 1-4 were classified as non-peptidergic C-fibers,

clusters 5-7 as peptidergic C-fibers, clusters 8-12 as peptidergic A-fibers, clusters 13-15 as low-

threshold mechanoreceptors A-fibers (A-LTMRs), and cluster 16 as an unknown A-fiber type

(Fig. 1E).

Based on the expression profiles of top molecular markers, which had either unique expression

pattern or well-known functions (Fig. 1F, S2H), we named these 16 hDRG neuron clusters using

a nomenclature system with the following format (Fig. 1G): 1) The "h" at the beginning of each

cluster name indicated "human"; 2) mouse nomenclature was used for conserved subtypes (i.e.

most non-peptidergic neurons and A-LTMRs); 3) human peptidergic neuron types were desig-

nated as hPEP.(marker gene). Briefly, for non-peptidergic C-fiber neurons, cluster 1 was named

hTRPM8, cluster 2 was C-fiber low-threshold mechanoreceptors (hC.LTMR), cluster 3 was type

I non-peptidergic nociceptors (hNP1), and cluster 4 was type II non-peptidergic nociceptor

(hNP2). For peptidergic C-fiber neurons, cluster 5 was named hPEP.SST, cluster 6

hPEP.TRPV1/A1.1, and cluster 7 hPEP.TRPV1/A1.2. For peptidergic A-fiber neurons, cluster 8

was named hPEP.PIEZO^h (the superscript 'h' means 'high'), cluster 9 hPEP.KIT, cluster 10

hPEP.CHRNA7, cluster 11 hPEP.NTRK3, and cluster 12 hPEP.0 (no distinctive molecular

marker). For A-LTMRs, cluster 13 was named Aδ low-threshold mechanoreceptors

(hAδ.LTMR), cluster 14 Aβ low-threshold mechanoreceptors (hAβ.LTMR), and cluster 15 pro-

prioceptors (hPropr). Cluster 16 was named hATF3.

To validate the clusters identified by Seurat analysis, we independently analyzed our data using

the graph-based clustering Conos[31] package. Cluster structure revealed by Conos analysis repro-

duced the Seurat results (Fig. S3A). As a third method to validate identified clusters, we em-

ployed a neural network-based probabilistic scoring module[15,18] that learned human cell type

features based on their molecular profiles (Fig. S3B-D). Namely, the accuracy score of our Seu-

rat clustering assignment by the learning module was near 90% (Fig. S3B), which meant that

most cells were accurately assigned to their corresponding clusters, confirming the accuracy of

clustering (Fig. S3C). Moreover, the cell type consistency was validated by probabilistic simi-

larity (Fig. S3D). Thus, three independent analysis methods confirmed the robustness of the clus-

tering and strongly supported the cell type assignment.

Using Conos we also performed co-clustering and label propagation with a recently published

single-nucleus RNA-seq dataset of hDRG neurons by Nguyen et al[23] (Fig. S4). This analysis

showed that while some clusters displayed a one-to-one match between the two datasets, such as

hTRPM8, hPEP.SST, hPEP.KIT, and hAd.LTMR, most did not have one-to-one match. This

mismatch could be caused by biological variations such as nucleus vs cytoplasm RNA species

and quantity[25] and technical differences (the increased resolution obtained by deep sequencing in

the present study or variability caused by the different technology platforms[26,32]). This mismatch

also indicated that our dataset generated novel molecular profiles and cell types different from

the single-nucleus RNA-seq results.

Cross-species comparison of DRG neuron types

Comparison of hDRG neuron types to those in model organisms helps to uncover the evolutional

conservation and divergence of DRG neuron populations, provides clues about functions of

hDRG neuron populations, and identifies potential species-specific hDRG neuron populations.

Here we performed a cross-species comparison between our human dataset, a mouse dataset

(Sharma)[16], and a macaque dataset (Kupari, SmartSeq2 dataset)[18]. To identify corresponding

clusters between human and previously published mouse and macaque datasets, we used three

different strategies: Conos pairwise co-clustering followed by label propagation (Fig. 2A-B &

S5A-B), probabilistic neural network learning (Fig. S5C-D), and machine-learning based hierar-

chical clustering of an integrated dataset of human and macaque (Fig. S5E). For characteristics

and details of different approaches, see Method section. In all these analyses, human non-PEP

DRG neuron cell types showed high correlation to those of mouse and macaque, including

hC.LTMR, hNP1, hNP2, hPEP.SST (called NP3 in mouse and monkey), hTRPM8, hAδ.LTMR,

hAβ.LTMR, and hProprioceptor neurons (Fig. 2A-B & S5). hPEP.TRPV1/A1.1 and

hPEP.TRPV1/A1.2 corresponded to macaque PEP1 (Usoskin nomenclature) and mouse subclass

PEP1.4/CGRP-ϵ. This conclusion was supported by both Conos label propagation analysis (Fig.

2A-B) and probabilistic neural network analysis (Fig. S5C-D), suggesting that these clusters to

represent C-fiber thermoreceptors and nociceptors. Notably, four types of mouse C-fiber PEP

(CGRP) nociceptors have been evidenced[18,33] (nomenclature from Emory & Ernfors/Sharma),

but our analysis indicated that mouse clusters PEP1.1/CGRP-α, PEP1.2/CGRP-β, and

PEP1.3/CGRP-γ might not be evolutionarily conserved, as they did not have equivalent types in

the human datasets (Fig. 2A & S5C). hPEP.CHRNA7 showed high correlation to mouse

PEP2/CGRP-ζ and macaque PEP2, while hPEP.KIT corresponded to mouse PEP3/CGRP-η and

macaque PEP3, suggesting these clusters functionally belong to A-fiber nociceptors (Fig. 2A-B

& S5C-D). Interestingly, hPEP.NTRK3, hPEP.PIEZO^h, and hPEP.0, which made about half of

human PEP nociceptors that we sequenced, did not have a one-to-one corresponding cell types in

either mouse or macaque (Fig. 2A-B & S5C-D). Since macaque PEP2 cluster in the high-quality

Smart-Seg2 macaque dataset contained only three neurons, we focused our interpretation on

analyses on mouse homologs for this cell type. hPEP.NTRK3 showed the greatest similarity to

mouse PEP2 (CGRP-ζ) by Conos propagation analysis (Fig. 2A). Probabilistic neural network

learning revealed similar scores to both mouse PEP2 (CGRP-ζ) and PEP3 (CGRP-η) (Fig. S5C).

Overall, it therefore seems that hPEP.NTRK3 represents a convergent mouse PEP2/3-like cell

type. hPEP.PIEZO^h showed some similarity to mouse PEP3 (CGRP-η) and macaque PEP1 and

PEP3 in co-clustering (Fig. 2A-B). Neural network learning, and hierarchical clustering, indicat-

ing this cell type to represent an A-fiber mechanosensory nociceptor (Fig. S5C-E). Thus,

hPEP.NTRK3 and hPEP.PIEZO^h represents diverged PEP2- and PEP3-like A-fiber nociceptors,

which likely have emerged as human species-specific sensory neuron types. hPEP.0, a type of

human PEP A-afferents, showed no similarity to any mouse DRG neurons but some relationship

to macaque PEP1 and PEP3 (Fig. 2A-B), suggesting that they might be primate specific PEP

nociceptors. A schematic overview of our conclusions based on the above analyses regarding the

cell-type homologs across mouse-macaque-human is illustrated in Fig. 2C.

Transcription factors play critical role in DRG neuron development and differentiation[34]. Thus,

we performed a transcription factor associated gene regulatory network analysis (TF-GRNs) us-

ing machine learning to identify shared and species-specific TF-GRNs contributing to the simi-

larities and differences between sensory neuron subtypes and species (Fig. 2D). Evolutionarily

conserved TF-GRNs defining C-fiber nociceptors, A-fiber nociceptors, A-LTMRs, TRPM8, C-

fiber pruriceptors/nociceptors (hNP1, hNP2, hPEP.SST), and C-LTMRs were observed (yellow

boxes in Fig. 2D) as well as species-specific networks, such as for C-fiber nociceptors, hTRPM8,

hNP1, hNP2, hPEP.SST and hC.LTMR (green boxes in Fig. 2D). Among cross-species con-

served transcriptional regulators, some were previously known to drive sensory neuron diversifi-

cation in mouse, including *ZEB2* in C-fiber nociceptors[35], *SHOX2* in A-LTMRs[36,37], *RUNX3* in

proprioceptors[38], *FOXP2* in TRPM8, *RUNX1* in NP1[39], *ZNF52* and *POU4F2* in C-LTMRs[16,40].

These transcription factors may contribute to the formation of different DRG neuron cell types

and regulate the conserved and species-specific gene expression in each cell population.

Similarities and differences of top marker genes across species

To reveal molecular differences among the corresponding cell types in human, macaque

(Kupari)[18], and mouse (Sharma)[16], we selected the top 10 marker genes from each hDRG neuron

population and compared them across species (Fig. S6). In general, the expression patterns of

these genes were more similar between human and macaque than between human and mouse,

reflecting the evolutionary distances between rodent, non-human primate, and human. Some

genes were expressed in the corresponding populations across all three species. For example,

TRPM8 was expressed in C-fiber cold-sensing neurons, and *IL31RA* was expressed in nocicep-

tive/pruriceptive population (NP3) as well as its corresponding human population hPEP.SST.

Some genes, such as *KCNV1*, a voltage-gated potassium channel, were specific for primate

Aδ.LTMR but had low or no expression in the corresponding mouse DRG neurons (Fig. S6).

Moreover, some marker genes were specific only for hDRG neuron types. For example,

Mechanosensory Transduction Mediator Homolog (*STUM*) was specifically expressed in

hTRPM8, and calsequestrin 2 (*CASQ2*) was specifically expressed in hC.LTMR (Fig. S6). Genes

specifically enriched in hDRG neurons may confer unique physiological and functional proper-

ties of the human somatosensory system. The observed differences from the top 10 marker genes

represent only the iceberg tip of the overall molecular differences between human and model

organism DRG neurons, highlighting the necessity of validating molecular targets in hDRG neu-

rons for translational studies.

Molecular marker expression and validation of C-fiber pruriceptors, thermoreceptors and

nociceptors

Based on the sequencing results, we established specific marker genes or their combination to

identify each type of hDRG neurons (Fig. S7A) and validated their expression using

RNASCOPE multiplex fluorescent *in situ* hybridization (FISH) (Fig. S8). We also deduced po-

tential functions of each hDRG neuron type based on the cross-species cell type comparison and

expression of molecules with known sensory functions.

hNP1 and hNP2 exclusively expressed *MRGPRX1* (Fig. 3A), a Mas-related GPCR family mem-

ber that could be activated by various pruritogens[41]. Similarly, primate specific bile acid receptor

MRGPRX4 for mediating human cholestatic itch[11,42] were detected by our single-soma dataset.

Enrichment of *MRGPRX4* and *MRGPRX3* (an orphan GPCR in the same family) in hNP2 but not

hNP1 helped to separate these two clusters (Fig. S7A). Other itch-related receptors, such as

HRH1 and *IL31RA* (Fig. 3A-B), were also expressed in both hNP1 and hNP2, suggesting that

these two populations function to detect various pruritogens and transmit the sensation of itch.

The co-expression of *MRGPRX1* and *HRH1* was validated by multiplex FISH (Fig. 3A). *PIEZO2*

was expressed at a higher level in hNP1 than in hNP2 neurons (Fig. S7B), suggesting that hNP1

neurons could be more mechanosensitive. Consistent with this molecular feature, recordings

from human afferents have revealed that some human histaminergic itch-sensing fibers are

mechanosensitive[43]. In mice, NP1 and NP2 afferents are well-characterized by specific expres-

sion of two different *Mrgpr* members: NP1 neurons (~20% of total DRG neurons) highly express

Mrgprd[44], while NP2 neurons (~5% of total DRG neurons) express *MrgprA3*[15]. In humans, how-

ever, *MRGPRD* was only expressed in a few NP1 neurons (Fig. S9A), so it was less useful for

marking human NP1 population, and *MRGPRA3* gene does not have an orthologue in the human

genome. In short, NP1 and NP2 populations likely have conserved physiological functions in

mediating itch sensation in both mice and humans. However, some key molecular receptors for

detecting pruritogens are different between the species, which may reflect evolutionary adapta-

tion to distinctive pruritogens encountered by human and mice in their living environments.

The hPEP.SST population displayed specific expression of the neuropeptide, somatostatin (*SST*),

and an enriched expression of *GFRA3,* a co-receptor of tyrosine kinase RET (Fig. 3B, S7A). We

also found another neuropeptide cholecystokinin (*CCK*)[45] enriched in this population (Fig. S7A).

The hPEP.SST cluster corresponded to mouse and macaque NP3 population (Fig. 2), which are

also marked by the expression of *SST*[15,16]. Given the previously established role of mouse NP3

neurons in itch sensation[46] and the high expression of itch-sensing receptors, such as *HRH1*,

IL31RA (Fig. 3A-B), hPEP.SST afferents could also mediate itch sensation, especially under

inflammatory conditions[15]. Neither *PIEZO1* nor *PIEZO2* (Fig. S9A) was detected in hPEP.SST

neurons, indicating that these afferents might not be mechanosensitive. Indeed, some human

histaminergic itch-sensing fibers are insensitive to mechanical forces[43]. A human feature of

hPEP.SST neurons was the co-expression of the peptidergic neuron marker, CGRP, which was

barely detected in the corresponding mouse NP3 neurons (Fig. S9B)[16,18].

The hTRPM8 population was distinguished from other cell types by the specific expression of a

novel molecular marker, *STUM,* and high-level expression of *TRPM8* (Fig. 3C). Almost all

STUM[+] neurons expressed *TRPM8,* which was validated by multiplex FISH (Fig. 3C). Since

TRPM8 is a receptor for cold temperature and cooling chemicals (such as menthol)[47-50], the

hTRPM8 population likely functions as cold- and menthol-sensing afferents. Notably, this newly

identified maker gene *STUM* was not detected in mouse *TRPM8[+]* neurons[16,33]. In macaque,

STUM was more broadly expressed in several clusters (Fig. S9C)[18]. Nevertheless, other molecu-

lar markers, such as *FOXP2* and *GPR26*, were shared among mouse[16,33], macaque[18], and human

TRPM8 cold-sensing neurons (Fig. S9A-C). Intriguingly, some hTRPM8 neurons also expressed

a low level of heat-sensing receptor *TRPV1* (Fig. 3D), suggesting that these neurons might also

be activated by heat stimuli. Consistently, human physiological recordings have identified cold-

sensitive C-fibers that also respond to heat[51]. Thus, some of the neurons in the hTRPM8 popula-

tion are likely to be polymodal cold-sensing afferents.

Two peptidergic C-fiber clusters displayed overlapping high expression of *TRPV1* and *TRPA1*

(Fig. 3D), which were therefore named as hPEP.TRPV1/A1.1 and hPEP.TRPV1/A1.2. Since

TRPV1 is activated by noxious heat and capsaicin, and *TRPA1* can be activated by noxious cold

and various noxious chemicals[52,53], they are likely to be C-fiber peptidergic thermoreceptors and

nociceptors, sensing noxious thermal and chemical stimuli and transmitting pain signals. From

the cross-species comparison, hPEP.TRPV1/A1.1 and hPEP.TRPV1/A1.2 were mostly similar to

macaque PEP1 and mouse subclass PEP1.4/CGRP-ε (Fig. 2A-C). The exact functional differ-

ences between these two quite similar populations are yet to be established, but it is tempting to

hypothesize that hPEP.TRPV1/A1.1 innervates the skin while hPEP.TRPV1/A1.2 innervates

deep tissues and visceral organs, because $CGRP^+$/$TRPV1^+$ afferents were observed in both the

human skin and deep tissues[54]. In contrast to the hPEP.TRPV1/A1.1 population,

hPEP.TRPV1/A1.2 neurons expressed PROKR2, and higher level of PTGER3, and prostaglan-

din I2 receptor (PTGIR) (Fig. S7E & S9A). These molecular markers have been found to be en-

riched in mouse viscera-innervating DRG neurons[55,56].

Molecular marker expression and validation of A-fiber peptidergic nociceptors

Five clusters of peptidergic A-fiber nociceptors were identified in our dataset. When compared to

mouse and macaque peptidergic populations, some clusters showed the greatest divergence (Fig.

2A-C), indicating that they might contain human specific sensory neuron types.

The hPEP.PIEZO^h cluster was named because they expressed the highest number of *PIEZO2*

transcripts among all PEP clusters (Fig. 4A), an expression level comparable to the most

mechanosensitive afferents, hC.LTMR and hAδ.LTMR. The hPEP.PIEZO^h neurons also ex-

pressed a relatively high levels of *PIEZO1* transcripts, though the overall expression of *PIEZO1*

in hDRG neurons was low (Fig. S9A). This cluster could be identified by its high expression of

PTGER3 but not *TRPA1* (Fig. 4A & 3D). No mouse DRG neuron population was highly

matched for the hPEP.PIEZO^h cluster. Nevertheless, the expression of a few unique marker

genes in this cluster could provide clues into its functions. The adrenoreceptor, *ADRA2C*, a spe-

cific molecular marker found in human sensory fibers innervating blood vessels[57-60], was mainly

detected in this cluster (Fig. S9A). In addition, the hPEP.PIEZO^h population expressed *GPR68*, a

membrane receptor sensing flow and shear forces in the vascular endothelia cells[61] (Fig. S9A).

Given the well-established functions of *PIEZO1* and *PIEZO2* in mediating neuronal sensing of

blood pressure and the baroreceptor reflex[62] and the expression of known function of *ADRA2C*

and *GPR68*, we proposed that some hPEP.PIEZO^h afferents innervate blood vessels and sense

the blood pressure or flow. This cluster also expressed a high level of *PTGIR* (Fig. S9A). Mouse

PTGIR$^+$ DRG neurons innervate the bladder[55], and *PIEZO2* expressed in these neurons is re-

quired for sensing the bladder pressure to coordinate urination[63]. Thus, some hPEP.PIEZO^h af-

ferents might also innervate bladder and play a similar role. Take all into consideration,

hPEP.PIEZO^h neurons might function in sensing mechanical forces from blood vessels and in-

ternal organs. Given that there was not a clear mouse DRG CGRP$^+$ population with high *PIEZO2*

expression from single-cell RNA-seq datasets, we speculated that hPEP.PIEZO^h neurons are

either human specific or greatly expanded in humans. A fundamental difference between human

and mouse is their body sizes (humans are more than 2000 times larger than mice[64]). This addi-

tion or expansion of hPEP.PIEZO^h neurons is likely to meet the challenge of mediating sensation

from internal organs and blood vessels in much larger human bodies.

The hPEP.KIT cluster had the specific expression of a receptor tyrosine kinase, *KIT*, and a medi-

um to low expression level of *PIEZO2* (Fig. 4B). In mouse DRG neurons, *Kit* is more broadly

expressed and found in four peptidergic clusters[16] (Fig. S9B). Cross-species analysis suggest that

this cluster mainly corresponded to the mouse PEP3/CGRP-η and macaque PEP3 population

(Fig. 2A-C), which are A-fiber high threshold mechanoreceptors (HTMRs), forming circumfer-

ential endings around hair follicles and mediating hair pulling pain[65,66]. A more recent study sug-

gests that mouse CGRP-η neurons mediates mechanosensation from the distal colon[67]. Thus, the

hPEP.KIT cluster likely functions as a population of fast-conducting mechano-nociceptors.

The third peptidergic A-fiber cluster, hPEP.CHRNA7, featured abundant expression of *CHRNA7*

but almost no expression of *PIEZO2* (Fig. 4C). Cross-species analysis revealed that this cluster

corresponds to mouse PEP2/CGRP-ζ and macaque PEP2 populations (Fig. 2A-C), which are also

marked by the unique expression of *CHRNA7*[16,18]. Interestingly, this cluster also expressed

PVALB (Fig. 1F), a molecular marker commonly used for proprioceptors in mouse and human.

Retrograde tracing from the mouse gastrocnemius muscle labeled *CHRNA7*+ DRG neurons[55],

suggesting that hPEP.CHRNA7 may contain muscle-innervating A-fiber nociceptive sensory

afferents.

hPEP.NTRK3 is a population of peptidergic A-fibers with high expression of *NTRK3* and

S100A4 (Fig. 4D). Neurons in this cluster expressed a low level of *PIEZO2* (Fig. 4B). Finally,

hPEP0 is a population of peptidergic A-fibers that expressed *CALCA* and a moderate level of

PIEZO2 but lack of other specific marker genes. Potential function of hPEP.NTRK3 and hPEP0

are currently unclear. They could be some types of A-fiber mechano-nociceptor[68].

Molecular marker expression and validation of C-LTMRs, A-LTMRs and an ATF3 popu-

lation

hC.LTMR is the putative human C-tactile nerve fibers. All cross-species neuron cluster compari-

son methods unequivocally identified C-LTMRs as conserved across all three species. Neverthe-

less, the specific marker gene *CASQ2* for hC.LTM (Fig. 5A) is not detected in either mouse or

macaque C-LTMRs[15,18]. Conversely, mouse C.LTMR cells are characterized by exclusive ex-

pression of tyrosine hydroxylase (TH) and *SLC17A8* (*VGLUT3*)[15] (Fig. S9B), which are barely

expressed in hDRG neurons (Fig. S9A). Thus, the molecular markers to identify C.LTMRs are

different between human and mouse. On the other hand, human, mouse, and macaque C.LTMRs

all had conserved expression of GFRA2, another co-receptor of RET, and a zinc finger transcrip-

tion factor ZNF521 (Fig. S9A-C). Multiplex FISH confirmed that *CASQ2*[+] cells expressed high

levels of *PIEZO2* (Fig. 5A). hC.LTMRs likely mediate innocuous affective touch sensation[70-72].

A-LTMRs were featured by large diameter somas and high expression of *NTRK2* and *NTRK3*

(Fig. 5B-C) but a lack of expressions of *SCN10A* and *PRDM12* (Fig. 4C & 1E), two genes highly

associated with human nociception[30,73,74]. We identified 4 clusters in this category. hAδ.LTMR

was named based on its high expression level of *NTRK2* and *PIEZO2*, but low level of *NTRK3*

(Fig. 5A-C), a molecular feature similar to the mouse Aδ-LTMRs (Fig. S9B). *KCNV1* was en-

riched in this cluster and could serve as a novel molecular marker for identifying this population

in hDRG neurons (Fig. 5B). hAβ.LTMR, for tactile touch, expressed higher level of *NTRK3* but

a lower level of *NTRK2* compared to hAδ.LTMRs (Fig. 5C). hPropr, for limb position sensing,

expressed a high level of a proprioceptor marker *PVALB* and *REEP5* (Fig. 5D).

We also identified a cluster named as hATF3 (Fig. 5E), which contained mainly large diameter

neurons and strongly corresponded the "unknown" cluster first identified in normal male mice by

the Sharma RNA-seq dataset[16]. This cluster in both human and mouse datasets showed a very

conserved molecular profile (Fig. S7A). One of the markers it expressed, *ATF3*, is a transcrip-

tional factor associated with sensory afferent injury, indicating that these cells may represent

neurons that are undergoing (or previously underwent) stress/damage/insult (Fig. 5E). $ATF3^+$

neurons also expressed high level of neuropeptide *ADCYAP1* (Fig. 5E). Since there were no

medical records indicating obvious nerve injuries in our human donors, and since the mouse data

came from naïve mice, we speculate that these neurons might reflect a very low level of sporadic

sensory afferent injury accumulated from daily lives. On the other hand, we cannot exclude the

possibility that $ATF3^+$ cluster represent a population of normal DRG neurons.

The single-soma deep RNA-seq dataset provides novel insights into the molecular and cel-

lular mechanisms underlying human sensation of itch and pain

Given the high number of unique transcripts per neuronal soma, our dataset is uniquely powerful

for molecular discovery. We identified a much higher number of membrane proteins, such as

GCPRs, ion channels, chemokine receptors, as well as neuropeptides, compared to 10x Visium

Spatial RNA-seq or single-nucleus RNA-seq datasets (Fig. 6A). These discoveries provided nov-

el insights into understanding molecular and cellular bases of physiological recording and psy-

chological experiment results regarding human itch and pain sensation.

Physiological recordings have identified at least two groups of C-fibers responding to different

pruritogens. One group responds to cowhage, a plant triggering intense itch in human[75], and

comprises mechano-sensitive polymodal units[43]. The other group responds to histamine with

sustained discharges but is mechanically insensitive[43]. However, the molecular and cellular bases

for these observations remain unclear. To understand these physiological properties, we analyzed

the expression of histamine and cowhage receptors, and the mechanoreceptor *PIEZO2* in the

three potential itch populations, hNP1, hNP2 and hPEP.SST. *PIEZO2* was highly expressed in

hNP1, with low expression in hNP2 and almost no detectable expression in hPEP.SST (Fig. 5A).

Protease-activated receptor *F2RL1*, the receptor mediating cowhage induced itch in humans[75],

was exclusively expressed in hNP1 (Fig. 6B). Thus, the hNP1 population likely contained C-

fiber pruriceptive afferents sensitive to both cowhage and mechanical stimuli. For histamine re-

ceptors, *HRH1* was expressed in all three itch populations, *HRH2* was expressed in hNP2 and

hPEP.SST but not in hNP1, *HRH3* had low expression in hNP1 and hPEP.SST, while *HRH4* was

not detected (Fig. 6B-C). Thus, hNP2 and hPEP.SST clusters are good candidates for histamine-

sensitive but mechano-insensitive itch-sensing C-fibers.

When intracutaneously applied into human skin, some chemicals tend to trigger itch, such as

histamine and prostaglandin E2 (PGE2), while others preferentially induce pain, such as seroto-

nin and bradykinin[76]. To explore the potential underlying mechanisms, we examined the expres-

sion of all known receptors for these chemicals in our dataset. We found that the receptors of

itch-inducing chemicals are enriched in itch sensing populations. For example, histamine recep-

tors *HRH1* and *HRH2*, and PGE2 receptor *PTGER2* were enriched in hNP1, hNP2 and

hPEP.SST putative itch-sensing neurons (Fig. S10A1). In contrast, serotonin receptors, *HTR1B*

and *HTR1F,* and bradykinin receptor *BDKRB2* were enriched in putative nociceptive populations

(Fig. S10A2). The differential expression patterns of these receptors might explain the different

sensory experience induced by these chemicals.

Utility of single-soma deep RNA-seq dataset for discovering novel drug targets and for ob-

taining insights into molecular targets of existing drugs

Our deep sequencing dataset can also provide novel insights into the mechanisms of existing

clinical drugs or for identifying potential novel drug targets. The expression patterns of promis-

ing drug targets, such as GPCRs, ion channels, chemokine and cytokine receptors, and neuropep-

tides, were analyzed (Fig. S11-S14). Here we analyzed itch-related receptors and molecules as an

example. A series of itch-sensing receptors have been identified in model organisms[3,46,79], but

which of these targets can be translated remains to be determined. A subset of these itch recep-

tors were indeed enriched in hNP1, hNP2 and hPEP.SST populations, such as the chloroquine

receptor *MRGPRX1*, bile acid receptor *MRGPRX4*, histamine receptor *HRH1*, leukotriene recep-

tor *CYSLTR2*, and interleukin receptors *IL31RA* and *OSMR* (Fig. 6B)[16,18]. However, some sus-

pected itch receptors did not exhibit expression enrichment in human itch populations. For ex-

ample, *HRH4* was reported to mediate histamine-dependent itch in mice[80], but expression of

HRH4 was barely detected in human itch-sensing populations (Fig. 6C), suggesting that *HRH4*

might not be directly involved in human histaminergic itch. The same was found true for *IL7R*

and *TLR7*[79], which were proposed to mediate non-histaminergic itch in mice (Fig. 6C). In addi-

tion to these known players, we identified other membrane receptors and signaling molecules in

human itch populations (Fig. 6D), including MRGPRX3, EDNRA, PTGDR, HTR3A, CHRNA3,

KCNG4, and PLCB3, which could represent novel anti-itch targets.

Gabapentin was originally developed for treating epilepsy and more recently used in the treat-

ment of neuropathic pain and chronic itch[81]. It inhibits neurotransmitter release by acting on α2δ-

1 and α2δ-2 voltage-dependent calcium channels *CACNA2D1* and *CACNA2D2*[81]. We found that

both receptors were broadly expressed in hDRG neurons (Fig. S10B), suggesting that one poten-

tial mechanism by which gabapentin could provide clinical benefit is through inhibiting synaptic

transmission of primary afferents, as has been shown in mice[82].

Opioids and derivatives activate opioid receptors to modulate pain and itch. Agonists of the μ-

opioid receptor (*OPRM1*) alleviate pain but elicit itch in humans and model organisms, whereas

agonists of the κ-opioid receptor (*OPRK1*) inhibit itch in humans[83,84]. Unlike mouse DRG neu-

rons, which did not5 display cell-type enriched expression patterns of opioid receptors (Fig.

S10C), we found that opioid receptors in hDRG neurons were present in some but not other neu-

ron populations. Transcripts of δ-opioid receptor (*OPRD1*) were preferentially expressed in itch

populations hNP1 and hNP2, while *OPRM1* was enriched in all hPEP clusters (Fig. S10C). Since

opioid receptors are inhibitory GPCRs, our results suggest that activation of *OPRM1* could di-

rectly inhibit human nociceptive afferents, while *OPRD1* could be a molecular target for inhibit-

ing itch transmission. On the other hand, *OPRK1* was barely detected in our dataset, suggesting

that *OPRK1* agonists may relieve itch through indirect or central mechanisms (Fig. S10C).

Immunostaining of sensory fibers in the human skin using molecular markers identified by

single-soma deep RNA-seq dataset

Although human peripheral sensory afferents can be identified and visualized using the pan neu-

ronal marker antibody against PGP9.5, different types of human sensory afferents cannot be dis-

tinguished due to a lack of specific molecular markers. Developing an antibody panel to label

different types of human sensory afferents would be invaluable for basic research, translational

studies, and diagnostics. To start this direction and to test whether some molecular markers iden-

tified by the single-soma RNA-seq are useful for labeling specific types of human sensory fibers,

we conducted immunostaining of sensory afferents with leg skin biopsies from three normal

adult donors (see Methods for detailed donor information).

SST is specifically expressed in hPEP.SST neurons (Fig. 3B). Consistent with this molecular

prediction, we found SST^+ sensory fibers in the human skin sections in the dermis, the epider-

mis-dermis junction, entering the epidermis (Fig. S15A-C), and near the hair follicle (Fig. S15D-

F). In addition, double immunostaining revealed that SST^+ sensory fibers were $CGRP^+$ and made

up a subset of $CGPR^+$ sensory fibers (Fig. S15A-F). These results confirmed that human SST^+

sensory afferents had CGRP proteins and belonged to the PEP but not NP afferents. Since CGRP

involves in neuro-immune interactions and other physiological functions[85], addition of this neu-

ropeptide in human SST^+ afferents suggests a potential gain of new functions during evolution.

hPEP.KIT neurons specifically express *KIT* transcripts. Strong KIT signals were observed in

continuous regenerative cells at the basal layers of the skin, sweat ducts, and hair follicles, serv-

ing as a positive control. A few KIT^+ and $PGP9.5^+$ sensory fibers were observed around hair fol-

licles (Fig. S15G-I). Consistent with our RNA-seq results, KIT^+ sensory fibers were also positive

for CGRP and NEFH (Fig. S15J-O), suggesting that they were peptidergic A fibers. In short,

these results support that our high-fidelity hDRG neuron single-soma transcriptome dataset is

useful for selecting specific molecular markers to label and visualize different types of peripheral

somatosensory afferents.

A novel strategy to study functions of human somatosensory afferents using single-soma

deep RNA-seq dataset-informed microneurography recordings

Molecules profiles and physiological properties of hDRG neurons are intrinsically linked: the

expressed molecules form the physical basis for physiological properties. Physiological record-

ings of human somatosensory afferents have been conducted for more than half a century, which

generated important parts of our knowledge on the human somatosensory system. Physiological

properties of somatosensory afferents generate an important foundation for informing molecular

cloning of critical receptors, such as TRPV1 and PIEZO2. Due to a lack of clear molecular com-

positions of human somatosensory afferents, the reverse experimental direction, from molecular

profiles of human sensory afferents to inform physiological recordings, has not been possible.

With the single-soma deep RNA-seq dataset, we were finally in a position to do so. Here we fo-

cused on two populations of human sensory afferents to demonstrate the feasibility and power of

using single-soma deep RNA-seq informed microneurography recordings to study the function

of human afferent subtypes.

As mentioned previously, the hPEP.KIT population was highly correlated to the mouse

PEP3/CGRP-η cluster[15,18] (Fig. 2A-C), a population of TRPV1-negative A-fiber fast-conducting

hair pull-sensitive mechano-nociceptors[86]. Our single-soma deep RNA-seq data revealed that

hPEP.KIT neurons express NEFH and PIEZO2, but not TRPV1 (Fig. 7A), suggesting that they

were A-fiber mechano-nociceptors without heat/capsaicin sensitivity. Interestingly, our dataset

also revealed that hPEP.KIT neurons expressed a low level of *TRPM8* (Fig. 7A), a feature that

has not been reported in mice. Multiplex FISH validated the co-expression of *KIT*, *PIEZO2* and

TRPM8 in hDRG neurons (Fig. S16A). Altogether, these molecular profiles suggest that

hPEP.KIT neurons may be A-fiber HTMRs, responsive to cooling but not heating. Thus, we hy-

pothesized that there exist some previously uncharacterized fast-conducting A-fiber HTMRs in

the human skin, which can be activated by cooling but not heating stimuli.

To test this idea, using the *in vivo* electrophysiological technique of microneurography, single-

unit axonal recordings were performed from the radial, antebrachial, and peroneal nerves of

healthy participants (Fig. 7B-D). The A-HTMRs (n=10) were identified by their insensitivity to

soft-brush stroking while responding to a rough brush and displaying high indentation thresholds

(≥4 mN); further, they had Aβ-range conduction velocities (>30 m/s, Fig. 7B-F). Remarkably, a

subtype of these heat-insensitive A-HTMRs (n=5) units responded to cooling (Fig. 7G-H). Com-

pared to mechanically evoked responses, those evoked by cooling although relatively modest but

reproducible with tested in triplicates for each recording (Fig. S16B). Furthermore, these cool-

ing-evoked responses persisted during the sustained phase (Fig. S16B). These observations con-

firmed our prediction that some human cutaneous A-HTMRs are cold-sensitive but heat-

insensitive. Based on the transcriptome dataset and current knowledge about molecular receptors

for heat and cold, hPEP.KIT seems to be the only population that has the molecular basis for this

distinct combination of physiological properties. Thus, we propose that the molecularly defined

hPEP.KIT population is correlated to the physiologically defined A-HTMRs with cold but not

heat sensitivity (Fig. 7I).

Human C-LTMRs are readily found during microneurography recordings from the upper limb

including the distal regions[87], but they seem to sparsely innervate the distal lower limb[88]. This is

consistent with our sequencing results wherein the hC.LTMRs constituted a small population of

neurons in lower thoracic and lumbar level DRGs (Fig. 1G). Unexpectedly, our sequencing data

revealed that human hC.LTMRs had almost no expression of the cold and menthol receptor

TRPM8 (Fig. 8A) though C.LTMRs display sensitivity to cooling in both humans and mice[89,90].

Another unexpected finding was that hC.LTMRs expressed *TRPV1* (Fig. 8A). Multiplex FISH

confirmed that $CASQ2^+$ hC.LTMRs were TRPM8$^-$ but TRPV1$^+$ (Fig. S17A). This expression

pattern suggested that hC.LTMRs might respond to heating and capsaicin, a novel physiological

property that has not been discovered in rodent or non-human primate models, but is predicted

not to respond to menthol. In human microneurography, the C-LTMRs (n=11) were identified by

their soft-brush sensitivity, low indentation threshold (in this case, ≤0.7 mN), and slow conduc-

tion velocity (~1 m/s, n=11, Fig. 7B-C, 8B-D). Consistent with mouse C-LTMRs, these human

counterparts responded to hair movement (Fig. 8D) and dynamic cooling (Fig. 8E). In two of

them, after having confirmed the cooling response, we applied menthol to their individual recep-

tive fields resulting in a cold sensation, but the recorded C-LTMRs, while still responsive to me-

chanical and thermal stimuli, were not activated by menthol (Fig. 8F). Remarkably,

microneurography recordings showed that a subset of human C-LTMRs responded to dynamic

heating (5 out of 11 units or 45%, example shown in Fig. 8G). This proportion of heat-sensitive

C-LTMRs aligns with findings in the rabbit (8 out of 18 units or 44%)[91]. In three C-LTMRs,

after having confirmed the heating response, we applied capsaicin to their receptive fields. Con-

sistent with the sequencing results, all three were activated by capsaicin (Fig. 8H). Relative to

mechanical stimulation, the C-LTMR responses to dynamic temperature changes were compara-

tively modest. However, these responses were reproducible, having been tested in triplicates for

each modality and recording (Fig. S16B-C). Interestingly, the C-LTMR has different cooling

response properties compared to A-HTMR cooling+ fibers. The cooling-evoked responses rapid-

ly diminished in C-TLMR but persisted during the sustained phase in A-HTMR cooling+ fibers.

The existence of a polymodal (mechano-heat-cold) C-LTMR type is novel and confirms the se-

quencing predictions. Furthermore, the cooling response must be mediated through a non-

TRPM8 mechanism.

For comparison, our recordings also identified C-HTMRs (n=11) that did not respond to soft-

brush stroking and hair movement (they responded to a rough brush), had high indentation

thresholds (in this case, ≥10 mN), and slow conduction velocities (~1 m/s, Fig. 7B-D & S17B-E).

Based on their temperature responses (tested in 9 units), a mix of C-fiber mechano-heat (n=6), C-

fiber mechano-cold (n=2), and C-fiber polymodal (mechano-heat-cold, n=1) subtypes were iden-

tified. An example of heating and capsaicin responses of a C-fiber mechano-heat nociceptor is

shown in Fig. S17F-G. Collectively, these results highlight the accuracy and utility of our single-

soma deep RNA-seq dataset of hDRG neurons in guiding and informing the functional character-

ization of human somatosensory afferents and the power of combining the two approaches. We

believe that this signifies a promising new research direction to link molecular and physiological

types and to discover novel functional properties of human somatosensory afferents.

Discussion

Despite the high relevance to human health, molecular and cellular mechanisms underlying normal and pathological human somatosensation remain largely elusive. In this study, we developed a novel LCM-based approach for single-soma deep RNA-seq and recovered more than 9000 unique transcripts per neuron over 1000 adult hDRG neurons. The sequenced hDRG neurons were clustered into 16 molecular groups, displaying similarities to and remarkable differences from macaque and mouse DRG neurons. As exemplified in the study, our dataset provides important and novel insights into human pain and itch sensory phenomena, explains mechanisms of drug effects, and represents a rich resource to identify new molecular targets for modulating the activity of itch- and pain-sensing primary afferents. Moreover, our dataset successfully guided histological labeling (Fig. S15) and physiological recording (Fig. 7, 8 & S17) of different types of human sensory afferents, leading to discovery of their new anatomical features and physiological properties, and serving as a common ground to connect them all together (Fig. 7I).

Single-cell RNA-seq of human DRG/TG neurons has been technically challenging. One main hurdle is to isolate sensory neurons from a large population of non-neuronal cells. The traditional enzymatic and mechanical dissociation method is incompatible with human DRG/TG neurons. Strategies, including spatial transcriptomics and single-nucleus RNA-seq[22,23,92], have generated some pioneering datasets characterizing the molecular profiles of human DRG and TG neurons. Nevertheless, the number of transcripts, sequencing depth, or single-cell resolution of the previous studies needs to be improved. Importantly, though nuclear and soma transcripts are intrinsically linked and overlapping to some extent, the soma contains much more transcripts both quan-

titatively and qualitatively. Soma transcripts are also one step closer than nuclear transcripts for

functions. Thus, soma transcripts are preferable, if available, for cell type clustering and func-

tional interpretation. In this study, we developed a new strategy by combining fresh frozen

hDRG tissues, cryo-section, laser capture microdissection of individual neuronal soma, and

Smart-seq2 deep sequencing. Fresh frozen tissue and cryosection techniques minimized

transcriptomic changes during the tissue transportation and single-cell isolation process. LCM

allowed isolation of DRG neuronal soma with minimal contamination from surrounding non-

neuronal cells, while preserving information about cellular morphology, and the Smart-seq2 pro-

tocol enabled a high recovery rate of mRNA molecules. Although LCM has been used for isolat-

ing a group of neurons for RNA-seq[6,7] or single TG neurons for genomic DNA analysis[93], the

successful application of LCM for single-cell transcriptomic analysis has not been achieved be-

fore. Thus, we have established a new method for single-cell RNA-seq of adult hDRG neurons.

Our approach should be readily applicable to other human neurons with large soma sizes, such as

other peripheral ganglia neurons, motor neurons, etc.

Different single-cell RNA-seq approaches, including single-nucleus RNA-seq[21,23,92], spatial

transcriptomics[22], and our LCM-based single-soma RNA-seq, have generated four datasets of

transcriptome profiles and cell type clusters of hDRG neurons. These datasets and results overlap

to some extent but also exhibit some significant differences (an example shown in Fig. S4). The

observed differences are likely caused by both biological and technical features associated with

each method. Given the high sequencing depth of transcripts from the neuronal soma, our ap-

proach is powerful for molecular discovery, especially for functional molecules expressed at a

low level. For example, our approach detected the specific expression of *MRGPRX1 and*

MRGPRX4, two important pruritogen-sensing GPCRs, in hNP1 and/or hNP2 neurons, while the

previous datasets barely or not detected these transcripts[22,23]. Our analysis of the expression of

receptors, ion channels, and neuropeptides, in human itch-sensing DRG neurons have also identi-

fied a set of potential new targets for modulating activities of these sensory afferents (Fig. 6).

In addition, our sequencing and microneurography results raise many interesting questions re-

garding the molecular receptors and afferent types involved in human cold and mechanical pain

sensation. For example, human hC.LTMRs rely on a non-TRPM8 cold receptor or another

mechanism for their cooling sensitivity[88]. We noticed a low expression level of *TRPA1*, which

also has cold sensitivity[94,95], within this population (Fig. S7E). Whether *TRPA1* or some current-

ly unknown cold receptors mediate cooling sensitivity in this population will be of interest for

future research. The discovery of the hPEP.KIT population indicated a potential role for PIEZO2

in human mechano-nociception. This population also responded to cooling, which is a curious

property of a large-diameter myelinated nociceptor. Our discovery of TRPM8 expression in the-

se neurons provides a molecular explanation for this unique property. In addition, a likely mo-

lecular type for C-HTMRs, which display a mixture of responsiveness to mechanical forces and

temperature, is the hPEP.TRPV1/A1.1 population. Since hPEP.TRPV1/A1.1 neurons have no

expression of *PIEZO2* or *PIEZO1* (Fig. S9A), a non-*PIEZO* mechanoreceptor may exist in these

neurons for mediating mechanical pain sensation. This is consistent with reports that patients

with *PIEZO2* loss-of-function mutations still have normal mechanical pain threshold and sensi-

tivity[68,96]. One suggested mechanical pain channel, TMEM120A (TACAN), is broadly expressed

in all types of hDRG neurons (Fig. S9A). This expression pattern does not support its purported

role as a mechanical pain receptor in humans. Both C-LTMRs and A-HTMRs responded to cool-

ing in human microneurography. While the responses were comparatively modest in contrast to

mechanically evoked responses, they were consistent on a trial-to-trial basis. This suggests that

the requisite circuit may already be in place, which could potentially have implications for un-

derstanding thermal hypersensitivities in pathological states. Indeed, there is indirect evidence

from human psychophysics and targeted pharmacology, pointing to the role of C-LTMRs in me-

diating acute cold allodynia[97] and the role of A fibers in signaling chronic cold allodynia[98]. In

short, our dataset serves as an important atlas for understanding sensory properties of hDRG af-

ferents and somatosensation more generally.

Insights into the relationship of the neuronal types between the mouse and human are essential,

as such knowledge is important for translation of pre-clinical discoveries and for inferring func-

tions of human sensory afferents. The three different analysis strategies used in this study com-

plement each other as they have different strength and weakness (see Methods for details). To

our satisfaction, despite the different advantages and disadvantages, results from these methods

were highly consistent with each other (Fig. 2 & S5). Our results suggested that many broader

functional groups of DRG sensory afferents are conserved across species, despite noticeably mo-

lecular differences (Fig. 2 & S6). The greatest divergence between mouse and hDRG neurons

was observed among C- and A-fiber nociceptors. Mice contain four C-fiber nociceptors (PEP1

neuron subtypes), two expressing TRPV1 but not TRPA1 or PIEZO2 (PEP1.1 and PEP1.2) and

another two which expresses all three channels (PEP1.3 and PEP1.4) (Fig. 2A & 2C). While not

much is known about mouse PEP1.2/CGRP beta, PEP1.1 represents a noxious heat sensory type,

terminating with free nerve endings in epidermis of hairy skin[99]. PEP1.3/CGRP gamma neurons

innervate mainly internal organs and are a silent nociceptor type, becoming active during in-

flammation[100]. No functional studies have been performed on PEP1.4 in the mouse. These neu-

rons are speculated to represent C-HTMRs which are functionally known to exist in the mouse[101].

In humans, it seems that this diversity of C-fiber nociceptors has conflated into two types, the

hPEP.TRPV1/A1.1 and hPEP.TRPV1/A1.2 (Fig. 2A & 2C), expressing both TRPV1/TRPA1

with or without PIEZO2, respectively. If similar to mice, different neuronal populations in hu-

man are somewhat specialized according localization of peripheral target, one of these types may

represent deeply innervating silent nociceptors, known to be present in human[102], and the other

involved in slow noxious thermal and burning cutaneous sensation and perhaps joint pain. One

thing to note is that our dataset contains transcriptomes of hDRG neurons from the lower thorac-

ic and lumbar levels, which mainly convey sensations from the leg and lower back. Thus, it re-

mains an open question whether humans have a population of internal organ-specific C-

nociceptors, like the mouse PEP1.3 cluster. This question will be resolved when sequencing

more hDRG neurons from the thoracic level.

For A-fiber nociceptors, only two types have been identified in mice. The PEP2/CGRP-zeta

mechano-heat nociceptive population expresses PIEZO2 and low levels of TRPV1[16,33], con-

sistent with heat activation only at very high temperatures[99], and the PEP3/CGRP-eta mechano-

nociceptive conveying fast and sharp pain, including hair pulling[44,103]. In this study, we identified

five types A-fiber nociceptors in hDRG neurons (Fig. 1G, 2A & 2C). Two populations have

clearly conserved features to mouse: hPEP.CHRNA7 showed molecular similarities to mouse

and macaque PEP2/CGRP-ζ, and hPEP.KIT to mouse and macaque PEP3/CGRP-η. The other

three populations displayed much greater divergence (i.e. hPEP.NTRK3, hPEP.PIEZO[h], and

hPEP.0). The hPEP.PIEZO[h] cluster is particularly interesting since these neurons express very

high levels of *PIEZO2* and *PIEZO1,* unlike any neuron types in the mouse. Our results suggest

that fast pain and interoception might have been an evolutionary preference in humans as com-

pared to rodents, an effect that may relate to significantly increased human body sizes.

Finally, in this study, we sequenced over 1000 DRG neurons from 6 DRGs at the thoracic (T11-

T12) and lumbar (L2-L5) levels of three Caucasian human donors with no obvious somatosenso-

ry or systematic diseases and substance uses (Supplementary tables 1 & 2). It is about ~150 to

200 neurons per DRG, which represent 1-2% of the total neurons within a hDRG. Thus, sequenc-

ing more neurons might give an improved representation. The existence of different types of RA

and SA Aβ-LTMRs and different types of proprioceptors in humans is well known, but our cur-

rent dataset did not have the resolution yet to separate them (Fig. 1G). With increased number of

neurons sequenced, we anticipate discovering an even greater heterogeneity among hDRG neu-

rons. In addition, DRGs at different spinal levels innervate different peripheral target tissues[29].

Sequencing DRG neurons from different spinal levels will help to uncover molecular and cellu-

lar mechanisms underlying physiological functions. Moreover, increased sampling from addi-

tional donors representing different demographics would also be critical for investigating sex,

race, and age-related differences. Last but not the least, pathological conditions greatly alter the

transcriptomic landscape[104,105]. Systematic comparison of molecular and cellular changes be-

tween donors at the baseline condition (like the screening criteria of this study) and those with

chronic itch or chronic pain would be of great value, if samples are available, for understanding

pathological mechanisms and for identifying molecular targets for effective treatments.

Reference:

1   Nascimento, A. I., Mar, F. M. & Sousa, M. M. The intriguing nature of dorsal root ganglion neurons: Linking structure with polarity and function. *Prog Neurobiol* **168**, 86-103, doi:10.1016/j.pneurobio.2018.05.002 (2018).

2   Purves, D. A., George J.; Fitzpatrick, David; Katz, Lawrence C.; LaMantia, Anthony-Samuel; McNamara, James O.; Williams, S. Mark. The Major Afferent Pathway for Mechanosensory Information: The Dorsal Column-Medial Lemniscus System. *Neuroscience. 2nd edition* (2001).

3   Azimi, E., Xia, J. & Lerner, E. A. Peripheral mechanisms of itch. *Itch-Management in Clinical Practice* **50**, 18-23 (2016).

4   Mills, S. E. E., Nicolson, K. P. & Smith, B. H. Chronic pain: a review of its epidemiology and associated factors in population-based studies. *Br J Anaesth* **123**, e273-e283, doi:10.1016/j.bja.2019.03.023 (2019).

5   Manglik, A. *et al.* Structure-based discovery of opioid analgesics with reduced side

effects. *Nature* **537**, 185-190, doi:10.1038/nature19112 (2016).

6   Yosipovitch, G., Rosen, J. D. & Hashimoto, T. Itch: From mechanism to (novel)

therapeutic   approaches.   *J   Allergy   Clin   Immunol*   **142**,   1375-1390,

doi:10.1016/j.jaci.2018.09.005 (2018).

7   Burma, N. E., Leduc□Pessah, H., Fan, C. Y. & Trang, T. Animal models of chronic pain:

advances and challenges for clinical translation. *Journal of neuroscience research* **95**,

1242-1256 (2017).

8   Hill, R. NK1 (substance P) receptor antagonists–why are they not analgesic in humans?

Trends in pharmacological sciences **21**, 244-246 (2000).

9   Haberberger, R. V., Barry, C., Dominguez, N. & Matusica, D. Human Dorsal Root

Ganglia. *Front Cell Neurosci* **13**, 271, doi:10.3389/fncel.2019.00271 (2019).

10   Middleton, S. J. *et al.* Studying human nociceptors: from fundamentals to clinic. *Brain*

144, 1312-1335 (2021).

11   Yu, H. *et al.* MRGPRX4 is a bile acid receptor for human cholestatic itch. *Elife* **8**, e48431

(2019).

12   Alemi, F. *et al.* The TGR5 receptor mediates bile acid–induced itch and analgesia. *The*

Journal of clinical investigation **123**, 1513-1530 (2013).

13   Tang, F. *et al.* mRNA-Seq whole-transcriptome analysis of a single cell. *Nat Methods* **6**,

377-382, doi:10.1038/nmeth.1315 (2009).

14   Hwang, B., Lee, J. H. & Bang, D. Single-cell RNA sequencing technologies and

bioinformatics pipelines. *Exp Mol Med* **50**, 96, doi:10.1038/s12276-018-0071-8 (2018).

15   Usoskin, D. *et al.* Unbiased classification of sensory neuron types by large-scale single-

cell RNA sequencing. *Nature neuroscience* **18**, 145-153 (2015).

16 Sharma, N. *et al.* The emergence of transcriptional identity in somatosensory neurons.

Nature **577**, 392-398 (2020).

17 Wang, K. *et al.* Single-cell transcriptomic analysis of somatosensory neurons uncovers

temporal development of neuropathic pain. *Cell research* **31**, 904-918 (2021).

18 Kupari, J. *et al.* Single cell transcriptomics of primate sensory neurons identifies cell

types associated with chronic pain. *Nature communications* **12**, 1-15 (2021).

19 Stuart, T. *et al.* Comprehensive integration of single-cell data. *Cell* **177**, 1888-1902.

e1821 (2019).

20 Mattei, D. *et al.* Enzymatic dissociation induces transcriptional and proteotype bias in

brain cell populations. *International journal of molecular sciences* **21**, 7944 (2020).

21 Jung, M. *et al.* Cross-species transcriptomic atlas of dorsal root ganglia reveals species-

specific programs for sensory function. *Nature Communications* **14**, 366 (2023).

22 Tavares-Ferreira, D. *et al.* Spatial transcriptomics of dorsal root ganglia identifies

molecular signatures of human nociceptors. *Science translational medicine* **14**, eabj8186

(2022).

23 Nguyen, M. Q., von Buchholtz, L. J., Reker, A. N., Ryba, N. J. & Davidson, S. Single-

nucleus transcriptomic analysis of human dorsal root ganglion neurons. *Elife* **10**, e71752

(2021).

24 Bhuiyan, S. *et al.* Harmonized cross-species cell atlases of trigeminal and dorsal root

ganglia. *bioRxiv*, 2023.2007. 2004.547740 (2023).

25 Ding, J. *et al.* Systematic comparison of single-cell and single-nucleus RNA-sequencing

methods. *Nature biotechnology* **38**, 737-746 (2020).

26 Picelli, S. *et al.* Full-length RNA-seq from single cells using Smart-seq2. *Nature*
protocols **9**, 171-181 (2014).

27 Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.
e3529 (2021).

28 Abraira, V. E. & Ginty, D. D. The sensory neurons of touch. *Neuron* **79**, 618-639,
doi:10.1016/j.neuron.2013.07.051 (2013).

29 Luo, L. *PRINCIPLES OF NEUROBIOLOGY*. 265-267 (Garland Science, 2016).

30 Chen, Y.-C. *et al.* Transcriptional regulator PRDM12 is essential for human pain
perception. *Nature genetics* **47**, 803-808 (2015).

31 Barkas, N. *et al.* Joint analysis of heterogeneous single-cell RNA-seq dataset collections.
Nature methods **16**, 695-698 (2019).

32 Jovic, D. *et al.* Single-cell RNA sequencing technologies and applications: A brief
overview. *Clin Transl Med* **12**, e694, doi:10.1002/ctm2.694 (2022).

33 Zeisel, A. *et al.* Molecular architecture of the mouse nervous system. *Cell* **174**, 999-1014.
e1022 (2018).

34 Hippenmeyer, S. *et al.* A developmental switch in the response of DRG neurons to ETS
transcription factor signaling. *PLoS biology* **3**, e159 (2005).

35 Jeub, M. *et al.* The transcription factor Smad-interacting protein 1 controls pain
sensitivity via modulation of DRG neuron excitability. *PAIN®* **152**, 2384-2398 (2011).

36 Scott, A. *et al.* Transcription factor short stature homeobox 2 is required for proper
development of tropomyosin-related kinase B-expressing mechanosensory neurons.
Journal of Neuroscience **31**, 6741-6749 (2011).

37 Abdo, H. *et al.* Dependence on the transcription factor Shox2 for specification of sensory

neurons conveying discriminative touch. *European Journal of Neuroscience* **34**, 1529-
1541 (2011).

38   Levanon, D. *et al.* The Runx3 transcription factor regulates development and survival of
TrkC dorsal root ganglia neurons. *The EMBO journal* **21**, 3454-3463 (2002).

39   Chen, C.-L. *et al.* Runx1 determines nociceptive sensory neuron phenotype and is
required for thermal and neuropathic pain. *Neuron* **49**, 365-377 (2006).

40   Vermeiren, S., Bellefroid, E. J. & Desiderio, S. Vertebrate sensory ganglia: common and
divergent features of the transcriptional programs generating their functional
specialization. *Frontiers in Cell and Developmental Biology* **8**, 587699 (2020).

41   Liu, Q. & Dong, X. The role of the Mrgpr receptor family in itch. *Pharmacology of itch*,
71-88 (2015).

42   Meixiong, J., Vasavda, C., Snyder, S. H. & Dong, X. MRGPRX4 is a G protein-coupled
receptor activated by bile acids that may contribute to cholestatic pruritus. *Proc Natl*
Acad Sci U S A **116**, 10525-10530, doi:10.1073/pnas.1903316116 (2019).

43   Namer, B. *et al.* Separate peripheral pathways for pruritus in man. *Journal of*
neurophysiology **100**, 2062-2069 (2008).

44   Dong, P. *et al.* TRPC3 is dispensable for β-alanine triggered acute itch. *Scientific reports*
7, 1-12 (2017).

45   Rehfeld, J. F. Cholecystokinin and the hormone concept. *Endocrine Connections* **10**,
R139-R150 (2021).

46   McNeil, B. & Dong, X. Peripheral mechanisms of itch. *Neuroscience bulletin* **28**, 100-
110 (2012).

47   Colburn, R. W. *et al.* Attenuated cold sensitivity in TRPM8 null mice. *Neuron* **54**, 379-

386 (2007).

48   Bautista, D. M. *et al.* The menthol receptor TRPM8 is the principal detector of

environmental cold. *Nature* **448**, 204-208 (2007).

49   Dhaka, A. *et al.* TRPM8 is required for cold sensation in mice. *Neuron* **54**, 371-378

(2007).

50   Samour, M. S., Nagi, S. S. & Mahns, D. A. Cav3. 2-expressing low-threshold C fibres in

human hairy skin contribute to cold allodynia—a non-TRPV1-and non-TRPM8-

dependent phenomenon. *Pain* **156**, 1566-1575 (2015).

51   Campero, M., Baumann, T., Bostock, H. & Ochoa, J. Human cutaneous C fibres

activated by cooling, heating and menthol. *The Journal of physiology* **587**, 5633-5652

(2009).

52   Caterina, M. J. *et al.* The capsaicin receptor: a heat-activated ion channel in the pain

pathway. *Nature* **389**, 816-824 (1997).

53   Bandell, M. *et al.* Noxious cold ion channel TRPA1 is activated by pungent compounds

and bradykinin. *Neuron* **41**, 849-857 (2004).

54   Defaye, M. *et al.* Gut-innervating TRPV1+ neurons drive chronic visceral pain via

microglial P2Y12 receptor. *Cellular and Molecular Gastroenterology and Hepatology*

13, 977-999 (2022).

55   Yang, F.-C. *et al.* Genetic control of the segregation of pain-related sensory neurons

innervating the cutaneous versus deep tissues. *Cell reports* **5**, 1353-1364 (2013).

56   Liu, S. *et al.* A neuroanatomical basis for electroacupuncture to drive the vagal–adrenal

axis. *Nature* **598**, 641-645 (2021).

57   Albrecht, P. J. *et al.* Pathologic alterations of cutaneous innervation and vasculature in

affected limbs from patients with complex regional pain syndrome. *Pain* **120**, 244-266

(2006).

58   Morelli, C. *et al.* Identification of a population of peripheral sensory neurons that

regulates blood pressure. *Cell reports* **35**, 109191 (2021).

59   Albrecht, P. J. *et al.* Excessive peptidergic sensory innervation of cutaneous arteriole–

venule shunts (AVS) in the palmar glabrous skin of fibromyalgia patients: Implications

for widespread deep tissue pain and fatigue. *Pain medicine* **14**, 895-915 (2013).

60   Bowsher, D. *et al.* Absence of pain with hyperhidrosis: a new syndrome where vascular

afferents may mediate cutaneous sensation. *PAIN®* **147**, 287-298 (2009).

61   Xu, J. *et al.* GPR68 senses flow and is essential for vascular physiology. *Cell* **173**, 762-

775. e716 (2018).

62   Zeng, W.-Z. *et al.* PIEZOs mediate neuronal sensing of blood pressure and the

baroreceptor reflex. *Science* **362**, 464-467 (2018).

63   Marshall, K. L. *et al.* PIEZO2 in sensory neurons and urothelial cells coordinates

urination. *Nature* **588**, 290-295 (2020).

64   Perlman, R. L. Mouse models of human disease: an evolutionary perspective. *Evolution,*

medicine, and public health **2016**, 170-176 (2016).

65   Willimann, B. & Trüeb, R. M. Hair pain (trichodynia): frequency and relationship to hair

loss and patient gender. *Dermatology* **205**, 374-377 (2002).

66   von Buchholtz, L. J. *et al.* Decoding cellular mechanisms for mechanosensory

discrimination. *Neuron* **109**, 285-298. e285 (2021).

67   Wolfson, R. L. *et al.* DRG afferents that mediate physiologic and pathologic

mechanosensation from the distal colon. *Cell* **186**, 3368–3385 (2023).

68   Nagi, S. S. *et al.* An ultrafast system for signaling mechanical pain in human skin. *Science advances* **5**, eaaw1297 (2019).

69   Schmidt, R. F. *Encyclopedia of Pain.* pp 2171–2173 (Spriner 2007).

70   Löken, L. S., Wessberg, J., McGlone, F. & Olausson, H. Coding of pleasant touch by unmyelinated afferents in humans. *Nature neuroscience* **12**, 547-548 (2009).

71   Olausson, H., Wessberg, J., McGlone, F. & Vallbo, Å. The neurophysiology of unmyelinated tactile afferents. *Neuroscience & Biobehavioral Reviews* **34**, 185-191 (2010).

72   Huzard, D. *et al.* The impact of C-tactile low-threshold mechanoreceptors on affective touch and social interactions in mice. *Science Advances* **8**, eabo7566 (2022).

73   Xue, Y., Chidiac, C., Herault, Y. & Gaveriaux-Ruff, C. Pain behavior in SCN9A (Nav1. 7) and SCN10A (Nav1. 8) mutant rodent models. *Neuroscience Letters* **753**, 135844 (2021).

74   Duan, G. *et al.* A SCN10A SNP biases human pain sensitivity. *Molecular pain* **12**, 1744806916666083 (2016).

75   Reddy, V. B., Iuga, A. O., Shimada, S. G., LaMotte, R. H. & Lerner, E. A. Cowhage-evoked itch is mediated by a novel cysteine protease: a ligand of protease-activated receptors. *Journal of Neuroscience* **28**, 4331-4335 (2008).

76   Schmelz, M. *et al.* Chemical response pattern of different classes of C-nociceptors to pruritogens and algogens. *Journal of neurophysiology* **89**, 2441-2448 (2003).

77   Dahlhamer, J. *et al.* Prevalence of Chronic Pain and High-Impact Chronic Pain Among Adults - United States, 2016. *MMWR Morb Mortal Wkly Rep* **67**, 1001-1006, doi:10.15585/mmwr.mm6736a2 (2018).

78 Tajiri, K. & Shimizu, Y. Recent advances in the management of pruritus in chronic liver

diseases. *World J Gastroenterol* **23**, 3418-3426, doi:10.3748/wjg.v23.i19.3418 (2017).

79 Dong, X. & Dong, X. Peripheral and central mechanisms of itch. *Neuron* **98**, 482-494

(2018).

80 Bell, J., McQueen, D. & Rees, J. Involvement of histamine H4 and H1 receptors in

scratching induced by histamine receptor agonists in BalbC mice. *British journal of*

pharmacology **142**, 374-380 (2004).

81 Calandre, E. P., Rico-Villademoros, F. & Slim, M. Alpha2delta ligands, gabapentin,

pregabalin and mirogabalin: a review of their clinical pharmacology and therapeutic use.

Expert review of neurotherapeutics **16**, 1263-1277 (2016).

82 Peng, C. *et al.* miR-183 cluster scales mechanical pain sensitivity by regulating basal and

neuropathic pain genes. *Science* **356**, 1168-1171 (2017).

83 Cowan, A., Kehner, G. B. & Inan, S. Targeting itch with ligands selective for κ opioid

receptors. *Pharmacology of itch*, 291-314 (2015).

84 Ko, M. H., Song, M., Edwards, T., Lee, H. & Naughton, N. The role of central μ opioid

receptors in opioid-induced itch in primates. *Journal of Pharmacology and Experimental*

Therapeutics **310**, 169-176 (2004).

85 Chu, C., Artis, D. & Chiu, I. M. Neuro-immune interactions in the tissues. *Immunity* **52**,

464-474 (2020).

86 Ghitani, N. *et al.* Specialized mechanosensory nociceptors mediating rapid responses to

hair pull. *Neuron* **95**, 944-954. e944 (2017).

87 Xu, S. *et al.* Mechanoreceptive Aβ primary afferents discriminate naturalistic social touch

inputs at a functionally relevant time scale. *bioRxiv*, 2023.2007. 2022.549516 (2023).

88   Löken, L. S., Backlund Wasling, H., Olausson, H., McGlone, F. & Wessberg, J. A

topographical and physiological exploration of C-tactile afferents and their response to

menthol and histamine. *Journal of Neurophysiology* **127**, 463-473 (2022).

89   Nordin, M. Low☐threshold mechanoreceptive and nociceptive units with unmyelinated

(C) fibres in the human supraorbital nerve. *The Journal of physiology* **426**, 229-240

(1990).

90   Middleton, S. J. *et al.* Nav1. 7 is required for normal C-low threshold mechanoreceptor

function in humans and mice. *Brain* **145**, 3637-3653 (2022).

91   Shea, V. & Perl, E. Sensory receptors with unmyelinated (C) fibers innervating the skin

of the rabbit's ear. *Journal of neurophysiology* **54**, 491-501 (1985).

92   Yang, L. *et al.* Human and mouse trigeminal ganglia cell atlas implicates multiple cell

types in migraine. *Neuron* **110**, 1806-1821. e1808 (2022).

93   Wang, K., Lau, T. Y., Morales, M., Mont, E. K. & Straus, S. E. Laser-capture

microdissection: refining estimates of the quantity and distribution of latent herpes

simplex virus 1 and varicella-zoster virus DNA in human trigeminal Ganglia at the

single-cell level. *Journal of virology* **79**, 14079-14087 (2005).

94   Bautista, D. M. *et al.* TRPA1 mediates the inflammatory actions of environmental

irritants and proalgesic agents. *Cell* **124**, 1269-1282 (2006).

95   Macpherson, L. J. *et al.* Noxious compounds activate TRPA1 ion channels through

covalent modification of cysteines. *Nature* **445**, 541-545 (2007).

96   Chesler, A. T. *et al.* The role of PIEZO2 in human mechanosensation. *New England*

Journal of Medicine **375**, 1355-1364 (2016).

97   Shaikh, S., Nagi, S. S., McGlone, F. & Mahns, D. A. Psychophysical investigations into

the role of low-threshold C fibres in non-painful affective processing and pain

modulation. *PLoS One* **10**, e0138299 (2015).

98   Jørum, E. & Opstad, P.-K. A 4-year follow-up of non-freezing cold injury with cold

allodynia and neuropathy in 26 naval soldiers. *Scandinavian journal of pain* **19**, 441-451

(2019).

99   Qi, L. *et al.* A DRG genetic toolkit reveals molecular, morphological, and functional

diversity of somatosensory neuron subtypes. *bioRxiv*, 2023.2004. 2022.537932 (2023).

100 Prato, V. *et al.* Functional and molecular characterization of mechanoinsensitive "silent"

nociceptors. *Cell reports* **21**, 3102-3115 (2017).

101 Boada, M. D. & Woodbury, C. J. Physiological properties of mouse skin sensory neurons

recorded intracellularly in vivo: temperature effects on somal membrane properties.

Journal of neurophysiology **98**, 668-680 (2007).

102 Buffington, C. T. Visceral pain in humans: lessons from animals. *Current pain and*

headache reports **5**, 44-52 (2001).

103 Arcourt, A. *et al.* Touch receptor-derived sensory information alleviates acute pain

signaling and fine-tunes nociceptive reflex coordination. *Neuron* **93**, 179-193 (2017).

104 Xiao, H.-S. *et al.* Identification of gene expression profile of dorsal root ganglion in the

rat peripheral axotomy model of neuropathic pain. *Proceedings of the National Academy*

of Sciences **99**, 8360-8365 (2002).

105 North, R. Y. *et al.* Electrophysiological and transcriptomic correlates of neuropathic pain

in human dorsal root ganglion neurons. *Brain* **142**, 1215-1226 (2019).

106 Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet. journal **17**, 10-12 (2011).

107 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013).

108 Hu, Y. *et al.* Neural network learning defines glioblastoma features to be of neural crest
perivascular or radial glia lineages. *Science Advances* **8**, eabm6340 (2022).

109 Shrikumar, A., Greenside, P. & Kundaje, A. in *International conference on machine*
learning. 3145-3153 (PMLR).

110 Huynh-Thu, V. A., Irrthum, A., Wehenkel, L. & Geurts, P. Inferring regulatory networks
from expression data using tree-based methods. *PloS one* **5**, e12776 (2010).

111 Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering. *Nature*
methods **14**, 1083-1086 (2017).

112 Vallbo, A., Olausson, H., Wessberg, J. & Kakuda, N. Receptive field characteristics of
tactile units with myelinated afferents in hairy skin of human subjects. *The Journal of*
physiology **483**, 783-795 (1995).

113 Vallbo, Å., Olausson, H. & Wessberg, J. Unmyelinated afferents constitute a second
system coding tactile stimuli of the human hairy skin. *Journal of neurophysiology* **81**,
2753-2763 (1999).

114 Wasner, G., Schattschneider, J., Binder, A. & Baron, R. Topical menthol—a human
model for cold pain by activation and sensitization of C nociceptors. *Brain* **127**, 1159-
1171 (2004).

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

Acknowledgements:

We thank all tissue donors and their families for the generous donations. The new knowledge

pieces generated from this study would not be possible without their precious gifts. We thank

NDRI for helping with hDRG procurement, Dr. Isaac Chen, Ms. Arianna Unger, and other

members at the UPenn (University of Pennsylvania) department of neurosurgery for helping with

patient DRG procurement for method optimization, the UPenn Skin Biology and Disease Re-

source-based Center (SBDRC) for helping with laser capture microdissection, and the Children's

Hospital of Philadelphia (CHOP) Center for Applied Genomics (CAG) for helping with sequenc-

ing. We thank Dr. Yulong Li for his encouragement in starting this new research direction and all

lab members in the Luo and Ma labs for their helps in conducting this project. This project is

supported by an internal funding from UPenn to Dr. Luo. Dr. Luo is also supported by NIH

fundings (R01 (R01NS083702), R34 (NS118411), U01 (EY034681), and P30 (AR069589)). Dr.

Ernfors is supported by the Swedish Research Council (2019-00761), Knut and Alice

Wallenbergs Foundation, ERC advanced grant (PainCells 740491). Dr. Olausson is supported by

Knut and Alice Wallenberg Foundation (grants KAW 2019.0047 and KAW 2019.0487). Dr.

Nagi is supported by the Swedish Research Council (2021-03054), Swedish Medical Society

(SLS), and ALF Grants, Region Östergötland.

1107

1108

1109

1110

1111

1112

Materials and Methods:

Human tissues and subjects

Human DRG tissues were procured from National Disease Research Interchange (NDRI). The

research application was approved by the NDRI Feasibility Committee (RLUW1 01). Six DRGs

between T11 to L5 of three human donors aged from 23- 61 years were used in this study. The

dissected DRG tissues from human donors were immediately imbedded in OCT, shipped to the

Luo lab on dry ice and stored in -80 □ until use. The information of DRGs and dis-identified

donors and screening criteria are summarized in the Supplementary Tables 1 & 2. As determined

by the University of Pennsylvania IRB, this study was exempted from the human subject re-

quirements for the Luo lab.

1123

The human skin biopsies were extracted from three healthy donors at the college of medicine,

University of Florida. This tissue procurement was approved by the university IRB (protocols

IRB201500232 and IRB202300291). The information of human skin biopsies and dis-identified

donors is summarized in the Supplementary Table 3. These three donors are members of one

family and have no noticeable abnormal somatosensation or peripheral neuropathy. All partici-

pants were provided written informed consent and signed the document.

1130

1131 *In vivo* recordings of peripheral sensory afferents of healthy human subjects were performed at

1132 Linköping University, Sweden. These subjects were recruited through social media. All partici-

1133 pants were provided written informed consent before the start of the experiment. The study was

1134 approved by the ethics committee of Linköping University (dnr 2020-04426) and complied with

1135 the revised Declaration of Helsinki.

1136

Laser capture microdissection of hDRG neurons

The hDRGs imbedded in OCT were cryosectioned (Cryostat Leica Cm1950) into 20 μm sections

and mounted onto Arcturus PEN Membrane Frame Slides (Applied biosystems, LCM0521). One

of every five consecutive sections was collected for laser capture microdissection to avoid re-

peated dissection of the soma from the same neuron in different sections. The slides were stored

in -80 ☐ until further use.

1143

On the day of laser capture microdissection, the slides were transferred to the SBDRC laser cap-

ture microdissection (LCM) core on dry ice. Before dissection, the section was briefly stained

with RNase free Arcturus™ Histogene™ staining solution (Applied biosystems, 12241-05) for

better visualization of neuronal soma: 70% cold EtOH for 30s; Histogene™ staining for 10s; 70%

cold EtOH for 30s; 95% cold EtOH for 30s; 100% cold EtOH for 30s and air dry for 2min. Then,

the slide was put onto laser capture microdissection microscope system (Leica LMD6) for the

neuronal soma dissection. The laser was calibrated, and the laser intensity was adjusted to

achieve best dissection efficiency. The dissected individual neuronal soma was collected in the

cap of a 200 μl PCR tube containing 4 μl lysis buffer[26]. The sequencing library was generated

following Smart-seq2 workflow[26]. The libraries passing through all quality controls were select-

ed for the final sequencing.

1155

Sequencing and sequence alignment

The libraries were pooled together (384 samples for one batch) and sequenced on NovaSeq 6000

platform by the Children's Hospital of Philadelphia (CHOP) Center for Applied Genomics

(CAG). Raw sequencing data was demultiplexed with bcl2fastq2 v.2.20 (Illumina) followed by

Tn5 transposon adapter sequences trimming with Cutadapt[106]. The processed reads were then

aligned to human genome (GRCh38 GENCODE as the reference genome, and GRCh38.104.

GTF as the annotation) using STAR v.2.7.9a49[107]. STAR quantMode GeneCounts was used to

quantify unique mapped reads to gene counts.

1164

Analysis of single-soma RNA-seq data of hDRG neurons using Seurat and R

R (version 4.1.2) and Seurat (version 4.0.5) were used for the single-cell RNA-seq analysis. Six

objects were created from the individual biological replicates. The data were normalized

(NormalizeData) after which 4500 most variable features were selected (FindVariableFeatures).

To mitigate batch effects between replicates, we used Seurat's integrated analysis approach that

transforms datasets into a shared space using pairwise correspondences (or "anchors")[19]. An-

chors were first defined using FindIntegrationAnchors (dims□=□1:30) and the data were then

integrated (IntegrateData) and scaled (ScaleData), followed by principal component analysis

(PCA) (RunPCA, npcs□=□50). For clustering, the final parameters were: RunUMAP, reduc-

tion□=□pca, dims□=□1:25; FindNeighbors, reduction□=□pca, dims□=□1:25; FindClusters,

resolution□=□3.4. Highly similar clusters without clearly distinguishable markers were merged

to produce the final 16 clusters.

1177

Analysis of single-soma RNA-seq data of hDRG neurons using Conos

For Conos[31] analysis, single-soma hDRG data were integrated using CCA space

$buildGraph(k□=□8, k.self=3, space□=□"CCA", ncomps=30, n.odgenes=2000, ver-

bose=TRUE, snn=T, snn.k□=□10). For human single-soma and human single-nucleus dataset,

co-integration was performed as $buildGraph(k□=□8, k.self=3, space□=□"CCA",

n.odgenes=2000, verbose=TRUE, snn=T, snn.k□=□10). For Conos co-clustering mouse

(Sharms) dataset was downsampled to max 300 cells per cluster, co-integration was performed as

$buildGraph(k□=□8, k.self=3, space□=□"CCA", n.odgenes=2000, verbose=TRUE,

snn=T,snn.k□=□10). Macaque (Kupari, SmartSeq2 dataset) was used for interspecies analysis.

For Conos co-clustering macaque/human graph was built as $buildGraph(k□=□4, k.self=3,

space□=□"CCA", ncomps = 30, n.odgenes=2000, snn=F, snn.k□=□10). For all UMAP plots in

Conos graphs were embedded as: $embedGraph(method□=□"UMAP", spread=15, seed = 3).

Label propagation ($propagateLabels) was run using method "diffusion".

1191

Methods used to elucidate cross-species cluster relationships

We used four different interspecies analysis approaches. First, Conos, which uses graph-based

dataset integration and was developed to co-cluster and compare datasets originating from differ-

ent RNA-seq platforms and species (Fig. 2A-B, S5A-B). Second, probabilistic neural network

analysis which is a variant of machine learning in which learning module is trained with one da-

taset and then testing other datasets for pattern recognition and probability output[18] (Fig. S5C-D).

Third, neural network based hierarchical clustering analysis (Fig. S5E). In the hierarchical clus-

tering analysis, each query neuron types, either human or macaque, is assigned weights of the

sensory-type associated patterns by a neural network, which was trained with gene patterns in-

cluding both species specific and shared cross-species features in the different sensory neuron

types. The weighted gene patterns were then used for dimensional reduction and nearest neigh-

bor analysis to infer the hierarchical relationship. Finally, we performed transcription factor as-

sociated gene regulatory network analysis across all three species using genes network modules

presumably driven by individual transcription factors (Fig.2D). These three methods have differ-

ent strengths and weaknesses. Conos finds shared principal components between integrated da-

tasets, but some species-specific features may be lost and lead to impaired statistical sufficiency

during integration and furthermore can be affected by the number of principal components and

nearest neighbor distance. Machine learning is based on gene expression and is not supervised by

shared latent space (i.e. common principal components). Each single reference dataset is used to

train the machine learning module and then tested by the other dataset. Thus, an advantage of

this method is that at the stage of machine learning, datasets are not integrated, and hence, prob-

ability calculations are not affected by principal components or nearest distance. Conversely, its

disadvantage is that since it emphasizes the features of each cell type, the learning accuracy and

reliability depend on the robustness of the reference or training dataset. For the third approach,

machine learning-based hierarchical clustering, we extracted the weight of cell type-specific fea-

tures to construct the latent space covering all cell types across species, whether shared or not.

With this strategy, we tried to obtain sufficient latent space, as compared to Conos, by training

and predicting every dataset independently and furthermore, parameterization was used to find

the most robust hierarchical clustering.

1221

1222   *Cell type probabilistic similarity estimation across cell types and the data integration across*

1223   *species*

1224   The assessment of cell type purity, the probabilistic similarity, and cell-type integration across

1225   species are performed using packages in a machine learning based single-cell analysis toolkit-

1226   scCAMEL, released separately at https://sccamel.readthedocs.io/.

1227

Probabilistic similarity estimation across cell types

The calculation of cell-type probabilistic score has been described in SWAPLINE package[108].

Briefly, a vanilla neural network model was built for cell-type classification. To train the model,

we removed the cell cycle–related genes and then computed the most variable features. In addi-

tion, we ranked the marker genes for each cell type by two heuristics for the cell-type specificity

of both fold change and enrichment score change. Subsequently, the ranked marker genes and

the most variable genes were merged, log-transformed, and scaled by min-max normalization for

learning models. The frame of the neural network model and the parameters have been described

in the SWAPLINE package. The learning accuracy of the neural-network classifier was inspect-

ed against epoch numbers and was estimated by k-fold cross-validation (k = 3). The learning rate

and learning epochs were selected according to the maximum point of the learning curve reach-

ing the accuracy plateaus. The probabilistic scores from mouse and macaque species against hu-

man reference were visualized in violin plot.

1241

Data integration across species

For the integration task, we applied interpretable neural-network learning. First, we took one

dataset from the dataset pool. We trained a neural-network classifier by learning the transcrip-

tional features of each cell-types in this dataset and then calculated the trained cells' probabilistic

scores against all cell-types. Subsequently, we used all other datasets as query datasets and calcu-

lated the probabilistic score of every cell in each query dataset via the trained classifier. Then, we

took another dataset from the dataset pool and repeated the training and prediction. We repeat the

training and prediction till every dataset has been used as a training reference for the predictions.

Here, we consider that the probabilistic score of each cell reflects the weighted gene patterns

representing each trained cell-type. Thus, we merged the probabilistic scores of all cells from all

trained and predicted datasets for the principal component analysis. The most significant princi-

pal components were determined by the elbow method and subsequently used as the latent space

for further downstream analysis. The tree plot was constructed with the parameter of 11 principal

components, 90 nearest neighbors, and correlation metric. The trained cell-type similarity was

calculated with the correlation distance and the average/UPGMA linkage and visualized in the

hierarchical heatmap.

1258

In parallel, we normalized the gene expression by interpretable learning. We transformed the

gene symbols of each species into the nomenclature in Homo sapiens. We estimated the features'

weights in each reference cell-type by using the DeepLift algorithm[109]. The gene expression of

each cell that has been learned or predicted in one trained reference dataset, was inferred by the

matrix multiplication between the features' weights and the cell-type probabilistic scores. And

the final gene*cell expression matrix was calculated by the average of non-empty values across

all datasets. Using this normalized expression matrix, we enriched the mostly co-expressed genes

via spearman correlation. These co-expressed genes were used for inferring the TF associated

gene patterns via a modified GENIE3, as described in[110,111]. The result was visualized as a hier-

archical heatmap.

1269

Multiplex FISH, confocal microscopy imaging, and quantification

OCT embedded freshly dissected human lumbar or thoracic DRG tissues were cryosectioned at

20 μm thickness and mounted on glass slides. The slides were stored in −80 °C to preserve

RNA integrity until use. RNAscope Fluorescent Multiplex Reagent Kit and RNASCOPE probes

for the targeted genes (Advanced Cell Diagnostics Inc.) were used for *multiplex FISH*.

RNAscope *in situ* hybridization was performed in accordance with the manufacturer's instruc-

tions. In brief, fresh frozen hDRG sections were fixed, dehydrated, and treated with protease.

The sections were then hybridized with the respective target probe for 2 hr at 40°C, followed by

two to three rounds of signal amplification. The sections were then mounted under coverslips,

sealed with nail polish, and stored in the dark at 4°C until imaged. A Leica SP5 confocal micro-

scope was used to capture images and ImageJ was used for image analysis. In some DRG neu-

rons, accumulation of lipofuscin in part of cells caused strong autofluorescence in all channels.

These signals were considered as non-specific background (labeled by asterisk) were excluded

for analysis. (See Fig. S9 for examples). The percentage of each cluster over all DRG neurons

could be a little bit overestimated due to the following two reasons: 1) Some marker genes or

marker gene combination may also label a small subset of other cell types; 2) An underestima-

tion in quantification of total neuronal numbers because some cells have neither multiple FISH

signals nor DAPI (4′,6-diamidino-2-phenylindole) nucleus staining signals.

1288

Human skin biopsy extraction, processing, and immunostaining

Dermal skin punch biopsies were performed as described herein. Briefly, 1 cc of lidocaine was

injected subdermally at each biopsy location (Supplementary table 3). A total of six, 3 mm der-

mal skin punch biopsies were performed on each patient. Excised skin was immediately placed

in 1.5 mL eppendorf tubes containing 4 degree Celsius 4% paraformaldehyde (PFA) solution that

was freshly prepared on the same day of the skin biopsy procedure. Biopsy tissue was fixed in 4%

PFA (dissolved in PBS) for exactly four hours at 4 degrees, followed by 2 X 30 minutes washes

in phosphate buffered saline (PBS) solution, and then cryoprotected using 1XPBS, 30% sucrose

at 4 degrees. These tissues in cold 1XPBS, 30% sucrose were overnight shipped to the lab of

Integrated Tissue Dynamics LLC.

1299

The skin biopsies were mounted in OCT and cryosectioned into 14 μm sections. Adjacent sec-

tions were collected by continuous slides. Immunofluorescence in this study was performed us-

ing combinations of mouse monoclional anti-human PGP9.5 (Protein Gene Product, CedarLane,

Burlington, Canada, 31A3, [source UltraClone Ltd, Isle of Wight, UK]; 1:200), sheep polyclonal

anti-human CGRP, mouse monoclonal anti-human NEFH (Sigma [ab142]; 1:400), rabbit anti-

human SST (ImmunoStar 20067, Hudson, WI, USA), and anti-human KIT. Slides were

preincubated in 1% bovine serum albumin and 0.3% Triton X-100 in PBS (PBS-TB) for 30

minutes and then incubated with primary antibodies diluted in PBS-TB overnight in a humid atmosphere at 4°C. Slides were then rinsed in excess PBS for 30 minutes and incubated for 2 hours at room temperature with the appropriate secondary antibodies diluted in PBS-TB. Following secondary antibody incubation, the sections were rinsed for 30 minutes in PBS and coverslipped under 90% glycerol in PBS. Images were collected using a 20X objective on an Olympus BX51-WI microscope equipped with conventional fluorescence filters (Cy3:528–553 nm excitation, 590–650 nm emission; Cy2/Alexa 488: 460–500 nm excitation, 510–560 nm emission), a Hamamatsu ER, DVC high-speed camera, linear focus encoder, and a 3-axis motorized stage system interfaced with Neurolucida software (MBF Bioscience, Essex, VT, USA).

In vivo electrophysiological recording of human peripheral sensory fibers

Single-unit axonal recordings (microneurography) were performed from the right posterior antebrachial cutaneous, radial, or superficial peroneal nerve of 41 healthy participants (19 males and 22 females; 19 to 41 years). Participants were comfortably seated in an adjustable chair with legs and arms stretched out (and hand pronated), supported by vacuum pillows, and covered in a blanket if they reported as feeling cold.

Under real-time ultrasound guidance (LOGIQ P9, GE Healthcare, Chicago, IL, USA), the target nerve was impaled with an insulated tungsten recording electrode (FHC Inc., Bowdoin, ME, USA). Adjacent to that, an uninsulated reference electrode was inserted just under the skin. A high-impedance preamplifier (MLT185 headstage) was attached to the skin near the recording electrode and used together with a low-noise high-gain amplifier (FE185 Neuro Amp EX, ADInstruments, Oxford, UK). Once the electrode tip was intrafascicular, single LTMRs were

1330   searched for by soft-brush stroking, and single HTMRs were searched for by coarse-brush strok-

1331   ing, pinching, and hair tugging in the fascicular innervation zone while making minute electrode

1332   adjustments.

1333

All recorded afferents were mechanically responsive and divided into subtypes based on estab-

lished criteria[68,112,113]. Mechanical threshold and receptive field size were determined using

Semmes-Weinstein monofilaments (nylon fiber; Aesthesio, Bioseb, Pinellas Park, FL, USA).

Mechanical threshold was defined as the weakest monofilament to which the unit responded to in

at least 50% of trials. Hair deflection was tested with a small pair of forceps, carefully avoiding

skin contact while manipulating the hair. Further, force measurements were performed to ensure

that no skin/hair pulling occurred. Conduction velocity of the recorded afferent was estimated

from latency responses to surface electrical stimulation of the receptive field (FE180 Stimulus

Isolator, ADInstruments, Oxford, UK). Electrically and mechanically evoked spikes were com-

pared on an expanded time scale to confirm they originated from the same unit. Thermal respon-

siveness was tested by placing a Peltier probe (7.4 x 12.2 mm, T09, QST.Lab, Strasbourg, France)

onto the receptive field. After recording baseline activity for at least 30 s (with the thermode in

contact with the receptive field) at a neutral temperature of 30°C, a series of cooling (down to

0°C) and warming (up to 50°C) stimuli were delivered at 30-s intervals. If needed, the thermode

was mounted on a stand for better stability.

1349

To test *TRPV1* expression, capsaicin (Capsina 0.075%, Bioglan AB, Malmö, Sweden) was topi-

cally applied to the receptive field. After 1 minute, the skin was wiped clean, and the emergence

of any spontaneous spiking activity from the recorded afferent was monitored. *TRPM8* expres-

sion was tested by placing an ethanol-soaked gauze pad (90% ethanol as control) onto the recep-

tive field followed by menthol solution (400 mg of 40% L-menthol dissolved in 90% ethanol,

Sigma-Aldrich, Inc., Schnelldorf, Germany[114]). The gauze pad was covered with an adhesive

film to prevent the evaporation of ethanol. After 5 minutes, the skin was wiped clean and the

emergence of any spontaneous spiking activity from the recorded afferent was monitored. Dur-

ing these procedures, we documented the participants' verbal descriptions of what they felt, and

if there was no obvious sensation, the procedure was repeated.

1360

Neural activity was sampled at 20 kHz and recorded using the ADInstruments data acquisition

system (LabChart software v8.1.24 and PowerLab 16/35 hardware PL3516/P, Oxford, UK), then

exported to Spike2 (v10.13, Cambridge Electronic Design Ltd., Cambridge, UK). Recorded ac-

tion potentials were carefully examined offline on an expanded time scale. Threshold crossing

was used to distinguish action potentials from noise with a signal-to-noise ratio of at least 2:1,

and spike morphology was confirmed by template matching. Recordings were discarded if mul-

tiple units were present or if spike amplitudes were not distinct from the noise, preventing secure

action potential identification.

1369

1370 **Figure Generation Software**

1371 Figures were generated in Powerpoint (Microsoft Office) and GraphPad Prism (v9, GraphPad

1372 Software Inc. La Jolla, CA, USA). Some cartoons were made partially in BioRender (BioRender,

1373 2022, RRID:SCR_018361).

1374

Data availability

The raw and processed datasets for the single-soma sequencing of hDRG neurons reported in this

study will be deposited into Broad Institute Single cell portal

(https://singlecell.broadinstitute.org/single_cell) once the manuscript is accepted for publication.

Macaque (Kupari) data is available at

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE165569

Mouse (Zeisel) DRG data is available at

http://loom.linnarssonlab.org/clone/Mousebrain.org.level6/L6_Peripheral_sensory_neurons.loom.

Mouse (Sharma) DRG data is available at

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE139088

1385

1386 **Code availability**

1387 Any custom code will be deposited to Github once the manuscript is accepted for publication.

1388 All analyses are based on previously published code and software.

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

1400

1401

1402

1403

1404

1405

1406

1407

1408

Figure 1. Developing a novel laser capture microdissection (LCM) based approach for sin-

gle-soma deep RNA-seq of hDRG neurons

(**A**) A diagram shows the overall workflow of this study. (left) A cartoon illustrates features as-

sociated with different strategies for single-cell RNA-seq of hDRG neurons. (Middle) One real

example of the laser dissection of a hDRG neuron soma is shown. (Right) Analyses and experi-

ments conducted in this study are summarized. Scale bar, 50 μm (cell) and 500 μm (cap)

(**B**) UMAP plot showing the clusters of 1066 hDRG neurons. (**C-D**) Violin plots showing total

number of detected genes (C) and the expression of neuronal marker *SLC17A6* (D). (**E**) The

grouping clusters based on the soma size, and the expression of *INA*, *NEFH*, *PRDM12*, and

CALCA. (**F**) UMAPs showing some canonical marker gene expression in each cluster. (**G**)

UMAP plot with names of each cluster.

1420

Fig. 2 Cross-species analysis of DRG neurons in human, macaque, and mouse

(**A, B**) Conos label propagation from mouse (Sharma) (**A,** combined Sharma & Usoskin nomen-

clature) and macaque (B) to hDRG neuron clusters showing the cell type correlation. For

UMAPs for correspondent co-integration, from which these results were inferred, see Fig. S5A-

B. (**C**) Summary of correspondence of DRG neuron clusters among three species. Solid lines

depict clear match, and dashed lines represent partial similarity. (**D**) Heatmap visualization of

cross-species-conserved and species-specific transcription factor associated gene patterns across

mouse, macaque, and human. Species are color coded in the right column. Yellow boxes, con-

served, green boxes, species-specific gene regulatory networks.

1430

Figure 3. Marker gene expression and validation in C-fiber pruriceptors, thermoreceptors,

and nociceptors

(**A-D**) Marker genes for specific labeling of each cluster and validation by multiplex FISH for

hNP1 and hNP2 (A), hPEP.SST(B), hTRPM8 (C), and hPEP.TRPV1/A1.1 and

hPEP.TRPV1/A1.2 (D). The fluorescent images show detected transcripts in one example hDRG

neuron (cell body outlined by the white dashed line. Circle charts next to images show quantifi-

cation: the arcs indicate the percentage of neurons positive for the given marker gene over all

sampled DRG neurons. The sector shaded areas indicate the approximate percentage of each cell

type over the total quantified hDRG neurons. N=2, A (199 neurons total), B (220 neurons total),

C (156 neurons total), D (202 neurons total). Scale bar, 50 μm.

1441

Figure 4. Marker gene expression and validation in A-fiber nociceptors

(**A-D**) Marker genes for specific labeling of each cluster and validation by multiplex FISH for

hPEP.PIEZO2[h] (A), hPEP.KIT (B), hPEP.CHRNA7 (C), and hPEP.NTRK3 (D). The fluorescent

images show the detected transcripts in one example hDRG neuron (cell body outlined by the

white dashed line). Circle charts next to images show quantification: the arcs indicate the per-

centage of neurons positive for the given marker gene over all sampled DRG neurons. The sector

shaded areas indicate the approximate percentage of each cell type over the total quantified

hDRG neurons. N=2, A (165 neurons total), B (173 neurons total), C (196 neurons total), D (191

neurons total). Scale bar, 50 μm.

1451

Figure 5. Marker gene expression and validation in C- and A-LTMRs

(**A- D**) Marker genes for specific labeling of each cluster and validation by multiplex FISH for

hC.LTMR (A), hAδ.LTMR (B), hAβ.LTMR (C), hPropr (D) and hATF3 (E). The fluorescent

images show the detected transcripts in one example hDRG neuron (cell body outlined by the

white dashed line). Circle charts next to images show quantification: the arcs indicate the per-

centage of neurons positive for the given marker gene over all sampled DRG neurons. The sector

shaded areas indicate the approximate percentage of each cell type over the total quantified

hDRG neurons. N=2, A (205 neurons total), B (183 neurons total), C (188 neurons total), D (198

neurons total), E (202 neurons total). Scale bar, 50 μm.

1461

Figure 6. Single-soma deep RNA-seq dataset is powerful for molecular discovery

(**A**) Comparison of the total detected number of GPCRs, ion channels, chemokine receptors, and

neuropeptides in single-soma, spatial, and single-nucleus RNA-seq datasets. (**B-C**) Expression of

putative itch receptors in hDRG neurons. Some receptors are highly enriched (B), while some

receptors are barely detected (C) in human itch populations. (**D**) Novel GPCRs, ion channels,

and other genes enriched in human itch populations.

1468

Figure 7. Molecular profile-informed single-unit microneurography recordings revealed novel physiological properties of a population of human A-HTMRs.

(**A**) Predicted physiological properties of hPEP.KIT sensory afferents based on single-soma deep RNA-seq data. (**B**) Receptive field locations of recorded single afferents from superficial peroneal (S. peroneal), posterior antebrachial cutaneous (PABCN), and radial nerve recordings (n = 47). (**C**) Distribution of mechanical (monofilament) thresholds for HTMRs and LTMRs in the recorded samples. (**D**) Individual and mean (±SEM) conduction velocities of different HTMR and LTMR types in response to surface electrical stimulation from upper and lower limbs (Field-LTMR: 40.3 ± 4.2 m/s, n=2; SA1-LTMR: 44.9 ± 2.6 m/s, n=3; SA2-LTMR: 44.9 ± 1.2 m/s, n=3; A-HTMR cooling-: 50.6 ± 4.8 m/s, n=5; A-HTMR cooling+: 48.9 ± 5.0 m/s, n=3; C-LTMR: 1.0 ± 0.05 m/s, n=8; C-HTMR: 0.7 ± 0.08 m/s, n=5). (**E**) Spike activities of a putative hPEP.KIT unit (A-HTMR cooling+) in response to repeated stimulations of the receptive field, superimposed on an expanded time scale. (**F-H**) Responses of an A-HTMR cooling+ unit to soft and coarse brushing (F), heating (G) and cooling (H). (**I**) Schematic showing the link between the histologically identified KIT$^+$/CGRP$^+$/NEFH$^+$ sensory afferents and the molecularly defined hPEP.KIT population, likely representing a type of heat/capsaicin-insensitive but cold-sensitive A-HTMRs.

1486

Figure 8. Molecular profile-informed single-unit microneurography recordings revealed novel physiological properties of a population of human C-LTMRs.

(**A**) Novel physiological properties of hC.LTMR sensory fibers predicted based on gene expression obtained from the single-soma deep sequencing. (**B**) Spike activity of a hC.LTMR unit in

response to repeated stimulations of the receptive field, superimposed on an expanded time scale.

(**C**) Individual and mean (±SEM) conduction velocities of different HTMR and LTMR types in

response to surface electrical stimulation from upper and lower limbs (the same plot from Fig.

7D) (**D-H**) Responses of a hC.LTMR unit to soft brushing and hair movement (D), cooling (E),

menthol (F), heating (G) and capsaicin (H). Spike activity of that hC.LTMR before and after

capsaicin application, overlaid on an expanded timescale (H). Conduction delay was adjusted

based on the latency of electrically triggered spiking for that recorded afferent. Note, different

scaling.

Figure 1

Figure 2

Figure 3

A | hNP1/hNP2

MRGPRX1 — HRH1

MRGPRX1 / HRH1 / Merged

Marker gene: MRGPRX1
Potential functions: Itch sensing, polymodal

38.7%
16.1%
24.1%

B | hPEP.SST

SST — IL31RA

SST / IL31RA / Merged

Marker gene: SST
Potential function: Inflammation, itch

30.5%
13.2%
17.3%

C | hTRPM8

TRPM8 — STUM

TRPM8 / STUM / Merged

Marker gene: STUM
Potential function: Cold sensing, polymodal

13.5%
12.2%
26.8%

D | hPEP.TRPV1/A1.1-hPEP.TRPV1/A1.2

TRPV1 — TRPA1

TRPV1 / TRPA1 / Merged

Marker gene: TRPV1+TRPA1
Potential function: Thermal and chemical pain

20.8%
19.3%
52.0%

Figure 4

Figure 5

Figure. 6

Figure 7

Figure 8

Supplementary Figure 1

A Cryo-sections → 70% EtOH (30s) → HistoGene™ staining (10s) → 70% EtOH (30s) → 95% EtOH (30s) → 100% EtOH (30s) → LCM

B

C

D **All DRG neurons**

Frequency (%) vs Diameter (µm)

E **Dissected DRG neurons**

Frequency (%) vs Diameter (µm)

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5

Supplementary Figure 6

Supplementary Figure 7

Supplementary Figure 8

Supplementary Figure 9

Supplementary Figure 10

Supplementary Figure 11

Supplementary Figure 12

Supplementary Figure 13

Supplementary Figure 14

Supplementary Figure 15

A — hPEP.KIT

B — Cooling

C — Heating