Control of meiotic entry by dual inhibition of a key mitotic transcription factor

Amanda J. Su[1], Siri C. Yendluri[1] and Elçin Ünal[1, §]

[1] Department of Molecular and Cell Biology, University of California, Berkeley, CA,

United States, 94720

[§] Lead contact: elcin@berkeley.edu

Keywords: meiosis, mitosis, transcription, cell cycle, LUTI, uORF, Ime1, Swi4, SBF,

Whi5, Rb, E2F

Abbreviations: SBF (Swi4/6 cell cycle box [SCB] Binding Factor); MBF (Mlu1 cell cycle

box [MCB] Binding Factor); LUTI (Long Undecoded Transcript Isoform); TF

(Transcription Factor)

HIGHLIGHTS

● SBF subunit Swi4 is downregulated at meiotic entry by the LUTI-based mechanism

● $SWI4^{LUTI}$ and Whi5 work together to inhibit SBF at meiotic entry

● SBF and Ime1 TFs antagonize each other, ensuring mutually exclusive cell states

● SBF-target G1 cyclins block meiosis by preventing Ime1-Ume6 interaction

SUMMARY

The mitosis to meiosis transition requires dynamic changes in gene expression, but whether and how the mitotic transcriptional machinery is regulated during this transition is unknown. In budding yeast, SBF and MBF transcription factors initiate the mitotic gene expression program. Here, we report two mechanisms that work together to restrict SBF activity during meiotic entry: repression of the SBF-specific Swi4 subunit through LUTI-based regulation and inhibition of SBF by Whi5, a functional homolog of the Rb tumor suppressor. We find that untimely SBF activation causes downregulation of early meiotic genes and delays meiotic entry. These defects are largely driven by the SBF-target G1 cyclins, which block the interaction between the central meiotic regulator Ime1 and its cofactor Ume6. Our study provides insight into the role of $SWI4^{LUTI}$ in establishing the meiotic transcriptional program and demonstrates how the LUTI-based regulation is integrated into a larger regulatory network to ensure timely SBF activity.

INTRODUCTION

A key aspect in understanding developmental programs and cell state transitions is mapping the interplay between transcription factors and their associated gene regulatory networks. In the budding yeast *Saccharomyces cerevisiae*, the transition from mitotic growth to meiotic differentiation is a crucial decision that is regulated by multiple inputs, such as nutrient availability, respiration competence, and cell identity. Under nutrient-limiting conditions, a diploid cell enters the meiotic program to produce four haploid gametes. The process of meiotic entry is tightly controlled by the master transcriptional regulator Ime1, as both extrinsic (e.g. nutrient status, extracellular pH) and intrinsic (e.g. mating type, mitochondrial function) cues are integrated at the *IME1* promoter (Kassir et

al., 1988; Honigberg, 2003; reviewed in van Werven and Amon, 2011). Once translated,

Ime1 is phosphorylated by the Rim11 and Rim15 kinases to promote its nuclear

localization and interaction with Ume6 (Vidan and Mitchell, 1997; Pnueli et al., 2004;

Malathi et al., 1999, 1997). In mitotically dividing cells, Ime1 target genes are repressed

by Ume6 through its association with the Sin3-Rpd3 histone deacetylase complex

(Kardosh and Struhl, 1997; Rundlett E. et al., 2003). However, under nutrient starvation,

entry into the meiotic program is initiated by the interaction between Ime1 and Ume6,

which together function as a transcriptional activator, culminating in the induction of early

meiotic genes (Bowdish et al., 1995; Harris and Ünal, 2023). Mitosis to meiosis transition

requires dynamic remodeling of the gene regulatory networks to maintain the mutual

exclusivity of these programs. While entry into the mitotic program is initiated by the

central transcription factors SBF and MBF (Spellman et al., 1998; Iyer et al., 2001),

whether and how these complexes are regulated during meiotic entry is unknown.

The molecular mechanisms regulating entry into the mitotic cell cycle, also known as

G1/S transition, are functionally conserved from yeast to metazoans (reviewed in Van

Den Heuvel and Dyson, 2008). SBF and MBF are heterodimeric transcription factors

composed of Swi4-Swi6 and Mbp1-Swi6 subunits, respectively (Figure 1A). Though there

is no sequence homology these transcription factors are functionally homologous to the

mammalian E2Fs. E2Fs are negatively regulated by the tumor suppressor protein Rb,

which is homologous to the budding yeast Whi5 that inhibits SBF in early G1 (De Bruin

et al., 2004; Costanzo et al., 2004; Hasan et al., 2014). Whi5-based SBF inhibition is

relieved by cyclin/CDK-dependent phosphorylation and subsequent re-localization of

Whi5 from the nucleus to the cytoplasm (Wagner et al., 2009). Although SBF and MBF

act in parallel during the G1/S transition (Spellman et al., 1998; Iyer et al., 2001), they

activate functionally specialized subsets of gene targets. SBF regulates the expression

of genes involved in budding and cell morphogenesis, while MBF-regulated genes are

involved in DNA replication and repair (Iyer et al., 2001; Simon et al., 2001). Despite the

well-characterized function in budding yeast mitotic growth, the regulation and function of

SBF and MBF during meiotic entry remains largely unknown. In contrast to mitotic

divisions where budding and DNA replication occur simultaneously, the meiotic program

requires DNA replication and repair, but not bud morphogenesis or asymmetric growth.

Accordingly, the regulation and subsequent activity of SBF and MBF are likely to be

divergent to establish meiotic entry.

The activity of SBF and MBF is regulated in part through subunit abundance, which in

turn controls expression of the G1/S regulon (Dorsey et al., 2018). For example,

overexpression of a hyperactive allele of *SWI4* can trigger premature entry into the cell

cycle (Sidorova and Breeden, 2002). Additionally, *SWI4* has also been shown to be

haploinsufficient and rate limiting during G1/S progression (Mcinerny et al., 1997). These

results suggest that the precise levels of SBF and MBF subunits are important for

activating the G1/S regulon at the correct time.

Three key observations support differential regulation of SBF and MBF during the meiotic

program: First, a meiotic mass spectrometry dataset suggests that Swi4 has dynamic

protein behavior, which is not observed for its counterparts Swi6 and Mbp1 (Cheng et al.,

2018). Second, meiotic cells express a non-canonical mRNA from the *SWI4* locus called

LUTI (Brar et al., 2012; Tresenrider et al., 2021), which stands for Long Undecoded

Transcript Isoform (Chen et al., 2017; Chia et al., 2017). Third, two SBF-specific targets,

namely the G1-cyclins *CLN1* and *CLN2*, have been shown to repress early meiotic gene

expression (Colomina et al., 1999).

LUTI-based gene regulation repurposes gene-activating transcription factors as

repressors by a two-pronged mechanism (Chen et al., 2017; Chia et al., 2017; Tresenrider

and Ünal, 2017; Tresenrider et al., 2021). First, transcription factor-dependent activation

of the LUTI promoter results in co-transcriptional silencing of the downstream canonical

gene promoter. Second, the coding sequence (CDS) within the LUTI is translationally

repressed due to competitive translation of the upstream open reading frames (uORFs)

in the LUTI-specific 5′ leader. Consequently, upregulation of the LUTI results in

downregulation of the canonical mRNA and corresponding protein. While LUTIs are

conserved from yeast to humans and have been identified in different cellular contexts

(Cheng et al., 2018; Van Dalfsen et al., 2018; Hollerer et al., 2019; Jorgensen et al., 2020),

their biological significance remains poorly understood. In fact, only a single LUTI has

been assigned a biological role so far with distinct phenotypic outcomes resulting from its

loss or gain of function (Chen et al., 2017; Chia et al., 2017).

In this study, we aimed to investigate the differential regulation of SBF and MBF during

the meiotic program and determine how the untimely activation of SBF impacts meiotic

entry. We found that overexpression of Swi4 results in the activation of SBF targets and

concomitant downregulation of the early meiotic genes. SBF targets Cln1 and Cln2 inhibit

meiotic entry by blocking the interaction between Ime1 and its cofactor Ume6.

Furthermore, the LUTI-based mechanism causes downregulation of Swi4 protein

synthesis and acts in conjunction with Whi5 to restrict SBF activity during meiosis.

Overall, our study reveals the functional role of a LUTI in establishing the early meiotic

transcriptome, demonstrates how the LUTI-based regulation is integrated into a larger

regulatory network to ensure timely SBF activity, and provides mechanistic insights into

how SBF misregulation impedes transition from mitotic to meiotic cell fate.

RESULTS

Swi4 is the sole downregulated subunit within the SBF and MBF complexes during

meiosis

While the SBF and MBF transcription factors have been heavily studied in the context of

the mitotic cell cycle, their involvement in regulating the meiotic transcriptional program is

not well understood. To understand the regulation and function of these complexes during

meiosis, we first monitored the levels of SBF and MBF subunits throughout a meiotic time

course. Cells were first grown in rich media overnight and were then transferred to pre-

sporulation media. After additional overnight growth, cells were shifted to sporulation

media (SPO) to induce meiosis, and samples were taken hourly for protein extraction and

immunoblotting to monitor the abundance of each subunit. Unlike the mitotic G1/S

transition (Kelliher et al., 2018), meiotic entry resulted in ~30% decrease in Swi4 levels

after 2 hours, while Mbp1 and Swi6 levels were increased (Figure 1B and 1C*). These

data indicate that Swi4 is the sole subunit within SBF and MBF whose level declines

during meiotic entry and are consistent with a published mass spectrometry dataset

(Cheng et al., 2018).

We next examined the expression of a set of well-characterized transcripts that are

regulated by SBF or MBF (Iyer et al., 2001; Simon et al., 2001; Bean et al., 2005; Harris

et al., 2013; Smolka et al., 2012) using a published meiotic RNA-seq dataset (Brar et al.,

2012). This analysis revealed that compared to mitotically dividing cells, most of the SBF-

specific target genes had either low or no expression during early meiosis (Figure 1D). In

contrast, most of the MBF targets displayed increased expression upon meiotic entry

(Figure 1D).

Based on our findings thus far, we propose that Swi4 levels are downregulated in meiosis

to ensure that SBF targets, including *CLN1* and *CLN2*, are turned off during meiotic entry.

If so, then higher Swi4 levels should lead to increased expression of SBF targets. To test

this possibility, we overexpressed *SWI4* by placing it under the regulation of the *ATG8*

promoter (*pATG8-SWI4*), which is highly expressed in meiosis (Brar et al., 2012). The

steady-state level of *pATG8*-driven Swi4 was 5 times higher than wild type (Figure S1A

and S1B). Using reverse transcription coupled with quantitative polymerase chain

reaction (RT-qPCR), we measured the transcript levels of well-characterized SBF targets

and observed a significant increase in the expression of *CLN1* and *CLN2* in *pATG8-SWI4*

cells relative to wild type (Figure 1E, $p = 0.0351$ [*CLN1*], $p = 0.0013$ [*CLN2*], two-tailed t-

test). In contrast, the MBF-specific targets *CDC21* and *RNR1* remained similar (Figure

1E, $p = 0.8488$ [*CDC21*], $p = 0.0859$ [*RNR1*], two-tailed t-test). These findings indicate

that upregulation of *SWI4* is sufficient to induce the expression of SBF-specific targets

CLN1 and *CLN2* in meiosis without affecting MBF-specific targets.

To test whether the downregulation of *SWI4* is functionally important for meiotic

progression, we used time-lapse fluorescence microscopy and visualized the kinetics of

meiotic divisions in *pATG8-SWI4* cells relative to wild type. By tracking the endogenous

histone H2B fused to the red fluorescent protein mCherry (Htb1-mCherry), we found that

SWI4 overexpression caused a significant delay in meiotic progression (Figure 1F, p =

0.0045, Mann-Whitney test). We conclude that downregulation of *SWI4* is necessary for

timely meiotic progression.

Regulation of Swi4 abundance is required for timely meiotic entry

Since Swi4 abundance, and by inference SBF activity, was decreased during the mitosis-

to-meiosis transition, we hypothesized that the meiotic progression delay observed in

pATG8-SWI4 cells was due to meiotic entry defects. To test this we monitored Ime1, a

meiosis-specific transcription factor and a master regulator of meiotic entry (Kassir et al.,

1988; Honigberg, 2003; reviewed in van Werven and Amon, 2011). To quantify the bulk

levels of Ime1 protein in meiosis, we performed immunoblotting. During meiotic entry (2

h in SPO), when Swi4 abundance was elevated, there was a 50% reduction in Ime1 levels

in *pATG8-SWI4* cells compared to wild type (Figure S1C and S1D). In parallel, we

monitored meiotic entry on a single-cell basis by measuring the localization of

endogenous Ime1 carrying an N-terminal green fluorescent protein tag (*GFP-IME1*;

Moretto 2018) and Htb1-mCherry. Compared to wild type where > 90% of the cells had

nuclear Ime1 following meiotic entry (2 h in SPO), only ~50% of *pATG8-SWI4* cells had

nuclear Ime1, which was significantly lower (Figure 2A and 2B, p = 0.0169, two-tailed t-

test).

Given that the increase in Swi4 levels coincided with a decrease in Ime1 protein

expression and nuclear localization, this raised the possibility that Swi4 interferes with

Ime1 function. To further investigate the relationship between these two transcription

factors at a single-cell level, we generated strains carrying endogenous fluorescent

protein tags for each transcription factor (*GFP-IME1, SWI4-mCherry*). Using DAPI as a

nuclear marker, we measured the mean nuclear intensity of each transcription factor

before (0 h) and after meiotic entry (4 h) (see Materials and Methods for details on image

quantification). In wild type, most cells exhibited decreased nuclear Swi4 and increased

nuclear Ime1 upon meiotic entry (Figure 2C-E). Conversely, in the *pATG8-SWI4* mutant,

there was a significant shift in the fraction of cells with higher nuclear Swi4 levels (Figure

2D and 2E, p < 0.0001, Mann-Whitney test,). Additionally, the cells with increased nuclear

Swi4 had reduced levels of nuclear Ime1 (Figure 2E, p < 0.0001, Mann-Whitney test).

This inverse relationship between Swi4 and Ime1 nuclear localization further indicates

that higher levels of Swi4 are antagonistic to Ime1 function.

Ime1 is necessary for the expression of early meiotic genes. Therefore, we next

characterized the changes in the early meiotic transcriptome upon *SWI4* overexpression

by mRNA-sequencing (mRNA-seq). Using DESeq2 (Love et al., 2014), we identified

differentially expressed genes in *pATG8-SWI4* mutant compared to wild type during

meiotic entry (2 h in SPO). To investigate general pathways being affected by increased

Swi4 levels in meiosis, we ran gene ontology analysis of statistically significant (padj <

0.05) output from DESeq2, which revealed that the genes with significantly increased

expression were involved in mitosis while those with significantly decreased expression

were involved in meiotic processes (Figure S2A and S2B). Regarding the SBF targets,

we noticed increased expression of many genes, including *CLN1* and *CLN2* (Figure 2F,

Supplemental Table 1). We focused on a cluster of previously identified early meiotic

genes involved in DNA replication and recombination (Brar et al., 2012) and observed

that more than 50% had a significant decrease in their expression upon *SWI4*

overexpression (padj < 0.05, Figure 2F, Supplemental Table 1). Finally, gene set

enrichment analysis was performed to test whether expression of either the early meiotic

gene set or SBF target gene set is enriched in wild type or *pATG8-SWI4* mutant cells

(Subramanian et al., 2005). This gene set level analysis revealed significant enrichment

of SBF target gene expression (NES = 2.88, p < 0.001, Figure 2G), as well as significant

disenrichment of early meiotic gene expression (NES = -2.88, p < 0.001, Figure 2G) in

the *pATG8-SWI4* mutant. Altogether, these findings demonstrate that the increased levels

of Swi4 during transition from mitotic to meiotic cell fate abruptly activates SBF and

disrupts the early meiotic transcriptome, highlighting the importance of *SWI4* regulation.

As an orthogonal approach, we performed live-cell imaging of Rec8, endogenously

tagged with GFP (Rec8-GFP), which is a meiosis-specific cohesin subunit and a direct

transcriptional target of Ime1 (Primig et al., 2000; Klein et al., 1999). Htb1-mCherry was

used as a nuclear marker. This analysis revealed a significant delay in Rec8-GFP nuclear

appearance in the *pATG8-SWI4* strain compared to wild type (Figure 2H and 2I, p =

0.0005, Mann-Whitney test). Furthermore, sporulation efficiency was decreased by ~20%

in the *pATG8-SWI4* strain relative to wild type (Supplemental Table 2). These findings

are consistent with the mRNA-seq data and further underscore the importance of *SWI4*

downregulation in establishing a robust meiotic cell fate.

Removal of the SBF targets Cln1 or Cln2 partially rescues the meiotic entry delay

in *pATG8-SWI4* mutants

Overexpression of the G1 cyclins *CLN1*, *CLN2* and *CLN3* has been previously shown to

inhibit meiotic entry (Colomina et al., 1999). Given that *CLN1* and *CLN2* are

transcriptional targets of SBF, we next determined whether the meiotic progression delay

in *pATG8-SWI4* cells could be due to increased G1 cyclin protein levels. We first

examined Cln1 and Cln2 protein levels using epitope-tagged alleles at their endogenous

loci (*CLN1-3V5* or *CLN2-3V5*) in wild-type and *pATG8-SWI4* cells. In response to *SWI4*

overexpression, we observed up to 2-fold and 10-fold increase in Cln1 and Cln2 protein

levels, respectively, corroborating the mRNA-seq data (Figure 3A-D).

To determine whether these two G1 cyclins are functionally responsible for the meiotic

progression delay observed in response to SBF misregulation, we performed time-lapse

fluorescence microscopy in cells carrying Rec8-GFP and Htb1-mCherry. We found that

deletion of either *CLN1* or *CLN2* significantly rescued the meiotic progression delay in the

pATG8-SWI4 mutant (p = 0.0111 [*cln1Δ*], p = 0.0478 [*cln2Δ*], Mann-Whitney test) (Figure

3E and 3F). The kinetics of meiotic entry in the absence of either Cln1 or Cln2 closely

resembled that of wild type (Figure S3A), indicating that the observed rescue was specific

to the meiotic defect resulting from elevated Swi4 levels. However, *pATG8-SWI4* cells

lacking either *CLN1* or *CLN2* were still delayed compared to wild type, suggesting

redundancy. We were unable to examine meiotic progression in the *cln1Δ cln2Δ* double

mutant due to its severe sickness in the SK1 background. Nevertheless, our analyses

establish a causal link demonstrating that both *CLN1* and *CLN2* contribute to the meiotic

defects arising from SBF misregulation.

Tethering of Ime1 to Ume6 is sufficient to overcome the meiotic block exerted by

G1 cyclin overexpression

Given the partial rescue of the meiotic delay in *pATG8-SWI4* mutants by the deletion of

individual G1 cyclins, we next investigated how the G1 cyclins interfere with meiosis. To

this end, we generated transgenes that expressed either *CLN1* or *CLN2* under the control

of the *pATG8* promoter for meiotic overexpression. While wild-type cells successfully

completed meiosis with more than 94% sporulation efficiency, *CLN2* overexpression

resulted in only 8.5% of cells forming gametes. Overexpression of *CLN1* also resulted in

a meiotic defect, albeit to a lesser extent than the *pATG8-CLN2* mutant (65% sporulation

efficiency, Figure 4A).

Since the meiotic defect was more pronounced in response to *CLN2* overexpression, we

decided to use the *pATG8-CLN2* mutant to explore how G1 cyclins counteract meiosis.

To assess meiotic entry, we performed fixed-cell imaging for GFP-Ime1 and found that

when *CLN2* was overexpressed, only 5% of the *pATG8-CLN2* cells displayed nuclear

GFP-Ime1 signal during early meiosis as opposed to >84% of wild-type cells (Figure 4B

and 4C, 2 h in SPO). In parallel, we measured *IME1* transcript and Ime1 protein levels.

In both cases, we observed ~30% decrease in abundance in the *pATG8-CLN2* mutant

relative to wild type (Figure S4A-C, 2 h in SPO). These data raise the possibility that the

meiotic entry defect observed in the *pATG8-CLN2* mutant arises from downregulation of

IME1 expression.

To test whether the increased levels of Ime1 can rescue the meiotic defect of *pATG8-*

CLN2 mutant, we replaced the endogenous *IME1* promoter with a copper-inducible *CUP1*

promoter (Figure 4D). We adapted a previously well-characterized overexpression allele

of *IME1*, *pCUP1-IME1,* (Berchowitz et al., 2013; Chia and van Werven, 2016) and

included a functional, N-terminal GFP tag to track Ime1's subcellular localization *(pCUP1-*

GFP-IME1). Use of the *CUP1* promoter was successful in elevating *IME1* transcript and

protein levels in the presence of *CLN2* overexpression (Figure S4D-F). Despite the

rescue of *IME1* expression, gamete formation was still severely perturbed (Figure 4A).

Using a single z-slice to measure mean nuclear intensity, we noticed that the intensity of

the nuclear Ime1 signal was significantly lower in the *pATG8-CLN2; pCUP1-GFP-IME1*

cells compared to the *pCUP1-GFP-IME1* control (Figure 4E and 4F, p < 0.0001, Mann-

Whitney test). This finding indicates that rescue of *IME1* expression did not also rescue

sporulation, further suggesting a defect in Ime1 nuclear localization. To test this

possibility, we utilized a nanobody trap strategy (Fridy et al., 2014) where we C-terminally

fused a single-domain anti-GFP antibody to Pus1, a constitutively nuclear localized

protein (*PUS1-αGFP*, Figure 4G). In this background, control strains carrying a *GFP-IME1*

allele sporulated efficiently, demonstrating that tethering of Ime1 to Pus1 does not

interfere with Ime1 function (Figure 4A). Furthermore, mean nuclear intensity of GFP-

Ime1 was indistinguishable between *PUS1-αGFP* and *pATG8-CLN2; PUS1-αGFP*,

indicating that nuclear localization was fully rescued (Figure 4H and 4I, p = 0.6563 [*PUS1-*

αGFP vs. *pATG8-CLN2; PUS1-αGFP*], Mann-Whitney test). Surprisingly, these cells still

failed to undergo meiosis (Figure 4A), suggesting that G1 cyclins interfere with Ime1

function at an additional step beyond misregulating its expression and localization.

Alternatively, G1 cyclins could disrupt a different meiotic factor in addition to Ime1.

To induce early meiotic genes, Ime1 must interact with another transcription factor called

Ume6 (Rubin-Bejerano et al., 1996). Since Ime1 itself does not possess a DNA-binding

domain, its binding to Ume6 is essential for targeting Ime1 to early meiotic gene

promoters (Smith et al., 1993; Rubin-Bejerano et al., 1996). To address whether the G1

cyclins might disrupt the interaction between Ime1 and Ume6, we fused the anti-GFP

nanobody to Ume6 (*UME6-αGFP*) in the *pATG8-CLN2* strain carrying a GFP tagged Ime1

(Figure 4J). This nanobody trap should lead to constitutive tethering of Ime1 to Ume6, as

evidenced by the rescue of Ime1 nuclear localization (Figure 4K and 4L, p = 0.035 [wild

type vs. *pATG8-CLN2 UME6-αGFP*] Mann-Whitney test). Under these conditions, the

sporulation defect of *pATG8-CLN2* mutant was rescued, reaching similar levels to wild

type (Figure 4A). Since *IME1* is expressed from its endogenous promoter in the *pATG8-*

CLN2; UME6-αGFP strain, these data suggest that overexpression of G1 cyclins results

in meiotic failure due to reduced Ime1-Ume6 interaction.

Our findings thus far highlight the biological significance of restricting SBF activity during

the transition from mitotic to meiotic cell fate. Among the SBF targets, G1 cyclins pose a

major block to meiotic entry by interfering with Ime1 function, critically at the level of Ime1-

Ume6 interaction. While our findings emphasize the importance of *SWI4* regulation to

ensure timely SBF activity, the question remains as to how SBF activity is downregulated

during transition from mitosis to meiosis.

Ime1-dependent expression of a LUTI from the *SWI4* locus leads to a reduction in

Swi4 protein levels during meiotic entry

Given the importance of *SWI4* downregulation in restricting SBF activity during mitotic to

meiotic cell fate transition, we next investigated the mechanism of *SWI4* downregulation.

A previous study identified a long undecoded transcript isoform (LUTI) expressed from

the *SWI4* locus in meiotic cells (Brar et al., 2012, Figure 5A). When *SWI4^{LUTI}* is expressed,

the canonical protein-coding *SWI4* transcript, *SWI4^{canon}*, is downregulated (Tresenrider et

al., 2021, Figure 5B), suggesting that LUTI expression restricts Swi4 protein levels and

thus SBF activity during meiotic entry.

To further investigate LUTI-based repression of *SWI4*, we examined the relationship

between *SWI4^{LUTI}* and *SWI4^{canon}* transcripts using single molecule RNA fluorescence in

situ hybridization (smFISH: Chen et al., 2017, 2018; Raj et al., 2008). Two distinct probes

were used: one, conjugated to Quasar 670 (Q670) and complementary to the *SWI4*

coding sequence (CDS) and the other, conjugated to CAL Fluor Red 590 (CF590) and

unique to the 5′ extended LUTI sequence. Accordingly, a spot where the two probe sets

colocalized indicated a $SWI4^{LUTI}$ transcript, whereas a spot marked with Q670 probe

alone highlighted a $SWI4^{canon}$ transcript. We used a well-established meiotic cell

synchronization system (Berchowitz et al., 2013) to investigate the precise temporal

expression of these two mRNA isoforms. In comparison to premeiotic state, meiotic cells

displayed a significant increase in $SWI4^{LUTI}$ transcripts (p < 0.0001, Mann-Whitney test)

as well as a significant decrease in $SWI4^{canon}$ transcripts (p = 0.0007, Mann-Whitney test)

(Figure 5C and 5D), thus confirming their inverse expression pattern.

To characterize the functional contribution of the LUTI to $SWI4$ downregulation, we

eliminated $SWI4^{LUTI}$ production ($\Delta LUTI$) by deleting its promoter (Tresenrider et al., 2021)

and used RNA blotting to visualize the $SWI4$ mRNA isoforms. In wild-type cells, $SWI4^{LUTI}$

was readily detectable after 1 hour in SPO, corresponding to early meiotic entry (Figure

5E). During the time points when $SWI4^{LUTI}$ was highly expressed, $SWI4^{canon}$ transcript

levels were lower, consistent with the smFISH data. In $\Delta LUTI$ cells, we observed an

increase in the abundance of $SWI4^{canon}$ mRNA (Figure 5E). This finding corroborates

previous reports, where LUTI expression leads to co-transcriptional silencing of the

canonical gene promoter, thereby silencing expression of the protein-coding transcript

(Chen et al., 2017; Chia et al., 2017; Van Dalfsen et al., 2018; Tresenrider et al., 2021;

Vander Wende et al., 2022). Finally, by immunoblotting, we observed an increase in Swi4

protein abundance in the $\Delta LUTI$ mutant compared to wild type (Figure 5E and 5F), further

indicating that $SWI4^{LUTI}$ expression downregulates Swi4 protein levels in meiosis.

SWI4^{LUTI} translation occurs within the upstream ORFs (uORFs) of its 5′ leader sequence

(Brar et al., 2012). If the uORFs repress productive translation of the *SWI4* CDS contained

within the LUTI, then their removal should result in increased Swi4 protein levels. To

determine whether the uORFs inhibit translation of the *SWI4* CDS, we mutated the start

codon of all seven uORFs within *SWI4^{LUTI}* from ATG to ATC (Δ*uORF*) and measured Swi4

protein levels using immunoblotting. Compared to wild type, the Δ*uORF* mutant had

higher levels of Swi4 protein (Figure 5G and 5H). RNA blotting confirmed that *SWI4^{canon}*

transcript levels remained similar between wild type and Δ*uORF* mutant, whereas

SWI4^{LUTI} levels were higher in the Δ*uORF* mutant compared to wild type (Figure 5G),

likely resulting from increased transcript stability due to bypass of nonsense mediated

decay (Tresenrider et al., 2021). Our findings indicate that the uORFs within the LUTI are

translated at the expense of *SWI4* CDS, thus halting Swi4 synthesis during early meiosis.

Our previous findings highlight an antagonistic relationship between Swi4 and Ime1

nuclear localization, whereby overexpression of *SWI4* leads to a decrease in Ime1

nuclear localization (Figure 2E). Given the previous finding that Ime1 activates

transcription of LUTIs in early meiosis (Tresenrider et al., 2021), we were curious whether

the reverse regulation could also occur, where Ime1 causes downregulation of Swi4

through activating *SWI4^{LUTI}* transcription. To test this possibility, we used an inducible

allele of *IME1* (*pCUP1-IME1*) (Chia and van Werven, 2016) and measured Swi4 protein

abundance. In the absence of *IME1* induction, Swi4 levels remained constant. However,

after 1 hour of *IME1* induction, Swi4 levels started to decrease dramatically (Figure 5I and

5J). Therefore, reduction in Swi4 levels is dependent on Ime1 rather than being driven by

nutrient deprivation in SPO, a condition that is known to trigger autophagy (Abeliovich

and Klionsky, 2001). To further assess whether $SWI4^{LUTI}$ is regulated by the Ime1-Ume6

transcription factor complex, we analyzed a published Ume6 ChIP-seq dataset

(Tresenrider et al., 2021) and found evidence for Ume6 binding at the $SWI4^{LUTI}$ promoter

(Figure 5K). Additionally, analysis of an mRNA-seq dataset from *UME6-T99N*

(Tresenrider et al., 2021), an allele of *UME6* that can no longer interact with Ime1 revealed

a dramatic reduction in $SWI4^{LUTI}$ expression in meiotic conditions (Figure S5). Together,

these data support the notion that Ime1-Ume6 complex induces the expression of

$SWI4^{LUTI}$, which in turn inhibits Swi4 protein synthesis through the combined act of

transcriptional and translational interference.

$SWI4^{LUTI}$ is integrated into a larger regulatory network to regulate SBF activity

during meiotic entry

Since the removal of $SWI4^{LUTI}$ resulted in an increase in Swi4 levels, we next wanted to

investigate how the loss of LUTI-based regulation affects meiotic entry. We performed

mRNA-seq in Δ*LUTI* mutant or wild-type cells during meiotic entry (2 h in SPO) and used

DESeq2 to identify differentially expressed genes. To our surprise, there was no obvious

increase in the expression of many SBF targets (Figure 6A). Namely, *CLN1* and *CLN2*

were both expressed at similar levels to wild type upon loss of $SWI4^{LUTI}$. Therefore, it

appears that disruption of the LUTI-based regulation alone is not sufficient to reactivate

SBF targets.

Since transition from the mitotic to meiotic program is a critical cell fate decision, it is likely

that additional players are in place to restrict SBF activity. In this case, even when the

LUTI-based regulation fails, SBF would remain largely inactive due to a backup

mechanism. In support of this notion and consistent with a previous report (Argüello-

Miranda et al., 2018), the SBF inhibitor Whi5 was expressed during early meiosis (0 h in

SPO) and localized to the nucleus (Figure S6A). During the mitotic G1 phase, Whi5

associates with promoter-bound SBF, thereby preventing the transcription of SBF target

genes (De Bruin et al., 2004; Costanzo et al., 2004). Shortly before the G1/S transition,

phosphorylation of Whi5 via the Cln3/CDK pathway activates SBF by promoting nuclear

export of Whi5 (De Bruin et al., 2004; Costanzo et al., 2004). However, in nutrient-

deprived conditions that favor meiosis such as nitrogen limitation, *CLN3* is translationally

repressed and any Cln3 protein that is expressed is also unstable (Parviz and Heideman,

1998; Gallego et al., 1997). Without Cln3/CDK, Whi5 is expected to inhibit SBF in a

constitutive manner.

To determine whether Whi5 and *SWI4^{LUTI}* act in parallel to restrict SBF activity, we

removed Whi5 from the nucleus using the anchor-away method, which enables

compartment-specific depletion of a target protein via inducible dimerization (Haruki et

al., 2008). Whi5 was tagged with FRB and ribosomal protein Rpl13a was tagged with

FKBP12. Upon addition of rapamycin, Rpl13a-FKBP12 and Whi5-FRB formed a

heterodimer, leading to successful nuclear exclusion of Whi5 (Figure 6B). Using Whi5

anchor away (*WHI5-AA)* alone or in combination with removal of LUTI-based *SWI4*

regulation (Δ*LUTI)*, we performed mRNA-seq to measure expression of SBF targets and

early meiotic genes. Similar to the Δ*LUTI* single mutant, *WHI5-AA* alone did not

significantly change the expression of SBF targets or early meiotic genes relative to wild

type (Figure 6C). However, loss of both modes of regulation resulted in increased

expression of SBF targets as well as a concomitant decrease in early meiotic transcripts

compared to wild type (Figure 6D, Supplemental Table 3). As expected, Swi4 levels were

similar to wild type in *WHI5-AA* but were elevated in Δ*LUTI* and Δ*LUTI*; *WHI5-AA* mutants

(Figure S6B). Gene set enrichment analysis revealed a significant enrichment of the SBF

regulon (NES = 1.95, p < 0.001) as well as significant disenrichment of the early meiotic

genes (NES = -3.39, p < 0.001) in the Δ*LUTI*; *WHI5-AA* double mutant (Figure S6C).

Finally, we monitored meiotic entry using time-lapse fluorescence microscopy in strains

carrying endogenously tagged Rec8-GFP in conjunction with Htb1-mCherry. In

agreement with our mRNA-seq data, loss of either LUTI- or Whi5-based repression of

SBF alone was not sufficient to cause a delay in meiotic entry (Figure 6E). However,

simultaneous perturbation of both pathways led to a significant delay in meiotic entry

(Figure 6E, p = 0.0112, Mann-Whitney test). These data further support the notion that

SWI4^{LUTI} is integrated into a larger regulatory network to regulate SBF activity during

meiotic entry, which includes Whi5-mediated repression of SBF.

DISCUSSION

In this study we describe two distinct mechanisms by which meiotic cells inhibit the activity

of SBF, a transcription factor complex crucial for the G1/S transition during the mitotic cell

cycle. The dual inhibition of SBF converges on its unique subunit Swi4 and is important

for proper transition from mitotic to meiotic cell fate. Swi4 inhibition occurs through: (1)

meiosis-specific expression of a LUTI from the *SWI4* locus, *SWI4^LUTI*, which

downregulates Swi4 protein levels, and (2) repression of Swi4 activity by Whi5, which

occurs even when the LUTI-based regulation is disrupted. Conditions that lead to

elevation of Swi4 levels at meiotic entry result in abrupt activation of the SBF and a

concomitant decline in the expression of early meiotic genes, leading to meiotic

progression delays. Additionally, we show that the SBF targets *CLN1* and *CLN2* are major

drivers of the meiotic entry defect resulting from SBF misregulation. The *CLN*-dependent

phenotypes can be largely rescued by tethering of the central meiotic regulator Ime1 to

its cofactor Ume6, suggesting that the primary reason for meiotic failure caused by these

G1 cyclins is due to a defect in Ime1-Ume6 interaction. Our findings reveal the functional

role of a LUTI in establishing the meiotic transcriptional program, demonstrate how the

LUTI-based regulation is integrated into a larger regulatory network to ensure timely SBF

activity, and provide mechanistic insights into how SBF misregulation impedes transition

from mitotic to meiotic cell fate (Figure 6F).

Rewiring of the G1/S regulon

It has long been known that entry into the mitotic cell cycle and meiotic differentiation are

regulated differently. During the mitotic cell cycle, budding yeast cells must reach a critical

size before commitment to division in late G1 phase, an event termed the "Start" (Hartwell

et al., 1974; Johnston et al., 1977). Start depends on the activation of the G1/S regulon

by SBF and MBF, which regulate ~200 genes that function in polarized growth,

macromolecular biosynthesis, DNA replication, and repair among other critical processes

(reviewed in Jorgensen and Tyers, 2004). In contrast, entry into meiosis requires the

transcriptional activator, Ime1, which is exclusively induced in diploid cells through the

integration of several extrinsic cues including nitrogen levels, carbon source and

extracellular pH (reviewed in van Werven and Amon, 2011). Ime1 initiates the first

transcriptional wave of early meiotic genes required for DNA replication, recombination,

and chromosome morphogenesis. Before our study, it was unclear whether and how

SBF-MBF activity is regulated during the transition from mitosis to meiosis, despite the

transcriptome-wide differences between these two developmental programs. We found

that during meiotic entry, the SBF-specific Swi4 subunit is downregulated, which indicates

that the activity of SBF, but not MBF, is constrained. Consistently, SBF-specific targets

display either low or no expression during early meiosis, while MBF targets are

upregulated upon meiotic entry. MBF targets are involved in processes shared between

mitosis and meiosis including DNA replication and repair, whereas SBF targets are

predominantly involved in processes unique to mitosis such as budding and cell wall

biosynthesis. This functional specialization might have enabled adaptations to

accommodate specific cell fates. Likewise, SBF-specific targets that are detrimental to

meiosis, such as the G1 cyclins (Colomina et al., 1999, and this study), might contribute

to the mutually exclusive nature of the mitotic and meiotic programs.

Absolute measurements of the concentrations of SBF and MBF in single cells during

mitotic G1 demonstrated that these transcription factors are sub-saturating with respect

to their target promoters in small cells, but that SBF and MBF levels increase as cells

grow, suggesting that their abundance is a limiting factor in activating the G1/S regulon

(Dorsey et al., 2018). Corroborating these findings, we show that overexpression of *SWI4*

as well as the simultaneous disruption of the two pathways that normally restrict SBF

activity at meiotic entry – namely LUTI and Whi5 – result in abrupt activation of the SBF

regulon and a concomitant decrease in early meiotic gene expression. However, MBF

targets do not seem to be affected in response to increased SBF activity. This is surprising

given that SBF and MBF share a common subunit, Swi6, and that an upsurge in Swi4

abundance might be expected to titrate Swi6 away from MBF. It is possible that meiosis-

specific regulators like Ime1 and Ime2 compensate to maintain MBF target expression

(Brush et al., 2012). Future studies are necessary to dissect the meiotic roles of MBF and

potential compensatory mechanisms.

Inhibition of G1 cyclins during meiotic entry and vice versa

We found that the meiotic entry delay due to untimely SBF activation can be partially

rescued by loss of either *CLN1* or *CLN2*, demonstrating that both cyclins are responsible

for the meiotic defects associated with SBF misregulation. In addition to *CLN1* and *CLN2*,

improper activation of SBF also leads to upregulation of *PCL1* during meiotic entry (Figure

2F and 6B). *PCL1* encodes a cyclin that interacts with the Pho85 CDK and is involved in

the regulation of polarized cell growth and morphogenesis as well as progression through

the cell cycle (Espinoza et al., 2016). Whether *PCL1* misexpression contributes to the

SBF-associated meiotic entry defects is an interesting avenue for future studies.

We and others have demonstrated that elevated levels of G1 cyclins inhibit meiotic entry.

Therefore, it is crucial to dissect the mechanisms that govern the downregulation of G1

cyclins during this process. Among the G1 cyclins that are known to inhibit meiotic entry

(Colomina et al., 1999), only the mechanism of *CLN3* restriction was previously known

(Parviz and Heideman, 1998; Gallego et al., 1997). Our study provides a mechanistic

understanding of how *CLN1* and *CLN2* remain quiescent during meiotic entry through the

combined act of LUTI-based and Whi5-dependent inhibition of SBF. Whether additional

mechanisms play a role in restricting SBF and/or *CLN1-CLN2* remains to be determined.

Our findings also shed light on the mechanism by which G1 cyclins prevent meiotic entry.

Previous work demonstrated that G1 cyclin overexpression leads to downregulation of

IME1 expression and inhibition of Ime1 nuclear localization (Colomina et al., 1999). Even

though we observed similar defects in *IME1* in response to *CLN2* overexpression,

increasing *IME1* expression or targeting Ime1 to the nucleus did not result in successful

meiosis (Figure 4). Instead, we found that restoring the interaction between Ime1 and its

cofactor Ume6 was sufficient to bypass the meiotic blockage exerted by Cln/CDK.

Collectively, our analyses suggest that the primary reason why G1 cyclins cause a meiotic

failure is due to a deficiency in Ime1-Ume6 interaction. A previous study reported that G1

cyclins do not affect the interaction between Ime1 and Ume6 (Colomina et al., 1999).

However, this conclusion was based on the use of a truncated Ime1 protein that only

contained the Ume6 interaction domain. Therefore CLN-dependent regulations on full-

length Ime1 might be missed in this context.

Future work could be aimed at dissecting how G1 cyclins affect the interaction between

Ime1 and Ume6 and whether their impact on Ime1's subcellular localization is primarily

due to G1 cyclin-dependent changes in Ime1-Ume6 interaction. Since there are no CDK

consensus phosphorylation sites on Ime1 itself, other players are likely to be involved.

Rim11 and Rim15 are potential candidates since these two kinases have been implicated

in Ime1-Ume6 phosphorylation as well as regulation of Ime1 localization and its

interaction with Ume6 (Rubin-Bejerano et al., 1996; Vidan and Mitchell, 1997; Bowdish et

al., 1994). Interestingly, Rim15 contains CDK consensus phosphorylation sites (Holt et

al., 2009; Moreno-Torres et al., 2017; Breitkreutz et al., 2010), Furthermore, Cln2/CDK

activity has been previously shown to inhibit Rim15 nuclear localization (Talarek et al.,

2010), thereby making it an attractive candidate for further investigation.

Antagonistic relationship between two key transcription factors, Swi4 and Ime1

Swi4 and Ime1 regulate each other in an antagonistic manner (Figure 6F), as supported

by several observations. First, overexpression of *SWI4* leads to a reduction in Ime1

nuclear localization (Figure 2B). Second, premature activation of SBF through *SWI4*

overexpression or simultaneous disruption of $SWI4^{LUTI}$ and Whi5 pathways results in a

significant delay in the chromosomal localization of Rec8, a direct transcriptional target of

Ime1 (Figure 2H, 2I and 6E). Additionally, many early meiotic genes that are also

transcriptional targets of Ime1 are downregulated under the same conditions (Figure 2F

and 6D). Third, *IME1* is inhibited by the G1 cyclins, which are themselves targets of

Swi4/SBF (Figure 4). The reverse regulation, where Ime1 inhibits Swi4, is also in place.

This occurs through the Ime1-dependent expression of $SWI4^{LUTI}$, which is necessary to

downregulate Swi4 levels (Figure 5, also reported in Tresenrider et al., 2021). The

antagonistic relationship between Swi4 and Ime1 is further evidenced by their mutually

exclusive pattern of nuclear localization at the single-cell level (Figure 2E). In summary,

our study highlights the multiple ways in which Swi4 and Ime1 regulate each other, which

likely plays a crucial role in cellular decision making between the mitotic and meiotic

transcriptional programs.

Intersection of LUTI-based gene repression with other regulatory pathways

While LUTI-based regulation is both necessary and sufficient to downregulate Swi4

levels, disrupting it alone is not enough to activate the SBF regulon. We found that the

LUTI-based mechanism works together with the Whi5 pathway to inhibit SBF activity

(Figure 6). As a result, only when both mechanisms are simultaneously disrupted, do cells

exhibit abrupt activation of SBF targets and subsequent meiotic entry defects. This two-

pronged inhibition of SBF activity is reminiscent of how meiotic cells prevent microtubule-

kinetochore interactions during prophase I (Miller et al., 2012). Specifically, LUTI-based

regulation represses the expression of a limiting outer kinetochore subunit, Ndc80 (Chen

et al., 2017). Even when the outer kinetochore assembles upon disruption of $NDC80^{LUTI}$,

microtubules are still unable to engage with the kinetochores due to restriction of Clb/CDK

activity. The loss of both regulations results in meiotic chromosome segregation defects

(Chen et al., 2017). Integrating LUTI-based repression into larger regulatory networks

likely ensures robustness in cellular decision-making, providing a fail-safe system.

However, combinatorial use of the LUTI-based mechanism with other regulatory

pathways poses a challenge to uncover their functional importance. Despite 8% of the

yeast genes being subject to LUTI-based regulation during meiosis (Cheng et al., 2018),

the biological significance of only two LUTIs, *NDC80^LUTI* and *SWI4^LUTI*, have been

uncovered thus far. Therefore, systematic studies aiming to dissect the biological roles of

LUTIs in meiosis and beyond would benefit from simultaneous perturbation approaches,

including synthetic genetic interactions.

Concluding remarks

Cell state changes are primarily governed by transcription factors. Our study highlights

how transition from mitotic to meiotic cell fate is ensured by inhibition of the SBF

transcription factor through two distinct mechanisms. The SBF regulon includes G1

cyclins, whose untimely expression blocks meiotic entry by disrupting the interaction

between Ime1 and Ume6, a key transcriptional co-activator of meiosis. SBF and Ime1-

Ume6 antagonize each other, thereby helping establish a mutually exclusive state

between the mitotic and meiotic transcriptional programs. Mammalian cells have

functional homologs to both sets of transcription factors. First, similar to Ume6, MEIOSIN

binds to the promoters of early meiotic genes and recruits Ime1-like STRA8 to activate

gene expression (Anderson et al., 2008; Kojima et al., 2019; Baltus et al., 2006; Ishiguro

et al., 2020). Second, SBF and MBF counterparts in mammalian cells are known as E2Fs,

nine of which control the G1/S transition (reviewed in Van Den Heuvel and Dyson, 2008).

Each E2F regulates a distinct sets of genes (Gaubatz et al., 2000). Furthermore, some

members of the E2F family are involved in tissue-specific regulation of cell fate (Julian et

al., 2016). However, the molecular mechanisms driving the rewiring of these complex

gene networks are not well understood. Our investigation could help shed light on the

regulatory mechanisms governing the interplay between mitotic and meiotic transcription

factors outside of yeast.

MATERIALS AND METHODS

Strain Construction and Cloning

All strains used in this study were derived from the SK1 background. Detailed information

about the strain genotypes can be found in Supplemental Table 4.

For *SWI4* (pUB1587) 1200 bases upstream of the ORF, *SWI4* ORF, C-terminal 3V5

epitope tag, and *SWI4* 3'UTR (1000 bases downstream of ORF) were cloned into a *LEU2*

single integration vector by Gibson assembly (NEB) (Gibson et al., 2009). LUTI promoter

deletion (Δ*LUTI*) strain (pUB1588) was similarly constructed with -1200 to -934 bases

upstream from ORF deleted. For the uORF mutant (Δ*uORF*) strain (pUB1734) all seven

ATG uORFs were mutated using gBlocks with the ATG>ATC mutations and cloned by

Gibson assembly into 3V5-tagged *LEU2* single integration vector. In all strains described

above, the endogenous *SWI4* gene is deleted using Pringle-based insertion of KanMX6

marker (Bähler et al., 1998).

For overexpression of *SWI4 (pUB1585)*, *CLN1 (pUB2144)*, and *CLN2 (pUB1899)*, the

relevant gene carrying a 3V5 epitope tag was cloned downstream of the *ATG8* promoter,

which is highly expressed during meiosis (Brar et al., 2012). A *LEU2* single integration

vector was used for cloning the fragments by Gibson assembly (*NEB*). In the strains

carrying overexpression transgenes, the wild-type alleles at the endogenous loci

remained intact. *pCUP1-GFP-IME1* allele was made with Pringle-based insertion of the

pCUP1 promoter upstream of *GFP-IME1*. *PUS1*-αGFP and *UME6*-αGFP were made by

Pringle-based insertion using a plasmid (pUB1707) gifted by the Lackner Lab. For *WHI5-*

AA the *WHI5-mCherry-FRB* allele was made by Pringle-based insertion using a plasmid

(pUB595) gifted by the Nasmyth lab. All plasmid sequences were confirmed by

sequencing.

All single integration plasmids were digested with Pme1 before transformation. Proper

integration was confirmed by PCR. The plasmids constructed in this study are listed in

Supplemental Table 5.

Sporulation Conditions

For meiotic experiments using *pCUP1-IME1/pCUP1-IME4* or *pCUP-IME1* system, cells

were synchronized as described in (Chia and van Werven, 2016). Briefly, after 24 h of

growth in YPD at room temperature, saturated cultures (OD_{600} > 10) were diluted to an

OD_{600} of 0.25 in BYTA (1% yeast extract, 2% bacto tryptone, 1% potassium acetate, and

1.02% potassium phthalate) for 16-18 h of growth at 30°C (OD_{600} of > 5). Cells were

washed with water twice before final resuspension in SPO with 0.5% potassium acetate

to an OD_{600} of 1.85. After 2 h in SPO, *IME1* and *IME4* were induced with 50 μM $CuSO_4$.

Sporulation efficiency was measured after 24 h in SPO. Anchor away meiotic experiments

was performed as described above with final 1 μM rapamycin (Millipore) added to BYTA

30 minutes before transfer to SPO and again 1 μM rapamycin added to SPO media.

In all other meiotic experiments, cells were prepared as in (Carlile and Amon, 2008).

Briefly, after 24 h of growth in YPD at room temperature, saturated cultures (OD_{600} > 10)

were diluted to an OD_{600} of 0.25 and inoculated in BYTA (1% yeast extract, 2% bacto

tryptone, 1% potassium acetate, and 1.02% potassium phthalate) for 16-18 h at 30°C

(OD_{600} of > 5). Cells were washed with water twice before final resuspension in SPO with

0.5% potassium acetate to an OD_{600} of 1.85. Sporulation efficiency was counted after 24

h in SPO.

UME6-αGFP meiotic experiments were prepared as in (Chia and van Werven, 2016).

Briefly, after 18 h of growth in YPD at room temperature, saturated cultures (OD_{600} > 10)

were diluted to an OD_{600} of 0.2 in reduced YPD (1% yeast extract, 2% peptone, 2%

dextrose, uracil [24 mg/L]). Reduced YPD was used instead of BYTA to prevent cells from

prematurely entering meiosis in BYTA due to the Ime1-Ume6 interaction from the GFP

nanobody. Cells were grown for ~6 h at 30°C until they reached an OD_{600} between 0.5

and 1.0. Cultures were then back diluted to OD_{600} of 0.1 and grown for 18 h at 30°C. Cells

were washed with water twice before final resuspension in SPO with 0.5% potassium

acetate to an OD_{600} of 1.85. Sporulation efficiency was counted after 24 h in SPO.

RNA Extraction for mRNA-seq, RT-qPCR, and RNA Blotting

RNA extraction was performed as described in (Tresenrider et al., 2021). Briefly, ~4

OD unit of cells were pelleted by centrifugation for 1 minute at 20,000 rcf and snap frozen

in liquid nitrogen. Cells were thawed on ice and resuspended in TES buffer (10 mM Tris

pH 7.5, 10 mM EDTA, 0.5% SDS). An equal volume of Acid Phenol:Chloroform:Isoamyl

alcohol (125:24:1; pH 4.7) was added to cells and incubated at 65°C for 45 min shaking

at 1400 rpm. Aqueous phase was transferred to a new tube with chloroform, vortexed for

30 sec, separated by centrifugation, and precipitated in isopropanol and sodium acetate

overnight at -20°C. Pellets were washed with 80% ethanol and resuspended in DEPC

water for 10 min at 37°C. Total RNA was quantified using a nanodrop.

mRNA Sequencing (mRNA-seq) and Analysis

RNA-seq libraries were generated with the NEXTflex™ Rapid Directional mRNA-Seq Kit

(NOVA-5138, Perkin Elmer). 10 µg of total RNA was used as input for all

libraries. AMPure XP beads (A63881, Beckman Coulter) were used to select fragments

between 200-500 bp. Libraries were quantified using the Agilent

4200 TapeStation (Agilent Technologies, Inc). Samples were submitted for 100 bp SE

sequencing by the Vincent J. Coates Genomics Sequencing Laboratory with a NovaSeq

SP 100SR.

Hisat2 (Kim et al., 2019) was used to align reads to map sequences to SK1 PacBio

genome. Quantification of RNA as transcripts per million was done StringTie (Pertea et

al., 2015). Fold-change quantification was performed by DESeq2 using default options

(version 1.34.0, (Love et al., 2014)).

For Gene Set Enrichment Analysis (GSEA) v4.3.2 was used to compare TPM values for

different gene sets. The early meiotic gene set was created from Figure 2 of (Brar et al.,

2012). The SBF regulon was from Figure 3 of (Iyer et al., 2001). GSEA was performed

on the desktop app with default settings expect "Collapse/Remap to gene symbols" was

set to "No_Collapse" and "Permutation type" was set to "gene_set".

For Gene Ontology (GO) Analysis, SGD Gene Ontology Slim Term Mapper was used for

GO analysis using Yeast Go-Slim: process GO Set. Used output from DE-Seq2 analysis

with a cutoff of padj (P value) < 0.05.

The two overlapping targets (*SWE1* and *TOS4*) between the early meiotic gene set and

SBF targets are not plotted on volcano plot or used for GSEA analysis.

Reverse Transcription-Quantitative Polymerase Chain Reaction (RT-qPCR)

5 μg of isolated total RNA was treated with DNase (TURBO DNA-free[TM] Kit). cDNA was

reverse transcribed following the Superscript III kit (ThermoFisher Scientific).

Quantification was performed with Absolute Blue qPCR Mix (ThermoFisher Scientific).

Meiotic samples were normalized to *PFY1,* and mitotic samples were normalized to

ACT1. Oligonucleotides are listed in Supplemental Table 6.

Protein Extraction and Immunoblotting

~4 OD_{600} of cells were collected and pelleted by centrifugation for 1 min at 20,000 rcf.

Pellet was resuspended in 5% (w/v) TCA for at least 15 min at 4°C. Cells were washed

with TE50 (50 mM Tris-HCl [pH 7.5], 1 mM EDTA) and then with 100% acetone. The cell

pellet was dried overnight and then lysed with glass beads in lysis buffer (Tris-HCl [pH

7.5], 1 mM EDTA, 2.75 mM DTT, protease inhibitor cocktail (cOmplete EDTA-free

[Roche]). Next 3x SDS sample buffer (187.5 mM Tris [pH 6.8], 6% β-mercaptoethanol,

30% glycerol, 9% SDS, 0.05% bromophenol blue) was added and the cell lysate was

boiled for 5 min at 95°C. Protein was separated by PAGE using 4-12% Bis-Tris Bolt gels

(Thermo Fisher) and transferred onto 0.45 µm nitrocellulose membranes.

Cln1-3V5, Cln2-3V5 and GFP-GFP tagged proteins were all transferred onto 0.45 µm

nitrocellulose membranes using a semi-dry transfer apparatus (Trans-Blot Turbo System

(Bio-rad)). Swi4-3V5 or untagged Swi4, Swi6, and Mbp1 proteins were all transferred onto

0.45 µm nitrocellulose membranes using a PROTEAN Tetra tank (BioRAD) filled with 25

mM Tris, 192 mM glycine, and 7.5% methanol. All blots were incubated at room

temperature with Odyssey Blocking Buffer (PBS) (LI-COR Biosciences).

Immunoblotting for Cln1-3V5, Cln2-3V5, and GFP-Ime1 was performed as previously

described in (Tresenrider et al., 2021). Briefly, mouse α-V5 antibody (R960-25, Thermo

Fisher) or mouse α-GFP antibody (632381, Takara) were diluted 1:2000 in Odyssey

Blocking Buffer (PBS) (LI-COR Biosciences) with 0.01% Tween. Rabbit α-hexokinase (α-

Hxk2) antibody (H2035, US Biological) was diluted to 1:20,000. Secondary antibodies

used were α-mouse antibody conjugated to IRDye 800CW (926-32212, LI-COR

Biosciences) and α-rabbit antibody conjugated to IRDye 680RD (926-68071, LI-COR

Biosciences). Secondary antibodies were diluted to 1:20,000 in Odyssey Blocking Buffer

(PBS) with 0.01% Tween.

For immunoblotting of Swi4, Swi6, and Mbp1. antibodies specific to each subunit were a

generous gift from the Andrews and deBruin labs (Andrews and Herskowitz, 1989; Harris

et al., 2013). α-Swi4, α-Swi6, α-Mbp1 antibodies were each diluted to 1:2000 in Odyssey

Blocking Buffer (PBS) (LI-COR Biosciences) with 0.01% Tween. Secondary antibodies

included a α-rabbit antibody conjugated to IRDye 800CW (926-32213, LI-COR

Biosciences) and a α-rabbit antibody conjugated to IRDye 680RD (926-68071, LI-COR

Biosciences).

Odyssey system (LI-COR Biosciences) was used to image the blots, and Image Studio

Lite (LI-COR Biosciences) was used for image quantification.

Northern (RNA) Blotting

For each blot, 10 µg of total RNA was dried in a Savant Speed Vac (SPD111V). RNA was

then resuspended and denatured in glyoxal/DMSO mix (1M deionized glyoxal, 50% v/v

DMSO, 10 mM sodium phosphate (NaPi) buffer [pH 6.8]) at 70°C for 10 min. RNA sample

was loaded into an agarose gel (1.1% [w/v] agarose in 0.01 M NaPi buffer) with loading

dye (10% v/v glycerol, 2 mM NaPi buffer [pH 6.8], ~0.25% w/v xylene cyanol, and orange

G) and ran for 3 h at 100 V with a Variable Speed Pump (BioRad) to circulate buffer during

the entire gel run.

RNA was transferred overnight to nylon membrane (Hybond-N+ [GE]) in SSC. Membrane

was crosslinked using a Stratalinker UV Crosslinker (Stratagene). Ribosomal RNA

(rRNA) bands were visualized with methylene blue staining and imaged on a Gel Doc

XR+ Molecular Imager with Image Lab software (BioRad).

Probe templates containing the T7 promoter were amplified using PCR. PCR product was

concentrated with MinElute Spin Columns (Qiagen) and then used for *in vitro* transcription

to generate a strand-specific RNA probe using a MaxiScript T7 Kit (Invitrogen) according

to the manufacturer's instructions, except cold UTP was replaced with α-P32 labeled UTP

(PerkinElmer). Excess nucleotides were removed with NucAway Spin Columns

(Invitrogen). Blots were blocked in ULTRAhyb Ultrasensitive Hybridization Buffer

(Invitrogen) for 1 h and then incubated with the α-P32 labeled probe overnight at 68°C.

Blots were then washed twice with low stringency wash buffer (2X SSC, 0.1% SDS) for

10 min and then washed twice with high stringency wash buffer (0.1X SSC, 0.1% SDS)

for 15 min. Blots were then exposed overnight on a storage phosphor screen (Molecular

Dynamics) and then imaged on a Typhoon phosphor-imaging system.

Fluorescence Microscopy

For time course imaging of cells expressing *GFP-IME1*, *SWI4-mCherry*, or *WHI5-*

mCherry, 500 µL of meiotic culture was fixed with a final concentration of 3.7%

formaldehyde (v/v) at room temperature for 15 min. Cells were then washed in 1 ml of

100 mM potassium phosphate [pH 6.4] and stored at 4°C in 20 µl of KPi Sorbitol solution

overnight (100 mM potassium phosphate [pH 7.5], 1.2M sorbitol). Cells were mounted on

a slide and imaged using DeltaVision Elite wide-field fluorescence microscope (GE

Healthcare) with a 60x/1.516 oil immersion objective. Deconvolution of images was done

with softWoRx imaging software (GE Life Sciences).

For live-cell imaging, cells at OD_{600} of 1.85 in conditioned SPO (filter-sterilized SPO

culture after ~5 h in 30°C) were sonicated and transferred to a concanavalin A (Sigma)

treated 96-well clear, flat bottom plate (Corning). Four z positions (2 µm step size) were

acquired per XY position. Acquisition was performed in a temperature-controlled chamber

at 30°C. Please refer to Supplemental Table 7 for acquisition settings.

Image Quantification

All image analysis was performed with FIJI (Schindelin et al., 2012). Maximum z-

projection are shown in figures and were modified using linear brightness and contrast

adjustments in FIJI.

To quantify localization of GFP-Ime1 and Rec8-GFP, z-slices containing the nucleus were

selected using the Htb1-mCherry signal. Max projection was created from these slices

and GFP-Ime1 was scored double-blinded as nuclear or not nuclear.

For mean nuclear intensity of GFP-Ime1 and Swi4-mCherry, an individual z-slice

containing the nucleus was selected and nuclear mask was generated using the Htb1-

mCherry signal. The nuclear mask was then used to quantify the mean nuclear intensity

of GFP-Ime1 signal.

Single molecule RNA FISH

smFISH was performed and quantified as previously described in (Chen et al., 2018). All

probes were ordered from (Biosearch Technologies). Unique region of $SWI4^{LUTI}$ was

visualized by twenty-eight 20-mer oligonucleotide probes coupled to CAL fluor Red 590.

Thirty-eight 20-mer probes coupled to Quasar 670 dye targeted to $SWI4^{canon}$. ~ 4 OD unit

of cells were fixed in final 3% formaldehyde (v/v) and incubated at room temperature for

20 minutes. Fixed samples were moved to 4°C to continue fixing overnight. Cells were

washed three times in cold Buffer B (0.1 M potassium phosphate [pH 7.5], 1.2 M sorbitol)

and resuspended in digestion buffer (Buffer B, 200 mM Vanadyl ribonucleoside complex

[VRC from NEB], zymolyse [zymolase 100T, MP Biomedicals]). Cells were digested at

30°C for 20 min and then gently washed with 1 mL of cold Buffer B and resuspended in

1 mL of 70% ethanol for 3.5-5 h.

Cells were then incubated in 1mL of 10% formamide wash buffer (10% formamide, 2X

SSC) at room temperature for 15 minutes. For hybridization, each probe set was added

(final concentration of 500 nM) to 20mM VRC and hybridization buffer (1% Dextran sulfate

[EMD Millipore], 1 mg/mL E. coli tRNA [Sigma], 2 mM VRC, 0.2 mg/mL BSA, 1X SSC,

10% formamide in nuclease-free water). Hybridization was done overnight at 30°C.

Samples were incubated in the dark for 30 min at 30°C in 1 mL of 10% formamide wash

buffer. Buffer was then removed, and cells were stained with DAPI and resuspended in

glucose-oxygen-scavenging buffer or GLOX buffer (10 mM Tris [pH 8.0], 2x SSC, 0.4%

glucose) without enzymes. Before imaging, GLOX solution with enzyme (1% v/v catalase,

1% v/v glucose oxidase (Sigma), 2 mM Trolox (Sigma)) was added to sample.

Images were acquired with the DeltaVision microscope as described in previous section

with filters: TRITC (EX542/27, EM597/45) for CAL Fluor Red 590 and CY5 (EX632/22,

EM679/34) for Quasar 670. Series of z-stacks (15–25 slices) were acquired with a step

size of 0.2 μm.

Matlab script (Chen et al., 2017, 2018) was run to quantify FISH spots with max intensity

projection of z-stacks. The same "signal" and "SNR" thresholds were applied to all the

images within a replicate.

Resource availability

All materials used in this study are available upon request from the corresponding author.

Sequencing data generated in this study are available at NCBI GEO under the accession

ID: GSE225963

The custom code used for the analysis is available in the following code repository:

https://github.com/elifesciences-publications/Chen_Tresenrider_et_al_2017

Acknowledgements:

We thank Gloria Brar, Nick Ingolia, Rebecca Heald, Kathlyeen Ryan, Jingxun Chen,

Anthony Harris, Andrea Higdon, Grant King, Jessica Leslie, Kate Morse, Emily Powers,

Cyrus Ruediger, Tina Sing, and Ben Styler for suggestions and comments on this

manuscript, David McSwiggen and Yuichiro Iwamoto for technical support, all members

of the Ünal and Brar labs and Jeremy Thorner for valuable discussions. This work is

supported by funds from the National Institutes of Health (R01 GM140005) and Astera

Institute to E.Ü. and a National Institutes of Health Traineeship (T32 GM007232) to A.J.S.

Author contributions:

A.J.S, conceptualization, data curation, formal analysis, investigation, methodology,

validation, visualization, and drafting of the manuscript. S.C.Y, data curation, formal

analysis, investigation, methodology, validation, visualization. E.Ü., conceptualization,

methodology, visualization, supervision, project administration, funding acquisition, and

drafting of the manuscript.

Figure Legends

Figure 1. Swi4 subunit of the SBF and SBF targets are downregulated during early
meiosis

A. A schematic of SBF and MBF complexes and the general functional groups of the

genes they regulate. **B.** Samples from strain UB35246 were collected between 0-6 hours

(h) in sporulation medium (SPO) and immunoblots were performed using α-Swi4, α-Swi6,

and α-Mbp1 respectively. Hxk2 was used a loading control. Representative blots from

one of two biological replicates are shown. **C.** Quantification of the immunoblots in (B).

The signal at each time point was first normalized to Hxk2 loading control and then to the

max signal. **D.** Scatterplot of RNA-seq data (RPKM) from (Brar et al., 2012) comparing 2

h in SPO vs. mitotic growth of well characterized SBF targets (pink) and MBF targets

(teal). **E.** Wild type (UB22199) and *pATG8-SWI4* (UB22226) cells were collected to

perform RT-qPCR for *CLN1, CLN2, CDC21,* and *RNR1* transcripts. Transcript abundance

was quantified using primer sets specific for each respective gene from three technical

replicates for each biological replicate. Quantification was performed in reference to *PFY1*

and then normalized to wild-type control. FC = fold change. Experiments were performed

twice using biological replicates, mean value plotted with range. Differences in wild type

versus *pATG8-SWI4* transcript levels at 2 h in SPO compared with a two-tailed t-test (*,

p = 0.0351 [*CLN1*]; **, p = 0.0013 [*CLN2*]; ns, p = 0.8488 [*CDC21*]; ns, p = 0.0859

[*RNR1*]). **F.** Live-cell imaging of strains containing the fluorescently tagged histone Htb1-

mCherry for wild type (UB32085) and *pATG8-SWI4* (UB32089). Experiments were

performed twice using biological replicates, mean value plotted with range. Differences in

meiotic progression tested by Mann Whitney test, two-tailed (**, p = 0.0045).

Figure 2. Regulation of Swi4 abundance is required for timely meiotic entry

A-B. Fixed imaging of cells marked with GFP-Ime1 and Htb1-mCherry. Wild type

(UB22199) and *pATG8-SWI4* (UB22226) cells were collected between 0-4 h in SPO. **A.**

Representative images with merge at 0 h and 2 h in SPO. Representative cells outlined.

Scale bar: 3 µm. **B.** Quantification as percent cells with nuclear Ime1. Experiments were

performed twice using biological replicates, mean value plotted with range. Total of 200

cells analyzed per strain. Differences in the fraction of cells with nuclear Ime1 was

compared using a two- tailed t-test (**, p = 0.0099 [1 h in SPO]; *, p = 0.0169 [2 h in SPO];

*, p = 0.0315 [3 h in SPO]; ns, two-tailed p = 0.4595 [4 h in SPO]). **C-E.** Fixed imaging of

cells marked with GFP-Ime1 and Swi4-mCherry. Wild type (UB31378) and *pATG8-SWI4*

(UB31381) cells were collected at 0 h and 4 h in SPO. **C.** Example images with merge at

4 h in SPO. Example cells outlined (*low nuclear GFP-Ime1 with high nuclear Swi4-

mCherry, **high nuclear GFP-Ime1 with low nuclear Swi4-mCherry). Scale bar: 3 µm. **D.**

Scatterplot of GFP-Ime1 mean nuclear intensity and Swi4-mCherry mean nuclear

intensity for wild type and *pATG8-SWI4* cells at 0 h in SPO. See Materials and Methods

for further details about image quantification. Dashed line is linear regression plotted for

each condition and strain. A total number of 269 cells were analyzed. **E.** Same as in (D)

but for wild type and. *pATG8-SWI4* cells at 4 h in SPO. A total number of 341 cells were

analyzed. Differences in mean nuclear GFP-Ime1 or Swi4-mCherry intensity between wild

type and *pATG8-SWI4* compared using a Mann-Whitney test, two-tailed (****, p < 0.0001

[wild type vs. *pATG8-SWI4* (GFP-Ime1)]; ****, p < 0.0001 [wild type vs. *pATG8-SWI4*

(Swi4-mCherry)]). **F.** Volcano plot of DE-Seq2 analysis for *pATG8-SWI4* versus wild type.

Dashed line indicates padj (p value) = 0.05. Analysis was performed using mRNA-seq

from two biological replicates. Wild type (UB22199) and *pATG8-SWI4* (UB22226) cells

were collected at 2 h in SPO. SBF targets (pink) (Iyer et al., 2001) and early meiotic genes

(blue) defined by (Brar et al., 2012). Darker pink or darker blue, labeled dots are well

studied targets in either gene set list. **G.** GSEA analysis of mRNA-seq comparing wild

type vs. *pATG8-SWI4* collected at 2 h in SPO. Vertical black bars represent the early

meiotic cluster from (Brar et al., 2012) or SBF cluster from (Iyer et al., 2001). The heatmap

indicates genes that are more enriched in *pATG8-SWI4* (red, left-side) or genes that are

enriched in wild type (blue, right-side). NES = normalized enrichment score. Enrichment

was determined by comparing *pATG8-SWI4* versus wild type. **H-I.** Live-cell imaging of

cells in meiosis marked by Rec8-GFP and nuclear marker Htb1-mCherry for wild type

(UB32085) and *pATG8-SWI4* (UB32089). **H.** Movie montage with example images

throughout meiosis for Rec8-GFP and Htb1-mCherry. Scale bar: 3 μm **I.** Quantification

as percent of cells that entered meiosis assayed by nuclear Rec8 appearance.

Experiments were performed using two biological replicates, mean value plotted with

range. A total number of 452 cells were analyzed. Differences in meiotic progression

compared by Mann Whitney test, two-tailed (***, p = 0.0005 [wild type vs. *pATG8-SWI4*]).

Figure 3. Removal of the SBF targets Cln1 or Cln2 partially rescues the meiotic

entry delay in the *pATG8-SWI4* mutant

A. Immunoblotting was performed on samples collected for wild type (UB29326) and

pATG8-SWI4 (UB29328) between 0-6 h in SPO using α-V5 antibody to track Cln1-3V5.

Hxk2 was used a loading control. Representative blots from one of two biological

replicates are shown. **B.** Quantification of (A). **C.** Same as in (A) but for wild type

(UB29330) and *pATG8-SWI4* (UB29332) cells using α-V5 antibody to track Cln2-3V5.

Hxk2 was used a loading control. Representative blots from one of two biological

replicates are shown. **D.** Quantification of (C). **E.** Live-cell imaging of meiotic cells marked

by Rec8-GFP and nuclear marker Htb1-mCherry, with the following genotypes: wild type

(UB32085), *pATG8-SWI4* (UB32089), and *pATG8-SWI4*; *cln1Δ* (UB34536).

Quantification of cells that entered meiosis assayed by the initial timing of nuclear Rec8

appearance. Experiments were performed using two biological replicates, mean value

plotted with range. A total number of 883 cells were analyzed. Differences in meiotic

progression compared by Mann-Whitney test, two-tailed (*, p = 0.0111 [*pATG8-SWI4* vs.

pATG8-SWI4; *cln1Δ*]). *cln1Δ* alone (not shown) has similar meiotic progression kinetics

relative to wild type. **F.** Same as (E) but with the following genotypes: wild type

(UB32085), *pATG8-SWI4* (UB32089), and *pATG8-SWI4*; *cln2Δ* (UB34165). A total

number of 610 cells were analyzed. Differences in meiotic progression compared by

Mann-Whitney test, two-tailed (*, p = 0.0478 [*pATG8-SWI4* vs. *pATG8-SWI4*; *cln2Δ*]).

cln2Δ alone (not shown) has similar meiotic progression kinetics relative to wild type.

Figure 4. Tethering of Ime1 to Ume6 is sufficient to overcome the meiotic block

exerted by G1 cyclin overexpression

A. Sporulation efficiency of cells at 24 h in SPO media wild type (UB22199), *pATG8-*

CLN1 (UB32820), *pATG8-CLN2* (UB25959), *pCUP-GFP-IME1* (UB34641), *pCUP1-GFP-*

IME1; pATG8-CLN2 (UB35057), *PUS1-αGFP* (UB35593), *PUS1-αGFP; pATG8-CLN2*

(UB35982), *UME6-αGFP* (UB35300), and *UME6-αGFP; pATG8-CLN2* (UB35177).

Experiments shown in this figure were performed using two biological replicates, mean

value plotted with range. Total of 200 cells counted per strain. See Supplementary Table

2 for statistics. **B-C.** Fixed imaging of cells marked with GFP-Ime1 and Htb1-mCherry.

Wild type (UB22199) and *pATG8-CLN2* (UB25959) cells were collected between 0-3 h in

SPO. **B.** Representative images with merge at 2 h in SPO. Representative cells outlined.

Scale bar: 3 μm. **C.** Quantification of cells with nuclear Ime1. Experiments were performed

using two biological replicates, mean value plotted with range. Total of 200 cells analyzed

per strain. Differences in percent of cells with nuclear Ime1 was compared by two-tailed

t-test (**, p = 0.00917 [1 h in SPO]; **, p = 0.0044 [2 h in SPO]; *, p = 0.0122 [3 h in SPO]).

D. Schematic depicting use of *pCUP1* promoter (*pCUP1-GFP-IME1*) to rescue Ime1

transcript and protein levels. **E-F** Fixed imaging of cells marked with GFP-Ime1 and Htb1-

mCherry. Cells with the following genotypes were collected at 2 h in SPO: wild type

(UB22199), *pATG8-CLN2* (UB35106), *pCUP1-GFP-IME1* (UB34641), and *pCUP1-GFP-*

IME1; pATG8-CLN2 (UB35057). **E.** Representative images with merge and

representative cells outlined. Scale bar: 3 µm. **F.** GFP-Ime1 mean nuclear intensity

measured for a single z-slice. A total number of 433 cells were analyzed. Differences in

mean nuclear intensity compared by Mann-Whitney test, two tailed, (****, p < 0.0001

[*pCUP1-IME1* vs. *pCUP1-IME1; pATG8-CLN2*]). **G.** Schematic of nanobody trap strategy

with *PUS1-αGFP* and GFP-Ime1 to rescue Ime1 nuclear localization in meiosis. **H-I.** Fixed

imaging of cells marked with GFP-Ime1 and Htb1-mCherry. Cells with the following

genotypes were collected at 2 h in SPO: wild type (UB22199), *pATG8-CLN2* (UB35106),

PUS1-αGFP (UB35593), and *PUS1-αGFP; pATG8-CLN2* (UB35982). **H.** Representative

images with merge and example cells outlined. Scale bar: 3 µm. **I.** GFP-Ime1 mean

nuclear intensity measured for a single z-slice. A total number of 934 cells were analyzed.

Differences in mean nuclear intensity compared by Mann-Whitney test, two-tailed, (****,

p < 0.0001 [wild type vs. *pATG8-CLN2*]; not significant (ns), p = 0.6563 [*PUS1-αGFP* vs.

pATG8-CLN2; PUS1-αGFP]; not significant (ns), p = 0.8881 [wildtype vs. *pATG8-CLN2;*

PUS1-αGFP]). **J.** Schematic of nanobody trap strategy with *UME6-αGFP* and GFP-Ime1

to rescue Ime1-Ume6 interaction in meiosis. **K-L** Fixed imaging of cells marked with GFP-

Ime1 and Htb1-mCherry. Cells with the following genotypes were collected at 2 h in SPO:

wild type (UB22199), *pATG8-SWI4* (UB35106), *UME6-αGFP* (UB35300), and *UME6-*

αGFP; pATG8-CLN2 (UB35177). **K.** Representative images with merge and

representative cells outlined. Scale bar: 3 μm. **L.** GFP-Ime1 mean nuclear intensity

measured for a single z-slice. A total number of 1220 cells were analyzed. Differences in

mean nuclear intensity compared by Mann-Whitney test, two-tailed (****, p < 0.0001 [wild

type vs. *pATG8-CLN2]*; ****, p < 0.0001 [*UME6-αGFP* vs. *pATG8-CLN2; UME6-αGFP*];

*, p = 0.0354 [wildtype vs. *pATG8-CLN2; UME6-αGFP*]).

1012

Figure 5. Ime1-dependent expression of a LUTI from the *SWI4* locus leads to a

reduction in Swi4 protein levels during meiotic entry

A. Genome browser views of RNA-seq data (Brar et al., 2012) of the *SWI4* locus. *SWI4^LUTI^*

transcription start site is ~1045bp upstream of *SWI4* ORF translation start site. **B.** A

schematic of LUTI-based gene regulation. Top: Mitotic growth, *SWI4^LUTI^* is repressed due

to Ume6-Rpd3-Sin3 complex and *SWI4^canon^* is induced by one or more transcription

factors including Ace2, Mbp1 and Swi5, leading to Swi4 protein production. Bottom:

Meiosis-specific expression of *SWI4^LUTI^* by Ime1-Ume6 leads to downregulation of Swi4

protein production due to combined effect of transcriptional and translation interference.

*SWI4^LUTI^* 5′ leader contains 7 AUG uORFs but only one is shown in the model for

simplicity. Schematic is adapted from (Tresenrider et al., 2021). **C.** Representative

smFISH images collected from premeiotic and meiotic cells for detecting *SWI4^canon^* and

*SWI4^LUTI^*. Cells with *pCUP1-IME1/pCUP1-IME4* meiotic synchronization system were

induced to enter meiosis with 50 μM $CuSO_4$ after 2 h in SPO. Premeiotic cells were

collected before *IME1/4* induction and meiotic cells were collected 2 h post *IME1/4*

induction from strain UB14273. Q 670 probes (green) hybridize to shared region within

SWI4 CDS. CF590 probes hybridize to the unique 5′ leader region of *SWI4^LUTI^* (depicted

on the schematic shown above the images). DNA was stained with DAPI. Scale bar: 3

µm. **D.** Quantification of smFISH shown in (C), plotted as relative frequency histograms

of cells with $SWI4^{canon}$ and $SWI4^{LUTI}$ transcripts per cell. Data pooled from two

independent biological replicates. Dashed line indicates median number of transcripts per

cell. Each histogram is normalized with maximum bin height being the same across all

histograms. A total number of 44 cells counted for premeiotic and 102 cells counted in

meiotic prophase. Differences in premeiotic versus meiotic were compared by Mann-

Whitney test, two-tailed (***, p = 0.0007 [$SWI4^{canon}$]; ****, p < 0.0001 [$SWI4^{LUTI}$]). **E.** RNA

blot performed on cells collected between 0-6 h in SPO. All strains carry a *SWI4-3V5*

tagged allele. Probe was specific for 3V5. Methylene blue staining of rRNA bands was

used as a loading control. Matched immunoblotting was performed against Swi4-3V5

using α-V5 and normalized to Hxk2 loading control for each sample. Cells collected are

wild type (UB22199) and Δ*LUTI* (UB23012). Representative blots from one of two

biological replicates are shown. **F.** Quantification of immunoblot in (E). **G.** Performed as

described in (E) for wild type (UB21386) and Δ*uORF* (UB23636) strains. Representative

blots from one of two biological replicates are shown. **H.** Quantification of immunoblot in

(G). **I.** Immunoblot using α-V5 performed on cells collected between 0-7 h in SPO from a

strain carrying *pCUP1-IME1* and *SWI4-3V5* alleles (UB34641). Swi4-3V5 abundance was

normalized to Hxk2 loading control. Cells were induced to enter meiosis with 50 µM

$CuSO_4$ after 2 h preincubation in SPO. Representative blots from one of two biological

replicates are shown. **J.** Quantification of immunoblot in (I). **K.** Genome browser view of

Ume6-ChIP at the *SWI4* locus (adapted from Tresenrider et al., 2021).

1052

Figure 6. *SWI4^LUTI^* **is integrated into a larger regulatory network to regulate SBF activity during meiotic entry**

A. Volcano plot of DESeq2 analysis for Δ*LUTI* versus wild type. Dashed line indicates padj (p-value) = 0.05. Analysis was performed with mRNA-seq in duplicate. Wild type (UB27083) and Δ*LUTI* (UB26874) collected at 2 h in SPO. SBF targets (pink) and early meiotic genes (blue) defined by (Iyer et al., 2001 and; Brar et al., 2012). Darker pink or darker blue, labeled dots are well studied targets in either gene set list. **B.** Top: Schematic of the anchor-away system using *WHI5-mCherry-FRB* (*WHI5-AA*) and *RPL13a-FKBP12* alleles. Bottom: Fixed imaging of cells marked with *WHI5-mCherry-FRB* (*WHI5-AA*) with DNA stained with DAPI. 1 µM rapamycin added at 0 h in SPO to induce nuclear exclusion of Whi5 (UB25431) strain collected at 0.5 h in SPO. Scale bar: 3 µm. Cells are rapamycin resistant due to mutated *TOR1* (*tor1-1*) and frp1Δ (yeast FKBP12 homolog) to reduce competition between for binding of Frb and Fkbp12. **C.** Same as in (A) but for wild type (UB27083) and *WHI5-AA* (UB25431) collected at 2 h in SPO. **D.** Same as in (A) but for wild type (UB27083) and Δ*LUTI; WHI5-AA* (UB25428) collected at 2 h in SPO. **E.** Live-cell imaging of cells in meiosis marked by Rec8-GFP and nuclear marker Htb1-mCherry. The following genotypes were imaged: wild type (UB35987), Δ*LUTI* (UB35989), *WHI5-AA* (UB35991), Δ*LUTI; WHI5-AA* (UB35989)**.** Quantification as percent of cells that entered meiosis assayed by nuclear Rec8 appearance. Experiments were performed using two biological replicates, mean value plotted with range. Differences in meiotic progression compared by Mann-Whitney test, two-tailed (*, p = 0.0112 [wild type vs. Δ*LUTI*; *WHI5-AA*]). **F.** Model of SBF regulation during meiotic entry. Ime1 downregulates Swi4 protein expression via induction of *SWI4^LUTI^* while Whi5 represses SBF activity in

1076 parallel to LUTI-based mechanism to prevent expression of SBF targets, including G1

1077 cyclins, which perturb meiotic entry via blocking interaction between Ime1 and its cofactor

1078 Ume6.

1079

1080
1081
1082
1083 **REFERENCES**

1084

Abeliovich, H., and D.J. Klionsky. 2001. Autophagy in Yeast: Mechanistic Insights and

Physiological Function. *Microbiol. Mol. Biol. Rev.* 65:463–479.

doi:10.1128/mmbr.65.3.463-479.2001.

Anderson, E.L., A.E. Baltus, H.L. Roepers-Gajadien, T.J. Hassold, D.G. De Rooij,

A.M.M. Van Pelt, and D.C. Page. 2008. Stra8 and its inducer, retinoic acid, regulate

meiotic initiation in both spermatogenesis and oogenesis in mice. *Proc. Natl. Acad.*

Sci. U. S. A. 105:14976–14980. doi:10.1073/pnas.0807297105.

Andrews, B.J., and I. Herskowitz. 1989. The yeast SWI4 protein contains a motif

present in developmental regulators and is part of a complex involved in cell-cycle-

dependent transcription. *Nature*. 342:830–833. doi:10.1038/342830a0.

Argüello-Miranda, O., Y. Liu, N.E. Wood, P. Kositangool, and A. Doncic. 2018.

Integration of Multiple Metabolic Signals Determines Cell Fate Prior to

Commitment. *Mol. Cell*. 71:733-744.e11. doi:10.1016/j.molcel.2018.07.041.

Bähler, J., J.Q. Wu, M.S. Longtine, N.G. Shah, A. McKenzie, A.B. Steever, A. Wach, P.

Philippsen, and J.R. Pringle. 1998. Heterologous modules for efficient and versatile

PCR-based gene targeting in Schizosaccharomyces pombe. *Yeast*. 14:943–951.

doi:10.1002/(SICI)1097-0061(199807)14:10<943::AID-YEA292>3.0.CO;2-Y.

Baltus, A.E., D.B. Menke, Y.C. Hu, M.L. Goodheart, A.E. Carpenter, D.G. De Rooij, and

D.C. Page. 2006. In germ cells of mouse embryonic ovaries, the decision to enter

meiosis precedes premeiotic DNA replication. *Nat. Genet.* 38:1430–1434.

doi:10.1038/ng1919.

Bean, J.M., E.D. Siggia, and F.R. Cross. 2005. High functional overlap between MluI

cell-cycle box binding factor and Swi4/6 cell-cycle box binding factor in the G1/S

transcriptional program in Saccharomyces cerevisiae. *Genetics*. 171:49–61.

doi:10.1534/genetics.105.044560.

Berchowitz, L.E., A.S. Gajadhar, F.J. van Werven, A.A. De Rosa, M.L. Samoylova, G.A.

Brar, Y. Xu, C. Xiao, B. Futcher, J.S. Weissman, F.M. White, and A. Amon. 2013. A

developmentally regulated translational control pathway establishes the meiotic

chromosome segregation pattern. *Genes Dev.* 27:2147–2163.

doi:10.1101/gad.224253.113.

Bowdish, K.S., H.E. Yuan, and A.P. Mitchell. 1994. Analysis of RIM11, a yeast protein

kinase that phosphorylates the meiotic activator IME1. *Mol. Cell. Biol.* 14:7909–

7919. doi:10.1128/mcb.14.12.7909-7919.1994.

Bowdish, K.S., H.E. Yuan, and A.P. Mitchell. 1995. Positive control of yeast meiotic

genes by the negative regulator UME6. *Mol. Cell. Biol.* 15:2955–2961.

doi:10.1128/mcb.15.6.2955.

Brar, G.A., M. Yassour, N. Friedman, A. Regev, N.T. Ingolia, and J.S. Weissman. 2012.

High-Resolution View of the Yeast Meiotic Program Revealed by Ribosome

Profiling. *Science (80-.).* 335:552–558. doi:10.1126/science.1215110.

Breitkreutz, A., H. Choi, J.R. Sharom, L. Boucher, V. Neduva, B. Larsen, Z.Y. Lin, B.J.

Breitkreutz, C. Stark, G. Liu, J. Ahn, D. Dewar-Darch, T. Reguly, X. Tang, R.

Almeida, Z.S. Qin, T. Pawson, A.C. Gingras, A.I. Nesvizhskii, and M. Tyers. 2010.

A global protein kinase and phosphatase interaction network in yeast. *Science (80-.*

). 328:1043–1046. doi:10.1126/science.1176495.

De Bruin, R.A.M., W.H. McDonald, T.I. Kalashnikova, J. Yates, and C. Wittenberg.

2004. Cln3 activates G1-specific transcription via phosphorylation of the SBF

bound repressor Whi5. *Cell*. 117:887–898. doi:10.1016/j.cell.2004.05.025.

Brush, G.S., N.A. Najor, A.A. Dombkowski, D. Cukovic, and K.E. Sawarynski. 2012.

Yeast ime2 functions early in meiosis upstream of cell cycle-regulated sbf and mbf

targets. *PLoS One*. 7:1–12. doi:10.1371/journal.pone.0031575.

Carlile, T.M., and A. Amon. 2008. Meiosis I Is Established through Division-Specific

Translational Control of a Cyclin. *Cell*. 133:280–291.

doi:10.1016/j.cell.2008.02.032.

Chen, J., D. Mcswiggen, and E. Ünal. 2018. Single Molecule Fluorescence In Situ

1139  Hybridization (smFISH) Analysis in Budding Yeast Vegetative Growth and

1140  Meiosis. 1–14. doi:10.3791/57774.

1141 Chen, J., A. Tresenrider, M. Chia, D.T. McSwiggen, G. Spedale, V. Jorgensen, H. Liao,

1142  F.J. Van Werven, and E. Ünal. 2017. Kinetochore inactivation by expression of a

1143  repressive mRNA. *Elife*. 6:1–31. doi:10.7554/eLife.27417.

1144 Cheng, Z., G.M. Otto, E.N. Powers, A. Keskin, P. Mertins, S.A. Carr, M. Jovanovic, and

1145  G.A. Brar. 2018. Pervasive, Coordinated Protein-Level Changes Driven by

1146  Transcript Isoform Switching during Meiosis. *Cell*. 172:910-923.e16.

1147  doi:10.1016/j.cell.2018.01.035.

1148 Chia, M., A. Tresenrider, J. Chen, G. Spedale, V. Jorgensen, E. Ünal, and F.J. van

1149  Werven. 2017. Transcription of a 5' extended mRNA isoform directs dynamic

1150  chromatin changes and interference of a downstream promoter. *Elife*. 6:1–23.

1151  doi:10.7554/eLife.27420.

1152 Chia, M., and F.J. van Werven. 2016. Temporal Expression of a Master Regulator

1153  Drives Synchronous Sporulation in Budding Yeast. *G3:*

1154  *Genes|Genomes|Genetics*. 6:3553–3560. doi:10.1534/g3.116.034983.

1155 Colomina, N., E. Garí, C. Gallego, E. Herrero, and M. Aldea. 1999. G1cyclins block the

1156  Ime1 pathway to make mitosis and meiosis incompatible in budding yeast. *EMBO*

1157  *J.* 18:320–329. doi:10.1093/emboj/18.2.320.

1158 Costanzo, M., J.L. Nishikawa, X. Tang, J.S. Millman, O. Schub, K. Breitkreuz, D. Dewar,

1159  I. Rupes, B. Andrews, and M. Tyers. 2004. CDK activity antagonizes Whi5, an

1160  inhibitor of G1/S transcription in yeast. *Cell*. 117:899–913.

1161  doi:10.1016/j.cell.2004.05.024.

1162 Van Dalfsen, K.M., S. Hodapp, A. Keskin, G.M. Otto, C.A. Berdan, A. Higdon, T.

1163  Cheunkarndee, D.K. Nomura, M. Jovanovic, and G.A. Brar. 2018. Global Proteome

1164  Remodeling during ER Stress Involves Hac1-Driven Expression of Long

1165  Undecoded Transcript Isoforms. *Dev. Cell*. 46:219-235.e8.

1166  doi:10.1016/j.devcel.2018.06.016.

1167 Dorsey, S., S. Tollis, J. Cheng, L. Black, S. Notley, M. Tyers, and C.A. Royer. 2018.

1168  G1/S Transcription Factor Copy Number Is a Growth-Dependent Determinant of

1169  Cell Cycle Commitment in Yeast. *Cell Syst.* 6:539-554.e11.

1170   doi:10.1016/j.cels.2018.04.012.

1171 Eisenberg, A.R., A.L. Higdon, I. Hollerer, A.P. Fields, I. Jungreis, P.D. Diamond, M.

1172   Kellis, M. Jovanovic, and G.A. Brar. 2020. Translation Initiation Site Profiling

1173   Reveals Widespread Synthesis of Non-AUG-Initiated Protein Isoforms in Yeast.

1174   *Cell Syst.* 11:145-160.e5. doi:10.1016/j.cels.2020.06.011.

1175 Espinoza, F.H., J. Ogas, I. Herskowitz, and D. Morgant. 2016. Cell Cycle Control by a

1176   Complex of the Cyclin HCS26 (PCL1) and the Kinase PHO85 Author (s): F .

1177   Hernan Espinoza , Joseph Ogas , Ira Herskowitz and David O . Morgan Published

1178   by : American Association for the Advancement of Science Stable URL : http://.

1179   266:1388–1391.

1180 Fridy, P.C., Y. Li, S. Keegan, M.K. Thompson, I. Nudelman, J.F. Scheid, M. Oeffinger,

1181   M.C. Nussenzweig, D. Fenyö, B.T. Chait, and M.P. Rout. 2014. A robust pipeline

1182   for rapid production of versatile nanobody repertoires. *Nat. Methods.* 11:1253–

1183   1260. doi:10.1038/nmeth.3170.

1184 Gallego, C., E. Garí, N. Colomina, E. Herrero, and M. Aldea. 1997. The Cln3 cyclin is

1185   down-regulated by translational repression and degradation during the G1 arrest

1186   caused by nitrogen deprivation in budding yeast. *EMBO J.* 16:7196–7206.

1187   doi:10.1093/emboj/16.23.7196.

1188 Gaubatz, S., G.J. Lindeman, S. Ishida, L. Jakoi, J.R. Nevins, D.M. Livingston, and R.E.

1189   Rempel. 2000. E2F4 and E2F5 play an essential role in pocket protein-mediated

1190   G1 control. *Mol. Cell.* 6:729–735. doi:10.1016/S1097-2765(00)00071-X.

1191 Gibson, D.G., L. Young, R.Y. Chuang, J.C. Venter, C.A. Hutchison, and H.O. Smith.

1192   2009. Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat.*

1193   *Methods.* 6:343–345. doi:10.1038/nmeth.1318.

1194 Harris, A., and E. Ünal. 2023. The transcriptional regulator Ume6 is a major driver of

1195   early gene expression during gametogenesis. *Genetics.* 225:1–21.

1196   doi:10.1093/genetics/iyad123.

1197 Harris, M.R., D. Lee, S. Farmer, N.F. Lowndes, and R.A.M. de Bruin. 2013. Binding

1198   Specificity of the G1/S Transcriptional Regulators in Budding Yeast. *PLoS One.*

1199   8:1–7. doi:10.1371/journal.pone.0061059.

1200 Hartwell, L.H., J. Culotti, J.R. Pringle, and B.J. Reid. 1974. Genetic control of the cell

division cycle in yeast. *Science*. 183:46–51. doi:10.1126/SCIENCE.183.4120.46.

Haruki, H., J. Nishikawa, and U.K. Laemmli. 2008. The Anchor-Away Technique: Rapid,

Conditional Establishment of Yeast Mutant Phenotypes. *Mol. Cell*. 31:925–932.

doi:10.1016/j.molcel.2008.07.020.

Hasan, M., S. Brocca, E. Sacco, M. Spinelli, P. Elena, L. Matteo, A. Lilia, and M.

Vanoni. 2014. A comparative study of Whi5 and retinoblastoma proteins : from

sequence and structure analysis to intracellular networks. 4:1–24.

doi:10.3389/fphys.2013.00315.

Van Den Heuvel, S., and N.J. Dyson. 2008. Conserved functions of the pRB and E2F

families. *Nat. Rev. Mol. Cell Biol.* 9:713–724. doi:10.1038/nrm2469.

Holt, L., B. Tuch, J. Villén, A. Johnson, S. Gygi, and D. Morgan. 2009. Global analysis

of Cdk1 substrate phosphorylation sites provides insights into evolution. *Science*

(80-.). 325:1682–1686.

Honigberg, S.M. 2003. Signal pathway integration in the switch from the mitotic cell

cycle to meiosis in yeast. *J. Cell Sci.* 116:2137–2147. doi:10.1242/jcs.00460.

Ishiguro, K. ichiro, K. Matsuura, N. Tani, N. Takeda, S. Usuki, M. Yamane, M.

Sugimoto, S. Fujimura, M. Hosokawa, S. Chuma, M.S.H. Ko, K. Araki, and H. Niwa.

2020. MEIOSIN Directs the Switch from Mitosis to Meiosis in Mammalian Germ

Cells. *Dev. Cell*. 52:429-445.e10. doi:10.1016/j.devcel.2020.01.010.

Iyer, V.R., C.E. Horak, P.O. Brown, D. Botstein, V.R. Iyer, M. Snyder, and C.S. Scafe.

2001. Genomic binding sites of the yeast cell-cycle transcription factors SBF and

MBF. *Nature*. 409:533–538. doi:10.1038/35054095.

Johnston, G.C., J.R. Pringle, and L.H. Hartwell. 1977. Coordination of Growth With Cell

Division. *Exp. Cell Res.* 105:79–98.

Jorgensen, P., and M. Tyers. 2004. How cells coordinate growth and division. *Curr. Biol.*

14:1014–1027. doi:10.1016/j.cub.2004.11.027.

Julian, L.M., Y. Liu, C.A. Pakenham, D. Dugal-Tessier, V. Ruzhynsky, S. Bae, S.Y. Tsai,

G. Leone, R.S. Slack, and A. Blais. 2016. Tissue-specific targeting of cell fate

regulatory genes by E2f factors. *Cell Death Differ.* 23:565–575.

doi:10.1038/cdd.2015.36.

Kadosh, D., and K. Struhl. 1997. Repression by Ume6 involves recruitment of a

complex containing Sin3 corepressor and Rpd3 histone deacetylase to target

promoters. *Cell*. 89:365–371. doi:10.1016/S0092-8674(00)80217-2.

Kassir, Y., D. Granot, and G. Simchen. 1988. IME1, a positive regulator gene of meiosis

in S. cerevisiae. *Cell*. 52:853–862. doi:10.1016/0092-8674(88)90427-8.

Kelliher, C.M., M.W. Foster, F.C. Motta, A. Deckard, E.J. Soderblom, M.A. Moseley, and

S.B. Haase. 2018. Layers of regulation on cell-cycle gene expression in the

budding yeast *Saccharomyces cerevisiae*. *Mol. Biol. Cell*. mbc.E18-04-0255.

doi:10.1091/mbc.E18-04-0255.

Kim, D., J.M. Paggi, C. Park, C. Bennett, and S.L. Salzberg. 2019. Graph-based

genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat.*

Biotechnol. 37:907–915. doi:10.1038/s41587-019-0201-4.

Klein, F., P. Mahr, M. Galova, S.B.C. Buonomo, C. Michaelis, K. Nairz, and K. Nasmyth.

1999. A Central Role for Cohesins in Sister Chromatid Cohesion, Formation of

Axial Elements, and Recombination during Yeast Meiosis Genetic and biochemical

analyses have identified. *Cell*. 98:91–103.

Kojima, M.L., D.G. De Rooij, and D.C. Page. 2019. Amplification of a broad

transcriptional program by a common factor triggers the meiotic cell cycle in mice.

Elife. 8:1–32. doi:10.7554/eLife.43738.

Love, M.I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15:1–21.

doi:10.1186/s13059-014-0550-8.

Malathi, K., Y. Xiao, and A.P. Mitchell. 1997. Interaction of yeast repressor-activator

protein Ume6p with glycogen synthase kinase 3 homolog Rim11p. *Mol. Cell. Biol.*

17:7230–7236. doi:10.1128/mcb.17.12.7230.

Malathi, K., Y. Xiao, and A.P. Mitchell. 1999. Catalytic roles of yeast GSK3β/shaggy

homolog Rim11p in meiotic activation. *Genetics*. 153:1145–1152.

doi:10.1093/genetics/153.3.1145.

Mcinerny, C.J., J.F. Partridge, G.E. Mikesell, D.P. Creemer, and L.L. Breeden. 1997. A

novel Mcml-dependent element in the WI4 , CLN3 , CDC6 , and CDC4S7

promoters activates M / Gi-specific transcription. *Genes Dev.* 1277–1288.

doi:10.1101/gad.11.10.1277.

Miller, M.P., E. Ünal, G.A. Brar, and A. Amon. 2012. Meiosis I chromosome segregation
is established through regulation of microtubule-kinetochore interactions. *Elife.*
2012:1–30. doi:10.7554/eLife.00117.

Moreno-Torres, M., M. Jaquenoud, M.P. Péli-Gulli, R. Nicastro, and C. De Virgilio. 2017.
TORC1 coordinates the conversion of Sic1 from a target to an inhibitor of cyclin-
CDK-Cks1. *Cell Discov.* 3:1–12. doi:10.1038/celldisc.2017.12.

Parviz, F., and W. Heideman. 1998. Growth-independent regulation of CLN3 mRNA
levels by nutrients in Saccharomyces cerevisiae. *J. Bacteriol.* 180:225–230.
doi:10.1128/jb.180.2.225-230.1998.

Pertea, M., G.M. Pertea, C.M. Antonescu, T.C. Chang, J.T. Mendell, and S.L. Salzberg.
2015. StringTie enables improved reconstruction of a transcriptome from RNA-seq
reads. *Nat. Biotechnol.* 33:290–295. doi:10.1038/nbt.3122.

Pnueli, L., I. Edry, M. Cohen, and Y. Kassir. 2004. Glucose and Nitrogen Regulate the
Switch from Histone Deacetylation to Acetylation for Expression of Early Meiosis-
Specific Genes in Budding Yeast. *Mol. Cell. Biol.* 24:5197–5208.
doi:10.1128/mcb.24.12.5197-5208.2004.

Primig, M., R.M. Williams, E.A. Winzeler, G.G. Tevzadze, A.R. Conway, S.Y. Hwang,
R.W. Davis, and R.E. Esposito. 2000. The core meiotic transcriptome in budding
yeasts. *Nat. Genet.* 26:415–423. doi:10.1038/82539.

Raj, A., P. van den Bogaard, S.A. Rifkin, A. van Oudenaarden, and S. Tyagi. 2008.
Imaging individual mRNA molecules using multiple singly labeled probes. *Nat.*
Methods. 5:877–879. doi:10.1038/nmeth.1253.

Rubin-Bejerano, I., S. Mandel, K. Robzyk, and Y. Kassir. 1996. Induction of meiosis in
Saccharomyces cerevisiae depends on conversion of the transcriptional represssor
Ume6 to a positive regulator by its regulated association with the transcriptional
activator Ime1. *Mol. Cell. Biol.* 16:2518–2526. doi:10.1128/mcb.16.5.2518.

Rundlett E., S., A. Carmen A., N. Suka, B. Turner M., and M. Grunstein. 2003.
Transcriptional repression byUME6 involves deacetylationof lysine5 of
histoneH4byRPD3. *Nature.* 426:181–186.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S.
Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.Y. Tinevez, D.J. White, V.

Hartenstein, K. Eliceiri, P. Tomancak, and A. Cardona. 2012. Fiji: An open-source platform for biological-image analysis. *Nat. Methods*. 9:676–682. doi:10.1038/nmeth.2019.

Sidorova, J.M., and L.L. Breeden. 2002. Precocious S-phase entry in budding yeast prolongs replicative state and increases dependence upon Rad53 for viability. *Genetics*. 160:123–136.

Simon, I., J. Barnett, N. Hannett, C.T. Harbison, N.J. Rinaldi, T.L. Volkert, J.J. Wyrick, J. Zeitlinger, D.K. Gifford, T.S. Jaakkola, and R.A. Young. 2001. Serial Regulation of Transcriptional Regulators in the Yeast Cell Cycle. 106:697–708.

Smith, H.E., S.E. Driscoll, R.A.L. Sia, H.E. Yuan, and A.P. Mitchell. 1993. Genetic evidence for transcriptional activation by the yeast IME1 gene product. *Genetics*. 133:775–784. doi:10.1093/genetics/133.4.775.

Smolka, M.B., F.M. Bastos De Oliveira, M.R. Harris, and R.A.M. De Bruin. 2012. The checkpoint transcriptional response: Make sure to turn it off once you are satisfied. *Cell Cycle*. 11:3166–3174. doi:10.4161/cc.21197.

Spellman, P., G. Sherlock, M. Zhang, V. Iyer, K. Anders, M. Eisen, P. Brown, D. Botstein, and B. Futcher. 1998. Comprehensive identification of cell cycle-regulated genes of the yeast Saccharomyces cerevisiae by microarray hybridization. *Mol. Biol. Cell*. 9:3273–3297. doi:10.1091/mbc.9.12.3273.

Subramanian, A., P. Tamayo, V.K. Mootha, S. Mukherjee, B.L. Ebert, M.A. Gillette, A. Paulovich, S.L. Pomeroy, T.R. Golub, E.S. Lander, and J.P. Mesirov. 2005. Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* 102:15545–15550. doi:10.1073/pnas.0506580102.

Talarek, N., E. Cameroni, M. Jaquenoud, X. Luo, S. Bontron, S. Lippman, G. Devgan, M. Snyder, J.R. Broach, and C. De Virgilio. 2010. Initiation of the TORC1-Regulated G0 Program Requires Igo1/2, which License Specific mRNAs to Evade Degradation via the 5′-3′ mRNA Decay Pathway. *Mol. Cell*. 38:345–355. doi:10.1016/j.molcel.2010.02.039.

Tresenrider, A., K. Morse, V. Jorgensen, M. Chia, H. Liao, F.J. van Werven, and E. Ünal. 2021. Integrated genomic analysis reveals key features of long undecoded

transcript isoform-based gene repression. *Mol. Cell*. 81:2231-2245.e11.

doi:10.1016/j.molcel.2021.03.013.

Tresenrider, A., and E. Ünal. 2017. One-two punch mechanism of gene repression: a

fresh perspective on gene regulation. *Curr. Genet.* 0:1–8. doi:10.1007/s00294-017-

0793-5.

Vidan, S., and A.P. Mitchell. 1997. Stimulation of yeast meiotic gene expression by the

glucose-repressible protein kinase Rim15p. *Mol. Cell. Biol.* 17:2688–2697.

doi:10.1128/mcb.17.5.2688.

Wagner, M. V., M.B. Smolka, R.A.M. de Bruin, H. Zhou, C. Wittenberg, and S.F. Dowdy.

2009. Whi5 regulation by site specific CDK-phosphorylation in Saccharomyces

cerevisiae. *PLoS One*. 4. doi:10.1371/journal.pone.0004300.

Vander Wende, H.M., M. Gopi, M. Onyundo, C. Medrano, T. Adanlawo, and G.A. Brar.

2022. Meiotic resetting of the cellular Sod1 pool is driven by protein aggregation,

degradation, and transient LUTI-mediated repression. *J. Cell Biol.* 222.

doi:10.1101/2022.06.28.498006.

van Werven, F.J., and A. Amon. 2011. Regulation of entry into gametogenesis. *Philos.*

Trans. R. Soc. B Biol. Sci. 366:3521–3531. doi:10.1098/rstb.2011.0081.

1342

1343

1344

Supplemental Figure Legends

S1. Swi4 and Ime1 protein abundance, Related to Figure 1

A. Wild type (UB22199) and *pATG8-SWI4* (UB22226) cells collected between 0-4 h in SPO. Immunoblot performed using α-V5 to quantify Swi4-3V5 abundance. Normalized to Hxk2 loading control. **B.** Quantification of the immunoblot in (A). **C.** Immunoblot using α-GFP to quantify GFP-Ime1 abundance. Normalized to Hxk2 loading control. **D.** Quantification of the immunoblot in (C).

S2. Gene ontology analysis for mRNA-seq comparing *pATG8-SWI4* to wild type, Related to Figure 2

A. Gene ontology analysis for genes that have significant increased expression by DESeq2 (see Materials and Methods) in *pATG8-SWI4* (UB2226) and wild type (UB22199*)*. Term size is the number of genes within a defined term. Enrichment score calculated for each term and plotted. **B.** same as in (A) but for genes that have significant decreased expression.

S3. Meiotic entry upon removal of either Cln1 or Cln2, Related to Figure 3

A. Live-cell imaging of meiotic cells marked by Rec8-GFP and nuclear marker Htb1-mCherry, with the following genotypes: wild type (UB32085), *pATG8-SWI4* (UB32089)*, cln2Δ* (UB35595), and *cln1Δ* (UB35597). Quantification of cells that entered meiosis assayed by the initial timing of nuclear Rec8 appearance. Experiments were performed using two biological replicates, mean value plotted with range. A total number of 800 cells were analyzed.

S4. Ime1 protein levels when C*LN2* is overexpressed and rescue of I*ME1* transcript levels and protein abundance with *pCUP1-IME1*, Related to Figure 4

A. Immunoblot was performed on wild type (UB22199) and *pATG8-CLN2* (UB35106) cells collected between 0-6 h in SPO. Immunoblotted using α-GFP to quantify GFP-Ime1 abundance. Normalized to Hxk2 loading control. Representative blots from one of two biological replicates are shown. All blots in this figure were performed the same way. **B.** Quantification of the immunoblot in (A). **C.** RT-qPCR was performed on *IME1* transcript

for samples collected in (A). Quantification was performed in reference to levels of meiotic housekeeping gene *PFY1* and then normalized to wild type. FC = fold change. Experiments were performed in duplicate, mean value plotted with range. **D.** Same as in (C) but for *pCUP1-GFP-IME1* (UB34641) and *pCUP1-GFP-IME1*; *pATG8-CLN2* (UB35057). **E.** Immunoblot was performed on samples from wild type (UB22199)*, pATG8-CLN2* (UB35106), *pCUP1-GFP-IME1* (UB34641), and *pCUP1-GFP-IME1*; *pATG8-CLN2* (UB35057) strains collected between 0-4 h in SPO. 50 μM $CuSO_4$ after 2 h in SPO was added to all cells. Representative blots from one of two biological replicates are shown. **F.** Quantification of the immunoblot in (E).

S5. Expression of *SWI4^LUTI^* in Ume6(T99N) mutant, Related to Figure 5

Fold change expression in wild type or Ume6(T99N) mutant between pre meiotic conditions and meiotic prophase (Tresenrider et al., 2021).

S6. Whi5 localization in early meiosis, Swi4 protein levels matching Figure 6A, 6C, 6D, and GSEA data for Δ*LUTI*; WHI5-AA, Related to Figure 6

A. Fixed imaging of cells marked with GFP-Ime1 and Whi5-mCherry with DNA stained with DAPI. Representative images with merge at 0 h and 2 h in SPO. Representative cells outlined. **B.** Left: Cells with the following genotypes were collected between 0-6 h in SPO: wild type (UB27083), Δ*LUTI* (UB26874), *WHI5-AA* (UB25431), Δ*LUTI; WHI5-AA* (UB25428). Immunoblotted using α-V5 to quantify Swi4-3V5 abundance. Normalized to Hxk2 loading control. Representative blots from one of two biological replicates are shown. Right: Quantification of blots. **C.** GSEA analysis of mRNA-seq comparing wildtype vs. Δ*LUTI; WHI5-AA (UB25428)* cells collected at 2 h in SPO. Vertical black bars represent early meiotic gene cluster from (Brar et al., 2012) or SBF cluster from (Iyer et al., 2001). The heatmap indicates genes that are more enriched in Δ*LUTI; WHI5-AA* (red, left-side) or genes that are enriched with wild type (blue, right-side). NES = normalized enrichment score. Enrichment was determined comparing wild type to Δ*LUTI; WHI5-AA.*

Supplement Table 1.

Table of differentially expressed genes (DESeq2) plotted in Figure 2F for *pATG8-SWI4* (UB22226) vs. wild type (UB22199).

Supplement Table 2.

Sporulation efficiency of cells at 24 h in SPO media for Rec8-GFP (UB32085*)*, Rec8-GFP; *pATG8-SWI4* (UB32089). 200 cells counted per strain. Second table is sporulation efficiency of cells at 24 h in SPO media for the following genotypes: wild type (UB22199), *pATG8-CLN1* (UB32820), *pATG8-CLN2* (UB25959), *pCUP-GFP-IME1* (UB34641), *pCUP1-GFP-IME1; pATG8-CLN2* (UB35057), *PUS1-αGFP* (UB35593), *PUS1-αGFP; pATG8-CLN2* (UB35982), *UME6-αGFP* (UB35300), and *UME6-αGFP; pATG8-CLN2* (UB35177).

Supplement Table 3.

Table of differentially expressed genes (DESeq2) plotted in Figure 6A, 6C, and 6D. First tab is *WHI5-AA*; Δ*LUTI* vs. wild type. Second tab is *WHI5-AA* vs. wild type. Third tab is Δ*LUTI* versus wild type.

Supplement Table 4.

The genotypes of the strains used in this study.

Supplement Table 5.

Plasmids used in this study.

Supplement Table 6.

Primers used for qPCR and RNA blotting in this study.

Supplemental Table 7.

Image acquisition information.

A

wild type
pATG8-CLN1
pATG8-CLN2
****

pCUP1-IME1
pCUP1-IME1;
pATG8-CLN2
**

PUS1-αGFP
PUS1-αGFP;
pATG8-CLN2
*

UME6-αGFP
UME6-αGFP;
pATG8-CLN2
ns

% sporulation

B

GFP-Ime1 | Htb1-mCherry | merge

wild type

pATG8-CLN2

2h in SPO

C

wild type   pATG8-CLN2

% cells with nuclear Ime1

time in SPO (h)

D

pCUP1   GFP-IME1

Ime1 level

GFP Ime1

?

GFP Ime1
Ume6
URS1   EMG

E

GFP-Ime1 | Htb1-mCherry | merge

pCUP1-IME1

pCUP1-IME1;
pATG8-CLN2

2 h post IME1 induction

F

pCUP1-IME1

pCUP1-IME1;
pATG8-CLN2

****

GFP-Ime1 mean nuclear intensity
(x10³, a.u.)

G

Ime1
GFP
αGFP
Pus1

Pus1 αGFP GFP Ime1
Ume6
URS1   EMG
?

cytoplasm   nucleus

H

GFP-Ime1 | Htb1-mCherry | merge

PUS1-αGFP

pATG8-CLN2;
PUS1-αGFP

2 h in SPO

I

wild type
pATG8-CLN2
PUS1-αGFP
PUS1-αGFP
pATG8-CLN2

****
ns
ns

GFP-Ime1 mean nuclear intensity
(x10³, a.u.)

J

Ime1
GFP
αGFP
Ume6

αGFP GFP Ime1
Ume6
URS1   EMG
?

cytoplasm   nucleus

K

GFP-Ime1 | Htb1-mCherry | merge

UME6-αGFP

pATG8-CLN2;
UME6-αGFP

2 h in SPO

L

wild type
pATG8-CLN2
UME6-αGFP
pATG8-CLN2;
UME6-αGFP

****
*
****

GFP-Ime1 mean nuclear intensity
(x10³, a.u.)