Pharmacologically targeting KRAS[G12D] in PDAC models:

tumor cell intrinsic and extrinsic impact

Vishnu Kumarasamy[1], Costakis Frangou[1], Jianxin Wang[1], Yin Wan[1], Andrew Dynka[1], Hanna Rosenheck[1], Prasenjit Dey[2], Ethan V. Abel[1], Erik S. Knudsen*[1], Agnieszka K. Witkiewicz*[1]

[1] Department of Molecular and Cellular Biology, Roswell Park Cancer Institute, Buffalo NY

[2] Department of Immunology, Roswell Park Cancer Institute, Buffalo NY

Running Title: Targeting G12D-mutant KRAS.

Keywords: MRTX1133, Cell cycle, KRAS, TME, organoid, PDX

*<u>Correspondence</u>

Agnieszka K. Witkiewicz
Department of Molecular and Cellular Biology
Roswell Park Cancer Institute, Buffalo NY
agnieszka.witkiewicz@roswellpark.org

Erik S. Knudsen
Department of Molecular and Cellular Biology
Roswell Park Cancer Institute, Buffalo NY
Erik.Knudsen@roswellpark.org

Abstract

Pancreatic ductal adenocarcinoma (PDAC) is an aggressive disease for which new therapeutic interventions are needed.   Here we assessed the cellular response to pharmacological KRAS inhibition, which target the central oncogenic factor in PDAC. In a panel of PDAC cell lines, pharmaceutical inhibition of KRAS[G12D] allele, with MRTX1133 yields variable efficacy in the suppression of cell growth and downstream gene expression programs in 2D culture. CRISPR screens identify new drivers for enhanced therapeutic response that regulate focal adhesion and signaling cascades, which were confirmed by gene specific knockdowns and combinatorial drug synergy. Interestingly, MRTX1133 was considerably more efficacious in the context of 3D cell cultures and *in vivo* PDAC patient-derived xenografts. In syngeneic models, KRAS[G12D] inhibition elicited potent tumor regression that did not occur in immune-deficient hosts. Further investigation using digital spatial profiling, single cell sequencing, and multispectral imaging revealed that the immunological response was associated with remodeling of the tumor microenvironment, suppression of neutrophils and influx of effector CD8[+] T-cells. Thus, both tumor cell intrinsic and extrinsic events contribute to response and credential KRAS[G12D] inhibition as promising strategy for a large percentage of PDAC tumors.

Main: In spite of significant effort, pancreatic ductal adenocarcinoma (PDAC) remains a largely-therapy recalcitrant disease for which overall five-year survival remains ~11%[1-4].   Genetic analyses indicates that PDAC tumors harbor multiple high-potency oncogenic and tumor

suppressive lesions[5-7]. While a myriad of targeted therapeutic approaches have been tested in

clinical trials, in many contexts rationally targeted therapies exploiting genetic features of PDAC

have failed to exhibit superiority to chemotherapy. For example, MEK inhibitors which would be

expected to have broad efficacy in KRAS-driven PDAC have shown little efficacy[8].   Thus,

developing new therapeutic strategies which target the genetics of PDAC remain a largely

unrealized opportunity.

PDAC is dominated by oncogenic mutation of KRAS occurring in >95% of tumors[9].

Mutant KRAS acts through multiple signaling pathways that contribute to deregulated

proliferation, invasion, and metastasis[10,11]. The signaling mediated by mutant KRAS initiates at

the membrane and is transduced by multiple effectors that represent independent drug targets.

In PDAC cell lines, there is a variable response to KRAS depletion which generally mediates a

cytostatic response[11,12].   However, the genetic ablation of KRAS can be tolerated and such

genetically modified PDAC cell lines can develop into tumors in xenograft models[13].   The

targeted deletion of KRAS in GEMM models can lead to tumor regression; however, KRAS-

independent tumors can evolve, with acquired resistance associated with the activation of

bypass pathways (e.g. EGFR, AKT, YAP) that abrogate the dependence[14-16].

The tumor microenvironment is particularly relevant in PDAC where the stromal

compartment can contribute to the bulk of the tumor volume[17,18].   This stroma has been

proposed to both serve tumor inhibitory and promoting functions[19,20]. In particular, it has been

believed that the stroma contributes to exclusion of functional anti-tumor immune functions, and

contributes to the lack of efficacy of immune checkpoint inhibitors and other forms of

immunotherapy[4,21].   Studies with the genetic modulation of KRAS have illustrated that tumor-

mediated events contribute to broad ranging effects on the tumor microenvironment[22].

KRAS inhibitors have been developed to the stage of FDA-approval in the context of

agents selectively directed against the G12C mutant allele[23].   This mutation of KRAS is present

at relatively low frequency (1-2%) in PDAC[10].   However, there is evidence that a G12C inhibitor

(ie. Sotorasib) can have clinical activity in PDAC[24], although there are clearly distinct

mechanisms through which resistance can emerge[25,26]   The most prevalent KRAS mutation in

PDAC is G12D[10,27].   Recently the agent MRTX1133 was described as a highly-selective and

potent inhibitor of G12D, which has been shown to have efficacy in PDAC models[28,29]. Here we

explored the functional response to G12D inhibition in a spectrum of models that define features

of sensitivity and resistance to this agent and underscore the importance of the tumor

microenvironment on therapeutic efficacy.

Results

Heterogeneous cellular response to MRTX1133 in PDAC models:

To define the impact of a pharmacological KRAS inhibitor, MRTX1133 we employed a panel of KRAS^{G12D} mutant PDAC cell lines that include both established and patient derived models [45]. Following the treatment with MRTX1133 we observed a highly varied response based on the proliferation of the cells as monitored by live cell imaging. The established PDAC cell lines, ASPC1 and HPAF-II and a patient derived cell line, 828, were sensitive to MRTX1133 and displayed a robust cytostatic response (Fig. 1A). Interestingly, the other patient derived PDAC cell lines, 519, 1222 and 3226 were largely resistant, as cellular proliferation continued despite the presence of MRTX1133 (Fig.1A & S1). The EC50 values for MRTX1133 in the sensitive models were at the range of 25-50 nM, while in the resistant models, the EC50 is greater than 1 µM (Fig. 1B). To validate the specificity of MRTX1133 to KRAS^{G12D} mutants, we evaluated the anti-proliferative effect of this drug in KRAS^{G12C} mutant cell lines MiaPaCa2 and a patient derived PDAC cell line, UM53 (Fig. S1). As a control, we included the KRAS^{G12C} inhibitor, MRTX849. Based on the growth curves, the proliferation of MiaPaCa2 cells was robustly inhibited by MRTX849, whereas the MRTX1133 had no effect, confirming that the KRAS^{G12D} is its intracellular target (Fig. S1). Although, in UM53 cell line, the response to MRTX849 is modest, there is a significant difference in response as compared to MRTX1133 (Fig. S1). To interrogate how different cellular mechanisms are impacted by MRTX1133 to mediate an anti-proliferative effect, transcriptome analysis was carried out in different PDAC cell lines in the presence and absence of the KRAS inhibitor. Based on differential gene expression analysis the sensitive models, ASPC1, HPAF-II and 828 displayed a relatively greater number of genes that were differentially expressed in the presence of MRTX1133 as compared to the resistant models 519, 827 and 1229 indicating that the resistant models are essentially devoid of a transcription output downstream from KRAS inhibition (Fig.1C & D). Gene set enrichment analysis (GSEA) on the cell lines indicated that the downregulated genes are associated with

E2F targets, MYC targets, G2/M check point and MTOR signaling (Fig. 1E). However, the

magnitude of repression differs among the models, which correlates with their sensitivity to

MRTX1133 (Fig. 1F). Since, the E2F target genes are known to regulate cell cycle progression

via RB inactivation, biochemical analysis was performed in the PDAC cell lines following the

treatment with MRTX1133. In the sensitive models, 828 and ASPC1 RB phosphorylation was

significantly inhibited, which further resulted in the suppression of cyclin A, which is an E2F

target (Fig. 1G). As expected, in the resistant models, 3226 and 519, RB remained

phosphorylated with modest impact on cyclin A, indicating that cell cycle remains ongoing in the

presence of the KRAS inhibitor (Fig. 1G). Consistent with the biochemical data, MRTX1133

significantly inhibited BrdU incorporation in the sensitive models confirming a robust cell cycle

arrest, while moderate or no effect was observed in the resistant models (Fig. S1). Interestingly,

the phosphorylation of ERK was inhibited across the PDAC models irrespective of their

sensitivities to MRTX1133, indicating that at least this downstream pathway of KRAS is

inhibited.   However, this was not sufficient to inhibit RB phosphorylation and expression of

downstream proteins (e.g. Cyclin A) in the resistant models (Fig. 1G). To confirm the inhibition

of ERK phosphorylation by MRTX1133 is a consequence of KRAS^{G12D} inhibition, we compared

its effect with MRTX849 on the MiaPaCa2 cells. Our data confirmed that inhibition of ERK

phosphorylation is not affected by MRTX1133 (Fig. S1). In conclusion, although different PDAC

models harbor oncogenic KRAS G12D mutation, there is heterogeneity in response to the

pharmacological inhibitor, which differentially impacts on cell cycle.

Genetic modifiers of the response to KRAS inhibitors:

To systematically and comprehensively identify critical genes involved in resistance and

sensitivity to KRAS inhibition, we employed a genome wide loss-of-function CRISPR/Cas9

library screen in a KRAS^{G12D} mutant PDAC cell line, ASPC1. Following the infection with the

CRISPR/Cas9 pooled library that contains 70,948 unique single guide RNAs (sgRNAs) that

target 18,053 genes with 4 sgRNAs per gene the cells were grown in the presence and absence

of MRTX1133 at low concentration (6.25 nM) for 11 passages (Fig. 2A). The sgRNAs that gets

depleted or enriched following the treatment with MRTX1133 imply that the loss of function of

their target genes could enhance or attenuate the cytostatic effect of the drug respectively. As

expected sgRNAs targeting genes that regulate ribosome biogenesis, proteasome degradation,

RNA metabolism and cell cycle were significantly depleted irrespective of drug treatment

following 11 passages, indicating the essential function of those genes in cell survival (Fig. S2).

Moreover, a pair-wise comparison between the replicates indicated a correlation coefficient of

approximately 0.9, illustrating the reproducibility of the screen (Fig. S2). Differential

representation of individual sgRNA was determined based on the fold change between vehicle

and MRTX1133 treated groups to define the genes that are positively and negatively selected

(Fig. 2B). REACTOME pathway analyses indicated that the negatively selected sgRNAs target

gene sets are highly enriched in different mitogenic pathways, including EGFR signaling, ERK

signaling cascade, and metabolic pathways (Fig. 2C). We further implemented the DrugZ

algorithm as previously described[38], to calculate the NormZ values for the sgRNAs targeting

each gene. Based on this analysis we identified 235 genes with NormZ values <-1.9 that were

significantly depleted in the presence of MRTX1133 (Fig. 2D). The identified genes either

directly regulate the function of KRAS such as SOS1 or GRB2, encompass parallel pathways to

drive downstream signaling (e.g. EGFR) or act as downstream effectors such as *BRAF,*

MAPK3, and *FOSL1*, suggesting that these genes contribute to resistance to MRTX1133 (Fig.

2D). The individual guides that target *EGFR* and *FOSL1*, revealed that at least 3 sgRNAs were

depleted in the presence of MRTX1133 (Fig. 2E). The functional roles of EGFR and FOSL1 on

the impact of MRTX1133 were further validated by gene specific depletion in ASPC1 cells.

Depletion of EGFR and FOSL1 with interfering RNA significantly cooperated with MRTX1133

and yielded potent growth suppression in ASPC1 cells as monitored by live cell imaging (Fig.

2F, Fig. S2). Among the sgRNAs that were positively selected following MRTX1133 treatment,

KEAP1 was most enriched, which is consistent with previous studies (Fig. 2D) [29,46].

Furthermore, sgRNAs targeting *ERRFI1* and *PTEN* which act as negative regulators of EGFR

singling and mTOR activity respectively were positively selected, supporting the functional

importance of the mitogenic signaling pathway in bypassing KRAS inhibition (Fig. 2D). Individual

guides targeting these genes further confirmed that MRTX1133 treatment increased the counts

of all four sgRNAs (Fig. 2E). Moreover, since MRTX1133 renders anti-proliferative effect via RB

activation, sgRNA targeting RB1 was positively selected, implying that loss of this gene could

attenuate the effect of KRAS inhibition (Fig. 2D). As a validation we employed a

pharmacological approach where the ASPC1 cell line was were exposed to different EGFR

inhibitors in combination with MRTX1133. Based on the relative growth rate it was evident that a

subset of EGFR inhibitors cooperated with MRTX1133 to induce a cytostatic response and

Gefitinib significantly enhanced the efficacy of MRTX1133 in ASPC1 cells (Fig. S2). While

ASPC1 is intrinsically sensitive to MRTX1133, it is expected that targeting EGFR-mediated

signaling pathway could hypersensitize the cells to KRAS inhibition.

To further identify gene sets that could reverse the sensitivity to KRAS inhibition

CRISPR/CAS9 screening was also performed in UM53, which is resistant to the KRAS[G12C]

inhibitor, MRTX849. Moreover, the use of this G12C model would allow us to define consistent

features of KRAS inhibition with mechanistically distinct agents. Integrin subunit beta 1 (ITGB1)

was identified as a negatively selected gene upon MRTX849 treatment (Fig. 2G). sgRNAS

targeting, *CAV1*, TAZ and TEAD2 that are associated with integrin-mediated

mechanotransdcution were also dropped out and the individual guides were depleted following

MRTX849 treatment in UM53 cells (Fig. 2G & H)[47,48]. Similarly, the individual guides against

ITGB1 in ASPC1 cells were selectively depleted in the presence of MRTX1133 (Fig. S2).

Further validation was performed using gene specific RNAi that target ITGB1 in combination

with KRAS inhibition in 4 different PDAC cell lines. Although ASPC1 is sensitive to MRTX1133,

deletion of ITGB1 modestly enhanced the efficacy of the drug (Fig. 2I). In the resistant cell lines,

3226, 1222 and UM53, ITGB1 deletion significantly reversed the sensitivity to KRAS inhibition

(Fig. 2I). Long term colony formation assay further confirmed that depletion of ITGB1 enhanced

the anti-proliferative effect of KRAS inhibition in 3226 and UM53 cell lines (Fig. S2). Overall our

data suggest that the focal adhesion to ECM might overcome the response to KRAS inhibition.

This observation is similar to a previous study, which has demonstrated that the anchorage

dependent growth of KRAS mutant models can attenuate the impact of KRAS inhibition [49].

Potent efficacy of MRTX1133 in 3D cell culture and in vivo settings:

Since anchorage dependent cell growth could overcome KRAS dependency, we

assessed the efficacy of MRTX1133 in organoids, derived from the PDAC cell lines, 3226 and

519 that are intrinsically resistant to KRAS inhibition [13,50]. Surprisingly, the 3226 and 519

organoids underwent a significant suppression of proliferation with MRTX1133 treatment at a

concentration which had no effect in 2D culture (Fig. 3A & B). We also evaluated the effect of

KRAS^{G12C} inhibitor, MRTX849, on organoids derived from UM53 cell line, which is also relatively

resistant to KRAS inhibition. Similar to the G12D models, MRTX849 significantly delayed the

growth of UM53 organoids (Fig. S3). Consistent with its efficacy against organoid growth,

MRTX1133 displayed a cytostatic effect on tumor growth *in vivo* against two different PDX

models (828 and 3226) (Fig. 3C & D). Since, the tumor microenvironment (TME) and features of

the immune system play a major role in the pathogenesis of PDAC, we additionally investigated

the effect of MRTX1133 in an immunocompetent setting[22].   We developed a syngeneic PDAC

model derived from *Kras$^{+/LSL-G12D}$;Pdx1-Cre* (KC) mice termed KC4568 (Fig. S4). We also

employed two additional mouse syngeneic models that were previously established from KPC

mouse (4662)[51] and the iKRAS^{G12D} model where the oncogenic KRAS^{G12D} is induced by

Doxycycline (AKB6)[52]. Similar to the human cell lines, the mouse models displayed differential

response to MRTX1133 in cell culture. BrdU incorporation in the 4662 and KC-4568 cell lines

was not inhibited in the presence of MRTX1133 indicating that these models are resistant to

KRAS inhibition (Fig. 3E & S3). However, the proliferation of iKRAS AKB6 cell line in the

presence of MRTX1133 was suppressed, which further resulted in the inhibition of BrdU

incorporation (Fig. 3E & S3). Interestingly, similar to the human cell lines, the 4662 model

possessed a more profound response to MRTX1133 when grown in organoids as compared to

the 2D cell culture (Fig. S3). The *in vivo* anti-tumor effect of MRTX1133 was examined in

C57BL6J mice, bearing tumors derived from the KC-4568 and AKB6 cell lines. KRAS inhibition

resulted in durable disease control in both the xenografts; however, unlike the PDX models,

there was significant tumor regression (Fig. 3F & S3). Furthermore, H&E staining revealed

major changes in the histology with fewer tumor cell population and enhanced collagen

deposition in both KC4568 and AKB6 xenografts, indicating that MRTX1133 alters TME (Fig.

3G).

Mechanistic impact of MRTX1133 in the tumor microenvironment:

To examine the molecular pathways that are perturbed by MRTX1133 in the tumor

component, which could further impact the TME, we performed single cell RNA sequencing on

pooled AKB6 tumor tissues that were treated with vehicle and MRTX1133.   The single cell

sequencing resolved multiple cell types including cancer cells, T-cells, Neutrophils,

macrophages, dendritic cells, and cancer-associated fibroblasts (CAFs) within the tumors (Fig.

S4 & S5). Among the different cell clusters, a particularly striking difference was selectively

observed in tumor cells, T cells and neutrophils following MRTX1133 treatment (Fig. 4A, Fig.

S4). The tumor population existed in two distinct subpopulations, which both express high levels

of epithelial marker genes such as *Krt7, Krt8* and *Krt19.* However, in one of the subpopulations

epithelial associated genes such as *Cdh1, Lamc2, Plec, Ecm1* were expressed at lower levels

suggesting that a subset of tumor cells is undergoing and epithelial to mesenchymal transition

(EMT) (Fig. 4B). Interestingly, both the cancer populations were dramatically decreased

following MRTX1333 treatment (Fig. 4C). Further examination of differentially expressed genes

in the tumor population revelaed that MRTX1133 resulted in upregulation of genes such as

Cxcl9, Irgm1, Irf1 that are associated with immune-related pathways, including the activation of

Interferon-γ signaling (Fig. 4D & E).

As a validation approach, we performed digital spatial profiling (DSP) whole

transcriptome analysis on paraffin embedded tumor tissues. The tissues were stained with

fluorescent tagged anti-PANCK and anti-CD45 antibodies that selectively recognizes the tumor

and the stromal population respectively along with a cocktail of antibodies conjugated with DNA

indexing oligonucleotides. Based on the fluorescent signal, digital masking was performed that

allows to demarcate the tumor and stromal cells (Fig. 4F). Multiple regions of interest (ROIs)

were selected and illuminated to release the oligos that are digitally counted and based on

which the differentially expressed genes were determined (Fig. 4G). Pathway analysis on the

downregulated genes against the Hallmark genesets in the Molecular Signatures Database

(MSigDB)[41-44] revealed a significant enrichment on pathways including hypoxia, glycolysis and

MTOR signaling, confirming the impact of KRAS inhibition (Fig. S6). Based on the normalized

counts, it was evident that MRTX1133 resulted in the suppression of the genes involved in

those pathways across all the selected ROIs (Fig. S6). Similar analysis on the upregulated

genes, displayed a high enrichment on different immune-related pathways, which is consistent

with the single cell sequencing data (Fig. S6). GSEA analysis further confirmed that the

upregulated genes were significantly enriched for Interferon-γ signaling, which resulted in the

upregulation of MHC antigen presentation molecules (Fig. 4H & Fig. 4G). The critical genes

involved in the interferon-like pathways such as *CD74, GBP2, H2-Ab1, CXCL10, IRF1 etc.* were

prominently upregulated by MRTX1133 across all the ROIs (Fig. 4I). GSEA analysis based on

the transcriptome data from ASPC1 and 828 cell lines revealed that MRTX1133-mediated

upregulated genes were also significantly enriched to the Interferon-γ signaling, indicating that

the impact of KRAS inhibition on the immune component is highly conserved between *in vitro*

and *in vivo* settings and tumor cell autonomous (Fig. 4J & S6). The expression of *HLA-DMA,*

HLA-E and *HLA-F,* which are involved in antigen presentation were significantly upregulated in

ASPC1, HPAF-II and 828 cells (Fig. 4K). Together these data provide evidence that the

pharmacological inhibition of KRAS upregulates the intra-tumoral antigen presentation and

Interferon-like signaling pathways that are known to enhance immune infiltration to the TME[53-55]

MRTX1133 broadly modulates the tumor immune microenvironment

We further analyzed the single cell RNA seq data to investigate the effect of MRTX1133

on other immune components in the TME. An important immune component that was

modulated by MRTX1133 is the neutrophil population, which was prominently decreased in the

drug treated group (Fig. 5A). Prior studies have shown that neutrophils promote tumor growth

via inhibiting the activation of effector CD8$^+$ T cells[56,57]. Based on the differential gene

expression in the neutrophils following MRTX1133 treatment, the downregulated genes were

associated with metabolism and hypoxia (Fig. S7). Expression of genes involved in these

pathways such as *Hilpda, Ldha, Eno1* were significantly downregulated by MRTX1133, which

could inhibit the survival of neutrophils and enhance T-cell activation (Fig. 5B & C). Additionally,

the increase in antigen presentation in tumor cells could also recruit T cells. Consistent to this

notion, MRTX1133 treatment notably increased the T-cell populations that comprise, CD4$^+$/

CD8$^+$ T cells and natural killer T cells (NKT) (Fig. 5D & Fig. S4). Among the different T-cell

population, a prominent increase was observed in the CD8$^+$ T cells (Fig. 5E). Increase in

interferon gama (*Ifng)* was selectively observed in the CD8$^+$ T-cells following MRTX1133

treatment and the upregulation of *Icos and CD69* was observed in both CD4$^+$ /CD8$^+$ T cells and

NKT cells, indicating the activation of effector T cells to induce anti-tumor immunity (Fig. 5D &

5F, Fig. S7)[22]. Moreover, the cytolytic function of T-cells was enhanced by MRTX1133, as

evident by the upregulation of *Prf1* in the CD8$^+$ T cells and *Klrb1b* in the NKT cells (Fig. 5D, Fig.

S7). To validate this observation, tumor tissues from the vehicle and MRTX1133 treated groups

were subjected to multi-spectral staining that comprises Pan-CK, DAPI and the T-cell marker,

CD8. Consistent to the single cell sequencing data, MRTX1133-treated tissues exhibited a

significantly higher CD8 expression in both the tumor and stroma confirming that MRTX1133

induces T-cell activation (Fig. 5G). CAFs and macrophages, which are also an important

immune component in the TME were not notably modulated by MRTX1133 (Fig. S7). Gene

expression of markers specific for iCAFS (*C1s1, Col14a1, Has1)* and myCAFs (*Col12a1)* were

not altered following MRTX1133 treatment (Fig. S7)[22]. We also evaluated the effect of

MRTX1133 on AKB6 xenografts developed in immune deficient NSG mice. Although

MRTX1133 induced a cytostatic response against tumor growth, the disease control was more

profound in immune-competent mice (Fig. 5H). Overall our data suggests that oncogenic KRAS

inhibition by MRTX1133 possess a major impact on TME and enhance cytotoxic T-cell

infiltration to induce a durable disease control.

Discussion:

The mutation of KRAS is considered the initiating oncogenic driver for the pathogenesis

of PDAC[9,58]. While multiple different KRAS mutant isoforms occur in PDAC, the G12D mutation

is the most common occurring ~40% of cases[29,59]. Multiple targeted therapies have been

interrogated that inhibit KRAS-mediated downstream effector pathways in clinical trials,

although the therapeutic benefits are limited[60,61]. While KRAS was considered undruggable for

many years, structure-based drug design approaches have led to the development of multiple

selective pharmacological inhibitors [28,62].

The current work demonstrates the cellular response of a potent KRAS[G12D] inhibitor,

MRTX1133 in different patient-derived PDAC cell lines and syngeneic models. The efficacy of

MRTX1133 *in vitro* cell culture assays is heterogeneous among the PDAC models and

characterized as extremely sensitive or largely totally refractory. Based on our transcriptome

analysis, the top pathway that is significantly inhibited in the sensitive models is the cell cycle

machinery, which remained largely unperturbed in the resistant models, suggesting the

involvement of parallel pathways that allow to bypass the impact of KRAS inhibition and drive

cell cycle.   Consistent with this concept key mediators of KRAS and RTK signaling were

identified as cooperating with pharmacological mutant KRAS inhibitors using CRISPR/Cas9

screening.   The genes identified included those directly associated with controlling RAS activity

that could represent activity through other RAS genes, as well as downstream effectors (e.g.

BRAF and MAPK3).   Conversely, EGFR was a key mediator of resistance that could be further

targeted pharmacologically to block the ability of cells to bypass mutant KRAS inhibition.

Genes whose loss associated with resistance are known to further deregulate signaling

pathways (e.g. PTEN) or limit effects on cell cycle (e.g. RB).   These findings advance a

number of combinatorial strategies that could be deployed with MRTX1133 to enhance cell

autonomous activity on tumor cell division.

Our CRISPR screen analysis further revealed a cadre of genes whose depletion potently

cooperated with KRAS inhibition (MRTX1133 and MRTX849) that are associated with adhesion

mediated signaling. This includes integrins (ITGB1 and ITGAV) and genes involved in

mechanotransdcution (CAV1, YAP, TEAD2)[47,48,63]. Our study further illustrates that the

resistance to MRTX1133 in 2D culture could be alleviated when the cells are grown in

organoids, and this phenomenon was observed in both human and mouse PDAC models.

Presumably, integrin-mediated mechano-transduction is limited in the Matrigel matrix, thus

growth of organoids is considerably inhibited following KRAS inhibition. It has been posited that

cell culture yields enhanced sensitivity to therapeutic agents while the organoids or xenografts

are considered a more stringent system to assess clinically relevant drug efficacy[64]. However, in

the context of KRAS inhibition, cells cultured as organoids are considerably more sensitive to

therapeutic targeting[13,49]. The different PDAC models were also evaluated as xenografts and a

prominent anti-tumor effect was observed following the treatment with MRTX1133.

Interestingly, prior studies using different targeted agents or genetic depletion of KRAS did not

indicate a drastic difference in efficacy between the 2D cell culture and organoids [32,65].

However, the data here highlight the importance of the 3D cell culture or organoids to be used in

an *in vitro* setting to more accurately predict the efficacy of KRAS inhibitors against tumor

growth.

Although MRTX1333 inhibited tumor growth in all the G12D mutant models tested, a

significant difference on disease control was observed between immune-deficient NSG strain

and an immune-competent C57BL/6 strain. While loss of KRAS was known to impact the TME,

our study illustrates that pharmacological inhibition of KRAS could yield a similar effect[22]. Single

cell sequencing revealed highly significant changes in TME following the treatment with

MRTX1133. Suppression of tumor cell population along with increase in cytolytic and effector

$CD8^+/CD4^+$ T cells and decrease in neutrophils implies that MRTX1133-mediated tumor

regression is associated with TME remodeling. In this study we elucidated the mechanism

through which MRTX1133 enhances the infiltration of effector T cells by utilizing a more

sophisticated unbiased approach, Digital spatial Profiling (DSP) to determine the differential

gene expression selectively on the tumor population. This technique is more advanced than the

conventional bulk RNA sequencing, because it spatially discriminates the tumor population from

the stromal cells[66]. Based on our analysis from both single cell RNA sequencing and DSP, the

tumor cells displayed a prominent upregulation of genes involved in interferon γ signaling and

MHC class II antigen presentation. These molecular events have been proposed to enhance T-

cell recognition, activation of effector $CD8^+$ T cells and infiltration to eliminate the cancer

cells[53,67-69]. Neutrophils, one of the major components in the TME are known to promote

tumorigenesis by suppressing the function of effector T-cells[57]. The survival of neutrophils is

mediated by oxidative mitochondrial metabolism that results in a hypoxic condition and

enhances Reactive Oxygen Species (ROS) production to inhibit effector T-cell activation and

suppress the anti-tumor immunity[56]. The impact of MRTX1133 on tumor cells downregulates

genes associated with hypoxia and metabolism in neutrophil population that impacts their

viability and promotes T-cell activation.

The impact on TME and activation of effector T cells are direct consequences of KRAS

inhibition, because in our prior studies the treatment with a MEK inhibitor in combination with

CDK4/6 inhibitor did not result in tumor regression rather the tumor escaped the acute cytostatic

effect [53]. The MRTX1133-mediated tumor regression is reversible because cessation of

treatment caused tumor recurrence in one of the syngeneic models (data not shown), which is

consistent with a recent study[59]. Moreover, our single cell sequencing data revealed an increase

in the exhaustion markers such as *Ctla4* and *Lag3* that suppress the anti-tumor immunity,

suggesting new combinatorial treatment modalities involving concurrent targeting of KRAS and

immunotherapy (Fig. S7).

In summary, this study highlights the efficacy of a KRAS[G12D] inhibitor, MRTX1133 in

PDAC models and demonstrates a differential response in 2D cell culture, organoids and *in*

vivo. While we demonstrate two distinct molecular components, focal adhesion and TME

remodeling that modulate the response to MRTX1133, it is limited to preclinical setting. Whether

the genes involved in those pathways could serve as biomarkers of response or resistance to

KRAS inhibitors in patient models need further investigation.

Methods:

Cell Culture and Therapeutic agents: Human primary PDAC cell lines 828, 519, 3226, 1222,

and 827 and mouse primary PDAC cell line, KC4568 were grown in Keratinocyte Serum Free

medium (KSF), supplemented with 2% FBS, EGF (0.2 ng/mL) and bovine pituitary extract (30

μg/mL) (Life Technologies, Carlsbad, CA) as described before[30]. The TC dishes were precoated

with collagen (Millipore, Burlington, MA). UM53, ASPC1 and HPAF-II cell lines were kindly

provided by Dr. Ethan Abel (Roswell Park Cancer Center) and were grown in RPMI medium,

supplemented with 10% FBS. Mia PaCa-2 cell line was cultured in DMEM media, supplemented

with 10% FBS. Mouse KPC PDAC cell line, 4662 was obtained from Dr. Vonderheide's

laboratory (University of Pennsylvania) and grown in RPMI media, supplemented with 10% FBS. The iKRAS AKB6 was kindly provided by Dr. Prasenjit Dey (Roswell Park Cancer Center) and cultured in RPMI medium, supplemented with 10% FBS. All cell lines were maintained in 37°C with 5% CO_2 and tested to be Mycoplasma free. ASPC1, Mia PaCa-2 and HPAF-II were authenticated by STR profiling. MRTX1133 was purchased from Chemietek (Cat # CT-MRTX1133) and dissolved in DMSO to get a final concentration of 10 mM. For *in vivo* work, MRTX1133 was kindly provided by Mirati Therapeutics. Gefitinib was purchased from SelleckChem.

Cell Proliferation Assay: To monitor the cell growth, we employed live cell imaging systems, IncuCyte S3 and CellCyte X. The cell lines were transfected to stably express H2B-GFP and seeded in 96 well TC dish in the presence and absence of test agents. The GFP counts were measured in real time that corresponds to proliferation rate.

Knockdown experiments: The cell lines were reverse transfected with gene specific RNAis that target *ITGB1, FOSL1 and EGFR* as described in previous study[31]. On-target plus human RNAi for *ITGB1* was purchased from Horizon Discovery (Cat # L-004506-00-0005). siRNAs for *EGFR* (Cat # S563) and *FOSL1* (Cat # S15583) were purchased from Thermo Scientific. Following transfection, the cells were treated with MRTX1133 and the cell proliferation was monitored using live cell imaging.

Western Blotting: Whole cell extracts were prepared as described in our previous studies [30,32]. The primary antibodies for pERK and pRB (S807/811) (Cat # 8516S) were purchased from Cell Signaling. β Actin antibody was purchased from R&D Systems (Cat # MAB8929). Cyclin A antibody was purchased from Sigma (Cat # C4710).

 Colony Formation assay: 3226 and UM53 cells were seeded in 6 well dish (1000 cells/well) and treated with MRTX1133 and MRX849 respectively in combination with ITGB1 RNAi. Colonies were allowed to form for 10 days and stained with Crystal Violet.

Organoid Cell culture: The H2B-GFP labelled 519, 3226 and UM53 cells were seeded in 50 %

Matrigel basement layer (4000 cells/well) and allowed to form Organoids. Following the

treatment with MRTX1133 and MRTX849, the growth of organoids were monitored using live

cell imaging that determined the GFP area.

Mice and patient-derived xenografts: NSG and C57/BL6 were maintained at Roswell Park

Cancer Center animal care facilities. All animal care, drug treatments and sacrifice were

approved by the Roswell Park Cancer Center Institutional Animal Care and Use Committee

(IACUC) in accordance with the NIH guide for the care and use of laboratory animals. Mice

were subcutaneously implanted with early passage 828 and 3226 PDX tumor fragments. AKB6

xenografts were developed in both NSG and C57/BL6 strains by subcutaneously injecting

5X10^6 cells/mouse. The entire experimental cohort was comprised of both male and female

mice. The mice were always maintained in Doxycycline water to induce the oncogenic KRAS.

KC4568 xenografts were developed in the C57/BL6 strain by subcutaneously injecting 5X10^6

cells/mouse. Once the tumor reached 200 mm^3, mice were randomized in a non-blinded

manner, comprising, Vehicle (n=5), and MRTX1133 (30 mg/kg) (n-5), administered

intraperitoneally once per day for three weeks. MRTX1133 was dissolved in 50 mM citrate

buffer and 10% Captisol. Tumor growth was monitored every other day using digital calipers

and the tumor volume was calculated using the formula (length*widith*width)/2 as described

before [30]. Mice were sacrificed and the tumors were embedded in paraffin for further analysis.

Mice body weights were monitored regularly to evaluate any adverse effects. Mice that died

during the course of treatment were excluded from analysis. The animal

Transcriptome Analysis: RNA was extracted using Quiagen RNeasyplus kit and analyzed via

the RNA6000 Nano assay, and with the Agilent 2200 TapeStation (Agilent, CA, USA) for

determination of an RNA Integrity Number (RIN), and only the cases with RIN > 7.0 were

included in this study. cDNA synthesis was performed using random hexamers to obtain full-

length, strand-specific representation of nonribosomal RNA transcripts. Targeted RNA

sequencing libraries were prepared with the DriverMap Human Genome-Wide Gene Expression

Profiling Sample Prep Kit hDM18Kv3 (Cellecta Inc., CA, USA). A defining feature of the

DriverMap method is the application of pre-designed multiplex PCR primer sets targeted to

specific 50-75⬜bp regions of all known protein-coding genes. Notably, each target-specific

primer consists of a complementary sequence to specific mRNA targets plus a universal primer

binding site (anchor). Ligation of oligonucleotides via PCR amplification introduces adaptors

required for sequencing and sample-specific "barcodes" that flank the target sequence and are

inserted into standard Illumina adaptors to permit dual-index sequencing and deconvolution of

sample-specific reads using standard Illumina software.

Briefly, to mitigate primer dimer formation, anchor PCR was performed with an initial hot

start at 95°C for 5 min, followed by 15 cycles of (95°C – 0.5 min, 68°C – 1 min, 72°C – 1 min),

and ended with a final 10 min extension at 72°C. The reaction products were confirmed on an

agarose gel in triplicate to assess replicability. PCR products were then purified by SPRI

(Agentcourt, 1:1 sample: reagent ratio) and quantified with the Qubit fluorescence assay (Qubit

dsDNA HS Assay Kit, ThermoFisher Scientific, MA, USA). Target-enriched RNA-seq libraries

were analyzed on an Illumina NextSeq 500 sequencer using a NextSeq500/550 High Output v2

Kit (75 cycles) according to the standard manufacturer's protocol (Illumina, CA, USA).

For RNA-seq, alignment was performed using STAR v2.7.10b, and this pipeline outputs gene-

level read counts directly[33]. Transcript abundance estimates for each sample were performed

using Salmon, an expectation-maximization algorithm using the UCSC gene definitions. Raw

read counts for all samples were normalized using the "weighted" Trimmed Mean of M-values

(TMM) approach in the Bioconductor package EdgeR[34]. After trimming the data [5% for the A

values, log ratio 0.3 for the M values to a reference array (the library whose upper quartile is

closest to the mean upper quartile)], scaling factors for each sample were generated using the

calcNormFactors function. Genes with an average of less than ten read counts across all

samples were excluded from further analysis. Following data integration, systematic bias was

corrected using ComBat, as described previously[35]. DE gene (DEG) analysis was performed

with EdgeR and selected based on p-value and log2 fold change (log2FC), respectively.

CRISPR screening: Human Toronto KnockOut (TKO) CRISPR library-version 3, containing

70,948 unique sgRNAs, which targets 18,053 genes was packaged in lentivirus according to the

previously published protocol[36]. ASPC1 and UM53 cells were infected with the TKOv3 lentiviral

particles at an MOI ~0.3 and the positive clones were selected using puromycin to achieve a

mutant pool comprising at least 200-fold coverage of the library. The resulting mutant pool was

grown in the absence and presence of KRASi at the EC25 concentration for at least 8

passages. Genomic DNA was extracted and subjected to sequencing library construction by

amplifying the gRNA inserts using a 2-Step PCR as described in the previously published

study[36]. Resulting libraries were further sequenced on an Illumina's NexSeq platform. Fastq files

were first subjected to adapter removal using Trim Galore (v0.6.7,

https://github.com/FelixKrueger/TrimGalore). Next, the MAGeCK[37] pipeline was used to process

the adapter trimmed fastq files and the resultant count files are used as input to DrugZ[38] python

script for chemogenetic interactions identification. An in-house developed R script was used to

creating plots for results visualization.

10X Single cell RNA sequencing: Tumors excised from AKB6 xenografts that were treated

with vehicle (n=2) and MRTX1133 (n=2) were digested using Liberase (Sigma Cat #

05401020001) to get single cells and sent to RPCI's Genomics Shared Resources for

sequencing using 10X Chromium and Illumina's NovaSeq instruments. The raw fastq data was

processed using 10X's cellranger pipeline (v6.1.2). The resultant 'filtered_feature_bc_matrix.h5'

files for treatment and vehicle control, respectively, were used as inputs for a custom R (v4.2.0)

script which utilizes the Seruat[39] R package (v4.2.0) from CRAN. We filtered each dataset with

the following criteria: percent.mt < 10, nFeature_RNA between 200 and 4000,

percent.largest.gene < 20. With these parameters, we obtained 9307 and 7204 single cells in

the treated and control samples, respectively. Since the treated sample resulted more single

cells, we down-sampled it by randomly selecting the same number of cells (7204) as in the

control. The two datasets were then integrated by using the FindIntegrationAnchors and

IntegrateData function from the Seurat suite. After this step, the standard Seurat workflow was

followed to cluster the combined dataset. We identified 20 clusters and annotated the clusters

with ScType[40]

Digital Spatial Profiling (DSP): We used NanoStrings GeoMx instrument and the Mouse NGS

Whole Transcriptome Atlas RNA (version 1.0) kit for ROIs selection and mRNA library

preparation. We used PanCK and CD45 antibodies cocktail for staining of cancer cell and

stromal segments respectively. The resulting mRNA samples are subjected to sequencing using

Illumina's NavoSeq instrument. Raw fastq files along with the configuration file are used as input

to the geomxngspipeline command line tool running on a Linux server. We used NanoString's

GeomxTools   Bioconductor   package

(https://www.bioconductor.org/packages/release/bioc/html/GeomxTools.html)   for downstream

data analyses. These include QC & pre-processing, data normalization, unsupervised clustering

and differential gene expression. For QC, we used values for these parameters:

minSegmentReads = 1000, percentTrimmed = 80, percentStitched = 80, percentAligned = 80,

percentSaturation = 50, minNegativeCount = 2, maxNTCCount = 1000, minNuclei = 100,

minArea = 5000, minLOQ = 2. For segments passed these QC parameters, we utilized Q3

normalization and mixed effect modeling for identifying differentially expressed genes in the

PanCK+ segments between treated and control.

Pathway analysis: We used the Enrichr[41-43] webserver and MSigDB[44] Hallmark 2020 dataset

for all pathway analysis in this paper.

Multi-spectral immunofluorescence (mIF) staining: Formalin-fixed Paraffin-embedded

(FFPE) 4 µm sections were cut and placed on charged slides. The multispectral

immunofluorescent (mIF) staining process involved serial repetitions of the following for each

biomarker: epitope retrieval/stripping with ER1 (citrate buffer pH 6, AR996, Leica Biosystems)

or ER2 (Tris-EDTA buffer pH9, AR9640, Leica Biosystems), blocking buffer (AKOYA

Biosciences,), primary antibody, Opal Polymer HRP secondary antibody (AKOYA Biosciences),

Opal Fluorophore (AKOYA Biosciences). All AKOYA reagents used for mIF staining come as a

kit (NEL821001KT). Spectral DAPI (AKOYA Biosciences) was applied once slides were

removed from the BOND. They were cover slipped using an aqueous method and Diamond

antifade mounting medium (Invitrogen ThermoFisher). The mIF panel consisted of the following

antibodies, CD8 (EPR20305, Abcam, 570), Pan Keratin (Wide Spectrum cytokeratin, Abcam,

480), and DAPI. Slides were imaged on the PhenoImager HT® Automated Quantitative

Pathology Imaging System (AKOYA Biosciences). Further analysis of the slides was performed

using inForm® Software v2.6.0 (AKOYA Biosciences).

Acknowledgements: The author thank all members of the laboratory group and

colleagues in the discussion and preparation of the manuscript. The research was

supported by a grant to AKW and ESK from National Cancer Institute (NCI).

Author contributions:

Study concept and design: VK, CF, EA, PD, ESK and AKW

Acquisition of data: VK, CF, JW, YW, AD and HR

Analysis and interpretation of data: VK, CF, JW, YW, AD, HR, ESK and AKW

Study supervision: ESK and AKW.

Competing Interests: Mirati Therapeutics provided MRTX1133 for *in vivo* work. The

study was written in the absence of input from them.

References

1   Yao, W., Maitra, A. & Ying, H. Recent insights into the biology of pancreatic cancer.
EBioMedicine **53**, 102655, doi:10.1016/j.ebiom.2020.102655 (2020).
2   Kleeff, J. *et al.* Pancreatic cancer. *Nat Rev Dis Primers* **2**, 16022,
doi:10.1038/nrdp.2016.22 (2016).
3   Neoptolemos, J. P. *et al.* Therapeutic developments in pancreatic cancer: current and
future perspectives. *Nat Rev Gastroenterol Hepatol* **15**, 333-348, doi:10.1038/s41575-
018-0005-x (2018).
4   Stott, M. C., Oldfield, L., Hale, J., Costello, E. & Halloran, C. M. Recent advances in
understanding pancreatic cancer. *Fac Rev* **11**, 9, doi:10.12703/r/11-9 (2022).
5   Witkiewicz, A. K. *et al.* Whole Exome Sequencing of Pancreatic Cancer:Genetic Diversity,
Prognostic Features, and Potential Therapeutic Targets. *Nature Communication* (2014).
6   Bailey, P. *et al.* Genomic analyses identify molecular subtypes of pancreatic cancer.
Nature **531**, 47-52, doi:10.1038/nature16965 (2016).
7   Knudsen, E. S., O'Reilly, E. M., Brody, J. R. & Witkiewicz, A. K. Genetic Diversity of
Pancreatic Ductal Adenocarcinoma and Opportunities for Precision Medicine.
Gastroenterology, doi:10.1053/j.gastro.2015.08.056 (2015).
8   Infante, J. R. *et al.* A randomised, double-blind, placebo-controlled trial of trametinib, an
oral MEK inhibitor, in combination with gemcitabine for patients with untreated
metastatic adenocarcinoma of the pancreas. *Eur J Cancer* **50**, 2072-2081,
doi:10.1016/j.ejca.2014.04.024 (2014).
9   Huang, L., Guo, Z., Wang, F. & Fu, L. KRAS mutation: from undruggable to druggable in
cancer. *Signal Transduct Target Ther* **6**, 386, doi:10.1038/s41392-021-00780-4 (2021).
10   Waters, A. M. & Der, C. J. KRAS: The Critical Driver and Therapeutic Target for Pancreatic
Cancer. *Cold Spring Harb Perspect Med* **8**, doi:10.1101/cshperspect.a031435 (2018).
11   Nollmann, F. I. & Ruess, D. A. Targeting Mutant KRAS in Pancreatic Cancer: Futile or
Promising? *Biomedicines* **8**, doi:10.3390/biomedicines8080281 (2020).
12   Zeitouni, D., Pylayeva-Gupta, Y., Der, C. J. & Bryant, K. L. KRAS Mutant Pancreatic
Cancer: No Lone Path to an Effective Treatment. *Cancers (Basel)* **8**,
doi:10.3390/cancers8040045 (2016).
13   Muzumdar, M. D. *et al.* Survival of pancreatic cancer cells lacking KRAS function. *Nat*
Commun **8**, 1090, doi:10.1038/s41467-017-00942-5 (2017).
14   Kapoor, A. *et al.* Yap1 activation enables bypass of oncogenic Kras addiction in
pancreatic cancer. *Cell* **158**, 185-197, doi:10.1016/j.cell.2014.06.003 (2014).
15   Viale, A. *et al.* Oncogene ablation-resistant pancreatic cancer cells depend on
mitochondrial function. *Nature* **514**, 628-632, doi:10.1038/nature13611 (2014).
16   Collins, M. A. *et al.* Oncogenic Kras is required for both the initiation and maintenance of
pancreatic cancer in mice. *J Clin Invest* **122**, 639-653, doi:10.1172/JCI59227 (2012).
17   Shinkawa, T., Ohuchida, K. & Nakamura, M. Heterogeneity of Cancer-Associated
Fibroblasts and the Tumor Immune Microenvironment in Pancreatic Cancer. *Cancers*
(Basel) **14**, doi:10.3390/cancers14163994 (2022).
18   Hosein, A. N., Brekken, R. A. & Maitra, A. Pancreatic cancer stroma: an update on
therapeutic targeting strategies. *Nat Rev Gastroenterol Hepatol* **17**, 487-505,
doi:10.1038/s41575-020-0300-1 (2020).

19   Rhim, A. D. *et al.* Stromal elements act to restrain, rather than support, pancreatic
ductal adenocarcinoma. *Cancer Cell* **25**, 735-747, doi:10.1016/j.ccr.2014.04.021 (2014).
20   Olive, K. P. *et al.* Inhibition of Hedgehog signaling enhances delivery of chemotherapy in
a mouse model of pancreatic cancer. *Science* **324**, 1457-1461, doi:1171362 [pii]
10.1126/science.1171362 (2009).
21   Li, X. *et al.* Immune checkpoint blockade in pancreatic cancer: Trudging through the
immune desert. *Seminars in cancer biology* **86**, 14-27,
doi:10.1016/j.semcancer.2022.08.009 (2022).
22   Ischenko, I. *et al.* KRAS drives immune evasion in a genetic model of pancreatic cancer.
Nat Commun **12**, 1482, doi:10.1038/s41467-021-21736-w (2021).
23   Nakajima, E. C. *et al.* FDA Approval Summary: Sotorasib for KRAS G12C-Mutated
Metastatic NSCLC. *Clin Cancer Res* **28**, 1482-1486, doi:10.1158/1078-0432.CCR-21-3074
(2022).
24   Strickler, J. H. *et al.* Sotorasib in KRAS p.G12C-Mutated Advanced Pancreatic Cancer. *The
New England journal of medicine* **388**, 33-43, doi:10.1056/NEJMoa2208470 (2023).
25   Ning, W., Marti, T. M., Dorn, P. & Peng, R. W. Non-genetic adaptive resistance to
KRAS(G12C) inhibition: EMT is not the only culprit. *Frontiers in oncology* **12**, 1004669,
doi:10.3389/fonc.2022.1004669 (2022).
26   Xue, J. Y. *et al.* Rapid non-uniform adaptation to conformation-specific KRAS(G12C)
inhibition. *Nature* **577**, 421-425, doi:10.1038/s41586-019-1884-x (2020).
27   Witkiewicz, A. K. *et al.* Whole-exome sequencing of pancreatic cancer defines genetic
diversity and therapeutic targets. *Nat Commun* **6**, 6744, doi:10.1038/ncomms7744
(2015).
28   Wang, X. *et al.* Identification of MRTX1133, a Noncovalent, Potent, and Selective
KRAS(G12D) Inhibitor. *Journal of medicinal chemistry* **65**, 3123-3133,
doi:10.1021/acs.jmedchem.1c01688 (2022).
29   Hallin, J. *et al.* Anti-tumor efficacy of a potent and selective non-covalent KRAS(G12D)
inhibitor. *Nat Med* **28**, 2171-2182, doi:10.1038/s41591-022-02007-7 (2022).
30   Knudsen, E. S. *et al.* Cell cycle plasticity driven by MTOR signaling: integral resistance to
CDK4/6 inhibition in patient-derived models of pancreatic cancer. *Oncogene*,
doi:10.1038/s41388-018-0650-0 (2019).
31   Knudsen, E. S. *et al.* CDK/cyclin dependencies define extreme cancer cell-cycle
heterogeneity and collateral vulnerabilities. *Cell Rep* **38**, 110448,
doi:10.1016/j.celrep.2022.110448 (2022).
32   Kumarasamy, V., Ruiz, A., Nambiar, R., Witkiewicz, A. K. & Knudsen, E. S. Chemotherapy
impacts on the cellular response to CDK4/6 inhibition: distinct mechanisms of
interaction and efficacy in models of pancreatic cancer. *Oncogene* **39**, 1831-1845,
doi:10.1038/s41388-019-1102-1 (2020).
33   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,
doi:10.1093/bioinformatics/bts635 (2013).
34   Robinson, M. D. & Oshlack, A. A scaling normalization method for differential expression
analysis of RNA-seq data. *Genome Biol* **11**, R25, doi:10.1186/gb-2010-11-3-r25 (2010).

35   Zhang, Y., Parmigiani, G. & Johnson, W. E. ComBat-seq: batch effect adjustment for
639   RNA-seq count data. *NAR Genom Bioinform* **2**, lqaa078, doi:10.1093/nargab/lqaa078
640   (2020).
641   36   Hart, T. *et al.* Evaluation and Design of Genome-Wide CRISPR/SpCas9 Knockout Screens.
642   *G3 (Bethesda)* **7**, 2719-2727, doi:10.1534/g3.117.041277 (2017).
643   37   Li, W. *et al.* MAGeCK enables robust identification of essential genes from genome-scale
644   CRISPR/Cas9 knockout screens. *Genome Biol* **15**, 554, doi:10.1186/s13059-014-0554-4
645   (2014).
646   38   Colic, M. *et al.* Identifying chemogenetic interactions from CRISPR screens with drugZ.
647   *Genome Med* **11**, 52, doi:10.1186/s13073-019-0665-3 (2019).
648   39   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587
649   e3529, doi:10.1016/j.cell.2021.04.048 (2021).
650   40   Ianevski, A., Giri, A. K. & Aittokallio, T. Fully-automated and ultra-fast cell-type
651   identification using specific marker combinations from single-cell transcriptomic data.
652   *Nat Commun* **13**, 1246, doi:10.1038/s41467-022-28803-w (2022).
653   41   Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment
654   analysis tool. *BMC Bioinformatics* **14**, 128, doi:10.1186/1471-2105-14-128 (2013).
655   42   Kuleshov, M. V. *et al.* Enrichr: a comprehensive gene set enrichment analysis web server
656   2016 update. *Nucleic Acids Res* **44**, W90-97, doi:10.1093/nar/gkw377 (2016).
657   43   Xie, Z. *et al.* Gene Set Knowledge Discovery with Enrichr. *Curr Protoc* **1**, e90,
658   doi:10.1002/cpz1.90 (2021).
659   44   Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set
660   collection. *Cell Syst* **1**, 417-425, doi:10.1016/j.cels.2015.12.004 (2015).
661   45   Knudsen, E. S. *et al.* Pancreatic cancer cell lines as patient-derived avatars: genetic
662   characterisation and functional utility. *Gut* **67**, 508-520, doi:10.1136/gutjnl-2016-313133
663   (2018).
664   46   Zeng, H. *et al.* Genome-wide CRISPR screening reveals genetic modifiers of mutant EGFR
665   dependence in human NSCLC. *Elife* **8**, doi:10.7554/eLife.50223 (2019).
666   47   Cooper, J. & Giancotti, F. G. Integrin Signaling in Cancer: Mechanotransduction,
667   Stemness, Epithelial Plasticity, and Therapeutic Resistance. *Cancer Cell* **35**, 347-367,
668   doi:10.1016/j.ccell.2019.01.007 (2019).
669   48   Cai, X., Wang, K. C. & Meng, Z. Mechanoregulation of YAP and TAZ in Cellular
670   Homeostasis and Disease Progression. *Front Cell Dev Biol* **9**, 673599,
671   doi:10.3389/fcell.2021.673599 (2021).
672   49   Janes, M. R. *et al.* Targeting KRAS Mutant Cancers with a Covalent G12C-Specific
673   Inhibitor. *Cell* **172**, 578-589 e517, doi:10.1016/j.cell.2018.01.006 (2018).
674   50   Fujita-Sato, S. *et al.* Enhanced MET Translation and Signaling Sustains K-Ras-Driven
675   Proliferation under Anchorage-Independent Growth Conditions. *Cancer Res* **75**, 2851-
676   2862, doi:10.1158/0008-5472.CAN-14-1623 (2015).
677   51   Beatty, G. L. *et al.* CD40 agonists alter tumor stroma and show efficacy against
678   pancreatic carcinoma in mice and humans. *Science* **331**, 1612-1616,
679   doi:10.1126/science.1198443 (2011).

52   Dey, P. *et al.* Oncogenic KRAS-Driven Metabolic Reprogramming in Pancreatic Cancer
Cells Utilizes Cytokines from the Tumor Microenvironment. *Cancer Discov* **10**, 608-625,
doi:10.1158/2159-8290.CD-19-0297 (2020).

53   Knudsen, E. S. *et al.* Targeting dual signalling pathways in concert with immune
checkpoints for the treatment of pancreatic cancer. *Gut* **70**, 127-138,
doi:10.1136/gutjnl-2020-321000 (2021).

54   Kalbasi, A. *et al.* Uncoupling interferon signaling and antigen presentation to overcome
immunotherapy resistance due to JAK1 loss in melanoma. *Sci Transl Med* **12**,
doi:10.1126/scitranslmed.abb0152 (2020).

55   Paschen, A., Melero, I. & Ribas, A. Central Role of the Antigen-Presentation and
Interferon-γ Pathways in Resistance to Immune Checkpoint Blockade. *Annual Review of*
Cancer Biology **6**, 85-102, doi:10.1146/annurev-cancerbio-070220-111016 (2022).

56   Rice, C. M. *et al.* Tumour-elicited neutrophils engage mitochondrial metabolism to
circumvent nutrient limitations and maintain immune suppression. *Nat Commun* **9**,
5099, doi:10.1038/s41467-018-07505-2 (2018).

57   Rogers, T. & DeBerardinis, R. J. Metabolic Plasticity of Neutrophils: Relevance to
Pathogen Responses and Cancer. *Trends Cancer* **7**, 700-713,
doi:10.1016/j.trecan.2021.04.007 (2021).

58   Uprety, D. & Adjei, A. A. KRAS: From undruggable to a druggable Cancer Target. *Cancer*
Treat Rev **89**, 102070, doi:10.1016/j.ctrv.2020.102070 (2020).

59   Kemp, S. B. *et al.* Efficacy of a Small-Molecule Inhibitor of KrasG12D in
Immunocompetent Models of Pancreatic Cancer. *Cancer Discov* **13**, 298-311,
doi:10.1158/2159-8290.CD-22-1066 (2023).

60   Schram, A. M. *et al.* A phase Ib dose-escalation and expansion study of the oral MEK
inhibitor pimasertib and PI3K/MTOR inhibitor voxtalisib in patients with advanced solid
tumours. *Br J Cancer* **119**, 1471-1476, doi:10.1038/s41416-018-0322-4 (2018).

61   Ko, A. H. *et al.* A Multicenter, Open-Label Phase II Clinical Trial of Combined MEK plus
EGFR Inhibition for Chemotherapy-Refractory Advanced Pancreatic Adenocarcinoma.
Clin Cancer Res **22**, 61-68, doi:10.1158/1078-0432.CCR-15-0979 (2016).

62   Hallin, J. *et al.* The KRAS(G12C) Inhibitor MRTX849 Provides Insight toward Therapeutic
Susceptibility of KRAS-Mutant Cancers in Mouse Models and Patients. *Cancer Discov* **10**,
54-71, doi:10.1158/2159-8290.CD-19-1167 (2020).

63   Pellinen, T. *et al.* ITGB1-dependent upregulation of Caveolin-1 switches TGFbeta
signalling from tumour-suppressive to oncogenic in prostate cancer. *Sci Rep* **8**, 2338,
doi:10.1038/s41598-018-20161-2 (2018).

64   Tiriac, H., Plenker, D., Baker, L. A. & Tuveson, D. A. Organoid models for translational
pancreatic cancer research. *Curr Opin Genet Dev* **54**, 7-11,
doi:10.1016/j.gde.2019.02.003 (2019).

65   Kumarasamy, V. *et al.* RB loss determines selective resistance and novel vulnerabilities
in ER-positive breast cancer models. *Oncogene*, doi:10.1038/s41388-022-02362-2
(2022).

66   Li, X. & Wang, C. Y. From bulk, single-cell to spatial RNA sequencing. *Int J Oral Sci* **13**, 36,
doi:10.1038/s41368-021-00146-0 (2021).

67   Hu-Lieskovan, S. *et al.* Improved antitumor activity of immunotherapy with BRAF and
MEK inhibitors in BRAF(V600E) melanoma. *Sci Transl Med* **7**, 279ra241,
doi:10.1126/scitranslmed.aaa4691 (2015).
68   Jorgovanovic, D., Song, M., Wang, L. & Zhang, Y. Roles of IFN-gamma in tumor
progression and regression: a review. *Biomark Res* **8**, 49, doi:10.1186/s40364-020-
00228-x (2020).
69   Axelrod, M. L., Cook, R. S., Johnson, D. B. & Balko, J. M. Biological Consequences of
MHC-II Expression by Tumor Cells in Cancer. *Clin Cancer Res* **25**, 2392-2402,
doi:10.1158/1078-0432.CCR-18-3200 (2019).

733
Figure Legends

Figure 1 Cellular response to MRTX1133 in a panel of established and patient-derived

KRAS[G12D] **mutant PDAC cell lines.** (A) Live cell imaging using IncuCyte S3 on the indicated

cell lines treated with different concentrations of MRTX1133. Cell proliferation was determined

based on the GFP counts. Graph represents mean and SD from triplicates. Experiments were

done at two independent times (B) EC50 values of MRTX1133 on different PDAC cell lines

based on its impact on the cell proliferation. (C) Volcano plots indicating the differentially

expressed genes on the indicated cell lines treated with MRTX1133 (100 nM) up to 48 H based

on Transcriptome analysis. Blue: represents genes that are significantly downregulated

($\log_2$Fold change<-1.2, -$\log_{10}$pVal >2). Red represents genes that are significantly upregulated

($\log_2$Fold change>1.2, -$\log_{10}$pVal >2). (D) Pie charts on the indicated cell lines indicate the total

number of genes that are upregulated ($\log_2$Fold change>1.2, -$\log_{10}$pVal >2) and downregulated

($\log_2$Fold change<-1.2, -$\log_{10}$pVal >2) in the presence of MRTX1133 (100 nM). (E) Heatmap

depicting the GSEA hallmarks signatures that are differentially altered on different cell lines in

the presence of MRTX1133 (100 nM). (F) Heatmap depicting the differential expression of the

E2F target genes and the genes involved in MTOR pathway from different cell lines. (G)

Immunoblotting on the indicated cell lines treated with different concentrations MRTX1133 up to

48 H.

Figure 2 Genome wide CRISPR/CAS9 library screening in PDAC models in the presence and absence of KRAS inhibitor. (A) Schema illustrating the workflow of the CRISPR/CAS9 screening. (B) Volcano plot indicating the genes whose sgRNAs are differentially selected in the presence of MRTX1133 (6.25 nM) in ASPC1 cells. (C) ENRICHR analysis to determine the pathways that are associated with the negatively selected genes from ASPC1 cells. (D) DrugZ analysis in ASPC1 cells indicating the genes that were negatively selected during the MRTX1133 (6.25 nM) treatment. (E) Effect of MRTX1133 on the individual guides targeting, *EGFR, ERRF1, FOSL1 and KEAP1.* (F) Live cell imaging to monitor the proliferation of ASPC1 cells following EGFR deletion in the absence and presence of MRTX1133. Error bars represent mean and SD from triplicates. ***p<0.0001 as determined by two-way ANOVA. (G) DrugZ analysis in UM53 cells indicating the genes that were negatively selected during the MRTX849 (500 nM) treatment. (H) Effect of MRTX849 on the individual guides targeting, *ITGB1, TAZ and CAV1* in UM53 cells*.* (I) Effect of ITGB1 knockdown in ASPC1, 326, 1222 and UM53 cells in the absence and presence of MRTX1133 and MRTX849 at the indicated concentrations on cell proliferation. Graphs indicate mean SD from triplicates and the experiment was done at 2 independent times. (***p<0.0001 as determined by two-way ANOVA).

Figure 3 *In vivo* efficacy of MRTX1133 on PDX and mouse derived syngeneic PDAC models. (A) Effect of MRTX1133 on the growth of organoids derived from 519 PDAC cell line in the absence and presence of MRTX1133 (500 nM). Growth of organoids were determined based on the object area using IncuCyte S3. Error bars represent mean and SEM from two different experiments that were done in triplicates. (*** p<0.0001 as determined by two-way ANOVA). (B) Representative images of organoids at day 0 and day 5 in the absence and presence of MRTX1133 (500 nM). (C) Relative tumor volume of 3226 and 828 PDX following vehicle and MRTX1133 (30 mg/kg) treatment, administered intraperitonially (q.i.d) for the indicated number of days. (*** p<0.0001 as determined by two-way ANOVA). (D) Representative images of tumors excised from 828 PDX that were treated with vehicle and

MRTX1133 (30 mg/kg) (E) Relative BrdU incorporation in two different syngeneic models,

KC4568 and AKB6 that harbor G12D KRAS mutation following the treatment with different

concentrations of MRTX1133 up to 72 H. Mean and SD were calculated from triplicates and the

experiments were done at three independent times. (F) Effect of MRTX1133 (30 mg/kg) on the

relative tumor growth of xenografts derived from KC4568 and AKB6 in C57/BL6 mice. (***

p<0.0001 as determined by two-way ANOVA). (G) Representative images of H&E and

Masson's Trichrome staining on tumor tissues excised from KC4568 and AKB6 xenografts that

were treated with vehicle and MRTX1133 (30 mg/kg) in C57/BL6 mice.

Figure 4: Mechanistic investigation on *in vivo* effect of MRTX1133. (A) Single-cell clustering

of vehicle and MRTX1133-treated AKB6 tumors to selectively indicate Tumor cells, T-cells and

neutrophils. (B) Seurat heatmap depicting the expression of genes that distinguish two different

tumor population. (C) Bar graph indicating the tumor cell population from the vehicle and

MRTX1133 samples. (p value was determined by Fisher extract test) (D) Volcano plot to

represent the differentially expressed genes following MRTX1133 treatment from the tumor

population. (E) ENRICHR analysis to identify pathways that are associated with the upregulated

genes from the tumor population. (F) Representative tumor images from the tissues excised

from AKB6 xenografts that were treated with Vehicle (n=2) and MRTX1133 (n=4) and stained

with PACNK, CD45 and DAPI. ROIs were selected for DSP analysis. Representative image of

an ROI, demarcating the tumor and stromal population based on PANCK and CD45 staining

respectively and the masking was performed appropriately. (G) Volcano plots illustrate the

differentially expressed genes, on the PANCK+ cells following MRTX1133 treatment on the

AKB6 xenografts. Blue: represents downregulated genes (log_2Fold change<-0.5). Red

represents upregulated genes (log_2Fold change>0.5). Antigen presenting genes were

significantly upregulated and indicated. (H) GSEA analysis identified the upregulated genes

following MRTX1133 treatment were significantly enriched for Interferon γ signaling. (I) Heat

map depicting the upregulated genes across multiple ROIs from the vehicle treated tumor

tissues (n=2) and MRTX1133 treated tumor tissues (n=4). (J) GSEA analysis based on

transcriptome analysis on ASPC1 and 828 cell lines, treated with MRTX1133 (100 nM) identified

that the upregulated genes were significantly enriched for Interferon γ signaling. (K) Column

graph indicates the relative expression of *HLA-DMA, HLA-E* and *HLA-F* from ASPC1 and 828

cell lines treated with MRTX1133 based on transcriptome analysis. Error bar represents mean

and SD. (***p<0.0001 as determine student t-test).

Figure 5: Single cell sequencing analysis on the immune component from AKB6

xenografts. (A) Column graph indicating the neutrophils population in the absence and

presence of MRTX1133. (p value was determine by Fisher extract test). (B) Volcano plot

indicating the downregulated and upregulated genes on the neutrophils population following

MRTX1133 treatment. (C) Violin plots indicating the expression of indicated genes within the

neutrophil population in the presence and absence of MRTX1133. (D) Violin plots to indicate the

differential expression of T-cell markers between the vehicle and MRTX1133-treated groups. (E)

Column graphs indicating the different T-cell population that comprise Memory CD8$^+$ T cells,

Effector CD8$^+$ T cells, NKT cells and Memory CD4$^+$ T cells following MRTX1133 treatment. P

values were determined by Fisher extract test. (F) Seurat feature plots to illustrate the

differential expression of representative T-cell markers from the vehicle and MRTX1133-treated

groups. (G) Representative images from the multispectral staining on the AKB6 tumors

(PanKeratin, CD8 and DAPI). Column graph represent the fraction of CD8-positive cells from

the stroma and tumor following the treatment with MRTX1133. Mean and SEM were determined

from 6 ROIs from vehicle (n=3) and MRTX1133 (n=3) treated tumor tissues. (***p<0.0001 and

**p<0.01 as determine by student t-test). (H) *In vivo* effect of MRTX1133 on the tumor growth of

AKB6 xenografts in NSG mice. Tumor weights from the AKB6 xenografts derived from NSG and

C57/BL6 strains treated with vehicle and MRTX1133. Mean and SEM were shown. (*** p<0001

as determined by student t test).

Fig. 1

Fig. 2

Fig. 4

Fig. 5