Vessel Density Mapping of Cerebral Small Vessels on 3D High Resolution Black Blood MRI

Mona Sharifi Sarabi[1], Samantha J. Ma[2], Kay Jann[1], John M. Ringman[3], Danny J.J. Wang[1], Yonggang Shi[1*]

[1]USC Stevens Neuroimaging and Informatics Institute, University of Southern California, Los Angeles, CA, USA

[2]Siemens Healthcare USA, Los Angeles, CA, USA

[3]Department of Neurology, University of Southern California, Los Angeles, CA, USA

*Corresponding to:

Yonggang Shi, PhD

Mark & Mary Stevens Neuroimaging and Informatics Institute

Keck School of Medicine

University of Southern California (USC)

2025 Zonal Avenue, Los Angeles, CA 90033

Email: yshi@loni.usc.edu

Abstract

Cerebral small vessels are largely inaccessible to existing clinical in vivo imaging technologies. This study aims to present a novel analysis pipeline for vessel density mapping of cerebral small vessels from high-resolution 3D black-blood MRI at 3T. Twenty-eight subjects (10 under 35 years old, 18 over 60 years old) were imaged with the T1-weighted turbo spin-echo with variable flip angles (T1w TSE-VFA) sequence optimized for black-blood small vessel imaging with iso-0.5mm spatial resolution at 3T. Hessian-based vessel segmentation methods (Jerman, Frangi and Sato filter) were evaluated by vessel landmarks and manual annotation of lenticulostriate arteries (LSAs). Using optimized vessel segmentation, large vessel pruning and non-linear registration, a semiautomatic pipeline was proposed for quantification of small vessel density across brain regions and further for localized detection of small vessel changes across populations. Voxel-level statistics was performed to compare vessel density between two age groups. Additionally, local vessel density of aged subjects was correlated with their corresponding gross cognitive and executive function (EF) scores using Montreal Cognitive Assessment (MoCA) and EF composite scores compiled with Item Response Theory (IRT). Jerman filter showed better performance for vessel segmentation than Frangi and Sato filter which was employed in our pipeline. Cerebral small vessels on the order of a few hundred microns can be delineated using the proposed analysis pipeline on 3D black-blood MRI at 3T. The mean vessel density across brain regions was significantly higher in young subjects compared to aged subjects. In the aged subjects, localized vessel density was positively correlated with MoCA and IRT EF scores. The proposed pipeline is able to segment, quantify, and detect localized differences in vessel density of cerebral small vessels based on 3D high-resolution black-blood MRI. This framework may serve as a tool for localized detection of small vessel density changes in normal aging and cerebral small vessel disease.

Keywords: High-resolution black-blood MRI, Turbo spin-echo with variable flip angles (TSE VFA), vessel segmentation, vessel density, small vessel disease

1. Introduction

Cerebral small vessel disease (cSVD) is often associated with neurodegenerative diseases, and it can contribute to the progression of cognitive decline and physical disabilities (Breteler et al., 1994; Debette et al., 2010; Mayda and DeCarli, 2009; Rosenberg et al., 2016; Wardlaw et al., 2013a). Neuroimaging plays a key role in characterizing cSVD by identifying various features linked to cSVD including recent small subcortical infarcts, lacunes, white matter hyperintensities, enlarged perivascular spaces, microbleeds, and brain atrophy (Blair et al., 2017; Vemuri et al., 2022; Wardlaw et al., 2013b). However, when these features become visible or detectable in structural imaging, they are already manifestations of significant deterioration caused by cSVD. Recent advanced imaging techniques have facilitated the assessment of pathophysiology by determining microstructural tissue integrity (Baykara et al., 2016) as well as vascular function through the measurement of cerebral blood flow, arterial stiffness, cerebrovascular reactivity, and blood brain barrier leakage (Jann et al., 2021; Shao et al., 2019; Sur et al., 2020; Yan et al., 2016). Because the underlying mechanisms of cerebral microvascular changes in conditions related to cSVD remain poorly understood, there is an increasing desire to detect the early features of cSVD in order to prevent or mitigate downstream disease-related tissue changes.

Considering that most clinical features of vascular neurodegenrative diseases are consequences of cSVD, it is essential to be able to directly visualize and evaluate the cerebral small vessels themselves. To date, however, these small vessels including arterioles, capillaries and venules are largely inaccessible to existing clinical in vivo imaging technologies. We recently proposed and optimized high resolution 3D black-blood MRI with sub-millimeter (iso-0.5mm) spatial resolution using T1-weighted turbo spin-echo with variable flip angles (T1w-TSE-VFA) at field strengths of 3 and 7 Tesla (Ma et al., 2019). The long echo train of TSE-VFA offers two main advantages for visualizing small vessels: 1) adequate flow suppression by inherent dephasing of flowing signals (black-blood MRI); 2) sub-millimeter spatial resolution (isotropic ~0.5–0.6 mm) and near whole-brain coverage in a clinically acceptable time (<10min). We demonstrated that optimized TSE-VFA is suitable for visualizing lenticulostriate arteries (LSAs) on the order of a few hundred microns, which can be manually segmented for the quantification of their branch number, length, and tortuosity (Cho et al., 2008; Kang et al., 2009; Ma et al., 2019).

The purpose of this study was to present a semiautomatic post-processing pipeline (**Figure 1**) for segmenting cerebral small vessels and further for mapping of small vessel density based on

Figure 1. The proposed small vessel processing pipeline based on black-blood MRI. (1) Vessel segmentation: (a) Input high-resolution black-blood MRI to the pipeline. (b) Preprocessing steps such as skull-stripping, bias correction and denoising by non-local means (NLM) applied on (a). (c) 3D vessel enhancement, various multiscale Hessian-based methods (Frangi et al., 1998, Sato et al., 2000, Jerman et al., 2016) have been evaluated for enhancement of small vessels; Jerman filter outperformed other candidate methods. (d) Binarized vessel mask (BVM) was obtained by thresholding Jerman's vesselness map using the optimal threshold. (e) Spatial masking was defined to construct region of interest (ROI) and classify small cerebral vessels. (f) Vessel density image (VDI) was calculated by diffusing the content of (e) to the entire image volume. (2) Co-registration of high-resolution black-blood MRI (g) to MPRAGE (h) for each subject with affine registration. 12 landmark points were incorporated to improve the robustness of co-registration. (3) VDI normalization: Mapping VDIs (l) from each subject to the MNI152 atlas (j) by combining the affine registration in (2) and the nonlinear registration between the MPRAGE (k) and the MNI152 atlas (i).

high resolution (isotropic ~0.5mm) black-blood MRI with near whole-brain coverage at 3T. This

pipeline included pre-processing steps (skull-stripping, bias correction and non-local means

(NLM) denoising), followed by optimized vessel segmentation and non-linear registration for

quantification of small vessel density across brain regions. The developed pipeline was evaluated

by comparison of localized vessel density between young and aged groups, as well as association

of vessel density in aged subjects with their neurocognitive scores at voxel-level.

2. Methods

2.1 Subjects and Imaging

The study was approved by the institutional review board of the University of Southern

California (USC) and in accordance with the principles of Declaration of Helsinki. Black-blood

MRI and T1w structural images (MPRAGE) were collected from 28 participants including young

control (N =10, 4 female, 27±3.5 years, age range [22,33]), and aged control (N =18, 14 female,

69.4±6 years, age range [60, 81]) groups. The subjects were screened for neurologic or psychiatric

disorders and were generally healthy. In 12 of the 18 aged subjects, standard neuropsychological

evaluation was performed. Images were acquired using a Siemens 3T MAGNETOM Prisma
scanner with a 32-channel head coil. The "black blood" contrast was attained with an optimized
T1w-TSE-VFA sequence (Ma et al., 2019) with the following parameters: TR/TE = 1000/12ms,
turbo factor = 44, matrix size = 756x896, resolution = 0.51x0.51x0.64 mm^3 interpolated to
0.5x0.5x0.5 mm^3, 160 sagittal slices and 10% oversampling, GRAPPA factor = 2; total imaging
time = 8:39 min. This sequence provided nearly whole brain coverage except that the sagittal FOV
was 80mm, and bilateral saturation bands were applied to suppress the out-of-volume signals from
temporal regions. Image quality grading of the high-resolution 3D black-blood MRI was
performed by two expert graders. Grading criteria was based on a 4-point Likert scale, considering
imaging artifacts such as motion, blurring, ringing, and wraparound. MRI images included in this
study had an image quality grade and vascular quality grade of 3 or higher. This dataset will be
referred to as high-resolution black blood-3T-28 (**HRBB-3T-28**) in the rest of the manuscript.
Additionally, for method validation purposes, 15 black-blood images from the volunteer cohort of
our previous LSA study (Ma et al., 2019) were used in this study. This cohort included a total of
15 healthy volunteers, with 9 participants (7 male, 27.2 ±3.5 years) between 22-33 years old and
6 participants (1 male, 64 ±2.4 years) older than 61 years of age, herein referred to as the young
and aged group, respectively. All image volumes in this LSA dataset were separated into left and
right hemispheres. This dataset will be referred to as **LSA-30** in the rest of the manuscript.

2.2 Pre-Processing

To prepare the images for vessel segmentation, initially, high-resolution black-blood MRI and
MPRAGE images were skull-stripped using the Statistical Parametric Mapping (SPM) software.
Due to intensity inhomogeneity and relatively low signal-to-noise ratio (SNR), skull-stripped
images of high-resolution black-blood MRI were further pre-processed by bias correction and
denoising via a non-local means (NLM) filter (Buades et al., 2005a). Due to the high computational
cost of NLM in 3D (Buades et al., 2005b), block-wise NLM was adopted in this study which
allows computationally efficient filtering without compromising the denoising result (Coupé et al.,
2008).

In this study, the NLM search window size, which has an impact on computational time and
visual quality of the results, was experimentally set to $3 \times 3 \times 3$ in a block of size $32 \times 32 \times 32$
within each 3D black-blood MRI volume. To avoid excessive denoising and preserve small vessel
details, the smoothing parameter h, which depends on the noise level, was estimated as 0.03.

Figures S1(a) and **(b)** in the supplementary material show an example of high-resolution black

block MRI of a young control subject before and after pre-processing, respectively.

2.3 Vessel Enhancement in 3D

To overcome the undesired intensity variations of vessel images and suppress non-vascular

structures, numerous filter-based vessel enhancement methods have been proposed (Agam et al.

2005, Wiemker et al. 2013, Krissian et al. 2000, Frangi et al. 1998, Sato et al. 2000, Li et al. 2003,

Erdt et al. 2008, Zhou et al. 2007, Law and Chung 2010, Jiang et al. 2006, Zhang et al. 2010), and

their performances have been significantly improved recently. Among these methods, Hessian-

based filters which are based on second order image derivatives have shown great success in

various clinical applications such as segmentation of liver vessels in abdominal CTA (Luu et al.,

2015), lung vessels in thoracic CT (Rudyanto et al., 2014), and computer-aided detection of

cerebral aneurysms in 3D rotational angiography, MRA and CTA (Hentschke et al., 2014). In this

category of methods, one recent method called the *Jerman filter*, which is based on the ratio of

Hessian matrix eigenvalues, has been proposed for enhancement of both 2D and 3D vasculatures

(Jerman et al., 2016). The authors demonstrated that the Jerman filter could overcome some

deficiencies of previous Hessian-based enhancement algorithms such as poor and non-uniform

responses for vessels with varying contrast and sizes, bifurcations and aneurysms. The

enhancement function in the Jerman filter is defined as:

122
$$V = \begin{cases} 0 & if \ \lambda_2 \leq 0 \wedge \lambda_\rho \leq 0 \\ 1 & if \ \lambda_2 \geq \frac{\lambda_\rho}{2} > 0 \\ \lambda_2^2(\lambda_\rho - \lambda_2)\left(\frac{3}{\lambda_2+\lambda_\rho}\right)^3 & otherwise. \end{cases} \tag{1}$$

123 where, λ_2 and λ_3 are the eigenvalues of the Hessian matrix, and λ_ρ is the regularized value of λ_3

124 and is defined below to ensure the robustness of vesselness measure to low magnitudes of

125 $\lambda_2 \ and \ \lambda_3$:

126
$$\lambda_{\rho(s)} = \begin{cases} \lambda_3 & if \lambda_3 > \tau \ max_x \ \lambda_3(x,s) \\ \tau \ max_x \ \lambda_3(x,s) & if 0 < \lambda_3 \leq \tau \ max_x \ \lambda_3(x,s) \\ 0 & otherwise. \end{cases} \tag{2}$$

where, τ is a cutoff threshold between zero and one. Higher value of τ increases the difference

between λ_2 and λ_3 magnitudes for low contrast structures. The proposed enhancement function's

response values range from 0 to 1.

In this study, according to the vascular anatomy of the brain and the study cohort that includes

aged control subjects with thin, tortuous and low contrast vessels, an ideal enhancement function

should exhibit robust, uniform, and high response in the presence of: i) varying noise and contrast;

ii) varying vascular morphology (e.g., multiscale vessels especially small vasculature, vessels with

different cross-sections (circular-elliptical)); iii) pathology (e.g., vessels thinning, high tortuosity

or vessel bends). To meet these expectations, the Jerman filter appears to be a proper choice. To

help justify our choice, the performance of the Jerman method will be compared with two widely

used enhancement functions: the Frangi (Frangi et al., 1998) and Sato (Sato et al., 2000) method.

We will present the comparison results among the candidate methods in **Section** 3.1.2.

Moreover, in multiscale Hessian-based methods, vessels' scale range and the threshold

value for binarizing the vesselness response are two important parameters to estimate. Inaccurate

assessment of these parameters will result in over or under segmentation of the vascular network

of interest. Therefore, to quantitatively evaluate the choice of parameters and compare the

Figure 2. Dataset for method validation. (1) Vessel and background landmark annotation on a selected sagittal slice of two representative subjects, where (A-1) is a healthy 29-year-old male, and (B-1) is a healthy 64-year-old female. Magnified view of vessel landmarks (red) and background points (green) of the two subjects are shown in (A-2) and (B-2), respectively. (2) Manual lenticulostriate artery (LSA) annotation by experts using ITK-SNAP. For an axial (C-1), coronal (D-1), and sagittal (E-1) slice of a Black-blood MRI, manual delineated labels are on the axial (C-2), coronal (D-2) and sagittal (E-2) view are plotted. Additionally, minimum intensity projections of high-resolution 3T MRI of LSAs for a healthy 25-year-old male (F-1), and a healthy 67-year-old female (F-2) are shown together with their matching 3D rendering of LSA labels in (G-1) and (G-2), respectively.

performance of the candidate methods, two sets of ground truth data were used. 1) manually delineated landmarks of cerebral small vessels from high-resolution black-blood MRIs (**Figure 2** (1)) that serve as the validation dataset for parameter optimization. This validation cohort was a subset of the HRBB-3T-28 subjects, comprised of five young controls and five aged control subjects. 2) 3D annotation of LSAs on unilateral LSA images of the LSA-30 dataset from the left and right hemispheres of 15 subjects (**Figure 2** (2)) were utilized as an independent testing dataset to further quantify the performance of the candidate vessel enhancement methods with optimized parameters estimated from the validation cohort. In the validation cohort, for each high-resolution black-blood MRI we annotated 2000 vascular and 2000 non-vascular landmarks (**Figure** 2) (drawn by experts with six years of experience in vascular image analysis). Vessel landmarks were mostly placed at the center and boundary of small and low contrast vasculature to increase the sensitivity of our evaluation in small vessel detection. The non-vascular landmarks were placed densely in the vicinity of the vessel's boundary and CSF region where over segmentation and false positive detection are most likely to occur. Representative examples of vessel and background landmarks from a young and aged control subjects are shown in (**Figure 2** (1) A-1, A-2), and (**Figure 2** (1) B-1, B-2), respectively.

To evaluate the segmentation accuracy of the candidate methods, the vessel segmentation was obtained by thresholding the vesselness map generated by each method, where the threshold value was carefully estimated by the Receiver Operating Characteristic (ROC) analysis using vessel landmarks of the validation dataset. Afterwards, the candidate methods with optimized parameters were tested on the independent LSA-30 dataset, where the LSAs were delineated manually by an experienced neuroradiologist. For an objective assessment of the segmentation performance, the following metrics were employed:

- Sensitivity (SEN) = TP/ (TP + FN).
- Specificity (SPE) = TN/ (TN + FP).
- Accuracy (ACC) = (TP + TN)/ (TP + TN + FP + FN).
- Average Hausdorff distance (AVD)

where TP is true positive, FP is false positive, TN is true negative, and FN is false negative. The AVD metric is a widely used performance measure for calculating the distance between two point sets. In medical image segmentation, AVD is utilized to compare the distance between ground-

Figure 3. Spatial distribution of arteries, veins, and dural sinuses in high-resolution black-blood MRI. (a) Sagittal, coronal, and axial slices with overlayed vessel annotations. (b) 3D-rendering of vascular networks in (a); in (B-1, B-2) vessels are color coded to show vascular classification, where red-colored vessels are small arteries and veins, green-colored vessels are dural sinuses, and yellow-colored vessels are MCAs. To obtain the final masked small vessels (B-3, B-4), dural sinuses and brainstem and cerebellum regions ((1) and (2) highlighted in B-1) were excluded.

truth and segmentation results, hence, smaller values indicate better performances (Taha and

Hanbury, 2015). In this study, the AVD metric was calculated only for LSA segmentation results

since we have complete annotation of these vessels.

2.4 Vessel Classification using Anatomical Masking

The vessels of interest in this study were small artery, arterioles, venules and small veins which

were detected by Hessian-based methods applied on skull-stripped high-resolution black-blood

MRI (0.5mm isotropic). The optimized scale (vessel radii) range for the study cohort was

empirically estimated as [0.1-0.4] voxel (i.e., 50-200um) based on ROC analysis as described in

section 2.3. To obtain the small vessels' mask, the resultant vascular structure (or BVM) of skull-

stripped high-resolution black-blood MRI was classified. More precisely, apart from a complete

small cerebrovascular network, the final BVM included full or partial detection of superficial

veins, dural sinuses (include superior sagittal sinus, straight sinus, transverse sinus, etc.) and

middle cerebral artery (MCA) (**Figure 3** (a)). To separate these vessels, a region of interest (ROI)

was defined via anatomical regions of MPRAGE images using the Freesurfer ASEG map. As

Figure 4. Representative examples of vessel density image (VDI) from two control subjects. (a) 3D-rendering of the BVM of a young control subject (A-1, A-2). (A-3) Corresponding VDI of the BVM in (A-1, A-2) before (A-3) and after normalization (A-4). (b) 3D-rendering of the BVM of an aged control subject (B-1, B-2). (B-3) Corresponding VDI of the BVM in (B-1, B-2) before (B-3) and after normalization (B-4).

shown in **Figure 3**, dural sinuses form the terminal of superficial venous system and distribute

without closely accompanying arteries (Rhoton, 2002). Therefore, to isolate dural sinuses,

superficial arteries and veins, a tentative mask (V_{ten}) was generated by morphological erosion of

the brain mask (consist of WM, GM, and CSF). Afterwards, to exclude regions with larger arteries

and highly variable small vessel detection due to low SNR or imaging artifacts, the final ROI

(**Figure 1**(e)) was constructed by eliminating cerebellum, brainstem, optic-chiasm and ventral

regions from V_{ten}. The ROI mask was then transformed to the black-blood image space using the

affine transformation to the MPRAGE image as will be described in **Section** 2.6.

2.5 Vessel Density Calculation

Vessel density images (VDI) were calculated to allow localized comparison of vascular

changes across individuals. This step is essential since the spreading pattern of small vessels in

high resolution black-blood MRI varies among different subjects, which makes the direct

comparison of cerebral small vessels at voxel-level a difficult task. To overcome this challenge,

we computed VDI of each subject by diffusing the content of BVM to the entire image volume.

More precisely, VDI was a smooth vessel map obtained by a convolution of the BVM with a 3D

kernel. The window size of the kernel was empirically set as [30 30 30] voxels for all subjects.

Representative VDI examples of a young and aged control subject are shown in **Figure 4**. In
the VDI representation, red regions denote high vessel density associated with dense vessel
distribution. Blue regions indicate lower vessel density. According to the VDI representation, we
can observe sparser vessel distribution in the aged subject (**Figure 4**(b)) and a denser vessel
network in the young subject especially in frontal regions **(Figure 4**(a)).

2.6 Mapping Vessel Density Across Subjects

Due to anatomical variation, different brain size and MRIs' orientations among subjects, it is
required to normalize the subjects' VDIs into common space to enable cross-subject comparison
at voxel-level. To this end, we proposed a two-step registration method (supplement **Figure S2**).
In the first step, to robustly co-register skull-stripped black-blood MRI and MPRAGE image pairs,
a 3D affine registration with 12 landmark points was performed (supplement **Figures S2**(a), and
(b)) using the Elastix software (Klein et al., 2010). The landmarks were manually selected from
the cerebrum, cerebellum, and brainstem regions to guide the co-registration between the two 3D
volumes with different field of views, where the black-blood MRI has near-whole brain coverage
while the MPRAGE scan has full brain coverage. In the second step, we computed the nonlinear
warp between the MNI152 Atlas (common space in this study) and the MPRAGE image
(supplement **Figures S2** (c), and (e)). By combining the affine and nonlinear warp, we obtained
the final transformation from the black-blood MRI to the MNI152 space.

To pool the VDIs from all subjects into the MNI152 space for statistical analysis, VDI was
first reversely transformed to MPRAGE space using previously computed affine transformation
(**Figure S2** (Step-1)), and then non-linearly transformed to the MNI Atlas using a B-spline
transformation (**Figure S2** (Step-2)). After non-linear registration, all VDIs were normalized and
transformed into the common space. Then, two-sample t-test was computed to compare VDIs of
the two groups at voxel-level and statistical threshold was set to $p<0.05$ two-tailed (uncorrected).

2.7 Regression Analysis between Vessel Density and Cognitive Scores

A prominent behavioral phenotype of cSVD is early executive dysfunction manifested by
impaired capacity to use complex information, to formulate strategies, and to exercise self-control,
with less pronounced episodic memory deficits compared to AD patients (Wallin et al., 2018).
Therefore, we used a composite score of executive function (EF) developed using IRT (Staffaroni
et al., 2020a) which offers advantages such as better reliability, fewer statistical comparisons, and
improved power to detect longitudinal change with smaller sample sizes (Gibbons et al., 2012;

Staffaroni et al., 2020b). IRT EF composite scores were calculated for the 12 aged subjects (age

range = [62, 81], Age mean±SD (years) = 68.2±7) who underwent neuropsychological assessment.

A correlation analysis was performed at each voxel between vessel density image and

neurocognitive scores including IRT EF composite score, IRT-Z EF score normalized by age,

gender and education, and MoCA score which reflects gross cognition. The resultant correlation

map was thresholded at $p < 0.05$ (two-tailed, uncorrected) to detect the voxels with a significant

association between vessel density and cognition scores.

3. Results

3.1 Validation Results of in vivo MRI

3.1.1 *Scale and threshold parameters estimation*

The scale range of the candidate hessian-based methods was estimated as [0.1-0.4] with a

step-size of 0.1 based on the vascular anatomy within ROI of the study cohort and was verified by

the validation cohort with vessel landmarks. The optimal threshold for each filter (e.g., Frangi,

Sato, Jerman) was determined by a point on the ROC curve closet to the optimal classifier (i.e.,

upper left corner of ROC plot). To robustly estimate this threshold, we used bootstrapped samples

of the validation cohort with 1000 iterations and performed ROC analysis, where the scale range

was fixed to the estimated value range of [0.1-0.4] voxels and the search window for the threshold

value was defined as [0.001-0.2] with the step size of 0.001.

3.1.2 *Performance evaluation using vessel landmarks and LSAs*

Evaluation and comparison of the multiscale filters with three enhancement functions:

Frangi (Frangi et al., 1998), Sato (Sato et al., 2000) and Jerman (Jerman et al., 2016) were

performed based on the annotated vessel landmarks of high-resolution black-blood MRIs (**Figure**

5) and tested using manual segmentation of LSAs (**Figure 6**), respectively. For all three methods,

Figure 5. Performance comparison of Hessian-based methods for small cerebral vessel segmentation. (Row-1) 3D-rendering result of a young control subject (Female, 25 years old) by (a) Jerman, (b) Frangi, (c) Sato filter with optimized parameters. (Row-2) Magnified view of the annotated white box in Row-1. (Row-3) 3D-rendering result of an aged control subject (Female, 70 years old) by (a) Jerman, (b) Frangi, (c) Sato filter with optimized parameters. (Row-4) Magnified view of the annotated white box in Row-2. Overall, Jerman filter has more uniform and continuous vessel segmentation result for both young and aged subjects.

the scale range of multiscale Hessian-based methods was set to the target range of [0.1-0.4].

Utilizing the ROC analysis, the optimal threshold value was identified as 0.072, 0.005 and 0.15

for Jerman, Frangi and Sato's filters, respectively. The three different thresholds for the candidate

methods were due to the difference in their enhancement functions that results in vesselness maps

with various range of values. We expect the winning method to generate more robust and

generalizable vessel segmentation for all small vessels including LSAs with the same parameter

setting.

Figure 6 Performance comparison of Hessian-based methods for LSA segmentation of the same subjects in Figure 5. 3D-rendering of the LSA manual segmentation (column-1), and the segmentation result of the young (Row-1) and aged (Row-3) subject by (a) Jerman, (b) Frangi, (c) Sato filters with the same parameters as in Figure 5. (Row-2, Row-4) Corresponding error map of the segmentation results in (Row-1) and (Row-3), respectively.

Qualitative comparison of ROI vessels using the candidate methods of a young and aged

subject is demonstrated in **Figure 5**. The results show that Jerman filter (**Figure 5**(a)) generated

more robust, uniform and continuous vessel segmentation results compared to the other methods.

Moreover, Frangi results had smaller, disconnected components in the vascular network

segmentation of both young (**Figure 5**(b)(B-1) and aged (**Figure 5**(b)(B-2)) subjects. With Sato's

method, the resultant BVM had more discontinuity compared to Jerman's and Frangi's methods.

For further assessment of the candidate methods, the output of LSA's segmentation of the same

young and aged subjects are demonstrated in **Figure 6**. The ground-truth of LSA delineation for

both subjects are illustrated in **Figure 6**(Column-1). Jerman's (**Figure 6**(a)) and Frangi's (**Figure 6**(b)) methods had comparable results for LSA segmentation. The error map visualization of both methods shows that most of the vessels were detected correctly for the representative subjects (**Figure 6**(A-2, B-2, A-4, B-4). Jerman's filter demonstrated less discontinuity compared to Frangi's method in the young subject (white arrows in **Figure 6**(A-1, B-1)). Sato's filter (**Figure 6**(c)) demonstrated lower detection rate compared to the other two methods mainly at the distal portion of the LSAs (white arrows in **Figure 6**(C-2, C-4)).

Quantitative results of the three methods based on the vessel landmarks of the validation dataset and LSA annotation of LSA-30 dataset are provided in **Table 1** and **Table 2**, respectively. According to **Table 1**, Jerman's method showed the best performance in all three metrics including the AUC of ROC curves, sensitivity (SE) and specificity (SP). Table 2 of LSA validation results also shows that Jerman's method outperformed the other two methods based on the four metrics including AUC, SE, SP and AVD.

Method	ACCU	SE	SP
Jerman	0.999	**0.873**	0.998
Frangi	0.994	0.844	0.994
Sato	0.995	0.749	0.995

Table 1. Summary of evaluation metrics for small vessel segmentation using vessel landmarks of validation dataset equally sampled from young control, aged control, and aged subjects with vascular risk factors.

Method	AUC	SE	SP	AVD
Jerman	**0.83**	**0.66**	0.99	**8.52**
Frangi	0.81	0.63	0.99	10.58
Sato	0.73	0.46	0.99	18.64

Table 2. Summary of evaluation metrics for LSA segmentation using manual annotation of young and aged subjects.

Figure 7 3D visualization of small vessels segmentation using Jerman method with optimized parameters. (a) young control, (b) aged control groups (second row) have sparser vessel network compared to young group (first row).

3.2 Qualitative Results in Young and Aged Subjects

Representative examples of 3D small vessel segmentation of high-resolution black-blood MRI using Jerman filter with optimized parameters are demonstrated in **Figure 7**. Four cases were selected from the young control (**Figure 7**(a)) and aged control (**Figure 7**(b)) group respectively. Overall, by qualitative comparison we can see that young control subjects had significantly higher vessel density compared to the aged group.

3.3 Quantitative Results

3.3.1 Global Mean Vessel Density

Figure 8 illustrates the comparison of mean small vessel density between the YC and AC groups. To account for variation of brain size across different individuals, small vessels' volume was normalized by the ROI size. More precisely, normalized vessel density measure of each subject equals to (small vessels volume/ ROI volume), where ROI is defined in **Section** 2.4. As shown in **Figure 8**, the result of two-way student t-test

Figure 8 Comparison of mean vessel density of small cerebral vessels in young (age 22-33 years) and aged (age > 60 years) groups. * indicates significance $p < 0.05$.

311 demonstrated that the YC group had significantly higher vessel density compared to the aged group
312 (p<0.05).
313

Figure 9. Areas with significant (p<0.05, uncorr.) higher vessel density in young controls (YC (N=10)) as compared to aged controls (AC (N=18)).

3.3.2   Localized Vessel Density Analysis

3.3.2.1 Group-wise Analysis of Age-dependent Vessel Density Changes in Normal Brains

Figure 9 demonstrates the quantitative results of voxelwise comparison between the young and
aged groups in the near-whole brain volume, where we projected the t-value maps onto a cortical
surface in the MNI152 space for better visualization. We observed widespread significant
reductions in vessel density in the aged group as compared to the young group. These regions
mainly include medial, superior and middle frontal gyrus, precentral gyrus, precuneus and cuneus.
321

3.3.2.2 Regression Analysis of Vessel Density Changes and Cognitive Scores

The normalized VDIs of twelve aged control subjects (age range = [62, 81], mean±SD (yrs) =
68.2±7) who underwent neuropsychological assessment were correlated with their cognitive and
executive function scores (such as IRT EF, IRT-ZScore of EF, and MoCA) and the resultant
correlation maps are shown in **Figure 10.** Consistent Associations across all cognitive scores were
observed in posterior cingulate cortex (PCC), dorsolateral prefrontal cortex (dlPFC) as well as in
medial temporal lobe.

Figure 10. Localization of significantly associations between VDI of AC (N=12) subjects and their corresponding IRT score, IRT-ZScore score and MoCA score.

4. Discussion

In this study, we developed, optimized and validated a comprehensive 3D analysis framework

for localized mapping of small vessel based on high resolution (isotropic ~0.5mm) black-blood

MRI with near whole-brain coverage of young and aged subjects at standard clinical magnetic

field strength of 3T. In our experimental results, we evaluated the performance of the proposed

framework in studying the small vessel density changes in normal aging, and in correlation with

cognitive scores. Overall, we observed a decreasing pattern in the small vessel density by aging,

and a positive correlation between small vessels density and cognitive scores such as IRT EF and

MoCA scores.

4.1 Clinical Value of Whole Brain Small Vessel Density Mapping

To date, limited in vivo techniques are available to directly assess cerebral small vessels across

the whole brain. Digital subtraction angiography (DSA), x-ray computed tomography

angiography, and time-of-flight MR angiography have been applied to observe large cerebral

vessels in clinical populations (Gotoh et al., 2012; Kammerer et al., 2017; Wardlaw et al., 2013b),

but the sensitivity to small vessels such as the perforating and pial arteries is moderate at best and

difficult to characterize and quantify at 3T. With the proposed 3D multimodal framework for
analyzing high-resolution black-blood MRI, the automated segmentation enables both
visualization and quantitative assessment of cerebral small vessels.

4.2 Segmentation and Quantification

This study was inspired by our previous work (Sarabi et al., 2020), where we developed a
3D retinal vessel density mapping approach for optical coherence tomography angiography
(OCTA) and demonstrated its efficacy in localized detection of microvascular changes and
monitoring retinal disease progression in different clinical applications such as normal aging and
diabetic retinopathy (DR). Despite some similarities between the two studies, with respect to two-
module framework design for fine-scale vessel segmentation and non-linear registration, there are
still some fundamental differences in the method development. 3D cerebral vascular mapping in
high resolution black-blood MRI images is more challenging compared to retinal vessels mapping
due to relatively poor vessel contrast mainly in cerebrospinal fluid (CSF) regions, varying degrees
of noise, inhomogeneous backgrounds, anatomical variations, and partial view of the brain in this
study. To tackle these challenges, initially high-resolution black-blood MRIs underwent several
preprocessing steps such as skull-stripping, bias correction and NLM denoising. NLM was
selected for denoising since previous works (Chen et al. 2011, Zhang et al. 2014) demonstrated
that this process preserves anatomical details while suppressing the noise. Since the
microvasculature involved in cSVD includes small arteries/arterioles (~10µm-1mm), capillaries
(<10µm), and venules (~10-50µm) (Charidimou et al., 2016), we employed multi-scale Hessian-
based method for the purpose of vessel segmentation. More precisely, multiple popular Hessian-
based methods (Frangi et al., 1998, Sato et al., 2000, Jerman et al., 2016) have been evaluated for
the enhancement of small vessels using clinical data. Our evaluation results demonstrated that the
Jerman filter was more robust to noise compared to other candidate methods, had higher and more
uniform responses for small vessels not only at vessel center but also at vessel periphery. To
robustly detect the small vessels, validation experiments were designed to determine the optimal
scale and threshold value via ROC analysis using a validation dataset with annotated vessel
landmarks. For localized small vessel density analysis across populations, first, a vessel density
image (VDI) was calculated by diffusing the content of the vessel mask to the entire image volume.
After that, we implemented a non-linear registration framework to pool the VDIs from all subjects
into the common space (MNI-152 Atlas). To this end, initially skull-stripped black-blood MRI and

MPRAGE image pairs were robustly co-registered using a 3D-Affine registration with 12
landmark points. Subsequently, a non-linear registration was performed between MPRAGE of
each subject and MNI-152 Atlas using B-Spline method. By utilizing these two deformation fields,
all VDIs were normalized to MNI-152 Atlas. Thus, together with the careful preprocessing of the
image data, the localized mapping of vessel density in different subjects and patient groups enabled
the analysis of region-specific changes across populations.

4.3 Aging and Association with Neurocognitive Scores

Our approach of vessel density mapping revealed decreasing small vessel density with aging which
is consistent with literature (Sadoun and Reed, 2003; Watanabe et al., 2020). Age-related cerebral
microvascular changes have been visualized using ultrasound-based microscopy in animals
(Lowerison et al., 2022), while retinal vessel density can be quantified using OCTA in humans
which was correlated with cognitive scores such as MoCA (Ashimatey et al., 2020; Sarabi et al.,
2020). Laser scanning confocal 3D immunofluorescent microscopy of young and aging vascular
beds in multiple mice and human organs, including brain tissue, has also demonstrated an age-
dependent decline in vessel density (Chen et al., 2021). To the best of our knowledge, this is the
first in vivo study on aging effects on cerebral small vessel density using MRI. We also explored
associations between localized vessel density and neurocognitive scores including IRT EF and
MoCA. IRT EF scores were chosen since early executive dysfunction has been shown to be a
prominent behavioral phenotype of cSVD (Wallin et al., 2018). The IRT-based score used in the
present study is sensitive to impairments in mild cognitive impairment (MCI), Alzheimer's disease
(AD), and frontotemporal dementia (Staffaroni et al., 2020b) and is associated with cerebral blood
flow of middle cerebral artery perforator territory (Jann et al., 2021). In the present study, we found
significant positive associations between regional vessel density and IRT EF scores in areas
associated with executive functions such as dlPFC and PCC. In addition, we found a similar pattern
of brain areas with significant positive associations between regional vessel density and MoCA
scores (Nissim et al., 2016; Yokosawa et al., 2020). These promising data suggest that the proposed
analysis framework may be useful as an early screening tool for cSVD to gauge overall health of
cerebral small vessels. Furthermore, it could serve as a biomarker to monitor disease progression
and/or response to interventions for vascular dementia.

4.4 Limitations of the Study

In a few subjects, near-whole brain coverage instead of full FOV caused wraparound of ear

tissue signal into the image at the brainstem. Saturation bands should be placed more carefully to

saturate the signal of tissue outside the FOV. Such high-resolution imaging is susceptible to motion

artifacts that cause blurring or ghosting in the images. Motion was minimized by padding the

subjects in the head coil and applying paper tape across the forehead skin for tactile feedback.

Motion compensated 3D TSE-VFA sequences may provide a robust method for small vessel

visualization and quantification (Hu et al., 2021). As a proof-of-concept study, we only included

a small cohort of young and aged subjects, therefore no correction of multiple comparisons was

performed for statistical analysis.

One potential confounding factor for segmenting small vessels on black-blood MRI is

perivascular space (PVS) which appears gray on T1w TSE-VFA images. We have compared LSA

and PVS in T1w and T2w TSE-VFA respectively, as shown in Figure S7 of (Ma et al., 2019), and

the locations of LSAs did not match those of PVS in the basal ganglia area. In contrast to PVS that

is primarily observed in white matter, the majority of segmented superficial small vessels are in

gray matter, as shown in **Figure 3**. The relationship between small vessels and PVS in terms of

their spatial distribution and quantity need to be evaluated in future studies.

5. Conclusions

We presented and evaluated a novel framework for segmenting and mapping brain small

vessels from high-resolution black-blood images acquired at 3T. Using filter-based segmentations

and non-linear registration, 3D mapping, and quantification of brain small vessels was

demonstrated to be feasible. This framework may serve as a promising tool for localized detection

of vessel density changes in patients with neurovascular diseases and in understanding the

correlation between vessel density and cognitive scores.

ACKNOWLEDGEMENT

This work was supported by National Institute of Health (NIH) grants UF1-NS100614, R01-

NS114382, R01-EB032169, R01-EB028297, R01-EB022744, RF1-AG064584, P30AG066530,

and P41-EB015922.

References:

Ashimatey, B.S., D'Orazio, L.M., Ma, S.J., Jann, K., Jiang, X., Lu, H., Wang, D.J.J., Ringman, J.M., Kashani, A.H., 2020. Lower retinal capillary density in minimal cognitive impairment among older Latinx adults. Alzheimers Dement (Amst) 12, e12071.

Baykara, E., Gesierich, B., Adam, R., Tuladhar, A.M., Biesbroek, J.M., Koek, H.L., Ropele, S., Jouvent, E., Initiative, A.s.D.N., Chabriat, H., 2016. A novel imaging marker for small vessel disease based on skeletonization of white matter tracts and diffusion histograms. Annals of Neurology 80, 581-592.

Blair, G.W., Hernandez, M.V., Thrippleton, M.J., Doubal, F.N., Wardlaw, J.M., 2017. Advanced neuroimaging of cerebral small vessel disease. Current treatment options in cardiovascular medicine 19, 1-17.

Breteler, M., van Amerongen, N.M., van Swieten, J.C., Claus, J.J., Grobbee, D.E., Van Gijn, J., Hofman, A., Van Harskamp, F., 1994. Cognitive correlates of ventricular enlargement and cerebral white matter lesions on magnetic resonance imaging. The Rotterdam Study. Stroke 25, 1109-1115.

Buades, A., Coll, B., Morel, J.-M., 2005a. A non-local algorithm for image denoising. 2005 IEEE Computer Society Conference on Computer Vision and Pattern Recognition (CVPR'05). IEEE, pp. 60-65.

Buades, A., Coll, B., Morel, J.-M., 2005b. A review of image denoising algorithms, with a new one. Multiscale modeling & simulation 4, 490-530.

Charidimou, A., Pantoni, L., Love, S., 2016. The concept of sporadic cerebral small vessel disease: A road map on key definitions and current concepts. Int J Stroke 11, 6-18.

Chen, J., Sivan, U., Tan, S.L., Lippo, L., De Angelis, J., Labella, R., Singh, A., Chatzis, A., Cheuk, S., Medhghalchi, M., Gil, J., Hollander, G., Marsden, B.D., Williams, R., Ramasamy, S.K., Kusumbe, A.P., 2021. High-resolution 3D imaging uncovers organ-specific vascular control of tissue aging. Science Advances 7, eabd7819.

Cho, Z.H., Kang, C.K., Han, J.Y., Kim, S.H., Kim, K.N., Hong, S.M., Park, C.W., Kim, Y.B., 2008. Observation of the lenticulostriate arteries in the human brain in vivo using 7.0T MR angiography. Stroke 39, 1604-1606.

Coupé, P., Yger, P., Prima, S., Hellier, P., Kervrann, C., Barillot, C., 2008. An optimized blockwise nonlocal means denoising filter for 3-D magnetic resonance images. IEEE transactions on medical imaging 27, 425-441.

Debette, S., Beiser, A., DeCarli, C., Au, R., Himali, J.J., Kelly-Hayes, M., Romero, J.R., Kase, C.S., Wolf, P.A., Seshadri, S., 2010. Association of MRI markers of vascular brain injury with incident stroke, mild cognitive impairment, dementia, and mortality: the Framingham Offspring Study. Stroke 41, 600-606.

Frangi, A.F., Niessen, W.J., Vincken, K.L., Viergever, M.A., 1998. Multiscale vessel enhancement filtering. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 130-137.

Gibbons, L.E., Carle, A.C., Mackin, R.S., Harvey, D., Mukherjee, S., Insel, P., Curtis, S.M., Mungas, D., Crane, P.K., Initiative, A.s.D.N., 2012. A composite score for executive functioning, validated in Alzheimer's Disease Neuroimaging Initiative (ADNI) participants with baseline mild cognitive impairment. Brain imaging and behavior 6, 517-527.

Gotoh, K., Okada, T., Satogami, N., Yakami, M., Takahashi, J.C., Yoshida, K., Ishii, A., Tanaka, S., Miyamoto, S., Togashi, K., 2012. Evaluation of CT angiography for visualisation of

the lenticulostriate artery: difference between normotensive and hypertensive patients.
The British Journal of Radiology 85, e1004-e1008.
Hentschke, C.M., Beuing, O., Paukisch, H., Scherlach, C., Skalej, M., Tönnies, K.D., 2014. A
system to detect cerebral aneurysms in multimodality angiographic data sets. Med Phys
41, 091904.
Hu, Z., van der Kouwe, A., Han, F., Xiao, J., Chen, J., Han, H., Bi, X., Li, D., Fan, Z., 2021.
Motion-compensated 3D turbo spin-echo for more robust MR intracranial vessel wall
imaging. Magn Reson Med 86, 637-647.
Jann, K., Shao, X., Ma, S.J., Cen, S.Y., D'Orazio, L., Barisano, G., Yan, L., Casey, M., Lamas,
J., Staffaroni, A.M., Kramer, J.H., Ringman, J.M., Wang, D.J.J., 2021. Evaluation of
Cerebral Blood Flow Measured by 3D PCASL as Biomarker of Vascular Cognitive
Impairment and Dementia (VCID) in a Cohort of Elderly Latinx Subjects at Risk of
Small Vessel Disease. Frontiers in Neuroscience 15.
Jerman, T., Pernus, F., Likar, B., Spiclin, Z., 2016. Enhancement of Vascular Structures in 3D
and 2D Angiographic Images. IEEE Trans Med Imaging 35, 2107-2118.
Kammerer, S., Mueller-Eschner, M., Berkefeld, J., Tritt, S., 2017. Time-resolved 3D rotational
angiography (4D DSA) of the lenticulostriate arteries: display of normal anatomic
variants and collaterals in cases with chronic obstruction of the MCA. Clinical
Neuroradiology 27, 451-457.
Kang, C.K., Park, C.A., Lee, H., Kim, S.H., Park, C.W., Kim, Y.B., Cho, Z.H., 2009.
Hypertension correlates with lenticulostriate arteries visualized by 7T magnetic
resonance angiography. Hypertension 54, 1050-1056.
Klein, S., Staring, M., Murphy, K., Viergever, M.A., Pluim, J.P., 2010. elastix: a toolbox for
intensity-based medical image registration. IEEE Trans Med Imaging 29, 196-205.
Lowerison, M.R., Sekaran, N.V.C., Zhang, W., Dong, Z., Chen, X., Llano, D.A., Song, P., 2022.
Aging-related cerebral microvascular changes visualized using ultrasound localization
microscopy in the living mouse. Sci Rep 12, 619.
Luu, H.M., Klink, C., Moelker, A., Niessen, W., van Walsum, T., 2015. Quantitative evaluation
of noise reduction and vesselness filters for liver vessel segmentation on abdominal CTA
images. Phys Med Biol 60, 3905-3926.
Ma, S.J., Sarabi, M.S., Yan, L., Shao, X., Chen, Y., Yang, Q., Jann, K., Toga, A.W., Shi, Y.,
Wang, D.J.J., 2019. Characterization of lenticulostriate arteries with high resolution
black-blood T1-weighted turbo spin echo with variable flip angles at 3 and 7 Tesla.
Neuroimage 199, 184-193.
Mayda, A., DeCarli, C., 2009. Vascular cognitive impairment: prodrome to VaD. Cambridge
University Press Cambridge, United Kingdom.
Nissim, N.R., O'Shea, A.M., Bryant, V., Porges, E.C., Cohen, R., Woods, A.J., 2016. Frontal
Structural Neural Correlates of Working Memory Performance in Older Adults. Front
Aging Neurosci 8, 328.
Rhoton, A.L., 2002. The cerebral veins. Neurosurgery 51, S159-205.
Rosenberg, G.A., Wallin, A., Wardlaw, J.M., Markus, H.S., Montaner, J., Wolfson, L., Iadecola,
C., Zlokovic, B.V., Joutel, A., Dichgans, M., 2016. Consensus statement for diagnosis of
subcortical small vessel disease. Journal of Cerebral Blood Flow & Metabolism 36, 6-25.
Rudyanto, R.D., Kerkstra, S., van Rikxoort, E.M., Fetita, C., Brillet, P.Y., Lefevre, C., Xue, W.,
Zhu, X., Liang, J., Öksüz, I., Ünay, D., Kadipaşaoğlu, K., Estépar, R.S., Ross, J.C.,
Washko, G.R., Prieto, J.C., Hoyos, M.H., Orkisz, M., Meine, H., Hüllebrand, M.,

Stöcker, C., Mir, F.L., Naranjo, V., Villanueva, E., Staring, M., Xiao, C., Stoel, B.C.,
Fabijanska, A., Smistad, E., Elster, A.C., Lindseth, F., Foruzan, A.H., Kiros, R., Popuri,
K., Cobzas, D., Jimenez-Carretero, D., Santos, A., Ledesma-Carbayo, M.J., Helmberger,
M., Urschler, M., Pienn, M., Bosboom, D.G., Campo, A., Prokop, M., de Jong, P.A.,
Ortiz-de-Solorzano, C., Muñoz-Barrutia, A., van Ginneken, B., 2014. Comparing
algorithms for automated vessel segmentation in computed tomography scans of the lung:
the VESSEL12 study. Med Image Anal 18, 1217-1232.
Sadoun, E., Reed, M.J., 2003. Impaired angiogenesis in aging is associated with alterations in
vessel density, matrix composition, inflammatory response, and growth factor expression.
J Histochem Cytochem 51, 1119-1130.
Sarabi, M.S., Khansari, M.M., Zhang, J., Kushner-Lenhoff, S., Gahm, J.K., Qiao, Y., Kashani,
A.H., Shi, Y., 2020. 3D Retinal Vessel Density Mapping with OCTA- Angiography.
IEEE J Biomed Health Inform PP.
Sato, Y., Westin, C.-F., Bhalerao, A., Nakajima, S., Shiraga, N., Tamura, S., Kikinis, R., 2000.
Tissue classification based on 3D local intensity structures for volume rendering. IEEE
Transactions on visualization and computer graphics 6, 160-180.
Shao, X., Ma, S.J., Casey, M., D'Orazio, L., Ringman, J.M., Wang, D.J.J., 2019. Mapping water
exchange across the blood–brain barrier using 3D diffusion-prepared arterial spin labeled
perfusion MRI. Magnetic Resonance in Medicine 81, 3065-3079.
Staffaroni, A.M., Asken, B.M., Casaletto, K.B., Fonseca, C., You, M., Rosen, H.J., Boxer, A.L.,
Elahi, F.M., Kornak, J., Mungas, D., J.H., K., 2020a. Development and Validation of the
Uniform Data Set (v3.0) Executive Function Composite Score (UDS3-EF). Alzheimer's
& Dementia, in press.
Staffaroni, A.M., Bajorek, L., Casaletto, K.B., Cobigo, Y., Goh, S.Y.M., Wolf, A., Heuer, H.W.,
Elahi, F.M., Ljubenkov, P.A., Dever, R., 2020b. Assessment of executive function
declines in presymptomatic and mildly symptomatic familial frontotemporal dementia:
NIH-EXAMINER as a potential clinical trial endpoint. Alzheimer's & Dementia 16, 11-
21.
Sur, S., Lin, Z., Li, Y., Yasar, S., Rosenberg, P., Moghekar, A., Hou, X., Kalyani, R., Hazel, K.,
Pottanat, G., Xu, C., van Zijl, P., Pillai, J., Liu, P., Albert, M., Lu, H., 2020. Association
of cerebrovascular reactivity and Alzheimer pathologic markers with cognitive
performance. Neurology 95, e962-e972.
Taha, A.A., Hanbury, A., 2015. Metrics for evaluating 3D medical image segmentation: analysis,
selection, and tool. BMC Med Imaging 15, 29.
Vemuri, P., Decarli, C., Duering, M., 2022. Imaging Markers of Vascular Brain Health:
Quantification, Clinical Implications, and Future Directions. Stroke 53, 416-426.
Wallin, A., Román, G.C., Esiri, M., Kettunen, P., Svensson, J., Paraskevas, G.P., Kapaki, E.,
2018. Update on Vascular Cognitive Impairment Associated with Subcortical Small-
Vessel Disease2. J Alzheimers Dis 62, 1417-1441.
Wardlaw, J.M., Smith, C., Dichgans, M., 2013a. Mechanisms of sporadic cerebral small vessel
disease: insights from neuroimaging. The Lancet Neurology 12, 483-497.
Wardlaw, J.M., Smith, E.E., Biessels, G.J., Cordonnier, C., Fazekas, F., Frayne, R., Lindley, R.I.,
O'Brien, J.T., Barkhof, F., Benavente, O.R., Black, S.E., Brayne, C., Breteler, M.,
Chabriat, H., DeCarli, C., de Leeuw, F.-E., Doubal, F., Duering, M., Fox, N.C.,
Greenberg, S., Hachinski, V., Kilimann, I., Mok, V., Oostenbrugge, R.v., Pantoni, L.,
Speck, O., Stephan, B.C.M., Teipel, S., Viswanathan, A., Werring, D., Chen, C., Smith,

C., van Buchem, M., Norrving, B., Gorelick, P.B., Dichgans, M., 2013b. Neuroimaging
standards for research into small vessel disease and its contribution to ageing and
neurodegeneration. The Lancet Neurology 12, 822-838.
Watanabe, C., Imaizumi, T., Kawai, H., Suda, K., Honma, Y., Ichihashi, M., Ema, M., Mizutani,
K.I., 2020. Aging of the Vascular System and Neural Diseases. Front Aging Neurosci 12,
557384.
Yan, L., Liu, C.Y., Smith, R.X., Jog, M., Langham, M., Krasileva, K., Chen, Y., Ringman, J.M.,
Wang, D.J.J., 2016. Assessing intracranial vascular compliance using dynamic arterial
spin labeling. NeuroImage 124, 433-441.
Yokosawa, K., Kimura, K., Takase, R., Murakami, Y., Boasen, J., 2020. Functional decline of
the precuneus associated with mild cognitive impairment: Magnetoencephalographic
observations. PLoS One 15, e0239577.

Supplementary material

Figure S1. Example high-resolution black blood MRI preprocessing shown on a selected sagittal scan. (a) Raw image of a young control subject. (b) preprocessed image using skull-stripping, bias correction and NLM denoising.

Figure S2. Flow chart of co-registration and VDI normalization. Step1: Co-registration of high-resolution black-blood MRI (a) and MPRAGE (b) image pairs, using 3D-Affine registration with 12 landmark points, where (a) is the fixed and (b) is the moving image. Step2) Nonlinear warp between the MNI-152 Atlas (c) and MPRAGE (e) image. VDI was first reversely transformed to MPRAGE space using affine transformation (f), and then non-linearly transformed to the MNI Atlas using a B-spline transformation (d).