Usefulness of Current sgRNA Design Guidelines and *in vitro* Cleavage Assays for Plant CRISPR/Cas Genome Editing: A Case in Eggplant (*Solanum melongena* L.)

Mark Gabriel S. Sagarbarria[1*] and John Albert M. Caraan[1]

[1]Genetics Laboratory, Institute of Plant Breeding, College of Agriculture and Food Science,

University of the Philippines, Los Baños, Laguna, Philippines

* Author for correspondence; e-mail: mssagarbarria@up.edu.ph

ABSTRACT

The advent of new plant breeding techniques such as genome editing ushers an unprecedented speed on the development of new crop varieties that can withstand agricultural challenges of the 21st century. Genome editing platforms, particularly the CRISPR/Cas9 system, generate highly specific mutations at the DNA sequence level, paving the way towards understanding the role of different genes in plant growth and development. The CRISPR/Cas9 system depends on the specificity of engineered single guide RNAs (sgRNAs) on the gene of interest. However, sgRNA design in plants can be challenging due to a multitude of design tools to choose from, many of which use guidelines that are based on animal experiments yet allow the use of plant genomes. Upon choosing sgRNAs, it is also unclear whether an *in vitro* assay, using a ribonucleoprotein (RNP) complex, is needed to validate the targeting efficiency of a particular sgRNA prior to *in vivo* delivery of the CRISPR/Cas9 system. Here, we demonstrate the *in vitro* and *in vivo* activity of four different sgRNAs that we selected based on an initial design pipeline that we followed. Our results suggest that an *in vitro* RNP cleavage may be a useful step to identify low efficiency sgRNAs, but not as a reliable basis to select and predict sgRNAs with highly efficient *in vivo* cleavage activity. Some sgRNAs that have moderate to high *in vitro* cleavage activity did not produce edits *in vivo,* questioning the necessity of using an *in vitro* assay step. Further analysis of our sgRNAs using other design algorithms did not reveal a consensus on the predicted efficiencies, aside from the CRISPR-P platform, which considers specific plant-validated criteria such as the presence of necessary secondary structures and appropriate base-pairing, which may be more useful in plants than scores predicted by other algorithms.

Key words: CRISPR/Cas9; *in vitro* assay; single guide RNA; *in vivo* genome editing, eggplant

INTRODUCTION

The CRISPR/Cas system has revolutionized the field of plant biology and plant breeding in recent years for its ability to produce precise and targeted induction of mutations at any region on the genome. Despite the development of other genome editing tools such as Zinc Finger Nucleases (ZFNs) or Transcription Activator-Like Effect Nuclease (TALENs), the CRISPR/Cas

system has been widely adopted by many researchers as it allows for rapid generation of gene

manipulation events in a simple, efficient, and specific manner with minimal to no off-target

effects (Gudeta 2019). The technology has progressed to even allow chromosomal deletions

(Ordon et al. 2016) and epigenetic changes (Papikian et al. 2019). This has led to the elucidation

of the functions of numerous genetic elements, which has been applied for crop improvement,

examples for which have been reviewed extensively (Schindele et al. 2020; Zhang et al. 2021).

Genome edits are brought about through the action of the CRISPR/Cas nucleases, which must

recognize a sequence downstream of the target, known as the protospacer-adjacent motif (PAM),

which is 5'-NGG-3' for *Streptococcus pyogenes* Cas9 (SpCas9), a commonly used Cas nuclease

in plants. However, targeting specificity is determined by the single guide RNA or sgRNA,

which usually consists of a 20-bp target-specific CRISPR RNA (crRNA) and the trans-activating

crRNA (tracrRNA) which provides the necessary secondary structure for the formation of the

DNA-sgRNA-Cas9 complex and activation of Cas9 (Jinek et al. 2012). Provided that Cas9

produces a double-stranded break (DSB) three bases upstream of the PAM site, a 20-bp region

upstream of any PAM within the gene of interest can designated as the 'crRNA' to guide Cas9 to

the target site. Afterwards, the crRNA can be cloned in appropriate vectors (ie., in a

Cas9/sgRNA expression vector) for plant transformation.

Theoretically, there are as many target sites in a single gene as there are the number of PAM

sequences. One of the many challenges is choosing which sgRNA will efficiently direct Cas9 to

generate double-stranded breaks (DSBs), and eventually produce the desired edits via the cell's

error-prone repair mechanisms. To aid in this, numerous algorithms have been developed to

predict the on-target efficiency of each sgRNA. However, effectiveness of these tools for plant

gene editing has been questioned recently (Naim et al. 2020). On the other hand, some have

recommended the use of an *in vitro* assay prior to delivery of sgRNAs *in vivo* (Holalu et al.

2019; Karmakar et al. 2021). However, this itself does not guarantee *in vivo* cleavage due to

numerous factors such as chromatin accessibility (Campenhout et al. 2019). As such, there is

currently no definitive protocol for sgRNA design in plants. Hence, we report our findings for

sgRNA efficiency to aid future genome editing studies in lesser researched crops such as

eggplant. In this study, we designed sgRNAs based on their ability to target multiple members of

the polyphenol oxidase gene family in eggplant (*SmelPPO*). We then assessed the activity of

each sgRNA *in vitro* prior to stable transformation and agroinfiltration. Afterwards, we

compared the on-target efficiency scores of each sgRNA using different online tools, as well as

design guidelines of sgRNAs validated for plants. Off-target mutations are less of a concern in

plants as undesired mutations can be eliminated through backcrossing and selection during the

breeding process (Tang et al. 2019). Hence, this study focused on the on-target activity of each

sgRNA.

MATERIALS AND METHODS

Design of sgRNAs

The sgRNAs were initially designed with the aid of the online web tool Benchling

(www.benchling.com), which uses the guide algorithm from Doench et al. (2016). The eggplant

polyphenol oxidase 5 gene (*SmelPPO5/*SMEL4.1_08g017350.1.01) obtained from eggplant

genome V4.1 (Barchi et al. 2021) was used as the input template for sgRNA design. A major

consideration in choosing sgRNAs was complementation to other eggplant PPO genes,

particularly in the conserved regions of the gene family (Shetty et al. 2011). Each sgRNA (from

the   highest   on-target   score)   was   manually   aligned   to   *SmelPPO   4*   and   *6*

(SMEL4.1_08g017940.1.01   and   SMEL4.1_08g017330.1.01)   to   determine   multiple   target

compatibility. Multiple targeting capability of each sgRNA was validated and off-targets were

determined using the CasOT software (Xiao et al. 2014). Each sgRNA chosen was able to target

at least two out of the three target eggplant PPO genes (*SmelPPO 4 – 6)*. We had sequenced

SmelPPO4 – 6 in our target genotype (*Solanum melongena* L. 'PHL 11424') to confirm

sequence similarity with the reference genome. In addition, we also screened the sgRNAs to

avoid internal interactions that may interfere with cleavage such as hairpins (Thyme et al. 2016).

In vitro CRISPR/Cas9 Cleavage Assay

DNA fragments containing the target site were amplified, purified, and eluted in Tris buffer (see

Supplementary Table 1 for the list of all primers used in this study). A ribonucleoprotein

reaction mix composed of synthesized sgRNAs (Macrogen, Korea; 250 ng), purified PCR

products (150 ng) and Cas9 protein (500 ng) (Macrogen, Korea) in 1X NEBuffer 3.1 (100 mM

NaCl, 50 mM Tris-HCl, 10 mM $MgCl_2$, 100 µg/ml BSA, pH 7.9) was prepared following a

minimum of 15:15:1 Cas9:sgRNA:DNA template molar reaction ratio (**Supplementary Table**

2). The reaction mixtures for each sgRNA were incubated at room temperature for 16 hours to

allow the formation of the CRISPR/Cas9 ribonucleoprotein (RNP) and to achieve maximum

cleavage activities. Subsequently, RNAse (10 ng) and Proteinase K (20 μg) were added into the

reaction mix and incubated at 37^{o}C for 15 minutes. The digestion products were separated in a

3.5% agarose TBE gel (1.25% Synergel gel clarifier + 1.0% agarose) and were analyzed using

Gel Analyzer v. 19.1 (Lazar and Lazar 2012). The cleavage efficiencies were assessed by

measuring the amount of digested products over the total amount of the initial DNA used for the

in vitro assay.

Preparation of CRISPR/Cas9 constructs

sgRNAs were expressed under the control of the *Arabidopsis thaliana* U6 promoter. Annealed

oligos of each sgRNA were cloned into pUC19::AtU6-tracrRNA via digestion at the *Bbs*I sites.

The sgRNA expression cassette was ligated into Cas9 expression vector pZH_PubiMMCas9

(Mikami et al. 2015) after digestion of both vectors with *Asc*I and *Pac*I to generate the final

Cas9/sgRNA all-in-one vector. Plasmids were sequenced to verify the insertion of each sgRNA.

Each Cas9/sgRNA all-in-one vector was transformed into *Agrobacterium tumefaciens* strain

'LBA4404'.

Stable transformation and plant regeneration

Agrobacterium-mediated transformation of eggplant (*Solanum melongena* L. 'PHL 11424')

cotyledons was carried out. An established eggplant regeneration protocol using zeatin riboside

(García-Fortea et al. 2020) was followed after agroinfection, but with the presence of selective

antibiotics and timentin. All culture conditions were at 25°C and under 16:8 photoperiod.

Eggplant seeds were surface sterilized with 10% bleach and sown in sterile bottles containing ½

strength Murashige and Skoog (MS) media with 1.5% sucrose and 0.45% vitro agar. Excised

cotyledons ($0.5 - 1\ cm^2$) from two-week old seedlings were wounded at all sides and inoculated

on sterile Petri dishes containing ½ strength MS media with 1.5% sucrose, 0.45% vitro agar, 2

mg/L zeatin riboside, and 100 μM acetosyringone for two days. An overnight culture of *A.*

tumefaciens LBA4404 harboring the Cas9/sgRNA vectors was resuspended in ½ strength MS

media with 1.5% sucrose and 100 μM acetosyringone. Volume was adjusted to reach an OD_{600}

equal to 0.5. In a sterile Petri dish, explants were submerged in the *Agrobacterium* solution for 5

minutes before drying onto a sterile paper towel and transferring back to the same media again

for a 3-day co-cultivation period. Explants were washed once with sterile distilled water and then

once with a 300 mg/L timentin solution before drying on a sterile paper towel and transferring to

selective media (½ strength MS, 1.5% sucrose, 0.45% vitro agar, 2 mg/L zeatin riboside, 100

mg/L timentin, 7.5 mg/L hygromycin, 200 mg/L PVP40). Shoot primordia were allowed to

develop for 3 – 4 weeks then explants were transferred to elongation media (½ strength MS,

1.5% sucrose, 0.45% vitro agar, 0.5 mg/L gibberellic acid, 100 mg/L timentin, 7.5 mg/L

hygromycin). Shoots around 2 cm in size were transferred to rooting media (½ strength MS,

1.5% sucrose, 0.45% vitro agar, 100 mg/L timentin, 5 mg/L hygromycin). Rooted shoots were

acclimatized in greenhouse conditions for 1 week prior to transfer to soil.

Agroinfiltration

An overnight culture of *A. tumefaciens* LBA4404 harboring the Cas9/sgRNA vectors was

resuspended in ½ strength MS media with 1.5% sucrose and 100 μM acetosyringone. Volume

was adjusted to reach an OD_{600} equal to 0.5. The *Agrobacterium* solution was resuspended in

agroinfiltration buffer (½ strength MS, 1.5% sucrose and 100 mg/L acetosyringone). A 1 mL

needleless syringe was used to carefully inject the abaxial surface of the third fully expanded leaf

of the 4-week-old eggplants. Infiltrated portions of the leaves were marked with a permanent

marker. The plants were kept in dim light conditions at 25°C under 16:8 photoperiod for 3 days

before DNA extraction of the infiltrated areas.

Mutation identification by T7E1 cleavage and heteroduplex mobility assays

Genomic DNA was extracted from acclimatized T_0 plants, from approximately 200 mg of leaf

discs (Paterson et al. 1993). The DNA pellet was resuspended in 40 – 50 μL of TE buffer and

stored at -20°C until use. DNA fragments containing the target sites were amplified and purified

before subjection to the T7 endonuclease I (T7E1) assay and heteroduplex mobility assay (Zhu et

al. 2014). For the T7E1 cleavage assay, the protocol provided by New England Biolabs was

followed. Purified PCR products amplified from T_0 plants and infiltrated leaf samples were

denatured and annealed in NEBuffer 2 with purified PCR products from wild-type samples. A

total of 75 - 150 ng of hybridized PCR products were digested with T7 endonuclease (New

England Biolabs, Catalogue # M0302S) for 30 minutes at 37°C. Digested products were

visualized in 2% agarose gel in 1X TBE after 30 minutes of electrophoresis at 100 V. Gels were

placed in 0.5% ethidium bromide solution for 10 minutes before visualization.

For the heteroduplex mobility assay (HMA), PCR products were denatured and annealed then

resolved by electrophoresis in non-denaturing polyacrylamide gels containing 5% acrylamide-

bisacrylamide (40% acrylamide/bis-acrylamide, 37.5:1), 1X TBE, ammonium persulfate, and

TEMED. After 75 minutes of electrophoresis at 100 V, gels were placed in 0.5% ethidium

bromide solution for 10 minutes before visualization.

RESULTS

Testing of sgRNA activity through an *in vitro* cleavage assay

Four (4) sgRNAs were chosen that could target multiple *SmelPPO* genes (**Table 1 and Figure

1).** Synthetic sgRNAs were used with Cas9 to form the ribonucleoprotein complexes tested for *in

vitro* cleavage efficiencies. Due to similar sizes of the expected cleavage products

(**Supplementary Table 3**), we took the amount for all observed digested bands in computation

of the cleavage efficiency (**Supplementary Table 4**).

Figure 1. **(A)** Linear sequence map of *SmelPPO5* showing the target sites of the designed sgRNAs at various gene domains. Target sites of the sgRNAs were labeled as follows: Yellow = Chloroplast Transit Peptide Domain (CTP); Pink = Near the Copper A Binding Domain (CuA); Blue and Cream = Copper B Binding Domain (CuB). **(B)** sgRNA target sites in multiple *SmelPPO* genes (1) sgRNA_1 targeting *SmelPPO*5 and 6, (2) sgRNA_2 targeting *SmelPPO* 4, 5, and 6, (3) sgRNA_3 targeting *SmelPPO* 4, 5, and 6, and (4) sgRNA_4 targeting *SmelPPO* 4, 5, and 6

181   **Table 1.** List of sgRNAs that could target multiple *SmelPPO* genes.

Gene Targets	Direction	Guide Name	Guide sequence	PAM	T_0 Code[1]
PPO5, PPO6	anti-sense	sgRNA 1	TCTATGAAGGAAAAGTTGAG	AGG	N/A
PPO4[2], PPO5, PPO6	anti-sense	sgRNA 2	GCGGATGCGGAGTTTAGTCA	TGG	T_0_A
PPO4, PPO5, PPO6	anti-sense	sgRNA 3	ATAGAAAACCGGGTCTAGAC	CGG	T_0_B
PPO4, PPO5, PPO6	sense	sgRNA 4	ATGAATGGAAAGCAATCGGA	GGG	T_0_C

182   [1]Code of T_0 plants transformed with the respective sgRNAs.
183   [2]Has one mismatch to PPO4 at the -20 position of the sgRNA (relative to PAM sequence)
184

From the observed digestion products of the RNP cleavage assay, all four (4) sgRNAs were able

to direct the Cas9 endonuclease to the target sites to perform DSBs, albeit at different

efficiencies (**Figure 2**). sgRNA_1 targeting the CTP domain was able to cleave its target genes

(*SmelPPO5* and *6*) at an *in vitro* efficiency rate of 85.15% (+++). sgRNA_2 targeting near the

CuA binding domain, on the other hand, obtained highly efficient *in vitro* cleavage activities on

SmelPPO5 and *6* at a rate of 84.70% (+++), but no cleavage activity was observed in *SmelPPO4*.

sgRNA_2 has a single mismatch (guanine-guanine) at 20[th] base upstream of the PAM for

SmelPPO4. Initially, we had expected that this mismatch would be tolerated by the

CRISPR/Cas9 system, as it has been reported that sgRNAs can exhibit function when the

mismatches occur at the most distal end from the PAM (Anderson et al. 2015). However, since

other targets are present that have perfect complementation, *SmelPPO5* and *6* may have been

preferred for cleavage. Meanwhile, sgRNA_3 targeting the CuB binding domain generated the

least efficient *in vitro* cleavage scores at a rate of 44.40% (+) among the four sgRNAs. This may

be because of nucleotide preferences of Cas9 particularly at the seed sequence (Doench et. al.,

2014). At the 1st base upstream of the PAM (position 20 relative to the target gene), Cas9 favors

a guanine residue the most while greatly disfavors a cytosine residue. In case of sgRNA_3, the

residue which immediately follows the PAM recognition sequence is cytosine. Lastly, we

observed that sgRNA_4 was able to produce highest *in vitro* efficiency score at a rate of 87.48%

(+++) across all target genes *SmelPPO genes* (4, 5, and 6). Since all the tested sgRNAs had

observed cleavage *in vitro*, we forwarded all sgRNAs in transformation experiments.

Figure 2. *In vitro* cleavage assay of the four (4) sgRNA/Cas9 ribonucleoprotein complex on the pooled PCR-amplified *Smel*PPO genes from Philippine Eggplant cv. 'Mistisa' (PH11424). (White arrows) Wild type *Smel*PPO (*4, 5, and* 6)DNA used as control. (Red arrows) Digested *Smel*PPO (*4, 5, and/or* 6) DNA after 16 hours of cleavage assay. Cleavage scores: (+) ≤60% cleavage; (++) 61-80% cleavage; (+++) ≥81% cleavage. *In vitro* cleavage activity only observed for *Smel*PPO5 and *Smel*PPO6

Editing the eggplant PPO gene family through *Agrobacterium*-mediated stable transformation and agroinfiltration

The construct, pZH_PubiMMCas9 (**Figure 3A**), which encodes a Cas9 protein under control of the maize ubiquitin promoter and an sgRNA expression cassette under the control of the *A. thaliana* U6 promoter, was used in transformation of eggplant cotyledon explants. A total of 9 plants were regenerated and acclimatized across sgRNA_2 - 4. All obtained regenerants tested positive for the presence of *Cas9* and *HPT* transgenes (**Supplementary Figure 1**), indicating successful transformation. The T7 endonuclease I (T7E1) assay and heteroduplex mobility assays (HMA) were used to detect mutations in all the target sites for each sgRNA. Mutations were only detected from plants transformed with sgRNA_4/T_0_C plants (**Figure 3C-D, Figure 3F-G, Figure 4A-B**). For plants transformed with sgRNA_2/T_0_A plants, multiple bands for each gene target were detected in the HMA even in control samples (**Figure 3D**). However, no cleavage products were observed in the T7E1 assay (**Figure 3C**). We attribute these to be 'shadow' bands that formed due to the different structural formations that may have formed in the PCR products that were separated in non-denaturing PAGE. Since these bands were also present on the control samples, but mismatch cleavage was not observed in the T7E1 assay, the multiple observed bands are most likely the same DNA fragment in different structural conformations. We had also observed that samples positive in the T7E1 assay were also positive in the HMA aside from T_0_C2 for *SmelPPO6*, which tested positive in the HMA but negative in the T7E1 assay. This could be because the T7E1 assay is considered less sensitive in accurately reporting the percentage of edited cells (Sentmanat et al. 2018), but is still useful as it does not overestimate or produce false positive results (Bell et al. 2014).

232   Unfortunately, no regenerants were obtained for sgRNA_1. We included a transient

233   transformation of sgRNA1 into eggplant leaves via agroinfiltration to determine if sgRNA_1

234   could produce edits *in vivo*. However, no edits were detected in both T7E1 and HMA (**Figure**

235   **4C-D**).

Figure 3. (A) Cas9/sgRNA all-in-one vector plasmid based on pZH_PubiMMCas9. (B) Purified PCR products of *SmelPPO4-6* in T0_A plants. (C) T7E1 assay of PCR products of *SmelPPO4-6* in T0_A plants hybridized with PCR products from wild-type plants in 2% agarose gel. (D) HMA of PCR products of of *SmelPPO4-6* in T0_A plants in 5% PAGE. (E) Purified PCR products of SmelPPO4-6 in T0_B and T0_C plants. (F) T7E1 assay of PCR products of *SmelPPO4-6* in T0_B and T0_C plants hybridized with PCR products from wild-type plants in 2% agarose gel. (G) HMA of PCR products of of *SmelPPO4-6* in T0_B and T0_C plants in 5% PAGE. Red arrows indicate presence of cleavage products for T7E1 assay and heteroduplexes from HMA.

Figure 4. (A) T7E1 assay of PCR products of *SmelPPO4-6* in T0_C plants hybridized with PCR products from wild-type plants in 2% agarose gel. **(B)** HMA of PCR products of of *SmelPPO4-6* in T0_C plants in 5% PAGE. (C) T7E1 assay of PCR products of *SmelPPO5-6* from sgRNA1 agroinfiltrated leaves hybridized with PCR products from wild-type plants in 2% agarose gel. (D) HMA of PCR products of of *SmelPPO5-6* from sgRNA1 agroinfiltrated leaves in 5% PAGE.

Comparison of *in vitro* and *in vivo* genome editing using secondary structure guidelines and sgRNA design scores

In order to understand the potential reasons that could have caused the discrepancy between the *in vitro* and *in vivo* cleavage/mutagenesis, we had investigated the predicted efficiency of each sgRNA across numerous design platforms and the secondary structure of each sgRNA. Five total tools were used to predict the efficiency of the chosen sgRNAs: 1) CRISPR-P (Liu et al. 2017), 2) CRISPRDB (Chen and Wang 2022), 3) CRISPOR (Concordet and Haeussler 2018), 4) CCTop (Stemmer et al. 2015), 5) CasDesigner (Park et al. 2015). The Doench et al. (2016) score was used for CRISPOR, since according to the CRISPOR manual, this performed better for U6 datasets. It is important to note that only CRISPOR supported the use of the eggplant reference genome to check for off-targets. Thus, only on-target scores were obtained from these platforms. Lastly, secondary structures and internal base-pairing of each sgRNA was checked using Mfold (www.unafold.org) (Zuker 2003), using guidelines from highly efficient sgRNAs in plants (Liang et al. 2016).

We also investigated the major characteristics of an effective sgRNA (Doench et al. 2016; Mohammadhassan et al. 2022), which are 1) number of targets/off-targets, 2) %GC content, 3) sgRNA length, 4) sgRNA sequence, and 5) secondary structure. For the off-targets, it is assumed that the same sgRNA across all targets and off-target sites would have competition for binding. However, all the tested sgRNAs have a high number of mismatches for potential off-targets, with sgRNA_4 being the lowest at two mismatches for one off-target site (**Table 2**). The CRISPR/Cas9 system has low tolerance for mismatches, with less than 5% of crRNAs being functional at sites with two mismatches (Anderson et al. 2015). Using the same assumption for number of targets, sgRNA_3 and 4 would be the least efficient with sgRNA_1 having higher

mutagenesis rates. However, sgRNA4 was the only sgRNA to produce mutagenesis *in vivo*

despite having the most potential for an off-target site. Therefore, the number of targets/off-

targets does not seem to be a major factor that would explain the observed mutagenesis in the

four sgRNAs. The second factor, %GC content is within the reported optimal range of 30 – 80%

(Liang et al. 2016). It would be difficult to attribute the vast difference in observed *in vitro* and *in*

vivo cleavage to a difference of 5 – 10% GC content, and so this may not be a major factor in

sgRNA editing efficiency. For sgRNA length, all sgRNA used are 20 nt long, which was found

to be the most efficient in rice (Liu et al. 2022). Thus, this should also not be a major factor for

consideration. In addition to sgRNA length, distance between the start codon and the PAM site

has been reported to have a significant impact on cleavage efficiency (Matson et al. 2019).

However, the distance between the non-functional sgRNA_3 and functional sgRNA_4 is on

average 65 nt across their three target sites, ruling out this factor. For sgRNA sequence, animal

studies have reported certain nucleotide positions attributed to more functional sgRNA (Doench

et al. 2014). But for sgRNAs validated in plants, it was found that there is no nucleotide

preference for functional sgRNAs (Liang et al. 2016), ruling out this factor. For sgRNA

secondary structure, *in silico* prediction of the secondary structure of each sgRNA revealed that

all four sgRNAs had a secondary structure that contained all the necessary stem loops (RAR,

SL2, and SL3) found in validated plant sgRNAs (Liang et al. 2016). However, all sgRNAs

except sgRNA_4 had excessive base-pairing between the guide sequence (crRNA) and the

scaffold (tracrRNA), exceeding the maximum of 12 bases paired for sgRNA_1 – 3 and more

than the maximum of 7 consecutive base pairs for sgRNA_1 (**Supplementary Table 5**). This

suggests that the base pairings within sgRNA_1 – 3 are favored and affected the base pairing of

the guide sequence with its target DNA, reducing mutagenesis rates *in vivo*. We compared these

results with scores for online prediction tools and observed that only CRISPR-P was able to give

low confidence scores to sgRNA_1 – 3 (**Table 2**), as only CRISPR-P considers these guidelines

for plant validated sgRNAs.

Table 2. Comparison of different factors affecting sgRNA efficiency in comparison to observed *in vitro* and *in vivo* cleavage results.

Targets	Name	GC (%)	Mismatch number[1]	Off-targets[2]	sgRNA Rankings					Liang et al. (2016) Guidelines[3]	*In vitro*	*In vivo*
					CRISPR-P	CRISPRDB	CRISPOR	CCTop	Cas-Designer			
PPO5, PPO6	sgRNA_1	35	3	5	0.0769	80.5	66	0.76	82	Fail	+++	No
PPO4, PPO5, PPO6	sgRNA_2	55	4	4	0.039	51.7	46	0.69	66.6	Fail	+++[4]	No
PPO4, PPO5, PPO6	sgRNA_3	45	3	1	0.0086	30.2	48	0.69	62.7	Fail	+	No
PPO4, PPO5, PPO6	sgRNA_4	40	2	1	0.6809	92.2	66	0.49	71.8	Pass	+++	Yes

[1]minimum number of mistaches between sgRNA sequences and possible off-target sequences
[2]Number of off-targets at that have the minimum mismatch number
[3]See Supplementary Table 2 for a detailed breakdown for each sgRNA
[4]*In vitro* cleavage activity only observed for *SmelPPO*5 and *SmelPPO*6

DISCUSSION

It has been previously suggested that sgRNA ranking tools for genome editing in plants were not

useful in avoiding less cleavable sites or selecting highly susceptible sites, and that it would be

more effective to choose PAM sites in coding exons for gene disruption rather than choose solely

from prediction scores (Naim et al. 2020). We took a similar approach in that we prioritized

sgRNAs that could target multiple members of the eggplant polyphenol oxidase (*SmelPPO*) gene

family and narrowing down the list with the ones with high on-target scores from our chosen

design platform, low number of off-targets, and checking for hairpin structures that could

interfere with cleavage. Due to other reports that sgRNA prediction algorithms are not able to

consistently predict functional sgRNAs (Gagnon et al. 2014; Thyme et al. 2016; Grainger et al.

2017), we subjected each sgRNA to an *in vitro* assay. All four of our chosen sgRNAs were able

to produce cleavage *in vitro,* which we had initially assumed would result in at least some level

of *in vivo* activity. Some reports had used an *in vitro* cleavage assay to screen for efficient

sgRNA prior to delivery of the Cas9/sgRNA in plants (Malnoy et al. 2016; Liang et al. 2017;

Bente et al. 2020). Although in zebrafish, sgRNA with at least some *in vitro* activity were found

to produce mutations *in vivo,* while those with no *in vitro* activity failed to produce mutations *in*

vivo (Grainger et al. 2017).

Activity *in vivo* can be further reduced as some genomic regions may not be susceptible to Cas9

cleavage due to limited chromatin accessibility (Uusi-Mäkelä et al. 2018). This is not considered

in current sgRNA design algorithms yet. For this study, we assume that *SmelPPO 4 – 6* are

available for cleavage by Cas9 as these are expressed in the plant (leaves, stems, flowers, fruits)

at relatively abundant levels in comparison to house-keeping tubulin genes (Shetty et al. 2011).

However, chromatin accessibility is a major determinant of Cas9 cleavage (Chung et al. 2020)

and should be considered in future genome editing experiments should data of plant epigenetic

landscapes become available.

Our observations in this study indicate that sgRNA with *in vitro* activity will not necessarily

produce mutations *in vivo* in eggplant. We observed that the *in vitro* assay could be useful in

predicting which sgRNAs will have low *in vivo* activity but could not discriminate the sgRNAs

that had high *in vitro* activity but low activity *in vivo*. We have narrowed down the major factor

for the discrepancy between *in vitro* and *in vivo* mutagenesis to be the presence of the needed

sgRNA secondary structures, as well as the number of base pairings within the sgRNA. Seeing

that some sgRNAs with moderate to high cleavage efficiency *in vitro* could not produce edits *in*

vivo, we began to think if the *in vitro* assay was a necessary step. Recently, it has been

mentioned that synthetic RNA guides do not seem to have a link between efficiency and

sequence at all (Haeussler and Concordet 2023). Our results support this statement so far, and we

suggest that resource-constrained laboratories that are interested in doing plant genome editing

work may choose to skip the *in vitro* step, provided they screen for the plant-validated

parameters in functional sgRNAs. Furthermore, we recommended the following pipeline for

sgRNA design in plants, whose reference genome may not necessarily be available in CRISPR-P

for the generation of specificity/off-target scores (**Figure 5**).

Figure 5. Suggested pipeline for sgRNA design in plants.

RECOMMENDATIONS

A major constraint of our study is that we had initially used Benchling for the design of our

initial list of sgRNAs. The Benchling platforms still follows the Doench et al. (2016) algorithm

which does not consider guidelines for plant validated sgRNAs. Perhaps if we used the CRIPSR-

P platform from the start, we would have had more success in finding efficient sgRNAs. In

addition, we had only used the T7E1 assay and HMA to detect mutations. Since these are PCR-

based assays, it is biased towards the more abundant template and small numbers of mutations

may not be detected (Zischewski et al. 2017). It could be possible that edits are present from

plants transformed with sgRNA1-3 but were not detected in the T7E1 assay and HMA. Further

mutation screening should be done in T_1 generations using more accurate methods such as

sequencing.

DECLARATIONS

Funding

This study was made possible through the support provided by the following institutions: the

Philippine Department of Science and Technology – Science Education Institute (DOST-SEI) for

the M.Sc. thesis grants of MSSagarbarria under the Accelerated Science and Technology Human

Resources Development Program (ASTHDRP) and JMCaraan under the Graduate Research and

Education Assistantship for Technology (GREAT) Program; the Philippine Council for

Agriculture, Aquatic and Natural Resources Research and Development (PCAARRD) project

grant to the University of the Philippines Los Baños (UPLB) led by Dr. Lourdes D. Taylo

(Project Code: N91832A) for the research funds. The research was also supported in part by

Plant Transgenic Design Initiative (PTraD) in the Gene Research Center at Tsukuba Plant

Innovation Research Center, University of Tsukuba, Japan.

Conflict of Interest

The authors declare that they have no conflict of interest.

Authors' contributions

Mark Gabriel S. Sagarbarria conceptualized the study and both authors contributed to the study

design. John Albert M. Caraan performed most of the *in vitro* experiments and construct

assembly. Mark Gabriel S. Sagarbarria performed the T7E1 assay and HMA. Transformation,

regeneration, material preparation, data collection and analysis were performed by both authors.

The first draft of the manuscript was written by Mark Gabriel S. Sagarbarria and all authors

commented on previous versions of the manuscript. All authors read and approved the final

manuscript.

Acknowledgements

We thank Dr. Kazuo N. Watanabe, who provided the sgRNA cloning vector and all-in-one

Cas9/sgRNA vector. We thank Dr. Desiree M. Hautea for her research supervision in the early

stages of this work. We also thank Ms. Rowena B. Frankie for her assistance with tissue culture

and Mr. Angelo Layos for assistance in DNA preparation and PCR.

REFERENCES

Anderson EM, Haupt A, Schiel JA, et al (2015) Systematic analysis of CRISPR–Cas9 mismatch tolerance
     reveals low levels of off-target activity. J Biotechnol 211:56–65.
     https://doi.org/10.1016/J.JBIOTEC.2015.06.427

Barchi L, Rabanus-Wallace MT, Prohens J, et al (2021) Improved genome assembly and pan-genome
     provide key insights into eggplant domestication and breeding. https://doi.org/10.1111/tpj.15313

Bell CC, Magor GW, Gillinder KR, Perkins AC (2014) A high-throughput screening strategy for
     detecting CRISPR-Cas9 induced mutations using next-generation sequencing. BMC Genomics
     15:1–7. https://doi.org/10.1186/1471-2164-15-1002/FIGURES/3

Bente H, Mittelsten Scheid O, Donà M (2020) Versatile in vitro assay to recognize Cas9-induced
     mutations. Plant Direct 4:e00269. https://doi.org/10.1002/PLD3.269

Campenhout C van, Cabochette P, Veillard A-C, et al (2019) Guidelines for optimized gene knockout
     using CRISPR/Cas9. Biotechniques 66:295–302. https://doi.org/10.2144/btn-2018-0187

Chen Y, Wang X (2022) Evaluation of efficiency prediction algorithms and development of ensemble
     model for CRISPR/Cas9 gRNA selection. Bioinformatics 38:5175–5181.
     https://doi.org/10.1093/BIOINFORMATICS/BTAC681

Chung CH, Allen AG, Sullivan NT, et al (2020) Computational Analysis Concerning the Impact of DNA
     Accessibility on CRISPR-Cas9 Cleavage Efficiency. Molecular Therapy 28:19–28.
     https://doi.org/10.1016/j.ymthe.2019.10.008

Concordet JP, Haeussler M (2018) CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing
     experiments and screens. Nucleic Acids Res 46:W242–W245.
     https://doi.org/10.1093/NAR/GKY354

Doench JG, Fusi N, Sullender M, et al (2016) Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. Nat Biotechnol 34:184–191. https://doi.org/10.1038/nbt.3437

Doench JG, Hartenian E, Graham DB, et al (2014) Rational design of highly active sgRNAs for CRISPR-Cas9–mediated gene inactivation. Nat Biotechnol 32:1262–1267. https://doi.org/10.1038/nbt.3026

Gagnon JA, Valen E, Thyme SB, et al (2014) Efficient mutagenesis by Cas9 protein-mediated oligonucleotide insertion and large-scale assessment of single-guide RNAs. PLoS One 9:5–12. https://doi.org/10.1371/journal.pone.0098186

García-Fortea E, Lluch-Ruiz A, Pineda-Chaza BJ, et al (2020) A highly efficient organogenesis protocol based on zeatin riboside for in vitro regeneration of eggplant. BMC Plant Biol 20:6. https://doi.org/10.1186/s12870-019-2215-y

Grainger S, Lonquich B, Oon CH, et al (2017) CRISPR guide RNA validation *in vitro*. Zebrafish 14:383–386. https://doi.org/10.1089/zeb.2016.1358

Gudeta D (2019) Genome Editing: Tools and Application in Plants. Open Access Journal of Microbiology & Biotechnology 4:. https://doi.org/10.23880/oajmb-16000135

Haeussler M, Concordet J-P (2023) CRISPOR Manual. http://crispor.tefor.net/manual/. Accessed 16 Feb 2023

Holalu S, Fang A, Blackman B, Benjamin Blackman M (2019) In vitro testing of guide RNA efficiency for CRISPR-mediated genome editing. https://doi.org/10.17504/PROTOCOLS.IO.8ZSHX6E

Jinek M, Chylinski K, Fonfara I, et al (2012) A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. Science (1979) 337:816–821. https://doi.org/10.1126/science.1225829

Karmakar S, Behera D, Baig MJ, Molla KA (2021) In Vitro Cas9 Cleavage Assay to Check Guide RNA Efficiency. pp 23–39

Lazar I, Lazar I (2012) Gel Analyzer 2010a: Freeware 1D gel electrophoresis image analysis software.

Liang G, Zhang H, Lou D, Yu D (2016) Selection of highly efficient sgRNAs for CRISPR/Cas9-based plant genome editing. Sci Rep 6:21451. https://doi.org/10.1038/srep21451

Liang Z, Chen K, Li T, et al (2017) Efficient DNA-free genome editing of bread wheat using CRISPR/Cas9 ribonucleoprotein complexes. Nature Publishing Group 8:1–5. https://doi.org/10.1038/ncomms14261

Liu H, Ding Y, Zhou Y, et al (2017) CRISPR-P 2.0: An Improved CRISPR-Cas9 Tool for Genome Editing in Plants. Mol Plant 10:530–532. https://doi.org/10.1016/J.MOLP.2017.01.003

Liu X, Yang J, Song Y, et al (2022) Effects of sgRNA length and number on gene editing efficiency and predicted mutations generated in rice. Crop J 10:577–581. https://doi.org/10.1016/J.CJ.2021.05.015

Malnoy M, Viola R, Jung M-H, et al (2016) DNA-free genetically edited grapevine and apple protoplast using CRISPR/Cas9 ribonucleoproteins. Front Plant Sci 7:1–9. https://doi.org/10.3389/fpls.2016.01904

Matson AW, Hosny N, Swanson ZA, et al (2019) Optimizing sgRNA length to improve target specificity
and efficiency for the GGTA1 gene using the CRISPR/Cas9 gene editing system. PLoS One
14:e0226107. https://doi.org/10.1371/JOURNAL.PONE.0226107

Mikami M, Toki S, Endo M (2015) Comparison of CRISPR/Cas9 expression constructs for efficient
targeted mutagenesis in rice. Plant Mol Biol 88:561–572. https://doi.org/10.1007/s11103-015-0342-
x

Mohammadhassan R, Tutunchi S, Nasehi N, et al (2022) The Prominent Characteristics of the Effective
sgRNA for a Precise CRISPR Genome Editing. CRISPR Technology - Recent Advances.
https://doi.org/10.5772/INTECHOPEN.106711

Naim F, Shand K, Hayashi S, et al (2020) Are the current gRNA ranking prediction algorithms useful for
genome editing in plants? PLoS One 15:e0227994. https://doi.org/10.1371/journal.pone.0227994

Ordon J, Gantner J, Kemna J, et al (2016) Generation of chromosomal deletions in dicotyledonous plants
employing a user-friendly genome editing toolkit. https://doi.org/10.1111/tpj.13319

Papikian A, Liu W, Gallego-Bartolomé J, Jacobsen SE (2019) Site-specific manipulation of Arabidopsis
loci using CRISPR-Cas9 SunTag systems. Nat Commun 10:729. https://doi.org/10.1038/s41467-
019-08736-7

Park J, Bae S, Kim JS (2015) Cas-Designer: a web-based tool for choice of CRISPR-Cas9 target sites.
Bioinformatics 31:4014–4016. https://doi.org/10.1093/BIOINFORMATICS/BTV537

Paterson AH, Brubabker CL, Wendel JF (1993) Genomic DNA suitable for RFLP or PCR analysis. Plant
Mol Biol Report 11:122–127

Schindele A, Dorn A, Puchta H (2020) CRISPR/Cas brings plant biology and breeding into the fast lane.
Curr Opin Biotechnol 61:7–14. https://doi.org/10.1016/J.COPBIO.2019.08.006

Sentmanat MF, Peters ST, Florian CP, et al (2018) A Survey of Validation Strategies for CRISPR-Cas9
Editing. Scientific Reports 2018 8:1 8:1–8. https://doi.org/10.1038/s41598-018-19441-8

Shetty SM, Chandrashekar A, Venkatesh YP (2011) Eggplant polyphenol oxidase multigene family:
Cloning, phylogeny, expression analyses and immunolocalization in response to wounding.
Phytochemistry 72:2275–2287. https://doi.org/10.1016/j.phytochem.2011.08.028

Stemmer M, Thumberger T, del Sol Keyer M, et al (2015) CCTop: An Intuitive, Flexible and Reliable
CRISPR/Cas9 Target Prediction Tool. PLoS One 10:e0124633.
https://doi.org/10.1371/JOURNAL.PONE.0124633

Tang J, Chen L, Liu Y-G (2019) Off-target effects and the solution. Nat Plants 5:341–342.
https://doi.org/10.1038/s41477-019-0406-z

Thyme SB, Akhmetova L, Montague TG, et al (2016) Internal guide RNA interactions interfere with
Cas9-mediated cleavage. Nat Commun 7:1–7. https://doi.org/10.1038/ncomms11750

Uusi-Mäkelä MIE, Barker HR, Bäuerlein CA, et al (2018) Chromatin accessibility is associated with
CRISPR-Cas9 efficiency in the zebrafish (*Danio rerio*). PLoS One 13:e0196238.
https://doi.org/10.1371/journal.pone.0196238

Xiao A, Cheng Z, Kong L, et al (2014) CasOT: A genome-wide Cas9/gRNA off-target searching tool.
482   Bioinformatics 30:1180–1182. https://doi.org/10.1093/bioinformatics/btt764

483   Zhang D, Zhang Z, Unver T, Zhang B (2021) CRISPR/Cas: A powerful tool for gene function study and
484   crop improvement. J Adv Res 29:207–221. https://doi.org/10.1016/j.jare.2020.10.003

485   Zhu X, Xu Y, Yu S, et al (2014) An efficient genotyping method for genome-modified animals and
486   human cells generated with CRISPR/Cas9 system. Sci Rep 4:. https://doi.org/10.1038/srep06420

487   Zischewski J, Fischer R, Bortesi L (2017) Detection of on-target and off-target mutations generated by
488   CRISPR/Cas9 and other sequence-specific nucleases. Biotechnol Adv 35:95–104.
489   https://doi.org/10.1016/J.BIOTECHADV.2016.12.003

490   Zuker M (2003) Mfold web server for nucleic acid folding and hybridization prediction. Nucleic Acids
491   Res 31:3406–3415. https://doi.org/10.1093/nar/gkg595

492