Title: Enhanced protective efficacy of a novel, thermostable, RBD-S2 vaccine formulation against SARS-CoV-2 and its variants

Authors: Nidhi Mittal[1], Sahil Kumar[2], Raju S Rajmani[1], Randhir Singh[3], Céline Lemoine[4], Virginie Jakob[4], Sowrabha BJ[3], Nayana Jagannath[3], Madhuraj Bhat[3], Debajyoti Chakraborty[1], Suman Pandey[3], Aurélie Jory[5], Suba Soundarya S.A.[5], Harry Kleanthous[6], Patrice Dubois[4], Rajesh P. Ringe[2*], Raghavan Varadarajan[1*]

Affiliations:

[1]Molecular Biophysics Unit (MBU), Indian Institute of Science; Bengaluru 560012, India

[2]Virology Unit, Institute of Microbial Technology, Council of Scientific and Industrial Research (CSIR); Chandigarh 160036, India

[3]Mynvax Private Limited; 3rd Floor, Brigade MLR Centre, No.50, Vani Vilas Road, Basavanagudi, Bengaluru 560004, India

[4]Vaccine Formulation Institute; Rue du Champ-Blanchod 4, 1228 Plan-les-Ouates, Switzerland

[5]National Centre for Biological Sciences, Tata Institute of Fundamental Research; Bengaluru 560065, India

[6]Bill and Melinda Gates Foundation; Seattle, WA, 98109, USA

*Corresponding authors: rajeshringe@imtech.res.in ; varadar@iisc.ac.in

Abstract: With the rapid emergence of variants of concern (VOC), the efficacy of currently licensed vaccines has reduced drastically. VOC mutations largely occur in the S1 subunit of Spike. The S2 subunit of SARS-CoV-2 is conserved and thus more likely to elicit broadly protective immune responses. However, the contribution of the S2 subunit in improving the overall efficacy of vaccines remains unclear. Therefore, we designed, characterized, and evaluated the immunogenicity and protective potential of a stabilized SARS-CoV-2 Receptor Binding Domain (RBD) fused to a stabilized S2. Designed immunogens were expressed as soluble proteins with approximately fivefold higher purified yield than the Spike ectodomain and formulated along with Squalene-in-water emulsion (SWE) adjuvant. S2 immunization failed to elicit a neutralizing immune response but significantly reduced lung viral titers in mice challenged with the heterologous Beta variant. In hamsters, SWE-formulated RS2 showed enhanced immunogenicity and efficacy relative to corresponding RBD and Spike formulations. Despite being based on the ancestral Wuhan strain of SARS-CoV-2, RS2 exhibited broad neutralization, including against Omicron variants (BA.1, BA.5 and BF.7), as well as the clade 1a WIV-1 and SARS-CoV-1 strains. RS2 sera also showed enhanced competition with both S2 directed and RBD Class 4 directed broadly neutralizing antibodies, relative to RBD and Spike elicited sera. When lyophilized, RS2 retained antigenicity and immunogenicity even after incubation at 37 °C for a month. The data collectively suggest that the RS2 immunogen is a promising modality to combat SARS-CoV-2 variants.

Keywords: Protein engineering, lyophilization, potency, protein design, COVID-19 vaccine, thermotolerant.

INTRODUCTION

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) virus is the causative agent of the COVID-19 pandemic [1]. A variety of SARS-CoV-2 vaccines were developed at an unprecedented pace to combat the pandemic [2–8]. Spike (S) is a ~ 200 kDa transmembrane glycoprotein composed of two functional subunits: the N-terminal S1 subunit, which mediates attachment to the host receptors, and the C-terminal S2 subunit, which facilitates membrane fusion [9–11]. Spike protein is highly immunogenic, and the receptor-binding domain (RBD) contains the major neutralizing antibody epitopes [12–18]. Most currently approved vaccines are based on the Spike immunogen, and a minority are based on RBD. RBD-based vaccines have been shown to elicit moderate to high-neutralizing antibody titers [19–22]. However, due to rapid virus evolution, various variants of concern (VOC) mutations have been identified primarily in the N-terminal domain (NTD) and RBD of the S1 subunit [23]. These mutations increase viral infectivity and induce immune evasion by evading virus neutralization. Requirements for low or ultracold temperature storage have acted as barriers to vaccine deployment in low-and middle-income countries (LMICs). There is an ongoing need for vaccination in vulnerable sections of the population, including pregnant women, those with co-morbidities and the elderly, both to protect these individuals and to minimize the potential for viral evolution in vulnerable hosts. Therefore, it remains important to develop and test vaccine candidates that confer a potent, protective, humoral immune response to a broad spectrum of SARS-CoV-2 variants and do not require a cold-chain for last mile distribution.

Compared to the S1 subunit, the S2 subunit of coronaviruses is more conserved and likely to elicit broadly protective antibodies [24]. Several cross-reactive monoclonal antibodies against the S2 stem helix region have been identified, and a few of them have been shown to neutralize SARS-CoV-2, thereby conferring *in vivo* protection [17,25–33]. Spike microarray analysis demonstrated that cross-reactive antibodies are elicited against the S2 stem helix region during natural infection [17]. The stem-helix epitope in the S2 region, residues (1142-1165) is well conserved, and antibodies that bind to this region neutralize diverse coronaviruses suggesting that immunogens containing this cryptic epitope might elicit pan-coronavirus immunity.

However, the protective efficacy of S2 as an immunogen remains unclear. It is also known that the bulk of the neutralizing response is directed against the RBD. While there are also neutralizing

antibodies directed against the NTD of the Spike, these have lower breadth than the RBD-directed

neutralizing antibodies, as the major NTD neutralizing epitope is mutated in VOCs [34–38].

To address these issues, we designed a stabilized S2-ectodomain, and genetic fusions of a

previously designed, stabilized RBD with S2 [39]. These are referred to as RS2 and S2R immunogens

depending on the order of connectivity of RBD and S2. Designed RS2 and S2R immunogens were

expressed with ~5.3-fold higher purified yield than stabilized Spike ectodomain in mammalian

cells. Sepivac SWE[TM] is an MF59 like oil-in-water emulsion adjuvant, subsequently referred to as

SWE. Two immunizations of SWE adjuvanted RS2 induced robust neutralizing immune responses

and conferred protection against SARS-CoV-2 variants in mice, showing superior immunogenicity

and protective efficacy, compared to stabilized RBD, and comparable immunogenicity to

stabilized Spike formulations. In hamsters, RS2 showed superior immunogenicity and protective

efficacy to a similarly adjuvanted, stabilized Spike formulation. When lyophilized, RS2 retained

antigenicity and immunogenicity even after incubation at 37 °C for a month. The data collectively

suggest that RS2 containing vaccine formulations are a promising modality to combat SARS-CoV-

2 variants.

RESULTS

Immunogen Design

For designing an S2-based SARS-CoV-2 immunogen, S2 subunit residues interacting with the S1

subunit were identified by accessibility calculations using the Naccess program on PDB 6VXX.

Accessible surface areas (ASAs) of all residues in Spike were calculated in the absence and

presence of the S1 subunit. Residues of the S2, which had a total side-chain ASA difference of >5

$Å^2$, were identified as S1 interacting residues. The following mutations were made to mask newly

exposed hydrophobic residues to prevent S2 aggregation in the absence of S1, namely F855S,

L861E, L864D, V976D, and L984Q [40,41].

Proline substitutions in the loop between HR1 (heptad repeat 1) and the central helix in fusion

proteins from viruses like SARS-CoV-2, SARS-CoV, MERS-CoV, and RSV are known to retain

prefusion conformation, and enhance expression yields [42–45]. More recently, HexaPro Spike, a S-

6P variant, with six proline substitutions (F817P, A892P, A899P, A942P, K986P, V987P)

displayed 9.8-fold increased protein expression and ~5 °C increase in T_m relative to S-2P (Spike

with two proline mutation) [46]. Hence, we additionally included all these six proline substitutions

in the present S2 immunogen (S2) comprising residues 698-1211 of the SARS-CoV-2 Spike. Since

RBD contains the major neutralizing antibody epitopes on Spike and S2 is well conserved, we

designed RS2 and S2R immunogens **(Fig. 1A)**. For this, an S2 fragment comprising residues 698-

1163 of the SARS-CoV-2 was genetically fused with our previously reported high-yielding,

thermostable RBD containing A348P, Y365W, and P527L mutations [20,22].

Biophysical characterization of S2, RS2, S2R and Spike

The designed immunogens were transiently expressed as secreted proteins in Expi293F

suspension cells. Recombinant proteins S2, RS2, and S2R were purified in high yields of ~120

mg/L, ~850 mg/L, and ~800 mg/L, respectively, using nickel affinity chromatography **(Fig. 1B)**.

The oligomeric state of the immunogens was determined using size exclusion chromatography

(SEC), and revealed that S2 exists as homogenous nonamers in the solution **(Fig. 1C)**. RS2 and

S2R exist as a mixture of monomers and trimers, and the calculated molecular weights were in

good agreement with the theoretical molecular weight **(Fig. 1D, E)**.

For comparison, the stabilized trimeric Spike ectodomain (1-1211) containing six proline

mutations (F817P, A892P, A899P, A942P, K986P, and V987P), along with the additional three

RBD stabilizing mutations (A348P, Y365W, and P527L) described above, was also expressed and

purified from Expi293F cells **(Fig 1A, B, F)** with a purified yield of ~150 mg/L [22,46].

The apparent melting temperature (T_m) and the thermal unfolding profile of designed immunogens

and Spike was determined using Nano-DSF. The RS2 and S2R displayed similar thermal

unfolding profiles and comparable T_m of ~ 50 °C **(Fig. 2A and B)**. The S2 immunogen exhibited

T_m of 52.2 °C while the stabilized trimeric spike demonstrated two T_m, T_{m1} of 50.4 °C and T_{m2} of

61.5 °C, which implied that trimeric Spike has different structural components of varying stability

(Fig. 2C and D). RS2, S2R, and S2, when subjected to 37 °C for 1 hour, exhibited similar thermal

unfolding profiles to those protein samples stored at 4°C **(Fig. 2E-G)**. However, the Spike showed

a slightly broadened thermal unfolding profile after incubation at 37 °C, with more noticeable

broadening at 50°C **(Fig. 2H)**. Moreover, RS2 and S2R were found to be stable even after

incubation at 50°C, although a slight decrease was observed. On the other hand, S2 was thermally

unstable at 50°C, which suggested that RS2 and S2R are more resistant to transient thermal stress

than Spike and S2. **(Fig. 2E-H)**. Compared to the other designed immunogens and Spike, RS2 was more rapidly digested by TPCK-trypsin at both 4 °C and 37 °C, **(Fig. 2H-K).** This data suggested that RS2 is more susceptible to proteolytic degradation than S2R, S2, and Spike, which showed highest proteolytic stability. Whether the apparent enhanced proteolytic resistance of Spike **(Fig. 2L)** is due to aggregation following initial proteolytic cleavage, remains to be elucidated.

The binding of S2, RS2, and S2R immunogens to its cognate receptor, ACE2-hFc, a panel of RBD conformation-specific (CR3022, S309, ADG-2, and H014) and S2-specific (B6 and CC40.8) antibodies were probed using surface plasmon resonance (SPR). Spike, RS2, and S2R bound well with ACE2-hFc, RBD-specific, and S2-specific antibodies **(Table 1 and 2)**. S2 binds only to S2-specific antibodies with high affinity. This indicated the proper folding of designed immunogens **(Table 2)**.

Immunogenicity and protective efficacy of S2R relative to S2 and RBD immunogens against heterologous challenge.

Since in the initial characterizations, S2R showed higher proteolytic stability and comparable thermal stability to RS2, the immunogenicity and protective efficacy of S2R, relative to RBD and S2 immunogens, was evaluated in hACE-2 expressing C57BL/6 transgenic mice. Our previously reported mammalian cell expressed, stabilized RBD containing A348P, Y365W, and P527L mutations was expressed and purified [20,22]. Mice were intramuscularly immunized with 2 μg immunogens (S2R, S2 or RBD) formulated with SWE in a prime-boost regimen 3 weeks apart. SWE is equivalent to MF59, a very safe adjuvant that has been used for many years in the context of human influenza vaccines [47]. While there are other more potent adjuvants available, stronger adjuvant mediated immune responses can be associated with unfavorable side effects [47]. Two weeks post-boost, RBD, S2 and Spike-specific IgG titers in sera of immunized mice were measured using ELISA. Relative to RBD and S2, the S2R immunogen elicited significantly higher RBD, S2 and Spike-specific ELISA endpoint titers **(Fig. 3A-C)**. S2R immunized mice elicited significantly higher neutralizing antibody titers against B.1 pseudovirus compared to RBD immunized mice, while the sera from S2 immunized mice failed to neutralize B.1 pseudovirus **(Fig. 3D)**. At this administered dose, sera from RBD-immunized mice did not show neutralization

against BA.1. However, S2R immunized mice sera showed neutralization against BA.1, BA.5, and

BF.7 albeit at significantly reduced levels **(Fig. 3E and F)**. Furthermore, these sera exhibited

substantial cross-neutralization against heterologous clade1a SARS-CoV-1 viruses, which

demonstrates the potential of S2R to elicit broadly protective antibodies against sarbecoviruses

(Fig. 3F). Although the neutralization ID_{50} was too low to be measured in mice immunized with

SWE formulated S2**,** addition of these S2 elicited sera enhanced neutralization potency of the

broadly neutralizing antibody S309 **(Fig. 3G)**.

Next, the protective efficacy of RBD, S2, and S2R formulations was assessed against the Beta

variant of SARS-CoV-2. Unimmunized-unchallenged, and unimmunized-Beta variant challenged

mice were used as control groups. Three weeks post boost, mice were intranasally challenged with

10^5 plaque-forming units (pfu) of Beta variant virus, and weight change was monitored for upto

five days. Only 33% of unimmunized mice survived, while 72% of S2 immunized mice survived

Beta-variant challenge. In contrast, all RBD and S2R immunized mice survived the Beta variant

challenge **(Fig. 3H)**. Post challenge, no weight change was seen in S2R immunized mice. Mice

immunized with RBD, and S2 showed a significant weight reduction of ~10 % and ~17%

respectively. As expected, no weight reduction was observed in the unimmunized group, while 22-

25 % weight reduction was seen in the unimmunized-Beta variant challenged group **(Fig. 3I)**. Mice

immunized with either RBD, S2 or S2R showed significantly lower lung viral titers than

unimmunized mice challenged with the Beta variant **(Fig. 3J)**. Despite lack of neutralization, lung

viral titers were significantly reduced in the S2-immunized group, suggesting that S2 provides

protection by non-neutralizing mechanisms. The lung tissue sections obtained from S2R

immunized mice showed clear interstitial spaces within lung epithelium and reduced immune cell

infiltration compared to unimmunized-Beta variant challenged group, RBD and S2 immunized

group **(Fig. 3K and L)**.

RS2 is more immunogenic than S2R in mice.

At the same time, we also compared the immunogenicity of RS2 and S2R in BALB/c mice.

BALB/c mice were vaccinated twice intramuscularly with either 20 µg of RS2 or S2R formulated

with SWE adjuvant. Two weeks after the second vaccination, RBD and Spike specific IgG titers

and neutralizing immune responses were measured in the sera samples of immunized mice. Both

RS2 and S2R immunized mice elicited equivalent RBD and Spike-specific ELISA endpoint titers

(Fig. S1 A and B). RS2 elicited significantly higher neutralizing titers against B.1 pseudovirus **(Fig. S1C)**. Neutralizing titers against Delta variant were also higher in RS2 immunized mice sera but differences did not reach statistical significance **(Fig. S1D)**. Thus, the data indicated that RS2 is more immunogenic than S2R. Hence for all further studies, including formulation stability and comparative immunization studies with Spike, RS2 was used.

RS2 induces an equivalent neutralizing immune response to Spike ectodomain in mice and protects against mouse-adapted SARS-CoV-2 challenge.

Considering that most currently licensed COVID-19 vaccines have the Spike as the sole immunogen, we compared the immunogenicity and protective efficacy of RS2 with the stabilized Spike ectodomain. BALB/c mice were intramuscularly immunized with 2 μg of RS2 or Spike formulated with SWE adjuvant in a prime-boost regimen 21 days apart. Fourteen days post-second immunization, both RS2 and Spike immunized mice elicited high RBD, S2 and Spike-specific IgG titers **(Fig. 4A-C)**. Although the neutralizing titers against B.1, Beta and Delta variant pseudoviruses were comparatively higher in RS2 immunized mice sera than Spike immunized mice sera, the differences did not reach statistical significance **(Fig. 4D-F)**. Sera from RS2 and Spike immunized mice also showed neutralization against BA.1, BA.5 and BF.7 pseudoviruses, albeit with lower titers **(Fig. 4G-I)**. Compared to Spike, RS2 also exhibited substantial cross-neutralization against heterologous clade1a SARS-CoV-1 and WIV-1 viruses, which demonstrates the potential of RS2 to elicit broadly protective antibodies against sarbecoviruses **(Fig. 4J and K)**. Three weeks post-boost, all mice were intranasally challenged with 10^5 pfu of mouse-adapted SARS-CoV-2 MA10 virus. One-day post-challenge, mice immunized with either RS2 or spike showed a slight body weight reduction of ~5 %, and from day two, all mice regained their initial weight. In contrast, unimmunized-MA10 challenge control mice showed ~25 % weight loss by day four, post-challenge. No weight change was seen in the unimmunized-unchallenged control group **(Fig. 4L)**. RS2 and Spike immunized mice showed significantly reduced lung viral titers compared to unimmunized-MA10 mice **(Fig. 4M)**. Analysis of lung tissue sections of mice immunized with RS2 showed clear lung epithelial interstitial spaces and lower immune cell infiltration compared to both Spike immunized and unimmunized MA10 challenged group **(Fig. 4N and O)**. This data suggests that RS2 elicits a broadly neutralizing humoral immune response and protects against SARS-CoV-2 MA10 virus challenge.

Thermal stability of RS2

As with our previously reported RBD immunogens, our newly designed RS2 immunogen showed identical thermal unfolding profiles before and after lyophilization and solubilization, suggesting that lyophilization did not affect the thermal stability of these immunogens **(Fig. S2A)** [20,22]. Further, the effect of transient thermal stress on the thermal stability of lyophilized RS2 was studied by performing nano-DSF. Following incubation at different temperatures (4 °C, 37 °C, 50 °C, 70 °C and 90 °C) for 60 min, no change in thermal unfolding profiles was observed **(Fig. S2B)**. Moreover, the T_m and conformational integrity of lyophilized RS2 protein remained unchanged after a month of storage at 37 °C **(Fig. S2C) (Table 3)**. In addition, the antigenicity of RS2 was assessed by performing SPR using a panel of RBD and S2-specific antibodies. Interestingly, following thermal stress, RS2 retained binding with the cognate receptor ACE2-hFc, RBD-specific antibodies (CR3022, S309), and S2-specific antibodies (B6, CC40.8). This suggests that lyophilized RS2 immunogen is resistant to transient thermal stress **(Fig. S2D-H)**. The stability of SWE adjuvanted RS2 in PBS buffer was also evaluated at 5 °C and 40 °C over a period of one month. Antigen integrity and physiochemical characterization of SWE adjuvanted RS2 formulations was carried out by performing ELISA, particle size, polydispersity, zeta potential, pH, osmolality and squalene content measurements. The RS2 formulation in SWE is stable at 5°C and 40 °C in both polypropylene tubes and glass vials for at least one month **(Fig. 5A-K)**.

Protective efficacy of lyophilized RS2 formulation after month long storage at 37 °C

The immunogenicity of the lyophilized RS2 protein stored at 37 °C for a month was evaluated in hACE-2 expressing C57BL/6 transgenic mice. Mice were intramuscularly immunized with 20 μg immunogen extemporaneously formulated with SWE adjuvant in a prime-boost regimen. Two weeks following the boost, the formulation showed high RBD, S2 and Spike-specific binding titers, and neutralizing titers against B.1, Beta, Delta, and BA.1 pseudoviruses **(Fig. 6A and B)**. The above formulation elicited equivalent neutralization titers to a non-lyophilized freshly prepared RS2 formulation stored at 4 °C, against B.1 and BA.1 pseudoviruses **(Fig. 6C)**. Twenty-one days following the boost, mice were challenged with 10^4 pfu of Beta and Delta variants of SARS-CoV-2. None of the unimmunized control mice survived Beta variant challenge, while 57% of unimmunized control mice survived Delta variant challenge and the remaining mice showed 20

% weight loss, nine days post-challenge. In contrast, all RS2 immunized mice survived the Beta and Delta variant SARS-CoV-2 challenges and showed no weight loss **(Fig. 6D-F)**. Lung viral titers of RS2-immunized mice challenged with Beta and Delta variants were below the detection limit and lung tissues showed minimal pathology **(Fig. 6G and H).** Lung viral titers and tissue sections from the unimmunized mice challenged with the Beta variant were not examined because none of the mice survived. These findings indicate that RS2 is stable and immunogenic even after storage at 37 °C for at least one-month.

RS2 shows superior immunogenicity and protective efficacy to stabilized Spike ectodomain in hamsters.

Protective efficacy of RS2 was compared with the stabilized Spike against challenge with the Beta variant of SARS-CoV-2 in Syrian hamsters. Female Syrian Golden hamsters were immunized with 5μg of stabilized Spike or RS2 formulated with SWE on day 0 and day 21, while the control group of hamsters was immunized with SWE adjuvant in PBS. Both Spike and RS2 immunized hamsters elicited high RBD, S2 and Spike-specific ELISA endpoint titers **(Fig. 7A-C)**. Notably, two immunizations with 5 μg SWE adjuvant formulated RS2 elicited significantly higher neutralizing titers against B.1, Beta, Delta and Omicron BA.1, BA.5 and BF.7 variants than corresponding titers elicited by SWE formulated Spike in hamsters **(Fig. 7D-I)**. Consistent with the BALB/c mice study, RS2 immunized hamsters exhibited substantially higher cross-neutralizing activity against clade 1a sarbecoviruses, WIV-1 and SARS-CoV-1 compared to those of Spike immunized hamsters (**Fig. 7H and K)**. Moreover, hamsters immunized with RS2 showed initial transient weight loss (up to 3 %) and regained weight at 3 days post Beta variant infection. In contrast, Spike-immunized hamsters showed comparatively higher lung viral titers and weight loss, while no weight regain was observed **(Fig. 7L and M)**. Both RS2 and Spike immunized hamsters showed significantly reduced lung viral titers compared to unimmunized-Beta challenged hamsters **(Fig. 7M)**. Analysis of RS2 and Spike immunized hamster lung tissue sections showed clear lung epithelial interstitial spaces and lower immune cell infiltration compared to the unimmunized Beta-challenged groups **(Fig. 7N and O)**. Overall, the data shows that RS2 is more immunogenic and efficacious than Spike in hamsters.

DISCUSSION

Vaccination has significantly reduced the global health burden caused by the COVID-19
pandemic. While vaccines have been proven to be highly efficacious against ancestral SARS-CoV-
2 stain, their efficacy has rapidly declined against new VOCs and periodic updating of vaccines
appears to be necessary. Most VOC mutations are identified in the receptor binding motif (RBM)
region of RBD. The substantial waning of serum-neutralizing antibody titers and the emergence
of variants with increased transmissibility and neutralizing antibody escape responses are
associated with increased breakthrough infection in vaccinated individuals. To curb infection and
sustain vaccine effectiveness, booster doses are required.

Various approaches have been deployed to broaden the protection breadth of SARS-CoV-2
vaccines, which include mosaic nanoparticle vaccine designs that display multiple RBDs from
different sarbecoviruses, and conferred broad protection against diverse coronaviruses, albeit with
a reduction in potency against some SARS-CoV-2 VOCs [48–50]. While promising, such nanoparticle
vaccine designs elicit titers against the nanoparticle scaffolds, and employ a large number of
antigens, which adds to manufacturing complexity. Alternatively, vaccine candidates based on
conserved antigens like S2 are also reported [51–53].

In the present study we designed a stabilized S2 ectodomain, and the genetic fusion of RBD and
S2. The RS2 and S2R contain the RBD component to elicit potent neutralizing antibodies, and the
S2 to increase immunogenicity and protective breadth. The novel RS2 and S2R immunogens
exhibited ~5.3-fold higher protein yields and were shown to be more stable to transient thermal
stress than a stabilized Spike (**Fig. 2E and H**).

All the currently licensed COVID-19 vaccines require refrigerated or frozen storage. Recently,
Ferritin displayed, alum adjuvanted Spike construct also displayed promising immunogenicity in
non-human primates and could be stored at 37°C for at least two weeks[54]. In the present study we
have shown that our RS2 vaccine candidate can retain its thermal stability and antigenicity without
any loss of immunogenicity and protective efficacy at 37 °C for at least a month. Encouragingly,
RS2 immunogen formulated with SWE was also shown to maintain its physico-chemical stability
at 40°C for up to one month, however the immunogenicity needs to be evaluated. This exceptional
stability of the RS2 candidate will facilitate distribution in low resource settings and will reduce
the cost associated with low or ultracold temperature storage and transportation.

Despite being based on the original Wuhan strain sequence, RS2 elicited neutralizing antibodies

against B.1, Beta, Delta, Omicron (BA.1, BA.5, BF.7) and clade 1a SARS-CoV-1 and WIV-1

pseudoviruses in mice and hamsters. While many SARS-CoV-2 immunogens exhibit high

immunogenicity in mice, immunogenicity is poorer in hamsters, and more predictive of

immunogenicity in humans[55–58]. Therefore, we chose to use the hamster animal model to further

probe for apparent differences in immunogenicity between RS2 and Spike immunized animals. In

hamsters, the RS2 immunogen elicited significantly higher neutralizing titers against all the tested

VOCs compared to Spike **(Fig.7D-K)**. Qualitatively, RS2 immunized animals also exhibited less

weight loss and lower lung viral titers compared to Spike immunized animals **(Fig.7H-J)**.

However, due to the limited number of animals in the study, the differences in weight loss and

lung viral titers between the two groups did not reach statistical significance. In addition, the sera

obtained from mice immunized with RS2 showed competition with S2X259, and B6 antibodies.

In contrast, the sera from mice immunized with Spike or RBD failed to show competition with

these antibodies **(Fig. S3)**. These findings suggest that RS2 sera specifically target highly

conserved epitopes located in the RBD and S2 stem helix, which are likely less accessible in Spike

or RBD based immunogens. In contrast to Spike where regions of RBD that contain neutralizing

epitopes are occluded in the down conformation, there no such conformational constraint in the

RBD-S2 fusions.

Like other vaccine formulations, it will be necessary to periodically update formulations based on

RS2. However, due to the lower mutation rate in the S2 region and higher yield, updating RS2

vaccine sequences to match circulating vaccines is expected to be easier than updating full length

Spike based vaccines.

In contrast to RS2, the S2 component alone induces high IgG binding titers but sera failed to

neutralize the viruses[52,53]. Addition of S2-elicited sera significantly enhanced neutralization

potency of the RBD directed, broadly neutralizing S309 antibody **(Fig. 3G).** This finding further

supports the role of S2 in boosting immunogenicity and enhancing the protective efficacy of the

vaccine designs based on genetic fusion of RBD and S2 as in RS2 and S2R.

Similar to previous reports, our study reaffirms that despite lack of significant neutralization titers,

lung viral titers were significantly reduced in S2 immunized animals. This indicates that either T-

cell mediated or S2 directed antibodies, could offer protection through non-neutralizing
mechanisms, but these were not evaluated in the present study. In future, further characterization
of cellular immune responses and the effector functions of S2 directed antibodies will be evaluated,
to provide insights into the diverse mechanisms underlying the observed reduction in viral loads.

In conclusion, our study demonstrates that RS2 has several advantages compared to Spike, based
on its improved immunogenicity in small animals, higher thermal tolerance, and substantially
greater purified yield. Currently a significant proportion of the world's population has been
immunized with vaccines with full-length Spike immunogens, and likely an even larger number
of people have acquired natural immunity through infection. In this scenario an RS2 vaccine could
still be used as a booster vaccine in vulnerable groups that will require annual vaccination and
could also be combined with other vaccine formulations for respiratory viruses. Given that the
virus is likely to continue to circulate in humans for the foreseeable future, it is important to
continue developing and improving vaccine technologies both to combat this evolving virus and
for future preparedness against other coronavirus pandemics. The RS2 vaccine design can be used
as a prototype for designing vaccines that target other coronaviruses, or potential future variants
of SARS-CoV-2.

MATERIALS AND METHODS

Expression and Purification of recombinant proteins

Designed immunogens S2, RS2, S2R and Spike are based on the ancestral Wuhan SARS-CoV-2 strain (GenBank Id: YP_009724390.1). Mammalian codon optimized genes for S2, RS2 and S2R were synthesized at GenScript Inc. The designed immunogens consisted a HRV3C protease site (LEVLFQGP) at the C-terminus of proteins to facilitate histidine tag removal after purification. Recombinant proteins S2, RS2, S2R, RBD, and Spike were transiently expressed in Expi293F™ cells according to the manufacturer's guidelines (Gibco, ThermoFisher Scientific). Briefly, Expi293F™ cells were maintained at a cell density of ~3×10^6 viable cells/mL in the Expi293F™ expression medium. Plasmid DNA and ExpiFectamine™ 293 reagent were diluted with Opti-MEM™ I reduced serum media and incubated at room temperature for 5 min. After 15-20 min, ExpiFectamine™ 293/Plasmid DNA complexes were slowly added to Expi293F™ cells. Eighteen hours post-transfection, ExpiFectamine™ 293 Transfection Enhancer 1 and 2 were added to the transfected cells. Six days post-transfection, culture was harvested, and proteins were purified from culture supernatant by nickel affinity chromatography. The culture supernatant was incubated with Ni-Sepharose 6 Fast Flow resin (GE Healthcare) for 6-8 h at 4 °C. Non-specific proteins were removed by passing twenty-column volumes of wash buffer (PBS containing 25 mM Imidazole, pH 7.4). Bound proteins were eluted from the Ni-NTA column using 500 mM imidazole in PBS buffer, pH 7.4, and were dialyzed against PBS buffer using a 10 kDa (MWCO) dialysis membrane. The purity of purified protein samples was analyzed on SDS-PAGE.

Size Exclusion Chromatography (SEC)

A Superdex-200 10/300 analytical column, equilibrated with PBS buffer, pH 7.4, was used for size exclusion chromatography-multi angle light scattering. 100-200 µg of purified protein samples were injected into the column, and protein peaks were resolved on a BioRad NGC chromatography system at a flow rate of 0.4 mL/min.

Differential Scanning Fluorimetry (nano-DSF)

A Prometheus NT.48 instrument was used to determine the thermal stability of immunogens. The thermal unfolding of immunogens was monitored from 20 °C to 95 °C at a scan rate of 1 °C/min [59].

Trypsin Proteolysis

The proteolytic stability of designed immunogens was studied by performing trypsin proteolysis. S2, RS2, and S2R were dialyzed against 50 mM Tris buffer (pH 7.5) containing 1 mM $CaCl_2$, followed by incubation with TPCK trypsin (protease) in a 1:50 molar ratio at 4 °C and 37 °C. At different time points, aliquots were taken out, and the reaction was quenched using 6X-SDS-loading dye. Collected samples were analyzed on 12 % SDS-PAGE.

Surface Plasmon Resonance (SPR)

The binding affinity of the S2, RS2, S2R and Spike immunogens to different RBD and S2-specific antibodies were measured using ProteOn XPR36 Protein Interaction Array V.3.1. The SPR sensor prism HC200M chip (Xantec bioanalytics) was first activated with EDC and sulfo-NHS. Then Protein G (10 μg/mL) was immobilized in the presence of 10 mM sodium acetate buffer, pH 4.0. Finally, excess sulfo-NHS esters were quenched using 1 M ethanolamine. 500 response units (R.U.) of ACE2-hFc, RBD specific monoclonal antibodies: CR3022, S309, ADG-2, H014, and S2-specific monoclonal antibodies: B6 and CC40.8 were immobilized. Different protein sample concentrations (100 nM, 50 nM, 25 nM, 12.5 nM, and 6.25 nM) were passed over the HC200M chip surface at a flow rate of 30 μL/min, followed by dissociation with PBS buffer containing 0.05% tween-20 (PBST). After each kinetic-binding assay, the chip was regenerated using 0.1 M Glycine-HCl, pH 2.7. Proteon Manager was used for fitting data to a simple 1:1 Langmuir interaction model to obtain kinetic parameters.

Formulation preparation with SWE adjuvant

The adjuvant SWE (squalene-in-water emulsion) was co-developed by the Vaccine Formulation Institute (Switzerland) and Seppic (France) and is available at GMP grade (Sepivac SWE™) under an open access model. Immunogens (S2, S2R or RS2) were formulated with SWE at 1:1 volume ratio. The formulation RS2 in combination with SWE was evaluated for 1 month stability measuring adjuvant physicochemical characteristics and antigen integrity. Measurements included: visual inspection, particle size and polydispersity (DLS), zeta potential (ELS), pH, osmolality, and squalene content (HPLC). Antigen integrity was evaluated by ELISA.

Immunization Studies

(BALB/c mice immunizations): Groups of 6-8 weeks old, female BALB/c mice (n=5) were

intramuscularly immunized with either 2 μg or 20 μg dose of SWE adjuvant formulated RS2

immunogen.

(hACE-2 expressing C57BL/6 transgenic mice immunizations): Groups of hACE-2 expressing

transgenic mice (n=6/7) were intramuscularly immunized with 2 μg and/or 20 μg dose of SWE

adjuvant formulated immunogens (S2, RS2, S2R, Spike, and RBD). B6N; DBA2-Tg(K18-

hACE2)3068Mgef/Blisc (acrc:21000509) mice strain was generated and provided by the Mouse

Genome Engineering Facility, NCBS Bangalore using the K18-hACE2 transgene plasmid, kindly

donated by Dr Paul B. McCray [60]

(Syrian Hamster Immunizations): Groups of five female golden Syrian hamsters were

intramuscularly immunized with either a 5 μg or 20 μg dose of SWE adjuvant-formulated

immunogens (RS2, Spike, and RBD).

SWE adjuvant-treated animals were used as control. The sera samples from immunized and

unimmunized animals were collected before prime (Day -1), 2 weeks post-prime (Day 14), and 14

days post-boost (Day 35) for endpoint ELISA and measuring neutralizing antibody titers.

Challenge Studies: Twenty-one days following boost immunizations, immunized animals were

intranasally challenged with either 10^5 pfu MA10, 10^5 pfu/ 10^4 pfu Beta, or 10^4 pfu Delta VOCs.

The unimmunized challenged animals were used as control. Weight change of immunized-

challenged animals, unimmunized-unchallenged animals (Unimmunized), and unimmunized-

virus challenged control groups were monitored and recorded for 5-9 days. Post-challenge, lungs

were harvested for viral titer estimation and histopathological examination were obtained on day

6 except for the lyophilized RS2 study, where lung samples were obtained on day 10, as previously

described [21]. For lung tissue histopathology scoring, we developed a scientific method using

Mitchison's virulence scoring system with some modifications, considering the consolidation of

lungs, severity of bronchial and alveolar inflammation, immune cell influx, and alveolar and

perivascular edema [21,61]. The histopathology scores were graded as 0-4 (4: Severe pathology; 3:

Moderate pathology; 2: mild pathology; 1: Minor/minimum pathology; 0: No pathology).

ELISA and Competition ELISA

As described previously, endpoint titers of serum-binding antibodies were determined using

ELISA [22]. Briefly, 96 well ELISA plates were coated with 4 µg/mL RBD (332-532 a.a) or Spike

(1-1208 a.a) and incubated at 25 °C for 2-3 h. Plates were washed with PBST, followed by

incubation and with blocking with 3 % skimmed milk (in PBST) at 25 °C for 1 h. Four-fold serially

diluted antisera raised against immunogens were added to wells and incubated at 25 °C for 1 h.

Following three washes with PBST, plates were incubated with 1:5000 diluted goat ALP-

conjugated anti-mouse IgG secondary antibody at 25 °C for 1 h. Plates were washed thrice with

PBST and incubated with pNPP liquid substrate at 37 °C for 30 min. Optical density (O.D.) was

measured at 405 nm. The ELISA endpoint titers were determined as the highest sera dilution with

an O.D. signal above 0.2 at 405 nm. Competition ELISA was performed as previously described

[22]. Briefly, 96 well plates (HIMEDIA, Cat# EP2-5X10NO) were coated with Expi293 cell

produced RS2 at 4 µg/mL concentration in 1x PBS (60 µl/well) and incubated for 2 h at 25 °C

under gentle shaking condition (300 rpm) on a thermomixer (Eppendorf, USA) and then plate was

transferred to 4 °C cold room for overnight. Next day each well was washed with of 1xPBST

(200µl/well) and then treated with blocking solution (100 µL 3% skimmed milk in 1xPBST) for

45 min at 25 °C, 300 rpm. The sera isolated from 5 animals in a group were used. Individual sera

against the mentioned antigens were added at 2-fold serial dilution with a starting dilution of 1:25

in blocking solution (60µL). Only blocking solution was added to the control wells. The plates

were then incubated for 1 h at 25°C, 300 rpm. Plates were provided with 3 additional washes with

1xPBST (200 µL of 1xPBST/well). An additional blocking step was also performed for 45 min

with blocking solution (100µL) incubated at 25°C, 300rpm. An excess of monoclonal antibody

either S2X259, or B6 were added (60µL at 20µg/mL) to their respective wells and incubated for

one hour at 25°C, 300rpm. Next, three washes were given (200 µL of PBST/well) to remove excess

unbound proteins. 50 µl/well Goat Anti-Human IgG Antibody, Alkaline Phosphatase conjugate

(Sigma-Aldrich Cat # AP112A, Lot # 3519874; diluted 1:5000 in blocking buffer) was added and

samples incubated for 1 hour at 25°C, 300 rpm. Plates were washed 3 times with 200 µL of

PBST/well. Finally, 50 µL/well of a 37 °C prewarmed alkaline phosphatase yellow (pNPP) liquid

substrate (Sigma-Aldrich, Cat # P7998, Lot # SLCJ1764) was added, and plates were incubated

for 30 minutes at 37 °C, 300 rpm. The chromogenic signal was measured at 405 nm using an

ELISA plate reader (Emax-Plus Microplate reader, Molecular Devices).

The percent competition was calculated using the following equation.

$$472 \qquad \% \text{ Competition} = \frac{[\text{Absorbance control} - \text{Absorbance Sera dilution}]}{[\text{Absorbance control}]} \text{ X } 100$$

where**,** *Absorbance control* is the absorbance at 405nm of ACE2-hFc, S2X259 or B6

protein/Antibody binding directly to RS2 protein in the absence of sera**,** Absorbance *sera dilution*

is the absorbance where the serum dilution is incubated with competitor agents like S2X259 or B6

protein. The % competition as a function of serum dilution was fit using a three-parameter

nonlinear least squares fit curve using Graph Pad Prism 10.0. The sera dilution at 50 % competition

on the fitted curve was termed as the IC_{50} competition titer.

Pseudoviral neutralization assay

Psuedoviral neutralization assays were performed as previously described [21]. The genes encoding

Spike proteins from VOC were synthesized at GenScript (USA) except for the SARS-CoV-1

pseudovirus which was obtained from Dr Kalpana Luthra at the All India Institute of Medical

Sciences, New Delhi. Pseudovirus neutralization titers (ID_{50}) were defined as the serum dilution

at which the infectivity of the virus is reduced to 50%.

Statistical Analysis:

Data analysis was performed using GraphPad Prism software 10.0.0. The ELISA binding,

neutralization titers, and pseudoviral virus neutralization titer data were analyzed with a two-tailed

Mann–Whitney test and non-parametric Kruskal–Wallis with Dunn's multiple, respectively.

Weight changes in mice and hamsters were analyzed with a two-tailed Student's t-test. (* indicates

$p < 0.05$, ** indicates $p < 0.01$, *** indicates $p < 0.001$, **** indicates $p < 0.0001$).

All mice and Hamster immunization studies were approved by the Institutional Animal Ethics

Committee (CAF/ETHICS/847/2021; CAF/ETHICS/887/2022). These were carried out at the

Central Animal Facility (CAF), Indian Institute of Science, according to CPCSEA and ARRIVE

guidelines.

REFERENCES

1. Li, Q. *et al.* Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia. *New England Journal of Medicine* **382**, 1199–1207 (2020).

2. Xia, S. *et al.* Effect of an Inactivated Vaccine Against SARS-CoV-2 on Safety and Immunogenicity Outcomes: Interim Analysis of 2 Randomized Clinical Trials. *JAMA* **324**, 951–960 (2020).

3. Jackson, L. A. *et al.* An mRNA Vaccine against SARS-CoV-2 — Preliminary Report. *New England Journal of Medicine* **383**, 1920–1931 (2020).

4. Keech, C. *et al.* Phase 1–2 Trial of a SARS-CoV-2 Recombinant Spike Protein Nanoparticle Vaccine. *New England Journal of Medicine* **383**, 2320–2332 (2020).

5. Logunov, D. Y. *et al.* Safety and immunogenicity of an rAd26 and rAd5 vector-based heterologous prime-boost COVID-19 vaccine in two formulations: two open, non-randomised phase 1/2 studies from Russia. *Lancet* **396**, 887–897 (2020).

6. Zhang, Y. *et al.* Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine in healthy adults aged 18–59 years: a randomised, double-blind, placebo-controlled, phase 1/2 clinical trial. *Lancet Infect Dis* **21**, 181–192 (2021).

7. Zhu, F. C. *et al.* Safety, tolerability, and immunogenicity of a recombinant adenovirus type-5 vectored COVID-19 vaccine: a dose-escalation, open-label, non-randomised, first-in-human trial. *The Lancet* **395**, 1845–1854 (2020).

8. Walsh, E. E. *et al.* Safety and Immunogenicity of Two RNA-Based Covid-19 Vaccine Candidates. *New England Journal of Medicine* **383**, 2439–2450 (2020).

9. Walls, A. C. *et al.* Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* **181**, 281-292.e6 (2020).

10. Shang, J. *et al.* Cell entry mechanisms of SARS-CoV-2. *Proc Natl Acad Sci U S A* **117**, 11727–11734 (2020).

11. Wang, Q. *et al.* Structural and Functional Basis of SARS-CoV-2 Entry by Using Human ACE2. *Cell* **181**, 894-904.e9 (2020).

12. Barnes, C. O. *et al.* SARS-CoV-2 neutralizing antibody structures inform therapeutic strategies. *Nature* **588**, 682–687 (2020).

13.   Wu, Y. *et al.* A noncompeting pair of human neutralizing antibodies block COVID-19 virus
binding to its receptor ACE2. *Science* **368**, 1274–1278 (2020).

14.   Huo, J. *et al.* Neutralization of SARS-CoV-2 by Destruction of the Prefusion Spike. *Cell*
Host Microbe **28**, 445-454.e6 (2020).

15.   Pinto, D. *et al.* Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
antibody. *Nature* **583**, 290–295 (2020).

16.   Brouwer, P. J. M. *et al.* Potent neutralizing antibodies from COVID-19 patients define
multiple targets of vulnerability. *Science* **369**, 643–650 (2020).

17.   Wang, C. *et al.* A conserved immunogenic and vulnerable site on the coronavirus spike
protein delineated by cross-reactive monoclonal antibodies. *Nat Commun* **12**, 1–15 (2021).

18.   Rogers, T. F. *et al.* Isolation of potent SARS-CoV-2 neutralizing antibodies and protection
from disease in a small animal model. *Science* **369**, 956–963 (2020).

19.   Thuluva, S. *et al.* Evaluation of safety and immunogenicity of receptor-binding domain-
based COVID-19 vaccine (Corbevax) to select the optimum formulation in open-label,
multicentre, and randomised phase-1/2 and phase-2 clinical trials. *EBioMedicine* **83**,
104217 (2022).

20.   Malladi, S. K. *et al.* Design of a highly thermotolerant, immunogenic SARS-CoV-2 spike
fragment. *Journal of Biological Chemistry* **296**, 100025 (2021).

21.   Malladi, S. K. *et al.* Immunogenicity and Protective Efficacy of a Highly Thermotolerant,
Trimeric SARS-CoV-2 Receptor Binding Domain Derivative. *ACS Infect Dis* **7**, 2546–2564
(2021).

22.   Ahmed, S. *et al.* A Stabilized, Monomeric, Receptor Binding Domain Elicits High-Titer
Neutralizing Antibodies Against All SARS-CoV-2 Variants of Concern. *Front Immunol* **12**,
5214 (2021).

23.   Harvey, W. T. *et al.* SARS-CoV-2 variants, spike mutations and immune escape. *Nat Rev*
Microbiol **19**, 409–424 (2021).

24.   Ng, K. T., Mohd-Ismail, N. K. & Tan, Y.-J. Spike S2 Subunit: The Dark Horse in the Race
for Prophylactic and Therapeutic Interventions against SARS-CoV-2. *Vaccines (Basel)* **9**,
178 (2021).

25.   Sauer, M. M. *et al.* Structural basis for broad coronavirus neutralization. *Nat Struct Mol*
Biol **28**, 478–486 (2021).

26.   Wec, A. Z. *et al.* Broad neutralization of SARS-related viruses by human monoclonal
antibodies. *Science* **369**, 731–736 (2020).

27.   Song, G. *et al.* Cross-reactive serum and memory B-cell responses to spike protein in SARS-
CoV-2 and endemic coronavirus infection. *Nat Commun* **12**, 2938 (2021).

28.   Li, W. *et al.* Structural basis and mode of action for two broadly neutralizing antibodies
against SARS-CoV-2 emerging variants of concern. *Cell Rep* **38**, 110210 (2022).

29.   Hsieh, C.-L. *et al.* Stabilized coronavirus spike stem elicits a broadly protective antibody.
Cell Rep **37**, 109929 (2021).

30.   Zhou, P. *et al.* A human antibody reveals a conserved site on beta-coronavirus spike proteins
and confers protection against SARS-CoV-2 infection. *Sci Transl Med* **14**, eabi9215 (2022).

31.   Hurlburt, N. K. *et al.* Structural definition of a pan-sarbecovirus neutralizing epitope on the
spike S2 subunit. *Commun Biol* **5**, 342 (2022).

32.   Pinto, D. *et al.* Broad betacoronavirus neutralization by a stem helix-specific human
antibody. *Science* **373**, 1109–1116 (2021).

33.   Jennewein, M. F. *et al.* Isolation and characterization of cross-neutralizing coronavirus
antibodies from COVID-19+ subjects. *Cell Rep* **36**, 109353 (2021).

34.   Iketani, S. *et al.* Antibody evasion properties of SARS-CoV-2 Omicron sublineages. *Nature*
604, 553–556 (2022).

35.   Liu, L. *et al.* Striking antibody evasion manifested by the Omicron variant of SARS-CoV-
2. *Nature* **602**, 676–681 (2022).

36.   Wang, P. *et al.* Antibody resistance of SARS-CoV-2 variants B.1.351 and B.1.1.7. *Nature*
593, 130–135 (2021).

37.   Voss, W. N. *et al.* Prevalent, protective, and convergent IgG recognition of SARS-CoV-2
non-RBD spike epitopes. *Science* **372**, 1108–1112 (2021).

38.   Du, L., Yang, Y. & Zhang, X. Neutralizing antibodies for the prevention and treatment of
COVID-19. *Cell Mol Immunol* **18**, 2293–2306 (2021).

39.   Ahmed, S. *et al.* A Stabilized, Monomeric, Receptor Binding Domain Elicits High-Titer
Neutralizing Antibodies Against All SARS-CoV-2 Variants of Concern. *Front Immunol* **12**,
5214 (2021).

40.   Bommakanti, G. *et al.* Design of an HA2-based escherichia coli expressed influenza
586   immunogen that protects mice from pathogenic challenge. *Proc Natl Acad Sci U S A* **107**,
587   13701–13706 (2010).

588   41.   Mallajosyula, V. V. A. *et al.* Influenza hemagglutinin stem-fragment immunogen elicits
589   broadly neutralizing antibodies and confers heterologous protection. *Proc Natl Acad Sci U
590   S A* **111**, E2514–E2523 (2014).

591   42.   Pallesen, J. *et al.* Immunogenicity and structures of a rationally designed prefusion MERS-
592   CoV spike antigen. *Proc Natl Acad Sci U S A* **114**, E7348–E7357 (2017).

593   43.   Krarup, A. *et al.* A highly stable prefusion RSV F vaccine derived from structural analysis
594   of the fusion mechanism. *Nat Commun* **6**, 8143 (2015).

595   44.   Kirchdoerfer, R. N. *et al.* Stabilized coronavirus spikes are resistant to conformational
596   changes induced by receptor recognition or proteolysis. *Sci Rep* **8**, 15701 (2018).

597   45.   Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation.
598   *Science (1979)* **367**, 1260–1263 (2020).

599   46.   Hsieh, C.-L. *et al.* Structure-based design of prefusion-stabilized SARS-CoV-2 spikes.
600   *Science* **369**, 1501–1505 (2020).

601   47.   O'Hagan, D. T., Ott, G. S., Van Nest, G., Rappuoli, R. & Del Giudice, G. The history of
602   MF59(®) adjuvant: a phoenix that arose from the ashes. *Expert Rev Vaccines* **12**, 13–30
603   (2013).

604   48.   Walls, A. C. *et al.* Elicitation of broadly protective sarbecovirus immunity by receptor-
605   binding domain nanoparticle vaccines. *Cell* **184**, 5432-5447.e16 (2021).

606   49.   Cohen, A. A. *et al.* Mosaic RBD nanoparticles protect against challenge by diverse
607   sarbecoviruses in animal models. *Science* **377**, eabq0839 (2022).

608   50.   Cohen, A. A. *et al.* Mosaic nanoparticles elicit cross-reactive immune responses to zoonotic
609   coronaviruses in mice. *Science* **371**, 735–741 (2021).

610   51.   Ma, Y. *et al.* SARS-CoV-2 Spike Stem Protein Nanoparticles Elicited Broad ADCC and
611   Robust Neutralization against Variants in Mice. *Small* **18**, 2200836 (2022).

612   52.   Ravichandran, S. *et al.* Antibody signature induced by SARS-CoV-2 spike protein
613   immunogens in rabbits. *Sci Transl Med* **12**, 3539 (2020).

614   53.   Ng, K. W. *et al.* SARS-CoV-2 S2–targeted vaccination elicits broadly neutralizing
615   antibodies. *Sci Transl Med* **14**, 3715 (2022).

54. Weidenbacher, P. A.-B. *et al.* A ferritin-based COVID-19 nanoparticle vaccine that elicits robust, durable, broad-spectrum neutralizing antisera in non-human primates. *Nat Commun* **14**, 2149 (2023).

55. Wørzner, K. *et al.* Adjuvanted SARS-CoV-2 spike protein elicits neutralizing antibodies and CD4 T cell responses after a single immunization in mice. *EBioMedicine* **63**, 103197 (2021).

56. Christensen, D. *et al.* SARS-CoV-2 spike HexaPro formulated in aluminium hydroxide and administered in an accelerated vaccination schedule partially protects Syrian Hamsters against viral challenge despite low neutralizing antibody responses. *Front Immunol* **14**, 941281 (2023).

57. Merkuleva, I. A. *et al.* Are Hamsters a Suitable Model for Evaluating the Immunogenicity of RBD-Based Anti-COVID-19 Subunit Vaccines? *Viruses* **14**, (2022).

58. Sia, S. F. *et al.* Pathogenesis and transmission of SARS-CoV-2 in golden hamsters. *Nature* **583**, 834–838 (2020).

59. Chattopadhyay, G. & Varadarajan, R. Facile measurement of protein stability and folding kinetics using a nano differential scanning fluorimeter. *Protein Science* **28**, 1127–1134 (2019).

60. McCray, P. B. *et al.* Lethal Infection of K18- hACE2 Mice Infected with Severe Acute Respiratory Syndrome Coronavirus . *J Virol* **81**, 813–821 (2007).

61. Mitchison, D. A. *et al.* A comparison of the virulence in Guinea-pigs of south indian and British tubercle bacilli. *Tubercle* **41**, 1–22 (1960).

Acknowledgments: NM acknowledges the Prime Minister Research Fellowship for her fellowship (PM/MHRD-20-17303.03). RV is a JC Bose Fellow of DST. DC acknowledges the Council of Scientific and Industrial Research for his fellowship (09/079(2837)/2019-EMR-I). We acknowledge Ms. Simran Srivastava for providing the sarbecoviruses spike constructs.

Funding: This work was funded in part by a grant to RV from the Bill and Melinda Gates Foundation (INV-005948). Funding for infrastructural support was from DST FIST, UGC Centre for Advanced study, MHRD, and the DBT IISc Partnership Program. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Author contributions:

       Conceptualization: RV, NM

       Methodology: RV, NM, SK, RSR, RS, SBJ, MB, DC, SP, RPR, CL, VJ, PMD, AJ, SSSA

       Investigation: NM, SK, RSR, RS, SBJ, NJ, MB, DC, SP, RPR, CL, VJ,

       Visualization: NM, SK, RSR, RS, SBJ, NJ, MB, DC, SP, RPR, CL, VJ

       Funding acquisition: RV

       Project administration: RV

       Supervision: RV

       Writing – original draft: NM

       Writing – review & editing: All authors

Competing interests: A provisional patent application has been filed for the RS2 formulations described in this manuscript. RV, NM, and RS are inventors. RV is a co-founder of Mynvax, RS, SBJ, NJ, MB, and SP are employees of Mynvax Private Limited. Other authors declare that they have no competing interests.

Data and materials availability: All data are available in the main text or the supplementary materials.

TABLES

668　**Table 1:** Kinetic parameters of RS2, S2R, and Spike for binding to different RBD conformation-

669　specific ligands in PBS pH 7.4 at 25 ºC. ND*: No dissociation.

Ligand	Parameter	Immunogens		
		RS2	S2R	Spike
ACE2-hFc	k_a (M^{-1}s^{-1})	$2.30 \pm 0.01 \times 10^5$	$5.15 \pm 0.10 \times 10^5$	$1.07 \pm 0.01 \times 10^5$
	k_d (s^{-1})	$3.90 \pm 0.02 \times 10^{-4}$	$1.49 \pm 0.01 \times 10^{-3}$	$3.04 \pm 0.07 \times 10^{-5}$
	K_D (M)	$1.71 \pm 0.01 \times 10^{-9}$	$2.99 \pm 0.06 \times 10^{-9}$	$2.88 \pm 0.16 \times 10^{-10}$
CR3022	k_a (M^{-1}s^{-1})	$3.43 \pm 0.02 \times 10^5$	$6.57 \pm 0.07 \times 10^5$	$6.17 \pm 0.11 \times 10^4$
	k_d (s^{-1})	$4.13 \pm 0.03 \times 10^{-4}$	$1.39 \pm 0.01 \times 10^{-3}$	$2.45 \pm 0.05 \times 10^{-4}$
	K_D (M)	$1.21 \pm 0.01 \times 10^{-9}$	$2.15 \pm 0.03 \times 10^{-9}$	$4.06 \pm 0.11 \times 10^{-9}$
S309	k_a (M^{-1}s^{-1})	$5.78 \pm 0.08 \times 10^4$	$5.31 \pm 0.05 \times 10^4$	$7.61 \pm 0.13 \times 10^4$
	k_d (s^{-1})	$9.87 \pm 0.26 \times 10^{-5}$	ND*	ND*
	K_D (M)	$1.73 \pm 0.05 \times 10^{-9}$	ND*	ND*
ADG-2	k_a (M^{-1}s^{-1})	$2.17 \pm 0.01 \times 10^5$	$2.26 \pm 0.01 \times 10^5$	$1.36 \pm 0.01 \times 10^5$
	k_d (s^{-1})	$4.51 \pm 0.23 \times 10^{-5}$	ND*	$6.88 \pm 0.06 \times 10^{-5}$
	K_D (M)	$1.05 \pm 0.05 \times 10^{-10}$	ND*	$5.10 \pm 0.20 \times 10^{-10}$
H014	k_a (M^{-1}s^{-1})	$4.52 \pm 0.03 \times 10^5$	$7.82 \pm 0.05 \times 10^5$	$7.38 \pm 0.13 \times 10^5$
	k_d (s^{-1})	$3.49 \pm 0.03 \times 10^{-4}$	$1.64 \pm 0.01 \times 10^{-3}$	$1.66 \pm 0.05 \times 10^{-4}$
	K_D (M)	$7.78 \pm 0.08 \times 10^{-10}$	$2.14 \pm 0.03 \times 10^{-9}$	$2.30 \pm 0.08 \times 10^{-9}$

671   **Table 2:** Kinetic parameters of S2, RS2, S2R, and Spike for binding to different S2-specific ligands

672   in PBS pH 7.4 at 25 ºC. ND*: No dissociation.

Ligand	Parameter	Immunogens			
		RS2	**S2R**	**Spike**	**S2**
B6	k_a ($M^{-1}s^{-1}$)	$2.53 \pm 0.05 \times 10^5$	$1.24 \pm 0.01 \times 10^5$	$3.16 \pm 0.07 \times 10^4$	$2.04 \pm 0.07 \times 10^5$
	k_d (s^{-1})	$7.32 \pm 0.36 \times 10^{-4}$	$1.53 \pm 0.8 \times 10^{-3}$	$3.48 \pm 0.25 \times 10^{-5}$	$2.50 \pm 0.25 \times 10^{-4}$
	K_D (M)	$2.30 \pm 0.12 \times 10^{-9}$	$1.24 \pm 0.12 \times 10^{-8}$	$1.12 \pm 0.08 \times 10^{-9}$	$1.22 \pm 0.08 \times 10^{-9}$
CC40.8	k_a ($M^{-1}s^{-1}$)	$1.62 \pm 0.01 \times 10^5$	$1.62 \pm 0.02 \times 10^5$	$1.19 \pm 0.02 \times 10^5$	9.11×10^4
	k_d (s^{-1})	$6.77 \pm 0.01 \times 10^{-4}$	$5.30 \pm 1.6 \times 10^{-4}$	ND*	ND*
	K_D (M)	$4.22 \pm 0.08 \times 10^{-9}$	$3.23 \pm 0.03 \times 10^{-9}$	ND*	ND*

676 **Table 3:** Kinetic parameters for binding of lyophilized and resolubilized RS2 after incubation of
677 lyophilized protein at 37 °C for 1 month, to different RBD-specific and S2-specific ligands in PBS
678 pH 7.4 at 25 °C. ND*: No dissociation.

Ligand	k_a ($M^{-1}s^{-1}$)	k_d (s^{-1})	K_D (M)
ACE-2hFc	$1.2 \pm 0.01 \times 10^5$	$3.2 \pm 0.01 \times 10^{-4}$	$2.7 \pm 0.01 \times 10^{-9}$
CR3022	$4.8 \pm 0.01 \times 10^5$	$7.9 \pm 0.4 \times 10^{-4}$	$1.6 \pm 0.01 \times 10^{-9}$
S309	$1.2 \pm 0.01 \times 10^5$	$3.2 \pm 0.01 \times 10^{-5}$	$2.4 \pm 0.01 \times 10^{-10}$
H014	$5.0 \pm 0.01 \times 10^5$	$6.8 \pm 0.4 \times 10^{-4}$	$1.3 \pm 0.01 \times 10^{-9}$
B6	$1.1 \pm 0.01 \times 10^5$	$1.6 \pm 0.01 \times 10^{-5}$	$1.2 \pm 0.01 \times 10^{-10}$
CC40.8	$1.1 \pm 0.01 \times 10^5$	ND*	ND*

 Figure Legends

Fig. 1. **Characterization of S2, RS2, S2R, and Spike immunogens. (A)** Schematic
representation of designed immunogen sequences. **(B)** Reducing SDS-PAGE profile of protein
samples. **(C-F)** SEC profile of purified **(C)** S2, **(D)** RS2, **(E)** S2R and **(F)** Spike immunogens.

Fig. 2. Equilibrium thermal unfolding, transient thermal stability and limited trypsin
proteolytic profiles of designed immunogens. (A-D) Thermal unfolding profiles. Apparent
melting temperature of **(A)** RS2, **(B)** S2R, **(C)** S2 and **(D)** Spike were measured using Nano-DSF.
(E-H) Transient thermal stability profiles. Protein samples were subjected to different
temperatures (4, 37, and 50 °C) for one hour. Thermal stability of **(E)** RS2, **(F)** S2R, **(G)** S2 and
(H) Spike was monitored using Nano-DSF. Normalized first derivative of fluorescence at 350 nm
is plotted as function of temperature. **(I-L)** Proteolytic stability profile. Coomassie stained SDS-
PAGE profiles of purified **(I)** RS2, **(J)** S2R, **(K)** S2 and **(L)** Spike subjected to TPCK-Trypsin
proteolysis at 37 °C and 4 °C.

Fig 3. Immunogenicity of RBD, S2 and S2R in hACE-2 expressing mice. Three groups of
hACE-2 expressing transgenic mice were primed and boosted with 2μg of RBD, S2 and S2R
respectively, followed by an intranasal challenge with 10^5 pfu of the beta variant of SARS-CoV-
2. **(A-C)** ELISA endpoint titers against RBD, S2, and spike ectodomain respectively two weeks
post-boost. **(D and E)** Neutralizing antibody titers elicited by RBD, S2, and S2R against B.1 and
BA.1 Omicron SARS-CoV-2 pseudovirus. No neutralization was seen with S2 immunized
animals. **(F)** Neutralizing antibody titers elicited by S2R against various pseudoviruses. Lines
connect the neutralizing titers for different variants in a sera sample from an individual animal
against different variants. **(G)** Neutralization curves of pooled S2 immunized mice sera, MAb
S309, and S2 immunized mice sera in presence of S309. The sera sample was tested in five
technical repeats. Each point represents the median of five independent values. **(H)** Survival
Curve. **(I)** Average weight changes up to 5 days post-challenge. **(J)** Lung viral titer. **(K)**
Histopathology scores of lungs. **(L)** Histology of lung tissue sections from unimmunized-
unchallenged control (UC), unimmunized- Beta variant challenged control (Unimmunized-Beta)
and mice immunized with RBD, S2 and S2R at 4X magnification. Titers are shown as geometric

mean with geometric SD. The ELISA binding, neutralization titer, lung viral titer and histopathology score data were analyzed with a two-tailed Mann–Whitney test and non-parametric Kruskal-Wallis test with Dunn's multiple correction. Pairwise weight changes were analyzed with a Multiple Student's t-test with Bonferroni Dunn's correction method. (ns indicates non-significant, * indicates $p < 0.05$, ** indicates $p < 0.01$, **** indicates $p < 0.0001$).

Fig. 4. Immunogenicity of RS2 and Spike in BALB/c mice. BALB/c mice were immunized twice with either 2μg RS2 or 2μg Spike, followed by intranasal challenge with 10^5 pfu of MA-10 mouse adapted SARS-CoV-2. Two weeks following the boost, RBD, S2 and Spike specific IgG and neutralizing titers in immunized mice sera were measured **(A-C)** ELISA endpoint titers against RBD, S2 and spike ectodomain, respectively. **(D-K)** Comparison of neutralizing antibody titers elicited by 2μg of RS2 and Spike against **(D)** B.1, **(E)** Beta, **(F)** Delta, **(G)** BA.1, **(H)** BA.5, **(I)** BF.7, **(J)** WIV-1, and **(K)** SARS-CoV-1 pseudoviruses. **(L)** Average weight changes upto six days post-MA10 challenge. **(M)** Lung viral titers **(N)** Histopathology scores of lungs. **(O)** Histology of lung tissue sections from unimmunized-unchallenged control (UC), Unimmunized-MA10 virus challenged control (Unimmunized MA-10), mice immunized with Spike or RS2 at 4X magnification. Titers are shown as geometric mean with geometric SD. The ELISA binding, neutralization titer, lung viral titer, and histopathology score data were analyzed with a two-tailed Mann–Whitney test. Pairwise weight changes were analyzed with a Multiple Student's t-test with Bonferroni Dunn's correction method. (ns indicates non-significant, * indicates $p < 0.05$, ** indicates $p < 0.01$, **** indicates $p < 0.0001$).

Fig. 5. Stability of RS2. Characterization of lyophilized and resolubilized RS2 after incubation at 37 °C for 1 month. **(A-H)** Physiochemical characterization of RS2 in 1X PBS with equal amount (v/v) of SWE adjuvant incubated at 5 and 40 °C in polypropylene (PP) and glass vials (GV) for one month. Adjuvant properties were measured on day 0, day 7, day 14 and day 30. **(A)** Particle size, **(B)** Polydispersity Index, **(C)** Zeta Potential, **(D)** pH, **(E)** Osmolality, **(F)** Squalene content. **(G and H)** Antigenic integrity of RS2 in PBS and SWE adjuvant was measured based on binding to CR3022 using ELISA. **(I)** Day 7, **(J)** Day 14, **(K)** Day 30. Freshly thawed RS2 sample without any external modification 'RS2 ctrl extemp' was used as a control for undegraded antigen.

Fig. 6. Immunogenicity of lyophilized RS2, that had been incubated at 37 ℃ for 1 month, in hACE-2 expressing transgenic mice. hACE-2 expressing transgenic mice immunized twice with 20μg of lyophilized RS2 that was previously incubated at 37℃ for over a month and then formulated in SWE adjuvant. This was followed by intranasal challenge with 10^4 pfu Beta and Delta variants. **(A)** ELISA endpoint titers against RBD, S2, and spike ectodomain. **(B)** Neutralizing antibody titers against B.1, Beta, Delta and BA.1 pseudoviruses. **(C)** Neutralizing antibody titers elicited by lyophilized RS2 subjected to 37℃ for over a month (Lyo) and non-lyophilized RS2 (Non-lyo) stored at 4 ℃ against B.1 and BA.1 Omicron SARS-CoV-2 pseudovirus **(D and E)** Average weight change upto nine days post-Beta and Delta virus challenge respectively. **(F)** Survival curve. **(G and H)** Lung viral titers in RS2 immunized mice, challenged with Beta VOC and Delta VOC respectively. **(I and J)** Histopathology scores of lungs. **(K)** Histology of lung tissue sections from unimmunized-unchallenged control (UC), mice immunized with 20μg RS2 challenged with Beta variants (RS2-Beta challenged), unimmunized Delta virus challenged control (Unimmunized-Delta), mice immunized with 20μg RS2 and challenged with Delta variant (RS2-Delta challenged), at 4X magnification. None of the unimmunized controls survived the Beta virus challenge (Unimmunized-Beta). Titers are shown as geometric mean with geometric SD. The ELISA binding, neutralization titer, lung viral titer and histopathology score data were analyzed with a two-tailed Mann–Whitney test and non-parametric Kruskal-Wallis test with Dunn's multiple correction. Pairwise weight changes were analyzed with a Multiple Student's t-test with Bonferroni Dunn's correction method. (ns indicates non-significant, * indicates $p < 0.05$, ** indicates $p < 0.01$, **** indicates $p < 0.0001$).

Fig. 7. Comparative protective efficacy of RS2 and spike in hamsters. Syrian hamsters were immunized twice with 5μg of RS2 or Spike, followed by intranasal challenge with 10^5 pfu of the Beta VOC. **(A-C)** ELISA endpoint titers against RBD, S2 and spike ectodomain, respectively. **(D-K)** Comparative neutralizing antibody titers elicited by 5μg of RS2 and spike against **(D)** B.1, **(E)** Beta, **(F)** Delta, **(G)** BA.1, **(H)** BA.5, **(I)** BF.7, **(J)** WIV-1 and , **(K)** SARS-CoV-1 pseudovirus. **(L)** Average weight change upto five days post-beta variant virus challenge. **(M)** Lung viral titers **(N)** Histology of lung tissue sections from unimmunized-unchallenged control (UC), unimmunized-Beta challenged control (Unimmunized-Beta) and hamsters immunized with Spike and RS2 at 4X magnification. **(O)** Histopathology scores of lungs. The ELISA binding,

neutralization titer, lung viral titer and, histopathology score data were analyzed with a two-tailed

Mann–Whitney test. Pairwise weight changes were analyzed with a Multiple Student's t-test with

Bonferroni Dunn's correction method. (ns indicates non-significant, * indicates $p < 0.05$, **

indicates $p < 0.01$, **** indicates $p < 0.0001$).

Supplementary Figure Legends

Fig. S1. Comparative immunogenicity of RS2 and S2R in mice. BALB/c were mice immunized

twice with 20µg of RS2 or S2R. Two weeks following boost vaccination RBD and Spike specific

IgG levels and neutralizing titers were measured in immunized mice sera samples. Comparative

ELISA endpoint titers against **(A)** RBD and **(B)** Spike ectodomain, respectively **(C and D)**

Comparative neutralizing antibody titers elicited by 20µg of RS2 and S2R against **(C)** B.1, **(D)**

Delta variant pseudoviruses. Titers are shown as geometric mean with geometric SD. The ELISA

binding and neutralization titers were analyzed with a two-tailed Mann–Whitney test. (ns indicates

non-significant, * indicates $p < 0.05$, ** indicates $p < 0.01$, **** indicates $p < 0.0001$).

Fig. S2. Effect of transient thermal stress on lyophilized RS2. RS2 protein was dialyzed in

water and then lyophilized. **(A)** Equilibrium thermal unfolding profile of RS2 before and after

lyophilization and resolubilization in PBS. Lyophilized RS2 was incubated for one hour at

different temperatures (4, 37, 50, 70 and 90°C). Following reconstitution in PBS buffer, thermal

stability of **(B)** RS2 was monitored using Nano-DSF. **(C)** Thermal stability of lyophilized and

resolubilized RS2 after incubation at 37 °C for 1 month. **(D-H)** In addition, RS2 samples were

characterized for binding of with panel of RBD **(D-F)** and S2 **(G and H)** specific antibodies using

SPR **(D)** ACE2-hFc, **(E)** CR3022, **(F)** S309, **(G)** CC40.8, and **(H)** B6.

Fig. S3. Competition of immunized mice sera with RBD and S2 specific monoclonal

antibodies. BALB/c mice were immunized with 20µg of RBD, Spike, or RS2. Top panel:

Schematic of assay. Bottom panel **(A and B)** ELISA competition titers against **(A)** S2X259 (Class

4) and **(B)** B6 (S2 helix) monoclonal antibodies for individual, immunized mice sera. Immunogens

used to elicit respective sera are listed on the x-axis and competition titers on the y-axis of each

bar graph.

Fig. 1.

Fig. 2.

Fig. 3

Fig. 4.

Fig. 5.

Fig. 6.

Fig. 7.

946   **Fig. S1.**

959 **Fig. S2.**

(A)   **(B)**

Fig. S3.