Purkinje-Enriched snRNA-seq in SCA7 Cerebellum Reveals Zebrin Identity Loss as a Central Feature of Polyglutamine Ataxias

Luke C. Bartelt[1,2,3], Pawel M. Switonski[4], Juliana Carvalho[3], Lisa A. Duvick[5], Daniel R. Scoles[6], Stefan M. Pulst[6], Harry T. Orr[5], Court Hull[7], Craig B. Lowe[3] & Albert R. La Spada[2,8,9]

[1]University Program in Genetics & Genomics, Duke University School of Medicine, Durham, NC 27710, USA

[2]Departments of Pathology & Laboratory Medicine, Neurology, and Biological Chemistry, University of California, Irvine; Irvine, CA 92697, USA

[3] Department of Molecular Genetics and Microbiology, Duke University School of Medicine, Durham, NC 27710, USA

[4]Department of Medical Biotechnology, Institute of Bioorganic Chemistry, Polish Academy of Sciences, 61-704 Poznan, Poland

[5]Institute for Translational Neuroscience, and Department of Laboratory Medicine and Pathology, University of Minnesota, Minneapolis, MN 55455, USA

[6]Department of Neurology, University of Utah, Salt Lake City, UT 84132, USA

[7]Department of Neurobiology, Duke University School of Medicine, Durham, NC 27710, USA

[8]Department of Neurology, Duke University School of Medicine, Durham, NC 27710, USA

[9]UCI Center for Neurotherapeutics, University of California, Irvine; Irvine, CA 92697, USA

Co-corresponding authors:
Craig B. Lowe, PhD
Department of Molecular Genetics and Microbiology
Duke University School of Medicine
Durham, NC 27710
craig.lowe@duke.edu

Albert R. La Spada, MD, PhD
Pathology & Laboratory Medicine, Neurology, and Biological Chemistry
UCI Center for Neurotherapeutics
University of California Irvine
Irvine, CA, USA 92697
alaspada@uci.edu

ABSTRACT

Spinocerebellar ataxia type 7 (SCA7) is a dominantly inherited neurodegenerative disorder caused by a CAG-polyglutamine repeat expansion in the ataxin-7 gene. SCA7 patients display retinal degeneration and cerebellar degeneration with a striking loss of Purkinje cell neurons as the disease progresses. The normal function of the ataxin-7 protein is closely tied to transcriptional regulation, as ataxin-7 is a core component of the STAGA co-activator complex. Using a genetic SCA7 knock-in mouse model, we profiled the gene expression signatures associated with SCA7 neurodegeneration at single-cell resolution, employing a novel Purkinje cell-enrichment method, followed by single-nucleus RNA-seq. We identified altered expression of inhibitory synapse genes, which upon detailed analysis of SCA7 transcriptome changes stemmed from dramatic dysregulation of Purkinje Cell zebrin-II (aldolase-C) subtypes. Indeed, zebrin-II gene regulatory networks were significantly impacted in presymptomatic mice, resulting in an obliteration of zebrin-II parasagittal striping at symptomatic stages. We also detected dysregulation of zebrin-II gene modules in human SCA7 patient cerebellum, accompanied by blunted zebrin-II expression. When we profiled cerebellar zebrin-II regulation in other SCA polyglutamine disease mouse models, we similarly observed a disintegration of zebrin-II striping in symptomatic animals, indicating that zebrin-II identity dysregulation could be a general mechanism of polyglutamine-driven ataxia. We then documented increased inhibitory synapses in SCA7 cerebellum, and detected slower spontaneous spiking frequencies and altered inhibitory postsynaptic currents in cerebellar slices from SCA7 mice, confirming the functional significance of this pathological synapse reorganization.

INTRODUCTION

Neurodegenerative diseases pose a significant burden on our society. Alzheimer's disease, Parkinson's disease, and ALS represent complex neurodegenerative diseases with a variable contribution of genetic and environmental factors, complicating efforts to isolate molecular mechanisms driving disease pathogenesis. Strictly inherited neurodegenerative diseases with a monogenic basis are ideally suited for studies into pathogenic mechanisms, because all affected patients share a common genetic origin. The CAG-polyglutamine repeat diseases are an important category of neurodegenerative diseases that share key features with more common, sporadic neurodegenerative disorders and for which highly representative mouse models have been generated and characterized[1–3]. Spinocerebellar ataxia type 7 (SCA7) is one CAG-polyglutamine (polyQ) repeat expansion disease which has been successfully modeled in mice by knock-in of an expanded 266 CAG repeat into the endogenous mouse ataxin-7 locus. SCA7 266Q knock-in mice develop early onset, rapidly progressive cerebellar and retinal degeneration[1], akin to juvenile-onset SCA7 in human patients[4–6].

While mouse models have implicated a lengthy list of biological pathways in the SCAs, transcription dysregulation has emerged as a predominant pathogenic theme across the CAG-polyQ repeat diseases, such that these disorders are categorized as "transcriptionopathies"[7]. This shared impact on transcription is typically mediated via the nuclear accumulation of toxic polyQ aggregates, which can sequester important transcription regulatory proteins, such as TATA-binding protein and CREB-binding protein[8,9]. Transcriptional mechanisms are particularly relevant in SCA7, as the ataxin-7 protein performs its primary role in the nucleus as a core component of the STAGA transcription co-activator complex[10,11], which facilitates RNA Polymerase II-mediated transcription genome-wide[12]. The STAGA complex has several functional domains, including a histone acetyltransferase module that catalyzes the deposition of the activating H3K9ac mark and a

deubiquitinase (DUB) module that removes H2BK120ub modifications – tuning both the structural and transcriptional properties of target genes[13-15]. Ataxin-7 sits in the DUB module, and polyQ repeat expansion leads to structural impairments of the STAGA complex and alters global levels of both H3K9ac and H2BK120ub modifications, resulting in widespread transcription dysregulation[16-18]. Interestingly, the transcriptional signatures associated with SCA7 cerebellar degeneration significantly correlate with differential gene expression changes in the cerebellum of SCA1 and SCA2 mouse models, implying the existence of common transcriptional impacts by different polyQ-expanded ataxin proteins, despite non-overlapping normal functions[19]. The basis for highly shared patterns of gene expression alteration in the polyglutamine cerebellar degenerations remains unknown.

The cerebellum is a highly organized, yet complex region of the central nervous system (CNS) with over 15 distinct cell types coordinating together to maintain proper cerebellar function[20]. The action of these cells is organized around the cerebellar Purkinje cell (PC), which is the largest and most metabolically active neuron in the CNS[21,22], and serves as the sole output neuron of the cerebellar cortex. Purkinje cell dysfunction and neurodegeneration are central to the development of ataxic phenotypes in SCA7 mice and human patients[23]. However, numerous studies have also demonstrated the critical involvement of Bergmann glia and cerebellar astrocytes in SCA7 Purkinje cell degeneration, indicating that there is a non-cell autonomous contribution to SCA7 Purkinje cell degeneration[24,25]. Furthermore, Purkinje neurons exist as two sub-types: zebrin-II-positive and zebrin-II-negative PCs, organized along parasagittal stripes in the cerebellum, each with distinct gene expression signatures, synaptic organization, electrophysiology, and circuitry[26-29].

To determine the transcriptional basis of SCA7 cerebellar degeneration at the single-cell level, we pursued an extensive single-nucleus RNA-seq analysis of SCA7 266Q knock-in mice at the

presymptomatic stage and early symptomatic stage of disease. Although single-cell and single-nucleus sequencing technologies offer the ability to isolate transcriptional signals of diverse cell types in tissues such as the cerebellum, standard methods suffer from unique challenges in the cerebellum due to the massive overabundance of granule cell neurons. To overcome this issue, we developed a simple and robust Purkinje cell nuclear enrichment protocol for use in single-nucleus sequencing assays, yielding an enrichment of Purkinje cells from 1% to >40% of the captured cell population in adult mouse cerebellum. The results of our snRNA-seq analysis of SCA7 266Q knock-in mouse cerebellum revealed significant expression alterations involving synapse organization genes, which we validated by directly examining synapse distribution and circuit function in SCA7 mice. In addition, we detected a striking alteration of SCA7 PC zebrin-II subtype specification at the level of gene expression, and upon anatomical analysis, we discovered a dramatic obliteration of zebrin-II parasagittal striping in SCA7 knock-in mice. When we evaluated zebrin-II parasagittal striping patterns in related polyglutamine SCAs, we discovered that this surprising dysregulation of PC zebrin-II subtypes extends to other polyglutamine disease mouse models, suggesting that dysfunctional zebrin-II subtype maintenance is a defining feature of polyglutamine ataxias.

RESULTS

Purkinje cell-enriched snRNA-seq yields a high-resolution map of the SCA7 cerebellum

To enrich for PCs, which are central to degeneration of the cerebellum in SCA7 and related ataxias, we developed a nuclear enrichment protocol which utilizes a modified nuclei isolation buffer with increased concentration of K_2SO_4 to preferentially increase the proportion of PC nuclei in suspension (**Figure S1**). With this PC nuclear enrichment method, we then pursued a single-nucleus RNA-seq (snRNA-seq) analysis of cerebella from SCA7 266Q knock-in mice and wild-type (WT) littermate controls (**Figure 1A**), at two successive timepoints: 5 weeks of age (presymptomatic) and 8.5 weeks of age (early symptomatic). To minimize batch effects and account for any sex differences, 2 male and 2 female mice for each genotype were barcoded and multiplexed with MULTI-seq cholesterol-modified oligonucleotides[30], and run in a single 10X GEM well. After quality control, we recovered 5334 and 8820 nuclear profiles from the 5-week and 8.5-week timepoints respectively, which were assigned to their animal of origin by MULTI-seq barcode counts. UMAP clustering separated nuclei into known resident cell types of the cerebellum with a significant enrichment of Purkinje cells to >40% of recovered nuclei (**Figure 1B**). Nuclei originating from WT and SCA7 mice displayed high levels of concordance in 5 week-old mice with relatively few Differentially Expressed Genes (DEGs)(**Supplementary Table 1**). However, at 8.5 weeks of age we identified dramatic transcriptional differences between WT and SCA7 mice, visually represented by the genotypes separately clustering in UMAP space and quantified by large numbers of DEGs in PCs, oligodendrocytes, Bergmann glia, and astrocytes (**Figure 1C-D; Supplementary Table 2**). Immediately apparent in the differential expression was a bias toward downregulation of genes across SCA7 cell types (**Figure 1E-F**). Furthermore, due to our controlled multiplexed experimental design, we were able to assess differential RNA abundance between SCA7 and WT animals. We observed a cell-type selective decrease in nuclear RNA abundance in affected cell-types, most apparent in PCs at both the 5-week and 8.5-week timepoints (**Figure S2**). Given

the known role of the STAGA complex in transcription initiation and the impact of nuclear polyQ

aggregates on gene regulatory complexes, proper differential expression testing is warranted for

analysis of this single-nucleus dataset, as standard log-normalization of gene counts may yield

misleading results in the context of global transcriptional effects[31]. Therefore, we performed

pseudobulk comparisons of gene counts between SCA7 and WT cell populations with DEseq2,

removing the standard assumption that total RNA abundance stays constant across groups being

compared[32,33].

Analysis of snRNA-seq DEGs confirms glial contribution to SCA7 due to loss of glial identity

Given the known contribution of *Gfa2*-lineage cell types to the non-autonomous Purkinje cell

degeneration in SCA7, we first wanted to investigate differential expression results for signs of Glial

dysfunction[24]. Indeed, across the three most frequent glial cell types we noticed a clear separation

of WT and SCA7 UMAP sub-clusters (**Figure 2A**) with a striking downregulation of key glial identity

genes represented among SCA7 DEGs: oligodendrocytes (*Mobp*, *Qk*, *Plp1*), Bergmann glia (*Sparcl1*),

and astrocytes (*Aqp4*, *Sparcl1*, *Slc1a2* [EAAT2]) (**Figure 2B**). Further analysis suggested that the

coordinate downregulation of glial identity genes in SCA7 cell clusters may stem from significantly

decreased RNA expression of the *Qk* gene, which encodes the Quaking RNA binding protein (QKI).

Loss of function of *Qk* is the genetic basis of the *quaking* mutant mouse, which shows severe

myelination defects leading to ataxia, tremor, and early death[34,35]. QKI is a critical regulator of glial

maturation and function through binding and stabilization of glial-specific transcripts[36]. Indeed,

SCA7 DEGs from glial clusters show a highly significant overlap with QKI target mRNAs recently

discovered by CLIP-seq[37] (**Figure 2C**). Our snRNA-seq results thus corroborate previous reports of

a glial contribution to SCA7[24], and provide at least one mechanistic hypothesis for the glial

dysfunction in SCA7 through the impaired activity of QKI.

Purkinje cell DEGs highlight altered inhibitory synapse organization in SCA7 cerebellum

When we considered the full snRNA-seq dataset, we observed that PCs contained the strongest transcriptional changes, with 145 DEGs in presymptomatic and 273 DEGs in symptomatic animals. Sub-setting and re-clustering of PC nuclei revealed a dramatic progression of SCA7 transcriptional dysregulation from 5 weeks to 8.5 weeks of age (**Figure 3A-D**), corresponding to the timing of motor symptom onset. To gain further insight into the biological impact of the progressively dysregulated SCA7 transcriptome, we performed Gene Ontology (GO) analysis to categorize the Purkinje cell DEGs by their biological process. GO analysis revealed a remarkable enrichment of genes involved with Synaptic Signaling and Synapse Organization (**Figure 3E**). Many cell-adhesion molecules were represented in this biological category (NRXN1, NLGN1, CNTN5, GRID2IP, HOMER3), suggesting that SCA7 PCs were remodeling synaptic inputs with surrounding cells upon disease onset (**Supplementary Table 2**). However, upon deeper analysis we determined that this enrichment of synaptic organization genes may be driven by a broader interference to PC subtype regulation.

The transcriptional differentiation of zebrin-II subtypes is severely diminished in the cerebellum of SCA7 mice

It has long been appreciated that the cerebellum of higher organisms, ranging from bony fish to mammals, exhibits an alternating pattern of gene expression of aldolase-C (*Aldoc)*, which upon immunostaining yields a distinct striping pattern, accounting for why aldolase-C came to also be known as 'zebrin-II'[27,38]. Numerous studies of the cerebellum have established that this gene expression patterning of zebrin-II positive PCs and zebrin-II negative PCs organizes the cerebellum into distinct functional zones, with divergent cellular morphology, electrophysiological properties, and circuitry[28,39]. When one group recently compared the transcriptomes of zebrin-II positive and

negative PCs in mouse cerebellum, they demonstrated that zebrin-negative PCs are specialized for synaptic plasticity during motor learning[40].

To determine if the transcriptional dysregulation occurring in SCA7 PC degeneration affects zebrin-II transcriptional signatures, we investigated sub-clusters of PC nuclei from SCA7 knock-in mice and WT mice in our snRNA-seq gene expression data. UMAP dimensional reduction at the early symptomatic time point displayed a highly divergent axis of SCA7 nuclear profiles that strongly correlated with zebrin-II expression, and revealed a marked decrease in the fraction of zebrin-II positive PCs (**Figure 4A**). To better assess the status of zebrin-II transcriptional signatures in SCA7 PCs, we initially sought to define a set of zebrin-II identity genes from WT PCs alone. Sub-clustering of WT PCs revealed a clear demarcation of the zebrin-II transcriptional signature, with UMAP clusters separating zebrin-II positive PCs from zebrin-II negative PCs (**Figure 4B**). When we performed differential expression analysis of the WT Zebrin-II PC subtypes, we identified 354 significant DEGs that accounted for the demarcation of zebrin-II positive and zebrin-II negative PCs in WT mouse cerebellum (**Figure 4C**). Strikingly, these WT zebrin-II axis genes showed highly significant overlap with SCA7 PC DEGs ($P < 1 \times 10^{-50}$), with 30% of SCA7 PC DEGs now identifiable as zebrin-II identity genes (**Figure 4D**). Treating this 'Zebrin-II axis' list as a biological category and then comparing FDR enrichment to the GO terms assessed in **Figure 3E** confirmed that the impact of SCA7 transcription dysregulation to the 'Zebrin-II-axis' genes far exceeds the highly significant effect on the Synapse Organization genes (**Figure 4E**).

We then assessed the transcriptional dysregulation of zebrin-II PC subtypes in SCA7 by sub-clustering SCA7 PCs alone and computing differential expression between clusters of zebrin-II positive and zebrin-II negative PCs. The SCA7 PC zebrin-II-axis showed significant abnormalities when compared to WT PC zebrin-II-axis in both presymptomatic and symptomatic animals. Indeed,

in 5 week-old SCA7 mice, the zebrin-II-axis gene expression signatures already display a clear trend toward the direction of zebrin identity loss, with log-fold expression of zebrin-II identity genes dampened in both the zebrin-II positive and zebrin-II negative PC subtypes (**Figure 4F**). When we plotted the log-fold expression values of zebrin-II axis genes for WT vs. SCA7 PCs and performed linear regression to quantify this trend, we calculated a decreased slope (m = 0.774) which diverges a line plotted with a slope (m = 1.0) for a theoretically perfect correlation (**Figure 4F**). This gene expression signature in presymptomatic SCA7 mice indicates that the presence of polyQ-expanded ataxin-7 in PC nuclei is likely interfering with the regulatory mechanisms that specify zebrin-II PC subtypes prior to the appearance of visible nuclear aggregates. Symptomatic 8.5 week-old SCA7 mice display progressive dysregulation of the zebrin-II axis, with the emergence of numerous significantly up- and down-regulated zebrin-II-identity genes, with even greater disruption of zebrin-II identity (**Figure 4G**).

Disruption of zebrin-II striping in symptomatic SCA7 mice and in related polyQ SCAs

Given the marked downregulation of the canonical zebrin-II marker gene *Aldoc* in the PCs of symptomatic SCA7 mice (**Figure 4C**), we hypothesized that the stereotypical parasagittal striping pattern of the cerebellum might be affected. To investigate the spatial expression pattern of zebrin-II, we performed immunohistochemistry on coronal cerebellar sections obtained from SCA7 266Q knock-in mice and WT littermates. Although immunostaining of zebrin-II in 5 week-old presymptomatic SCA7 mice revealed normal patterning with robust striped expression in the posterior lobules, we documented a dramatic, near-complete loss of the zebrin-II striping pattern in 8.5 week-old SCA7 mice, despite limited evidence of PC degeneration based upon robust calbindin-1 expression (**Figure 5A**). To quantify this phenotype, we developed an image analysis pipeline to compute the amplitude of zebrin-II stripe expression across lobule VIII of the cerebellum, and

verified that zebrin-II striping integrity is virtually absent by this time point in SCA7 mice (**Figure 5B**).

A number of studies have noted striking similarities between the transcriptional alterations in the cerebellum of various polyQ SCAs, correlating findings from SCA1, SCA2, and SCA7[41,42]. To determine if this transcriptional pathology includes the normal developmentally programmed parasagittal striping of the cerebellum, we examined zebrin-II spatial expression in the cerebellum of SCA1 175Q knock-in mice and SCA2 127Q transgenic mice, two extensively studied mouse models believed to be representative of their cognate human diseases[2,3]. Zebrin-II immunostaining of coronal cerebellar sections similarly revealed an obliteration of the normal parasagittal striping pattern in both early symptomatic SCA1 and SCA2 mice (**Figure 5B-C**). To determine if this phenomenon might also occur in Huntington's disease (HD), which is characterized by high-level cerebellar expression of polyQ protein – but without cerebellar degeneration, we then performed zebrin-II immunostaining of coronal cerebellar sections from a well-known HD transgenic mouse model – N171-82Q[43], and remarkably we observed a perfectly normal parasagittal striping pattern in the cerebellum of early symptomatic HD N171-82Q mice (**Figure 5B-C**). These results indicate that disrupted zebrin-II identity is a shared feature of polyQ SCAs.

Human SCA7 Cerebellum displays Zebrin-II Dysregulation in Molecular Layer Interneurons
While the presence of disrupted zebrin-II expression patterning in SCA7 model mice is a compelling finding, an important question is whether this phenomenon is relevant to the disease process in human patients. To address this issue, we performed multiplexed snRNA-seq on post-mortem cerebellum samples obtained from five SCA7 patients and four unaffected controls, and applied our Purkinje-enriched nuclear isolation method and MULTI-seq barcoding as described earlier. When we analyzed the human snRNA-seq data using a standard Seurat normalization and clustering

pipeline, we were unable to detect any PC nuclei based upon cell type identification of UMAP clusters by marker gene expression (**Figure 6A**). As it turns out, this was not unexpected, as several studies of post-mortem human cerebellum have encountered this identical technical issue, and have failed to recover PC nuclei to date[44,45]. However, when we reviewed the UMAP clustering of our human cerebellum snRNA-seq results, we noted significant enrichment of both Type-I and Type-II Molecular Layer Interneurons (MLI), which revealed significant gene expression alterations between SCA7 patients and unaffected controls (**Figure 6B**). Similar to our observation in SCA7-266Q mice, SCA7 human tissues showed a dramatic cell type selective decrease in RNA abundance, with outsized impacts on MLIs (**Supplemental Figure S4**). Of the 836 DEGs in MLI-1 cells, *Aldoc* ranked as the 13th strongest downregulated gene, confirming that marked dysregulation of zebrin-II gene expression is a prominent feature of the disease process in the cerebellum of human SCA7 patients (**Figure 6C-D**).

Our detection of altered zebrin-II expression in human SCA7 MLIs raises a provocative question: Are MLIs also subject to a developmental program of alternating gene expression linked to PC parasagittal striping? Indeed, a number of previous studies have suggested that both the morphological and electrophysiological properties of MLIs appear bounded to parasagittal zones, generating functional modular circuits with zebrin-II positive and negative PCs[46–48]. To definitively evaluate the gene expression properties of MLIs, we reanalyzed MLIs from the Kozareva et.al. Transcriptomic Mouse Cerebellar Atlas to determine if zebrin-II and its co-regulated genes display a gene expression gradient akin to PCs. This analysis revealed that both MLI-1 nuclei and MLI-2 nuclei from the Mouse Cerebellar Atlas form robust subclusters with a detectable zebrin-II gene expression gradient (**Figure 6E, Supplemental Figure S5**). Using subclusters with 'high' and 'low' expression of *Aldoc*, we performed differential expression analysis to delineate the entire list of genes defining this gradient. This analysis yielded 1,347 significant DEGs for MLI-1 and 1,437

significant DEGs for MLI-2 nuclei, with strongly overlapping gene sets that form a gene expression

gradient correlated with *Aldoc* (**Figure 6E, Supplemental Figure S5**). To evaluate the relationship

between the MLI gene expression changes detected in human SCA7 patients and MLI gene

expression patterning detected in mouse cerebellum, we compared the human SCA7 MLI-1 DEGs

with the DEGs defining mouse MLI-1 Zebrin subclusters, and we noted an extremely high degree of

enrichment ($P = 1.02 \times 10^{-147}$), indicating that MLI-1 patterning genes are disproportionally

impacted in SCA7 patients (**Figure 6F**). Interestingly, the 4th strongest DEG defining mouse MLI

subclusters is *Aldolase-a*, whose expression resembles the *Aldoc* expression patterning observed in

PCs. To assess the biological relevance of this finding, we immunostained mouse cerebellar

sections for *Aldolase*-a and confirmed the existence of an *Aldolase-a* striping pattern that clearly

parallels the alternating expression pattern of zebrin-II (**Figure 6G**). Finally, to examine the

expression level of PC zebrin-II identity genes in human SCA7 patient postmortem cerebellum, we

pursued qRT-PCR analysis on bulk cerebellar RNA (in lieu of PCs in single-cell RNA sequencing),

and we confirmed significant downregulation of PC zebrin-II identity genes in SCA7 patients

(**Figure 6H**).

PC inhibitory synapse activity is increased in the cerebellum of SCA7 model mice

Although changes in the expression of zebrin-II subtype genes and the resulting disappearance of

zebrin-II parasagittal stripes in SCA7 model mice are remarkable, an important question is in how

these findings alter cerebellar function and thereby contribute to SCA7 cerebellar degeneration.

Since the Gene Ontology Biological Function categories most dysregulated in SCA7 PCs are

overwhelmingly related to Synapse Organization (**Figure 3E**), and Zebrin-II subtypes are known to

differ in synaptic functions and electrophysiology[28], we hypothesized that the breakdown of Zebrin

subtype maintenance would have downstream consequences on PC synapses. To address this

hypothesis, we assessed PC synapse organization by performing quantitative

immunohistochemistry of both excitatory and inhibitory synapses in the cerebellar molecular layer. Purkinje cell dendrites receive synaptic input through three distinct synapses marked by the presynaptic vesicular proteins VGAT (inhibitory)[49], VGLUT1 (excitatory)[50], and VGLUT2 (excitatory)[51]. VGLUT1 marks excitatory synapses originating from the highly dense parallel fibers, which emanate from granule cell neurons and elicit high frequency simple spikes in PCs[52,53], while VGLUT2 marks excitatory synapses formed by the climbing fibers, which originate from the inferior olive and show a tight 1:1 relationship with PCs[54]. Climbing fiber synapses operate at a lower frequency and generate the complex spikes in PCs[55]. We stained WT and SCA7 cerebella with antibodies against VGLUT2 paired with calbindin-1 to mark PC dendrites and quantified synaptic puncta in lobule VII of the cerebellar vermis. At both the presymptomatic and symptomatic stages, climbing fiber synapses in SCA7 mice appeared normal, with VGLUT2 synaptic density and innervation along the length of the PC dendrite matching what is seen in WT mice (**Supplemental Figure S6A-D**), in agreement with a previous report[42]. To profile inhibitory synapses, we stained for presynaptic VGAT and postsynaptic Gephyrin proteins, and performed calbindin-1 co-staining to differentiate inhibitory synapses on PC dendrites from synapses on the surrounding molecular layer inhibitory interneurons (MLI). We observed a marked increase in the number of inhibitory synapses – both on PC dendrites and on surrounding MLI – in 8.5 week-old symptomatic SCA7 mice (**Figure 7A-B**). In 5 week-old pre-symptomatic SCA7 mice, immunostaining quantification indicated a similar number of both PC and non-PC inhibitory synapses when compared to 5 week-old WT littermate controls, though there was a trend toward increased numbers of inhibitory synapses in SCA7 cerebellum (**Figure 7C-D**). These findings confirm that PC gene expression changes observed by snRNA-Seq result in neuroanatomical alterations in cerebellum synapse organization in SCA7 model mice.

To determine the functional consequences of altered inhibitory synapse organization in symptomatic SCA7 mice, we directly examined Purkinje neuron electrical activity in acute cerebellar slices. Unlike previous studies that assessed the intrinsic pace-making activity of PCs and documented evidence of irregular spiking[42,56], here we focused on local circuits upstream of PC firing by performing whole-cell voltage clamp recordings to measure spontaneous inhibitory post-synaptic currents (sIPSCs). Analysis of sIPSCs uncovered significantly increased inhibitory synaptic amplitudes in SCA7 PCs from 8.5 week-old mice (**Figure 8A**), with no change in firing frequency compared to WT littermates. Excitatory post-synaptic currents (sEPSCs) in SCA7 PCs displayed normal amplitudes and decreased sEPSC frequency (**Figure 8B**). We then collected cell-attached recordings of PC action potentials and quantified the spiking rate and regularity through inter-event-intervals. SCA7 PCs displayed a significant decrease in spiking rates (**Figure 8C**), reflecting the shift in excitatory/inhibitory synaptic balance on the PC dendrites.

DISCUSSION

We, and others, have identified transcription dysregulation as a central feature of SCA7 disease pathogenesis, stemming from the realization that ataxin-7 is a core component of the STAGA co-activator complex, and that mutant polyQ-ataxin-7 protein gets incorporated into the STAGA complex and therein disrupts the function of transcription factors that depend upon STAGA for transactivation of their target genes[10,17]. We previously performed bulk RNA-seq analysis on cerebellum from presymptomatic and early symptomatic SCA7 mice, and documented a role for altered calcium regulation in SCA7 cerebellar degeneration, linking the calcium dysregulation to dysfunction of Sirtuin-1, a master regulator of gene expression and cellular function[56]. While this work yielded key insights into the molecular basis of SCA7, there are two major limitations in the biological interpretation of such bulk RNA-seq datasets: 1) RNA reads cannot be assigned to a particular cell type, and 2) the cerebellum is dominated by the abundant granule cell neurons (GCNs), as GCNs comprise >90% of cells in the cerebellum and thereby dilute signals from non-GCN cell types. As Purkinje cell (PC) neurons integrate neural circuit information to provide critical output to the deep cerebellar nuclei and thence to the entire motor system of the brain, and PCs undergo degeneration in SCA7, evaluation of PC gene expression is crucial to obtain a thorough understanding of the SCA7 disease process. Here we introduced use of a novel PC nuclear enrichment protocol, and successfully applied it to perform snRNA-seq to interrogate cerebellar degeneration in SCA7 266Q knock-in mice. By evaluating both presymptomatic and early symptomatic SCA7 mice in comparison to age- and sex-matched WT littermate controls, we discovered alterations in PC gene expression which revealed changes in synapse organization and developmental patterning that likely underlie SCA7 disease pathogenesis.

The cerebellum is a complex brain structure that plays a critical role in motor coordination and learning, functionally organized into parasagittal modules marked by zebrin-II. While zebrin-II

expression is typically appreciated as a marker of PC subtypes, these functional modules extend to the surrounding MLIs, which we have now characterized by the differential expression of over 1000 genes. While the significance of zebrin-II parasagittal expression striping is still not fully understood, it has been implicated in several functions, including the control of motor learning, regulation of synaptic plasticity, and organization of cerebellar circuitry[29,57]. The development of the cerebellum is an incredibly refined and meticulous process that extends well into the postnatal period before completion in humans and mice. PCs can be initially recognized in the developing mouse brain at embryonic day 10.5 (E10.5), with a second layer of PCs emerging at E12.5[58–61]. Based on single-cell studies, at least 4 "embryonic" clusters of PCs can be identified in the developing cerebellum with a specified set of marker genes unique to this developmental window[62]. Although SCA7 mice retain parasagittal striping of the cerebellum at 5 weeks of age, evidence for gene expression alterations in zebrin identity genes are already present at this time point (**Figure 4**). These findings indicate that altered development of the cerebellum is a defining feature of SCA7 disease, and thus contribute to an emerging view of polyglutamine neurodegeneration where altered development is critical for setting the stage for later disease pathogenesis. For example, in SCA1, the timing of onset of mutant ataxin-1 gene expression is a powerful determinant of Purkinje cell degeneration, as delaying the expression of polyQ-ataxin-1 from postnatal day 2 (P2) to postnatal day 10 (P10) markedly protects against later cerebellar degeneration, even if much higher levels of polyQ-ataxin-1 protein expression occur starting at P10[63]. Furthermore, recent studies of Huntington's disease (HD), another CAG-polyQ disorder, indicate that defects in the development of the cortex can be detected in human patients and model mice, underscoring a likely role for altered neurodevelopment in HD pathogenesis[64].

To determine if disintegration of zebrin-II expression patterning is restricted to SCA7 or is a feature of other polyglutamine diseases, we immunostained cerebellar sections from two closely related

polyQ SCA mouse models and a representative HD mouse model, and we documented disruption of

parasagittal striping in the cerebellum of early symptomatic SCA1 and SCA2 mice, but observed

normal preservation of parasagittal striping in the cerebellum of HD mice. Hence, it appears that

the effect of polyQ disease proteins on zebrin-II parasagittal striping correlates with the cerebellar

disease process, as HD mice do not display cerebellar degeneration in this brain region[65]. Previous

studies have emphasized that SCA1, SCA2, and SCA7 display considerable overlap in transcriptional

alterations in the degenerating cerebellum in highly representative mouse models[42,66]. It is

therefore tempting to speculate that the disruption of zebrin-II identity could be at the core of the

transcriptional pathology underlying these polyQ SCAs. Dysregulation of zebrin identity genes in

multiple polyQ SCAs suggests a common mechanism of disease pathogenesis and could serve as a

potential therapeutic target for the treatment of these disorders. Future studies are needed to map

the extent of overlap between altered zebrin-II identity genes in the various SCAs and determine

the mechanistic basis for these shared abnormalities. Overall, these findings provide novel insights

into the molecular mechanisms underlying polyQ ataxias and could have important implications for

the development of new therapies for these devastating diseases.

To determine if the zebrin-II dysregulation observed in SCA7 mice is relevant to the disease process

in human patients, we obtained post-mortem cerebellum from five SCA7 patients and four

unaffected human controls, and performed snRNA-seq on nuclei isolated from these samples.

Although we could not obtain PCs, which is the typical experience based upon all other published

reports to date[44,45], we did detect MLIs and noted prominent zebrin-II dysregulation in SCA7

patients, which led us to evaluate MLIs and document alternating expression patterns in this

cerebellar cell type. When we directly measured the expression levels of zebrin-II identity genes in

bulk RNA samples obtained from SCA7 and control human cerebella, we observed decreased

expression of a subset of these marker genes in SCA7 patients. Together, these results from SCA7

human cerebellar tissues further support the notion that dysregulated zebrin identity could be one of the foremost driving pathological transcriptional mechanisms leading to neurodegeneration and neuron dysfunction in SCA7, and therefore could be considered a new pathological hallmark of polyQ SCAs.

One of the most striking observations emerging from the SCA7 mouse snRNA-seq dataset was a dramatic upregulation in the expression of cell adhesion molecules involved in inhibitory synapse formation. To determine if increased expression of synapse organization genes in SCA7 PCs was paralleled by functional changes of biological relevance, we sought to quantify synapses and examine synapse activity in SCA7 cerebellum. We documented that early symptomatic SCA7 mice display markedly increased numbers of inhibitory synapses in the molecular layer of the cerebellum, accompanied by significantly increased inhibitory synaptic amplitudes in PCs upon cerebellar slice electrophysiology. These findings indicate a dramatic process of synapse reorganization in the degenerating cerebellum in SCA7. Whether this increase in inhibitory synapse function represents an attempt at compensation to alleviate excessive aberrant electrical activity or is driving the disease process in the cerebellum remains to be determined. However, there is reason to expect that increased inhibitory synapse formation resulting in excessive inhibitory synaptic input to PCs could be deleterious. In the ax^J mouse, loss of function of ubiquitin-specific protease 14 results in increased levels of the GABA(A) receptor on PCs and correspondingly increased inhibitory postsynaptic currents (IPSCs), which may contribute to cerebellar dysfunction and degeneration[67]. Similarly, GABA transporter subtype 1 knock-out mice develop ataxia due to impaired uptake of GABA in the synaptic cleft, resulting in slower decay of IPSCs at both PC and cerebellar granule neuron synapses[68]. Although these findings suggest that increased inhibitory synapse numbers and inhibitory interneuron synaptic input in SCA7 PCs is part of the disease process, it will be necessary to modulate GABA(A) receptor activity in SCA7 mice using genetic and

pharmacological interventions to determine if this alteration is compensatory or contributory to

disease. Differentiating between these two scenarios will dictate whether an effective treatment

strategy for SCA7 cerebellar degeneration will be to block or to enhance inhibitory synaptic

transmission, and will have broader application to related polyQ SCAs.

MATERIALS & METHODS

Animals

SCA7-266Q mice (B6.129S7-Atxn7[tm1Hzo]/J ; JAX#:008682) were maintained on a C57BL/6J background at Duke University DLAR facility and at UC Irvine ULAR facility. All animal work adhered to NIH guidelines under IACUC approved protocols at Duke, UC Irvine, University of Minnesota, and University of Utah.

snRNA-seq Tissue Preparation and Sequencing

Snap frozen cerebellar tissues from two male and two female SCA7-266Q mice and paired WT littermates were used for snRNA-seq experiments. Nuclear isolation with Purkinje nuclei enrichment was performed using a modified protocol based closely on Kozareva et.al. 2021, by increasing K_2SO_4 concentration to 2.8g/500mL Dissociation Buffer which causes Granule Neuron nuclei to aggregate and fall out of solution. The supernatant will then contain a nuclear homogenate with elevated Purkinje cell proportions. Decreasing K_2SO_4 concentration to 1.4g/500mL in the Dissociation Buffer eliminated this effect. However, this concentration should be determined by the user, as other labs have observed varying K_2SO_4 levels to induce aggregates. After nuclei were purified, enriched for PCs, and stained with Hoechst 33342, individual wells representing nuclei from individual animals were separately labeled with MULTI-seq Cholesterol-Modified Oligos (CMOs) containing unique sample barcodes as described in McGinnis et.al. 2019. Barcoded nuclei from each animal were then sorted into a single collection tube using a BD FACSAria II cell sorter from the Hoechst singlet population before loading onto the 10X Genomics Chromium Controller with the 3' Gene Expression v3.1 kit. Sequencing libraries for snRNA and MULTI-seq barcodes were prepared separately from the cDNA as described in McGinnis et.al. 2019, and sequenced on an Illumina Novaseq 6000. Identical processing steps were performed on post-mortem human

cerebellar tissues (SCA7 n=5, Unaffected Controls n=4), but were instead sequenced on an Illumina Nextseq 2000.

snRNA-seq QC and Analysis

Sequencing FASTQ files were demultiplexed before running through the 10X Genomics CellRanger *count* pipeline (v4.0.0) to generate a feature filtered barcode matrix containing UMI filtered gene counts for each cell barcode. The Seurat R package (v4.2.0) was used to normalize counts, identify variable features, scale data, and cluster nuclei with UMAP dimensional reduction. MULTI-seq barcode FASTQs were run through the deMULTIplex R package () to generate sample specific barcode counts for each nucleus, remove doublets and ambiguous nuclei, and classify each nucleus to it's animal of origin. Filtered and labeled nuclear clusters were then annotated to their cell type of origin using top cluster specific marker genes with Seurat::FindMarkers, referencing known cell type marker genes and the Mouse Transcriptomic Cerebellar Atlas (Kozareva et.al. 2021). The same processing steps were performed on post-mortem human cerebellar snRNA-seq data.

Cell-type sub-clustering and DEseq2 Differential Expression

Cell types of interest were further investigated by subsetting into individual Seurat objects before re-normalization, re-scaling, and re-clustering with UMAP reduction. Cell Subtypes were investigated based on unbiased UMAP clusters to determine if SCA7 transcriptional dysregulation had outsized impacts on specific cell populations or cell states. Seurat::FindMarkers with the specified setting 'test.use = "DESeq2"' was used to compute differential expression between WT and SCA7 nuclei to pull from raw counts for pseudobulk comparisons. Cell Type DEG lists are provided in **Supplemental Data Table 1 (5wk DEGs) & Supplemental Data Table 2 (8.5wk DEGs)**.

Zebrin-axis Quantification and Overlap with SCA7 PC DEGs

In Purkinje Cells we determined that there was significant divergence in SCA7 transcriptional profiles in populations of Zebrin-positive and Zebrin-negative PCs. To accurately characterize these cell populations, 8.5wk Wild-Type PC nuclei were subset into a new Seurat object before re-normalization, re-scaling, and re-clustering with UMAP reduction. UMAP clusters definitively characterized nuclei into clusters of ZebrinII+ or ZebrinII- PCs, which were compared using Seurat::FindMarkers (significance cutoffs: adjusted p_val < 0.05 ; |log2FC| > 0.25) to generate a gene list representing the Zebrin transcriptional axis (**Supplemental Data Table 3**). These Zebrin-axis genes were intersected with the SCA7 Purkinje Cell DEGs and statistical enrichment was calculated using a Hypergeometric test with a Bonferonni correction parameter of 13,078 matching the number of Gene Ontology Biological Process terms for multiple testing.

Zebrin-axis WT and SCA7 Cross-comparison

The same steps described above to compute the WT Zebrin-axis were performed identically for 8.5wk SCA7 PC nuclei without any log2FC cutoffs. Differential expression outputs from WT and SCA7 Zebrin-axes were then directly compared by log2FC values for each gene and plotted together before running a linear regression to determine expression trends between WT and SCA7 Zebrin-axes.

Quantitative Synapse IHC

To quantify cerebellar synapses, SCA7-266Q mice and WT littermates were perfused with 4% PFA after clearing with PBS and brains were post-fixed in 4% PFA overnight. Brains were then cryoprotected by soaking in 10% sucrose/PBS, 20% sucrose/PBS, and 30% sucrose/PBS in 24hr increments before mounting cerebellar hemispheres in OCT and snap-freezing in a liquid nitrogen isopentane bath. A Leica CM3050S Cryostat was used to prepare 30μm sagittal slices from the

cerebellar vermis which were cryopreserved in 50% Glycerol : 30% Sucrose TBS. Synapse staining

was then performed by washing slices 3X in TBST (0.2% Triton-X) for 10 minutes, blocking with

10% NGS in TBST at room temp for 1hr, and staining with primary antibody cocktail in 10% NGS-

TBST overnight at 4°C. Slices were then washed 3X in TBST for 10 minutes, and stained with

secondary fluorescent antibodies at a concentration of 1:300 for 2hrs in 10% NGS-TBST at room

temp. Finally, slices were stained with Hoescht 33342 (1:5000) for 10 minutes in TBST and washed

2X for 10 minutes in TBST before mounting to microscope slides in Prolong Glass Antifade

Mountant. Slides were imaged on a Zeiss 880 Inverted Confocal microscope at appropriate

magnification (20X VGlut2 climbing fibers, 63X VGlut1 parallel fibers and VGat-Gepherin inhibitory

synapses) using Z-stack settings to collect 15 images with 0.34 μm step size in cerebellar Lobule VII.

The PunctaAnalyzer ImageJ tool[69] was used to separate the Z-stack images into 5 sets of 3

consecutive slices (~1 μm total) and compute puncta overlap between pre-synaptic and post-

synaptic markers, and averaging the co-localized synapse numbers from each slice to generate a

final synapse count for the Field of View.

Cerebellar Slice Electrophysiology

Purkinje Cell electrophysiology was performed in 250μm acute cerebellar slices in the coronal

plane of the posterior vermis in Lobule VII as described in Tsai et.al. 2012[70]. Briefly, slices were cut

in ice-cold Cutting Solution consisting of (mM): 130 K-Gluconate, 15 KCl, 0.05 EGTA, 20 HEPES, 25

Glucose (pH 7.4, osmolarity 320) before transferring to artificial cerebrospinal fluid (aCSF)

containing (mM): 125 NaCl, 26 NaHCO$_3$, 1.25 NaH$_2$PO$_4$, 2.5 KCl, 25 Glucose, 1 MgCl, 2 CaCl$_2$ (pH 7.3,

osmolarity 310). Slices were equilibrated with carbogen (95% O$_2$ and 5% CO$_2$) in aCSF for 20

minutes before moving to room temp with constant supply of carbogen. Recordings were taken in

aCSF without addition of synapse or action potential inhibiting agents with borosilicate patch

pipettes pulled to a resistance of 3-7 MΩ filled with an internal solution containing (mM): 140 Cs-

Gluconate, 15 HEPES, 0.5 EGTA, 2 TEA-Cl, 2 MgATP, 0.3 NaGTP, 10 phosphocreatine-Tris$_2$, 2 QX

314-Cl (pH 7.2). Action Potentials were recorded in a cell-attached configuration with a Gigaohm

resistance seal before breaking into the membrane for whole-cell voltage clamp recordings of

Spontaneous Post-Synaptic Currents. sEPSCs were recorded at -75mV and sIPSCs were recorded at

+10mV. Membrane potentials were not corrected for the liquid junction potential.

Zebrin-II IHC and Amplitude Analysis

Zebrin-II immunohistochemistry experiments were performed similarly as described above for

Quantitative Synapse IHC with some modifications. Coronal 100μm slices were taken from the

posterior cerebellum, and secondary antibody incubation time was increased to 4hrs. Images were

taken at a 10X magnification on the Zeiss 880 Inverted Confocal microscope with a tile-scan to

capture Zebrin-II and Calbindin immunofluorescence across the entire cerebellar slice. Zebrin

striping amplitudes were then analyzed by taking xy-intensity values across the Purkinje and

Molecular layer of the medial 3 Zb+ stripes (P1+,P2+) and medial 4 Zb- stripes (P1-,P2-) of

cerebellar Lobule VIII. The signal was collapsed by taking the median intensity value along the x-

axis, and a Loess polynomial regression was calculated to generate a signal trace across the region.

Local minima and maxima were identified on the regression trace, and signal amplitudes were

calculated for each consecutive stripe before computing the average Zebrin amplitude from these

values.

Quantitative Real-Time PCR (qRT-PCR)

Bulk RNA was purified from post-mortem human cerebellar tissues by pulverizing tissue with a

cold mortar and pestle on dry ice. Roughly 25mg of pulverized tissue was then lysed with TRIzol

(ThermoFisher) and purified with the Direct-zol RNA MiniPrep Plus kit (Zymo Research). Total

RNA was then treated with TURBO DNase (ThermoFisher) and reverse transcribed to cDNA with

SuperScript IV VILO (ThermoFisher). Taqman probes (ThermoFisher) against target genes with

FAM dye were used to amplify cDNA in Taqman Gene Expression Master Mix (ThermoFisher) and

compared to endogenous 18S rRNA with a Taqman VIC probe to generate Delta CT (dCt) values.

Relative expression fold changes were calculated using the 2^{-ddCt} method against the mean dCt value

for unaffected controls as reference. Two-tailed t-test was used to compute significance between

SCA7 and unaffected controls.

REFERENCES

1. Yoo, S. Y. *et al.* SCA7 knockin mice model human SCA7 and reveal gradual accumulation of mutant ataxin-7 in neurons and abnormalities in short-term plasticity. *Neuron* **37**, 383–401 (2003).

2. Watase, K. *et al.* A long CAG repeat in the mouse Sca1 locus replicates SCA1 features and reveals the impact of protein solubility on selective neurodegeneration. *Neuron* **34**, 905–919 (2002).

3. Hansen, S. T., Meera, P., Otis, T. S. & Pulst, S. M. Changes in Purkinje cell firing and gene expression precede behavioral pathology in a mouse model of SCA2. *Hum. Mol. Genet.* **22**, 271–283 (2013).

4. van de Warrenburg, B. P. C. *et al.* Striking anticipation in spinocerebellar ataxia type 7: the infantile phenotype. *J. Neurol.* **248**, 911–914 (2001).

5. Ansorge, O. *et al.* Ataxin-7 aggregation and ubiquitination in infantile SCA7 with 180 CAG repeats. *Ann. Neurol.* **56**, 448–452 (2004).

6. Trang, H. *et al.* Massive CAG Repeat Expansion and Somatic Instability in Maternally Transmitted Infantile Spinocerebellar Ataxia Type 7. *JAMA Neurol.* **72**, 219 (2015).

7. La Spada, A. R. & Taylor, J. P. Polyglutamines Placed into Context. *Neuron* **38**, 681–684 (2003).

8. Harjes, P. & Wanker, E. E. The hunt for huntingtin function: interaction partners tell many different stories. *Trends Biochem. Sci.* **28**, 425–433 (2003).

9. Li, S.-H. & Li, X.-J. Huntingtin-protein interactions and the pathogenesis of Huntington's disease. *Trends Genet. TIG* **20**, 146–154 (2004).

10. Helmlinger, D. *et al.* Ataxin-7 is a subunit of GCN5 histone acetyltransferase-containing complexes. *Hum. Mol. Genet.* **13**, 1257–1265 (2004).

11. Zhao, Y. *et al.* A TFTC/STAGA module mediates histone H2A and H2B deubiquitination, coactivates nuclear receptors, and counteracts heterochromatin silencing. *Mol. Cell* **29**, 92–101 (2008).

12. Martinez, E. *et al.* Human STAGA Complex Is a Chromatin-Acetylating Transcription Coactivator That Interacts with Pre-mRNA Splicing and DNA Damage-Binding Factors In Vivo. *Mol. Cell. Biol.* **21**, 6782–6795 (2001).

13. Jenuwein, T. & Allis, C. D. Translating the Histone Code. *Science* **293**, 1074–1080 (2001).

14. Minsky, N. *et al.* Monoubiquitinated H2B is associated with the transcribed region of highly expressed genes in human cells. *Nat. Cell Biol.* **10**, 483–488 (2008).

15. Wojcik, F. *et al.* Functional crosstalk between histone H2B ubiquitylation and H2A modifications and variants. *Nat. Commun.* **9**, 1394 (2018).

16. McMahon, S. J., Pray-Grant, M. G., Schieltz, D., Yates, J. R. & Grant, P. A. Polyglutamine-expanded spinocerebellar ataxia-7 protein disrupts normal SAGA and SLIK histone acetyltransferase activity. *Proc. Natl. Acad. Sci.* **102**, 8478–8482 (2005).

17. Palhan, V. B. *et al.* Polyglutamine-expanded ataxin-7 inhibits STAGA histone acetyltransferase activity to produce retinal degeneration. *Proc. Natl. Acad. Sci.* **102**, 8472–8477 (2005).

18. Köhler, A., Schneider, M., Cabal, G. G., Nehrbass, U. & Hurt, E. Yeast Ataxin-7 links histone deubiquitination with gene gating and mRNA export. *Nat. Cell Biol.* **10**, 707–715 (2008).

19. David, G. *et al.* Cloning of the SCA7 gene reveals a highly unstable CAG repeat expansion. *Nat. Genet.* **17**, 65–70 (1997).

20. Kozareva, V. *et al.* A transcriptomic atlas of mouse cerebellar cortex comprehensively defines cell types. *Nature* **598**, 214–219 (2021).

21. Howarth, C., Peppiatt-Wildman, C. M. & Attwell, D. The Energy Use Associated with Neural Computation in the Cerebellum. *J. Cereb. Blood Flow Metab.* **30**, 403–414 (2010).

22. Howarth, C., Gleeson, P. & Attwell, D. Updated Energy Budgets for Neural Computation in the Neocortex and Cerebellum. *J. Cereb. Blood Flow Metab.* **32**, 1222–1232 (2012).

23. Horton, L. C. *et al.* Spinocerebellar Ataxia Type 7: Clinical Course, Phenotype-Genotype Correlations, and Neuropathology. *Cerebellum Lond. Engl.* **12**, 176–193 (2013).

24. Custer, S. K. *et al.* Bergmann glia expression of polyglutamine-expanded ataxin-7 produces neurodegeneration by impairing glutamate transport. *Nat. Neurosci.* **9**, 1302–1311 (2006).

25. Furrer, S. A. *et al.* Spinocerebellar ataxia type 7 cerebellar disease requires the coordinated action of mutant ataxin-7 in neurons and glia, and displays non-cell-autonomous bergmann glia degeneration. *J. Neurosci. Off. J. Soc. Neurosci.* **31**, 16269–16278 (2011).

26. Oscarsson, O. Functional units of the cerebellum - sagittal zones and microzones. *Trends Neurosci.* **2**, 143–145 (1979).

27. Brochu, G., Maler, L. & Hawkes, R. Zebrin II: A polypeptide antigen expressed selectively by purkinje cells reveals compartments in rat and fish cerebellum. *J. Comp. Neurol.* **291**, 538–552 (1990).

28. Zhou, H. *et al.* Cerebellar modules operate at different frequencies. *eLife* **3**, e02536 (2014).

29. Cerminara, N. L., Lang, E. J., Sillitoe, R. V. & Apps, R. Redefining the cerebellar cortex as an assembly of non-uniform Purkinje cell microcircuits. *Nat. Rev. Neurosci.* **16**, 79–93 (2015).

30. McGinnis, C. S. *et al.* MULTI-seq: sample multiplexing for single-cell RNA sequencing using lipid-tagged indices. *Nat. Methods* **16**, 619–626 (2019).

31. Squair, J. W. *et al.* Confronting false discoveries in single-cell differential expression. *Nat. Commun.* **12**, 5692 (2021).

32. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

33. Junttila, S., Smolander, J. & Elo, L. L. Benchmarking methods for detecting differential states between conditions from multi-subject single-cell RNA-seq data. *Brief. Bioinform.* **23**, bbac286 (2022).

34. Sidman, R. L., Dickie, M. M. & Appel, S. H. MUTANT MICE (QUAKING AND JIMPY) WITH DEFICIENT MYELINATION IN THE CENTRAL NERVOUS SYSTEM. *Science* **144**, 309–311 (1964).

35. Chen, T. & Richard, S. Structure-Function Analysis of Qk1: a Lethal Point Mutation in Mouse quaking Prevents Homodimerization. *Mol. Cell. Biol.* **18**, 4863–4871 (1998).

36. Takeuchi, A. *et al.* Identification of Qk as a Glial Precursor Cell Marker that Governs the Fate Specification of Neural Stem Cells to a Glial Cell Lineage. *Stem Cell Rep.* **15**, 883–897 (2020).

37. Sakers, K. *et al.* Loss of Quaking RNA binding protein disrupts the expression of genes associated with astrocyte maturation in mouse brain. *Nat. Commun.* **12**, 1537 (2021).

38. Ahn, A. H., Dziennis, S., Hawkes, R. & Herrup, K. The cloning of zebrin II reveals its identity with aldolase C. *Development* **120**, 2081–2090 (1994).

39. Xiao, J. *et al.* Systematic Regional Variations in Purkinje Cell Spiking Patterns. *PLOS ONE* **9**, e105633 (2014).

40. Chen, X. *et al.* Transcriptomic mapping uncovers Purkinje neuron plasticity driving learning. *Nature* **605**, 722–727 (2022).

41. Niewiadomska-Cimicka, A., Hache, A. & Trottier, Y. Gene Deregulation and Underlying Mechanisms in Spinocerebellar Ataxias With Polyglutamine Expansion. *Front. Neurosci.* **14**, (2020).

42. Niewiadomska-Cimicka, A. *et al.* SCA7 Mouse Cerebellar Pathology Reveals Preferential Downregulation of Key Purkinje Cell-Identity Genes and Shared Disease Signature with SCA1 and SCA2. *J. Neurosci. Off. J. Soc. Neurosci.* **41**, 4910–4936 (2021).

43. Schilling, G. *et al.* Intranuclear Inclusions and Neuritic Aggregates in Transgenic Mice Expressing a Mutant N-Terminal Fragment of Huntingtin. *Hum. Mol. Genet.* **8**, 397–407 (1999).

44. Tejwani, L. *et al.* Longitudinal single-cell transcriptional dynamics throughout neurodegeneration in SCA1. 2021.10.22.465444 Preprint at https://doi.org/10.1101/2021.10.22.465444 (2021).

45. Siletti, K. *et al.* Transcriptomic diversity of cell types across the adult human brain. 2022.10.12.511898 Preprint at https://doi.org/10.1101/2022.10.12.511898 (2022).

46. Gao, W., Chen, G., Reinert, K. C. & Ebner, T. J. Cerebellar Cortical Molecular Layer Inhibition Is Organized in Parasagittal Zones. *J. Neurosci.* **26**, 8377–8387 (2006).

47. Consalez, G. G. & Hawkes, R. The compartmental restriction of cerebellar interneurons. *Front. Neural Circuits* **6**, 123 (2012).

48. Zhou, J. *et al.* Purkinje cell neurotransmission patterns cerebellar basket cells into zonal modules defined by distinct pinceau sizes. *eLife* **9**, e55569 (2020).

49. Chaudhry, F. A. *et al.* The Vesicular GABA Transporter, VGAT, Localizes to Synaptic Vesicles in Sets of Glycinergic as Well as GABAergic Neurons. *J. Neurosci.* **18**, 9733–9750 (1998).

50. Wojcik, S. M. *et al.* An essential role for vesicular glutamate transporter 1 (VGLUT1) in postnatal development and control of quantal size. *Proc. Natl. Acad. Sci.* **101**, 7158–7163 (2004).

51. Hioki, H. *et al.* Differential distribution of vesicular glutamate transporters in the rat cerebellar cortex. *Neuroscience* **117**, 1–6 (2003).

52. Häusser, M. & Clark, B. A. Tonic synaptic inhibition modulates neuronal output pattern and spatiotemporal synaptic integration. *Neuron* **19**, 665–678 (1997).

53. Raman, I. M. & Bean, B. P. Resurgent sodium current and action potential formation in dissociated cerebellar Purkinje neurons. *J. Neurosci. Off. J. Soc. Neurosci.* **17**, 4517–4526 (1997).

54. De Zeeuw, C. I. *et al.* Microcircuitry and function of the inferior olive. *Trends Neurosci.* **21**, 391–400 (1998).

55. Davie, J. T., Clark, B. A. & Häusser, M. The Origin of the Complex Spike in Cerebellar Purkinje Cells. *J. Neurosci.* **28**, 7599–7609 (2008).

56. Stoyas, C. A. *et al.* Nicotinamide Pathway-Dependent Sirt1 Activation Restores Calcium Homeostasis to Achieve Neuroprotection in Spinocerebellar Ataxia Type 7. *Neuron* **105**, 630-644.e9 (2020).

57. Apps, R. & Hawkes, R. Cerebellar cortical organization: a one-map hypothesis. *Nat. Rev. Neurosci.* **10**, 670–681 (2009).

58. Miale, I. L. & Sidman, R. L. An autoradiographic analysis of histogenesis in the mouse cerebellum. *Exp. Neurol.* **4**, 277–296 (1961).

59. Goffinet, A. M. The embryonic development of the cerebellum in normal and reeler mutant mice. *Anat. Embryol. (Berl.)* **168**, 73–86 (1983).

60. Yuasa, S., Kawamura, K., Ono, K., Yamakuni, T. & Takahashi, Y. Development and migration of Purkinje cells in the mouse cerebellar primordium. *Anat. Embryol. (Berl.)* **184**, 195–212 (1991).

61. Hatten, M. E. & Heintz, N. Mechanisms of neural patterning and specification in the developing cerebellum. *Annu. Rev. Neurosci.* **18**, 385–408 (1995).

62. Apsley, E. J. & Becker, E. B. E. Purkinje Cell Patterning-Insights from Single-Cell Sequencing. *Cells* **11**, 2918 (2022).

63. Serra, H. G. *et al.* RORalpha-mediated Purkinje cell development determines disease severity in adult SCA1 mice. *Cell* **127**, 697–708 (2006).

64. Capizzi, M. *et al.* Developmental defects in Huntington's disease show that axonal growth and microtubule reorganization require NUMA1. *Neuron* **110**, 36-50.e5 (2022).

65. Mangiarini, L. *et al.* Exon 1 of the HD Gene with an Expanded CAG Repeat Is Sufficient to Cause a Progressive Neurological Phenotype in Transgenic Mice. *Cell* **87**, 493–506 (1996).

66. Bushart, D. D. & Shakkottai, V. G. Ion channel dysfunction in cerebellar ataxia. *Neurosci. Lett.* **688**, 41–48 (2019).

67. Lappe-Siefke, C. *et al.* The Ataxia (axJ) Mutation Causes Abnormal GABAA Receptor Turnover in Mice. *PLOS Genet.* **5**, e1000631 (2009).

68. Chiu, C.-S. *et al.* GABA Transporter Deficiency Causes Tremor, Ataxia, Nervousness, and Increased GABA-Induced Tonic Conductance in Cerebellum. *J. Neurosci.* **25**, 3234–3245 (2005).

69. Ippolito, D. M. & Eroglu, C. Quantifying Synapses: an Immunocytochemistry-based Assay to Quantify Synapse Number. *J. Vis. Exp. JoVE* 2270 (2010) doi:10.3791/2270.

70. Tsai, P. T. *et al.* Autistic-like behaviour and cerebellar dysfunction in Purkinje cell Tsc1 mutant

   mice. *Nature* **488**, 647–651 (2012).

ACKNOWLEDGEMENTS

Anti-Zebrin-II antibody was generated and generously provided by Dr. Richard Hawkes and Dr. Carol Armstrong, University of Calgary. Post-mortem Human Tissues were provided by L. Ranum (Center for NeuroGenetics, University of Florida) and M. Perkins (University of Michigan Brain Bank). This work was supported by grants from the N.I.H. (Genomic Innovator Award to C.B.L.; R01 EY014061 and R35 NS122140 to A.R.L.S.), and the Holland-Trice Scholars Award (L.C.B).

Author Contributions

L.C.B, and A.R.L.S. provided the conceptual framework for the study. L.C.B, A.R.L.S, C.B.L, and C.H. designed the experiments. L.C.B. performed the experiments and analyzed the data. L.C.B, C.B.L, and A.R.L.S. wrote the manuscript.

Declarations

The authors have nothing to declare.

Figure Legends

Figure 1: Purkinje-Enriched snRNA-seq Yields a High-Resolution Map of SCA7 Cerebellum. A) Experimental schematic detailing Purkinje-enriched snRNA-seq method combined with MULTI-seq CMO barcoding for multiplexing biological replicates into a single 10X GEM well, applied to 2-male and 2-female WT or SCA7-266Q cerebella. B) Relative cell type abundance ratios quantified from snRNA-seq samples does not find any significant cell type ratio differences between WT and SCA7 samples by two-tailed t-test. C) UMAP dimensional reduction of 5wk pre-symptomatic Purkinje-enriched snRNA-seq data with points colored by genotype of origin. (coral = SCA7; teal = WT). D) UMAP dimensional reduction for 8.5wk symptomatic timepoint. E) Differential Expression results per cell type between SCA7 and WT nuclei for the 5wk timepoint using pseudobulk DEseq2. F) Differential Expression results per cell type for 8.5wk timepoint. Each point represents a significant DEG (cutoffs: adjusted p_val < 0.05 ; |log2-fold change| > 0.25)

Figure 2: Signatures of Glial Identity Loss Identified in SCA7-266Q Animals. A) UMAP dimensional reduction for Astrocytes, Bergmann Glia, and Oligodendrocytes at the 8.5wk timepoint after re-clustering. Points colored genotype of origin (coral = SCA7; teal = WT). B) Volcano Plots showing differential expression results between SCA7 and WT nuclei in Astrocytes, Bergmann Glia, and Oligodendrocytes. Log2FC and –$\log_{10}$(p-value) for each gene is shown with significant DEGs in red (cutoffs: adjusted p_val < 0.05 ; |log2-fold change| > 0.25). D) Venn-Diagram showing overlap of significant DEGs from each glial cell type with CLIP-seq targets of the QK RNA-binding protein (Sakers et.al. 2021). Hypergeometric test used to calculate overlap enrichment. Unadjusted p-value for overlaps shown along with Gene IDs.

Figure 3: Progressive Transcriptional Dysregulation in SCA7 Purkinje Cells Highlights Changes in Synaptic Signaling. A) UMAP dimensional reduction for Purkinje Cell nuclei after

subsetting and re-clustering at 5wk time point. B) Volcano Plot showing differential expression results between presymptomatic SCA7 and WT Purkinje nuclei with pseudobulk DEseq2. Significant DEGs shown in red (cutoffs: adjusted p_val < 0.05 ; |log2-fold change| > 0.25). C) UMAP dimensional reduction for Purkinje Cell nuclei after subsetting and re-clustering at 8.5wk time point. D) Volcano Plot showing differential expression results between symptomatic SCA7 and WT Purkinje nuclei with pseudobulk DEseq2. Significant DEGs in red (cutoffs: adjusted p_val < 0.05 ; |log2-fold change| > 0.25). E) Gene Ontology categories for 8.5wk symptomatic DEGs (273 DEGs) calculated with ShinyGO v0.77. Biological Process categories are sorted by -Log10(FDR) with vertical cutoff lines drawn at equivalent adjusted p-values of (dotted: p=0.01, dashed: p=0.001, solid: p=0.0001).

Figure 4: Zebrin-II Subtype Dysregulation Explains Highest Proportion of SCA7 Purkinje Cell DEGs. A) UMAP dimensional reduction for 8.5wk Purkinje Cells shown alongside normalized Aldoc (Zebrin-II) expression for each single-nucleus. B) WT Purkinje Cells subset and re-clustered to find Zebrin-positive and Zebrin-negative clusters. C) Volcano Plot showing differential expression between WT Zebrin-positive and Zebrin-negative clusters. Wilcoxon Rank Sum test was used and significant genes are shown in red (cutoffs: adjusted p_val < 0.05 ; |log2FC| > 0.25). C) Venn-Diagram showing overlap of 8.5wk SCA7 DEGs in PCs with Significant Genes of the WT Zebrin-Axis. Statistical enrichment calculated with Hypergeometric test and adjusted with multiple test correction of 13,078 representing number of Gene Ontology Biological Process Categories (adjusted p_val = 2.65e-51***). E) Gene Ontology Biological Processes from Figure 3B re-plotted with Zebrin-Axis gene category added. F) WT Zebrin-axis DEGs calculated from 5wk PCs (as in panel B) and plotted against 5wk SCA7 Zebrin-axis DEGs by Log2FC value. Linear Regression was applied to find correlation of Log2FC values from WT and SCA7 (slope = 0.744). (Points colored red if Log2FC

difference between WT and SCA7 is greater than 0.25). G) WT and SCA7 Zebrin-axis comparative plot for 8.5wk PCs. (slope = 0.833)

Figure 5: Zebrin-II Parasagittal Stripes are Lost in SCA7 and Related Polyglutamine Ataxias. A) Immunohistochemistry in coronal cerebellar slices from 8.5wk WT and SCA7 animals stained for Zebrin-II/Aldoc (green) and Calbindin (blue). B) Representative images of Zebrin staining intensity across the central 3 stripes of Cerebellar Lobule VIII paired with median pixel intensity values across the measured area. Intensity values were fit with a Loess Polynomial Regression, before measuring the signal amplitude of each stripe and averaging to find the mean Zebrin intensity for each animal. Models and ages tested: WT 5wk (n=3), SCA7 5wk (n=3), WT 8.5wk (n=3), SCA7 8.5wk (n=3), SCA1-175Q 12wk (N=3), SCA2-127Q 12wk (n=2) and HD-N171-82Q 13wk (N=3). WT littermates for SCA1, SCA2 and HD models were also analyzed for Zebrin stripe intensity shown grouped with WT 8.5wk. Amplitudes for each group were compared to respective WT controls with a one-tailed T-test. (SCA7 8.5wk **p = 0.0065; SCA1-175Q 12wk **p= 0.0083; SCA2-127Q 12wk, *p= 0.016)

Figure 6: Human SCA7 Cerebellar Tissues Show Degradation of Zebrin-Related Subtypes in Molecular Layer Interneurons. A) snRNA-seq UMAP clusters from Human Post-mortem Cerebellar tissues from 5 SCA7 and 4 Healthy controls colored by cell type. B) UMAP clusters colored by patient disease status. C) Differential expression results comparing SCA7 to Healthy controls for each recovered cell type using pseudobulk DEseq2. D) Molecular Layer Interneuron sub-clustering results showing strong co-clustering of SCA7 nuclei, and strong down-regulation of Zebrin-II/ALDOC. Panels are colored by UMAP cluster (top-left), Disease Status (top-right), Sample Donor (bottom-left), Normalized ALDOC expression (bottom-right). E) UMAP sub-clustering of MLI-1 cells from the Mouse Cerebellar Atlas colored by UMAP cluster (top-left), Normalized Aldoc

expression (top-right), Normalized Aldoa expression (bottom-left), and volcano plot of resulting differential expression results from Aldoc-high vs. Aldoc-low clusters (bottom right). F) Venn-Diagram showing overlap of SCA7 Human MLI-1 DEGs with Mouse MLI-1 Stripe DEGs. Statistical enrichment of overlap computed with hypergeometric test shown with corrected p-value (p=1.02e-147). G) IHC representative image in 5wk WT mouse cerebellum showing striped expression of Aldoc along with Aldoa. H) qRT-PCR for Zebrin Identity genes on bulk RNA extracted from human SCA7 (n=4) and Healthy control (n=3) cerebella. Fold change shown relative to mean Ct value for Healthy control samples, statistics calculated with one-tailed t-test.

Figure 7: Symptomatic SCA7-266Q Mice Contain Increased Inhibitory Synapses. A) Inhibitory Synapse IHC in the molecular layer of Lobule VII in 8.5wk WT and SCA7-266Q mice (63X magnification; green = Vgat, red = Gephyrin, blue = Calb1). Image in-sets contain arrows pointing to yellow co-localized signal of intact inhibitory synapses. White arrows show synapses on Calb1 PC dendrites. Orange arrow delineates a non-PC synapse. B) Quantification of 8.5wk inhibitory synapses by intact synapse numbers in a 15um z-scan. Inhibitory synapses overlapping Calb1 were classified as PC-synapses, or non-PC synapses if not overlapping Calb1. (n=6 animals; two images per animal; 8.5wk PC Synapses **p=0.001, 8.5wk non-PC Synapses *p=0.021) C) Inhibitory Synapse IHC for 5wk WT and SCA7-266Q mice. D) Quantification of 5wk Inhibitory Synapses.

Figure 8: Increased Inhibitory Post-Synaptic Currents in SCA7 Purkinje Cells Leads to Reduced Spiking Rates. A) Spontaneous Inhibitory Post-Synaptic Currents (sIPSC) were recorded from Purkinje cells in acute cerebellar slices from 8.5wk symptomatic SCA7-266Q animals and WT littermates. Whole-cell voltage-clamped recordings were measured at +10mV from n=6 WT and n=15 SCA7 Purkinje Cells. Amplitudes and Inter-Event-Intervals were measured for each sIPSC event before randomly sampling an equal number from each cell, shown plotted in a cumulative

histogram. Statistics performed with Wilcoxson Rank Sum test. (IPSC amplitudes ***p=2.48x10^{-13})

B) Spontaneous Excitatory Post-Synaptic Currents (sEPSC) were recorded, voltage-clamped at -75mV from n=18 WT and n=17 SCA7 Purkinje Cells. Amplitudes and Inter-Event-Intervals were measured for each sEPSC and randomly sampled, shown plotted in a cumulative histogram. (EPSC inter-event-intervals ***p=2.20x10^{-16}) C) Cell-attached recordings were taken from n=14 WT and n=14 SCA7 Purkinje Cells. Action Potentials were analyzed to find inter-event-intervals and randomly sampled from each cell to produce a cumulative histogram. Statistics performed with Wilcoxon Rank Sum test (***p=2.53x10^{-10}) D) The Coefficient of Variation in recorded action potentials was measured for each cell, as well as the CV2 statistic. No significant differences in spiking regularity were found.

Supplementary Figure legends

Supplemental Figure S1: Purkinje Cell Nuclear Enrichment Method. A) Images of nuclear extraction buffers with standard concentration of potassium sulfate generating a homogenous suspension of cerebellar nuclei (left), and enriched concentrations of potassium sulfate leading to precipitation of granule neuron nuclei (right). B) Immunohistochemistry from sagittal cerebellar slices of INTACT Sun1-GFP mice display nuclear envelopes labeled with GFP. White arrows point to Purkinje Cell nuclei which are notably larger than surrounding cell types. C) Flow Cytometry of nuclei isolated from INTACT Sun1-GFP mice with Standard and K_2SO_4 Enriched extraction buffers. Side-scatter (SSC-A) and GFP fluorescence separate cerebellar cell types by nuclear size and granularity. K_2SO_4 Enriched extraction buffers leads to a selective enrichment of GFP+++ nuclei. D) qRT-PCR from sorted total nuclei vs. GFP+++ gate validates enriched expression of Purkinje Cell markers (n=3)

Supplemental Figure S2: snRNA-seq Cell Type Marker Genes. A) Seurat::FeaturePlots showing canonical Purkinje Cell marker *Pcp2* and non-canonical marker *Ryr1* which showed highest specificity to Purkinje Cells in our dataset. B) Dot Plot showing top 5 marker genes for all cell types recovered. C) Feature plots visualizing cluster specific expression of top markers for Astrocytes (*Slc1a2*), Bergmann Glia (*Slc1a3*), Golgi Cells (*Gabrg3*), Granule Neurons (*Chn2*), Oligodendrocytes (*Mbp*), Molecular Layer Interneurons (*Cacna2d3*)

Supplemental Figure S3: Cell Type Selective Decrease in RNA Abundance in SCA7-266Q snRNA-seq. A)   Violin Plot showing RNA counts per nucleus (nCount_RNA) for the 5wk timepoint split by cell type and genotype. Statistical differences between SCA7 and WT were calculated for each cell type by Wilcoxon Rank Sum test. (p-value cutoffs: * < 0.05, ** < 0.01, *** < 0.001, **** < 0.0001) B) Violin Plot showing RNA counts for the 8.5wk timepoint. C) Violin Plot showing RNA counts for 8.5wk Purkinje Cells split by animal. D) Violin Plot showing # of Genes detected per nucleus (nFeature_RNA) for the 8.5wk timepoint split by cell type and genotype.

Supplemental Figure S4: Cell Type Selective Decrease in RNA Abundance in Human SCA7 snRNA-seq. A) Violin Plot showing RNA counts per nucleus (nCount_RNA) for the Human Post-mortem Cerebellum snRNA-seq split by cell type and disease status. Statistical differences between SCA7 and Healthy controls were calculated for each cell type by Wilcoxon Rank Sum test. (p-value cutoffs: *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001) B) Violin Plot showing RNA counts for Type-I Molecular Layer Interneurons. C) Violin Plot showing RNA counts for Granule Neurons.

Supplemental Figure S5: Subclusters of MLI-1 and MLI-2 Nuclei from Transcriptomic Mouse Cerebellar Atlas Show Zebrin-II Gene Expression Patterning. A) UMAP plots showing re-clustered MLI-1 and MLI-2 nuclei from Kozareva et.al. Transcriptomic Mouse Cerebellar Atlas. B)

Feature Plots showing normalized expression of *Aldoc* in MLI-1 and MLI-2 clusters. Dotted lines highlight *Aldoa*-'high' and *Aldoa*-'low' subclusters. C) Feature Plots showing normalized expression of *Aldoa* in MLI-1 and MLI-2 clusters. D) Venn-Diagram showing overlap of Zebrin DEGs between *Aldoa*-'high' and *Aldoa*-'low' subclusters in MLI-1 and MLI-2.

Supplemental Figure S6: VGlut2 Climbing Fiber Synapses Remain Unchanged in Early Symptomatic SCA7-266Q Animals. A) Quantitative Synapse IHC representative images for excitatory climbing fiber synapses in cerebellar Lobule VII in 5wk mice. (20X magnification; magenta = VGlut2; cyan = Calb1) B) Quantification of 5wk climbing fiber synapses by puncta density and percent innervation along full length of the PC dendrite. C) Quantitative Synapse IHC representative images for excitatory climbing fiber synapses in cerebellar Lobule VII in 8.5wk mice. (20X magnification; magenta = VGlut2; cyan = Calb1) D) Quantification of 8.5wk climbing fiber synapses by puncta density and percent innervation along full length of the PC dendrite.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

WT – 8.5wk
SCA7 – 8.5wk
Zebrin II/Aldoc
Calbindin

WT – 5wk
SCA7 – 5wk
WT – 8.5wk
SCA7 – 8.5wk
SCA1 – 12wk
SCA2 – 12wk
HD – 13wk

Zebrin Stripe Amplitudes

Mean Zebrin Amplitude

WT (5wk)
SCA7 (5wk)
WT (8.5wk)
SCA7 (8.5wk)
SCA1 (12wk)
SCA2 (12wk)
HD (13wk)

Figure 6

Figure 7

Figure 8

Supplementary Figure S1

Supplementary Figure S2

Supplementary Figure S3

Supplementary Figure S4

Supplementary Figure S5

Supplementary Figure S6