Inferring compound heterozygosity from large-scale exome sequencing data

Michael H. Guo[1,2,*], Laurent C. Francioli[2,3,*], Sarah L. Stenton[2,3,4], Julia K. Goodrich[2,3], Nicholas A. Watts[2,3], Moriel Singer-Berk[2], Emily Groopman[2,4], Philip W. Darnowsky[2,3], Matthew Solomonson[2,3], Samantha Baxter[2], gnomAD Project Consortium[†], Grace Tiao[2,3], Benjamin M. Neale[2,3,5], Joel N. Hirschhorn[2,6,7,8], Heidi L. Rehm[2,3,9], Mark J. Daly[2,3,10], Anne O'Donnell-Luria[2,3,4,9], Konrad J. Karczewski[2,3,9], Daniel G. MacArthur[2,3,11,12], and Kaitlin E. Samocha[2,3,9‡]

[1] Department of Neurology, Hospital of the University of the Pennsylvania, Philadelphia, PA, USA

[2] Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[3] Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA

[4] Division of Genetics and Genomics, Boston Children's Hospital, Boston, MA

[5] Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[6] Department of Genetics, Blavatnik Institute, Harvard Medical School, Boston, MA, USA

[7] Division of Endocrinology, Boston Children's Hospital, Boston, MA, USA

[8] Center for Basic and Translational Obesity Research, Boston Children's Hospital, Boston, MA, USA

[9] Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA

[10] Institute for Molecular Medicine Finland, (FIMM) Helsinki, Finland

[11] Centre for Population Genomics, Garvan Institute of Medical Research, UNSW Sydney, Sydney, Australia

[12] Centre for Population Genomics, Murdoch Children's Research Institute, Melbourne, Australia

* These authors contributed equally: Michael H. Guo and Laurent C. Francioli

† List of authors and their affiliations appear at the end of the paper

‡ Correspondence should be addressed to K.E.S. (samocha@broadinstitute.org)

Abstract

Severe recessive diseases arise when both the maternal and the paternal copies of a

gene carry, or are impacted by, a damaging genetic variant in the affected individual. When a

patient carries two different potentially causal variants, accurate diagnosis requires determining

that these two variants occur on different copies of the chromosome (i.e., are in *trans*) rather

than on the same copy (i.e., in *cis*). However, current approaches for determining phase,

beyond parental testing, are limited in clinical settings. We developed a strategy for inferring

phase for rare variant pairs within genes, leveraging haplotype patterns observed in exome

sequencing data from the Genome Aggregation Database (gnomAD v2, n=125,748). When

applied to trio data where phase is known, our approach estimates phase with high accuracy,

even for very rare variants (frequency $<1\times10^{-4}$), and also correctly phases 95.2% of variant pairs

in a set of 293 patients carrying presumed causal compound heterozygous variants. We provide

a public resource of phasing estimates from gnomAD, including phasing estimates for coding

variants across the genome and counts per gene of rare variants in *trans*, that can aid

interpretation of rare co-occurring variants in the context of recessive disease.

Introduction

17   Determination of haplotypic phase has important implications in clinical genetics,

particularly in the diagnosis of recessive diseases that result from disruption of both copies of a

gene. The disrupting bi-allelic variants can be either homozygous, where the same variant is

present on both copies, or compound heterozygous, where two different variants are present on

the two copies of the gene. Compound heterozygous variants present a challenge in genetic

diagnosis because two variants observed within a gene in an individual can occur in *trans* or in

cis, and only the former scenario results in compound heterozygosity. However, parental data

are often not readily available for phasing (which is the typical situation in panel testing) or

parents may not be available for follow-up testing, and short-read next generation sequencing

largely cannot directly distinguish whether variant pairs are in *trans* or in *cis.* Thus, there is an

important need for other approaches to accurately, easily, and cheaply determine phase of

variant pairs.

29   The genetic relationship between a pair of variants on a haplotype can be disrupted by

one of two processes: meiotic recombination and recurrent mutations. Meiotic recombination

occurs more frequently in "hotspot" regions, and the probability of a recombination event

occurring increases with distance between two variants[1]. A recurrent germline mutation event

affecting a variant on a haplotype can also disrupt the genetic relationship of the variants on the

haplotype. Rates of recurrent mutations are dependent on mutation type (e.g., transition versus

transversion) and epigenetic marks (particularly CpG methylation), among other factors[2–6].

Thus, the rates of both meiotic recombination and mutation have important implications in

determining the phase of variants.

38   There are several approaches for directly inferring phase for variant pairs observed in an

individual. Phase may be determined directly using data from sequencing reads. However, for

data from typical short-read sequencing technologies such as Illumina, read-based phasing

methods are generally only possible for variants in close proximity to each other[7], although

some variant pairs at longer distances can be phased with more sophisticated algorithms[8–10].

Long-read sequencing technologies allow for direct determination of phase for variant pairs at

longer genetic distances, but these technologies are more expensive and have not yet been

widely applied in clinical settings[11,12]. There are also laboratory-based molecular methods for

determining phase of variant pairs, but these methods are low-throughput and technically

challenging[13]. While phase can be determined based on transmission of variants from parents

to offspring, this approach increases cost and may pose other logistical and ethical challenges,

and is not always an option if parents are deceased, unavailable (e.g., living far away or

incarcerated), unknown in the case of adoption, or unwilling to participate. These direct phasing

approaches all thus present critical limitations for determining phase of variant pairs within an

individual in a clinical setting.

Alternative, indirect approaches to phasing rely on statistical methods applied to

population data (reviewed in Tewhey et al.[14] and Browning and Browning[15]). These approaches

leverage genetic data from large numbers of unrelated or distantly related individuals, and rely

on identifying shared haplotypes among individuals in a population. However, these methods

require a large number of reference samples (typically $n \sim 10^5$-10^6 individuals) and are

computationally intensive to perform. These approaches perform less well for rare variants and

cannot be applied to exome sequencing data, which does not provide enough density of

surrounding variants to allow for accurate phasing. Despite these limitations, these population-

based approaches are attractive because they do not require sequencing of additional family

members or application of expensive sequencing approaches.

In this work, we sought to address existing challenges of phasing in clinical settings,

particularly with regard to rare variants seen in exome sequencing data. We implement an

approach that leverages the principles of population-based phasing by estimating haplotype

patterns from a large reference population and using these patterns to infer variant phase in an

individual. We cataloged the haplotype patterns of rare coding variants within genes using the

Genome Aggregation Database (gnomAD), which performed aggregation and joint genotyping

of exome sequencing data from 125,748 individuals[16]. We then demonstrate that we can

leverage these data to generate a resource for phasing rare coding variants observed in an

individual, and identify factors that influence the accuracy of our approach. Additionally, we

provide statistics for how often different types of variants are observed, split by allele frequency

and mutational consequence, in *trans* within gnomAD to provide a background rate

contextualization when observing biallelic rare variants in rare disease cases. Finally, we

disseminate these resources in a user-friendly fashion via the gnomAD browser for community

use.

Results

Inference of phase in gnomAD

We sought to address the challenges of phasing individual samples in clinical settings by

applying the principles of population-based phasing. Specifically, we leveraged the fact that

haplotypes are usually shared across individuals in a population to infer phase of variants in an

individual (**Figure 1A**). If two variants are in *cis* in many individuals in a population, then they

are likely to be *cis* in any given individual's DNA. Similarly, if two variants are in *trans* in other

individuals in a population, then they are likely to be *trans* in any given individual's DNA. This

later scenario also provides information that the variant combination may be tolerated in *trans*

since it has been found in an individual in gnomAD. We reasoned that by generating phasing

estimates from a large reference population, we could infer the phase of variants observed in an

individual.

Predicting the phase of a given pair of variants in an individual first requires that we

estimate the haplotype frequencies in the population for a given pair of variants. To estimate

haplotype frequencies, we used exome sequencing samples from gnomAD v2, a large

sequencing aggregation database[16]. In total, there were 125,748 exome sequencing samples

after rigorous sample and variant quality control (Methods). There are several key advantages

of using gnomAD. First, samples in gnomAD undergo uniform processing and variant-calling,

mitigating the impact of technical artifacts. Second, with over 250,000 chromosomes in

gnomAD, the database provides sufficient sample sizes to estimate haplotype frequencies

below 1×10^{-5}. Lastly, gnomAD offers significant genetic ancestral diversity, allowing results of

our study to be applied beyond samples with European genetic ancestry.

We focus in this study on pairs of rare coding variants occurring in the same gene, which

are of the greatest interest in the context of Mendelian conditions. We required both variants to

have a global minor allele frequency in gnomAD exomes <5% and required variants to be

coding, flanking intronic (from position -1 to -3 in acceptor sites, and +1 to +8 in donor sites) or

in the 5'/3' UTRs. This encompassed 20,921,100 pairs of variants across 19,685 genes. We

performed estimates based on all exome sequencing samples in gnomAD v2, as well as

separate estimates within each of seven genetic ancestry groups, which ranged from 5,076

individuals of Ashkenazi Jewish genetic ancestry to 63,369 for non-Finnish European genetic

ancestry.

For each pair of variants, we first generated pairwise genotype counts in gnomAD, with

nine possible pairwise genotypes for each pair of variants (**Figure 1A**). We then applied the

Expectation-Maximization (EM) algorithm to each pair of variants to generate haplotype

frequency estimates based on the observed pairwise genotype counts[17]. For a given pair of

variants observed in an individual, the probability of two variants being in *trans* (P_{trans}) is the

probability of inheriting each of the haplotypes that contain only one of the two variant alleles.

Validation of phasing estimates using trio data

To estimate the accuracy of our approach, we analyzed variants in a set of 4,992 trios

that were exome sequenced and jointly processed with gnomAD, but which are independent of

the samples included in the gnomAD release. The trio structure allowed us to accurately

determine phase by transmission in these samples. We estimated the genetic ancestry of each

individual in the trios by projecting on the principal components of ancestry in the gnomAD v2

samples (**Figure S1**). In order to maximize the number of variant pairs for testing, we utilized

the variant pairs observed in either parent and used the child's genotype to phase the parental

genotypes. Separately, we used our population-based method leveraging gnomAD data to

estimate phase for every pair of rare coding variants observed in either of the parents. Across

the 4,992 trios, we identified 434,326 pairs of variants where both variants were observed at

least once in gnomAD and thus were amenable to phasing using our approach. 62,363 pairs of

variants were seen in multiple individuals, resulting in 371,963 unique pairs of variants. As

described above, we calculated the probability of being in *trans* (P_{trans}) based on the haplotype

frequency estimates in gnomAD. We found a bimodal distribution of P_{trans} scores; that is, the

majority of probabilities were either very high (>0.99; suggesting a high likelihood of being in

trans), or they were very low (<0.01; suggesting a high likelihood of being in *cis*) (**Figure S2**).

Using the trio phasing-by-transmission data as a gold standard, we generated receiver-

operator curves for distinguishing whether a variant pair is likely in *trans* and found that our

approach achieved high sensitivity and specificity (AUC ranging from 0.905 to 0.998 across the

component genetic ancestry groups) (**Figure 1B**) and high precision and recall (**Figure S3**).

We next defined P_{trans} thresholds for classifying variants as being in *cis* versus *trans* (see

Methods for additional details). To set these thresholds, we first binned variant pairs observed in

the trio data based on their P_{trans}. For each bin, we calculated the proportion of trio variant pairs

that were in *cis* or *trans* based on trio phasing-by-transmission. The P_{trans} threshold for variant

pairs in *trans* was defined as the minimum P_{trans} such that ≥90% of variant pairs in that bin were

in *trans* based on trio phasing-by-transmission. Similarly, the P_{trans} threshold for variants in *cis*

was defined as the maximum P_{trans} such that ≥90% of variant pairs in that bin were in *cis* based

on trio phasing by transmission. This resulted in P_{trans} thresholds of ≤0.10 and ≥0.55 as the

threshold for variants in *cis* and *trans*, respectively (**Figure 1C**).

Based on these P_{trans} thresholds, the overall phasing accuracy was 93.5%. The accuracy

for variants that are in *cis* based on trio data was 92.1%, and the accuracy for variants in *trans*

was 94.7%. Only 6.72% of variants had indeterminate P_{trans} (0.10< P_{trans} <0.55). Together, these

results suggest that our approach can generate accurate phasing estimates.

Figure 1: **A)** Schematic of phasing approach. **B)** Receiver-operator curves for use of P_{trans} for distinguishing between variant pairs on same versus opposite haplotypes. Separate lines are shown for each genetic ancestry group. **C)** (bottom) Proportion of variant pairs in each P_{trans} bin that are on opposite haplotypes. Each point represents variant pairs with P_{trans} bin size of 0.01. Blue dashed line at 10% indicates the P_{trans} threshold at which ≥90% of variant pairs in bin are on the same haplotype (P_{trans} ≤0.10). Red dashed line at 90% indicates the P_{trans} threshold at which ≥90% of variant pairs in bin are on opposite haplotypes (P_{trans} ≥0.55). Calculations are performed using variant pairs where each variant in the pair has allele count > 100. P_{trans} estimates are population-specific. (top) Histogram showing number of variant pairs in each P_{trans} bin that are on the same haplotype (red) or opposite haplotypes (blue).

Accuracy of phasing across allele frequencies

Since the variants that are most likely to be of interest in clinical genetics are rare, we

assessed the accuracy of phasing at different allele frequency (AF) bins. Using P_{trans} estimates

derived from non-Finnish European samples, we found high accuracy (i.e., proportion correct

classifications) ranging from 0.76 to 0.99 across pairs of AF bins (**Figure 2**). In general,

accuracy remained high across allele frequencies for variant pairs in *trans* based on trio phasing

data. For variant pairs in *cis* based on trio phasing data, accuracy was high when variant pairs

were of similar allele frequencies. However, accuracy was much lower, as low as 0.62, for

variant pairs where one variant was much more common than the other. Variant pairs where

both variants are singletons were phased well for variants in *trans* based on the trio phasing

data (0.97). Given the lack of information, we do not report the phasing estimates for

singleton/singleton variant pairs in *cis* (see Discussion).

Figure 2

Figure 2: Phasing accuracy at different allele frequency bins for all variants (left), variants on

opposite haplotypes (middle), and same haplotype (right). Shading of squares and numbers in

each square represent the accuracy. Y-axis labels refer to the more frequent variant in each

variant pair. X-axis labels refer to the rarer variant in each variant pair. Here, accuracy is the

proportion of correct classifications (i.e., correct classifications / all classifications). Accuracies

are calculated for variants seen in non-Finnish European samples based on population-specific

P_{trans} calculations.

Accuracy of phasing across genetic ancestry groups

In the above analyses, we used P_{trans} estimates calculated from the genetic ancestry in

which the variant was seen in the trio data. We next asked if using all samples in gnomAD to

calculate P_{trans} ("cosmopolitan") would improve accuracy given larger sample sizes from which

to calculate P_{trans}, with the caveat that using the full set of gnomAD samples would result in

some genetic ancestry mismatching. Across genetic ancestry groups, we found that accuracy

was similar when using population-specific ancestry estimates as compared to cosmopolitan

estimates (**Figure 3**). However, for certain genetic ancestry groups such as African (AFR) and

East Asian (EAS), accuracy was notably lower when using cosmopolitan estimates as

compared to population-specific estimates specifically for variants in *trans* in these populations;

for example, the phasing accuracy for variants in trans in the AFR ancestry group is close to 1

when using AFR-specific P_{trans} estimates, but drops to ~0.5 when cosmopolitan P_{trans} estimates

are used. These results suggest that cosmopolitan estimates generally achieve similar accuracy

as population-specific estimates, though we do identify certain scenarios where more caution is

required.

Figure 3

Figure 3: Phasing accuracy using population-specific P_{trans} estimates (light blue) or

cosmopolitan P_{trans} estimates (medium blue). Accuracies are shown separately for variants in

trans (left) and variants in *cis* (right)

Effect of variant distance and mutation rates on phasing accuracy

One factor that is predicted to affect the accuracy of our phasing estimates is the

frequency of meiotic recombination between pairs of variants, as recombination events would

disrupt the haplotype configuration of the variant pairs. To further explore the impact of

recombination, we plotted the accuracy of our phasing estimates as a function of physical

distance between variant pairs. We found that for variants in *trans,* the accuracy of phasing was

maintained across physical distances. However, for variant pairs in *cis,* the accuracy rapidly

decreased with longer physical distances (**Figure 4A**). Nonetheless, within the scale of typical

gene lengths (less than 100,000 base pairs), phasing accuracy was relatively preserved.

Since physical distance is only a proxy for recombination frequency, we also performed

this analysis using interpolated genetic distances (**Figure 4B**). We found again that variants in

trans had preserved phasing accuracy across genetic distances, while variants in *cis* had

phasing estimates that decreased substantially with genetic distance, particularly at distances

greater than 0.1 centiMorgan.

We also tested the effect of recombination by examining a set of 20,319 multinucleotide

variants (MNVs), which are pairs of genetic variants in *cis* that are very close together in

physical distance (≤2 bp) and thus have minimal opportunity for recombination between them.

These variants have previously been accurately phased using physical read data[18,19]. When

examining this set of MNVs, we found that the phasing accuracy using our approach was

98.9%, with only 0.60% that were phased incorrectly (the remaining 0.45% had indeterminate

phasing estimates). We found that the vast majority of MNVs that were phased incorrectly using

P_{trans} had disparate AFs between the two variants, similar to what we had observed above (data

not shown).

Figure 4

Figure 4: A) Phasing accuracy (y-axis) as a function of physical distance (in base pairs on $\log_{10}$

scale) between variants (x-axis). Red represents variants on the same haplotype, and blue

represents variants on opposite haplotypes. **B)** Same as **A,** except the x-axis shows genetic

distance (in centiMorgans). Accuracies for **A)** and **B)** are based on variant pairs seen in all

populations and utilize population-specific P_{trans} estimates.

Another potential driver of inaccurate phasing is recurrent germline mutations, which

depend in part on the type of mutation. Transitions have higher mutation rates than

transversions[2,20]. Furthermore, CpG transitions have the highest mutation rates among single

nucleotide changes, with mutation rates increasing with higher methylation rates at the CpG

sites[16]. To better understand the impact of mutation rates, we classified each single nucleotide

variant (SNV) in the trio data as a transversion, non-CpG transition, or CpG transition. For CpG

transitions, we further classified the SNV as having low, medium, or high DNA methylation as

before[16]. We then calculated phasing accuracy as a function of combinations of mutation types

using the trio data (**Figure S4**). We found similar accuracy for transversions and transitions

(~0.96). However, we found that CpG sites had lower phasing accuracy, particularly for variant

pairs in *cis.* The phasing accuracies were lower at medium and high methylation CpG sites

(0.88-0.94) than they were for low methylation sites (0.93-0.97). These results are consistent

with recurrent mutations contributing to inaccurate phasing estimates.

Demonstration in a cohort of patients with Mendelian disorders

To demonstrate our approach in a clinically relevant situation, we turned to a set of 627

patients from the Broad Institute Center for Mendelian Genetics (CMG)[21]. All patients had a

confident or strong candidate genetic diagnosis of a Mendelian condition based on carrying two

rare variants in a recessive disease gene consistent with the patient's phenotype. Across the

627 diagnoses, we were able to estimate phase for the 293 patients in which both variants were

present in gnomAD (**Supplemental Table 1**). Of these 293 variant pairs, our phasing method

indicated 279 (95.2%) to be in *trans*, nine (3.1%) to be in *cis*, and five (1.7%) were

indeterminate (P_{trans} >0.10 and <0.55 or singleton-singleton variant in the same individual). Of

the nine variant pairs indicated to be in *cis*, six originated from patients with proband-only

sequencing. For these patients, the responsible clinician was contacted to ensure phenotype

overlap with the disease gene and to pursue parental Sanger sequencing for confirmatory

phasing by transmission, where possible. The remaining three variant pairs indicated to be in *cis*

originated from patients with trio-sequencing, which confirmed the variants to be in *trans*, and

our inferred phase to be incorrect. Despite a small number of errors, overall, the results suggest

that our phasing approach is highly accurate in clinical scenarios in patients with suspected

Mendelian conditions and can be informative for a large fraction (just under 50% in our cohort)

of candidate diagnoses, dependant on presence of both variants in gnomAD.

Bi-allelic predicted damaging variants

Next, we tabulated for each gene the number of individuals in gnomAD who carry two

rare heterozygous variants, stratified by the predicted phase (e.g., in *trans,* unphased, and in

cis), allele frequency, and the predicted functional consequence of the least damaging variant in

the pair. For comparison, we tabulated individuals with homozygous variants in the same

manner. We classified predicted functional consequences as predicted loss-of-function (pLoF),

missense with deleteriousness scored by REVEL[22] in line with recent ClinGen

recommendations[23], and synonymous. These data are available for gnomAD v2 as a

downloadable table with the counts of individuals by phase for each gene across a total of 26

consequences and five allele frequency thresholds.

Overall, the number of individuals with rare, compound heterozygous (in *trans*),

predicted damaging variants was low (median 0 individuals per gene with compound

heterozygous loss-of-function variants at 1% allele frequency, range 0-9) and only occurred in a

small number of genes (**Figure 5** and **Figure S5**). Twenty eight genes carried compound

heterozygous pLoF variants (in 56 individuals) and an additional four genes carried compound

heterozygous variants with at least a strong REVEL missense predicted consequence (in six

individuals) at 1% allele frequency. The vast majority of these genes have not, to date, been

associated with disease (**Figure 5B**). Manual curation of the pLoF variants resulted in seven

high confidence "human knock-out" genes (*ARHGEF37, CCDC66, FAM81B, FYB2, GNLY,*

RBKS, and *SDSL*). These genes are not associated with Mendelian disease nor are they known

to be essential (see Methods for additional details). In the remaining 21 of the 28 genes with

compound heterozygous pLoF variants, true loss-of-function was found to be uncertain or

unlikely following manual curation, due, for example, to the variant falling in the last exon of the

gene, in a weakly conserved exon, or in the minority of transcripts (**Supplemental Table 2**). We

previously manually curated all homozygous pLoF variants in gnomAD[16]. The absence of rare

compound heterozygous "human knock-out" events in essential genes was expected given that

gnomAD is largely depleted of individuals with severe, early-onset conditions.

A

B

Figure 5: A) Proportion of genes with one or more individuals in gnomAD carrying predicted compound heterozygous (in *trans*) variants or a homozygous variant at ≤1% and ≤5% allele frequency stratified by predicted functional consequence. **B)** Number of genes with ≥1 individual in gnomAD carrying compound heterozygous (in *trans*) or homozygous predicted damaging variants at ≤1% allele frequency, stratified by predicted functional consequence and Mendelian disease-association in the Online Mendelian Inheritance in Man database (OMIM). In total, 28 genes (25 non-disease, 2 AD, and 1 AR) carried predicted compound heterozygous loss-of-function variants at ≤1% allele frequency, only seven of which were high confidence "human knock-out" events following manual curation. For predicted compound heterozygous variants, both variants in the variant pair must be annotated with a consequence at least as severe as the consequence listed (i.e., a compound heterozygous loss-of-function variant would be counted under the LoF category but also included with a less deleterious variant under the other

categories). All homozygous pLoF variants previously underwent manual curation as part of

Karczewski et al[16]. AF, allele frequency; comp het, compound heterozygous; hom, homozygous;

AD, autosomal dominant; AR, autosomal recessive.

Generation of public resource

To make our resource widely usable to both clinicians and researchers, we have

calculated and released pairwise genotype counts and phasing estimates for each pair of rare

coding variants occurring in the same gene for gnomAD. These genotype counts are phasing

estimates shown for all pairs of variants within a gene where both variants have global minor

allele frequency (or population-specific frequency) in gnomAD exomes <5%, and are either

coding, flanking intronic (from position -1 to -3 in acceptor sites, and +1 to +8 in donor sites) or

in the 5'/3' UTRs. We have integrated these data into the gnomAD browser so that clinicians

and researchers can easily look up a pair of variants to obtain the genotype counts, haplotype

frequency estimates, P_{trans} estimates, and likely co-occurrence pattern (**Figure 6A**). These

results are shown for each individual genetic ancestry group and across all genetic ancestries in

gnomAD v2. In addition, the data are available as a downloadable table for all variants.

A

Overview

Population	Samples consistent with variants appearing on different haplotypes	Samples consistent with variants appearing on the same haplotype	Samples consistent with either co-occurrence pattern	Likely co-occurrence pattern
African/African-American	1,060	0	3	Different haplotypes
Latino/Admixed American	300	0	1	Different haplotypes
Ashkenazi Jewish	66	0	0	Different haplotypes
East Asian	13	0	0	No prediction*
European (Finnish)	697	0	0	No prediction*
European (non-Finnish)	1,283	0	0	Different haplotypes
Other	76	0	0	Different haplotypes
South Asian	31	0	0	Different haplotypes
All	**3,526**	**0**	**4**	**Different haplotypes**

Note Only samples covered at both variant sites are included in this table.

* A likely co-occurrence pattern cannot be calculated in some cases, such as when only one of the variants is observed in a population.

Details

Select a population in the overview table to view genotype counts for that population.

Genotype Counts

1-55523855-G-A

		G/G	G/A	A/A
	G/G	94,121	1,180	44
1-55505647-G-T	G/T	2,273	4	0
	T/T	29	0	0

■ Samples consistent with variants appearing on different haplotypes.
■ Samples consistent with variants appearing on the same haplotype.
■ Samples consistent with either co-occurrence pattern.

Based on their co-occurrence pattern in gnomAD, these variants are likely found on different haplotypes in most individuals in gnomAD.

Note Only samples covered at both variant sites are included in this table.

Estimated Haplotype Counts

1-55523855-G-A

		G	A
1-55505647-G-T	G	191,695	1,272
	T	2,335	0

The estimated probability that these variants occur in different haplotypes is >99%.

Note Probability values are not well calibrated, particularly where both variants are extremely rare. Interpret with caution. Please see our blog post on variant co-occurrence for accuracy estimates and additional detail.

B

Figure 6: A) Sample gnomAD browser output for two variants (1-55505647-G-T and 1-

55523855-G-A) in the gene *PCSK9*. On the top, a table subdivided by genetic ancestry group

displays how many individuals in gnomAD from that genetic ancestry are consistent with the two

variants occurring on different haplotypes (*trans*), and how many individuals are consistent with

their occurring on the same haplotype (*cis*). Below that, there is a 3x3 table that contains the 9

possible combinations of genotypes for the two variants of interest. The number of individuals in

gnomAD that fall in each of these combinations are shown and are colored by whether they are

consistent with variants falling on different haplotypes (red) or the same haplotype (blue), or

whether they are indeterminate (purple). The estimated haplotype counts for the four possible

haplotypes for the two variants as calculated by the EM algorithm is displayed on the bottom

right. The probability of being in *trans* for this particular pair of variants is > 99%. **B)** Variant co-

occurrence tables on the gene landing page. For each gene (*GBA1* shown), the top table lists

the number of individuals carrying pairs of rare heterozygous variants by inferred phase, allele

frequency, and predicted functional consequence. The number of individuals with homozygous

variants are tabulated in the same manner and presented as a comparison below. Allele

frequency thresholds of ≤ 5%, ≤ 1%, and ≤ 0.5% are displayed across six predicted functional

consequences (combinations of pLoF, various evidence strengths of predicted pathogenicity for

missense variants, and synonymous variants). Both variants in the variant pair must be

annotated with a consequence at least as severe as the consequence listed (i.e., pLoF + strong

missense also includes pLoF + pLoF).

Furthermore, on the landing page of each gene in the gnomAD v2 browser, we have

incorporated counts tables detailing the number of individuals carrying two rare variants

stratified by allele frequency, and functional consequence. The first table counts individuals

carrying two rare heterozygous variants by predicted phase (in *trans*, unphased, and in *cis*) and

the second table counts individuals carrying homozygous variants (**Figure 6B**). We envision

that these data will aid the medical genetics community in interpreting the clinical significance of

co-occurring variants in the context of recessive conditions. The data for all genes are also

available as a downloadable table within gnomAD v2.

Discussion

In this work, we have leveraged a large exome sequencing cohort to estimate haplotype

frequencies for pairs of rare variants within genes, and demonstrate that these haplotype

frequency estimates can be utilized to predict phase of pairs of variants. Overall, we achieve

high accuracy across a range of allele frequencies and across genetic ancestries and

demonstrate that our approach is able to distinguish variants that are likely compound

heterozygous in a clinical setting. Finally, we freely disseminate our results in an easy-to-use

browser for the community.

Our phasing approach was accurate across a range of allele frequencies (even for

singleton variants) and across genetic ancestry groups. However, we identified meiotic

recombination as one key driver of decreased phasing accuracy. We found that for variants in

cis, phasing accuracy diminished linearly with genetic distance, consistent with the role of

meiotic recombination in decreasing phasing accuracy. However, for variants in *trans,* phasing

accuracy was maintained across genetic and physical distances between pairs of variants. This

result is intuitive, as for rare variants, a recombination event is much more likely to disrupt a

haplotype comprised of two variants than to bring two rare variants onto the same haplotype.

These results suggest the need for caution when using these phasing estimates for variants at

long distances from each other where the phasing prediction is *cis.*

We also compared population-specific estimates with phasing estimates derived from

samples across all genetic ancestry groups in gnomAD v2 ("cosmopolitan"). While population-

specific phasing estimates are more likely to match the haplotypes seen in a given individual,

they utilize information from fewer samples in gnomAD. We found that, in general, population-

specific estimates were similar in accuracy to using cosmopolitan estimates. For individuals of

African genetic ancestry, however, we found that use of cosmopolitan estimates resulted in

much lower phasing accuracy than the use of African-specific estimates. This is consistent with

the observation that there are more unique haplotypes seen in individuals of African genetic

ancestry and/or older haplotypes in individuals of African genetic ancestry for which

recombination is more likely to have occurred[24]. Moreover, there are other genetic ancestry

groups not currently represented in gnomAD for which we expect this phasing approach to have

lower accuracy than in the well-represented populations.

Pairs of singleton variants pose a unique challenge. When a pair of singleton variants is

observed in different individuals in a population, this provides evidence that the variants are on

different haplotypes. However, if a pair of singleton variants is observed in the same individual

in the population, we cannot readily distinguish whether the variants are on the same haplotype

or different haplotypes as we lack information from other individuals in the population for

singleton variants. For this reason, we have chosen to not report phasing estimates for singleton

variant pairs that are observed in the same individuals in gnomAD. Nonetheless, using our trio

data, 93% of these singleton variant pairs observed in the same individual in gnomAD were in

cis based on our trio validation data.

Our work focuses on the challenging scenario of determining phase for rare variants

identified during sequencing of rare disease patients in the absence of parental data. Exome

and targeted gene panel sequencing poses a unique challenge for population-based phasing

given the sparsity of variants, precluding the use of common non-coding variants as a "scaffold"

for phasing. One recent work performed population-based phasing of rare variants from exome

sequencing data by combining the exome data with SNP genotyping arrays[25]. However, SNP

genotyping data are not usually generated in conjunction with a sequencing test. Given this

limitation of population-based phasing of exome sequencing data, we chose to apply the EM

algorithm on pairs of rare variants. Rare variants, which are of the greatest interest in Mendelian

diseases, are also challenging to phase using population-based approaches given the small

numbers of shared haplotypes from which to make phasing estimates in the population. Recent

methods have shown accurate phasing of rare variants using genome sequencing data[25–27], but

relies on a large genome reference panel. In our work, there were limited numbers of genome

sequences available for use in a population-based phasing approach. However, as the numbers

of genome sequencing samples increases in future releases of gnomAD, this may represent a

tractable and more accurate approach for phasing of rare variants.

Utilizing phase estimates to tabulate the number of individuals in gnomAD with two rare

predicted in *trans* variants by gene, we found that there are only a small number of "human

knock-out" genes affected by predicted compound heterozygous (in *trans*) loss-of-function

variants, and that this number is substantially lower than is observed for homozygous loss-of-

function variants. These compound heterozygous "human knock-out" events occurred in genes

that are not known to be essential, an expected finding given that gnomAD is largely depleted of

individuals with severe and early-onset conditions. When analyzing the 23,667 individuals that

carry two pLoF variants with allele frequency ≤ 1%, we predict that in 20,706 (88%) of those

individuals, the variant pair is in *cis* and only a small fraction (~0.2%) are confidently predicted

to be in *trans*. This may be counter-intuitive, so warrants emphasis: when a pair of rare pLoF

variants is observed in the same gene in an individual from a general population sample, it is

vastly more likely that these variants are carried on the same haplotype than that the individual

is a genuine "knock-out" due to compound heterozygosity.

To aid the medical genetics community in interpreting the clinical significance of rare co-

occurring variants in the context of recessive disease, we have released these counts across a

spectrum of variant consequences (loss-of-function, missense, and synonymous) and allele

frequencies by gene in the gnomAD browser. These provide background frequencies–to the

best of our ability–of compound heterozygous rare damaging variants. These background

frequencies can be used to assess the probability that a given variant pair identified in a patient

may have occurred by chance. We note, however, that our ability to identify rare variant pairs in

trans in gnomAD v2 is limited by the fact this dataset was used for training. We find that our

ability to detect in *trans* variant pairs extends only to an allele frequency of 0.5%, with all rarer

combinations being dominated by unphased predictions and very few predictions of *in trans*

variant pairs. The per-gene variant co-occurrence resource developed and released here is

therefore to be considered a first step in this space. We plan to use the predictions from this

dataset (gnomAD v2) on newer versions of gnomAD with additional samples, where we can

confidently predict rare variant pairs that are in *trans*.

Beyond our current restriction to predicting within the gnomAD dataset, there are several

other important limitations to our work. First, we have only reported phasing estimates for rare

coding and intronic flanking/UTR variant pairs within genes in order to limit the computational

burden. We believe that these are the variant pairs of most interest to the medical genetics

community, but acknowledge that phase with deeper intronic variation will become important as

more genome sequencing is performed. Second, while there was a broad range of genetic

ancestral diversity represented in our samples, future studies would benefit from even larger

sample sizes, especially for genetic ancestry groups not well represented in our present study.

Finally, we have only tested our phasing accuracy in a clinical setting in a retrospective manner

and future prospective studies will be needed to confirm the clinical utility of our approach.

Methods

gnomAD characteristics and data processing

In this work, we used exome sequencing data from the gnomAD v2.1 dataset (n =

125,748 individuals). These data were uniformly processed, underwent joint variant calling, and

rigorous quality control, as described in Karczewski et al.[16]. Briefly, we aggregated ~200k

exome sequences and ~20k genome sequences, primarily from case-control studies of

common adult-onset conditions, and applied a BWA-Picard-GATK pipeline[28]. Using Hail

(https://github.com/hail-is/hail), we then removed samples that (1) failed population- and

platform-specific quality control, (2) had second-degree or closer relations in the dataset, (3) did

not have appropriate consent for release, and (4) had known severe, early-onset conditions. For

variant quality control, we trained a random forest on site-level and genotype-level metrics (e.g.,

the quality by depth, QD), and demonstrated that it achieved both high precision and recall for

both common and rare variants.

We subsetted the final cleaned gnomAD dataset for variants with global allele frequency

in gnomAD exomes <5% that were either coding, flanking intronic (from position -1 to -3 in

acceptor sites, and +1 to +8 in donor sites) or in the 5'/3' UTRs. This encompassed 20,921,100

pairs of variants across 19,685 genes.

Analysis in this manuscript was performed using Hail version 0.2.105[29], and analysis

code is available at https://github.com/broadinstitute/gnomad_chets.

Haplotype estimates

Consider two variants, A and B. A and B represent the major alleles, and a and b

represent the respective minor alleles. There are thus 9 pairwise genotypes for A and B: AABB,

AaBB, aaBB, AABb, AaBb, aaBb, AAbb, Aabb, and aabb. Of these pairwise genotypes, only the

phase for the double heterozygote (AaBb) is unknown. From these 9 possible genotypes, there

are four possible haplotype configurations: AB, Ab, aB, and ab.

For each pair of variants, we applied the expectation-maximization (EM) algorithm[17] to

estimate haplotype frequencies from genotype counts. The initial conditions of the EM algorithm

were set by partitioning the doubly heterozygous (AaBb) genotype counts equally between the

AB|ab and Ab|aB haplotype configurations. The EM algorithm was run until convergence or until

the absolute value of the difference between consecutive maximum likelihood function values

was less than 1×10^{-7}. We calculated haplotype frequencies based on genotypes present within

the same population ("population-specific") or using all samples from gnomAD ("cosmopolitan").

Haplotype frequency estimates were performed using Hail.

We then calculate P_{trans} as the likelihood that any given pair of doubly heterozygous

variants (AaBb) in a patient is compound heterozygous (Ab|aB). P_{trans} can be calculated simply

from the haplotype frequency estimates (AB, Ab, aB, and ab):

$$P_{trans} = ((Ab \times aB))/(AB \times ab + Ab \times aB)$$

Thus, P_{trans} simply represents the probability that the patient is compound heterozygous by

inheriting both the Ab and aB haplotypes.

Determination of P_{trans} cutoffs

To determine P_{trans} cutoffs for classifying variants as occurring in *cis* or *trans*, we binned

variant pairs in increments of 0.01 of P_{trans}. For each bin, we calculated the proportion of variant

pairs in that bin that are compound heterozygous based on phasing by trio data. The P_{trans} for

compound heterozygous variants was determined as the minimum P_{trans} such that 90% of

variants in the bin are compound heterozygous based on trio data. The P_{trans} for variants in *cis*

was determined as the maximum P_{trans} such that 90% of variants in the bin are in *cis* based on

trio data. For these calculations, we used only variants where both variants had a population

allele count greater than 100. We used population-specific P_{trans} estimates.

Trio validation data

Part of the original joint calling of gnomAD included exome sequencing data from full

parent-child trios. Since most of these trios came from cohorts where the child had a severe,

early-onset condition, the trios were removed from the final release of gnomAD but were

similarly processed as the released samples. Here, we made use of 4,992 trios that were

processed with gnomAD but not released. Having access to parental information allows us to

perform phase-by-transmission and accurately determine whether two co-occurring variants in

the same gene are in *cis* or in *trans*.

First, we estimated genetic ancestry of each individual in the trios by using ancestry

inference estimates from the full gnomAD dataset, as previously described[16]. Briefly, we

selected bi-allelic variants that passed all hard filters, had allele frequencies in a joint exome

and genome callset > 0.001, and high joint call rates (> 0.99). The variants were then LD-

pruned ($r^2 = 0.1$) and used in a principal component analysis (PCA). We previously used

samples with known genetic ancestry to train a random forest on the first 20 principal

components (PCs), and assigned samples to a genetic ancestry group based on having a

random forest probability > 0.9. For the trios in this cohort, we projected their PCs for genetic

ancestry onto the same gnomAD v2 samples to infer the population used here (**Figure S1**).

We then phased the trio data using the Hail *phase_by_transmission*

(https://hail.is/docs/0.2/experimental/index.html#hail.experimental.phase_by_transmission)

function, which uses Mendelian transmission of alleles to infer haplotypes in trios for all sites

that aren't heterozygous in all members of the trio. Assigning haplotypes in trios based on

parental genotype has traditionally been the gold standard, has switch error rates below 0.1%,

and importantly errors aren't dependent on the allele frequency of the variants phased[30]. To

maximize our confidence in the genotypes and phasing, we filtered genotypes to include only

those with genotype quality (GQ) > 20, depth > 10 and allele balance > 0.2 for heterozygous

genotypes prior to phasing. Sex chromosomes were excluded. Using these criteria, we ended

up with 434,326 variant pairs where both of the variants were observed at least once in

gnomAD and could thus be used to evaluate accuracy of our method.

CpG analysis

CpG analysis

Single nucleotide variants seen in the trio data were divided into transitions and

transversions. Transitions were further subdivided into those that are CpG mutations (5'-CpG-3'

mutating to 5'-TpG-3') and those that are not. For each CpG transition, we calculated the mean

DNA methylation values across 37 tissues in ENCODE[16]. We then stratified CpG transitions into

3 levels: low (missing or < 0.2), medium (0.2–0.6), and high (>0.6) methylation. Phasing

accuracy–here, the proportion correct (correct classifications/all classifications)–was then

calculated for pairwise combinations of transversions, non-CpG transitions, low methylation

CpG transitions, medium methylation CpG transitions, and high methylation CpG transitions. All

SNVs were included in the analysis and population-specific EM estimates were used.

Calculating accuracy as a function of genetic distance

To estimate the genetic distance between pairs of genetic variants, we interpolated

genetic distances between variant pairs using a genetic map from HapMap2[31]

(https://github.com/joepickrell/1000-genomes-genetic-maps). We downloaded a pre-generated

HapMap2 genetic map representing average over recombination rates in the CEU, YRI, and

ASN populations. We then ran interpolate_maps.py (downloaded from

https://github.com/joepickrell/1000-genomes-genetic-

maps/blob/master/scripts/interpolate_maps.py) for all variant pairs in the phasing trio data. As

above, accuracy is the proportion of correct classifications.

MNV analysis

We obtained multi-nucleotide variant pairs for which read-back phasing had previously

been calculated[18]. We included only multi-nucleotide variant pairs where each constituent

variant was analyzed in our study. Phasing estimates were calculated using cosmopolitan EM

estimates.

Rare disease patient analysis

627 patients from the Broad Institute Center for Mendelian Genetics (CMG)[21] with a

confident or strong candidate genetic diagnosis of a Mendelian condition were selected for

analysis. Each patient carried two presumed bi-allelic variants in an autosomal recessive

disease gene consistent with the patient's phenotype. For 293 of the patients, both variants

were present in gnomAD and phase was predicted. Trio-sequencing (i.e., sequencing of the

proband and the two unaffected biological parents) had been performed for 168 of the 293

patients. For fully sequenced trios, we were able to confirm phasing of the two variants via

phase-by-transmission.

Determining counts of individuals with two rare, damaging variants

Variants were annotated with the worst consequence on the canonical transcript by the

Ensembl Variant Effect Predictor (VEP)[32]. Putative LoF variants were annotated with LOFTEE[16],

and only high confidence LoF variants were counted as "pLoF". Missense variants were

annotated with REVEL[22]. REVEL scores ≥0.932 were counted as "strong_revel_missense",

≥0.773 as "moderate_to_strong_revel_missense", ≥0.644 as

"supporting_to_strong_revel_missense" in line with recent ClinGen recommendations[23].

Variant pairs were annotated with predicted phase based on the P_{trans} thresholds. All

singleton-singleton variant pairs (AC = 1) and variant pairs with an indeterminate P_{trans} values

(>0.1 and <0.55) were annotated as unphased.

Five allele frequency thresholds were selected for analysis and variant pairs were filtered

based on the highest global allele frequency and, where available, the "popmax" allele

frequency of each variant in gnomAD (i.e., the highest allele frequency information for the non-

bottlenecked population - excluding Ashkenazi Jewish, European Finnish, and "Other"

populations): 0.5%, 1%, 1.5%, 2%, and 5%. Further, all variant pairs containing a variant with an

allele frequency >5% in a bottlenecked population were filtered out.

The number of individuals carrying a variant pair (irrespective of phase) and the number

indicated to be compound heterozygous (in *trans*), unphased (indeterminate), and on the same

haplogroup (in *cis*) were counted gene-wise by allele frequency threshold and combined

functional consequences (26 consequences). This counting was repeated twice, once restricting

individuals to be counted in only one phase group, prioritizing in *trans* over unphased and

unphased over in *cis* (displayed in the "variant co-occurrence" gnomAD browser feature), and

once allowing individuals to be counted in multiple phase groups, if carrying multiple variant

pairs in the same gene with different phase predictions.

Essential gene lists

The following essential gene lists were queried for the presence of the true "human

knock-out" genes identified in this study:

● 2,454 genes essential in mice from Georgi et al. 2013[33]

● 553 pan-cancer core fitness genes from Behan et al., 2019[34]

- 360 core essential genes from genomic perturbation screens from Hart et al. 2014[35]
- 684 genes essential in culture by CRISPR screening from Hart et al. 2017[36]
- 1,075 genes annotated by the ADaM analysis of a large collection of gene dependency profiles (CRISPR-Cas9 screens) across human cancer cell lines from Vinceti et al. 2021[37]

Code

The code used to estimate P_{trans} estimates for variant pairs and to determine the number of individuals carrying rare, compound heterozygous variants can be found at:

https://github.com/broadinstitute/gnomad_chets

Data availability

We provide both web-based look up tools and downloads for the data generated here. A look-up tool to find the likely co-occurrence pattern between two rare (global allele frequency in gnomAD exomes <5%) coding, flanking intronic (from position -1 to -3 in acceptor sites, and +1 to +8 in donor sites) or 5'/3' UTR variants can be found at:

https://gnomad.broadinstitute.org/variant-cooccurrence

Additionally, we display the per-gene counts tables that detail the number of individuals with two rare variants, stratified by allele frequency and functional consequence, on each gene's main page. One table details counts of individuals with two heterozygous variants and includes predicted phase, while the second details individuals with homozygous variants. Both can be found by clicking on the "Variant Co-occurrence" tab on each gene's main page.

All variant co-occurrence tables can be downloaded from:

https://gnomad.broadinstitute.org/downloads#v2-variant-cooccurrence

Author contributions

M.H.G., L.C.F., S.L.S., J.K.G., A.O., K.J.K., D.G.M., and K.E.S. conceived and designed experiments. M.H.G., L.C.F., S.L.S., and J.K.G. performed the analyses. N.A.W., P.W.D., and M.S. developed visualizations for the web browser. E.G. and M.S-B. performed variant curation. G.T., B.M.N., J.N.H., H.L.R., M.J.D., A.O., and K.J.K. provided data and analysis suggestions. J.N.H., D.G.M., and K.E.S. supervised the work. M.H.G., L.C.F., S.L.S., J.K.G., and K.E.S.

completed the primary writing of the manuscript with input and approval of the final version from

all other authors.

Acknowledgements

We would like to thank all members of the gnomAD team for helpful comments and

suggestions, and to particularly recognize the members of the gnomAD methods and browser

teams who worked hard over many years to provide cleaned datasets, easy-to-use browsers,

and visualizations. This work was supported by NHGRI U24HG011450, UM1HG008900, and

U01HG011755.

Supplementary Figures

Figure S1: Principal component analysis (PCA) plot for the full gnomAD v2 cohort (left) and specifically for the trios (right, trios in black) included in this paper. The top row shows PC1 vs PC2, the middle row shows PC3 vs PC4, and the bottom row shows PC5 vs PC6. Genetic ancestry group labels for the global gnomAD populations were done as described in Karczewski et al. 2020[16].

Figure S2: Histogram of P$_{trans}$ values for variant pairs on the same haplotype (top) or opposite haplotypes (bottom). P$_{trans}$ bin size of 0.01.

Figure S3: Precision-recall curves for use of P_{trans} for distinguishing between variant pairs on same versus opposite haplotypes. Separate lines are shown for each ancestry.

Figure S4: Phasing accuracy for transversions, non-CpG transitions, and CpG transitions. CpG transitions are further stratified by degree of DNA methylation (low, medium, or high) as in Karczewski et al 2020. Shading of squares and numbers in each square represent phasing accuracy. Phasing accuracies are based on variant pairs seen in all populations and utilize population-specific P_{trans} estimates.

Figure S5: A) Proportion of genes with one or more individuals in gnomAD carrying two rare variants at ≤1% and ≤5% allele frequency stratified by predicted functional effect and phase. For comp het (in *trans*), unphased, and in *cis*, both variants in the variant pair must be annotated with a consequence at least as severe as the consequence displayed. **B)** Number of individuals per gene in gnomAD carrying two rare variants at ≤1% and ≤5% allele frequency stratified by predicted functional effect and phase. For compound heterozygous (in *trans*) both variants in the variant pair must be annotated with a consequence at least as severe as the consequence displayed. comp het, compound heterozygous; hom, homozygous.

Supplemental Tables

Table S1. CMG diagnostic variants. In this table, we provide details about the presumed bi-allelic causal variants from 293 individuals from the Broad Institute Center for Mendelian Genetics. For each variant pair, we provide the gene symbol ("gene_name"), information about the position and alleles of both variants, whether both of the variants were singletons in gnomAD ("singleton_singleton") and seen in the same individual or not, the estimated P_{trans} value, the predicted phase based on the P_{trans} value ("phase_prediction"), and the known phase from phase by transmission when available.

Table S2. Manual curation results for compound heterozygous loss-of-function variants. Here, we provide the variant curation information for the 28 genes that have predicted compound heterozygous loss-of-function variants with allele frequency ≤ 1%. For every predicted compound heterozygous variant pair, we provide the gene symbol, maximum allele frequency in the gnomAD exomes from the two variants ("variant_pair_max_af"), the number of individuals who carry the variant pair ("n_individuals"), information about the position and alleles of both variants, any manual curation flags e.g., mapping error for the variants, and the final loss-of-function curation for both variants as well as the variant pair ("high_confidence_human_knock_out").

References

1. McVean, G. A. T. *et al.* The fine-scale structure of recombination rate variation in the human genome. *Science* **304**, 581–584 (2004).
2. Rahbari, R. *et al.* Timing, rates and spectra of human germline mutation. *Nat. Genet.* **48**, 126–133 (2016).
3. Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**, 285–291 (2016).
4. Ségurel, L., Wyman, M. J. & Przeworski, M. Determinants of mutation rate variation in the human germline. *Annu. Rev. Genomics Hum. Genet.* **15**, 47–70 (2014).
5. Hodgkinson, A. & Eyre-Walker, A. Variation in the mutation rate across mammalian genomes. *Nature Reviews Genetics* vol. 12 756–766 Preprint at https://doi.org/10.1038/nrg3098 (2011).
6. Carlson, J. *et al.* Extremely rare variants reveal patterns of germline mutation rate heterogeneity in humans. *Nat. Commun.* **9**, 3753 (2018).
7. Wang, Q. *et al.* Landscape of multi-nucleotide variants in 125,748 human exomes and 15,708 genomes. *Nat. Commun.* **11**, 2539 (2020).
8. Bansal, V., Halpern, A. L., Axelrod, N. & Bafna, V. An MCMC algorithm for haplotype assembly from whole-genome sequence data. *Genome Res.* **18**, 1336–1346 (2008).
9. Patterson, M. *et al.* WhatsHap: Weighted Haplotype Assembly for Future-Generation Sequencing Reads. *J. Comput. Biol.* **22**, 498–509 (2015).
10. Hager, P., Mewes, H.-W., Rohlfs, M., Klein, C. & Jeske, T. SmartPhase: Accurate and fast phasing of heterozygous variant pairs for genetic diagnosis of rare diseases. *PLoS Comput. Biol.* **16**, e1007613 (2020).
11. Maestri, S. *et al.* A Long-Read Sequencing Approach for Direct Haplotype Phasing in Clinical Settings. *Int. J. Mol. Sci.* **21**, (2020).
12. Mantere, T., Kersten, S. & Hoischen, A. Long-Read Sequencing Emerging in Medical Genetics. *Front. Genet.* **10**, 426 (2019).
13. Snyder, M. W., Adey, A., Kitzman, J. O. & Shendure, J. Haplotype-resolved genome sequencing: experimental methods and applications. *Nat. Rev. Genet.* **16**, 344–358 (2015).
14. Tewhey, R., Bansal, V., Torkamani, A., Topol, E. J. & Schork, N. J. The importance of phase information for human genomics. *Nat. Rev. Genet.* **12**, 215–223 (2011).
15. Browning, S. R. & Browning, B. L. Haplotype phasing: existing methods and new developments. *Nat. Rev. Genet.* **12**, 703–714 (2011).
16. Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature* **581**, 434–443 (2020).
17. Excoffier, L. & Slatkin, M. Maximum-likelihood estimation of molecular haplotype frequencies in a diploid population. *Mol. Biol. Evol.* **12**, 921–927 (1995).
18. Wang, Q. *et al.* Author Correction: Landscape of multi-nucleotide variants in 125,748 human exomes and 15,708 genomes. *Nat. Commun.* **12**, 827 (2021).
19. Kaplanis, J. *et al.* Exome-wide assessment of the functional impact and pathogenicity of multinucleotide mutations. *Genome Res.* **29**, 1047–1056 (2019).
20. Lynch, M. Rate, molecular spectrum, and consequences of human mutation. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 961–968 (2010).
21. Baxter, S. M. *et al.* Centers for Mendelian Genomics: A decade of facilitating gene discovery. *Genet. Med.* **24**, 784–797 (2022).
22. Ioannidis, N. M. *et al.* REVEL: An Ensemble Method for Predicting the Pathogenicity of Rare Missense Variants. *Am. J. Hum. Genet.* **99**, 877–885 (2016).
23. Pejaver, V. *et al.* Evidence-based calibration of computational tools for missense variant pathogenicity classification and ClinGen recommendations for clinical use of PP3/BP4 criteria. *bioRxiv* 2022.03.17.484479 (2022) doi:10.1101/2022.03.17.484479.

24. Mathieson, I. & McVean, G. Demography and the age of rare variants. *PLoS Genet.* **10**, e1004528 (2014).
25. Hofmeister, R. J., Ribeiro, D. M., Rubinacci, S. & Delaneau, O. Accurate rare variant phasing of whole-genome and whole-exome sequencing data in the UK Biobank. *bioRxiv* 2022.10.19.512867 (2022) doi:10.1101/2022.10.19.512867.
26. Sharp, K., Kretzschmar, W., Delaneau, O. & Marchini, J. Phasing for medical sequencing using rare variants and large haplotype reference panels. *Bioinformatics* **32**, 1974–1980 (2016).
27. Browning, B. L., Tian, X., Zhou, Y. & Browning, S. R. Fast two-stage phasing of large-scale sequence data. *Am. J. Hum. Genet.* **108**, 1880–1890 (2021).
28. Van der Auwera, G. A. *et al.* From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinformatics* **43**, 11.10.1–11.10.33 (2013).
29. Hail Team. Hail 0.2.105-acd89e80c345. *GitHub* https://github.com/hail-is/hail/commit/acd89e80c345.
30. Choi, Y., Chan, A. P., Kirkness, E., Telenti, A. & Schork, N. J. Comparison of phasing strategies for whole human genomes. *PLoS Genet.* **14**, e1007308 (2018).
31. International HapMap Consortium *et al.* A second generation human haplotype map of over 3.1 million SNPs. *Nature* **449**, 851–861 (2007).
32. McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol.* **17**, 122 (2016).
33. Georgi, B., Voight, B. F. & Bućan, M. From mouse to human: evolutionary genomics analysis of human orthologs of essential genes. *PLoS Genet.* **9**, e1003484 (2013).
34. Behan, F. M. *et al.* Prioritization of cancer therapeutic targets using CRISPR-Cas9 screens. *Nature* **568**, 511–516 (2019).
35. Hart, T., Brown, K. R., Sircoulomb, F., Rottapel, R. & Moffat, J. Measuring error rates in genomic perturbation screens: gold standards for human functional genomics. *Mol. Syst. Biol.* **10**, 733 (2014).
36. Hart, T. *et al.* Evaluation and Design of Genome-Wide CRISPR/SpCas9 Knockout Screens. *G3* **7**, 2719–2727 (2017).
37. Vinceti, A. *et al.* CoRe: a robustly benchmarked R package for identifying core-fitness genes in genome-wide pooled CRISPR-Cas9 screens. *BMC Genomics* **22**, 828 (2021).

Genome Aggregation Database Consortium

Maria Abreu[1], Carlos A. Aguilar Salinas[2], Tariq Ahmad[3], Christine M. Albert[4,5], Jessica Alföldi[6,7], Diego Ardissino[8], Irina M. Armean[6,7,9], Gil Atzmon[10,11], Eric Banks[12], John Barnard[13], Samantha M. Baxter[6], Laurent Beaugerie[14], Emelia J. Benjamin[15,16,17], David Benjamin[12], Louis Bergelson[12], Michael Boehnke[18], Lori L. Bonnycastle[19], Erwin P. Bottinger[20], Donald W. Bowden[21,22,23], Matthew J. Bown[24,25], Steven Brant[26], Sarah E. Calvo[6,27], Hannia Campos[28,29], John C. Chambers[30,31,32], Juliana C. Chan[33], Katherine R. Chao[6], Sinéad Chapman[6,7,34], Daniel Chasman[4,35], Siwei Chen[6,7], Rex L. Chisholm[36], Judy Cho[20], Rajiv Chowdhury[37], Mina K. Chung[38], Wendy K. Chung[39,40,41], Kristian Cibulskis[12], Bruce Cohen[35,42], Ryan L. Collins[6,27,43], Kristen M. Connolly[44], Adolfo Correa[45], Miguel Covarrubias[12], Beryl Cummings[6,43], Dana Dabelea[46], Mark J. Daly[6,7,47], John Danesh[37], Dawood Darbar[48], Joshua Denny[49], Stacey Donnelly[6], Ravindranath Duggirala[50], Josée Dupuis[51,52], Patrick T. Ellinor[6,53], Roberto Elosua[54,55,56], James Emery[12], Eleina England[6], Jeanette Erdmann[57,58,59], Tõnu Esko[6,60], Emily Evangelista[6], Yossi Farjoun[12], Diane Fatkin[61,62,63], Steven Ferriera[64], Jose Florez[35,65,66], Laurent C. Francioli[6,7], Andre Franke[67], Martti Färkkilä[68], Stacey Gabriel[64], Kiran Garimella[12], Laura D. Gauthier[12], Jeff Gentry[12], Gad Getz[35,69,70], David C. Glahn[71,72], Benjamin Glaser[73], Stephen J. Glatt[74], David Goldstein[75,76], Clicerio Gonzalez[77], Julia K. Goodrich[6], Leif Groop[78,79], Sanna Gudmundsson[6,7,80], Namrata Gupta[6,64], Andrea Haessly[12], Christopher Haiman[81], Ira Hall[82], Craig Hanis[83], Matthew Harms[84,85], Mikko Hiltunen[86], Matti M. Holi[87], Christina M. Hultman[88,89], Chaim Jalas[90], Thibault Jeandet[12], Mikko Kallela[91], Diane Kaplan[12], Jaakko Kaprio[79], Konrad J. Karczewski[6,7,34], Sekar Kathiresan[27,35,92], Eimear Kenny[89,93], Bong-Jo Kim[94], Young Jin Kim[94], George Kirov[95], Zan Koenig[6], Jaspal Kooner[31,96,97], Seppo Koskinen[98], Harlan M. Krumholz[99], Subra Kugathasan[100], Soo Heon Kwak[101], Markku Laakso[102,103], Nicole Lake[104], Trevyn Langsford[12], Kristen M. Laricchia[6,7], Terho Lehtimäki[105], Monkol Lek[104], Emily Lipscomb[6], Christopher Llanwarne[12], Ruth J.F. Loos[20,106], Steven A. Lubitz[6,53], Teresa Tusie Luna[107,108], Ronald C.W. Ma[33,109,110], Daniel G. MacArthur[6,111,112], Gregory M. Marcus[113], Jaume Marrugat[55,114], Alicia R. Martin[6], Kari M. Mattila[105], Steven McCarroll[34,115], Mark I. McCarthy[116,117,118], Jacob McCauley[119,120], Dermot McGovern[121], Ruth McPherson[122], James B. Meigs[6,35,123], Olle Melander[124], Andres Metspalu[125], Deborah Meyers[126], Eric V. Minikel[6], Braxton D. Mitchell[127], Vamsi K. Mootha[6,128], Ruchi Munshi[12], Aliya Naheed[129], Saman Nazarian[130,131], Benjamin M. Neale[6,7], Peter M. Nilsson[132], Sam Novod[12], Anne H. O'Donnell-Luria[6,7,80], Michael C. O'Donovan[95], Yukinori Okada[133,134,135], Dost Ongur[35,42], Lorena Orozco[136], Michael J. Owen[95], Colin Palmer[137], Nicholette D. Palmer[138], Aarno Palotie[7,34,79], Kyong Soo Park[101,139], Carlos Pato[140], Nikelle Petrillo[12], William Phu[6,80], Timothy Poterba[6,7,34], Ann E. Pulver[141], Dan Rader[130,142], Nazneen Rahman[143], Heidi L. Rehm[6,27], Alex Reiner[144,145], Anne M. Remes[146], Dan Rhodes[6], Stephen Rich[147,148], John D. Rioux[149,150], Samuli Ripatti[79,151,152], David Roazen[12], Dan M. Roden[153,154], Jerome I. Rotter[155], Valentin Ruano-Rubio[12], Nareh Sahakian[12], Danish Saleheen[156,157,158], Veikko Salomaa[159], Andrea Saltzman[6], Nilesh J. Samani[24,25], Kaitlin E. Samocha[6,7,27], Jeremiah Scharf[6,27,34], Molly Schleicher[6], Heribert Schunkert[160,161], Sebastian Schönherr[162], Eleanor Seaby[6], Cotton Seed[7,34], Svati H. Shah[163], Megan Shand[12], Moore B. Shoemaker[164], Tai Shyong[165,166], Edwin K. Silverman[167,168], Moriel Singer-Berk[6], Pamela Sklar[169,170,171], J. Gustav Smith[152,172,173], Jonathan T. Smith[12], Hilkka Soininen[174], Harry Sokol[175,176,177], Matthew Solomonson[6,7], Rachel G. Son[6], Jose Soto[12], Tim Spector[178], Christine Stevens[6,7,34], Nathan Stitziel[82,179], Patrick F. Sullivan[88,180], Jaana Suvisaari[159], E. Shyong Tai[181,182,183], Michael E. Talkowski[6,27,34], Yekaterina Tarasova[6], Kent D. Taylor[155], Yik Ying Teo[181,184,185], Grace Tiao[6,7], Kathleen Tibbetts[12], Charlotte Tolonen[12], Ming Tsuang[186,187], Tiinamaija Tuomi[79,188,189], Dan Turner[190], Teresa Tusie-Luna[191,192], Erkki Vartiainen[193], Marquis Vawter[194], Christopher Vittal[6,7], Gordon Wade[12], Arcturus Wang[6,7,34], Qingbo Wang[6,133], James S. Ware[6,195,196], Hugh Watkins[197], Nicholas A. Watts[6,7], Rinse K. Weersma[198], Ben Weisburd[12], Maija Wessman[79,199], Nicola Whiffin[6,200,201], Michael W. Wilson[6,7], James G. Wilson[202], Ramnik J. Xavier[203,204], Mary T. Yohannes[6]

[1]University of Miami Miller School of Medicine, Gastroenterology, Miami, USA
[2]Unidad de Investigacion de Enfermedades Metabolicas, Instituto Nacional de Ciencias Medicas y Nutricion, Mexico City, Mexico
[3]Peninsula College of Medicine and Dentistry, Exeter, UK
[4]Division of Preventive Medicine, Brigham and Women's Hospital, Boston, MA, USA
[5]Division of Cardiovascular Medicine, Brigham and Women's Hospital and Harvard Medical School,

Boston, MA, USA

[6]Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[7]Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA

[8]Department of Cardiology University Hospital, Parma, Italy

[9]European Molecular Biology Laboratory, European Bioinformatics Institute, Wellcome Genome Campus, Hinxton, Cambridge, UK

[10]Department of Biology Faculty of Natural Sciences, University of Haifa, Haifa, Israel

[11]Departments of Medicine and Genetics, Albert Einstein College of Medicine, Bronx, NY, USA

[12]Data Science Platform, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[13]Department of Quantitative Health Sciences, Lerner Research Institute Cleveland Clinic, Cleveland, OH, USA

[14]Sorbonne Université, APHP, Gastroenterology Department Saint Antoine Hospital, Paris, France

[15]NHLBI and Boston University's Framingham Heart Study, Framingham, MA, USA

[16]Department of Medicine, Boston University Chobanian and Avedisian School of Medicine, Boston, MA, USA

[17]Department of Epidemiology, Boston University School of Public Health, Boston, MA, USA

[18]Department of Biostatistics and Center for Statistical Genetics, University of Michigan, Ann Arbor, MI, USA

[19]National Human Genome Research Institute, National Institutes of Health Bethesda, MD, USA

[20]The Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[21]Department of Biochemistry, Wake Forest School of Medicine, Winston-Salem, NC, USA

[22]Center for Genomics and Personalized Medicine Research, Wake Forest School of Medicine, Winston-Salem, NC, USA

[23]Center for Diabetes Research, Wake Forest School of Medicine, Winston-Salem, NC, USA

[24]Department of Cardiovascular Sciences, University of Leicester, Leicester, UK

[25]NIHR Leicester Biomedical Research Centre, Glenfield Hospital, Leicester, UK

[26]John Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

[27]Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA

[28]Harvard School of Public Health, Boston, MA, USA

[29]Central American Population Center, San Pedro, Costa Rica

[30]Department of Epidemiology and Biostatistics, Imperial College London, London, UK

[31]Department of Cardiology, Ealing Hospital, NHS Trust, Southall, UK

[32]Imperial College, Healthcare NHS Trust Imperial College London, London, UK

[33]Department of Medicine and Therapeutics, The Chinese University of Hong Kong, Hong Kong, China

[34]Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[35]Department of Medicine, Harvard Medical School, Boston, MA, USA

[36]Northwestern University Feinberg School of Medicine, Chicago, IL, USA

[37]University of Cambridge, Cambridge, England

[38]Departments of Cardiovascular, Medicine Cellular and Molecular Medicine Molecular Cardiology, Quantitative Health Sciences, Cleveland Clinic, Cleveland, OH, USA

[39]Department of Pediatrics, Columbia University Irving Medical Center, New York, NY, USA

[40]Herbert Irving Comprehensive Cancer Center, Columbia University Medical Center, New York, NY, USA

[41]Department of Medicine, Columbia University Medical Center, New York, NY, USA

[42]McLean Hospital, Belmont, MA, USA

[43]Division of Medical Sciences, Harvard Medical School, Boston, MA, USA

[44]Genomics Platform, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[45]Department of Medicine, University of Mississippi Medical Center, Jackson, MI, USA

[46]Department of Epidemiology Colorado School of Public Health Aurora, CO, USA

[47]Institute for Molecular Medicine Finland, (FIMM) Helsinki, Finland

[48]Department of Medicine and Pharmacology, University of Illinois at Chicago, Chicago, IL, USA

[49]Vanderbilt University Medical Center, Nashville, TN, USA

[50]Department of Genetics, Texas Biomedical Research Institute, San Antonio, TX, USA

[51]Department of Biostatistics, Boston University School of Public Health, Boston, MA, USA

[52]National Heart Lung and Blood Institute's Framingham Heart Study, Framingham, MA, USA

[53]Cardiac Arrhythmia Service and Cardiovascular Research Center, Massachusetts General Hospital,

Boston, MA, USA

[54]Cardiovascular Epidemiology and Genetics Hospital del Mar Medical Research Institute, (IMIM) Barcelona Catalonia, Spain

[55]CIBER CV Barcelona, Catalonia, Spain

[56]Departament of Medicine, Medical School University of Vic-Central, University of Catalonia, Vic Catalonia, Spain

[57]Institute for Cardiogenetics, University of Lübeck, Lübeck, Germany

[58]German Research Centre for Cardiovascular Research, Hamburg/Lübeck/Kiel, Lübeck, Germany

[59]University Heart Center Lübeck, Lübeck, Germany

[60]Estonian Genome Center, Institute of Genomics University of Tartu, Tartu, Estonia

[61]Victor Chang Cardiac Research Institute, Darlinghurst, NSW, Australia

[62]Faculty of Medicine, UNSW Sydney, Kensington, NSW, Australia

[63]Cardiology Department, St Vincent's Hospital, Darlinghurst, NSW, Australia

[64]Broad Genomics, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[65]Diabetes Unit and Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA

[66]Programs in Metabolism and Medical & Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[67]Institute of Clinical Molecular Biology, (IKMB) Christian-Albrechts-University of Kiel, Kiel, Germany

[68]Helsinki University and Helsinki University Hospital Clinic of Gastroenterology, Helsinki, Finland

[69]Bioinformatics Program MGH Cancer Center and Department of Pathology, Boston, MA, USA

[70]Cancer Genome Computational Analysis, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[71]Department of Psychiatry and Behavioral Sciences, Boston Children's Hospitaland Harvard Medical School, Boston, MA, USA

[72]Harvard Medical School Teaching Hospital, Boston, MA, USA

[73]Department of Endocrinology and Metabolism, Hadassah Medical Center and Faculty of Medicine, Hebrew University of Jerusalem, Israel

[74]Department of Psychiatry and Behavioral Sciences, SUNY Upstate Medical University, Syracuse, NY, USA

[75]Institute for Genomic Medicine, Columbia University Medical Center Hammer Health Sciences, New York, NY, USA

[76]Department of Genetics & Development Columbia University Medical Center, Hammer Health Sciences, New York, NY, USA

[77]Centro de Investigacion en Salud Poblacional, Instituto Nacional de Salud Publica, Mexico

[78]Lund University Sweden, Sweden

[79]Institute for Molecular Medicine Finland, (FIMM) HiLIFE University of Helsinki, Helsinki, Finland

[80]Division of Genetics and Genomics, Boston Children's Hospital, Boston, MA, USA

[81]Lund University Diabetes Centre, Malmö, Skåne County, Sweden

[82]Washington School of Medicine, St Louis, MI, USA

[83]Human Genetics Center University of Texas Health Science Center at Houston, Houston, TX, USA

[84]Department of Neurology Columbia University, New York City, NY, USA

[85]Institute of Genomic Medicine, Columbia University, New York City, NY, USA

[86]Institute of Biomedicine, University of Eastern Finland, Kuopio, Finland

[87]Department of Psychiatry, Helsinki University Central Hospital Lapinlahdentie, Helsinki, Finland

[88]Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

[89]Icahn School of Medicine at Mount Sinai, New York, NY, USA

[90]Bonei Olam, Center for Rare Jewish Genetic Diseases, Brooklyn, NY, USA

[91]Department of Neurology, Helsinki University, Central Hospital, Helsinki, Finland

[92]Cardiovascular Disease Initiative and Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[93]Charles Bronfman Institute for Personalized Medicine, New York, NY, USA

[94]Division of Genome Science, Department of Precision Medicine, National Institute of Health, Republic of Korea

[95]MRC Centre for Neuropsychiatric Genetics & Genomics, Cardiff University School of Medicine, Cardiff, Wales

[96]Imperial College, Healthcare NHS Trust, London, UK

[97]National Heart and Lung Institute Cardiovascular Sciences, Hammersmith Campus, Imperial College

London, London, UK

[98]Department of Health THL-National Institute for Health and Welfare, Helsinki, Finland

[99]Section of Cardiovascular Medicine, Department of Internal Medicine, Yale School of Medicine, Center for Outcomes Research and Evaluation Yale-New Haven Hospital, New Haven, CT, USA

[100]Division of Pediatric Gastroenterology, Emory University School of Medicine, Atlanta, GA, USA

[101]Department of Internal Medicine, Seoul National University Hospital, Seoul, Republic of Korea

[102]The University of Eastern Finland, Institute of Clinical Medicine, Kuopio, Finland

[103]Kuopio University Hospital, Kuopio, Finland

[104]Department of Genetics, Yale School of Medicine, New Haven, CT, USA

[105]Department of Clinical Chemistry Fimlab Laboratories and Finnish Cardiovascular Research Center-Tampere Faculty of Medicine and Health Technology, Tampere University, Finland

[106]The Mindich Child Health and Development, Institute Icahn School of Medicine at Mount Sinai, New York, NY, USA

[107]National Autonomous University of Mexico, Mexico City, Mexico

[108]Salvador Zubirán National Institute of Health Sciences and Nutrition, Mexico City, Mexico

[109]Li Ka Shing Institute of Health Sciences, The Chinese University of Hong Kong, Hong Kong, China

[110]Hong Kong Institute of Diabetes and Obesity, The Chinese University of Hong Kong, Hong Kong, China

[111]Centre for Population Genomics, Garvan Institute of Medical Research and UNSW Sydney, Sydney, Australia

[112]Centre for Population Genomics, Murdoch Children's Research Institute, Melbourne, Australia

[113]University of California San Francisco Parnassus Campus, San Francisco, CA, USA

[114]Cardiovascular Research REGICOR Group, Hospital del Mar Medical Research Institute, (IMIM) Barcelona, Catalonia, Spain

[115]Department of Genetics, Harvard Medical School, Boston, MA, USA

[116]Oxford Centre for Diabetes, Endocrinology and Metabolism, University of Oxford, Churchill Hospital Old Road Headington, Oxford, OX, LJ, UK

[117]Welcome Centre for Human Genetics, University of Oxford, Oxford, OX, BN, UK

[118]Oxford NIHR Biomedical Research Centre, Oxford University Hospitals, NHS Foundation Trust, John Radcliffe Hospital, Oxford, OX, DU, UK

[119]John P. Hussman Institute for Human Genomics, Leonard M. Miller School of Medicine, University of Miami, Miami, FL, USA

[120]The Dr. John T. Macdonald Foundation Department of Human Genetics, Leonard M. Miller School of Medicine, University of Miami, Miami, FL, USA

[121]F. Widjaja Foundation Inflammatory Bowel and Immunobiology Research Institute Cedars-Sinai Medical Center, Los Angeles, CA, USA

[122]Atherogenomics Laboratory University of Ottawa, Heart Institute, Ottawa, Canada

[123]Division of General Internal Medicine, Massachusetts General Hospital, Boston, MA, USA

[124]Department of Clinical Sciences University, Hospital Malmo Clinical Research Center, Lund University, Malmö, Sweden

[125]Estonian Genome Center, Institute of Genomics, University of Tartu, Tartu, Estonia

[126]University of Arizona Health Science, Tuscon, AZ, USA

[127]Department of Medicine, University of Maryland School of Medicine, Baltimore, MD, USA

[128]Howard Hughes Medical Institute and Department of Molecular Biology, Massachusetts General Hospital, Boston, MA, USA

[129]International Centre for Diarrhoeal Disease Research, Bangladesh

[130]Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

[131]Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

[132]Lund University, Dept. Clinical Sciences, Skåne University Hospital, Malmö, Sweden

[133]Department of Statistical Genetics, Osaka University Graduate School of Medicine, Suita, Japan

[134]Laboratory of Statistical Immunology, Immunology Frontier Research Center (WPI-IFReC), Osaka University, Suita, Japan

[135]Integrated Frontier Research for Medical Science Division, Institute for Open and Transdisciplinary Research Initiatives, Osaka University, Suita, Japan

[136]Instituto Nacional de Medicina Genómica, (INMEGEN) Mexico City, Mexico

[137]Medical Research Institute, Ninewells Hospital and Medical School University of Dundee, Dundee, UK

[138]Wake Forest School of Medicine, Winston-Salem, NC, USA

[139]Department of Molecular Medicine and Biopharmaceutical Sciences, Graduate School of Convergence Science and Technology, Seoul National University, Seoul, Republic of Korea

[140]Department of Psychiatry Keck School of Medicine at the University of Southern California, Los Angeles, CA, USA

[141]Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, MD, USA

[142]Children's Hospital of Philadelphia, Philadelphia, PA, USA

[143]Division of Genetics and Epidemiology, Institute of Cancer Research, London, SM, NG

[144]University of Washington, Seattle, WA, USA

[145]Fred Hutchinson Cancer Research Center, Seattle, WA, USA

[146]Medical Research Center, Oulu University Hospital, Oulu Finland and Research Unit of Clinical Neuroscience Neurology University of Oulu, Oulu, Finland

[147]Center for Public Health Genomics, University of Virginia, Charlottesville, VA, USA

[148]Department of Public Health Sciences, University of Virginia, Charlottesville, VA, USA

[149]Research Center Montreal Heart Institute, Montreal, Quebec, Canada

[150]Department of Medicine, Faculty of Medicine Université de Montréal, Québec, Canada

[151]Department of Public Health Faculty of Medicine, University of Helsinki, Helsinki, Finland

[152]Broad Institute of MIT and Harvard, Cambridge, MA, USA

[153]Department of Biomedical Informatics Vanderbilt, University Medical Center, Nashville, TN, USA

[154]Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA

[155]The Institute for Translational Genomics and Population Sciences, Department of Pediatrics, The Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center, Torrance, CA, USA

[156]Department of Biostatistics and Epidemiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

[157]Department of Medicine, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA, USA

[158]Center for Non-Communicable Diseases, Karachi, Pakistan

[159]National Institute for Health and Welfare, Helsinki, Finland

[160]Deutsches Herzzentrum, München, Germany

[161]Technische Universität München, Germany

[162]Institute of Genetic Epidemiology, Department of Genetics and Pharmacology, Medical University of Innsbruck, 6020 Innsbruck, Austria

[163]Duke Molecular Physiology Institute, Durham, NC

[164]Division of Cardiovascular Medicine, Nashville VA Medical Center, Vanderbilt University School of Medicine, Nashville, TN, USA

[165]Division of Endocrinology, National University Hospital, Singapore

[166]NUS Saw Swee Hock School of Public Health, Singapore

[167]Channing Division of Network Medicine, Brigham and Women's Hospital, Boston, MA, USA

[168]Harvard Medical School, Boston, MA, USA

[169]Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[170]Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[171]Institute for Genomics and Multiscale Biology, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[172]The Wallenberg Laboratory/Department of Molecular and Clinical Medicine, Institute of Medicine, Gothenburg University and the Department of Cardiology, Sahlgrenska University Hospital, Gothenburg, Sweden

[173]Department of Cardiology, Wallenberg Center for Molecular Medicine and Lund University Diabetes Center, Clinical Sciences, Lund University and Skåne University Hospital, Lund, Sweden

[174]Institute of Clinical Medicine Neurology, University of Eastern Finad, Kuopio, Finland

[175]Sorbonne Université, INSERM, Centre de Recherche Saint-Antoine, CRSA, AP-HP, Saint Antoine Hospital, Gastroenterology department, F-75012 Paris, France

[176]INRA, UMR1319 Micalis & AgroParisTech, Jouy en Josas, France

[177]Paris Center for Microbiome Medicine, (PaCeMM) FHU, Paris, France

[178]Department of Twin Research and Genetic Epidemiology King's College London, London, UK

[179]The McDonnell Genome Institute at Washington University, Seattle, WA, USA
[180]Departments of Genetics and Psychiatry, University of North Carolina, Chapel Hill, NC, USA
[181]Saw Swee Hock School of Public Health National University of Singapore, National University Health System, Singapore
[182]Department of Medicine, Yong Loo Lin School of Medicine National University of Singapore, Singapore
[183]Duke-NUS Graduate Medical School, Singapore
[184]Life Sciences Institute, National University of Singapore, Singapore
[185]Department of Statistics and Applied Probability, National University of Singapore, Singapore
[186]Center for Behavioral Genomics, Department of Psychiatry, University of California, San Diego, CA, USA
[187]Institute of Genomic Medicine, University of California San Diego, San Diego, CA, USA
[188]Endocrinology, Abdominal Center, Helsinki University Hospital, Helsinki, Finland
[189]Institute of Genetics, Folkhalsan Research Center, Helsinki, Finland
[190]Juliet Keidan Institute of Pediatric Gastroenterology Shaare Zedek Medical Center, The Hebrew University of Jerusalem, Jerusalem, Israel
[191]Instituto de Investigaciones Biomédicas, UNAM, Mexico City, Mexico
[192]Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico
[193]Department of Public Health Faculty of Medicine University of Helsinki, Helsinki, Finland
[194]Department of Psychiatry and Human Behavior, University of California Irvine, Irvine, CA, USA
[195]National Heart & Lung Institute & MRC London Institute of Medical Sciences, Imperial College, London, UK
[196]Royal Brompton & Harefield Hospitals, Guy's and St. Thomas' NHS Foundation Trust, London, UK
[197]Radcliffe Department of Medicine, University of Oxford, Oxford, UK
[198]Department of Gastroenterology and Hepatology, University of Groningen and University Medical Center Groningen, Groningen, Netherlands
[199]Folkhälsan Institute of Genetics, Folkhälsan Research Center, Helsinki, Finland
[200]National Heart & Lung Institute and MRC London Institute of Medical Sciences, Imperial College London, London, UK
[201]Cardiovascular Research Centre, Royal Brompton & Harefield Hospitals NHS Trust, London, UK
[202]Department of Physiology and Biophysics, University of Mississippi Medical Center, Jackson, MS, USA
[203]Program in Infectious Disease and Microbiome, Broad Institute of MIT and Harvard, Cambridge, MA, USA
[204]Center for Computational and Integrative Biology, Massachusetts General Hospital, Boston, MA, USA

Authors received funding as follows:
Emelia J Benjamin: Framingham Heart Study 75N92019D00031; and R01HL092577
Matthew J. Bown: British Heart Foundation awards CS/14/2/30841 and RG/18/10/33842
Josée Dupuis: National Heart Lung and Blood Institute's Framingham Heart Study Contract (HHSNI); National Institute for Diabetes and Digestive and Kidney Diseases (NIDDK) R DK
Martti Färkkilä: State funding for university level health research
Laura D. Gauthier: Intel, Illumina
Stephen J. Glatt: U.S. NIMH Grant R MH
Leif Groop: The Academy of Finland and University of Helsinki: Center of Excellence for Complex Disease Genetics (grant number 312063 and 336822), Sigrid Jusélius Foundation; IMI 2 (grant No 115974 and 15881)
Mikko Hiltunen: Academy of Finland (grant 338182) Sigrid Jusélius Foundation the Strategic Neuroscience Funding of the University of Eastern Finland
Chaim Jalas: Bonei Olam
Jaakko Kaprio: Academy of Finland (grants 312073 and 336823)
Jacob McCauley: National Institute of Diabetes and Digestive and Kidney Disease Grant R01DK104844
Yukinori Okada: JSPS KAKENHI (19H01021, 20K21834), AMED (JP21km0405211, JP21ek0109413, JP21gm4010006, JP21km0405217, JP21ek0410075), JST Moonshot R&D (JPMJMS2021)
Michael J. Owen: Medical Research Council UK: Centre Grant No. MR/L010305/1, Program Grant No. G0800509
Aarno Palotie: the Academy of Finland Center of Excellence for Complex Disease Genetics (grant numbers 312074 and 336824) and Sigrid Jusélius Foundation

John D. Rioux: National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK; DK062432), from the Canadian Institutes of Health (CIHR GPG 102170), from Genome Canada/Génome Québec (GPH-129341), and a Canada Research Chair (#230625)

Samuli Ripatti: the Academy of Finland Center of Excellence for Complex Disease Genetics (grant number) Sigrid Jusélius Foundation

Jerome I. Rotter: Trans-Omics in Precision Medicine (TOPMed) program was supported by the National Heart, Lung and Blood Institute (NHLBI). WGS for "NHLBI TOPMed: Multi-Ethnic Study of Atherosclerosis (MESA)" (phs001416.v1.p1) was performed at the Broad Institute of MIT and Harvard (3U54HG003067-13S1). Core support including centralized genomic read mapping and genotype calling, along with variant quality metrics and filtering were provided by the TOPMed Informatics Research Center (3R01HL-117626-02S1; contract HHSN268201800002I). Core support including phenotype harmonization, data management, sample-identity QC, and general program coordination were provided by the TOPMed Data Coordinating Center (R01HL-120393; U01HL-120393; contract HHSN268201800001I). We gratefully acknowledge the studies and participants who provided biological samples and data for MESA and TOPMed. JSK was supported by the Pulmonary Fibrosis Foundation Scholars Award and grant K23-HL-150301 from the NHLBI. MRA was supported by grant K23-HL-150280, AJP was supported by grant K23-HL-140199, and AM was supported by R01-HL131565 from the NHLBI. EJB was supported by grant K23-AR-075112 from the National Institute of Arthritis and Musculoskeletal and Skin Diseases.The MESA project is conducted and supported by the National Heart, Lung, and Blood Institute (NHLBI) in collaboration with MESA investigators. Support for MESA is provided by contracts 75N92020D00001, HHSN268201500003I, N01-HC-95159, 75N92020D00005, N01-HC-95160, 75N92020D00002, N01-HC-95161, 75N92020D00003, N01-HC-95162, 75N92020D00006, N01-HC-95163, 75N92020D00004, N01-HC-95164, 75N92020D00007, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, N01-HC-95169, UL1-TR-000040, UL1-TR-001079, and UL1-TR-001420. Also supported in part by the National Center for Advancing Translational Sciences, CTSI grant UL1TR001881, and the National Institute of Diabetes and Digestive and Kidney Disease Diabetes Research Center (DRC) grant DK063491 to the Southern California Diabetes Endocrinology Research Center

Edwin K. Silverman: NIH Grants U01 HL089856 and U01 HL089897

J. Gustav Smith: The Swedish Heart-Lung Foundation (2019-0526), the Swedish Research Council (2017-02554), the European Research Council (ERC-STG-2015-679242), Skåne University Hospital, governmental funding of clinical research within the Swedish National Health Service, a generous donation from the Knut and Alice Wallenberg foundation to the Wallenberg Center for Molecular Medicine in Lund, and funding from the Swedish Research Council (Linnaeus grant Dnr 349-2006-237, Strategic Research Area Exodiab Dnr 2009-1039) and Swedish Foundation for Strategic Research (Dnr IRC15-0067) to the Lund University Diabetes Center

Kent D. Taylor: Trans-Omics in Precision Medicine (TOPMed) program was supported by the National Heart, Lung and Blood Institute (NHLBI). WGS for "NHLBI TOPMed: Multi-Ethnic Study of Atherosclerosis (MESA)" (phs001416.v1.p1) was performed at the Broad Institute of MIT and Harvard (3U54HG003067-13S1). Core support including centralized genomic read mapping and genotype calling, along with variant quality metrics and filtering were provided by the TOPMed Informatics Research Center (3R01HL-117626-02S1; contract HHSN268201800002I). Core support including phenotype harmonization, data management, sample-identity QC, and general program coordination were provided by the TOPMed Data Coordinating Center (R01HL-120393; U01HL-120393; contract HHSN268201800001I). We gratefully acknowledge the studies and participants who provided biological samples and data for MESA and TOPMed. JSK was supported by the Pulmonary Fibrosis Foundation Scholars Award and grant K23-HL-150301 from the NHLBI. MRA was supported by grant K23-HL-150280, AJP was supported by grant K23-HL-140199, and AM was supported by R01-HL131565 from the NHLBI. EJB was supported by grant K23-AR-075112 from the National Institute of Arthritis and Musculoskeletal and Skin Diseases.The MESA project is conducted and supported by the National Heart, Lung, and Blood Institute (NHLBI) in collaboration with MESA investigators. Support for MESA is provided by contracts 75N92020D00001, HHSN268201500003I, N01-HC-95159, 75N92020D00005, N01-HC-95160, 75N92020D00002, N01-HC-95161, 75N92020D00003, N01-HC-95162, 75N92020D00006, N01-HC-95163, 75N92020D00004, N01-HC-95164, 75N92020D00007, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, N01-HC-95169, UL1-TR-000040, UL1-TR-001079, and UL1-TR-001420. Also supported in part by the National Center for Advancing Translational Sciences, CTSI grant UL1TR001881, and the National Institute of Diabetes and Digestive and Kidney Disease Diabetes Research Center (DRC) grant

DK063491 to the Southern California Diabetes Endocrinology Research Center
Tiinamaija Tuomi: The Academy of Finland and University of Helsinki: Center of Excellence for Complex Disease Genetics (grant number 312072 and 336826), Folkhalsan Research Foundation, Helsinki University Hospital, Ollqvist Foundation, Liv och Halsa foundation; NovoNordisk Foundation
Teresa Tusie-Luna: CONACyT Project 312688
James S. Ware: Sir Jules Thorn Charitable Trust [21JTA], Wellcome Trust [107469/Z/15/Z], Medical Research Council (UK), NIHR Imperial College Biomedical Research Centre
Rinse K. Weersma: The Lifelines Biobank initiative has been made possible by subsidy from the Dutch Ministry of Health Welfare and Sport the Dutch Ministry of Economic Affairs the University Medical Centre Groningen (UMCG the Netherlands) the University of Groningen and the Northern Provinces of the Netherlands

No conflicts of interest to declare