Lysine-Cysteine-Lysine (KCK) tag changes ParB action *in vitro* but not *in vivo.*

Miranda Molina[1,2,4], Lindsey E. Way[3,4], Zhongqing Ren[3], Qin Liao[3], Xindan Wang[3]*, HyeongJun Kim[1,2]*

[1]Biochemistry and Molecular Biology Program, University of Texas Rio Grande Valley, Edinburg, Texas, United States of America

[2]Department of Physics and Astronomy, University of Texas Rio Grande Valley, Edinburg, Texas, United States of America

[3]Department of Biology, Indiana University, 1001 E 3rd St, Bloomington, Indiana, United States of America

[4]These authors contributed equally.

*Corresponding authors

E-mail: xindan@indiana.edu (XW); hyeongjun.kim@utrgv.edu (HK)

Abstract

Due to the enhanced labeling capability of maleimide-based fluorescent probes in *in vitro* experiments, lysine-cysteine-lysine (KCK) tags are frequently added to proteins for visualization. Here we show that, although no noticeable changes were detected from *in vivo* fluorescence imaging and chromatin immunoprecipitation (ChIP) assays, the KCK-tag substantially altered DNA compaction rates by *Bacillus subtilis* ParB protein in *in vitro* single-molecule DNA flow-stretching experiments. Furthermore, our measurements and statistical analyses demonstrate that the KCK-tags also altered the ParB protein's response to nucleotide (cytidine triphosphate CTP or its nonhydrolyzable analog CTPγS) binding and the presence of the specific DNA binding sequence (*parS*). Remarkably, the appended KCK-tags are capable of even reversing the trends of DNA compaction rates upon different experimental conditions. DNA flow-stretching experiments for both fluorescently-labeled ParB proteins and ParB proteins with an N-terminal glutamic acid-cysteine-glutamic acid (ECE) tag support the notion that electrostatic interactions between charges on the tags and the DNA backbone are an underlying cause of the protein's property changes. While it is typically assumed that the short KCK-tag minimally perturbs protein function, our results demonstrate that this assumption must be carefully tested when using tags for protein labeling.

Introduction

Single-molecule DNA flow-stretching is a powerful method to study the actions of DNA-binding proteins. Maleimide-conjugated fluorescent dyes have been widely used to label proteins via covalent conjugation to surface-exposed cysteines[1]. Despite this specificity and convenience, labeling all desired cysteines with maleimide dyes is not always achieved. The reaction efficiency between the thiol group on cysteine and the maleimide moiety of a fluorescent dye can be increased by flanking the cysteine with two positively charged lysine residues. It was revealed that the neighboring lysine residues decrease pKa of the cysteine residue, thereby increasing thiol-maleimide reactivity[2–5]. Thus, appending the lysine-cysteine-lysine (KCK) tag to a protein has been a popular and extensively used method due to its superior fluorescence labeling efficiency[6–12]. In this

study, we report that DNA compaction by the DNA-binding protein ParB is artificially enhanced by KCK-tags in single-molecule assays *in vitro*, producing misleading results.

Results and discussion

The ParABS DNA partitioning system is a broadly conserved segregation machinery for bacterial chromosomes and plasmids. ParB binds to *parS* sequences and spreads to neighboring regions[13,14] to form a nucleoprotein complex, which is translocated by ParA[13,14]. *In vivo*, ParB spreading is evident by two approaches: fluorescence microscopy in which fluorescently-tagged ParB proteins form foci in live cells and chromatin immunoprecipitation (ChIP) assays in which ParB protein associates with 10-20 kb DNA regions encompassing *parS*[13,14]. Importantly, it was recently discovered that ParB protein is a novel enzyme that utilizes cytidine triphosphate (CTP) to modulate ParB spreading[15–17].

KCK-tags increase BsParB's DNA compaction rates *in vitro*. To elucidate the roles of CTP in the action of ParB protein, we purified apyrase-treated wild-type *Bacillus subtilis* ParB (BsParB(WT)) protein (Supplementary Fig. 1) and employed single-molecule DNA flow-stretching assays with a lambda DNA substrate (Fig. 1a). The speed of DNA compaction by BsParB(WT) was measured by tracking the positions of a fluorescent quantum dot labeled at one DNA end[12] (Fig. 1b). In the presence of 50 nM BsParB(WT), we observed robust DNA compaction in the absence of CTP as previously shown[12] (Fig. 1b). Interestingly, both CTP and CTPγS (a non-hydrolyzable CTP analog) inhibited DNA compaction by 39-fold and 149-fold, respectively (Fig. 1c), implying counter-productive roles of CTP binding in DNA compaction. Next, we purified BsParB(WT) with the KCK-tag at its N-terminus (hereafter "KCK-BsParB(WT)") (Supplementary Fig. 1). We observed that DNA compaction by the KCK-BsParB(WT) was robust without CTP albeit slower than BsParB(WT) (Fig. 1c). However, inclusion of CTP or CTPγS led to strikingly increased DNA compaction rate (10.5-fold and 19.4-fold for CTP and CTPγS, respectively) in KCK-BsParB(WT) compared with BsParB(WT) (Fig. 1c). Since batch-to-batch variations in purified proteins only lead to up to two-fold differences for DNA compaction rates in our experience, these dramatic changes prompted us to investigate further.

Supplementary Fig. 1. Purified BsParB proteins. InstantBlue Coomassie-stained SDS-PAGE gel of recombinant *Bacillus subtilis* ParB proteins used in this study. The protein ladder (Gold Biotechnology, #P007-500) is on the left.

Next, we examined the effect of a *parS* sequence on DNA compaction rates by utilizing an engineered lambda

DNA harboring one *parS* in the middle (hereafter, "*parS* DNA")[12]. Although *parS* DNA compaction by BsParB(WT)

without any nucleotides was about three-quarters that of lambda DNA (Supplementary Fig. 2a, c), in the

presence of CTP or CTPγS, the compaction rates of *parS* DNA decreased by 16-fold and 50-fold, respectively

(Supplementary Fig. 2a). Furthermore, KCK-BsParB(WT) exhibited substantial boosts in the *parS* DNA

compaction rates in the presence of CTP (4.9-fold) or CTPγS (7.2-fold) compared with BsParB(WT)

(Supplementary Fig. 2a). Thus, the KCK-tag enhanced BsParB(WT)'s DNA compaction rate (compared to the

untagged BsParB(WT)) when nucleotides are present on both lambda DNA and *parS* DNA.

Given that ParB protein's CTP binding pocket resides at the N-terminal domain (NTD) and the NTD is

implicated to be the DNA-entry gate[16–18], we questioned if the unexpected compaction rate increases also occur

when KCK is tagged at the C-terminal of BsParB(WT) protein (hereafter, "BsParB(WT)-KCK") (Supplementary

Fig. 1). Indeed, like KCK-BsParB(WT), BsParB(WT)-KCK also showed much faster compaction with CTP

compared with BsParB(WT) (Fig. 1c for the lambda DNA and Supplementary Fig. 2a Fig for the *parS* DNA).

Thus, KCK enhanced BsParB(WT)'s DNA compaction rate when appended to either terminus.

Fig. 1 *in vitro* quantitative and qualitative BsParB compaction rate changes by the KCK-tags. a Schematic of single-molecule DNA flow-stretching assays. **b** An example of DNA compaction by 50 nM BsParB(WT) protein (top) and the definition of compaction rate (bottom). **c-d** Lambda DNA compaction rates by 50 nM (**c**) wild-type and (**d**) R80A mutant proteins. Numbers indicate compaction rate fold increases. Error bars: SEM. **e** Top: The Mann-Whitney test (the Wilcoxon rank sum test) *p*-value color scheme. Bottom: Mann-Whitney test comparisons for compaction rates by wild-type BsParB and its KCK-

versions. **f** Mann-Whitney test comparisons for BsParB(R80A) and its KCK-versions. (**e-f**) Cyan, green, and yellow boxes highlight qualitative protein property changes due to the KCK-tags for visual aids. (**c-f**) See Tab 1 in the Supplementary File for detailed sample number (*N*) information.

Supplementary Fig. 2. KCK-tags lead to quantitative and qualitative compaction rate changes. (**a**) *parS* DNA compaction rates by BsParB(WT), KCK-BsParB(WT), and BsParB(WT)-KCK proteins in the absence and presence of 1 mM CTP or 1 mM CTPγS. Error bars: s.e.m., The numbers indicate compaction rate fold changes. (**b**) *parS* DNA compaction rates by BsParB(R80A), KCK-BsParB(R80A), and BsParB(R80A)-KCK proteins in the absence and presence of 1 mM CTP or 1 mM CTPγS. Error bars: s.e.m., The numbers indicate compaction rate fold changes. (**c**) For direct comparisons, the compaction rates shown in (**a**),

(**b**) and Fig. 1c, d are consolidated. Error bars: s.e.m. (**a-c**) See Tab 1 in the Supplementary

File for detailed sample number (*N*) information.

KCK-tags alter BsParB's DNA compaction properties in response to *parS* and nucleotides. To assess

how the BsParB(WT) protein and its KCK-tagged variants respond to different nucleotides and a *parS* site, Mann-

Whitney tests were performed for compaction rates with all possible permutations (Fig. 1e. Also see

Supplementary Fig. 3a). The Mann-Whitney tests revealed that, without any nucleotide or with 1 mM CTP,

BsParB(WT) was responsive to the existence of *parS* (*p*<0.001) while BsParB(WT)-KCK did not make

statistically significant compaction rate changes with *parS* (*p*≥0.05) (See cyan boxes in Fig. 1e). We note that

the KCK-tags not only changed compaction rates (Fig. 1c, d and Supplementary Fig. 2a-c) but also reversed the

trend of compaction. Specifically, without nucleotides, when a *parS* site was added to DNA, BsParB(WT)'s

compaction rate was slowed down, but KCK-BsParB(WT)'s compaction rate was increased (Supplementary Fig.

2c). These results show that the KCK-tag alters the DNA-compaction ability both quantitatively and qualitatively.

Supplementary Fig. 3. KCK-tags lead to qualitative compaction rate changes. Since not all results pass the Shapiro-Wilk normality test, we employed the Mann-Whitney tests to compare DNA compaction rates in Fig 1e, f. However, Welch's t-test results are still informative as long as there are not extreme outliers and there are enough (>25) data points [19]. Indeed, the Welch's t-test results provided here are very similar to the ones from the Mann-Whitney tests. (**a**) Top: The Welch's t-test *p*-value color scheme. Bottom: The Welch's t-test comparisons for compaction rates by wild-type BsParB and its KCK-versions. (**b**) The Welch's t-test comparisons for BsParB(R80A) and its KCK-versions. (**a-b**) Cyan, green and yellow boxes highlight qualitative protein property changes due to the KCK-tags for visual aids. See Tab 1 in the Supplementary File for detailed sample number (*N*) information.

The KCK-tag alters the action of BsParB R80A mutant. We next investigated whether the compaction rate

differences induced by the KCK-tag are limited only to the wild-type BsParB. The R80A mutant of BsParB has

been shown to abolish proper *in vivo* sporulation, localization, and spreading along with *in vitro* lambda DNA

compaction in the absence of nucleotides[12,20,21]. Surprisingly, without nucleotides, although its DNA compaction

rate was 18.2-fold lower than BsParB(WT) (Supplementary Fig. 2c), BsParB(R80A) (Supplementary Fig. 1) was

still capable of compacting the lambda DNA (Fig. 1d), contradicting a previous report[12]. Although both studies

are using the same assay, in our study, we supplemented magnesium ions to our buffer as a cofactor of CTP

and used apyrase during our protein purification to remove residual CTPs. Since BsParB(R80A) is deficient in

CTP hydrolysis[16], it is possible that CTPs were co-purified with BsParB(R80A) in the previous study[12]. Consistent

with our speculation, in the absence of Mg^{2+} and the presence of CTP, BsParB(R80A)'s compaction rate was

reduced dramatically (Supplementary Fig. 4), providing an explanation for the undetectable compaction by

BsParB(R80A) in the previous study. Next, we wondered whether a KCK tag alters BsParB(R80A)'s action on

DNA. Indeed, with lambda DNA, the compaction rates of both KCK-BsParB(R80A) and BsParB(R80A)-KCK

were substantially increased for all tested nucleotides (Fig. 1d). When *parS* DNA was used as a substrate,

compaction rate increases by KCK tags (*p*<0.0001) were also noted (Supplementary Fig. 2b). The visualized

Mann-Whitney comparison charts for DNA compaction rates highlight that BsParB(R80A), KCK-BsParB(R80A),

and BsParB(R80A)-KCK respond differently to different nucleotides and the presence of *parS* (See green and

yellow boxes in Fig. 1f. Also see Supplementary Fig. 3b.)

Supplementary Fig. 4. BsParB(R80A) DNA compaction rates in different conditions.
DNA compaction rates by 50 nM BsParB(R80A) on lambda DNA and *parS* DNA in the
presence and absence of magnesium ions ($[MgCl_2]$ = 2.5 mM), CTP (1 mM), and CTPγS (1
mM). DNA Compaction rates in the presence of CTP and in the absence of magnesium ions
are highlighted in red. The absence of magnesium and presence of CTP could explain why
the previous study[12] did not detect DNA compaction by BsParB(R80A). Error bars: s.e.m., ****
denotes $p<0.0001$. See Tabs 1 and 2 in the Supplementary File for detailed sample number
(*N*) information.

The effects of KCK-tags on protein action are limited to *in vitro* assays but not *in vivo.* The different effects

of KCK tags in DNA compaction *in vitro* prompted us to systematically test the effect of KCK tag on BsParB's or

BsParB(R80A)'s localization and spreading *in vivo*. We first generated eight GFP fusions to the ParB variants

with KCK tags at the C- or N-terminus of the protein and performed fluorescence microscopy (Fig. 2a). Consistent

with previous findings that R80A abolishes ParB spreading[12], BsParB(WT) formed foci in the cells, while

BsParB(R80A) had diffused localization on the DNA. Interestingly, KCK tags at the C- or N-terminus did not alter

the localization of ParB(WT) or ParB(R80A) (Fig. 2a). In a complementary approach, we analyzed the *in vivo*

spreading of ParB variants on the genome by chromatin immunoprecipitation (ChIP-seq) assays using anti-ParB

antibodies (Fig. 2b). We observed that BsParB(WT) spread to a ~20 kb region surrounding the *parS* site, but

BsParB(R80A) did not spread. These results are consistent with previously published data[12]. Importantly, having

a KCK tag at the C- or N-terminus did not affect the spreading of BsParB(WT) or BsParB(R80A). We also show

that the KCK-tagged proteins have similar expression levels compared to the matched untagged controls

(Supplementary Fig. 5a, b). These experiments demonstrate that the KCK tag does not affect BsParB's functions

in vivo. Thus, the effects of KCK tags on BsParB(WT) and BsParB(R80A) are specific to *in vitro* experiments.

Fig. 2 KCK tags do not affect *in vivo* BsParB localization or spreading. a Localization of fluorescently tagged ParB(WT) and ParB(R80A) (green). The nucleoid is labeled with HBsu-mCherry (red), and phase-contrast images are shown in gray. Scale bar represents 2 μm. **b** ChIP-seq of wild-type and mutant ParB association with a region of the *B. subtilis* chromosome from 354° to 360° (3960–4033 kb of strain the PY79 genome). Red dotted lines indicate the positions of the four *parS* sites. The number of reads were normalized by the total number of reads per sample. Whereas wild-type ParB spreads several kilobases from *parS* sites, the R80A mutant is restricted to the immediate vicinity of each *parS* site. KCK tags at the N-terminus or C-terminus of ParB or R80A do not change the property of the variants.

Supplementary Fig. 5. KCK tags do not significantly alter the level of ParB or R80A. (**a**) Western blot of GFP-tagged ParB variants. Although GFP-ParB levels are higher than ParB-GFP levels, the R80A mutation or KCK tag does not change the protein levels. SigA levels are shown to control for loading. (**b**) Western blot of ParB variants. The R80A mutation or KCK tag does not dramatically change the protein levels. Asterisk indicates the ParB band. SigA levels are shown to control for loading.

Charges on the KCK-tag contribute to the *in vitro* protein property changes. This finding prompted us to

understand the mechanism by which the KCK-tag boosts the DNA compaction rate of ParB protein *in vitro.* One

possibility for the compaction rate increase is that more BsParB proteins were recruited onto DNA due to

interactions between the positively-charged KCK-tag and the negatively-charged DNA backbone. Alternatively,

the KCK-tag could impact the subsequent action of the BsParB proteins while the level of the initial protein

recruitment is intact. To obtain insight into these two possibilities, we directly visualized the recruitment of

untagged and KCK-tagged BsParB(R80A) proteins onto lambda DNA. Proteins were nonspecifically labeled with

the NHS-ester version of Cyanine3 fluorescent dye, and the moment of the very first labeled protein's arrival into

the camera's field-of-view was evident by increase in background intensity (Fig. 3a). In this approach, background-subtracted integrated fluorescence intensity on DNA is directly proportional to the amount of BsParB protein recruited onto the DNA. The microscopy showed that the background-subtracted integrated fluorescence intensities with KCK-BsParB(R80A) and BsParB(R80A)-KCK were higher than those with BsParB(R80A) (p<0.0001) (Fig. 3b). Thus, our data show that the KCK-tags enhanced protein loading and increased compaction rates with a caveat that our experimental approaches do not address if the KCK tags impact on subsequent protein action after being recruited onto DNA.

To obtain another line of insight, we prepared recombinant wild-type and R80A mutant BsParB proteins where a negatively-charged glutamic acid-cysteine-glutamic acid (ECE)-tag is N-terminally appended. If electrostatic interactions between the appended tags and DNA backbone contribute to *in vitro* artifacts, slower compaction rates are expected with ECE-tagged BsParB proteins (hereafter "ECE-BsParB") due to repulsive forces between negative charges. As expected, DNA compactions by ECE-BsParB(R80A) were noticeably inefficient. The compaction rates by ECE-BsParB(R80A) are significantly lower (p<0.0001) than those by BsParB(R80A) regardless of the presence of the *parS* DNA sequence and CTP (Fig. 3c). Consistent with this observation, the ECE-BsParB(WT) protein also exhibits inefficient DNA compaction compared with its BsParB(WT) counterpart in the absence of any nucleotides (Supplementary Fig. 6).

Next, we investigated any *in vivo* property changes caused by N-terminally appended ECE-tag. Fluorescence microscopy experiments show that the ECE-tag does neither abolish the *in vivo* fluorescence foci formation with the wild-type BsParB protein nor lead to the formation of clear foci with the R80A mutant BsParB (Fig. 3d). Additionally, ChIP-seq assays using anti-ParB antibodies indicate that wild-type BsParB proteins spread to a ~20 kb regions around the *parS* site and the R80A mutant does not spread regardless of the presence of the ECE-tag (Fig. 3e). All *in vivo* results consistently demonstrate that the KCK and ECE tags appended to BsParB proteins do not have noticeable impacts. The effects of the tags are only limited to *in vitro* assays, and electrostatic interactions between charged residues on the tag and the DNA backbone are at least partly responsible for the *in vitro* effects.

Fig. 3 Charged amino acid residues in the appended tag are responsible for *in vitro* compaction rate changes. a An example of time-trajectories for mean background intensity and integrated fluorescence intensity on a DNA. The time point when the mean background intensity starts to increase is defined as t=0. DNA flow-stretching experiments were performed with fluorescently-labeled proteins. **b** Integrated fluorescence intensities on lambda DNAs by cyanine3-labeled BsParB(R80A) (*N*=51), KCK-BsParB(R80A) (*N*=40), and BsParB(R80A)-KCK (*N*=26) measured at different time points. Error bars: s.e.m., **** denotes

p<0.0001. **c** Lambda and *parS* DNA compaction rates by BsParB(R80A) and ECE-BsParB(R80A) both in the presence and absence of CTP. Error bars: s.e.m., **** denotes *p*<0.0001. See Tab 1 in the Supplementary File for detailed sample number (*N*) information. **d** Localization of fluorescently tagged BsParB(WT), BsParB(R80A), and their ECE-tagged versions (green). Red: the nucleoid labeled with HBsu-mCherry. Gray: phase-contrast images. Scale bar represents 2 μm. **e** ChIP-seq of ECE-tagged wild-type (left) and mutant ParB (right) association with a region of the *B. subtilis* chromosome from 354° to 360° (3960–4033 kb of strain the PY79 genome). Red dotted lines indicate the positions of the four *parS* sites. The number of reads were normalized by the total number of reads per sample. Whereas ECE-ParB(WT) spreads several kilobases from *parS* sites, the ECE-R80A mutant is restricted to the immediate vicinity of each *parS* site.

Supplementary Fig. 6. Lambda and *parS* DNA compaction rates by BsParB(WT) and ECE-BsParB(WT) both in the presence and absence of CTP. Error bars: s.e.m., **** denotes *p*<0.0001. See Tab 1 in the Supplementary File for detailed sample number (*N*) information.

179   In summary, we report that, although KCK tagging did not change the *in vivo* behavior of BsParB(WT) or

180   BsParB(R80A), it dramatically altered DNA compaction rates at the single-molecule level *in vitro*. Importantly,

181   the KCK-tag affected not only quantitative compaction rates but also qualitative behaviors of the protein against

182   different nucleotide statuses and the presence of a *parS* sequence. DNA flow-stretching assays with

fluorescently-labeled proteins and ECE-tagged BsParB proteins suggest that electrostatic interactions are, at least partly, a cause of *in vitro* property changes.

Deep understanding of any biological system requires both *in vitro* and *in vivo* approaches. Our study reveals that addition of short amino acid tags may produce misleading *in vitro* results despite normal functionality *in vivo.* Additionally, our results raise a possibility that fluorescent dyes conjugated to a DNA-binding protein result in altered *in vitro* protein activities due to electrostatic interactions between charges on the fluorescent probe and those on the DNA backbone. Whenever adding a small amino acid tag is desired for *in vitro* experiments, careful controls must be performed to ensure that this does not perturb the activity of the protein.

Methods

Plasmid constructions for *in vitro* single-molecule assays. Plasmids harboring coding sequences of His6-SUMO-BsParB(WT) (pTG011)[12], His6-SUMO-KCK-BsParB(WT) (pTG042)[12], His6-SUMO-BsParB(R80A) (pTG037)[12], and His6-SUMO-KCK-BsParB(R80A) (pTG044)[12] were generous gifts from Thomas Graham. Site-directed mutagenesis were performed to generate plasmids harboring coding sequences of His6-SUMO-BsParB(WT)-KCK (m0067) and His6-SUMO-BsParB(R80A)-KCK (m0069) using oHK050F and oHK050R as primers. The plasmid harboring coding sequences of His6-SUMO-ECE-BsParB(WT) (m0064) were generated using oHK048F and oHK048R as primers and m0043 as a substrate. Contrary to other plasmids, the plasmid harboring coding sequences of His6-SUMO-ECE-BsParB(R80A) (m0070) was generated by following the vendor-supplied NEBuilder HiFi DNA Assembly Master Mix (NEB E2621S, Ipswich, MA) protocol. First, the His6-SUMO-BsParB(WT) plasmid (pTG011 = m0041) was linearized and the majority of SUMO-BsParB(WT) coding sequences were removed by PCR using oHK038F and oHK038R as primers. Then, gfHK009 and gfHK010 were used as gene fragments with both containing 23 bp overlaps. After NEBuilder HiFi DNA assembly, NEB 5-alpha competent *E. coli* cells (NEB C2987H, Ipswich, MA) were transformed with the reaction mixture. The sequences were confirmed using T7, oHK023, oHK024, oHK025, and oHK026 oligos. See Tabs 7-8 in the Supplementary File for their sequences.

Protein expression and purification. Rosetta2(DE3)pLysS competent cells (EMD Millipore, Burlington, MA) transformed with a plasmid were cultured overnight at 37˚C in the presence of 100 μg/mL ampicillin and 20 μg/mL chloramphenicol. 1 L of LB medium with 80 μg/mL ampicillin was inoculated with the overnight culture and grown at 37˚C until the OD_{600} reached 0.4-0.6. Protein expression was induced with 500 μM isopropyl-β-D-thiogalactoside (IPTG), and the culture was shaken at 30˚C for an additional 4 hours. The cells were harvested by centrifugation at 4˚C. The cell pellets were resuspended in PBS buffer and spun at 5,000 g. They were resuspended in ParB lysis buffer (20 mM Tris, pH 8.0, 1 M NaCl, 50 mM imidazole, 5 mM 2-mercaptoethanol), supplemented with 0.1 mM phenylmethylsulfonyl fluoride (PMSF) and a protease inhibitor cocktail (Roche, Basel, Switzerland) (Total volume: 45 mL), and flash-frozen. BsParB proteins were purified based on a two-step tandem purification method as previously described[12] but with some modifications. Briefly, after thawing the harvested cells, additional 0.9 mM PMSF (total 1.0 mM PMSF), 50 mg/mL lysozyme, 3 μL of universal nuclease (Thermo Fisher Scientific 88701, Waltham, MA), and 5 mM 2-mercaptoethanol were added, and it was left in ice for 30 minutes. Cells were lysed by sonification and centrifuged twice in an FA-6x50 rotor: first at 11,000 g for 30 minutes, then at 20,133 g for 30 minutes. The clarified supernatant was incubated with Ni-NTA agarose beads (Qiagen, Hilden, Germany) for 1 hour in the presence of 1 unit of apyrase (NEB, Ipswich, MA) and 5 mM $MgCl_2$, to help minimize cellular NTPs that may otherwise be co-purified, and 1 tablet of cOmplete Mini EDTA-free protease inhibitor cocktail (Roche, Basel, Switzerland). The Ni-NTA agarose resin was washed with lysis buffer (supplemented with 5 mM $MgCl_2$) followed by ParB salt-reduction buffer (20 mM Tris, pH 8.0, 350 mM NaCl, 50 mM imidazole, 5 mM $MgCl_2$, 5 mM 2-mercaptoethanol). The proteins were manually eluted ten times with 1.5 mL of ParB elution buffer (20 mM Tris, pH 8.0, 350 mM NaCl, 250 mM imidazole, 5 mM $MgCl_2$, 5 mM 2-mercaptoethanol).

The peak fractions of ParB protein were pooled and treated with His6-Ulp1 protease to remove the N-terminal His6-SUMO tag[12]. The pooled proteins and His6-Ulp1 protease were dialyzed together overnight at 4˚C against ParB dialysis/storage 1 buffer (20 mM Tris, pH 8.0, 350 mM NaCl, 10 mM imidazole, 5 mM 2-mercaptoethanol, 1 mM $MgCl_2$, 10% glycerol). After centrifuging the dialyzed proteins at maximum speed for 10 minutes, the supernatant was allowed to interact with the Ni-NTA resin for at least 1 hour at 4˚C. Then, the flowthrough was collected. 0.5 mL of the ParB dialysis/storage 1 buffer to the Ni-NTA resin column was added multiple times, and the eluents were collected. Running an SDS-polyarcylamide (SDS-PAGE) gel indicated that the flowthrough and

the eluent fractions contained ParB protein, while the cleaved His6-SUMO and His6-Ulp1 remained in the resin.

The flowthrough and the peak fractions were pooled and dialyzed against ParB dialysis/storage 2 buffer (20 mM

Tris, pH 8.0, 350 mM NaCl, 10% glycerol), where 5 mM 2-mercaptoethanol was included in case of KCK-tagged

protein purifications. The protein concentration was measured by a NanoDrop One spectrophotometer (Thermo

Scientific, Waltham, MA) using 32.58 (kDa) and 7,450 (M^{-1} cm^{-1}) as its molecular weight and extinction coefficient,

respectively. The purified proteins (Supplementary Fig. 1) were run on a precast polyacrylamide gel (Bio-Rad,

Hercules, CA) with Tris/Glycine/SDS running buffer (Bio-Rad, Hercules, CA). InstantBlue Coomassie protein

stain (Abcam, Cambridge, United Kingdom) was used to stain for the polyacrylamide gel. The gel image was

obtained using UVP UVsolo touch gel documentation system (Analytik Jena, Jena, Germany) and provided in

the Supplementary Fig. 1 without any image processing.

DNA and Quantum-dot preparations. One end of bacteriophage lambda DNA (or *parS* DNA[12]) was labeled

with a biotin to tether the DNA onto the single-molecule microfluidic flowcell, and the other end was labeled with

a digoxigenin to attach a quantum dot (Fig. 1a) as previously described[22,23]. Briefly, Lambda-BL1Biotin and

Lambda-Dig2 oligos (Tab 8 in the Supplementary File) were treated with T4 polynucleotide kinase (PNK) (NEB,

Ipswich, MA) for phosphorylation at 37˚C for 1 hour. A 15-fold molar excess of the phosphorylated Lambda-

BL1Biotin oligo was introduced for annealing to a 12-base 5' single-stranded overhang on one end of lambda

DNA (or *parS* DNA[12]). The mixture of DNA and oligo was incubated at 65˚C for 10 minutes and slowly cooled

down, and then ligated by T4 ligase for 2 hours at room temperature. The other end of the lambda DNA (or *parS*

DNA) was tagged with a digoxigenin by supplementing a 60-fold molar excess of the phosphorylated Lambda-

Dig2 oligo at 45˚C. After 30-minute incubation, the mixture was slowly back to room temperature followed by a

2-hour ligation step at room temperature. Since the sequences of Lambda-BL1Biotin and Lambda-Dig2 oligos

are complementary to each other, it is important to remove unreacted excess oligos. After running a 0.4%

agarose gel overnight at 4˚C, the desired DNA band was excised and put into a dialysis tube. Applying an electric

field allowed DNAs to leave the excised agarose gel, but DNAs were confined to the dialysis tube volume. DNAs

were collected, and ethanol precipitation was performed to recover doubly-tagged lambda DNAs (or *parS* DNAs)

in EB buffer (10 mM Tris-Cl, pH 8.5).

As we previously did[22–24], anti-digoxigenin antibody-conjugated quantum dot 605 (Invitrogen, Waltham, MA)

was prepared following Invitrogen's Qdot 605 antibody conjugation kit (Q22001MP) manual. However, since this kit was discontinued, all the kit components were separately purchased including Qdot 605 ITK amino (PEG) quantum dots (Invitrogen Q21501MP). For the antibody, anti-digoxigenin fab fragments (Roche 11214667001, Basel, Switzerland) were used.

BsParB protein labeling with fluorescent dyes. BsParB(R80A) proteins were incubated with sulfo-Cyanine3 NHS ester dye (Lumiprobe 11320, Hunt Valley, MD) at 4°C overnight. Labeled protein was separated from free dye using Micro Bio-Spin P-6 gel columns (Bio-Rad 7326221, Hercules, CA). Each labeled protein and Cyanine3 dye concentrations were measured three times using Nanodrop, and the averaged values were used as final concentrations. The protein labeling efficiencies were 30.1%, 32.0%, and 30.0% for BsParB(R80A), KCK-BsParB(R80A), and BsParB(R80A)-KCK, respectively. These numbers correspond to about 0.6 Cyanine3 dyes per each BsParB protein dimer.

Single-molecule flow-stretching assays. Surface-passivated coverglasses were prepared by aminopropyl silanization and PEGylation (PEG: polyethylene glycol) as previously described[22,23]. A microfluidic flow cell was constructed from a quartz plate (Technical Glass Product, Paineville, OH) adhered to the PEGylated coverglass via double-sided tape (Grace Bio-Labs, Bend, OR) with rectangular cuts that make up the flow cell channels. Inlet and outlet tubing were inserted through holes on the quartz plate and made air-tight with epoxy[22,23]. In-depth description of single-molecule flow-stretching assays was already provided in previous publications[22–24]. Briefly, about 4% of the PEG on the surface-passivated coverglass contains biotins that serve as a neutravidin binding platform. Pre-mixed quantum dot-labeled biotinylated lambda DNA (or *parS* DNA) was introduced to allow the DNA surface tethering. For experiments with labeled proteins, quantum dot incubation with biotinylated DNA is omitted. After washing unbound DNAs and quantum dots, an intended concentration of BsParB protein was flowed in (with and without nucleotides). Unless otherwise stated, the buffer composition was 10 mM Tris, pH 7.5, 100 mM NaCl, and 2.5 mM $MgCl_2$. For the experiments without magnesium ions, the 2.5 mM $MgCl_2$ was omitted. CTPγS was custom-synthesized (Jena Bioscience, Jena, Germany). The single-molecule imaging was performed on a semi-custom microscope with a 532-nm laser (Coherent, Santa Clara, CA) built upon the IX-83 total internal reflection fluorescence (TIRF) microscope (Evident Scientific, Olympus, Waltham, MA). The images

were recorded every 200 milliseconds with 100-millisecond exposure time using the Micro-Manager software[25].

Regions-of-interest (ROIs) of DNA compaction events were determined using ImageJ (FIJI) software, and the

positions of quantum dots as a function of time were determined by Gaussian-fitting-based custom-written Matlab

software codes[23]. The compaction rate measurements were taken from distinct samples (quantum dot-bound

DNAs).

Bacterial strains and growth. *Bacillus subtilis* strains were derived from the prototrophic strain PY79[26]. Cells

were grown in defined rich Casein Hydrolysate (CH) medium[27] at 37˚C. Strain, plasmids, oligonucleotides, and

next-generation sequencing samples used in this study can be found in Tabs 6-9 in the Supplementary File.

Fluorescence microscopy. Fluorescence microscopy was performed using a Nikon Ti2 microscope (Nikon

Instruments, Melville, NY) equipped with Plan Apo 100x/1.45NA phase contrast oil objective and an sCMOS

camera. Images were cropped and adjusted using MetaMorph software. Final figure preparation was performed

in Adobe Illustrator.

ChIP-seq. Chromatin immunoprecipitation (ChIP) was performed as described previously[28,29]. Briefly, cells were

crosslinked using 3% formaldehyde for 30 min at room temperature and then quenched using 125 mM glycine,

washed using PBS, and lysed using lysozyme. Crosslinked chromatin was sheared to an average size of 250

bp by sonication using Qsonica Q800R2 water bath sonicator. The lysate was precleared using Protein A

magnetic beads (GE Healthcare/Cytiva 28951378, Marlborough, MA) and was then incubated with anti-ParB

antibodies[30] overnight at 4˚C. The next day, the lysate was incubated with Protein A magnetic beads for 1h at

4˚C. After washes and elution, the immunoprecipitate was incubated at 65˚C overnight to reverse the crosslinks.

The DNA was further treated with RNaseA, Proteinase K, extracted with PCI, resuspended in 100 µl EB and

used for library preparation with the NEBNext Ultra II kit (E7645). The library was sequenced using Illumina

NextSeq500 (Illumina, San Diego, CA) at IU Center for Genomics and Bioinformatics. The sequencing reads

were mapped to *B. subtilis* PY79 genome (NCBI Reference Sequence NC_022898.1) using CLC Genomics

Workbench (Qiagen, Hilden, Germany). We note that the genome coordinate of this genome is shifted compared

to the *B. subtilis* 168 genome (NC000964) used in our previous study[12]. Sequencing reads were normalized by

the total number of reads, plotted and analyzed using R.

Immunoblot analysis. Exponentially growing cells were collected and resuspended in lysis buffer (20 mM Tris

pH 7.0, 1 mM EDTA, 10 mM MgCl$_2$, 1 mg/ml lysozyme, 10 µg/ml DNase I, 100 µg/ml RNase A, 1 mM PMSF and

1% proteinase inhibitor cocktail (Sigma-Aldrich P-8340, St. Louis, MO) to a final OD$_{600}$ of 10 for equivalent loading.

The cell resuspensions were incubated at 37˚C for 10 min for lysozyme treatment, followed by the addition of an

equal volume of 2x Laemmli Sample Buffer (Bio-Rad 1610737, Hercules, CA) containing 10% β-Mercaptoethanol.

Samples were heated for 15 min at 65˚C prior to loading. Proteins were separated by precast 4-20%

polyacrylamide gradient gels (Bio-Rad 4561096, Hercules, CA) and electroblotted onto mini PVDF membranes

using Bio-Rad Transblot Turbo system and reagents (Bio-Rad 1704156, Hercules, CA). The membranes were

blocked in 5% nonfat milk in phosphate-buffered saline (PBS) with 0.5% Tween-20, then probed with anti-ParB

(1:5000)[30] or anti-SigA (1:10,000)[31] diluted into 3% BSA in 1x PBS-0.05% Tween-20. Primary antibodies were

detected using Immun-Star horseradish peroxidase-conjugated goat anti-rabbit antibodies (Bio-Rad 1705046,

Hercules, CA) and Western Lightning Plus ECL chemiluminescence reagents as described by the manufacturer

(Perkin Elmer NEL1034001, Waltham, MA). The signal was captured using ProteinSimple Fluorchem R system.

The intensity of the bands was quantified using ProteinSimple AlphaView software.

Plasmid construction for *in vivo* experiments.

pWX1092 [*pelB::Psoj-spo0J(ΔparS)-mgfpmut3 tet*] was constructed by an isothermal assembly reaction

containing three fragments: 1) pWX516 digested with HindIII and BamHI, and gel purified; 2) *spo0J (ΔparS)*

amplified from pWX563[12] using oWX2974 and oWX2975; 3) *mgfpmut3* amplified from pWX563[12] using oWX2976

and oWX2977. pWX516 contains *pelB::Psoj (tet)*. The construct was sequenced using oWX507, oWX669, and

oWX670.

pWX1093 [*pelB::Psoj-KCK-spo0J(ΔparS)-mgfpmut3 tet*] was constructed by an isothermal assembly reaction

containing three fragments: 1) pWX516 digested with HindIII and BamHI, and gel purified; 2) *KCK-spo0J (ΔparS)*

amplified from pWX563[12] using oWX2978 and oWX2975; 3) *mgfpmut3* amplified from pWX563[12] using oWX2976

and oWX2977. pWX516 contains *pelB::Psoj (tet)*. The construct was sequenced using oWX507, oWX669, and

oWX670.

pWX1103 [*pelB::Psoj-mgfpmut3-spo0J-R80A(ΔparS)-KCK cat*] was constructed by an isothermal assembly

reaction containing two PCR products: 1) pWX611 amplified using oWX3001 and oWX418; 2) pWX611 amplified

using oWX3002 and oWX2071. This procedure introduced the R80A mutation to pWX611[12], which is *pelB::Psoj-*

mgfpmut3-spo0J(ΔparS)-KCK cat. The construct was sequenced using oWX507, oWX669, and oWX670.

pWX1104 [*pelB::Psoj-spo0J-R80A(ΔparS)-mgfpmut3 tet*] was constructed by an isothermal assembly reaction

containing two PCR products: 1) pWX1092 amplified using oWX3001 and oWX418; 2) pWX1092 amplified using

oWX3002 and oWX2071. This procedure introduced the R80A mutation to pWX1092. The construct was

sequenced using oWX507, oWX669, and oWX670.

pWX1105 [*pelB::Psoj-KCK-spo0J-R80A(ΔparS)-mgfpmut3 tet*] was constructed by an isothermal assembly

reaction containing two PCR products: 1.) pWX1093 amplified using oWX3001 and oWX418; 2) pWX1093

amplified using oWX3002 and oWX2071. This procedure introduced the R80A mutation to pWX1093. The

construct was sequenced using oWX507, oWX669, and oWX670.

pWX1106 [*pelB::Psoj-soj-spo0J-R80A(ΔparS)-KCK cat*] was constructed by an isothermal assembly reaction

containing two PCR products: 1) pWX612 amplified using oWX3001 and oWX418; 2) pWX612 amplified using

oWX3002 and oWX2071. This procedure introduced the R80A mutation to pWX612[12], which is *pelB::Psoj-soj-*

spo0J(ΔparS)-KCK cat. The construct was sequenced using oWX507, oWX1086, and oML77.

pWX1107 [*pelB::Psoj-KCK-spo0J(ΔparS) tet*] was constructed by an isothermal assembly reaction containing

two PCR products: 1) pWX1093 amplified using oWX3004 and oWX418; 2) pWX1093 amplified using oWX3003

and oWX2071. This procedure introduced a stop codon and removed *mgfpmut3* from pWX1093. The construct

was sequenced using oWX507 and oML85.

pWX1108 [*pelB::Psoj-KCK-spo0J-R80A(ΔparS) tet*] was constructed by an isothermal assembly reaction

containing two PCR products: 1) pWX1107 amplified using oWX3001 and oWX418; 2) pWX1107 amplified using

oWX3002 and oWX2071. This procedure introduced the R80A mutation to pWX1107. The construct was

sequenced using oWX507 and oML85.

pWX1167 *[pelB::Psoj-ECE-spo0J(ΔparS) tet]* was constructed by an isothermal assembly reaction containing

two PCR products: 1) pWX1107 amplified using oWX3197 and oWX418; 2) pWX1107 amplified using oWX3198

and oWX2071. This procedure introduced the ECE tag and removed the KCK tag from pWX1107. The construct

was sequenced using oWX507 and oML85.

pWX1168 *[pelB::Psoj-ECE-spo0J-R80A(ΔparS) tet]* was constructed by an isothermal assembly reaction

containing two PCR products: 1) pWX1108 amplified using oWX3197 and oWX418; 2) pWX1108 amplified using

oWX3198 and oWX2071. This procedure introduced the ECE tag and removed the KCK tag from pWX1108.

The construct was sequenced using oWX507 and oML85.

pWX1169 *[pelB::Psoj-ECE-spo0J(ΔparS)-mgfpmut3 tet]* was constructed by an isothermal assembly reaction

containing two PCR products: 1) pWX1093 amplified using oWX3197 and oWX418; 2) pWX1093 amplified using

oWX3198 and oWX2071. This procedure introduced the ECE tag and removed the KCK tag from pWX1093.

The construct was sequenced using oWX507, oWX669, and oWX670.

pWX1170 *[pelB::Psoj-ECE-spo0J-R80A(ΔparS)-mgfpmut3 tet]* was constructed by an isothermal assembly

reaction containing two PCR products: 1) pWX1105 amplified using oWX3197 and oWX418; 2) pWX1105

amplified using oWX3198 and oWX2071. This procedure introduced the ECE tag and removed the KCK tag from

pWX1105. The construct was sequenced using oWX507, oWX669, and oWX670.

Strain construction

B. subtilis strains were generated by successive transformations of plasmids or genomic DNA.

Statistics and reproducibility

Not all measurement groups passed the normality test (See Tab 3 in the Supplementary File). Therefore, in this study, we report the results of nonparametric Mann-Whitney test in Fig. 1a, f. However, we obtained similar results from two-sided Welch's t-test (Supplementary Fig. 3a, b) since the t-test results are still valid when the sample sizes are large (>25) and there are not extreme outliers[19]. All the statistical analyses (Shapiro-Wilk normality test, Mann-Whitney test, and two-sided Welch's t-test due to different variances and sample sizes) for DNA compaction rates were performed using Prism software (GraphPad, San Diego, CA). The exact sample sizes (*N*), mean, and standard error of the mean are provided in Tabs 1 and 2 in the Supplementary File. The normality test results are available in Tab 3 in the Supplementary File. Tabs 4 and 5 in the Supplementary File show the exact *p*-values for comparing wild-type (and its KCK-tagged versions) and R80A mutant (and its KCK-tagged versions) compaction rates, respectively. The reproducibility of single-molecule experiments for each experimental condition was checked by performing the same experiments at least three times.

Data availability

A list of figures that have associated raw data can be found from Tabs 6 and 9 in the Supplementary File. Single-molecule analysis data can be found in Tabs 1-5 in the Supplementary File. The datasets generated during and/or analyzed during the current study are available from the corresponding authors on reasonable request.

Code availability

The Matlab codes used in single-molecule data are available from our previous publication[23]. Alternatively, the codes will be available from the corresponding author (H.K.) upon request.

References

1.   Toseland, C. P. Fluorescent labeling and modification of proteins. *J Chem Biol* **6**(3), 85-95 (2013).

2.   Jensen, K. S., Pedersen, J. T., Winther, J. R. & Teilum, K. The pKa value and accessibility of cysteine residues are key determinants for protein substrate discrimination by glutaredoxin. *Biochemistry* **53**, 2533–2540 (2014).

3. Lobocki, M. *et al.* High-Yield Site-Specific Conjugation of Fibroblast Growth Factor 1 with
Monomethylauristatin e via Cysteine Flanked by Basic Residues. *Bioconjug Chem* **28**, 1850–1858
(2017).

4. Parente, A., Merrifield, B., Geraci, G. & D'Alessio, G. Molecular Basis of Superactivity of Cysteine
Residues 31 and 32 of Seminal Ribonuclease. *Biochemistry* **24**, 1098–1104 (1985).

5. Nawrocka, D., Krzyscik, M. A., Opaliński, Ł., Zakrzewska, M. & Otlewski, J. Stable fibroblast growth
factor 2 dimers with high pro-survival and mitogenic potential. *Int J Mol Sci* **21**, 4108 (2020).

6. Larson, A. G. *et al.* Liquid droplet formation by HP1α suggests a role for phase separation in
heterochromatin. *Nature* **547**, 236–240 (2017).

7. Sallee, N. A. *et al.* The pathogen protein EspFU hijacks actin polymerization using mimicry and
multivalency. *Nature* **454**, (2008).

8. Ginosyan, A. A., Grintsevich, E. E. & Reisler, E. Neuronal drebrin A directly interacts with mDia2 formin
to inhibit actin assembly. *Mol Biol Cell* **30**, 646–657 (2019).

9. Son, S. *et al.* Molecular height measurement by cell surface optical profilometry (CSOP). *Proc Natl Acad
Sci U S A* **117**, 14209–14219 (2020).

10. Hansen, S. D. & Mullins, R. D. VASP is a processive actin polymerase that requires monomeric actin for
barbed end association. *Journal of Cell Biology* **191**, 571–584 (2010).

11. van den Ent, F., Johnson, C. M., Persons, L., de Boer, P. & Löwe, J. Bacterial actin MreB assembles in
complex with cell shape protein RodZ. *EMBO Journal* **29**, 1081–1090 (2010).

12. Graham, T. G. W. *et al.* ParB spreading requires DNA bridging. *Genes Dev* **28**, 1228–1238 (2014).

13. Badrinarayanan, A., Le, T. B. K. & Laub, M. T. Bacterial chromosome organization and segregation.
Annu Rev Cell Dev Biol **31**, 171–199 (2015).

14. Jalal, A. S. B. & Le, T. B. K. Bacterial chromosome segregation by the ParABS system. *Open Biol* **10**,
200097 (2020).

15. Osorio-Valeriano, M. *et al.* ParB-type DNA Segregation Proteins Are CTP-Dependent Molecular
Switches. *Cell* **179**, 1512-1524.e15 (2019).

16. Soh, Y. M. *et al.* Self-organization of parS centromeres by the ParB CTP hydrolase. *Science (1979)* **366**,
1129–1133 (2019).

17. Jalal, A. S., Tran, N. T. & Le, T. B. ParB spreading on DNA requires cytidine triphosphate in vitro. *Elife*
9, 1–24 (2020).

18. Antar, H. *et al.* Relief of ParB autoinhibition by parS DNA catalysis and recycling of ParB by CTP
hydrolysis promote bacterial centromere assembly. *Sci Adv* **7**, esbj2854 (2021).

19. le Cessie, S., Goeman, J. J. & Dekkers, O. M. Who is afraid of non-normal data? Choosing between
parametric and non-parametric tests. *European journal of endocrinology* vol. 182 E1–E3 Preprint at
https://doi.org/10.1530/EJE-19-0922 (2020).

20. Song, D., Rodrigues, K., Graham, T. G. W. & Loparo, J. J. A network of cis and trans interactions is
required for ParB spreading. *Nucleic Acids Res* **45**, 7106–7117 (2017).

21. Autret, S., Nair, R. & Errington, J. Genetic analysis of the chromosome segregation protein SPoOj of
Bacillus subtilis: Evidence for separate domains involved in DNA binding and interactions with SOj
protein. *Mol Microbiol* **41**, 743–755 (2001).

22. Kim, H. & Loparo, J. J. Observing Bacterial Chromatin Protein-DNA Interactions by Combining DNA
Flow-Stretching with Single-Molecule Imaging. in *Bacterial chromatin, Methods in Molecular Biology*
277–299 (Humana Press (Springer), 2018). doi:https://doi.org/10.1007/978-1-4939-8675-0_15.

23. Kim, H. & Loparo, J. J. Multistep assembly of DNA condensation clusters by SMC. *Nat Commun* **7**,
10200 (2016).

24. Kim, H. *et al.* The gene-silencing protein MORC-1 topologically entraps DNA and forms multimeric
assemblies to cause DNA compaction. *Mol Cell* **75**, 700–710 (2019).

25. Edelstein, A. D. *et al.* Advanced methods of microscope control using µManager software. *J Biol*
Methods **1**, e10 (2014).

26. Youngman, P. J., Perkins, J. B. & Losick, R. Genetic transposition and insertional mutagenesis in
Bacillus subtilis with Streptococcus faecalis transposon Tn917. Proc. Nati. Acad. Sci. USA 80, 2305-
2309 (1983).

27. Harwood, C. R. & Cutting, S. M. *Molecular Biological Methods for Bacillus*. (Wiley, 1990).

28. Wang, X., Brandão, H. B., Le, T. B. K., Laub, M. T. & Rudner, D. Z. Bacillus subtilis SMC complexes
juxtapose chromosome arms as they travel from origin to terminus. *Science (1979)* **527**, 524–527
(2017).

29. Wang, X. *et al.* Condensin promotes the juxtaposition of DNA flanking its loading site in Bacillus subtilis.
Genes Dev **29**, 1661–1675 (2015).

30. Chi-hong Lin, D., Anne Levin, P. & Grossman, A. D. Bipolar localization of a chromosome partition
protein in Bacillus subtilis. *Proc. Nati. Acad. Sci. USA* **94**(9), 4721-4726 (1997).

31. Fujita, M. Temporal and selective association of multiple sigma factors with RNA polymerase during
sporulation in Bacillus subtilis. *Genes to Cells* **5**, 79–88 (2000).

Acknowledgements

We thank the Kim and Wang labs for discussions and support, Xheni Karaboja for assisting with ChIP-

seq and microscopy experiments, Patrick Sheets for assisting with strain building, Candice Etson for

providing the *E. coli* strain containing *parS*-lambda DNA, Thomas Graham for providing plasmids for

BsParB wild-type and R80A mutant, Alan Grossman for strains and antibodies, and the Indiana University

Center for Genomics and Bioinformatics for high throughput sequencing. We thank Thomas Graham for

reading the manuscript and providing insightful feedback. Support for this work comes from National

Institutes of Health R01GM141242 (X.W.) and R35GM143093 (H.K.).

Author contributions

506 M.M. and H.K. purified proteins, performed single-molecule experiments, and analyzed data. L.E.W.

507 constructed plasmids and strains, performed microscopy, and immunoblot analysis. Z.R., Q.L., and X.W.

508 performed ChIP-seq and analysis. X.W. designed, analyzed, and supervised the *in vivo* experiments. H.K.

509 designed, analyzed data, and supervised the *in vitro* experiments. M.M. and L.E.W. contributed to writing

510 the method sections of the paper. X.W. and H.K. wrote the paper with input from all authors.

Competing interests

513 All authors declare the absence of any competing interests.