***Wolbachia* rescue of several *Drosophila melanogaster bag of marbles* protein coding mutants**

Miwa Wenzel[1] and Charles F. Aquadro[1]

[1]Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY, USA

[*]Corresponding author

Email:

Abstract

The *D. melanogaster* protein coding gene *bag of marbles* (*bam*) plays a key role in early male

and female reproduction by forming complexes with partner proteins to promote differentiation

in gametogenesis. Like another germline gene, *Sex lethal*, *bam* genetically interacts with the

endosymbiont *Wolbachia*, as *Wolbachia* rescues the reduced fertility of a *bam* hypomorphic

mutant. Here, we explored the specificity of the *bam-Wolbachia* interaction by generating 22

new *bam* mutants. We find *Wolbachia* rescues all six mutants of partially reduced fertility, but

none of the four with severely reduced fertility. There is no specificity between the rescue and

the known binding regions of *bam*, suggesting *Wolbachia* does not interact with one singular

bam partner to rescue fertility. We further tested if *Wolbachia* interacts with *bam* in a non-

specific way, by increasing *bam* levels or acting upstream in germline stem cells. However, a

fertility assessment of a *bam* RNAi knockdown mutant reveals that *Wolbachia* rescue is specific

to functionally mutant *bam* alleles and we find no obvious evidence of *Wolbachia* interaction

with germline stem cells in *bam* mutants.

Author Summary

Reproduction in the *Drosophila melanogaster* fruit fly is dependent on the *bag of marbles* (*bam*)

gene, which acts early in the process of generating eggs and sperm. Mutations to this gene

negatively impact the fertility of the fly, causing it to be sterile or have less progeny.

Interestingly, the bacteria *Wolbachia*, which resides within reproductive cells across a wide

range of insects, partially restores the fertility of several *bam* mutants that have mutations

affecting the resultant Bam protein that is produced. The protein function of Bam is further

suggested to be important by the lack of fertility rescue for a fly that has a fertility defect due to

low expression of a non-mutated Bam protein. Previous work makes similar conclusions about

Wolbachia with another reproductive gene, *Sex lethal* (*Sxl*), highlighting the potential for rescue

of fertility mutants to occur in a similar way across different genes. An understanding of the

ways in which *Wolbachia* can affect host reproduction provides us with context with which to

frame *Wolbachia*'s impact on host genes, such as *bam* and *Sxl*.

Introduction

The *bag of marbles* (*bam*) gene plays a critical role in female and male gametogenesis in *D. melanogaster.* In the stem cell niche of the ovaries and testes lies a collection of germline stem cells (GSCs) that asymmetrically divide to self-renew and produce differentiating daughter cells. In females, *bam* acts as a master regulator and its expression induces daughter cell differentiation [1]. In males, *bam* functions later in gametogenesis where it initiates terminal differentiation of the spermatogonia that eventually become sperm [1, 2, 3, 4].

In female *D. melanogaster, bam* is largely repressed in GSCs by signaling it receives from the stem cell niche [5, 6, 7]. Bam that escapes silencing in GSCs is antagonized by eukaryotic translation initiation factor 4A (eIF4A) [8]. When a GSC divides, a daughter cell is displaced from the stem cell niche, allowing *bam* to now be expressed. In the daughter cell, Bam functions to bring together various proteins that repress self-renewal and promote differentiation, leading to the establishment of a cystoblast.

In both male and female *D. melanogaster, bam* null mutants are completely sterile and have a tumorous cytological phenotype [1, 9]. A partial loss-of-function mutant (*bam^{bw}*) has reduced fertility in females and is sterile in males [10, 11]. In females, the fertility defect of the *bam^{bw}* mutant is rescued by the endosymbiont bacteria *Wolbachia pipientis* [11].

Wolbachia is estimated to inhabit 50% or more of insect species and is an obligate bacterium in filarial nematodes [12, 13, 14]. Across both arthropods and nematodes, *Wolbachia* manipulates reproduction to promote its own maternal transmission (reviewed in [15, 16]). For example, *Wolbachia* famously induces cytoplasmic incompatibility, in which a female's progeny are not viable if they do not harbor the same *Wolbachia* strain as the male parent. *Wolbachia* is also known to induce parthenogenesis, feminization, and male-killing of host progeny, all of

which promote its maternal transmission. Thus, the rescuing of the hypomorphic *bam* mutant

aligns with the evolutionary interests of *Wolbachia.*

Wolbachia also rescues the sterility of hypomorphs of another GSC gene, *Sex lethal* (*Sxl*)

[17]. Starr and Cline [17] proposed that the rescue is dependent on the point mutation induced in

the *Sxl* mutant and not on the severity of the mutant, though they suggested that rescue cannot

happen if *Sxl* protein levels are too low. Of note is that the *Sxl* mutants they studied were in

different genetic backgrounds, which may have affected the results observed and the conclusions

made. Such background effects were observed with *bam* in [18], where females with the same

L255F *bam* mutation had strikingly different fertility across two different genetic backgrounds of

D. melanogaster. Nevertheless, further research into one of the *Sxl* mutants, *Sxl^{f4}*, showed that

the rescue is partially due to a restoration of GSCs by the *Wolbachia* protein toxic manipulator of

oogenesis (TomO) [19].

Mutants of additional germline genes *meiotic P26* (*mei-P26*), *ovarian tumor* (*otu*), and

sans fille (*snf*) are not rescued by *Wolbachia,* despite having similar severity of an egg-absence

phenotype to some *Sxl* mutants [17]. These results suggest the *Wolbachia* interaction may be

specific to a functional pathway in female *D. melanogaster,* possibly through a direct, physical

interaction with a gene product like Sxl that is expressed in the germline.

In this study, we explore the *bam Wolbachia* interaction, building off the findings for the

bam^{bw} hypomorphic mutant. The *bam^{bw}* hypomorphic allele is a L255F change in a gene region

critical for *bam* function. Thus far, four functional regions of *bam* have been documented: The

CAF40 binding motif (residues 13-36) [20], the Benign gonial cell neoplasm (Bgcn) binding

region (residues 100-350) [21], the COP9 signalosome 4 (Csn4) binding region (residues 151-

350) [21], and a ubiquitin binding region (residues 200-250) [22]. *Bam* also has a PEST domain

(residues 402-434), which marks the protein for rapid degradation [1]. *Bam*'s specific binding

regions and their interactions with binding partners are illustrated in Figures 1a and 1b

respectively.

Figure 1. *Bam* is involved in several processes that regulate the differentiation of GSCs in female *D. melanogaster*. (a) Schematic of the five known binding or functional regions of *bam* are colored across the coding sequence exons of the gene. Each black vertical bar represents the location of a new mutant generated with alanine replacements in this study. (b) In GSCs (left), Bam is repressed and antagonized by eIF4A. In the differentiating daughter cells (right), Bam is expressed and coordinates other molecules to inhibit the self-renewal pathway or promote the differentiation pathway. Bam has been shown to genetically and physically interact with additional molecules that are not shown.

To repress GSC self-renewal, Bam forms a complex with Bgcn, Meiotic-P26 (Mei-P26),

Brain tumor (Brat), a deadenylase, and either Argonaute-1 (Ago1) or Sxl to destabilize *nanos*

(*nos*) mRNA [23, 24, 25]. The destabilization of *nos* mRNA may occur through Bam's

interaction with the CAF40 which is part of the CCR4-NOT deadenylase complex [20, 26].

Thus, in GSCs, *nos* mRNA is translated to promote GSC self-renewal; in differentiating daughter

cells, *nos* mRNA is destabilized by the Bam complex, inhibiting *nos* translation and the self-

renewal pathway.

Bam also represses the self-renewal pathway through its interaction with Csn4 [21]. In

GSCs, the COP9 complex is composed of nine Csn subunits. When Bam is expressed in

differentiating daughter cells, it breaks up the complex by sequestering Csn4, thereby inhibiting

the GSC self-renewal pathway.

Additionally, Bam promotes differentiation by forming a complex with Otu to

deubiquitinate Cyclin A (CycA) [22]. In GSCs, CycA is degraded through a process involving

the ubiquitin-conjugating enzyme Effete [27]. In differentiating GSC daughter cells, it is

proposed that Bam binds to both Otu and to proteins such as CycA, which have ubiquitin chains

that mark them for degradation. When in a complex with Bam, Otu deubiquitinates CycA,

stabilizing CycA levels and thus promoting GSC daughter cell differentiation.

Here, we have generated 22 new *bam* mutants and analyzed the genetic interaction

between these mutants and *Wolbachia* to better understand the extent of the *bam-Wolbachia*

interaction. We are partly motivated by the observation of rapidly diversifying amino acids at

bam across the *Drosophila* phylogeny [28, 29, 30]. We have previously hypothesized that

Wolbachia could contribute to this rapid evolution observed at *bam* [11, 29, 30, 31]. Intriguingly,

recent results for *Sxl* could be consistent with a parallel hypothesis for that gene, as the excess

nonsynonymous substitutions observed in *Sxl* cluster near the locations of the alleles that

genetically interact with *Wolbachia* [32].

In this study, we find that *Wolbachia* genetically interacts with alanine point mutants

throughout the *bam* gene, suggesting that *Wolbachia* rescue of *bam* mutants is not specific to any

single binding/functional region of *bam.* We find no evidence for two hypotheses of how

Wolbachia could rescue *bam* in a non-specific manner: There is no effect of *Wolbachia* on a

hypomorph-like *bam* knockdown mutant, nor is there an effect of *Wolbachia* on the frequency of

GSC mitosis that occurs upstream of *bam.*

Materials and Methods

Design of putative *bam* hypomorph alleles

22 *bam* mutant alleles were designed with a Venus fluorescent tag (Fig. 1a), described

further below. The known Bam binding residues in the CAF40 binding motif were targeted for

replacement [20], producing independent alleles with mutations of L17A, M24A, and L28A. The

remaining mutant alleles were designed through an alanine scanning approach [33] that targeted

the known functional and/or binding regions of *bam.* Strings of charged residues were changed

to alanine by choosing an alanine codon with similar usage bias in *D. melanogaster* as that of the

original residue [34]. Additionally, we designed an allele to recreate the *bam^bw^* L255F mutation

as a transgene with Venus tag. All new alleles are listed in Supplemental file S1, with the

original and mutated codons.

Generation of transgenic flies

Transgenic fly lines carrying distinct putative hypomorphic *bam* alleles and a control

transgenic fly line were generated using PhiC-mediated integration [35]. Approximately 1.5 kb

upstream and 700 bp downstream regions of *bam* were cloned from genomic DNA of a *D.*

melanogaster bam::Venus line [36]. This construct is similar to that of the *D. melanogaster bam*

YFP line from [11] but contains a 36bp glycine-glycine-serine linker sequence between the 3'

end of the *bam* coding sequence and the Venus tag to allow for proper folding of both proteins.

The *bam::Venus* sequence was cloned into a pMiniT 2.0 vector, using the NEB PCR cloning kit.

Mutant alleles were subsequently made in the established *bam::Venus* pMiniT 2.0 plasmid using

the NEB Q5 Site-Directed Mutagenesis Kit, with primers designed online using NEBase

Changer (Supplemental table S1) [37]. Briefly, mutagenic primers were used to amplify the

bam::Venus pMiniT 2.0 plasmid using with a PCR cycle of 98°C for 30 seconds; 25 cycles of

98°C for 10 seconds, 57-72°C annealing for 30 seconds, 72°C for 3.5 minutes; and 72°C for 2

minutes. PCR products were treated with kinase, ligase, and DpnI before being transformed into

competent cells. The resultant plasmids were subcloned into the pCasper\attB vector, a gift from

Dan Barbash that was used by [11] for PhiC transgenesis. Plasmids were verified by sequencing.

Plasmids were injected into embryos of a *w*, *nos*-int; P{CaryP}attP40 *D. melanogaster*

line by GenetiVision. G_0 flies were mated with a w^{1118} isogenic line and F_1 progeny were

screened for successful transgenesis by selecting for those with mini-white eye color. Mutant

lines were identified by Sanger sequencing.

All transgenic flies were assessed for function in a *bam* null background. Thus, the

genotype that was assessed in all experiments was *w*; [w^{+mc}, *bam::Venus*]/+; *bam*^{Δ59}/*bam*^{Δ86},

where *bam::Venus* represents the control *bam* allele or one of the 22 mutant alleles. For the

remainder of this paper, "transgenic females" refers to flies of this genotype. *Wolbachia* status of

the assessed flies (uninfected or *Wolbachia* infected) was established by maternal inheritance

when crossing in the *bam* null alleles.

Fertility assays of female transgenic flies

Virgin transgenic females and virgin CantonS males were collected over 48 hours and

aged two days. At least 30 individual crosses of one female and two males were set up on yeast-

glucose food (minimum of 30, maximum of 88 crosses were set per genotype tested). All crosses

were kept in an incubator at 25°C with a 12-hour light/dark cycle. Parents were removed on the

seventh day. Progeny were counted every other day over seven days, starting on the day of first

eclosion. If the female parent died by the time of removal, data for that cross was discarded. Five

independent fertility assays were conducted to cover all 22 mutants, with a transgenic control (*w*;

[*w^{+mc}*, *bam::Venus*]/+; *bam^{Δ59}*/*bam^{Δ86}*) present in each assay.

The total progeny after seven days was summed and estimation statistics [38] were

performed to determine first, if a fertility defect was present for a mutant, and second, if there

was a *Wolbachia* rescue effect. Estimation statistics and plots were generated using the dabestr

package in R (v0.3.0). While this offers an alternative statistical test that allows us to easily

visualize effect size, we also report p-values for ease and convention. P-values were calculated

through the associated website as results of a two-sided permutation t-test [39], as they were not

provided in the R package.

Cytological observations of the ovaries of female transgenic flies

For cytology of transgenic females with fertility defects and control females, flies were

aged 2-3 days and then dissected and prepared for confocal microscopy as in [11, 30]. Ovaries

were dissected in ice-cold 1x PBS and fixed in 4% paraformaldehyde (Electron Microscopy

Sciences) for 15 minutes. Tissue was rinsed 3x and washed 4x, respectively, with PBST (1X

PBS, 0.1% Triton-X 100), blocked in PBTA (1X PBS, 0.1% Triton-X100, 3% BSA) (Alfa

Aesar), and then incubated with primary antibodies overnight. The rinse and wash steps were

repeated, followed by an overnight incubation with a goat anti-serum to minimize fluorescent

background binding, and then a 1.5-hour incubation with secondary antibodies. The rinse and

wash steps were repeated and tissue was mounted in ProLong Diamond AntiFade (Invitrogen).

The primary antibodies that were used were anti-Vasa (anti-rat, 1:20, Developmental

Studies Hybridoma Bank (DSHB)), anti-Hts-1B1 (anti-mouse, 1:40, DSHB), and anti-GFP (anti-

rabbit, 1:200, Invitrogen Cat No: G10362). Secondary antibodies that were used were Alexa488

(goat anti-rat, Invitrogen Cat No: A-11006), Alexa568 (goat anti-mouse, Invitrogen Cat No: A-

11031), and Alexa633 (goat anti-rabbit, Invitrogen Cat No: A-21070), all at 1:500.

A Zeiss i880 confocal microscope was used for all imaging with 405, 488, and 568nm

laser lines at 63X (Plan-Apochromat 1.4 NA, oil) (Cornell BRC Imaging Core Facility). Images

were edited using Fiji [40].

RNAi knockdown of *bam*

We performed reciprocal crosses between a *Wolbachia* infected *nos*-Gal4 and an

uninfected UAS-*bam*[HMS00029] line (a TriP *bam* shRNA line) (Bloomington Drosophila Stock

Center lines 4442 and 33631, respectively) to generate the *bam* knockdown lines.

Bam expression was evaluated by RT-qPCR. RNA of four samples consisting of three 1-

2 day old female flies each were prepared using the NEB Monarch Total RNA Miniprep Kit.

RT-qPCR was performed with 10µL reactions (5µL Luna Universal One-Step Reaction Mix

(2X), 0.5µL Luna WarmStart RT Enzyme Mix (20X), 0.4µL forward primer, 0.4µL reverse

primer, 2μL template RNA, 1.7μL water), using three technical replicates and a standard curve.

Reverse transcription was performed at 55°C for 10 minutes, followed by initial denaturation at

95°C for 1 minute, and then 40 cycles of 95°C for 10 seconds and 60°C for 30 seconds. Primers

included 5'GCGCAATCGAAACGGAAACT3' and 5'GAGTAGCGGTGCTCCAGATC3' for

bam and 5'CCGCTTCAAGGGACAGTATC3' and 5'CAATCTCCTTGCGCTTCTTG3' for the

reference gene *Ribosomal protein L32* (*Rpl32*) (*Rpl32* primers are from [41]). Relative

expression to a wildtype *bam* control was done by the method of [42].

A fertility assay was performed as described above, but at 29°C to increase the activity of

the Gal4-UAS system. Estimation statistics [38] were used to compare the mean fertility of the

uninfected and infected *bam* knockdown mutants.

Additionally, an assay for nurse-cell-positive egg chambers was performed, as has been

previously done for a supplemental assessment of *bam*'s function [11, 18]. Briefly, ovaries of 1-2

day old flies were fixed in 4% paraformaldehyde, rinsed and washed with PBST as described

above, and then mounted in ProLong Glass with NucBlue (ThermoFisher). Ovaries were imaged

on a Zeiss i880 confocal microscope at 40X and 63X (Plan-Apochromat 1.4 NA, oil), using a

405nm laser line (Cornell BRC Imaging Core Facility). Egg chambers with any identifiable

large, polyploid nuclei of nurse cells indicated an event of successful Bam function and were

counted as nurse-cell-positive egg chambers. Statistically significant results were determined

with the Wilcoxon signed rank test.

Analysis of GSC mitosis frequency

Nurse cell assays were performed as described above to confirm that the L255F bam

mutation that we created as a transgene produced a partial *bam* mutant phenotype and *Wolbachia*

rescue like that of the original *bam^{bw}* hypomorph.

Then, we assessed GSC mitosis frequency in the transgenic control (*w*; [*w^{+mc}*,

bam::Venus]/+; *bam^{Δ59}*/*bam^{Δ86}*) and the transgenic L255F mutant (*w*; [*w^{+mc}*, *bam^{L255F}::Venus*]/+;

bam^{Δ59}/*bam^{Δ86}*) by comparing the number of actively dividing GSCs indicated by mitosis marker

pH3 to the number of non-dividing GSCs. Flies were aged 2-3 days at 25°C, at which time

ovaries were dissected, immunostained and prepared for confocal microscopy, and imaged as

described above.

Primary antibodies that were used were anti-Vasa (anti-rat, 1:20, DSHB), anti-GFP (anti-

rabbit, 1:200, Invitrogen Cat No: G10362), and anti-pH3 (anti-rabbit, 1:200, Millipore Sigma Cat

No: 06-570). The same secondary antibodies listed above (Alexa488, Alexa568, and Alexa633)

were used at 1:500.

For each genotype, a Fisher's exact test was used to determine if there were more

mitotically active GSCs in the *Wolbachia* infected germariums than in the uninfected

germariums.

Results

22 mutants were generated, ten with *bam* female fertility defects

22 novel mutants were generated as *bam* transgenes with alanine replacements and a

Venus fluorescent tag. They were distributed across the five functional domains/regions of the

bam gene (Fig 1a). Each mutant was assayed for fertility defects and results were compared to a

control transgenic line of a wildtype *bam* gene tagged with Venus. Mutants were characterized

by comparing the average number of progeny produced by females with and without *Wolbachia*

infection.

Of the 22 *bam* alanine-scan mutants generated, ten had fertility defects in uninfected

females (Fig 2, Supplemental fig S1, Supplemental table S2). These included all three of the

mutants that targeted known binding residues in the CAF40 binding motif of *bam*, as well as

mutants that targeted residues in the Bgcn, Csn4, and ubiquitin binding regions. There were four

null-like fertility mutants that produced no progeny when uninfected (transgenic Venus-tagged

bam^{L17A}, *bam*$^{H180A, D181A, H182A}$, *bam*$^{K225A, E227A, R228A}$, and *bam*$^{E320A, H321A, R322A}$). There were six

hypomorph-like fertility mutants that produced less progeny on average than the control when

uninfected, which included three that exhibited severe fertility defects (transgenic Venus-tagged

bam^{L28A}, *bam*$^{E152A, K153A, H155A}$, and *bam*$^{R234A, R235A}$) and three that exhibited mild fertility defects

(transgenic Venus-tagged *bam*^{M24A}, *bam*$^{D249A, E250A}$, and *bam*$^{E336A, R337A}$). The mean effect size

between the number of progeny of the mutants with fertility defects compared to the control

ranged from a minimum of -28.2 to a maximum of -74.3 (Fig 2, Supplemental fig S1).

That these new *bam* mutations that produce fertility defects are distributed throughout the

gene demonstrates that multiple regions of the gene and protein are critical for *bam* function.

Additionally, the variation in severity of fertility defects for mutations within one binding or

functional region shows that these phenotypes are allele specific.

	control	E25A, E26A, H27A	H180A, D181A, H182A	K225A, E227A, R228A	E417A, E418A, E419A	control	K127A, E128A	R133A, K134A, K136A	E320A, H321A, R322A	E336A, R337A	control	E165A, H166A,	E246A, K247A	K270A, K272A	R300A, R301A
mean	70.270	100.231	0	0	87.917	79.267	102.000	63.875	0	37.880	81.095	81.369	73.250	85.421	73.273
p-value		0	0	0	0.01		0.03	0.24	0	0.001		0.96	0.27	0.38	0.27
effect size		30	-70.3	-70.3	17.6		22.7	-15.4	-79.3	-41.4		0.27	-7.85	4.33	-7.82

	control	D13A, D14A	D18A, H19A, K22A	R234A, R235A	D249A, E250A	D263, K266	control	L17A	M24A	L28A	K108A, K109A, R111A	E152A, K153A, H155A
mean	92.538	72.214	71.913	2.706	38.690	75.132	74.280	0	36.893	3.200	59.933	3.667
p-value		0.26	0.18	0	0	0.07		0	0	0	0.06	0
effect size		5.35	5.05	-64.2	-28.2	8.27		-74.3	-37.4	-71.1	-14.3	-70.6

Figure 2. Fertility of transgenic mutants compared to the respective *bam::Venus* transgenic control. Data is colored according to the documented functional region that the mutation is located in. Below the jitter plots are the mean, two-sided permutation t-test p-value, and the average effect size between the mutant and the respective control. Gray panels indicate mutants with fertility defects (negative average effect size and p<0.05).

We also assayed the ovarian cytology of the mutants with fertility defects. Our transgenic

bam::Venus control had no tumorous germaria and proper Bam expression (Fig 3a-b). In parallel

with the novel mutants we generated, we recreated the *bam^bw^* hypomorphic mutation as a

transgenic *bam^L255F^::Venus* mutant. As has been previously characterized for this L255F *bam*

mutation in a *bam* null background, we see a tumorous germarium with some cells containing a

spectrosome (Fig 3c) [8, 10, 21]. We also found that the overproliferating GSC-like cells in this

hypomorph expressed Bam, and the lack of Bam expression in the anterior most GSCs suggested

that there was no ectopic expression of Bam or loss of GSCs.

The ten new *bam* fertility mutants displayed cytological tumorous phenotypes like that

observed in the *bam^L255F^::Venus* mutant, with the severities largely consistent with the severity

of their fertility defects (Fig 3d-f, Supplemental fig S2). When uninfected, the null-like fertility

mutants consistently had more severe tumorous germariums and severe tumorous stage one egg

chambers, like that in Fig 3d and 3e. The hypomorph-like fertility mutants had more mild

tumorous defects, like that in Fig 3f, and more wildtype egg chambers. Of note, the

bam^M24A^::Venus mutant, which had a mild fertility defect, had no tumorous germariums

(Supplemental fig S2e). The fertility counts of females of this line are consistent with the

cytological phenotype, as all but three flies had upwards of 20 progeny over the course of the

fertility assay. All mutants, including the null-like fertility mutants, stained successfully with

anti-GFP, indicating the presence of Bam::Venus protein.

Figure 3. Wildtype and transgenic *D. melanogaster* ovarioles stained with anti-Hts-1B1 (cyan), anti-Vasa (magenta), and anti-GFP (yellow). (a) CantonS wildtype control. (b) Transgenic *bam::Venus* control. (c) Transgenic *bam^{L255F}::Venus* mutant. (d-f) Representative images of cytological defects observed in the transgenic *bam* mutants with fertility defects, with arrowheads indicating spectrosomes, arrows indicating branched fusomes, and asterisks indicating the apical tip of the germarium where GSCs are identifiable as cells not expressing Bam::Venus. (d) Mutant E152A, K153A, H155A. (e) Mutant K255A, E227A, R228A. (f) Mutant E336A, R337A. Scale bar for all images is 20μM.

The twelve *bam* mutants that did not have fertility defects when uninfected were also

distributed across the *bam* gene (Fig 2, Supplemental fig S1, Supplemental table S2). These were

the transgenic Venus tagged *bam*$^{D13A, D14A}$, *bam*$^{D18A, D19A, K22A}$, *bam*$^{E25A, E26A, H27A}$, *bam*$^{K108A, K109A,}$

^{R111A}, *bam*$^{K127A, E128A}$, *bam*$^{R133A, K134A, K136A}$, *bam*$^{E165A, H166A}$, *bam*$^{E246A, K247A}$, *bam*$^{D263A, K266A}$,

bam$^{K270A, K272A}$, *bam*$^{R300A, R301A}$, and *bam*$^{E417A, E418A, E419A}$. Three of these mutants (*bam*$^{E25A, E26A,}$

^{H27A}, *bam*$^{E417A, E418A, E419A}$, and *bam*$^{K127A, E128A}$) had greater fertility on average than their

respective control ($p < 0.05$). Mutants without fertility defects include those with replacements at

highly conserved amino acid sites and replacements at poorly conserved sites (Supplemental file

S1). These results identify residues that are not critical to *bam*'s function in differentiation of

GSC daughter cells in *D. melanogaster* oogenesis, at least in laboratory conditions.

***Wolbachia* rescues all of the female hypomorph-like fertility mutants**

To determine if *Wolbachia* rescues the new *bam* mutants with fertility defects, we

compared fertility of uninfected and infected transgenic mutants. The transgenic *bam*::Venus

control showed a slight increase in fertility with *Wolbachia* infection across each independent

fertility assay ($p < 0.05$, Fig 4, Supplemental table S2). Thus, for a mutant to be classified with

Wolbachia rescue of fertility, the increase in mean effect size had to be greater than that

observed in the control.

Of the four null-like fertility mutants that produced no progeny when uninfected, three

produced some progeny when infected with *Wolbachia* (transgenic Venus-tagged *bam*^{L17A},

bam$^{H180A, D181A, H182A}$, and *bam*$^{E320A, H321A, R322A}$) ($p < 0.05$); the fourth null-like mutant (transgenic

Venus-tagged *bam*$^{E320A, H32A, R322A}$) did not produce progeny regardless of infection status (Fig 4,

Supplemental table S2). The slight production of progeny in the three null-like fertility mutants

with *Wolbachia* is consistent with *Wolbachia* infection increasing fertility of transgenic flies but

cannot be classified as *Wolbachia* rescue here because of the small effect size.

All six hypomorph-like mutants that had reduced, but not zero, fertility compared to the

control when uninfected and had an increase in average fertility with *Wolbachia* infection, with a

mean effect size greater than that of the control (p<0.05, Fig 4, Supplemental table S2). The three

mutants with severe fertility defects (transgenic Venus-tagged *bam*^{L28A}, *bam*$^{E152A, K153A, H155A}$, and

bam$^{R234A, R235A}$) saw an approximate 20-fold increase in the average number of progeny per

female when infected with *Wolbachia*. The three mutants with mild fertility defects (transgenic

Venus-tagged *bam*^{M24A}, *bam*$^{D249A, E250A}$, and *bam*$^{E336A, R337A}$) saw an approximate 2.5-fold increase

in the average number of progeny per female when infected with *Wolbachia,* bringing them to a

fertility level similar to that of the infected control.

Figure 4. Fertility of transgenic control and transgenic *bam* mutants with fertility defects as uninfected and infected with *Wolbachia*. Two-sided permutation t-test p-values and the average effect size between the uninfected (blue) and infected with *Wolbachia* (red) data are shown below each data set. Mutant E320A, H321A, R322A was completely sterile with and without *Wolbachia* and thus is not shown. Gray panels highlight the mutants that are classified as having *Wolbachia* rescue due having an average effect size greater than that observed in the respective control and p<0.05.

Overall, we find that *Wolbachia* rescues the reduced fertility of the hypomorph-like

fertility mutants across all the documented binding regions of *bam* and throughout the predicted

3D protein structure (Fig 5) [43, 44]. These results show that the *Wolbachia* rescue is not specific

to the L255F *bam* mutation and that *Wolbachia* interacts with *bam* in a manner that is not

specific to the known binding regions of the gene.

Figure 5. The locations of the mutations generated and their effect on fertility and
***Wolbachia* rescue across the documented binding and functional regions of *bam*.** Shown in
(a) a linear schematic of the gene and (b) the predicted protein structure.

We plotted the mean progeny for all transgenic females (both uninfected and *Wolbachia*

infected) to easily compare the above fertility results across all mutants (Fig 6). When comparing

the mean fertility for each mutant between the uninfected and *Wolbachia* infected lines, we find

that the data falls roughly along a logarithmic curve, suggesting there is a relationship between

the severity of uninfected mutants and the increase in fertility seen with *Wolbachia* infection.

However, we cannot discount the possibility of the increase in fertility being due to an allele-

specific interaction with *Wolbachia*.

378
Figure 6. Mean fertility comparison of all transgenic lines generated with and without
Wolbachia. The control lines are transgenic *bam::Venus* (black) and the transgenic mutants are
labeled by the first residue that was changed to alanine (orange for mutants with fertility defects
with *Wolbachia* rescue, red for mutants with fertility defects without *Wolbachia* rescue, and blue
for mutants without fertility defects). Figure shows data across five independent fertility assays
with five different controls.
385

GSC mitosis in a *bam* hypomorphic mutant is not affected by *Wolbachia*

To determine if the increase in fertility of *bam* mutants is due to an increase in

asymmetric division by GSCs, we quantified GSC mitosis in the *bam::Venus* control and the

bam^{L255F}::Venus mutant. The *bam^{L255F}* transgenic hypomorph recapitulates the severity of the

endogenous *bam^{bw}* hypomorph, though shows slightly reduced rescue compared to the *bam^{bw}*

hypomorph (Supplemental fig S3). This difference is likely due to the differences in the genetic

backgrounds of these two otherwise identical mutations.

Cells in the apical tip of the germarium that expressed the pH3 mitosis marker but not the

Venus-tagged Bam protein were identified as GSCs actively undergoing mitosis (Supplemental

fig S4). The rate of GSC mitosis was not different between *Wolbachia* infected and uninfected

flies within both the transgenic control and the transgenic hypomorph (p=1.0) (Table 1). Overall,

the *bam^{L255F}::Venus* transgenic hypomorph had fewer GSC mitotic events compared to the

transgenic control, but it was not statistically significant (p=0.259 with *Wolbachia* and p=0.448

without *Wolbachia*).

	Transgenic control		Transgenic *bam* hypomorph	
	uninfected	*Wolbachia* infected	uninfected	*Wolbachia* infected
# of germariums with pH3 positive GSCs	5	9	2	4
# of germariums without pH3 positive GSCs	111	173	105	173

401 **Table 1. Number of mitotically active (pH3 positive) and quiescent GSCs**. GSC counts are
402 shown for the transgenic control (*bam::Venus*) and a transgenic hypomorph (*bam^{L255F}::Venus*)
403 when uninfected and infected with *Wolbachia.*

Wolbachia does not rescue a *bam* knockdown phenotype

To determine if *Wolbachia* can rescue the phenotype of a reduced expression, but functional Bam protein in females, we tested for the rescue of a *bam* RNAi knockdown mutant by performing a reciprocal cross using the UAS-Gal4 system. We assessed both the number of egg chambers that have a nurse cell present (nurse-cell-positive egg chambers) per ovary in wildtype and *bam* knockdown mutants, as in [11, 18], as well as the fertility of the knockdown mutants.

We find that the *bam* knockdown mutants have fewer nurse-cell-positive egg chambers per ovary than the wildtype lines regardless of *Wolbachia* status ($p<0.05$) (Fig 8a, Supplemental fig S5). *Bam* knockdown mutants with *Wolbachia* had fewer nurse-cell-positive egg chambers on average than *bam* knockdown mutants without *Wolbachia* ($p<0.05$) but had similar levels of fertility over a period of seven days ($p = 0.69$, Fig 8b). This difference is likely due to the narrow window of time captured in the nurse-cell-positive assay compared to the fertility assay. These results show that *Wolbachia* does not rescue the tumorous mutant phenotype or the fertility of a *bam* knockdown mutant.

Additionally, there is no increase in *bam* expression in the knockdown mutant with *Wolbachia* compared to the uninfected knockdown mutant (Fig 8c). This shows that *Wolbachia* does not increase the expression levels of *bam,* consistent with the lack of rescue that we observe. All together, these results suggest that *Wolbachia* rescue is specific to partial-loss-of-function protein coding *bam* mutants.

Figure 7. *Bam* knockdown down by RNAi in the UAS-Gal4 system. *Bam* wildtype 1 and 2
are the UAS and Gal4 stocks, respectively; *bam* knockdowns are the nos-Gal4/+; UAS-
bam[HMS00029]/+ reciprocal F1 progeny of the stocks. (a) The number of nurse-cell-positive egg
chambers per ovary. *** = significant at p<0.001; NS = not significant. (b) Fertility of the *bam*
knockdown mutants as uninfected and *Wolbachia* infected. Jitter plots (left) show the raw data as
dots with horizontal lines representing the mean values of each dataset. The mean difference
between uninfected and infected lines are plotted as a bootstrap sampling distribution with a 95%
confidence interval (right). (c) *Bam* expression on a log2 scale relative to *bam* wildtype 1. Each
bar represents the dataset of four samples consisting of three female flies, and shows the mean
value with standard error.

Discussion

Bam is one of a few genes that has been shown to genetically interact with *Wolbachia* in

D. melanogaster. In female and male adults, *bam*'s molecular role is to bring together other

proteins into complexes, which then carry out functions vital to the maintenance or progression

of gametogenesis. *Bam*'s central and extensive role in early reproduction motivates a question

about the specificity of the *bam Wolbachia* interaction.

We targeted *bam*'s role of protein binding and minimized affecting the overall

conformation of the Bam protein by replacing charged residues with alanine throughout *bam*'s

documented binding regions [33]. We find no specificity to the presence/absence of rescue or the

magnitude of fertility rescue for the different functional regions of the linear *bam* sequence or

with the AlphaFold-predicted Bam protein structure. The strongest evidence for this comes from

our finding that *Wolbachia* rescues fertility defects of the known CAF40 binding residues

(bam^{M24A} and bam^{L28A}), as well as fertility defects of several mutations not involved in CAF40

binding. This suggests that *Wolbachia* does not rescue the specific interaction between Bam and

a single one of its known protein partners in *bam* fertility mutants.

While it is possible that the rescued mutants' amino acid sites are involved in a direct

interaction between Bam and an unmapped binding partner (e.g., Otu [22]), we tested two

alternative hypotheses for how *Wolbachia* could be rescuing *bam* fertility mutants. The first

proposes that *Wolbachia* induces a fertility rescue by affecting *bam*'s expression. It has

previously been shown that there is no difference in *bam* expression in the bam^{bw} hypomorph as

uninfected and infected with *Wolbachia* [11]. Here, we tested the ability of *Wolbachia* to rescue

the mutant phenotype of a wildtype *bam* gene that is reduced in expression via RNAi.

Importantly, we have a knockdown mutant that displays the hypomorph phenotype of tumorous

germariums with some wildtype egg chambers. If the knockdown phenotype is too severe,

Wolbachia may have no effect or only a minimal effect as evidenced by the null-like fertility

mutants generated in this study (transgenic Venus-tagged *bam*^{L17A}, *bam*$^{H180A, D181A, H182A}$,

bam$^{K225A, E227A, R228A}$, and *bam*$^{E320A, H321A, R322A}$) and *Wolbachia*'s lack of interaction with the *bam*

null mutant documented previously [11]. With the hypomorph-like-phenotype *bam* knockdown

mutant (*nos*-Gal4/+; UAS-*bam*^{HMS00029}/+), we find that *Wolbachia* does not rescue the number of

nurse-cell-positive egg chambers or affect the fertility of female flies; it also does not increase

the expression level of *bam.*

The second hypothesis proposes that *Wolbachia* increases the rate of GSC mitosis, which

increases the output of reproduction. Previous studies have shown *Wolbachia* to interact with

GSCs, including research in *D. mauritiana,* where the increased fertility of *Wolbachia* infected

females is attributed to the increased rate of GSC mitosis [45]. Other evidence of *Wolbachia*'s

interaction with GSCs includes research on the *Sxl*^{f4} *D. melanogaster* mutant that has shown that

Wolbachia restores the number of GSCs that were lost due to ectopic *bam* expression [19] and

the necessity for *Wolbachia* for GSC maintenance in filarial nematodes [46].

We find that *Wolbachia* does not increase the rate of GSC mitosis in the *bam*^{L255F}*::Venus*

hypomorph, and we do not observe any obvious ectopic Bam expression in the GSC niche in any

of our transgenic *bam* mutants. These results are consistent with what we see with the *bam*

knockdown results, as an increase in GSC mitosis rate would likely be reflected in an increase in

progeny in the *Wolbachia* infected *bam* knockdown females.

Thus, while it is possible for there to be a subtle *Wolbachia*-GSC interaction that we have

not yet assessed, collectively, our results suggest that a large component of the *Wolbachia bam*

interaction is specific to mutant phenotypes that are the result of a disruption of *bam* function.

Biochemical assays are needed to firmly conclude that the *Wolbachia bam* interaction is specific

to *bam* functions, but it is notable that this proposal is akin to that suggested for *Wolbachia's*

interaction with *Sxl* [47]. The observations that *Wolbachia* cannot rescue the reduced egg count

of a *snf* mutant, which has no *Sxl* expression in the germline, and a *Sxl^{f5,fHa}* mutant, which has a

transposon insertion in the second intron [17] parallels the *bam* knockdown results of our study.

It is possible that the fertility rescue of *bam* mutants occurs through the same (unknown)

mechanism by which the rescue of *Sxl* mutants occurs. This would be a process beyond the

documented restoration of GSC number in *Sxl^{f4}* mutants by *Wolbachia* factor TomO, as it was

found that TomO is not sufficient to restore fertility of the *Sxl^{f4}* mutants [19].

Bam and Sxl have been proposed to partner together for two distinct functions. Together,

with Bgcn, Mei-P26, Brat, Ago1, and a deadenylase complex, they repress the translation of the

stem cell maintenance factor *nos* to determine GSC daughter cell fate [24, 25]. In this sense,

Wolbachia could increase stability of host protein complexes, through, for example, use of

molecular chaperones to help fold mutant Bam or Sxl proteins. It has also been suggested that

Bam and Sxl function together to maintain female sexual identity in the adult germline [48], as

failure to regulate *nos* does not cause the tumorous phenotype observed in *Sxl* and *bam* mutants

[24]. Here, *Wolbachia* could restore proper female sexual identity to the germline by repressing

testes specific gene expression.

Alternatively, it is possible that *Wolbachia* interacts with *bam* and *Sxl* independently.

Bam has a number of other genetic and physical interacting partners that offer functional

pathways for which *Wolbachia* could be interacting with. For example, Bam promotes

differentiation of GSC daughter cells by facilitating deubiquitination of CycA with Otu [22].

Most recently, Bam itself has been shown to interact with E3 ligase Cullin-2, which ubiquitinates

Bam and leads to Bam's controlled expression [49]. Critical ubiquitin binding residues in Bam

have been identified which offer another opportunity to assess the extent of *Wolbachia*'s

interaction.

While more research is needed to determine if the *Wolbachia* interaction is specific to

Bam-Sxl complexes or if *Wolbachia* interacts with *bam* independently of *Sxl*, our study suggests

that *Wolbachia* could play a role by stabilizing the complexes or restoring binding affinity

between Bam and its partners to restore *bam* function in *bam* protein-coding mutants.

515   REFERENCES

516   1. McKearin DM, Spradling AC. *Bag-of-marbles*: a *Drosophila* gene required to initiate both

517   male and female gametogenesis. Genes Dev. 1990;4: 2242-2251.

518   2. Gönczy P, Matunis E, DiNardo S. *bag-of-marbles* and *benign genial cell neoplasm* act in the

519   germline to restrict proliferation during *Drosophila* spermatogenesis.

520   Development 1997;124(21): 4361-4371.

521   3. Insco ML, Leon A, Tam CH, McKearin DM, Fuller MT. Accumulation of a differentiation

522   regulator specifies transit amplifying division number in an adult stem cell lineage. Proc

523   Natl Acad Sci. 2009;106(52): 22311-22316.

524   4. Insco ML, Bailey AS, Kim J, Olivares GH, Wapinski OL, Tam CH, et al. A self-limiting

525   switch based on translational control regulates the transition from proliferation to

526   differentiation in an adult stem cell lineage. Cell Stem Cell 2012;11: 689-700.

527   5. Chen D, McKearin D. Dpp signaling silences *bam* transcription directly to establish

528   asymmetric divisions of germline stem cells. Curr Biol. 2003;13(20): 1786–1791.

529   6. Chen D, McKearin DM. A discrete transcriptional silencer in the bam gene determines

530   asymmetric division of the *Drosophila* germline stem cell. Development 2003;130:

531   1159–1170.

532   7. Song X, Wong MD, Kawase E, Xi R, Ding BC, McCarthy JJ, et al. Bmp signals from niche

533   cells directly repress transcription of a differentiation-promoting gene, *bag of marbles*, in

534   germline stem cells in the *Drosophila* ovary. Development 2004;131: 1353-1364.

535   8. Shen R, Weng C, Yu J, Xie T. eIF4A controls germline stem cell self-renewal by directly

536   inhibiting BAM function in the *Drosophila* ovary. Proc Natl Acad Sci USA 2009;106:

537   11623–11628.

9. McKearin D and Ohlstein B. A role for the *Drosophila* Bag-of-marbles protein in the

differentiation of cystoblasts from germline stem cells. Development 1995;121: 2937–

2947.

10. Ohlstein B, Lavoie CA, Vef O, Gateff E, Mckearin DM. The *Drosophila* cystoblast

differentiation factor, *benign gonial cell neoplasm*, is related to DExH-box proteins and

interacts genetically with *bag-of-marbles*. Genetics 2000;155: 1809-1819.

11. Flores HA, Bubnell JE, Aquadro CF, Barbash DA. The *Drosophila bag of marbles* gene

interacts genetically with *Wolbachia* and shows female-specific effects of divergence.

PLoS Genet. 2015;11: e1005453.

12. Hilgenboecker K, Hammerstein P, Schlattmann P, Telschow A, Werren JH. How many

species are infected with *Wolbachia*? — a statistical analysis of current data. FEMS

Microbiol Lett. 2008;281: 215-220.

13. Weinert LA, Araujo-Jnr EV, Ahmed MZ, Welch JJ. The incidence of bacterial

endosymbionts in terrestrial arthropods. Proc R Soc B. 2015;282(1807): 20150249.

14. Taylor MJ, Bandi C, and Hoerauf A. *Wolbachia* bacterial endosymbionts of filarial

nematodes. Adv Parasitol. 2005;60: 245-284.

15. Werren JH, Baldo L, Clark ME. *Wolbachia*: master manipulators of invertebrate biology. Nat

Rev Microbiol. 2008;6: 741-751.

16. Kaur R, Shropshire JK, Cross KL, Leigh B, Mansueto AJ, Steward V, et al. Living in the

endosymbiotic world of *Wolbachia*: A centennial review. Cell Host Microbe 2021;29(6):

879-93.

17. Starr DJ, Cline TW. A host-parasite interaction rescues *Drosophila* oogenesis defects. Nature

2002;418: 76-79.

18. Bubnell JE, Fernandez-Begne P, Ulbing CKS, Aquadro CF. Diverse *w*Mel variants of

Wolbachia pipientis differentially rescue fertility and cytological defects of the *bag of*

marbles partial loss of function mutation in *Drosophila melanogaster.* G3 (Bethesda)

2021;11(12).

19. Ote M, Ueyama M, Yamamoto D. *Wolbachia* protein TomO targets *nanos* mRNA and

restores germ stem cells in *Drosophila sex-lethal* mutants. Curr Biol. 2016;26: 2223-

2232.

20. Sgromo A, Raisch T, Bawankar P, Bhandari D, Chen Y, Kuzuoğlu-Öztürk D, et al. A

CAF40-binding motif facilitates recruitment of the CCR4-NOT complex to mRNAs

targeted by Drosophila Roquin. Nat Commun. 2017;8: 14307.

21. Pan L, Wang S, Lu T, Weng C, Song X, Park JK, et al. Protein competition switches the

function of COP9 from self-renewal to differentiation. Nature 2014;514: 233-236.

22. Ji S, Li C, Hu L, Liu K, Mei J, Luo Y, et al. Bam-dependent deubiquitinase complex can

disrupt germ-line stem cell maintenance by targeting cyclin A. Proc Natl Acad Sci USA

2017;114: 6316-6321.

23. Li Y, Minor NT, Park JK, McKearin DM, Maines JZ. Bam and Bgcn antagonize Nanos-

dependent germ-line stem cell maintenance. Proc Natl Acad Sci USA. 2009;106(23):

9304–9309.

24. Chau J, Kulnane LS, Salz HK. *Sex-lethal* enables germline stem cell differentiation by down-

regulating Nanos protein levels during *Drosophila* oogenesis. Proc Natl Acad Sci USA.

2012;109: 9465–9470.

25. Malik S, Jang W, Kim JY, Kim C. Mechanisms ensuring robust repression of the *Drosophila*

female germline stem cell maintenance factor Nanos via posttranscriptional regulation.

The FASEB Journal 2020;34: 11421-11430.

26. Fu Z, Geng C, Wang H, Yang Z, Wenig C, Li H, et al. Twin promotes the maintenance and

differentiation of germline stem cell lineage through modulation of multiple pathways.

Cell Rep. 2015;13: 1366-1379.

27. Chen D, Wang Q, Huang H, Xia L, Jiang X, Kan L, et al. Effete-mediated degradation of

Cyclin A is essential for the maintenance of germline stem cells in *Drosophila*.

Development 2009;136: 4133-4142.

28. Civetta A, Rajakumar SA, Brouwers B, Bacik JP. Rapid evolution and gene-specific patterns

of selection for three genes of spermatogenesis in *Drosophila*. Mol Biol Evol. 2006;23:

655-662.

29. Bauer DuMont VL, Flores HA, Wright MH, Aquadro CF. Recurrent positive selection at

bgcn, a key determinant of germ line differentiation, does not appear to be driven by

simple coevolution with its partner protein *bam*. Mol Biol Evol. 2007;24: 182–191.

30. Bubnell JE, Ulbing CKS, Fernandez Begne P, Aquadro CF. Functional divergence of the

bag-of-marbles gene in the *Drosophila melanogaster* species group. Mol Biol Evol.

2022;39(7).

31. Flores HA, Bauer DuMont VL, Fatoo A, Hubbard D, Hijji M, Barbash DA, et al. Adaptive

evolution of genes involved in germline stem cells in *Drosophila melanogaster* and *D.*

simulans. G3 (Bethesda) 2015;5: 583-592.

32. Bauer DuMont VL, White SL, Zinshteyn D, Aquadro CF. Molecular population genetics of

Sex-lethal (*Sxl*) in the *Drosophila melanogaster* species group: a locus that genetically

interacts with *Wolbachia pipientis* in *Drosophila melanogaster.* G3 (Bethesda)

2021;11(8).

33. Cunningham BC, Wells JA. High-resolution epitope mapping of hGH-receptor interactions

by alanine-scanning mutagenesis. *Science* 1989;244: 1081-1085

34 Nakamura Y. Kisarazu (Japan): Kazusa DNA Research Institute. Codon Usage Database.

[last accessed 2021 Oct 12].

https://www.kazusa.or.jp/codon/cgi-bin/showcodon.cgi?species=7227

35. Groth AC, Fish M, Nusse R, Calos MP. Construction of transgenic *Drosophila* by using the

site-specific integrase from phage phiC31. Genetics 2004;166: 1775-1782.

36. Bubnell JE. Functional evolution of the *bag of marbles* gene in *Drosophila.* PhD thesis,

Cornell University. 2020. Available from https://hdl.handle.net/1813/103228

37. New England Biolabs. 2021. NEBaseChanger. [last accessed 2021 Oct 13].

https://nebasechanger.neb.com/

38. Ho J, Tumkaya T, Aryal S, Choi H, Claridge-Chang A. Moving beyond P values: data

analysis with estimation graphics. Nat Methods. 2019;16: 565-566.

39. Claridge-Chang A, Ho J. Estimation Statistics: Analyze your data with effect Sizes. C2017-

2021. [accessed 2022 Sept 20]. https://www.estimationstats.com/#/

40. Shindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an

open-source platform for biological-image analysis. Nature Methods 2012;9(7): 676-682.

41. Chrostek E, Marialva MSP, Esteves SS, Weinert LA, Martinez J, Jiggins FM, et al.

Wolbachia variants induce differential protection to viruses in *Drosophila melanogaster*:

a phenotypic and phylogenomic analysis. PLoS Genet. 2013;9: e1003896.

42. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-

PCR. Nucleic Acids Res. 2001;29: e45.

43. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate

protein structure prediction with AlphaFold. Nature 2021;596: 583-589.

45. Fast EM, Toomey ME, Panaram K, Desjardins D, Kolaczyk ED, Frydman HM. *Wolbachia*

enhance *Drosophila* stem cell proliferation and target the germline stem cell niche.

Science 2011;334: 990-992.

46. Foray V, Pérez-Jiménez MM, Fattouh N, Landmann F. *Wolbachia* control stem cell behavior

and stimulate germline proliferation in filarial nematodes. Dev Cell. 2018;45: 198-

211.e3.

47. Ote M, Yamamoto D. Impact of *Wolbachia* infection on *Drosophila* female germline stem

cells. Curr Opin Insect Sci. 2020;27: 8-15.

48. Shapiro-Kulnane L, Smolko AE, Salz HK. Maintenance of *Drosophila* germline stem cell

sexual identity in oogenesis and tumorigenesis. Development 2015;142(6): 1073-1082.

49. Cai Q, Yan J, Duan R, Zhu Y, Hua Y, Liao Y, et al. E3 ligase Cul2 mediates *Drosophila*

early germ cell differentiation through targeting Bam. Dev Biol. 2023;493: 103-108.

44. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold

Protein Structure Database: massively expanding the structural coverage of protein-

sequence space with high-accuracy models. Nucleic Acids Res. 2021;50: D439-D444.

Supplemental figure S1. Full estimation plots for fertility data of transgenic mutants compared to the *bam::Venus* transgenic control. Fertility assays were done in five independent rounds. Below the jitter plots are the resampled bootstrap sampling distributions, with the mean differences represented by the black dots and the 95% confidence intervals represented by the vertical lines.

Supplemental figure S2. Germariums of controls and all transgenic *bam* mutants with fertility defects when uninfected with *Wolbachia*. Stained with anti-Hts-1B1 (cyan), anti-Vasa (magenta), and anti-GFP (yellow). (a) CantonS, (b) *bam::Venus* control, (c) *bam^{L255F}::Venus*, (d) *bam^{L17A}::Venus*, (e) *bam^{M24A}::Venus*, (f) *bam^{L28A}::Venus*, (g) *bam^{E152A, K153A, H155A}::Venus*, (h) *bam^{H180A, D181A, H182A:}::Venus*, (i) *bam^{R234A, R235A}::Venus*, (j) *bam^{D249A, E250A}::Venus*, (k) *bam^{K255A, E227A, R228A}::Venus*, (l) *bam^{E320A, H321A, R322A}::Venus*, (m) *bam^{E336A, R337A}::Venus*. Scale bar is 20μM for all images.

665 **Supplemental figure S3. Comparison of original L255F mutant (bam^{bw}) and the transgenic**
666 **L255F mutant with a Venus tag ($bam^{L255F}::Venus$) by the number of nurse-cell-positive egg**
667 **chambers per ovary.** ** = significant at p<0.01; *** = significant at p<0.001; NS = not
668 significant.

Supplemental figure S4. GSC mitosis in the transgenic *bam::Venus* control and transgenic *bam^{L255F}::Venus* hypomorph. (a) No GSC mitosis and (b) active GSC mitosis in the *bam::Venus* control germarium. (c) No GSC mitosis and (d) active GSC mitosis in the *bam^{L255F}::Venus* hypomorphic mutant.

Supplemental figure S5. Ovaries of a *bam* wildtype and *bam* RNAi knockdown fly. (a)
Ovaries of a *bam* wildtype fly. All egg chambers are results of successful Bam function. (b)
Mutant ovaries of a nos-Gal4/+; UAS-*bam*[HMS00029]/+ uninfected fly. Arrows indicate nurse-cell-
positive egg chambers that represent successful Bam function. Scale bar for both images is
100μM.

Bam mutations	Mutagenesis forward primer	Mutagenesis reverse primer	Annealing temp. (°C)
D13A, D14A,	TGAGGGCAACgccgccCAGCAGTTGG	GGACACACGTCACGTGCATTAAG	70
L17A	CGACCAGCAGgctGACCACAATTTTAAGCAG	TCGTTGCCCTCAGGACAC	65
D18A, D19A, K22A,	ttttgccCAGATGGAGGAGCATTTGG	ttagcggcCAACTGCTGGTCGTCGTT	64
M24A	TTTTAAGCAGgctGAGGAGCATTTGGCC	TTGTGGTCCAACTGCTGG	62
E25A, E26A, H27A,	cgcTTTGGCCTTAATGGTGGAAG	gcggCCATCTGCTTAAAATTGTGG	60
L28A	GGAGGAGCATgctGCCTTAATGG	ATCTGCTTAAAATTGTGGTC	58
K108A, K109A, R111A,	tcggcaTCCGGAGGCGTGTTGGTC	ggcggcGGCCGGAGAACCGTCGAAT	71
K127A, E128A	GCAACTGCAGgccgctAATGTGTGGAACCGGAAGAGTAAAG	TTCTGCTTTGGCCCGGTG	66
R133A, K134A, K136A,	agtgctGGCTCTGCGTCCGCGGAT	ggctgcGTTCCACACATTTTCCTTCTGCAGTTGC	71
E152A, K153A, H155A	ctggcaATGATTGGTCTGCACGGC	agcggcAATAGTTATGGGCAGTTTCTCAATATTATC	65
E165A, H166A,	CAATAGCTTAgccgctAACGCCGTGC	TTTAGATAAGGAGCTTTTTATTATG	58
H180A, D181A, H182A,	cgctCTGACCGCCGATTTGGGC	gcagcCAGGGATCTGAACAGATTCATCAAAC	66
K225A, E227A, R228A	gccgctTTCCTTGTCCAGCAGCGC	tacagcGGTGCACAGCACCTGCAG	69
R234A, R235A	TGTCCAGCAGgctgctACCTTGGAGG	AGGAAGCGCTCTACTTTG	61
E246A, K247A,	CTTCGATTTCgccgctTACGACGAGTGTGACAAGTTG	TGGCGATTCGCCTCCAAG	65
D249A, E250A	gccgctTTCCTTGTCCAGCAGCGC	tacagcGGTGCACAGCACCTGCAG	61
L255F	TGACAAGTTGtTTAAGGGTTTC	CACTCGTCGTATTTCTCG	58
D263A, K266A,	ttcgctCTGCTTTTAAAGCCCAAAATG	gttggcCAAATAGGATGCGAAACC	59
K270A, K272A,	cgctATGCGCAATCGAAACGGA	ggggcTAAAAGCAGTTTGAAGTTGTCC	62
R300A, R301A,	AATTGGTCTGgctgccTGGATCAAGGCTGCGCATC	AGCAATCTCTCCATGCGC	62
E320A, H321A, R322A	tgctTACTCCGGGGCCATGACC	gcggcCAGATCCATTTCCCAGTTAAATACGTG	67
E336A, R337A	GTCGTTGAACgccgcaGCCATCCTTTTGTCC	TTGTGGCTTTCGGTCATG	59
E417A, E418A, E419A.	cgccTTTGAGGAGACCGAGGAAGTGC	gcggcCGACGCTGACGGCTGCTC	70

681   **Supplemental table S1. Mutagenesis primers used to generate new *bam* mutants.** Each row

682   lists the corresponding primer sequences and annealing temperature used in the mutagenesis

683   PCR step of the cloning procedure to generate the respective amino acid mutant.

Transgenic *bam* allele	Uninfected mean fertility	Infected mean fertility	Uninfected sample size	Infected sample size	Assay round
bam$^{D13A, D14A}$*::Venus*	72.2142857	90.3571429	28	28	two
bam^{L17A}*::Venus*	0.0000000	1.9600000	43	50	five
bam$^{D18A, H19A, K22A}$*::Venus*	71.9130435	90.8421053	23	19	one
bam^{M24A}*::Venus*	36.8928571	94.0909091	28	22	five
bam$^{E25A, E26A, H27A}$*::Venus*	100.2307692	102.0645161	26	31	one
bam^{L28A}*::Venus*	3.2000000	71.0000000	50	31	five
bam$^{K108A, K109A, R111A}$*::Venus*	59.9333333	98.6800000	30	25	five
bam$^{K127A, E128A}$*::Venus*	102.0000000	119.1111111	17	9	three
bam$^{R133A, K134A}$*::Venus*	63.8750000	107.5000000	16	10	three
bam$^{E152A, K153A, H155A}$*::Venus*	3.6666667	67.3421053	45	38	five
bam$^{E165A, H166A}$*::Venus*	81.3684211	100.8000000	19	10	four
bam$^{H180A, D181A, H182A}$*::Venus*	0.0000000	0.7714286	34	35	one
bam$^{K225A, E227A, R228A}$*::Venus*	0.0000000	0.9200000	29	25	one
bam$^{R234A, R235A}$*::Venus*	2.7058824	75.4814815	34	27	two
bam$^{E246A, K247A}$*::Venus*	73.2500000	104.8888889	20	18	four
bam$^{D249A, E250A}$*::Venus*	38.6896552	88.173913	29	23	two
bam$^{D263A, K266A}$*::Venus*	75.1320755	88.1111111	53	45	two
bam$^{K270A, K272A}$*::Venus*	85.4210526	99.5384615	19	13	four
bam$^{R300A, D301A}$*::Venus*	73.2727273	102.4583333	22	24	four
bam$^{E320A, H321A, R322A}$*::Venus*	0.0000000	0.0000000	41	39	three
bam$^{E336A, R337A}$*::Venus*	37.8800000	98.9375000	25	16	three
bam$^{E417A, E418A, E419A}$*::Venus*	87.9166667	111.4347826	36	23	one
bam::Venus	70.2702703	108.9200000	37	25	one
bam::Venus	66.8666667	92.5384615	30	26	two
bam::Venus	79.2666667	116.9000000	15	10	three
bam::Venus	81.0952381	101.1875000	21	16	four
bam::Venus	74.2800000	102.6086957	25	23	five

685 **Supplemental table S2. Mean adult progeny per female seven days after first progeny**
686 **eclosed, as uninfected and infected with *Wolbachia*.** *Bam* alleles are listed in order of location
687 of mutations from N- to C-terminal, with mean fertility of and sample size of surviving females
688 at the end of the assay. The corresponding assay in which the transgenic flies were evaluated is
689 indicated by the fertility assay round.