***Wolbachia* infection at least partially rescues the fertility and ovary defects of several new**

***Drosophila melanogaster bag of marbles* protein-coding mutants**

Miwa Wenzel[1*] and Charles F. Aquadro[1*]

[1]Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY, USA

[*]Corresponding author

Email: mjw349@cornell.edu; cfa1@cornell.edu

**Abstract**

The *D. melanogaster* protein coding gene *bag of marbles* (*bam*) plays a key role in early male

and female reproduction by forming complexes with partner proteins to promote differentiation

in gametogenesis. Like another germline gene, *Sex lethal*, *bam* genetically interacts with the

endosymbiont *Wolbachia*, as *Wolbachia* rescues the reduced fertility of a *bam* hypomorphic

mutant. Here, we explored the specificity of the *bam-Wolbachia* interaction by generating 22

new *bam* mutants, with ten mutants displaying fertility defects. Nine of these mutants trend

towards rescue by the *w*Mel *Wolbachia* variant, with eight statistically significant at the fertility

and/or cytological level. In some cases, fertility was increased a striking 20-fold. There is no

specificity between the rescue and the known binding regions of *bam*, suggesting *w*Mel does not

interact with one singular *bam* partner to rescue the reproductive phenotype. We further tested if

*w*Mel interacts with *bam* in a non-specific way, by increasing *bam* transcript levels or acting

upstream in germline stem cells. A fertility assessment of a *bam* RNAi knockdown mutant

reveals that *w*Mel rescue is specific to functionally mutant *bam* alleles and we find no obvious

evidence of *w*Mel interaction with germline stem cells in *bam* mutants.

**Author Summary**

Reproduction in the *Drosophila melanogaster* fruit fly is dependent on the *bag of marbles* (*bam*)

gene, which acts early in the process of generating eggs and sperm. Mutations to this gene

negatively impact the fertility of the fly, causing it to be sterile or have fewer progeny.

Interestingly, we find that the bacteria *Wolbachia*, which resides within reproductive cells across

a wide range of insects, partially restores the fertility and ovary phenotype of several *bam*

mutants of which the resultant Bam protein is altered from wildtype. The protein function of

Bam is further suggested to be important by the lack of rescue for a fly that has a fertility defect

due to low expression of a non-mutated *bam* gene. Previous work makes similar conclusions

about *Wolbachia* with another reproductive gene, *Sex lethal* (*Sxl*), highlighting the potential for

rescue of fertility mutants to occur in a similar way across different genes. An understanding of

the ways in which *Wolbachia* can affect host reproduction provides us with context with which

to frame *Wolbachia*'s impact on host genes, such as *bam* and *Sxl,* and consider the evolutionary

implications of *Wolbachia*'s infection in *D. melanogaster* fruit flies.

## Introduction

The *bag of marbles* (*bam*) gene plays a critical role in female and male gametogenesis in *D. melanogaster.* In the stem cell niche of the ovaries and testes lies a collection of germline stem cells (GSCs) that asymmetrically divide to self-renew and produce differentiating daughter cells. In females, *bam* acts as a master regulator and its expression induces daughter cell differentiation [1]. In males, *bam* functions later in gametogenesis where it initiates terminal differentiation of the spermatogonia that eventually become sperm [1, 2, 3, 4].

In female *D. melanogaster, bam* is largely repressed in GSCs by signaling it receives from the stem cell niche [5, 6, 7]. Bam that escapes silencing in GSCs is antagonized by eukaryotic translation initiation factor 4A (eIF4A) [8]. When a GSC divides, a daughter cell is displaced from the stem cell niche, allowing *bam* to now be expressed. In the daughter cell, Bam functions to bring together various proteins that repress self-renewal and promote differentiation, leading to the establishment of a cystoblast.

In both male and female *D. melanogaster, bam* null mutants are completely sterile and have a tumorous cytological phenotype [1, 9]. A partial loss-of-function mutant (*bam*$^{bw}$) has reduced fertility in females and is sterile in males [10, 11]. In females, the fertility defect of the *bam*$^{bw}$ mutant is rescued by both *w*Mel and *w*MelCS variants of the endosymbiont bacteria *Wolbachia pipientis* [11, 12].

*W. pipientis*, commonly referred to as *Wolbachia*, is estimated to inhabit 50% or more of insect species and is an obligate bacterium in filarial nematodes [13, 14, 15]. Across both arthropods and nematodes, *Wolbachia* manipulates reproduction to promote its own maternal transmission (reviewed in [16, 17]). For example, *Wolbachia* famously induces cytoplasmic incompatibility, in which a female's progeny may have reduced viability if they do not harbor

the same *Wolbachia* strain as the male parent. *Wolbachia* is also known to induce

parthenogenesis, feminization, and male-killing of host progeny, all of which promote its

maternal transmission. The extent of these parasitic phenotypes varies across *Wolbachia* variants

and their hosts, as for example, *w*Mel does not exhibit a strong cytoplasmic incompatibility

effect in *D. melanogaster* [18, 19, 20]. Rescuing the fertility defect of the hypomorphic *bam*

mutant aligns with the evolutionary interests of *Wolbachia.*

*Wolbachia* also rescues the sterility of hypomorphs of another GSC gene, *Sex lethal* (*Sxl*),

in *D. melanogaster* (*Wolbachia* variant unspecified) [21]. It has thus been proposed that the

rescue is dependent on the point mutation induced in the *Sxl* mutant and not on the severity of the

mutant, though the researchers suggest that rescue cannot happen if *Sxl* protein levels are too low.

Of note is that the *Sxl* mutants they studied were in different genetic backgrounds, which may

have affected the results observed and the conclusions made. Such background effects were

observed with *bam* in [22], where females with the same L255F *bam* mutation had strikingly

different fertility across two different genetic backgrounds of *D. melanogaster*. Nevertheless,

further research into one of the *Sxl* mutants, *Sxl$^{f4}$*, showed that the rescue is partially due to a

restoration of GSCs by the *Wolbachia* protein toxic manipulator of oogenesis (TomO) [23].

Mutants of additional germline genes *meiotic P26* (*mei-P26*), *ovarian tumor* (*otu*), and

*sans fille* (*snf*) are not rescued by *Wolbachia,* despite having similar severity of an egg-absence

phenotype to some *Sxl* mutants (*Wolbachia* variant unspecified) [21]. These results suggest the

*Wolbachia* interaction may be specific to a functional pathway in female *D. melanogaster,*

possibly through a direct, physical interaction with a gene product like Sxl that is expressed in

the germline.

In this study, we explore the *bam-w*Mel interaction, building off the findings for the

*bam^{bw}* hypomorphic mutant. The *bam^{bw}* hypomorphic allele is a L255F change in a gene region

critical for *bam* function. Thus far, four functional regions of *bam* have been documented in *D.*

*melanogaster*: The CAF40 binding motif (residues 13-36) [23], the Benign gonial cell neoplasm

(Bgcn) binding region (residues 100-350) [24], the COP9 signalosome 4 (Csn4) binding region

(residues 151-350) [24], and a ubiquitin binding region (residues 200-250) [25]. *Bam* also has a

PEST domain (residues 402-434), which marks the protein for rapid degradation [1]. *Bam*'s

specific binding regions and their interactions with binding partners are illustrated in Figures 1a

and 1b respectively.

93
**Figure 1. *Bam* is involved in several processes that regulate the differentiation of GSCs in**
**female *D. melanogaster*.** (a) Schematic of the five known binding or functional regions of *bam*
are colored across the coding sequence exons of the gene. Each black vertical bar represents the
location of a new mutant generated with alanine replacements in this study. (b) In GSCs (left),
Bam is repressed and antagonized by eIF4A. In the differentiating daughter cells (right), Bam is
expressed and coordinates other molecules to inhibit the self-renewal pathway or promote the
differentiation pathway. Bam has been shown to genetically and physically interact with
additional molecules that are not shown.
102

To repress GSC self-renewal, Bam forms a complex with Bgcn, Meiotic-P26 (Mei-P26),

Brain tumor (Brat), a deadenylase, and either Argonaute-1 (Ago1) or Sxl to destabilize *nanos*

(*nos*) mRNA [26, 27, 28]. The destabilization of *nos* mRNA may occur through Bam's

interaction with the CAF40 which is part of the CCR4-NOT deadenylase complex [23, 29]. Thus,

in GSCs, *nos* mRNA is translated to promote GSC self-renewal; in differentiating daughter cells,

*nos* mRNA is destabilized by the Bam complex, inhibiting *nos* translation and the self-renewal

pathway.

Bam also represses the self-renewal pathway through its interaction with Csn4 [24]. In

GSCs, the COP9 complex is composed of nine Csn subunits. When Bam is expressed in

differentiating daughter cells, it breaks up the complex by sequestering Csn4, thereby inhibiting

the GSC self-renewal pathway.

Additionally, Bam promotes differentiation by forming a complex with Otu to

deubiquitinate Cyclin A (CycA) [25]. In GSCs, CycA is degraded through a process involving

the ubiquitin-conjugating enzyme Effete [30]. In differentiating GSC daughter cells, it is

proposed that Bam binds to both Otu and to proteins such as CycA, which have ubiquitin chains

that mark them for degradation. When in a complex with Bam, Otu deubiquitinates CycA,

stabilizing CycA levels and thus promoting GSC daughter cell differentiation.

Here, we have generated 22 new *bam* mutants across the different binding and functional

regions of *bam.* We analyzed the genetic interaction between these mutants and the *w*Mel variant

of *Wolbachia* to better understand the extent of the *bam-w*Mel interaction. We are partly

motivated by the observation of rapidly diversifying amino acids at *bam* across the *Drosophila*

phylogeny [31, 32, 33]. *Bam* appears novel to the *Drosophila* genus [30], and we have

previously hypothesized that *Wolbachia* could contribute to the rapid evolution observed at *bam*

due to the observed *bam-Wolbachia* genetic interaction and the fact that both *Wolbachia*

infection and positive selection at *bam* occur intermittently throughout the *Drosophila* phylogeny

[11, 32, 33, 34]. While there is no congruence between the patterns of infection and positive

selection at *bam* across species (p = 1.0, Supplemental Table S1), we cannot disregard the

potential for historic infections that may have driven positive selection at *bam.*

In this study, we find that *w*Mel genetically interacts with alanine point mutants

throughout the *bam* gene, suggesting that *w*Mel rescue of *bam* mutants is not specific to any

single binding/functional region of *bam.* We find no evidence for two hypotheses of how *w*Mel

could rescue *bam* in a non-specific manner: There is no effect of *w*Mel on a hypomorph-like *bam*

knockdown mutant, nor is there an effect of *w*Mel on the frequency of GSC mitosis that occurs

upstream of *bam*. Thus, further consideration on how *Wolbachia* genetically interacts with *bam*

is valuable for future endeavors that examine potential co-evolution of *bam* and *Wolbachia.*

**Materials and Methods**

**Design of putative *bam* hypomorph alleles**

A Venus-tagged *bam* allele was designed to facilitate easy analysis of Bam protein

expression and to serve as a control to the subsequent 22 Venus-tagged *bam* mutant alleles (Fig.

1a), all of which are described further below. The known Bam binding residues in the CAF40

binding motif were targeted for replacement [23], producing independent alleles with mutations

of L17A, M24A, and L28A. The remaining mutant alleles were designed through an alanine

scanning approach [35] that targeted the known functional and/or binding regions of *bam.*

Strings of charged residues were changed to alanine by choosing an alanine codon with similar

usage bias in *D. melanogaster* as that of the original residue [36]. Additionally, we designed an

allele to recreate the *bam*[bw] L255F mutation as a transgene with Venus tag. All new alleles are

listed in Supplemental file S1, with the original and mutated codons.

**Generation of transgenic flies**

Transgenic fly lines carrying distinct putative hypomorphic *bam::Venus* alleles and a

control *bam::Venus* transgenic fly line were generated using PhiC-mediated integration [37].

Approximately 1.5 kb upstream and 700 bp downstream regions of *bam* were cloned from

genomic DNA of a *D. melanogaster bam::Venus* line [38]. This construct is similar to that of the

*D. melanogaster bam* YFP line from [11] but contains a 36bp glycine-glycine-serine linker

sequence between the 3' end of the *bam* coding sequence and the Venus tag to allow for proper

folding of both proteins. The *bam::Venus* sequence was cloned into a pMiniT 2.0 vector, using

the NEB PCR cloning kit. Mutant alleles were subsequently made in the established *bam::Venus*

pMiniT 2.0 plasmid using the NEB Q5 Site-Directed Mutagenesis Kit, with primers designed

online using NEBase Changer (Supplemental table S2) [39]. Briefly, mutagenic primers were

used to amplify the *bam::Venus* pMiniT 2.0 plasmid using a PCR cycle of 98°C for 30 seconds;

25 cycles of 98°C for 10 seconds, 57-72°C annealing for 30 seconds (specified per primer set in

Supplemental table S2), 72°C for 3.5 minutes; and 72°C for 2 minutes. PCR products were

treated with kinase, ligase, and DpnI before being transformed into competent cells. The

resultant plasmids were subcloned into the pCasper\attB vector, a gift from Dan Barbash that

was used by [11] for PhiC transgenesis. Plasmids were verified by sequencing.

Plasmids were injected into embryos of a *w*, *nos*-int; P{CaryP}attP40 *D. melanogaster*

line by GenetiVision. $G_0$ flies were mated with a *w*[1118] isogenic line and $F_1$ progeny were

screened for successful transgenesis by selecting for those with mini-white eye color. Mutant

lines were identified by Sanger sequencing.

All transgenic flies were assessed for function in a *bam* null background. Thus, the

genotype that was assessed in all experiments was *w*; [$w^{+mc}$, *bam::Venus*]/+; $bam^{\Delta 59}/bam^{\Delta 86}$,

where *bam::Venus* represents the control *bam* allele or one of the 22 mutant alleles. For the

remainder of this paper, "transgenic females" and "transgenic *bam* mutants" refer to flies of this

genotype. *Wolbachia* status of the assessed flies (uninfected or *w*Mel-infected) was established

by maternal inheritance when crossing in the *bam* null alleles.

180
**Fertility assays of female transgenic flies**

Virgin transgenic females and virgin CantonS males were collected over 48 hours and

aged two days. At least 30 individual crosses of one female and two males were set up on yeast-

glucose food (minimum of 30, maximum of 88 crosses were set per genotype tested). All crosses

were kept in an incubator at 25°C with a 12-hour light/dark cycle. Parents were removed on the

seventh day. Progeny were counted every other day over seven days, starting on the day of first

eclosion. If the female parent died by the time of removal, data for that cross was discarded. Five

independent fertility assays were conducted to cover all 22 mutants, with a transgenic control (*w*;

[$w^{+mc}$, *bam::Venus*]/+; $bam^{\Delta 59}/bam^{\Delta 86}$) present in each assay.

The total progeny after seven days was summed and estimation statistics [40] were

performed to determine first, if a fertility defect was present for a mutant, and second, if there

was a *w*Mel rescue effect. Estimation statistics and plots were generated using the dabestr

package in R (v4.0.1). While this offers an alternative statistical test that allows us to easily

visualize effect size, we also report p-values for ease and convention. P-values were calculated

through the associated website as results of a two-sided permutation t-test [41], as they were not

provided in the R package.

**Nurse-cell-positive egg chamber assessment of female transgenic flies**

For the ten transgenic mutants that displayed a fertility defect, an assay for nurse-cell-

positive egg chambers was also performed, as has been previously done for a supplemental

assessment of *bam*'s function [11, 12]. Briefly, ovaries of 1-2 day old flies were fixed in 4%

paraformaldehyde, rinsed and washed with PBST as described above, and then mounted in

ProLong Glass with NucBlue (ThermoFisher). Ovaries were imaged on a Zeiss i880 confocal

microscope at 40X and 63X (Plan-Apochromat 1.4 NA, oil), using a 405nm laser line (Cornell

BRC Imaging Core Facility). Egg chambers with any identifiable large, polyploid nuclei of nurse

cells indicated a successful event of Bam's differentiation function and were counted as nurse-

cell-positive egg chambers. Estimation statistics were used to assess effect size and a t-test was

used for statistical significance as described above for fertility of the female transgenic flies.

Correlation between nurse-cell-positive egg chamber data and fertility data was performed with

Kendall's rank correlation in R (v4.0.1).

**Cytological observations of the ovaries of female transgenic flies**

For cytology of the ten transgenic females with fertility defects and the control females,

flies were aged 2-3 days and then dissected and prepared for confocal microscopy as in [11, 33].

Ovaries were dissected in ice-cold 1x PBS and fixed in 4% paraformaldehyde (Electron

Microscopy Sciences) for 15 minutes. Tissue was rinsed 3x and washed 4x, respectively, with

PBST (1X PBS, 0.1% Triton-X 100), blocked in PBTA (1X PBS, 0.1% Triton-X100, 3% BSA)

(Alfa Aesar), and then incubated with primary antibodies overnight at 4°C. The rinse and wash

steps were repeated, followed by an overnight incubation with a goat anti-serum at 4°C to

minimize fluorescent background binding, and then a 1.5-hour incubation with secondary

antibodies. The rinse and wash steps were repeated and tissue was mounted in ProLong Diamond

AntiFade (Invitrogen).

The primary antibodies that were used were anti-Vasa (anti-rat, 1:20, Developmental

Studies Hybridoma Bank (DSHB)), anti-Hts-1B1 (anti-mouse, 1:40, DSHB), and anti-GFP (anti-

rabbit, 1:200, Invitrogen Cat No: G10362). Secondary antibodies that were used were Alexa488

(goat anti-rat, Invitrogen Cat No: A-11006), Alexa568 (goat anti-mouse, Invitrogen Cat No: A-

11031), and Alexa633 (goat anti-rabbit, Invitrogen Cat No: A-21070), all at 1:500.

A Zeiss i880 confocal microscope was used for all imaging with 405, 488, and 568nm

laser lines at 63X (Plan-Apochromat 1.4 NA, oil) (Cornell BRC Imaging Core Facility). Images

were edited using Fiji [42].

**RNAi knockdown of *bam***

We performed reciprocal crosses between a *w*Mel-infected *nos*-Gal4 and an uninfected

UAS-*bam*^HMS00029 line (a TriP *bam* shRNA line) (Bloomington *Drosophila* Stock Center lines

4442 and 33631, respectively) to generate the *bam* knockdown lines.

*Bam* expression was evaluated by RT-qPCR. RNA of four samples consisting of three 1-

2 day old female flies each were prepared using the NEB Monarch Total RNA Miniprep Kit.

RT-qPCR was performed with 10μL reactions (5μL Luna Universal One-Step Reaction Mix

(2X), 0.5μL Luna WarmStart RT Enzyme Mix (20X), 0.4μL forward primer, 0.4μL reverse

primer, 2μL template RNA, 1.7μL water), using three technical replicates and a standard curve.

Reverse transcription was performed at 55°C for 10 minutes, followed by initial denaturation at

95°C for 1 minute, and then 40 cycles of 95°C for 10 seconds and 60°C for 30 seconds. Primers

included 5'GCGCAATCGAAACGGAAACT3' and 5'GAGTAGCGGTGCTCCAGATC3' for

*bam* and 5'CCGCTTCAAGGGACAGTATC3' and 5'CAATCTCCTTGCGCTTCTTG3' for the

reference gene *Ribosomal protein L32* (*Rpl32*) (*Rpl32* primers are from [43]). Relative

expression to a wildtype *bam* control was done by the method of [44].

A fertility assay and a nurse-cell-positive egg chamber assay were performed as

described above, but at 29°C to increase the activity of the Gal4-UAS system. Estimation

statistics [40] were used to compare the mean fertility of the uninfected and infected *bam*

knockdown mutants. A Wilcoxon signed rank test was used to determine the statistical

significance of the mean nurse-cell-positive egg chambers between the uninfected and infected

*bam* knockdown mutants.

**Analysis of GSC mitosis frequency**

Nurse-cell-positive egg chamber assays were performed as described above to confirm

that the L255F bam mutation that we created as a transgene produced a partial *bam* mutant

phenotype and *w*Mel rescue like that of the original $bam^{bw}$ hypomorph.

Then, we assessed GSC mitosis frequency in the transgenic control (*w*; [$w^{+mc}$,

*bam::Venus*]/+; $bam^{\Delta 59}/bam^{\Delta 86}$) and the transgenic L255F mutant (*w*; [$w^{+mc}$, $bam^{L255F}::Venus$]/+;

$bam^{\Delta 59}/bam^{\Delta 86}$) by comparing the number of actively dividing GSCs indicated by mitosis marker

pH3 to the number of non-dividing GSCs. Flies were aged 2-3 days at 25°C, at which time

ovaries were dissected, immunostained and prepared for confocal microscopy, and imaged as

described above.

Primary antibodies that were used were anti-Vasa (anti-rat, 1:20, DSHB), anti-GFP (anti-

rabbit, 1:200, Invitrogen Cat No: G10362), and anti-pH3 (anti-rabbit, 1:200, Millipore Sigma Cat

No: 06-570). The same secondary antibodies listed above (Alexa488, Alexa568, and Alexa633)

were used at 1:500.

For each genotype, a Fisher's exact test was used to determine if there were more

mitotically active GSCs in the *w*Mel-infected germariums than in the uninfected germariums.

**Identification of b*am* sequence amino acid divergences**

*Bam* nucleotide sequences from the *D. melanogaster* Zambia population of the

*Drosophila* Population Genomics Project [45] were extracted for use in the MK test. Sequences

that had any ambiguous nucleotides (Ns) were removed, leaving 187 individual sequences for

our analyses. The *D. simulans* FlyBase reference sequence (r2.02) was used as an outgroup.

Sequences were aligned using PRANK v.170427 [46]. Nucleotide location of synonymous and

nonsynonymous divergences between *D. melanogaster* and *D. simulans* were identified using a

custom R script (v4.0.1) and mapped against the locations of the alanine mutations generated in

this study.

**Results**

**22 mutants were generated, ten with *bam* female fertility defects**

22 novel mutants were generated as *bam* transgenes with alanine replacements and a

Venus fluorescent tag. They were distributed across the five functional domains/regions of the

*bam* gene (Fig 1a). Each mutant was assayed for fertility defects and results were compared to a

control transgenic line of a wildtype *bam* gene tagged with Venus. Mutants were characterized

by comparing the average number of progeny produced by females with and without $w$Mel

infection.

Of the 22 *bam* alanine-scan mutants generated, ten had fertility defects in uninfected

females (Fig 2, Supplemental fig S1, Supplemental table S3). These included all three of the

mutants that targeted known binding residues in the CAF40 binding motif of *bam*, as well as

mutants that targeted residues in the Bgcn, Csn4, and ubiquitin binding regions. There were four

null-like fertility mutants that produced no progeny when uninfected (transgenic Venus-tagged

$bam^{L17A}$, $bam^{H180A,\ D181A,\ H182A}$, $bam^{K225A,\ E227A,\ R228A}$, and $bam^{E320A,\ H321A,\ R322A}$). There were six

hypomorph-like fertility mutants that produced less progeny on average than the control when

uninfected. The six hypomorph-like fertility mutants included three that exhibited severe fertility

defects (transgenic Venus-tagged $bam^{L28A}$, $bam^{E152A,\ K153A,\ H155A}$, and $bam^{R234A,\ R235A}$) and three

that exhibited mild fertility defects (transgenic Venus-tagged $bam^{M24A}$, $bam^{D249A,\ E250A}$, and

$bam^{E336A,\ R337A}$). The mean effect size between the number of progeny of the mutants with

fertility defects compared to the control ranged from a minimum of -28.2 to a maximum of -74.3

(Fig 2, Supplemental fig S3).

**Figure 2. Fertility of transgenic mutants compared to the respective *bam::Venus* transgenic control.** Data is colored according to the documented functional region that the mutation is located in. Below the jitter plots are the mean, two-sided permutation t-test p-value, and the average effect size between the mutant and the respective control. Gray panels indicate mutants with fertility defects (negative average effect size and p<0.05).

To verify that the fertility defects are a result of *bam*'s defective function in regulating GSC daughter cell differentiation, we counted the number of nurse-cell-positive egg chambers per ovary (Fig 3, Supplemental fig S2, Supplemental table S4), as well as looked for the presence of tumorous germariums in uninfected flies (Fig 4). The presence of an egg chamber with large

polyploid nurse cells indicates successful differentiation of a GSC daughter cell. Interestingly,

for three of the four null-like fertility mutant transgenic flies (transgenic Venus-tagged $bam^{L17A}$,

$bam^{H180A, D181A, H182A}$, and $bam^{K225A, E227A, R228A}$) contain egg chambers with nurse cells, indicating

rare occurrences of differentiation that did not translate to adult progeny in the fertility assays.

Thus, these mutants are only null-like in their fertility but are not true null alleles in the assessed

genetic background.

Two mutants with mild fertility defects (transgenic Venus-tagged $bam^{D249A, E250A}$ and

$bam^{M24}$) had mean nurse-cell-positive egg chamber counts that trended lower than the control,

but with small effect sizes (-5.54, p=0.086 and –6.28, p=0.023; respectively). Overall, fertility

and the number of nurse-cell-positive egg chambers correlated across the ten fertility-defective

$bam$ mutants (Kendall's rank correlation τ=0.828, Supplemental fig S3).

326
**Figure 3. Number of nurse-cell-positive egg chambers for transgenic mutants compared to**
**the *bam::Venus* control.** Data is colored according to the documented functional region that the
mutation is located in. Below the jitter plots are the mean, two-sided permutation t-test p-value,
the average effect size between the mutant and the respective control, and the corresponding
fertility classification as described in the text. Gray panels indicate mutants with nurse-cell-
positive egg chamber counts that are less than the control (negative average effect size and
p<0.05).

Our transgenic *bam::Venus* control had no tumorous germaria and had proper Bam

expression (Fig 4a-b). In parallel with the novel mutants we generated, we recreated the *bam^bw*

hypomorphic mutation as a transgenic *bam^L255F^::Venus* mutant. As has been previously

characterized for this L255F *bam* mutation in a *bam* null background, we see a tumorous

germarium with some cells containing a spectrosome (Fig 4c) [8, 10, 24]. We also found that the

overproliferating GSC-like cells in this hypomorph expressed Bam, and the lack of Bam

expression in the anterior most GSCs suggested that there was no ectopic expression of Bam in

the anterior part of the germarium that would cause a loss of GSCs.

The ten new *bam* fertility defective mutants largely displayed cytological tumorous

phenotypes like that observed in the *bam^L255F^::Venus* mutant (Fig 4d-f, Supplemental fig S4).

Tumorous phenotypes included severely tumorous germariums with no egg chambers in the

ovariole (Fig 4d); tumorous germariums with Bam expression in tumorous cells immediately

outside the germarium (Fig 4e); and tumorous germariums with some tumorous egg chambers

with no Bam expression (Fig 4f). Intriguingly, the *bam^M24A^::Venus* mutant, which had a mild

fertility defect, had no tumorous germariums (Supplemental fig S4e). All mutants, including the

severe mutants, stained successfully with anti-GFP, indicating the presence of Bam::Venus

protein.

351
**Figure 4. Wildtype and transgenic *D. melanogaster* ovarioles stained with anti-Hts-1B1**
**(cyan), anti-Vasa (magenta), and anti-GFP (yellow).** (a) CantonS wildtype control. (b)
Transgenic *bam::Venus* control. (c) Transgenic *bam*[L255F]*::Venus* mutant. (d-f) Representative
images of cytological defects observed in the transgenic *bam* mutants with fertility defects, with
arrowheads indicating spectrosomes, arrows indicating branched fusomes, and asterisks
indicating the apical tip of the germarium where GSCs are identifiable as cells not expressing
Bam::Venus. (d) Mutant E152A, K153A, H155A. (e) Mutant K255A, E227A, R228A. (f)
Mutant E336A, R337A. Scale bar for all images is 20μM.
360

That these ten new *bam* phenotypic mutants are distributed throughout the gene

demonstrates that multiple regions of the gene and protein are critical for *bam* function.

Additionally, the variation in severity of defects for mutations within one binding or functional

region shows that these phenotypes are allele specific.

The twelve *bam* mutants that did not have fertility defects when uninfected were also

distributed across the *bam* gene (Fig 2, Supplemental fig S1, Supplemental table S3). These were

the transgenic Venus tagged $bam^{D13A, D14A}$, $bam^{D18A, D19A, K22A}$, $bam^{E25A, E26A, H27A}$, $bam^{K108A, K109A,}$

$^{R111A}$, $bam^{K127A, E128A}$, $bam^{R133A, K134A, K136A}$, $bam^{E165A, H166A}$, $bam^{E246A, K247A}$, $bam^{D263A, K266A}$,

$bam^{K270A, K272A}$, $bam^{R300A, R301A}$, and $bam^{E417A, E418A, E419A}$. Three of these mutants ($bam^{E25A, E26A,}$

$^{H27A}$, $bam^{E417A, E418A, E419A}$, and $bam^{K127A, E128A}$) had greater fertility on average than their

respective control ($p<0.05$). Mutants without fertility defects include those with replacements at

highly conserved amino acid sites and replacements at poorly conserved sites and are found

throughout the gene (Supplemental fig S5, Supplemental file S1). These results identify residues

that are not critical to *bam*'s function in differentiation of GSC daughter cells in *D. melanogaster*

oogenesis, at least in laboratory conditions.

***w*Mel clearly rescues 8 of 10 fertility defective mutants**

To determine if *w*Mel rescues the new *bam* mutants with fertility defects, we performed

both the fertility assessment and nurse-cell-positive egg chamber assessment on the ten

transgenic mutants as uninfected and infected with *w*Mel. When infected with *w*Mel, the

transgenic *bam*::Venus control showed a slight increase in fertility across each independent

fertility assay ($p<0.05$, Fig 5, Supplemental table S3) and in the number of nurse-cell-positive

egg chambers (p<0.05, Fig 6). Thus, for a mutant to be classified with $w$Mel rescue, the increase

in mean effect size had to be greater than that observed in the control.

Of the four null-like fertility mutants that produced no progeny when uninfected, three

produced some progeny when infected with $w$Mel and showed an increase in nurse-cell-positive

egg chambers (transgenic Venus-tagged $bam^{L17A}$, $bam^{H180A,\ D181A,\ H182A}$, and $bam^{E320A,\ H321A,\ R322A}$)

(p<0.05). While the slight production of progeny cannot be classified as $w$Mel fertility rescue

because of the small effect size for the three mutants, there is rescue at the cytological nurse-cell

level for transgenic $bam^{L17A}$*::Venus* and $bam^{E320A,\ H321A,\ R322A}$::Venus. Transgenic $bam^{H180A,\ D181A,}$

$^{H182A}$*::Venus* trends towards rescue at the nurse-cell level, but has a small effect. The fourth null-

like fertility mutant ($bam^{E320A,\ H32A,\ R322A}$*::Venus*) did not produce progeny or nurse-cell-positive

egg chambers regardless of infection status (Fig 4, Supplemental table S2).

All six hypomorph-like fertility mutants that had an increase in average fertility (p<0.05,

Fig 5, Supplemental fig S6, Supplemental table S3) and average number of nurse-cell-positive

egg chambers with $w$Mel infection (p<0.05, Fig 6, Supplemental fig S7, Supplemental table S4),

with a mean effect size greater than that of the control ). The three mutants with severe fertility

defects (transgenic Venus-tagged $bam^{L28A}$, $bam^{E152A,\ K153A,\ H155A}$, and $bam^{R234A,\ R235A}$) saw an

approximate 20-fold increase in the average number of progeny per female when infected with

$w$Mel. The three mutants with mild fertility defects (transgenic Venus-tagged $bam^{M24A}$, $bam^{D249A,}$

$^{E250A}$, and $bam^{E336A,\ R337A}$) saw an approximate 2.5-fold increase in the average number of

progeny per female when infected with $w$Mel, bringing them to a fertility level similar to that of

the infected control (p>0.05 for all). The cytological data likewise showed an increase in nurse-

cell-positive egg chambers in $w$Mel-infected flies that brought the total numbers to be

comparable to the infected control (p>0.05 for all, except *bam*<sup>*E152A, K153A, H155A*</sup>*::Venus* where

p=0.042).

407
408

**Figure 5. Fertility of transgenic control and transgenic *bam* mutants with fertility defects as**
**uninfected and infected with *w*Mel.** Two-sided permutation t-test p-values and the average
effect size between the uninfected (blue) and infected with *w*Mel (red) data are shown below
each data set. Mutant E320A, H321A, R322A was completely sterile with and without *w*Mel and
thus is not shown. Gray panels highlight the mutants that are classified as having *w*Mel rescue at
the fertility level due having an average effect size greater than that observed in the respective
control and p<0.05.

| mean | 42.333 | 53.000 | 0.176 | 21.227 | 36.053 | 57.000 | 16.926 | 52.053 | 14.524 | 44.565 | 0.132 | 7.273 | 0.037 | 18.125 | 20.966 | 51.409 | 36.792 | 52.75 | 15.464 | 54.4 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| p-value | 0.005 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0.0058 | | 0 | |
| effect size | 10.7 | | 21.1 | | 20.9 | | 35.1 | | 30.0 | | 7.14 | | 18.1 | | 30.4 | | 16.0 | | 38.9 | |

**Figure 6. Number of nurse-cell-positive egg chambers of the transgenic *bam* mutants with fertility defects, as uninfected and infected with *w*Mel.** Two-sided permutation t-test p-values and the average effect size between the uninfected (blue) and infected with *w*Mel (red) data are shown below each jitter plot data set. Mutant E320A, H321A, R322A had no nurse-cell-positive egg chambers with and without *w*Mel and thus is not shown. Gray panels highlight the mutants that are classified as having *w*Mel rescue at the nurse-cell level, due to having an average effect size greater than that observed in the control and p<0.05.

Overall, we find that *w*Mel rescues the reduced fertility and/or nurse-cell-positive egg chambers of eight mutants across all the documented binding regions of *bam* and throughout the predicted 3D protein structure (Fig 7) [47, 48]. Of the last two mutants, the mutant that contains some nurse-cell-positive egg chambers when uninfected trends towards rescue and the mutant that is null-like at the fertility and nurse-cell level does not. These results show that the *w*Mel rescue is not limited to the L255F *bam* mutation and that *w*Mel interacts with *bam* in a manner that is not specific to the known binding regions of the gene.

**Figure 7. The locations of the mutations generated and their effect on fertility and *w*Mel rescue across the documented binding and functional regions of *bam*.** Shown in (a) a linear schematic of the gene and (b) the predicted protein structure. * indicates fertility defect mutants in which there was no *w*Mel rescue at the fertility and the nurse-cell-positive egg chamber level.

We plotted the mean progeny for all transgenic females (both uninfected and *w*Mel-

infected) to easily compare the above fertility results across all mutants (Fig 8). When comparing

the mean fertility for each mutant between the uninfected and *w*Mel-infected lines, we find that

the data falls roughly along a logarithmic curve, suggesting there is a relationship between the

severity of uninfected mutants and the increase in fertility seen with *w*Mel infection. However,

we cannot discount the possibility of the increase in fertility being due to an allele-specific

interaction with *w*Mel or variation in *Wolbachia* abundance.

454
**Figure 8. Mean fertility comparison of all transgenic lines generated with and without**
**_w_Mel.** The control lines are transgenic *bam::Venus* (black) and the transgenic mutants are
labeled by the first residue that was changed to alanine (orange for mutants with hypomorph-like
fertility defects, red for mutants with null-like fertility defects, and blue for mutants without
fertility defects). Figure shows data across five independent fertility assays with five different
controls.
461

**GSC mitosis in a *bam* hypomorphic mutant is not affected by *w*Mel**

To determine if the increase in fertility and the number nurse-cell-positive egg chambers

of *bam* mutants by *w*Mel is due to an increase in asymmetric division by GSCs, we quantified

GSC mitosis in the *bam::Venus* control and the *bam^{L255F}::Venus* mutant. The *bam^{L255F}* transgenic

hypomorph recapitulates the severity of the endogenous *bam^{bw}* hypomorph, though shows

slightly reduced rescue compared to the *bam^{bw}* hypomorph (Supplemental fig S8). This

difference is likely due to the differences in the genetic backgrounds of these two otherwise

identical mutations.

Cells in the apical tip of the germarium that expressed the pH3 mitosis marker but not the

Venus-tagged Bam protein were identified as GSCs actively undergoing mitosis (Supplemental

fig S9). The rate of GSC mitosis was not different between *w*Mel-infected and uninfected flies

within both the transgenic control and the transgenic hypomorph (p=1.0) (Table 1). Overall, the

*bam^{L255F}::Venus* transgenic hypomorph had fewer GSC mitotic events compared to the

transgenic control, but it was not statistically significant (p=0.259 with *w*Mel and p=0.448

without *w*Mel).

| | Transgenic control | | Transgenic *bam* hypomorph | |
|---|---|---|---|---|
| | uninfected | *w*Mel-infected | uninfected | *w*Mel-infected |
| # of germariums with pH3 positive GSCs | 5 | 9 | 2 | 4 |
| # of germariums without pH3 positive GSCs | 111 | 173 | 105 | 173 |

**Table 1. Number of mitotically active (pH3 positive) and quiescent GSCs**. GSC counts are
shown for the transgenic control (*bam::Venus*) and a transgenic hypomorph (*bam^{L255F}::Venus*)
when uninfected and infected with *w*Mel.

### *w*Mel does not rescue a *bam* knockdown phenotype

To determine if *w*Mel can rescue the phenotype of a reduced expression, but functional Bam protein in females, we tested for the rescue of a *bam* RNAi knockdown mutant by performing a reciprocal cross using the UAS-Gal4 system. We assessed both the number of egg chambers that have a nurse cell present (nurse-cell-positive egg chambers) per ovary in wildtype and *bam* knockdown mutants, as in [11, 12], as well as the fertility of the knockdown mutants.

We find that the *bam* knockdown mutants have fewer nurse-cell-positive egg chambers per ovary than the wildtype lines regardless of *w*Mel status ($p<0.05$) (Fig 9a, Supplemental fig S10). *Bam* knockdown mutants with *w*Mel had fewer nurse-cell-positive egg chambers on average than *bam* knockdown mutants without *w*Mel ($p<0.05$) but had similar levels of fertility over a period of seven days ($p = 0.69$, Fig 9b). This difference is likely due to the narrow window of time captured in the nurse-cell-positive assay compared to the fertility assay. These results show that *w*Mel does not rescue the tumorous mutant phenotype or the fertility of a *bam* knockdown mutant.

Additionally, there is no increase in *bam* expression in the knockdown mutant with *w*Mel compared to the uninfected knockdown mutant (Fig 9c). This shows that *w*Mel does not increase the expression levels of *bam,* consistent with the lack of rescue that we observe. All together, these results suggest that *w*Mel rescue is specific to partial-loss-of-function protein coding *bam* mutants.

**Figure 9. *Bam* knockdown by RNAi in the UAS-Gal4 system.** *Bam* wildtype 1 and 2 are the
UAS and Gal4 stocks, respectively; *bam* knockdowns are the nos-Gal4/+; UAS-*bam*[HMS00029]/+
reciprocal F1 progeny of the stocks. (a) The number of nurse-cell-positive egg chambers per
ovary. *** = significant at p<0.001; NS = not significant. (b) Fertility of the *bam* knockdown
mutants as uninfected and *w*Mel-infected. Jitter plots (left) show the raw data as dots with
horizontal lines representing the mean values of each dataset. The mean difference between
uninfected and infected lines are plotted as a bootstrap sampling distribution with a 95%
confidence interval (right). (c) *Bam* expression on a log2 scale relative to *bam* wildtype 1. Each
bar represents the dataset of four samples consisting of three female flies, and shows the mean
value with standard error.

**Discussion**

The critical reproductive gene *bam* is one of a few genes that has been shown to

genetically interact with *Wolbachia* in *D. melanogaster* [11, 18]. In female and male adults,

*bam*'s molecular role is to bring together other proteins into complexes, which then carry out

functions vital to the maintenance or progression of gametogenesis. *Bam*'s central and extensive

role in early reproduction, as well as *Wolbachia*'s potential for evolutionary impact motivates a

question about the specificity of the *bam-Wolbachia* interaction. Here, we first discuss our

results considering the molecular nature of the *bam-Wolbachia* interaction and then follow up

with a discussion on implications for the *D. melanogaster* host and *w*Mel *Wolbachia* variant

from an evolutionary perspective.

We targeted *bam*'s role of protein binding and minimized affecting the overall

conformation of the Bam protein by replacing charged residues with alanine throughout *bam*'s

documented binding regions [35]. We find no specificity to the presence/absence of rescue or the

magnitude of fertility rescue for the different functional regions of the linear *bam* sequence or

with the AlphaFold-predicted Bam protein structure. The strongest evidence for this comes from

our finding that *w*Mel rescues fertility defects of the known CAF40 binding residues (*bam*$^{M24A}$

and *bam*$^{L28A}$), as well as fertility defects of several mutations not involved in CAF40 binding.

This suggests that *w*Mel does not rescue the specific interaction between Bam and a single one of

its known protein partners in *bam* fertility mutants. Of note, however, is the possibility that the

fertility defects have arisen from a disruption of Bam protein structure despite the practice to

mitigate this with an alanine-scanning approach. Additionally, it is possible that the rescued

mutants' amino acid sites are involved in a direct interaction between Bam and an unmapped

binding partner (e.g., Otu [25]).

The severe and mild mutants were largely rescued to the level of the infected control

females. Thus, the severity of the mutation likely plays a large role in *w*Mel's ability to have a

substantial impact on host reproductive success. The biochemical nature of the allele interactions

and the effect of *w*Mel titer on host reproductive success are areas to be further investigated.

Additionally, some egg chambers were noticeably mutant in their cytology, which may explain

discrepancies between the nurse-cell-positive egg chamber data and the fertility data. As *bam*

also plays a role in mitotic divisions of the germline [9], nurse-cell-positive egg chambers may

be present, but with an incorrect number of nurse cells.

We tested two alternative hypotheses for how *w*Mel could be rescuing *bam* fertility

mutants. The first proposes that *w*Mel induces a fertility rescue by affecting *bam*'s expression. It

has previously been shown that there is no difference in *bam* expression in the *bam^{bw}* hypomorph

as uninfected and infected with *w*Mel [11]. Here, we tested the ability of *w*Mel to rescue the

mutant phenotype of a wildtype *bam* gene that is reduced in expression via RNAi. Importantly,

we have a knockdown mutant that displays the hypomorph phenotype of tumorous germariums

with some wildtype egg chambers. If the knockdown phenotype is too severe, *w*Mel may have

no effect as evidenced by the null-like fertility mutant generated in this study (transgenic

*bam^{E320A, H321A, R322A}::Venus*) and *w*Mel's lack of interaction with the *bam* null mutant

documented previously [11]. With the hypomorph-like-phenotype *bam* knockdown mutant (*nos-*

Gal4/+; UAS-*bam^{HMS00029}*/+), we find that *w*Mel does not rescue the number of nurse-cell-

positive egg chambers or affect the fertility of female flies; it also does not increase the

expression level of *bam.*

The second hypothesis proposes that *w*Mel increases the rate of GSC mitosis, which

increases the output of reproduction. Previous studies have shown *w*Mel to interact with GSCs,

including preliminary research in a *mei-P26* D. melanogaster mutant that shows increased GSC

mitosis in *w*Mel-infected germariums compared to uninfected germariums [49]. Other evidence

of *w*Mel's interaction with GSCs includes research on the $Sxl^{f4}$ *D. melanogaster* mutant that has

shown that *w*Mel restores the number of GSCs that were lost due to ectopic *bam* expression [22]

and the necessity for *w*Mel for GSC maintenance in filarial nematodes [50].

We find that *w*Mel does not increase the rate of GSC mitosis in the $bam^{L255F}::Venus$

hypomorph, and we do not observe any obvious ectopic Bam expression in the GSC niche in any

of our transgenic *bam* mutants. These results are consistent with what we see with the *bam*

knockdown results, as an increase in GSC mitosis rate would likely be reflected in an increase in

progeny in the *w*Mel-infected *bam* knockdown females.

Thus, while it is possible for there to be a subtle *w*Mel-GSC interaction that we have not

yet assessed, collectively, our results suggest that a large component of the *w*Mel *bam* interaction

is specific to mutant phenotypes that are the result of a disruption of *bam* function. Biochemical

assays are needed to firmly conclude that the *w*Mel *bam* interaction is specific to *bam* functions,

but it is notable that this proposal is akin to that suggested for *Wolbachia's* interaction with *Sxl*

[51]. The observations that *Wolbachia* cannot rescue the reduced egg count of a *snf* mutant,

which has no *Sxl* expression in the germline, and a $Sxl^{f5,fHa}$ mutant, which has a transposon

insertion in the second intron [21] parallels the *bam* knockdown results of our study.

It is possible that the fertility rescue of *bam* mutants occurs through the same (unknown)

mechanism by which the rescue of *Sxl* mutants occurs. This would be a process beyond the

documented restoration of GSC number in $Sxl^{f4}$ mutants by *Wolbachia* factor TomO, as it was

found that TomO is not sufficient to restore fertility of the $Sxl^{f4}$ mutants [22].

Bam and Sxl have been proposed to partner together to repress the translation of the stem

cell maintenance factor *nos* to determine GSC daughter cell fate [27, 28] and to maintain female

sexual identity in the adult germline [52]. Alternatively, it is possible that *Wolbachia* interacts

with *bam* and *Sxl* independently. Bam has a number of other genetic and physical interacting

partners that offer functional pathways for which *Wolbachia* could be interacting with. For

example, Bam promotes differentiation of GSC daughter cells by facilitating deubiquitination of

CycA with Otu [25].

Regardless of the manner in which *w*Mel molecularly interacts with *bam,* that *w*Mel

increases fertility of almost all *bam* protein-coding mutants generated in this study has broader

implications for *bam* and *Wolbachia's* population and evolutionary dynamics. A recent study

modeling potential *bam-Wolbachia* interactions suggests that *w*Mel's ability to mediate the

negative fitness consequences of protein-coding mutants like that which is observed here, is not

enough to generate the magnitude of positive selection currently observed at *bam* [53]. The

nonsynonymous amino acid replacements that are prevalent in *bam* across several *Drosophila*

species [31-33] may instead be due to arms race dynamics between *bam* and *Wolbachia*, an

evolutionary process independent of *Wolbachia,* or some combination of the two.

For *Wolbachia*, the increased reproductive success for almost all *bam* protein-coding

mutants with *w*Mel infection is an addition to *w*Mel's arsenal of promoting its transmission and

spread through populations. It is particularly advantageous for *w*Mel, as, unlike some other

*Wolbachia* variants, it exhibits minimal cytoplasmic incompatibility in field-caught *D.*

*melanogaster* [19]. Ultimately, that *w*Mel rescues fertility defects of multiple *bam* protein-coding

mutants, as well as evidence that *Wolbachia* rescues reproductive defects of other germline genes

such as *Sxl* and *mei-P26*, may allow for opportunities for an obligate symbiotic relationship

between *D. melanogaster* host and *w*Mel to develop.

**Supplemental file S1.** Summary of all generated transgenic mutants. Included for each mutant

are the amino acid site(s) and the corresponding functional and/or binding region; the original

codon and residue; the number of species that the residue is conserved across from the seven

species (*D. melanogaster*, *D. simulans, D. yakuba*, *D. santomea, D. teissieri*, *D. erecta*, and *D.*

*eugracilis*) multiple alignment of [38]; the alanine codon that replaced the original codon

sequence; and the results of the fertility and nurse-cell-positive egg chamber experiments.

**Acknowledgements**

We thank Dan Barbash for sharing the pCasper\attB vector and are grateful to Eric Alani

for advice on generating alanine-scanning mutants. We thank Mariana Wolfner and Andy Clark

for insightful discussions and review of the manuscript. We also thank Jackie Bubnell, Catherine

Kagemann, and Luke Arnce for thoughtful feedback throughout this project. This work was

supported by National Institutes of Health grant R01-GM095793 to C.F.A. and by NYSTEM

C029155 and NIH S10OD018516 grants to the Cornell Institute of Biotechnology Imaging

Center for the Zeiss i880 LSM880 microscope.

REFERENCES

1. McKearin DM, Spradling AC. *Bag-of-marbles*: a *Drosophila* gene required to initiate both male and female gametogenesis. Genes Dev. 1990;4: 2242-2251.

2. Gönczy P, Matunis E, DiNardo S. *bag-of-marbles* and *benign genial cell neoplasm* act in the germline to restrict proliferation during *Drosophila* spermatogenesis. Development 1997;124(21): 4361-4371.

3. Insco ML, Leon A, Tam CH, McKearin DM, Fuller MT. Accumulation of a differentiation regulator specifies transit amplifying division number in an adult stem cell lineage. Proc Natl Acad Sci. 2009;106(52): 22311-22316.

4. Insco ML, Bailey AS, Kim J, Olivares GH, Wapinski OL, Tam CH, et al. A self-limiting switch based on translational control regulates the transition from proliferation to differentiation in an adult stem cell lineage. Cell Stem Cell 2012;11: 689-700.

5. Chen D, McKearin D. Dpp signaling silences *bam* transcription directly to establish asymmetric divisions of germline stem cells. Curr Biol. 2003;13(20): 1786–1791.

6. Chen D, McKearin DM. A discrete transcriptional silencer in the bam gene determines asymmetric division of the *Drosophila* germline stem cell. Development 2003;130: 1159–1170.

7. Song X, Wong MD, Kawase E, Xi R, Ding BC, McCarthy JJ, et al. Bmp signals from niche cells directly repress transcription of a differentiation-promoting gene, *bag of marbles*, in germline stem cells in the *Drosophila* ovary. Development 2004;131: 1353-1364.

8. Shen R, Weng C, Yu J, Xie T. eIF4A controls germline stem cell self-renewal by directly inhibiting BAM function in the *Drosophila* ovary. Proc Natl Acad Sci USA 2009;106: 11623–11628.

645    9. McKearin D and Ohlstein B. A role for the *Drosophila* Bag-of-marbles protein in the

646           differentiation of cystoblasts from germline stem cells. Development 1995;121: 2937–

647           2947.

648    10. Ohlstein B, Lavoie CA, Vef O, Gateff E, Mckearin DM. The *Drosophila* cystoblast

649           differentiation factor, *benign gonial cell neoplasm*, is related to DExH-box proteins and

650           interacts genetically with *bag-of-marbles*. Genetics 2000;155: 1809-1819.

651    11. Flores HA, Bubnell JE, Aquadro CF, Barbash DA. The *Drosophila bag of marbles* gene

652           interacts genetically with *Wolbachia* and shows female-specific effects of divergence.

653           PLoS Genet. 2015;11: e1005453.

654    12. Bubnell JE, Fernandez-Begne P, Ulbing CKS, Aquadro CF. Diverse *w*Mel variants of

655           *Wolbachia pipientis* differentially rescue fertility and cytological defects of the *bag of*

656           *marbles* partial loss of function mutation in *Drosophila melanogaster*. G3 (Bethesda)

657           2021;11(12).

658    13. Hilgenboecker K, Hammerstein P, Schlattmann P, Telschow A, Werren JH. How many

659           species are infected with *Wolbachia*? — a statistical analysis of current data. FEMS

660           Microbiol Lett. 2008;281: 215-220.

661    14. Weinert LA, Araujo-Jnr EV, Ahmed MZ, Welch JJ. The incidence of bacterial

662           endosymbionts in terrestrial arthropods. Proc R Soc B. 2015;282(1807): 20150249.

663    15. Taylor MJ, Bandi C, and Hoerauf A. *Wolbachia* bacterial endosymbionts of filarial

664           nematodes. Adv Parasitol. 2005;60: 245-284.

665    16. Werren JH, Baldo L, Clark ME. *Wolbachia*: master manipulators of invertebrate biology. Nat

666           Rev Microbiol. 2008;6: 741-751.

17. Kaur R, Shropshire JK, Cross KL, Leigh B, Mansueto AJ, Steward V, et al. Living in the

endosymbiotic world of *Wolbachia*: A centennial review. Cell Host Microbe 2021;29(6):

879-93.

18. Solignac M, Vautrin D, Rousset F. 1993. Widespread occurrence of the proteobacteria

*Wolbachia* and the partial cytoplasmic incompatibility in *Drosophila melanogaster*. *C R*

*Acad Sci Paris* 317:461-470.

19. Hoffmann AA, Hercus M, Dagher H. 1998. Population dynamics of the *Wolbachia* infection

causing cytoplasmic incompatibility in *Drosophila melanogaster*. *Genetics* 148:221-231.

20. Reynolds KT, Hoffmann AA. 2002. Male age, host effects and the weak expression or non-

expression of cytoplasmic incompatibility in *Drosophila* strains infected by maternally

transmitted *Wolbachia*. *Genet Res*. 80:79-87.

21. Starr DJ, Cline TW. A host-parasite interaction rescues *Drosophila* oogenesis defects. Nature

2002;418: 76-79.

22. Ote M, Ueyama M, Yamamoto D. *Wolbachia* protein TomO targets *nanos* mRNA and

restores germ stem cells in *Drosophila sex-lethal* mutants. Curr Biol. 2016;26: 2223-2232.

23. Sgromo A, Raisch T, Bawankar P, Bhandari D, Chen Y, Kuzuoğlu-Öztürk D, et al. A

CAF40-binding motif facilitates recruitment of the CCR4-NOT complex to mRNAs

targeted by Drosophila Roquin. Nat Commun. 2017;8: 14307.

24. Pan L, Wang S, Lu T, Weng C, Song X, Park JK, et al. Protein competition switches the

function of COP9 from self-renewal to differentiation. Nature 2014;514: 233-236.

25. Ji S, Li C, Hu L, Liu K, Mei J, Luo Y, et al. Bam-dependent deubiquitinase complex can

disrupt germ-line stem cell maintenance by targeting cyclin A. Proc Natl Acad Sci USA

2017;114: 6316-6321.

26. Li Y, Minor NT, Park JK, McKearin DM, Maines JZ. Bam and Bgcn antagonize Nanos-

dependent germ-line stem cell maintenance. Proc Natl Acad Sci USA. 2009;106(23):

9304–9309.

27. Chau J, Kulnane LS, Salz HK. *Sex-lethal* enables germline stem cell differentiation by down-

regulating Nanos protein levels during *Drosophila* oogenesis. Proc Natl Acad Sci USA.

2012;109: 9465–9470.

28. Malik S, Jang W, Kim JY, Kim C. Mechanisms ensuring robust repression of the *Drosophila*

female germline stem cell maintenance factor Nanos via posttranscriptional regulation.

The FASEB Journal 2020;34: 11421-11430.

29. Fu Z, Geng C, Wang H, Yang Z, Wenig C, Li H, et al. Twin promotes the maintenance and

differentiation of germline stem cell lineage through modulation of multiple pathways.

Cell Rep. 2015;13: 1366-1379.

30. Chen D, Wang Q, Huang H, Xia L, Jiang X, Kan L, et al. Effete-mediated degradation of

Cyclin A is essential for the maintenance of germline stem cells in *Drosophila*.

Development 2009;136: 4133-4142.

31. Civetta A, Rajakumar SA, Brouwers B, Bacik JP. Rapid evolution and gene-specific patterns

of selection for three genes of spermatogenesis in *Drosophila*. Mol Biol Evol. 2006;23:

655-662.

32. Bauer DuMont VL, Flores HA, Wright MH, Aquadro CF. Recurrent positive selection at

*bgcn*, a key determinant of germ line differentiation, does not appear to be driven by

simple coevolution with its partner protein *bam*. Mol Biol Evol. 2007;24: 182–191.

33. Bubnell JE, Ulbing CKS, Fernandez Begne P, Aquadro CF. Functional divergence of the

*bag-of-marbles* gene in the *Drosophila melanogaster* species group. Mol Biol Evol.

2022;39(7).

34. Flores HA, Bauer DuMont VL, Fatoo A, Hubbard D, Hijji M, Barbash DA, et al. Adaptive

evolution of genes involved in germline stem cells in *Drosophila melanogaster* and *D.*

*simulans.* G3 (Bethesda) 2015;5: 583-592.

35. Cunningham BC, Wells JA. High-resolution epitope mapping of hGH-receptor interactions

by alanine-scanning mutagenesis. *Science* 1989;244: 1081-1085

36 Nakamura Y. Kisarazu (Japan): Kazusa DNA Research Institute. Codon Usage Database.

[last accessed 2021 Oct 12].

https://www.kazusa.or.jp/codon/cgi-bin/showcodon.cgi?species=7227

37. Groth AC, Fish M, Nusse R, Calos MP. Construction of transgenic *Drosophila* by using the

site-specific integrase from phage phiC31. Genetics 2004;166: 1775-1782.

38. Bubnell JE. Functional evolution of the *bag of marbles* gene in *Drosophila.* PhD thesis,

Cornell University. 2020. Available from https://hdl.handle.net/1813/103228

39. New England Biolabs. 2021. NEBaseChanger. [last accessed 2021 Oct 13].

https://nebasechanger.neb.com/

40. Ho J, Tumkaya T, Aryal S, Choi H, Claridge-Chang A. Moving beyond P values: data

analysis with estimation graphics. Nat Methods. 2019;16: 565-566.

41. Claridge-Chang A, Ho J. Estimation Statistics: Analyze your data with effect Sizes. C2017-

2021. [accessed 2022 Sept 20]. https://www.estimationstats.com/#/

42. Shindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an

open-source platform for biological-image analysis. Nature Methods 2012;9(7): 676-682.

43. Chrostek E, Marialva MSP, Esteves SS, Weinert LA, Martinez J, Jiggins FM, et al.

*Wolbachia* variants induce differential protection to viruses in *Drosophila melanogaster*:

a phenotypic and phylogenomic analysis. PLoS Genet. 2013;9: e1003896.

44. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-

PCR. Nucleic Acids Res. 2001;29: e45.

45. Lack JB, Cardeno CM, Crepeau MW, Taylor W, Corbett-Detig RB, Stevens KA, Langley

CH, Pool JE. The *Drosophila* genome nexus: A population genomic resource of 623

*Drosophila melanogaster* genomes, including 197 from a single ancestral range

population. Genetics 2015;199:1229-1241.

46. Löytynoja A. Phylogeny-aware alignment with PRANK. *Methods Mol Biol*. 2014;1079:155-

170.

47. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate

protein structure prediction with AlphaFold. Nature 2021;596: 583-589.

48. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold

Protein Structure Database: massively expanding the structural coverage of protein-

sequence space with high-accuracy models. Nucleic Acids Res. 2021;50: D439-D444.

49. Russell SL, Ruelas Castillo J, Sullivan WT. *Wolbachia* endosymbionts manipulate GSC self-

renewal and differentiation to reinforce host fertility. BioRxiv 2022.

https://doi.org/10.1101/2022.12.15.520626

50. Foray V, Pérez-Jiménez MM, Fattouh N, Landmann F. *Wolbachia* control stem cell behavior

and stimulate germline proliferation in filarial nematodes. Dev Cell. 2018;45: 198-211.e3.

51. Ote M, Yamamoto D. Impact of *Wolbachia* infection on *Drosophila* female germline stem

cells. Curr Opin Insect Sci. 2020;27: 8-15.

52. Shapiro-Kulnane L, Smolko AE, Salz HK. Maintenance of *Drosophila* germline stem cell

sexual identity in oogenesis and tumorigenesis. Development 2015;142(6): 1073-1082.

53. Shen R, Wenzel M, Messer PW, Aquadro CF. A model of functionall buffered deleterious

mutations can lead to signatures of positive selection. 2023. bioRxiv.

https://doi.org/10.1101/2022.02.14.480440

54. Mateos M, Castrezana SJ, Nankivell BJ, Estes AM, Markow TA, Moran NA. 2006. Heritable

endosymbionts of *Drosophila*. *Genetics* 174:363-376.

55. Merçot H, Charlat S. 2004. *Wolbachia* infections in *Drosophila melanogaster* and *D.*

*simulans*: polymorphism and levels of cytoplasmic incompatibility. In: Capy, P., Gibert,

P., Boussy, I. (eds) Drosophila melanogaster, Drosophila simulans: So Similar, So

Different. Contemporary Issues in Genetics and Evolution, vol 11. Springer, Dordrecht.

56. Zabalou S, Charlat S, Nirgianaki A, Lachaise D, Merçot H, Bourtzis K. 2004. Natural

*Wolbachia* infections in the *Drosophila yakuba* species complex do not induce

cytoplasmic incompatibility but fully rescue the wRi modification. *Genetics* 167(2):827-

34.

57. Bourtzis K, Nirgianaki A, Markakis G, Savakis C. 1996. *Wolbachia* infection and

cytoplasmic incompatibility in *Drosophila* species. *Genetics* 144(3):1063-73.

58. Ravikumar H, Prakash BM, Sampathkumar S, Puttaraju HP. 2011. Molecular subgrouping of

*Wolbachia* and bacteriophage WO infection among some Indian *Drosophila* species. *J*

*Genet*. 90:507-510.

59. Kopp A, Barmina O. 2005. Evolutionary history of the *Drosophila bipectinata* species

complex. *Genet Res*. 85:23-46.

60. Brown, J. 2021. Microbiomes in the context of insect communities [PhD thesis, University of

South Bohemia]. Theses.cz. https://theses.cz/id/17gyx3/

61. Richardson KM, Schiffer M, Griffin PC, Lee SF, Hoffmann AA. 2016. Tropical *Drosophila*

*pandora* carry *Wolbachia* infections causing cytoplasmic incompatibility or male killing.

*Evolution* (8):1791-802.

62. Haine ER, Pickup NJ, and Cook JM. 2005. Horizontal transmission of *Wolbachia* in a

*Drosophila* community. *Ecol Entomol.* 30:464-472.