Cell type-specific control of secondary cell wall formation by Musashi-type translational regulators in Arabidopsis

Alicia Kairouani,[1]* Dominique Pontier,[1]* Claire Picart,[1] Fabien Mounet,[2] Yves Martinez,[3] Lucie Le-Bot,[4] Mathieu Fanuel,[4,5] Philippe Hammann,[6] Lucid Belmudes,[7] Rémy Merret,[1] Jacinthe Azevedo,[1] Marie-Christine Carpentier,[1] Dominique Gagliardi,[8] Yohann Couté,[7] Richard Sibout,[4] Natacha Bies-Etheve,[1@] Thierry Lagrange[1@]

[1]Laboratoire Génome et Développement des Plantes, Université de Perpignan via Domitia, CNRS, UMR5096, Perpignan Cedex, France.

[2]Laboratoire de Recherche en Sciences Végétales, Université de Toulouse III, CNRS, INP, UMR5546, Castanet-Tolosan, France.

[3]FRAIB-CNRS plateforme imagerie, Castanet-Tolosan, France.

[4]Biopolymères Interactions Assemblages, UR1268 BIA, INRAE, Nantes, France.

[5]PROBE research infrastructure, BIBS Facility, INRAE, Nantes, France.

[6]Plateforme Protéomique Strasbourg Esplanade de CNRS, Université de Strasbourg, Strasbourg, France.

[7]Université Grenoble Alpes, INSERM, UA13 BGE, CNRS, CEA, FR2048, Grenoble, France.

[8]Institut de Biologie Moléculaire des Plantes, BMP, CNRS, Université de Strasbourg, Strasbourg, France.

*These authors contributed equally to this work.

@ Correspondence to: Natacha Bies-Etheve, Thierry Lagrange (ORCiD: 0000-0003-3090-0918).

Email: natacha.etheve@univ-perp.fr; lagrange@univ-perp.fr

Deciphering the mechanism of secondary cell wall/SCW formation in plants is key to

understanding their development and the molecular basis of biomass recalcitrance.

Although transcriptional regulation is essential for SCW formation, little is known

about the implication of post-transcriptional mechanisms in this process. Here we report

that two *bonafide* RNA-binding proteins homologous to the animal translational

regulator Musashi, MSIL2 and MSIL4, function redundantly to control SCW formation

in Arabidopsis. MSIL2/4 interactomes are similar and enriched in proteins involved in

mRNA binding and translational regulation. MSIL2/4 mutations alter SCW formation

in the fibers, leading to a reduction in lignin deposition, and an increase of 4-*O*-

glucuronoxylan methylation. In accordance, quantitative proteomics of stems reveal an

overaccumulation of glucuronoxylan biosynthetic machinery, including GXM3, in the

***msil2/4* mutant stem. We showed that MSIL4 immunoprecipitates *GXM* mRNAs,**

suggesting a novel aspect of SCW regulation, linking post-transcriptional control to the

regulation of SCW biosynthesis genes.

Introduction

As the main constituent of the terrestrial plant biomass, lignocellulose represents a huge reservoir of fixed carbon and a renewable resource for bioproducts and energy that is essential for human usages (Meents et al., 2018). Lignocellulose biomass enclosed mainly within secondary cell wall (SCW) reinforces vessels for long-distance transport and provides physical properties to fibers allowing upright growth of the plants (Meents et al., 2018; Scheller and Ulvskov, 2010). SCWs are composite material made of high molecular weight biopolymers, including cellulose, hemicelluloses, and lignin, as well as cell wall proteins whose composition and structure can differ markedly between dicots and monocots (Scheller and Ulvskov, 2010; Yokoyama and Nishitani, 2004). Cellulose is synthesized at the plasma membrane by the cellulose synthase complex and is composed of linear chains of β-(1–4) glucans that aggregate to form highly crystalline cellulose microfibrils (Scheller and Ulvskov, 2010; Yokoyama and Nishitani, 2004). Xylans, the main hemicellulosic polysaccharides in SCW, are synthesized in the Golgi apparatus by a complex biosynthetic machinery and comprise a group of polysaccharides that share a common backbone of β-(1,4)-linked xylose (Xyl) units, but differs by the presence of side chains, whose nature is dependent on tissues and species (Scheller and Ulvskov, 2010). In particular, glucuronoxylan, the dominant xylan in the SCW of dicots, is notably substituted at O-2 position with α-D-glucuronic acid (GlcA) or (4-O-methyl)- α-D-glucuronic acid (MeGlcA) groups, and with O-acetyl moieties at C-2 or C-3 positions (Scheller and Ulvskov, 2010). By interacting with the cellulose and lignin, xylans contribute to the strengthening of the SCW, and are considered to be one of the main factors contributing to the biomass recalcitrance to enzymatic hydrolysis (Yokoyama and Nishitani, 2004; Kumar et al., 2016). Finally, lignin polymers are made up of H, G, and S units which are the results of the oxidation and the polymerisation of the p-coumaryl, coniferyl and Synapyl alcohols, respectively. These monomers, also called monolignols are

synthesized on the ER surface/cytoplasm prior to transportation to the cell wall. Lignins

confer stiffness, strength and hydrophobicity to the SCW (Scheller and Ulvskov, 2010;

Yokoyama and Nishitani, 2004; Kumar et al., 2016).

The metabolic load imposed by the coordinated production of SCW biopolymers

emphasizes the importance for plants of having a precise control over temporal and spatial

expression of the corresponding SCW biosynthetic genes. Based on founding studies

performed in the Arabidopsis model, it is generally recognized that the SCW synthesis is

primarily controlled at a transcriptional level by a multilayer and interconnected network of

transcription factors/TFs (Taylor-Teeples et al., 2015; Zhang et al., 2018). Key players of this

regulatory network are related NAC (NAM- ATAF1/2-CUC) and MYB-type TFs that induce,

in a redundant and combinatorial manner, the expression of SCW biosynthetic genes by

binding their promoter region (Taylor-Teeples et al., 2015). In addition, a family of Class III

homeodomain leucine zipper (HD-ZIPIII) TFs, whose members control several aspects of

Arabidopsis development, was also shown to interact with SCW regulatory network,

contributing to xylem cell specification and SCW synthesis (Taylor-Teeples et al., 2015;

Zhang et al., 2018). Although this transcriptional level of control can gain complexity in

perennial plants, functional orthologs of the main regulators were identified in many different

species, suggesting that SCW deposition proceeds via a conserved mechanism in all vascular

plants (Myburg et al., 2014; Zhong and Ye, 2014). During the last decade, engineering the

SCW regulatory network to deposit modified lignin and/or improve polysaccharide

composition became a promising strategy to optimize biomass processability (Sibout et al.,

2005; Whitehead et al., 2018). However, despite huge progresses in understanding the

regulation of SCW synthesis, the engineering of SCW regulatory network to improve biomass

yield and digestibility has proven difficult and often impairs plant growth because of altered

vascular tissue development (Bonawitz and Chapple, 2013; Wang et al., 2016).

Downstream of transcriptional regulations by TFs, RNA-binding proteins (RBPs) are

essential post-transcriptional modulators of gene expression across all kingdoms of life (Singh

et al., 2015). However, although a large cohort of RBPs exists in plants (Reichel et al., 2016),

there is little information about RBP-mediated posttranscriptional regulation of SCW

biosynthetic genes. So far, only few microRNAs (miRNAs) families have been implicated in

the post-transcriptional regulation of genes involved in both regulatory and enzymatic aspects

of SCW biosynthesis (Zhang et al., 2018). In particular, miR165/166 were shown to target

HD-ZIPIII transcriptional regulators, having a direct impact on the transcriptional network

associated with SCW biosynthesis (Zhang et al., 2018). In the other hand,

miRNA397/857/408 were shown to affect lignin content *via* the targeting of laccase/LAC

genes, which control lignin polymerization from monolignols precursors (Wang et al., 2014;

Zhao et al., 2015). In addition, two related tandem CCCH zinc-finger proteins, which exhibit

both DNA and RNA binding abilities, have been proposed to modulate SCW formation by

regulating the expression of genes associated with cell wall metabolism (Chai et al., 2015).

Despite these mounting evidences, our current knowledge of the role of post-transcriptional

regulators in SCW synthesis is still very limited.

In this work, we report that two RRM-domain containing RNA-binding proteins

homologous to the animal translational regulator Musashi, Musashi-like2/MSIL2 and

Musashi-like4/MSIL4, function redundantly to control various aspects of development,

including the stiffness of the inflorescence stem in Arabidopsis. We show that RRM-

dependent RNA-binding activity is essential for MSIL2/4 functions *in vivo*, and that MSIL2/4

interactomes are similar, being enriched in proteins involved in 3'-UTR binding and

translational regulation. MSIL2/4 mutations alter the formation of SCW in the interfascicular

fiber cells, leading to a reduction in lignin deposition, and a change in the decoration pattern

of glucuronoxylan that is associated with an increase of 4-*O*-methylation of GlcA substituent.

In accordance, quantitative mass-spectrometry-based protein analysis reveals an

overaccumulation of glucuronoxylan biosynthetic machinery, including the GlucuronoXylan

Methyltransferase3/GXM3, in the *msil2/4* mutant stem. We show that MSIL4

immunoprecipitates GXM3 mRNA in vivo, likely regulating its expression at a translational

level. Our results demonstrate that MSILs regulate SCW synthesis in interfascicular fiber

cells and point to a novel aspect of SCW regulation linking translational repression to

regulation of SCW biosynthesis genes.

Results

Musashi-like MSIL2/4 proteins redundantly control specific Arabidopsis development

processes.

While many studies have implicated the MSIs in the control of gene expression during

cellular proliferation, cell fate determination and cancer in animals (Fox et al., 2015; Kudinov

et al., 2017), little is known about the presence and activity of MSI-type proteins in plants. In

this work, we describe an RNA-binding protein family in Arabidopsis, whose members

hereafter named as *MUSASHI-Like1* to *4/MSIL1-4*, share both sequence, domain organization,

and model-based structural similarities with animal MSIs (Figure 1A and B). Phylogenetic

analysis revealed that Arabidopsis indeed harbors a clade of seven *MSIL*-type genes with a

prominent sub-clade harboring the *MSIL1-4* genes (Figure 1B). Notably, a neighboring sub-

clade contains two genes, *RBGD2* and *RGBD4* (Figure 1B), that encode for heat-inducible

RBPs implicated in the response to heat stress in Arabidopsis (Zhu et al., 2022). MSIL1-4

proteins share a common domain architecture consisting of two N-terminal RNA recognition

motifs (RRM1 and RRM2) that are followed by a poorly conserved, intrinsically unstructured,

carboxy-terminal region (Figure 1A and Figure1-figure supplement 1A). The survey of the

gene expression ARAPORT11 databases (https://araport.org/; http://fgcz-pep2pro.uzh.ch)

revealed that the *MSIL1-4* genes are widely expressed in Arabidopsis, with *MSIL1* showing

the lowest levels of expression (Figure 1C). Database searches further indicated that *MSIL*

orthologs are widely distributed in land plants, from bryophytes to angiosperms (Figure1-

figure supplement 1B). To gain a mechanistic understanding of MSIL function in Arabidopsis,

we determined the subcellular localization of these proteins by performing fluorescence

microscopy on GFP-tagged versions of MSIL2 and MSIL4 (MSIL2G and MSIL4G). Both

proteins showed a diffuse cytoplasmic distribution in root cells of stable Arabidopsis

transgenic plants that contrasts with the nucleus and cytoplasmic distributions of the free GFP

protein control (Figure 1D).

To further investigate the function of the MSILs in Arabidopsis, we raised specific

antibodies against non-conserved regions of MSIL1-4 and characterized T-DNA insertion

msil mutant lines that lack full-length *MSIL* mRNA and MSIL proteins as judged by RT-PCR

and western blot experiments, respectively (Figure1-figure supplement 2A-C). However, none

of the four characterized single *msil* mutants showed any obvious phenotype (Figure1-figure

supplement 3A and data not shown). To assess the potential functional redundancy between

MSIL genes, we crossed the single *msil* mutants together and characterized the six possible

double mutant combinations (Figure1-figure supplement 3A). Only the *msil2*-1/*msil4*-1

(*msil2/4*) double mutant showed discernable developmental abnormal phenotypes, including

enlarged and curled leaves, early leaf senescence and a pendant inflorescence stem (Figure 1E

and F and Figure1-figure supplement 3A-C). To assess whether MSIL1 and MSIL3 also

contribute to Arabidopsis development in the *msil2/4* double mutant background, we

generated a *msil1*-1/*msil2*-1/*msil3*-1/*msil4*-1 (*msil1/2/3/4*) quadruple mutant that shows no

particular or aggravated developmental phenotypes with respect to the *msil2/4* double mutant

(Figure1-figure supplement 3A-C). To test the functional redundancy of MSIL2 and MSIL4

proteins *in vivo*, we transformed the *msil2/4* double mutant with constructs coding either for a

Flag/HA-tagged version of MSIL2 (MSIL2F) or MSIL4 (MSIL4F), and selected two

independent transgenic plants expressing similar and near physiological levels of the tagged

protein, as evaluated by western blot using our home-made antibodies (Figure1-Figure

supplement 3D). The ectopic expression of either MSIL2F or MSIL4F proteins was able to

rescue all the phenotypes associated with the MSIL2/4 defect (Figure 1E and F and Figure1-

Figure supplement 3E). This confirms that MSIL2F and MSIL4F are functional proteins and

more importantly that MSIL2 and MSIL4 redundantly control various aspects of development

in Arabidopsis.

RRM-dependent RNA-binding activity is essential for MSIL2/4 functions *in vivo*.

Animal MSIs interact with target transcripts through their RRM domains and mutations in

these domains have been shown to impair MSI activity *in vivo* (Chavali et al., 2017; Nguyen

et al., 2020). MSILs are very similar to metazoan MSIs over their RRM sequences (Figure 1A

and Figure 2-figure supplement 1A), a closeness of sequences also observed when homology

models of the MSIL4 RRM motifs were generated using SWISS-MODEL (Biasani et al.,

2014) and AlphaFold2 (Jumper et al. 2021) servers (Fig. 2A and Figure 2-figure supplement

1B). Homology-based structure prediction proposed that these domains adopt the

characteristic RRM fold with a four-stranded β-sheet structure bearing the conserved

phenylalanine residues involved in the specific recognition of RNA bases (Daubner et al.,

2013) (Figure 2A and Figure 2-figure supplement 1A). This supports the idea that MSILs

have retained the ability to interact with RNA, a notion further supported by the identification

of MSIL2/4 proteins in the experimentally determined Arabidopsis mRNA-binding proteome

(Reichel et al., 2016). To examine the requirement for the interaction of MSIL with RNA in

developmental control, we first generated an RNA-binding mutant form of MSIL4,

MSIL4^{RRM}, by introducing phenylalanine to aspartate (F→D) mutations in both RRM

domains (Figure 2B). Protein-RNA interaction assays using RNA homopolymers

immobilized on agarose beads confirmed that the MSIL4, but not MSIL4^{RRM}, exhibited

intrinsic RNA binding activity *in vitro* (Figure 2C). To assess the impact of these mutations *in*

vivo, the MSIL4G and MSIL4G^{RRM} constructs were expressed under the control of the *MSIL4*

endogenous promoter into the *msil2/4* mutant, and two independent transformants expressing

near physiological levels of the WT (*MSIL4G*-3 and *MSIL4G*-8), and mutant (*MSIL4G^{RRM}*-3

and *MSIL4G^{RRM}*-10) proteins were selected for further complementation analysis (Figure 2-

figure supplement 2A). MSIL4G, but not MSIL4G^{RRM} proteins were able to rescue the

developmental defects incurred by *MSIL2/4* mutations (Figure 2D and Figure 2-figure

supplement 2B-C), indicating that these proteins are likely to exert their functions by

interacting with RNA targets *in vivo*.

MSIL2/4 protein interactomes are enriched in proteins involved in 3'-UTR binding and

translational regulation.

Animal MSIs modulate mRNA expression mostly at a translational level through binding to

conserved motifs in the 3'-UTR of the target mRNA and further interactions with various

mRNA binding partners (Fox et al., 2015; Weill et al., 2017; Cragle et al., 2019). To further

characterize the components of the MSIL2/4 network in Arabidopsis, we performed affinity

purification coupled to LC-MS/MS, as described in Scheer et al. (Scheer et al., 2021), using

the complemented *MSIL2F1* and *MSIL4F1* lines. Consistent with their functional redundancy,

MSIL2F1 and MSIL4F1 exhibit a similar protein interaction network, with 27 proteins

significantly enriched in both MS analyses (Figure 3A and B, and Supplementary file 1).

Examination of the PANTHER database (http://www.pantherdb.org) identified that

MSIL2F1/4F1 are part of a protein-protein interaction network that was principally enriched in

4 categories of GO molecular functions, namely poly(A) binding protein (GO:0008143),

mRNA 3-UTR binding (GO:0003730), single-stranded RNA binding (GO:0003727), and

translation initiation factor activity (GO:0003743), highlighting the relationship between

MSIL2/4 and the mRNA binding and regulatory proteome (Figure 3C).

Many of the proteins present in the MSIL2/4 interactome, such as PABs, PUMs,

UBP1 are involved in translation regulation and are targeted to stress granule (SGs)

components, a membraneless cytoplasmic organelle that formed consecutive to the global

inhibition of translation caused by heat-stress (Chantarachot and Bailey-Serres, 2018;

Scutenaire et al., 2018). To assess the potential functional link between MSIL and those

proteins, we crossed the MSIL2G/4G plants with a functional PAB2-RFP/PAB2R line

(Scutenaire et al., 2018), and assessed the localization of the fusion proteins before and after

heat-stress in both parent and crossed lines. Fluorescence analysis revealed that MSIL2G/4G,

like PAB2R, show a dispersed cytoplasmic signal under normal condition, and are recruited

to cytoplasmic granules that largely overlap with PAB2R-containing SG upon heat stress as

judged by the yellow-orange appearance in the merged figure (Figure 3D and Figure 3-figure

supplement 1), and by an independent analysis using the Colocalization Threshold plugin

from ImageJ software (data not shown). Treatment with cycloheximide (CHX), a translation

inhibitor that prevent SG assembly by trapping mRNAs at polysomes (Mahboubi and Stochaj,

2017), confirmed that those foci correspond to *bonafide* SGs (Figure 3E). These results were

consistent with the RNA-binding protein-based signature and the nature of MSIL

interactomes, further supporting a functional relationship between MSILs and the translation

machinery in Arabidopsis.

MSIL2/4 proteins regulate the molecular architecture of SCW in Arabidopsis fibers.

Pendant stem inflorescence is a common phenotypic trait of Arabidopsis mutants defective in

SCW biosynthesis, that results from a lack of mechanical support and rigidity (Zhang et al.,

2018; Mitsuda et al., 2007). Given the current lack of knowledge of the role of post-

transcriptional mechanisms in the control of SCW biosynthesis in plants, we henceforth

focused our analysis on the pendant stem phenotype of the *msil2/4* double mutant. Survey of

the Arabidopsis inflorescence stem tissue-specific transcriptome database (Shi et al., 2021)

indicated that the *MSIL2/4* genes are highly expressed in the SCW-forming interfascicular

fiber (*if*) and xylem vessel (*xy*) cells (Figure 4A). Toluidine blue O staining and microscopic

analysis of Col-0 and *msil2/4* mutant stem cross-sections revealed no major changes in the

architecture of both the interfascicular and xylary fibers in the mutant stem. Notably, xylem

cells deformation or collapse, which are a characteristic of mutants deficient in SCW

synthesis, were not observed in the *msil2/4* mutant (Figure 4B). However, the examination of

the cell wall thickness in cross sections of stems revealed that the interfascicular fiber cells

display a significantly thinner SCW in *msil2/4* mutant (Figure 4C). UV autofluorescence and

phloroglucinol staining of the Col-0, *msil2/4*, *msil2/4+MSIL2*F1 and *msil2/4+MSIL4*F1 stem

sections revealed a reduced lignin signal in the interfascicular fibers of the *msil2/4* mutant

compared to Col-0 or complemented plants (Figure 4D and Figure 4-figure supplement 1A).

In agreement with these observations, a significant decrease of 30% in AcBr lignin content

was observed in the *msil2/4* mutant compared to the Col-0 control plants (Figure 4E). Despite

a tendency to decrease, the difference of lignin content was not significant in the

complemented lines *msil2/4+MSIL2*F1 and *msil2/4+MSIL4*F1 compared to Col-0 (Figure 4E).

To further investigate potential changes in the lignin structure and composition in the Col-0

and *msil2/4* mutant, we evaluated the impact of MSIL2/4 deficiency on lignin composition by

thioacidolysis, a degradation method allowing the determination of the relative amounts of H,

G and S lignin units linked by beta-O-4 linkages. Total monomer yield did not significantly

differ between the Col-0 and the *msil2/4* mutant (Figure 4-figure supplement 1B), suggesting

that the structure of the lignin polymer is not significantly affected in the mutant. Further

examination of β-glucan composition using the fluorescent dye calcofluor white (Hert and

Schnepf, 1980) revealed an increased staining in the interfascicular fiber cells of *msil2/4*

mutant, supporting changes in the composition/accessibility of cellulose (Figure 4F and

Figure 4-figure supplement 1C).

To extend this analysis, we then subjected purified Col-0 and *msil2/4* SCW materials

to Fourier-Transform Infrared (FTIR) spectroscopy (Alonso-Simón et al., 2011). PLS-DA

analyses performed on FT-IR absorbance data clearly separated *msil2/4* mutant lines from

Col-0 plants, using the first two components (explaining 74% and 10% of total variability,

respectively) (Figure 4-figure supplement 1D, left panel). The large-scale chemotyping of

CW composition in stem by FTIR spectroscopy suggested that not only lignin deposition was

impaired in *msil2/4* mutants but also polysaccharides composition (Figure 4-figure

supplement 1D, right panel; Supplementary file 2). The lowest absorbance values in *msil2/4*

lines were observed in the region 1800-1500cm^{-1}, mostly associated to the absorbance of

lignin related compounds. On the contrary, a higher absorbance was detected in *msil2/4*

samples within 1050-900 cm^{-1} region, mostly associated to polysaccharides compounds

(Alonso-Simón et al., 2011). Because the optimal interaction between the SCW lignin and

polysaccharide polymers is believed to contribute to SCW recalcitrance, we used

saccharification as a proxy to compare cell wall sugars accessibility between Col-0, *msil2/4*

and complemented mutant lines (Figure 4G). A higher amount of reducing sugars from native

cell wall (without pre-treatment/-Tr) was released using *msil2/4* samples compared to Col-0

samples. After cell wall loosening using alkali pre-treatment/+Tr, an even more important

amounts of sugars (+5%) was released from double mutants CW enzymatic digestion (Figure

4G). Together, our data indicate that MSIL2/4 regulate the molecular architecture of SCW in

the interfascicular fibers, contributing significantly to the setting of biomass recalcitrance in

Arabidopsis.

The accumulation and activity of the glucuronoxylan decoration machinery are mis-regulated in *msil2/4* mutant.

To elucidate the impact of the MSIL2/4 on the expression of SCW-related biosynthesis genes in the Arabidopsis inflorescence stem, we performed mRNA sequencing (mRNA-seq) on both wild-type and *msil2/4* mutant backgrounds. In total, we found 234 genes to be significantly differentially regulated (DEG) in the *msil2/4* mutant background (with stringent thresholds having a logarithm of fold change >1.5 or <-1.5 and false discovery rate (FDR $\leq$ 0.05), including 156 up-regulated and 78 down-regulated mRNAs (Figure 5-figure supplement 1A, and Supplementary file 3). Interestingly, genes encoding regulatory or enzymatic components of the SCW biosynthesis pathway were absent from the DEG list, and their expression was not significant changed in *msil2/4* mutant (Figure 5-figure supplement 1A). Gene Ontology analysis of DEGs revealed significantly enriched signaling pathways (FDR $\leq$0.05) involved in plant defense or responses to biotic and abiotic stresses (Figure 5-figure supplement 1B), encoding products that are preferentially targeted to the extracellular region and the cell wall (Figure 5-figure supplement 1C).

The low representation of SCW biosynthesis genes in the DEG list suggests that MSIL2/4 could regulate the SCW formation at a post-transcriptional/translational level, a hallmark of MSI function in animal models (Fox et al., 2015; Kudinov et al., 2017). To investigate proteome-level changes that could potentially occur upon MSIL2/4 mutations, we performed a comparative proteomic analysis of Col-0 and *msil2/4* inflorescence stems using MS-based quantitative proteomics, and represented ≈4300 reliably identified and quantified proteins in a volcano plot according to their statistical p-value and their relative difference of abundance. This analysis revealed 267 proteins with significant abundances in the *msil2/4* mutant versus Col-0 inflorescence stems (fold change $\geq$2, Benjamini-Hochberg FDR < 1%),

including 156 up-regulated, and 111 down-regulated proteins (Figure 5A and Supplementary

file 4). While no specific GO terms were enriched in the set of down-regulated proteins, the

inspection of the up-regulated proteins revealed a strong enrichment in enzymes involved in

the glucuronoxylan biosynthetic pathway (Figure 5A and B, and Supplementary file 4).

Notably, the upregulated glucuronoxylan-related proteins included glycosyltransferases

required for the xylan backbone synthesis (IRX9/10/14), as well as enzymes involved in its

substitution (ESK1, GUX1, and GXM3) (Scheller and Ulvskov, 2010) (Figure 5A-C). Visual

inspection of the RNA-seq aligned reads of the up-regulated glucuronoxylan-related genes

using the Integrative Genomics Viewer (IGV) and quantitative RT-PCR analysis of the

glucuronoxylan-related decoration genes confirmed that the observed proteomic changes in

msil2/4 were not associated with significant variations in mRNA levels (Figure 5-figure

supplement 2A-B).

The outcomes of our histological and proteomic analyses suggest potential variations

in polysaccharide content and/or composition in the SCW of *msil2/4* mutant. To address this

point, we carried out monosaccharide composition analysis on either unpretreated (-Tr) or

acid-pretreated (+Tr) WT and *msil2/4* stem cell walls. The levels of glucose (Glc) and xylose

(Xyl), which are the building blocks of cellulose and xylan polysaccharides in the SCW were

not substantially affected in the *msil2/4* mutant (Figure 5D). Similarly, no major changes were

observed in the levels of the primary cell wall-related monosaccharides, galactose, rhamnose,

arabinose, mannose, fucose (Figure 5D), indicating that the MSIL2/4 mutations did not

significantly altered cellulose or glucuronoxylan levels in the SCW.

To assess possible changes in glucuronoxylan decoration in *msil2/4* mutant, we

performed cytochemical analysis on Col-0 and *msil2/4* stem sections using two antixylan

antibodies (LM10 and LM11), whose binding specificities depend on the degree of

glucuronoxylan substitution (McCartney et al., 2006). LM11, which binds with similar

affinity both unsubstituted and substituted glucuronoxylan, shows a similar signal in the

interfascicular and xylary fibers of Col-0 and *msil2/4* mutant stems, confirming that the

glucuronoxylan content was not significantly modified in *msil2/4* compared to Col-0 (Figure

5E). In contrast, LM10, which binds less efficiently the substituted glucuronoxylan

(McCartney et al., 2006), shows a significant signal decrease in the interfascicular fibers of

msil2/4 mutant (Figure 5E and Supplementary file 5). Together, our data indicate that the

MSIL2/4 mutations impact the accumulation and activity of the glucuronoxylan decoration

machinery in the interfascicular fiber cells of Arabidopsis stem.

MSILs restrain the degree of 4-O-methylation of glucuronoxylan in Arabidopsis.

In Arabidopsis, the glucuronoxylan backbone is decorated predominantly by acetyl, GlcA,

and MeGlcA groups, that are deposited in a regular and controlled manner by a specific

enzymatic machinery (Scheller and Ulvskov, 2010; Grantham et al., 2017; Wierzbicki et al.,

2019). Interestingly, the ESK1, GUX2, and GXM3 enzymes, which are involved in the

deposition of the acetyl and GlcA groups, as well as the methylation of GlcA group,

respectively, are over-accumulated in the *msil2/4* stem proteomics, providing a possible

explanation for the modification in glucuronoxylan decoration observed in the interfascicular

fiber cells of the *msil2/4* mutant. To investigate further the variations in glucuronoxylan

decoration incurred by the *msil2/4* mutation, we analyzed the patterns of xylanase-released

SCW oligosaccharide using matrix-assisted laser desorption ionization-time-of-flight mass

spectrometry (MALDI-TOF) (Ropartz et al., 2011). Consistent with previous observations

(Mortimer et al., 2010; Urbanowicz et al., 2012), MALDI-TOF analysis of Col-0

glucurononoxylan oligosaccharides showed the release of xylo-oligosaccharide pairs, evenly

spaced by their degrees of polymerization and acetylation, bearing a GlcA (m/z

1705.5/1747.5) or methylated GlcA (m/z 1719.5/1761.5) substitution group (Figure 6A).

Interestingly, the level of GlcA branched xylo-oligomers was strongly decreased in the

msil2/4 double mutant, and this whatever the complexity of the oligomers considered (Figure

6A and Figure 6-figure supplement 1A). Importantly, the molecular phenotypes observed in

the *msil2/4* mutant were rescued in both *msil2/4+MSIL*2F1 and *msil2/4+MSIL4*F1

complemented lines, confirming the specificity of the mutation (Figure 6A). In agreement

with the MALDI–TOF outcome, quantification of the acidic sugar levels in unpretreated (-

Tr), or acid pretreated (+Tr) WT and *msil2/4* stem samples confirmed that the release of GlcA

moiety was strongly reduced in *msil2/4* double mutant, as opposed to the galacturonic

acid/GalA group located on pectins (Figure 6B).

It has been previously proposed that the intermediate level of glucuronoxylan

methylation observed in Arabidopsis is due to an intrinsic rate-limiting activity of GXM

enzymes *in vivo* (Yuan et al., 2014). We noticed that in addition to GXM3, a second member

of the GXM family, GXM1 (Yuan et al., 2014), tends also to be over-expressed in *msil2/4*

stem although at a fold change of 1.77 slightly below the fixed significance threshold in the

proteomic analysis (Figure 5A and Supplementary file 4). This observations make us wonder

whether the apparent decrease of GlcA branched xylooligosaccharides in *msil2/4* could be due

to an overmethylation of the GlcA substituent. To address this question, we performed the

xylanase digestion and MALDI-TOF analysis in presence of an added spike-in

oligosaccharide control (Pentaacetyl-chitopentaose) for normalization of the MALDI_TOF

data (Figure 6-figure supplement 1A). Upon normalization, the MALDI-TOF data indicated

that the decrease of the peaks corresponding to the GlcA-branched xylooligosaccharides was

correlated with a increase in corresponding meGlcA branched xylooligosaccharides, (Figure

6C and Figure 6-figure supplement 1B, Supplementary file 6), suggesting that the MSIL2/4

mutations unleash the activity of the GXM1/3 in stem. RNA immunoprecipitation assays

further confirmed that the *GXM3* and *GXM1* mRNAs are efficiently pulled-down from

MSIL4G-3, but not MSIL4G^{RRM}-3 plant extracts (Figure 6D), suggesting that the MSIL2/4

interaction network controls the 4-*O*-methylation of glucuronoxylan by repressing the activity

of the GXM1/3 mRNAs in stem.

Discussion

RNA-binding proteins (RBPs) are essential components of the gene regulation machinery that

govern the fate and expression of cellular RNA at post-transcriptional levels, including

processing, splicing, base modification, translation and degradation (Singh et al., 2015).

Sequence-based bioinformatic, reverse and forward genetic, and affinity-based proteomic

analyses have converged to show that plants, like animals and fungi, harbor a large diversity

of RBPs whose functions in gene regulation and plant development are ill-defined (Reichel et

al., 2016). In this study, we have identified an RRM motif-containing RBP family, hereafter

named as MSIL, whose members share sequence, structural and functional similarities with

the animal translational regulator Musashi/MSI. We found that two of the four MSIL

members, MSIL2 and MSIL*4*, function redundantly to control various aspects of Arabidopsis

development, including leaf senescence, morphology of rosette leaves, and the rigidity of the

inflorescence stem. Our data suggest that the pleiotropy associated with the MSIL2/4

mutation likely results from the multifunctionality of these proteins in Arabidopsis

development, a situation similar to the complex phenotypic traits previously associated to

MSI mutations in animals (Fox et al., 2015; Kudinov et al., 2017; Nguyen et al., 2020).

Although the molecular bases underlying the *msil2/4* phenotypic variation remain unclear, the

presence of MSILs in the high confidence Arabidopsis mRNA-binding proteome (Reichel et

al., 2016), suggests that these proteins are likely to exert control over development by

targeting a complex repertoire of mRNAs. Regardless of the mechanism, our data support a

cell/tissue-context specific role for MSIL and provide a resource for further studies on the

mechanisms of MSIL regulation in plants.

426   Our data also provide evidence that the RRM domains of the MSIL2/4 proteins exhibit

RNA-binding ability and are essential for the activity of these proteins *in vivo*, an observation

that is corroborated by the presence of the MSIL2/4 proteins in an experimentally determined

Arabidopsis mRNA-binding proteome (Reichel et al., 2016). Further supporting a role for

MSIL2/4 in mRNA-related transactions, affinity purification-mass spectrometry analysis

revealed that these proteins share a common network of protein-protein interactions with

well-known mRNA-binding proteins, including the PAB2/4/8, RBP45/47, and the translation

factor EIF4G. Notably, the animal polyA-binding protein/PABP (homolog of the plant PABs)

was previously shown to be a functional partner of MSI that is essential for it activity in

translation repression (Fox et al., 2015; Cragle et al. 2019). Interestingly, the MSIL2/4-

associated proteins also include the phylogenetically related RBGD2/4 proteins, two

constitutively expressed RBPs that are involved in heat resistance in Arabidopsis (Zhu et al.,

2022). The comparison of the MSIL2/4 and RBGD2/4 interactomes indicates only a partial

overlap, suggesting a certain level of subfunctionalization. In accordance with this

observation, the MSILs and RBGDs clearly diversify in function, since the *rbgd2/4* double

mutant exhibits no developmental phenotypes under normal growth conditions.

442   A penetrant developmental phenotype associated with the MSIL2/4 mutations is a loss

in stem rigidity that we could trace back to a defect in SCW formation, an important reservoir

of fixed carbon and a renewable resource of major environmental and economic importance

(Meents et al., 2018). Yet, with the exception of few microRNAs involved in the control of

lignin polymer synthesis (Zhang et al., 2018; Wang et al., 2014; Zhao et al., 2015), no

information about RBPs-mediated posttranscriptional regulation of SCW biosynthetic genes

was previously reported. In this study, we show that MSIL2 and MSIL4, function redundantly

to promote SCW formation in the interfascicular fibers and that their ability to bind RNA is essential to fulfill this function. Our conclusions are supported by independent approaches which highlight changes of SCW thickness, lignin content, and xylan decoration profile in the interfascicular fiber cells of the *msil2/4* mutant. The fiber-specific effect of the MSIL2/4 mutations on SCW formation, is further supported by our evidence showing that the *msil2/4* mutant grows normally and lacks the xylem collapses usually observed in more pleiotropic SCW deficient mutants. Previous studies have shown that specific master transcriptional regulators, such as NST1 and NST3, control cell-lineage specific formation of SCW formation in Arabidopsis (Taylor-Teeples et al., 2015; Mitsuda et al., 2007). Our study extends this observation by showing that RBP-mediated post-transcriptional mechanisms are also involved in cell-type specific control of SCW formation in Arabidopsis. Notably, the presumed role of MSIL2/4 in SCW formation in the interfascicular fiber cells contrasts with the rather constitutive expression of the corresponding *MSIL2/4* genes, suggesting as previously discussed, that MSIL2/4 activities depend on cell-context. This is in accordance with the reported cell-type specific activities of the ubiquitous MSI2 in cell renewal, cell differentiation and cancer control in animals (Fox et al., 2015; Kudinov et al., 2017). Understanding how the activity of the *MSIL2/4* genes is controlled by the cellular context will be essential from a basic point of view, providing a unique model for studying RBP regulation in specific cell type in plants.

From the set of genes found to be misregulated in the *msil2/4* stem transcriptome, none could be functionally associated with SCW biosynthesis. In contrast, we found that the MSIL2/4 mutation upregulates a battery of plant defense-related genes, known to be involved in systemic acquired resistance (SAR). In this regard, the constitutive activation of defense pathways has been well documented in mutant plants defective in cell wall formation, including the SCW (Ellis et al. 2002; Miedes et al., 2014; Liu et al. 2023), an observation that

tends to support the idea that the transcriptome changes incurred by the MSIL2/4 disruption

could be indirect. This hypothesis was supported by several lines of evidence: 1) the level of

activation of defense genes in *msil2/4* is low compared to that usually observed in plants

developing SAR or in mutant plants showing an auto-activation of plant immunity (Li et al.,

2010; Wu et al., 2012); 2) SAR-dependent activation of defense genes is controlled mostly at

a transcriptional level (Hussain et al., 2018); 3) the stem phenotypes displayed by *msil2/4*

mutant have not been reported, to our knowledge, in the numerous plant-pathogen studies

published so far; 4) the top enriched proteins in the stem proteome of *msil2/4* are players of

the SCW biosynthesis pathway but not plant-defense signaling. That being said, we can't

exclude that the activation of immune-related genes, that include many cell wall-modifying

enzymes, additively contributes to the *msil2/4* stem phenotype. Further studies will be

required to clarify this issue.

Our data reveal that the accumulation of the mRNA of SCW biosynthesis enzymes

and regulators are unaffected in the *msil2/4* mutant, suggesting that the MSIL2/4 proteins act

at a translational level to regulate the SCW formation. In this regard, the stem proteomics

gives us a glimpse into the changes in protein abundance that could account for the observed

msil2/4 SCW phenotype. Indeed, our MS data reveal that proteins related to the

glucuronoxylan biosynthesis machinery, including enzymes involved both in the synthesis

(IRX9/10/14) and substitution (GUX1, ESK1, and GXM3) of the xylan backbone, are

significantly over-accumulated in the *msil2/4* mutant. Knowing that the synthesis of the

structurally complex glucuronoxylan polysaccharide requires the coordinated production and

action of these enzymes in the Golgi apparatus, our results suggests that MSIL2/4 is exerting

a multilevel control on the glucuronoxylan pathway, a mechanism reminiscent of the multi-

nodal control displays by MSI2 on the TGF-β and translation pathways in animal cells (Park

et al. 2014; Duggimpudi et al., 2018).

From a mechanistic point of view, however, our data raise several intriguing questions.

In particular, the extent of glucuronoxylan-related protein abundance changes in *msil2/4* is at

odds with the minor changes observed to glucuronoxylan composition. In particular, while the

proteomic changes can correlate with the observed increase in glucuronoxylan methylation, it

is not associated as expected with an increase in global glucuronoxylan content. One possible

explanation for this observation could be that the activities of the GUX, ESK, IRXs

glucuronoxylan biosynthesis enzymes, in contrast to that of GXM3, are not rate-limiting for

glucuronoxylan synthesis *in vivo*. Alternatively, in light with prior studies that have shown

that the steady-state levels of glucuronoxylan and cellulose are correlated within the SCW

(Wu et al., 2009; Ye and Zhong, 2022), one could imagine that mechanisms acting

downstream of glucuronoxylan *de novo* synthesis step contribute to fine tune the final content

in glucuronoxylan in the SCW. Comparing the levels of glucuronoxylan produced in the

Golgi apparatus versus the ones deposited in the SCW in both WT ve*rs*us msil2/4 plants will

be necessary to ascertain this hypothesis. In contrast, the concomitance between the

glucuronoxylan hypermethylation and the GXM3 overaccumulation phenotypes observed in

msil2/4 clearly support the previous assumption that GXM-mediated methylation is a rate-

limiting step in glucuronoxylan that can be overcome by GXM protein overexpression (Yuan

et al., 2014). In fine, our data support the idea that transcriptional and post-transcriptional

regulation act in a concerted manner to fine tune the GXM3 and glucuronoxylan methylation

levels in the stem. We currently posit that MSIL2/4 act in addition to the transcriptional

network to adjust the final level of GXM enzymes by regulating the translation of the *GXM*

mRNAs (Fig. 7). Future biochemical and functional studies will be necessary to precisely

understand the mechanism by which MSIL2/4 proteins impact gene expression during SCW

synthesis in Arabidopsis.

Our data indicate that, in addition to the reported change in glucuronoxylan

methylation, the MSIL2/4 mutations is also associated with a specific decrease of lignin

content in the SCW of the interfascicular fiber cells, whose origin remains unclear. On the

basis of the previous observations, we envision two possible mechanisms by which the

MSIL2/4 mutations could impact the lignin deposition in the fiber cells. In the first model, we

propose that MSIL2/4 positively control the lignin synthesis pathway in the interfascicular

cells by promoting the translation of lignin biosynthesis genes (Fig. 7). In this regard,

although no GO enrichment could be observed for lignin biosynthesis related genes in the

proteomic analysis, two lignin biosynthetic enzymes, CAD4 and PAL2, are significantly

down-accumulated in *msil2/4* mutant (Supplementary file 4), an observation that could

account for the observed decrease in lignin content. However, previous published biochemical

and genetic analyses do not support this conclusion, as the single *cad4* and *pal2* knock-out

mutants display no significant alteration of lignin content due to genetic redundancy (Sibout

et al., 2005; Rohde et al., 2004). Moreover, thioacidolysis analyses showed that the ratio or

degree of polymerization of the different lignin monomers remain unchanged in the *msil2/4*

mutant, reinforcing the idea that the lignin biosynthesis pathway is functional in this mutant,

leaving open the question of the specificity of MSIL2/4 action on the lignin pathway (Fig.7).

An alternative explanation for the decrease in lignin phenotype observed in the *msil2/4*

mutant would be that it results from an indirect cascade effect primarily triggered by the

msil2/4-dependent changes in glucuronoxylan methylation. In this regard, it has been

proposed that the 4-*O*-methylation of GlcA substituent results in the xylan being more

hydrophobic, possibly impacting its interaction with lignin and the capacity of lignin to

assemble in the SCW (Urbanowicz et al., 2012; Busse-Wicher et al., 2013; Kang et al., 2019;

Terret and Dupree, 2019) (Fig. 7). However, a 35S promoter-dependent overexpression study

of GXMs in Arabidopsis previously failed to report any discernable developmental or

molecular phenotypes associated to a glucuronoxylan hypermethylation (Yuan et al., 2014).

In the light of these results and despite it is necessary to elucidate deeper the role of MSIL on

other SCW genes, our work provides evidence that they are new players in the regulation of

SCW and may consequently impact biomass digestibility, with possible implications for

improving feedstock for biorefinery.

Methods

Plant material. *Arabidopsis thaliana* Col-0 and mutants *msil1-1* (*GABI_462A04*), *msil2-1*

(*Salk_066670*), *msil3-1* (*Salk_002477*) and *msil4-1* (*Salk_094167*) obtained from the SALK

Institute genomic analysis laboratory were used in this study. PAB2-RFP/PAB2R line has

been provided by Cécile Bousquet-Antonelli (LGDP, Université Perpignan). The primers

used to genotype lines are listed in Supplementary file 7. Double mutants and quadruple

mutants were generated by crossing plants. Plants were either grown in soil or cultivated *in*

vitro on plates containing 2.20 g/l synthetic Murashige and Skoog (MS) (Duchefa) medium

and 0.5g/l MES, pH 5.7, and 7 g/l agar. To breakdown dormancy, seeds were incubated for 48

h at 6°C in the dark. Germination and development were performed in growth chambers, at

20°C, 60–75% hygrometry with a 16-h light/8-h dark photoperiod (100 µE m -2 s-1 light

[fluorescent bulbs with white 6500K spectrum, purchased from Sylvania]). For *in vitro*

culture, the seeds were surface-sterilized before been sown on plates, incubated for 48 h at

6°C in the dark, and placed in a growth cabinet at 20°C with a 16-h-d/8-h-dark cycle and 120

µE m -2 s-1 light (LEDs with white 4500K spectrum, purchased from Vegeled).

Transgenic lines construction. MSIL2F- and MSIL4F-tagged versions were produced from

Phusion-generated genomic DNA PCR products (promoter and coding region) using primers

TL976(SalI)-TL924(PstI) for MSIL2F and TL974(SalI)-TL975(BamHI) for MSIL4F. PCR

products were cloned in the CTL235 binary vector containing a Flag-HA tag in front of a

NosT terminator and the hygromycin resistance gene driven by a 35S promoter. To obtain the

MSIL2G and MSIL4G constructs used for fluorescence microscopy, the same PCR products

were cloned in the CTL579 binary vector in front of a EGFP tag. The obtained plasmids were

then used to transform *mgl2/4* plants via Agrobacterium transformation. MSIL4-

GFP/MSIL4G-tagged version used for complementation and RIP corresponds to the fusion of

a genomic PCR product containing *MSIL4* promoter (primers TL3527(HindIII)-TL3528(SalI))

fused with a second PCR cDNA fragment (primers TL3529(SalI)-TL3530(BamHI)) cloned

initially in pGEM T Easy (Promega). After sequencing the fusion DNA has been cloned in the

binary vector CTL579 containing GFP cDNA sequence. The MSIL4G-tagged construct was

finally introduced in *msil2/4* plants via Agrobacterium transformation. The rrm mutated

MSIL4G^{RRM} version was produced as MSIL4G except that the cDNA fragment has been

gene-synthetized (Genecust) to modify RNA binding ability by having aspartic instead of

phenylalanine at position 9, 49 and 51 in RRM1 domain and in position 111, 151 and 153 in

RRM2 domain. Transgenic seeds selection was performed in the presence of 30 μg/l

hygromycin and the progenies were screened by PCR.

Protein extraction and immunodetection. Total plant extracts (up to 100 mg) were ground

in liquid nitrogen and proteins were extracted according to Hurkman and Tanaka, 1986.

Before migration, SDS–PAGE loading buffer was added and Coomassie staining was used to

calibrate loadings. Proteins were separated on SDS/ PAGE gels and blotted onto Immobilon-P

PVDF membrane (Cat. No. IPVH00010; MerckMillipore). Protein blot analysis was

performed using the Immobilon Western Chemiluminescent HRP Substrate (Cat. No.

WBKLS0500; MerckMillipore). Specific MSIL antibodies were raised in rabbits against

defined   epitopes   (Epitope   MSIL1   (SCDGTSSTFGYNRIPS),   MSIL2

(RLQEYFGKYGDLVE),   MSIL3   (GYGVKPEVRYSPAVGN)   and   MSIL4

(TWRSPTPETEGPAPFS) (Eurogentec).

Protein production and purification. The MSIL4 RRM binding domains were amplified

using primers TL4128 and TL4129 either from MSIL4G construct for the wild type version or

from MSIL4GΔr for the mutated version. PCR products were then cloned in pET41 to obtain

His-tagged RRM domains. Tagged proteins were produced by induction in Escherichia coli

BL21 cells. Cultures of 200 mL of an ampicillin-resistant (100 mg mL–1) colony were grown

at 37°C and induced by 100mM isopropyl-b-D-galactopyranoside in the exponential phase

(optical density 0.5 at 500 nm). After induction, bacteria were harvested by centrifugation and

pellets were resuspended in 5 ml of binding buffer (His-bind buffer kit (Millipore), proteins

were extracted using a Constant cell disrupter system (Constant Systems) with a disruption

pressure of 2.35kbar and then purified using His-bind buffer kit (Millipore) following

manufacturer's recommendations.

RNA binding assays. Binding assays were performed using homopolymer RNA conjugated

to CNBr-activated Sepharose beads (Sigma **C9142**). Either poly(A), poly(G), poly(C) or poly

(U) from Mercks (respectively P9403, P4903, P4404 and P9528) were used following

Sigma's recommendations. 0.7 to 1 mg of RNA were used for 200 mg sepharose beads. 0.45

mg of purified protein (MSIL4 RRM binding domain or mutated version) in 0.5 ml binding

buffer (100 mM NaCl, 10 mM Tris-HCl, pH 7.5, 2.5 mM MgCl2, 2.5 % Triton X-100) were

incubated with 5 mg RNA conjugated Sepharose beads for 1 hour. After incubation, beads

were washed 5 times for 10 min each in 0.5 ml of binding buffer. Beads were then pelleted at

12,000 rpm, resuspended in SDS sample buffer, boiled for 20 min, and pelleted again.

Supernatants were transferred to fresh tubes and loaded on a 12% SDS-PAGE minigel for

electrophoresis (Bio-Rad) and protein blot analysis.

RNA immunoprecipitation. RIP was performed as described previously in Merret et al.,

2017) using 400 mg of stem powder as starting material.

Expression analysis. Total RNA was isolated using the TRI re-agent (Cat. No. TR-118;

Euromedex) according to the manufacturer's instructions. Genomic DNA was then digested

out using the RQ1 RNase-Free DNase (Cat. No. M6101; Promega). cDNAs were obtained

from 400 ng of RQ1-treated RNA using 1 µl of GoScript Reverse Transcriptase (Cat. No.

A5003; Promega) in a 20-µl final volume reaction using random primers (Cat. No. C1181;

Promega) in the presence of 20 units of RNasin (Recombinant Ribonuclease Inhibitor, Cat.

No. C2511; Promega) and dNTP mix at a final concentration of 0.5 mM of each dNTP (Cat.

No. U1511; Promega). Semi-quantitative RT–PCR amplifications were performed on 1 µl of

cDNA in a 12.5-µl reaction volume to start with. The amplification of EF1αtranscripts was

used to equilibrate. qRT–PCR was performed on a Light Cycler 480 II machine (Roche

Diagnostics) by using the Takyon No ROX SYBR MasterMix blue dTTP kit (Cat. No. UF-

NSMT-B0701; Eurogentec). Each amplification reaction was set up in a 10-µl reaction

containing each primer at 300 nM and 1 µl of RT template in the case of total RNA with a

thermal profile of 95°C for 10 min and 40 amplification cycles of 95°C, 15 s; 60°C, 60 s.

Relative transcript accumulation was calculated using the ΔΔCt methodology (Livak and

Schmittgen, 2001), using ACTIN2 as internal control. Average ΔΔCt represents three

experimental replicates with standard errors. PolyA+ RNAs were purified from RNA

obtained by the TRI Reagent extraction, using the PolyATtract mRNA Isolation System III

(Cat. No. Z5310; Promega) according to the manufacturer's instructions. cDNAs were then

synthetized according to the above protocol using 50 ng of polyA+ RNA as starting material.

Semiquantitative and quantitative RT–PCRs were performed as described above with a 1/25

dilution of cDNAs from polyA+ RNA as template for qRT–PCR. The primers used for RT

and qRT-PCR are listed in Supplementary file 7.

RNA Sequencing. Total RNA were extracted from the basal stem section of mature Col-0

and *msil2/4* mutant plants using the RNeasy Plant Mini Kit (Qiagen, Cat.#74904) and treated

onto the column with the RNase-Free DNase Set (Qiagen, Cat.#79254). Two replicates were

performed per plant lines. PolyA purification, library preparation using the Illumina TruSeq

stranded mRNA kit and library quality controls were performed by the BioEnvironment

Illumina Platform. The mRNA sequencing of Col-0 and msil2/4 mRNAs was done with

single-reads, 1x 75pb and $30–35x10^6$ reads per library.

RNASeq analysis. For each library, $30–35x10^6$ reads were obtained with 85–90% of the

bases displaying a Q-score $\geq$ 30, with a mean Q-score of 38 as assessed with Fastqc

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Reads were mapped against the

TAIR 10 genome using gtf annotation file and default parameters of TopHat2 v2.0.7

(Langmead and Salzberg, 2012). Assembly and transcript quantification were performed with

Cufflinks v2.2.1 (Kim et al., 2013). Finally, the low expressed transcripts - less than 1 RPKM

(reads per kilobase per million mapped reads) in one of the libraries were filtered out. The

differential analysis was conducted by Cuffdiff software, belonging to the Trapnell suite.

Foldchange (FC) was determined as the ratio between the normalized read counts between

mutant and wild-type.

Fluorescence detection and tissue colorations. GFP detections were performed on *in vitro*

grown one week seedling roots fixed in fixation buffer (50mM Pipes, 5mM EGTA, 5mM

MgSO4, pH7 added with 2% para-formaldehyde and 0.2% Triton). After 5 min vacuum

infiltration and 10 min agitation at room temperature, roots were washed twice with fixation

buffer and stored at 4°C until confocal observation. Confocal imaging was done using an

Zeiss Axio Observer Z1 LSM 700. Roots were observed in water and for GFP detection

samples were excited at 488 nm and the signal captured at 495–555 nm. For RFP detection,

excitation was performed at 555 nm and the signal captures at 600-700 nm. Images were

analysed with ImageJ. At least 15 samples per line were analyzed.

Stem coloration were performed from 5 cm length basal stem pieces conserved in

ethanol 80%. Stems were rehydrated first in ethanol 50% and later in water and then

embedded in agarose 5% before being transversally cut using a vibratome (Leica vt 1000s) to

obtain 120 micron thickness sections stored in water. The presence of lignin was determined

by staining with phloroglucinol for 30 seconds, giving a red coloration in the presence of

lignin cinnamaldehyde groups. Sections were observed using an inverted microscope (Leitz

DMRIBE, Leica Micro-systems, Wetzlar, Germany) and images were registered using a CCD

camera (Color Collview, Photonic Science, Milham, UK).

For cellulose detection, the brightener Calcofluor White ST [4,4'-bis(anilino-6-bis(2-

hydroxyethyl)-amino-s-triazine-ylamino)-2,2'-stilbene disulfonic acid; Cyanamide Co.,

Bound Brook, N.J] was used at a final concentration of 0.1%. The samples were incubated

with calcofluor for 3 min, washed three times with water before being observed. For

detection, excitation was performed at 405 nm and the signal captures at 434-496 nm. For

auto-fluorescence analysis stem sections were excited at 405 nm and the signal captures at

410-470 nm for blue. Pictures have been analyzed with LAS X (Leica Application Suite X).

Biochemical analyses of SCWs. Arabidopsis stems (5 cm sections at the bottom part of the

stems) were harvested from Col0 and *msil2/4* double mutant as four independent biological

replicates. Soluble extractives were eliminated as described by Ployet et al. (Ployet et al.,

2018). Briefly, stems were freeze dried for 48h, ground using a Mixer Mill MM 400 (Retsch),

and extracted by hot solvents (successively water, ethanol, ethanol/toluene (1:1 v/v) and

acetone) to obtain extractive-free stem residues (ESR). The determination of acetyl bromide

(AcBr) lignin was performed on 10mg of ESR as described by Ployet et al. (Ployet et al.,

2019). All analyses were done in triplicate. Fourier transform infrared spectroscopy (FT-IR)

analysis was performed on 100-200 mg of ESR. Spectra were recorded from ten technical

replicates, in the range 400–4000 cm−1 with a 4 cm-1 resolution and 32 scans per spectrum

using an attenuated total reflection (ATR) Nicolet IS50 FT-IR spectrometer (Thermo Fisher,

Illkirch-Graffenstaden, France) equipped with a deuterated-triglycine sulfate (DTGS) detector.

Spectra analyses were performed as described in Dai et al., 2020. Saccharification yield was

estimated with or without alkali pretreatment, as described by Van Acker et al., 2016.

Reducing sugar concentration was assessed with dinitrosalicylic acid (DNS) reagent using

10µl of the supernatant after 9 h of reaction. Enzyme activity was assessed at 0.25 filter paper

units (FPU) ml^{-1} in our conditions. The Boxplots were generated using ggplot2 R package (R

v2022.07.1).

Mass spectrometry (MS)-based proteomic analyses. Proteins from total inflorescence stem

extracts of three biological replicates of Col-0 and *msil2/4* mutant plants were solubilized in

Laemmli buffer and heated for 10 min at 95°C. They were then stacked in the top of a 4-12%

NuPAGE gel (Invitrogen), stained with Coomassie blue R-250 (Bio-Rad) before in-gel

digestion using modified trypsin (Promega, sequencing grade) as previously described

(Casabona et al., 2013). The resulting peptides were analyzed by online nanoliquid

chromatography coupled to MS/MS (Ultimate 3000 RSLCnano and Orbitrap Exploris 480,

Thermo Fisher Scientific) using a 180-min gradient. For this purpose, the peptides were

sampled on a precolumn (300 µm x 5 mm PepMap C18, Thermo Scientific) and separated in

a 75 µm x 250 mm C18 column (Aurora Generation 2, 1.6µm, IonOpticks). The MS and

MS/MS data were acquired using Xcalibur 4.4 (Thermo Fisher Scientific).

Peptides and proteins were identified by Mascot (version 2.8.0, Matrix Science)

through concomitant searches against the Viridiplantae database (from Uniprot, Arabidopsis

thaliana (thale-cress) taxonomy, 136447 sequences) and a homemade database containing the

sequences of classical contaminant proteins found in proteomic analyses (human keratins,

trypsin..., 126 sequences). Trypsin/P was chosen as the enzyme and two missed cleavages

were allowed. Precursor and fragment mass error tolerances were set at respectively at 10 and

20 ppm. Peptide modifications allowed during the search were: Carbamidomethyl (C, fixed),

Acetyl (Protein N-term, variable) and Oxidation (M, variable). The Proline software

(Bouyssié et al., 2020) (version 2.2.0) was used for the compilation, grouping, and filtering of

the results (conservation of rank 1 peptides, peptide length $\geq$ 6 amino acids, false discovery

rate of peptide-spectrum-match identifications < 1% (Couté et al., 2020), and minimum of

one specific peptide per identified protein group). MS data have been deposited to the

ProteomeXchange Consortium via the PRIDE partner repository (Perez-Riverol et al., 2022).

Proline was then used to perform a MS1 label-free quantification of the identified protein

groups based on razor and specific peptides.

Statistical analysis was performed using the ProStaR software (Wieczorek et al., 2017)

based on the quantitative data obtained with the four biological replicates analyzed per

condition. Proteins identified in the contaminant database, proteins identified by MS/MS in

less than two replicates of one condition, and proteins quantified in less than three replicates

of one condition were discarded. After log2 transformation, abundance values were

normalized using the variance stabilizing normalization (vsn) method, before missing value

imputation (SLSA algorithm for partially observed values in the condition and DetQuantile

algorithm for totally absent values in the condition). Statistical testing was conducted with

limma, whereby differentially expressed proteins were selected using a log2(Fold Change)

cut-off of 1 and a p-value cut-off of 0.00513, allowing to reach a false discovery rate close to

1% according to the Benjamini-Hochberg estimator. Proteins found differentially abundant

but identified by MS/MS in less than two replicates, and detected in less than three replicates,

in the condition in which they were found to be more abundant were invalidated (p-value = 1).

MALDI-TOF MS analyses of the patterns of xylanase-released SCW oligosaccharide. 10

mg of dried alcohol insoluble cell wall residues were incubated with 10U.ml^{-1} of endo-1,4-D-

xylanase (E-XYLNP) from MEGAZYME previously desalted with centrifulgal filters

(Amicon Ultra-0.5ml) in 1 ml during 12hrs at 40°C. After boiling during 5 min to inactivate

enzyme activity, samples where filtred with 50KD centrifulgal filters (Amicon Ultra-0.5ml).

As positive control, 2 mg of wheat AX (MEGAZYMZE) were treated in a similar manner.

The samples were then analyzed by matrix-assisted laser desorption ionization-time of flight

(MALDI-TOF) MS using DMA/DHB (N,N-dimethylaniline/2,5-dihydroxybenzoic acid)

matrix (Ropartz et al., 2011). The samples (1 µL) were deposited and then covered by the

matrix (1 µL) on a polished steel MALDI target plate. MALDI measurements were then

performed on a rapifleX MALDI-TOF spectrometer (Bruker Daltonics, Bremen, Germany)

equipped with a Smartbeam 3D laser (355 nm, 10000 Hz) and controlled using the Flex

Control 4.0 software package. The mass spectrometer was operated with positive polarity in

reflectron mode. Spectra were acquired in the range of 560–3200 *m/z*. The detected species

correspond to ions in the form of sodium adducts.

Thioacidolysis. This method is adapted from Lapierre et al., 1995. Briefly, 10 mg of alcohol

insoluble cell wall residues were incubated in 3ml of dioxane with ethanethiol (10%), BF3

etherate (2.5%) containing 0.1% of heinecosane C21 diluted in CH2Cl2 at 100°C during 4

hours. 3ml of NaHCO3 (0.2 M) were added after cooling and mixed prior the addition of 0.1

mL of HCl (6 M). The tubes were wortexed after addition of 3 mL of dichloromethane and

the whole lower organic phase was collected in a new tube before concentration under

nitrogen atmosphere to approximately 0.5 ml. Then, 10 μL were trimethylsilylated (TMS)

with 100 μL of N,O-bis(trimethylsilyl) trifluoroacetamide and 10 μL of ACS-grade pyridine.

The trimethylsilylated samples were injected (1 μL) onto an Agilent 5973 Gas

Chromatography–Mass Spectrometry system. Specific ion chromatograms reconstructed at

m/z 239, 269 and 299 were used to quantify H, G and S lignin monomers respectively and

compared to the internal standard at m/z 57, 71, 85).

Cell wall sugar analysis. Neutral monosaccharide sugar content was determined by gas

chromatography after acid hydrolysis and conversion of monomers into alditol acetates as

described in Hoebler et al., 1989, and Blakeney et al., 1983. Gas Chromatography was

performed on a DB 225 capillary column (J&W Scientific, Folsorn, CA, USA; temperature

205 °C, carrier gas H_2). Calibration was made with standard sugar solution and inositol as

internal standard.

For acid sugar analysis, the uronic acid content was automatically quantified after

hydrolysis of polysaccharides in concentrated sulfuric acid (18 M) containing sodium

tetraborate (12.5 mM) or not, followed by m-hydroxydiphenyl colorimetric determination

(Blumenkrantz et al., 1973; Thibault, 1979). Galacturonic and Glucuronic acid were used as

standard for quantification.

Immunohistochemistry. Samples of Arabidopsis tissues were dehydrated in a successive

ethanol series (20%, 40%, 60%, 80%, 95%, and 100%) and embedded in LR White resin

(Electron Microscopy Sciences). Thin sections (1 μm) were placed on Teflon-coated slides

(Electron Microscopy Sciences), blocked in phosphate-buffered saline, 2% Tween, and 1%

bovine serum albumin for 2 h (PBST-BSA), and labeled overnight (12 h) at 4 °C with primary

antibody diluted in PBST-BSA. Sections were washed with PBST and incubated at room

temperature for 2 h with a secondary antibody diluted in PBST-BSA. Slides were then

washed with deionized water and dried under a stream of dry air. Primary antibodies and

dilutions were as follows:, LM10 and LM11 (1:10, v/v; Plant Probes for both). The secondary

antibody was a goat anti-rat IgG coupled to the fluorescent dye Alexa Fluor 555 (Molecular

Probes) and was used at a 1:100 (v/v) dilution. Observations were carried out using a

LeicaDM 6000microscope with a TCS SP8 scanning confocal system using a 40 x

apochromatic oil immersion objective to visualized Alexa555 fluorescence (excitation: 561

nm; emission: 565–645 nm). Acquisition settings, including laser power,photomultiplier gain,

field of view (X,Y dimension), Z-step, and pixel size of the image, were strictly identical to

ensure reliable comparisons between plant material (i.e. the wild type versus mutants).

Data availability

The high-throughput sequencing data generated in this study are accessible at NCBI's Gene

Expression Omnibus (GEO) via GEO Series accession number GSE223732. The mass

spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via

the PRIDE partner repository with the dataset identifier PXD040207 (proteomic identification

of MSIL-interacting partners), and PXD040020 (proteomic analysis of total inflorescence

stem extracts).

References

Alonso-Simón A, Garcia-Angulo P, Mélida H, Encina A, Álvarez JM, Acebes JL (2011) **The**

use of FTIR spectroscopy to monitor modifications in plant cell wall architecture caused

by cellulose biosynthesis inhibitors *Plant Signaling & Behavior* **6**, 1104-1110 (2011).

Biasani M, Bienert S, Waterhouse A, Arnold K, Studer G, Schmidt T, Kiefer F, Gallo

Cassarino T, Bertoni M, Bordoli L, Schwede T (2014) **SWISS-MODEL: modelling protein**

tertiary and quarternary structure using evolutionary information *Nucleic Acids*

Research **42**: W252-258.

Blakeney AB, Harris PJ, Henry RJ, Stone BA (1983) **A simple and rapid preparation of**

alditol acetates for monosaccharide analysis *Carbohydrate Research* **113**: 291–299.

Blumenkrantz N, Asboe-Hansen G (1973) **New method for quantitative determination of**

uronic acids *Analytical Biochemistry* **54**: 484-489.

Bonawitz ND, Chapple C (2013) **Can genetic engineering of lignin deposition be**

accomplished without an unacceptable yield penalty? *Current Opinion in Biotechnology*

24, 336-343.

Bouyssié D, Hesse AM, Mouton-Barbosa E, Rompais M, Macron C, Carapito C, Gonzalez de

Peredo A, Couté Y, Dupierris V, Burel A, Menetrey JP, Kalaitzakis A, Poisat J, Romdhani A,

Burlet-Schiltz O, Cianférani S, Garin J, Bruley C (2020) **Proline: an efficient and user-**

friendly software suite for large-scale proteomics *Bioinformatics* **36**: 3148-3155.

Busse-Wicher M, Li A, Silveira RL, Pereira CS, Tryfona T, Gomes TC, Skaf MS, Dupree P

(2013) **Evolution of xylan substitution patterns in gymnosperms and angiosperms:**

implications for xylan interaction with cellulose *Plant Physiology* **171:** 2418-2431.

Casabona MG, Vandenbrouck Y, Atree I, Couté Y (2013) **Proteomic characterization of**

Pseudomonas aeruginosa PAO1 inner membrane *Proteomics* **13**: 2419-2423.

Chai G, Kong Y, Zhu M, Yu L, Qi G, Tang X, Wang Z, Cao Y, Yu C, Zhou G (2015)

Arabidopsis C3H14 and C3H15 have overlapping roles in the regulation of secondary

wall thickening and anther development *Journal of Experimental Botany* **66**: 2595-2609.

Chantarachot T, Bailey-Serres J (2018) **Polysomes, stress granules, and processing bodies:**

a dynamic triumvirate controlling cytoplasmic mRNA fate and function *Plant Physiology*

176: 254-269.

Chavali PL, Stojic L, Meredith LW, Joseph N, Nahorski MS, Sanford TJ, Sweeney TR,

Krishna BA, Hosmillo M, Firth AE, Bayliss R, Marcelis CL, Lindsay S, Goodfellow I,

Woods CG, Gergely F (2017) **Neurodevelopmental protein Musashi-1 interacts with the**

Zika genome and promotes viral replication *Science* **357**: 83-88.

Couté Y, Bruley C, Burger T (2020) **Beyond target-decoy competition: stable validation of**

peptide and protein identifications in mass spectrometry-based discovery proteomics

Analytical Chemistry **92:** 14898-14906.

Cragle CE, MacNicol MC, Byrum SD, Hardy LL, Mackintosh SG, Richardson WA, Gray NK,

Childs GV, Tackett AJ, MacNicol AM (2019) **Musashi interaction with poly(A)-binding**

protein is required for activation of target mRNA translation *Journal of Biological*

Chemistry **294:** 10969-10986.

Dai Y, Hu G, Dupas A, Medina L, Blandels N, Clemente HS, Ladouce N, Badawi M,

Hernandez-Raquet G, Mounet F, Grima-Pettenati J, Cassan-Wang H. Dai, Y (2020)

Implementing the CRISPR/Cas9 technology in Eucalyptus hairy roots using wood-

related genes *International Journal of Molecular Sciences* **21:** 3408.

Daubner GM, Cléry A, Allain FH (2013) **RRM-RNA recognition: NMR or**

crystallography… and new findings *Current Opinion in Structural Biology* **23:**100-108.

Weill L, Belloc E, Castellazzi CL, Mendez R (2017) **Musashi 1 regulates the timing and**

extent of meiotic mRNA translational activation by promoting the use of specific CPEs

Nature Structural and Molecular Biology **24**: 672-681.

Duggimpudi S, Kloetgen A, Maney SK, Münch PC, Hezaveh K, Shaykhalishahi H, Hoyer W,

McHardy AC, Lang PA, Borkhardt A, Hoell JI (2018) **Transcriptome-wide analysis**

uncovers the targets of the RNA-binding protein MSI2 and effects of MSI2's RNA-

binding activity on IL-6 signaling *Journal of Biological Chemistry* **293**: 15359-15369.

Eliss C, Karafyllidis I, Wasternack C, Turner JG (2002) **The Arabidopsis cev1 links cell**

wall signaling to jasmonate and ethylene responses *The plant cel* **14**:1557-1566.

Fox RG, Park FD, Koechlein CS, Kritzik M, Reya T (2015) **Musashi signaling in stem cells**

and cancers *Annual Review of Cell and Developmental Biology* **31**: 249-267.

Grantham NJ, Wurman-Rodrich J, Terrett OM, Lyczakowski JJ, Stott K, Iuga D, Simmons TJ,

Durand-Tardif M, Brown SP, Dupree R, Busse-Wicher M, Dupree P (2017) **An even pattern**

of xylan substitution is critical for interaction with cellulose in plant cell walls *Nature*

Plants **11**: 859-865.

Herth W, Schnepf E (1980) **The fluorochrome, calcofluor white, binds oriented to**

structural polysaccharide fibrils *Protoplasma* **105**: 129-133.

Hoebler C, Barry J-L, David A, Delort-Laval J (1989) **Rapid acid hydrolysis of plant cell**

wall polysaccharides and simplified quantitative determination of their neutral

monosaccharides by gas-liquid chromatography *Journal of Agricultural and Food*

Chemistry **37:** 360–365.

Hurkman WJ, Tanaka CK (1986) **Solubilization of plant membrane proteins for analysis**

by two-dimensional gel electrophoresis Plant Physiology 81: 802-806.

Hussain RMF, Sheikh AH, Haider I, Quareshy M, Linthorst HJM (2018) **Arabidopsis**

WRKY50 and TGA transcription factors synergistically activate expression of PR1

Frontiers in Plant Science **9**:930.

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K,

Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A,

Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E,

Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O,

Senior AW, Kavukcuoglu K, Kohli P, Hassabis D (2021) **Highly accurate protein structure**

prediction with AlphaFold *Nature* **596**:583-589.

Kang X, Kirui A, Dickwella Widanage MC, Mentink-Vigier F, Cosgrove DJ, Wang T (2019)

Lignin-polysaccharide interactions in plant secondary cell walls revealed by solid-state

NMR *Nature Communications* **10:** 347.

Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL (2013) **TopHat2: accurate**

alignment of transcriptomes in the presence of insertions, deletions and gene fusions

Genome Biology **14:** R36.

Kudinov AE, Karanicolas J, Golemis EA, Boumber Y (2017) **Musashi RNA-binding**

proteins as cancer drivers and novel therapeutic targets *Clinical Cancer Research* **23**:

2143-2153.

Kumar M, Campbell L, Turner S (2016) **Secondary cell walls: biosynthesis and**

manipulation *Journal of Experimental Botany* **67**: 515-531.

Langmead B, Salzberg SL (2012) **Fast gapped-read alignment with Bowtie 2** *Nature*

Methods **9:** 357-359.

Lapierre C, Pollet B, Rolando C (1995) **New insights into the molecular architecture of**

hardwood lignins by chemical degradative methods *Research on Chemical Intermediates*

21: 397-412.

Li Y, Li S, Bi D, Cheng YT, Li X, Zhang Y (2010) **SRF1 negatively regulates plant NB-**

LRR resistance protein accumulation to prevent autoimmunity *PloS pathogens*

6:e1001111.

Liu C, Yu H, Voxeur A, Rao X, Dixon RA (2023) **FERONIA and wall-associated kinases**

coordinate defense induced by lignin modification in plant cell walls *Science Advances*

9:eadf7714.

Livak KJ, Schmittgen TD (2001) **Analysis of relative gene expression data using real-time**

quantitative PCR and the 2(-Delta Delta C(T)) method *Methods* **25:**402-408.

Mahboubi H, Stochaj U (2017) **Cytoplasmic stress granules: Dynamic modulators of cell**

signaling and disease *Biochimica et Biophysica Acta Molecular Basis of Disease* **1863:** 884-

895.

McCartney L, Blake AW, Flint J, Bolam DN, Boraston AB, Gilbert HJ, Knox JP (2006)

Differential recognition of plant cell walls by microbial xylan-specific carbohydrate-

binding modules *Proceeding of the National Academy of Sciences USA* **103:** 4765-4770.

Meents MJ, Watanabe Y, Samuels AL (2018) **The cell biology of secondary cell wall**

biosynthesis *Ann Bot* **121**: 1107-1125.

Merret R, Carpentier MC, Favory JJ, Picart C, Descombin J, Bousquet-Antonelli C, Tillard P,

Lejay L, Deragon JM, Charng YY (2017) **Heat shock protein HSP101 affects the release of**

ribosomal protein mRNAs for recovery after heat shock *Plant Physiology* **174**: 1216-1225.

Miedes E, Vanholme R, Boerjan W, Molina A (2014) **The role of the secondary cell wall in**

plant resistance to pathogens *Frontiers in Plant Science* **5**: 358.

Mitsuda N, Iwase A, Yamamoto H, Yoshida M, Seki M, Shinozaki K, Ohme-Takagi M (2007)

NAC transcription factors, NST1 and NST3, are key regulators of the formation of

secondary walls in Woody tissues of Arabidopsis The *Plant Cell* **19**: 270-280.

Mortimer JC, Miles GP, Brown DM, Zhang Z, Segura MP, Weimar T, Yu X, Seffen KA,

Stephens E, Turner SR, Dupree P (2010) **Absence of branches from xylan in Arabidopsis**

***gux* mutants reveals potential for simplification of lignocellulosic biomass** *Proceeding of*

the National Academy of Sciences USA **107:** 17409-17414.

Myburg AA, Grattapaglia D, Tuskan GA, Hellsten U, Hayes RD, Grimwood J, Jenkins J,

Lindquist E, Tice H, Bauer D, Goodstein DM, Dubchak I, Poliakov A, Mizrachi E, Kullan

AR, Hussey SG, Pinard D, van der Merwe K, Singh P, van Jaarsveld I, Silva-Junior OB,

Togawa RC, Pappas MR, Faria DA, Sansaloni CP, Petroli CD, Yang X, Ranjan P,

Tschaplinski TJ, Ye CY, Li T, Sterck L, Vanneste K, Murat F, Soler M, Clemente HS, Saidi

N, Cassan-Wang H, Dunand C, Hefer CA, Bornberg-Bauer E, Kersting AR, Vining K,

Amarasinghe V, Ranik M, Naithani S, Elser J, Boyd AE, Liston A, Spatafora JW,

Dharmwardhana P, Raja R, Sullivan C, Romanel E, Alves-Ferreira M, Külheim C, Foley W,

Carocha V, Paiva J, Kudrna D, Brommonschenkel SH, Pasquali G, Byrne M, Rigault P,

Tibbits J, Spokevicius A, Jones RC, Steane DA, Vaillancourt RE, Potts BM, Joubert F, Barry

K, Pappas GJ, Strauss SH, Jaiswal P, Grima-Pettenati J, Salse J, Van de Peer Y, Rokhsar DS,

Schmutz J (2014) **The genome of Eucalyptus grandis** *Nature* **510**: 356-362.

Nguyen DTT, Lu Y, Chu KL, Yang X, Park SM, Choo ZN, Chin CR, Prieto C, Schurer A,

Barin E, Savino AM, Gourkanti S, Patel P, Vu LP, Leslie CS, Kharas MG (2020)

HyperTRIBE uncovers increased MUSASHI-2 RNA binding activity and differential

regulation in leukemic stem cells *Nature Communications* **11**: 2026.

Park SM, Deering RP, Lu Y, Tivnan P, Lianoglou S, Al-Shahrour F, Ebert BL, Hacohen N,

Leslie C, Daley GQ, Lengner CJ, Kharas MG (2014) **Musashi-2 controls cell fate, lineage**

bias, and TGF-b signailing in HSCs *The Journal of Experimental Medecine* **211**:71-87.

Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S, Kamatchinathan S,

Kundu DJ, Prakash A, Frericks-Zipper A, Eisenacher M, Walzer M, Wang S, Brazma A,

Vizcaíno JA (2022) **The PRIDE database and related tools and resources in 2019:**

improving support for quantification data *Nucleic Acids Research* **47:** D442-D450.

Ployet R, Soler M, Carocha V, Ladouce N, Alves A, Rodrigues JC, Harvengt L, Marque C,

Teulières C, Grima-Pettenati J, Mounet F (2018) **Long cold exposure induces**

transcriptional and biochemical remodelling of xylem secondary cell wall in Eucalyptus

Tree Physiology **38**: 409-422.

Ployet R, Veneziano Labate MT, Regiani Cataldi T, Christina M, Morel M, San Clemente H,

Denis M, Favreau B, Tomazello Filho M, Laclau JP, Labate CA, Chaix G, Grima-Pettenati J,

Mounet F (2019) **A systems biology view of wood formation in Eucalyptus grandis trees**

submitted to different potassium and water regimes *The New Phytologist* 223: 766-782.

Reichel M, Liao Y, Rettel M, Ragan C, Evers M, Alleaume AM, Horos R, Hentze MW,

Preiss T, Millar AA (2016) **In planta determination of the mRNA-binding proteome of**

Arabidopsis etiolated seedlings The *Plant Cell* **28**: 2435-2452.

Rohde A, Morreel K, Ralph J, Goeminne G, Hostyn V, De Rycke R, Kushnir S, Van

Doorsselaere J, Joseleau JP, Vuylsteke M, Van Driessche G, Van Beeumen J, Messens E,

Boerjan W (2004) **Molecular phenotyping of the pal1 and pal2 mutants of Arabidopsis**

thaliana reveals far-reaching consequences on phenylpropanoid, amino acid, and

carbohydrate metabolism *The Plant cell* **16**: 2749-2771.

Ropartz D, Bodet PE, Przybylski C, Gonnet F, Daniel R, Fer M, Helbert W, Bertrand D,

Rogniaux H (2011) **Performance evaluation on a wide set of matrix-assisted laser**

desorption ionization matrices for the detection of oligosaccharides in a high-throughput

mass spectrometric screening of carbohydrate depolymerizing enzymes *Rapid*

Communications in Mass Spectrometry **25:** 2059-2070.

Scheer H, de Almeida C, Ferrier E, Simonnot Q, Poirier L, Pflieger D, Sement FM, Koechler

S, Piermaria C, Krawczyk P, Mroczek S, Chicher J, Kuhn L, Dziembowski A, Hammann P,

Zuber H, Gagliardi D (2021) **The TUTase URT1 connects decapping activators and**

prevents the accumulation of excessively deadenylated mRNAs to avoid siRNA

biogenesis *Nature Communications* **12:** 1298.

Scheller HV, Ulvskov P (2010) **Hemicelluloses** *Annual Review in Plant Biology* **61:** 263-289.

Scutenaire J, Deragon JM, Jean V, Benhamed M, Raynaud C, Favory JJ, Merret R, Bousquet-

Antonelli C (2018) **The YTH domain protein ECT2 is an m^{6}A reader required for**

normal trichome branching in Arabidopsis *The Plant Cell* **30**: 986-1005.

Shi D, Jouannet V, Agustí J, Kaul V, Levitsky V, Sanchez P, Mironova VV, Greb T (2021)

Tissue-specific transcriptome profiling of the Arabidopsis inflorescence stem reveals

local cellular signatures The *Plant Cell* **33:** 200-223.

Sibout R, Eudes A, Mouille G, Pollet B, Lapierre C, Jouanin L, Séguin A (2005)

CINNAMYL ALCOHOL DEHYDROGENASE-C and -D are the primary genes

involved in lignin biosynthesis in the floral stem of Arabidopsis *Plant Cell* **17**: 2059-2076.

Singh G, Pratt G, Yeo GW, Moore MJ (2015) **The clothes make the mRNA: past and**

present trends in mRNP fashion *Annual Review of Biochemistry* **84**, 325-354.

Taylor-Teeples, M., Lin L., de Lucas M, Turco G, Toal TW, Gaudinier A, Young NF,

Trabucco GM, Veling MT, Lamothe R, Handakumbura PP, Xiong G, Wang C, Corwin J,

Tsoukalas A, Zhang L, Ware D, Pauly M, Kliebenstein DJ, Dehesh K, Tagkopoulos I, Breton

G, Pruneda-Paz JL, Ahnert SE, Kay SA, Hazen SP, Brady SM (2015) **An Arabidopsis gene**

regulatory network for secondary cell wal synthesis *Nature* **517**: 571-575.

Terret OM, Dupree P (2019) **Covalent interactions between lignin and hemicelluloses in**

plant secondary cell walls *Current Opinion in Biotechnology* **56:** 97-104.

Thibault JF (1979) **Automated-method for the determination of pectic substances**

Lebensmittel-Wissenschaft Und-Technologie **12:** 247-251.

Urbanowicz BR, Peña MJ, Ratnaparkhe S, Avci U, Backe J, Steet HF, Foston M, Li H,

O'Neill MA, Ragauskas AJ, Darvill AG, Wyman C, Gilbert HJ, York WS (2012) **4-O-**

methylation of glucuronic acid in Arabidopsis glucuronoxylan is catalyzed by a domain

of unknown function family 579 protein *Proceeding of the National Academy of Sciences*

USA **109**, 14253-14258.

Van Acker R, Vanholme R, Piens K (2016) **Saccharification protocol for small-scale**

lignocellulosic biomass samples to test processing of cellulose into glucose *Bio-Protocol*. **6**:

1701.

Wang CY, Zhang S, Yu Y, Luo YC, Liu Q, Ju C, Zhang YC, Qu LH, Lucas WJ, Wang X,

Chen YQ (2014) **MiR397b regulates both lignin content and seed number in Arabidopsis**

via modulating a laccase involved in lignin biosynthesis *Plant Biotechnology Journal* **12**:

1132-1142.

Wang Y, Fan C, Hu H, Li Y, Sun D, Wang Y, Peng L (2016) **Genetic modification of plant**

cell walls to enhance biomass yield and biofuel production in bioenergy crops

Biotechnology Advances **34**: 997-1017.

Weill L, Belloc E, Castellazzi CL, Mendez R (2017) **Musashi 1 regulates the timing and**

extent of meiotic mRNA translational activation by promoting the use of specific CPEs.

Nature Structural and Molecular Biology **24:** 672-681.

Whitehead C, Ostos Garrido FJ, Reymond M, Simister R, Distelfeld A, Atienza SG, Piston F,

Gomez LD, McQueen-Mason SJ (2018) **A glycosyl transferase family 43 protein involved**

in xylan biosynthesis is associated with straw digestibility in *Brachypodium distachyon*

The *New Phytologist* **218**: 974-985.

Wieczorek S, Combes F, Lazar C, Giai Gianetto Q, Gatto L, Dorffer A, Hesse AM, Couté Y,

Ferro M, Bruley C, Burger T (2017) **DAPAR & ProStaR: software to perform statitical**

analyses in quantitative discovery proteomics *Bioinformatics* **33:** 135-136.

Wierzbicki MP, Maloney V, Mizrachi E, Myburg AA (2019) **Xylan in the middle:**

understanding xylan biosynthesis and its metabolic dependencies toward improving

wood fiber for industrial processing *Frontiers in Plant Science* **25:** 176.

Wu AM, Rihouey C, Seveno M, Hörnblad E, Singh SK, Matsunaga T, Ishii T, Lerouge P,

Marchant A (2009) **The Arabidopsis IRX10 and IRX10-LIKE glycosyltransferases are**

critical for glucuronoxylan biosynthesis during secondary cell wall formation *Plant*

Journal **57**: 718-731.

Wu Y, Zhang D, Chu JY, Boyle P, Wang Y, Brindle ID, De Luca V, Després C (2012) **The**

Arabidopsis NPR1 protein is a receptor for the plant defense hormone salicylic acid *Cell*

Reports **1**:639-647.

Ye Z-H, Zhong E (2022) **Outstanding questions on xylan biosynthesis** *Plant Science* 325:

111476.

Yokoyama R, Nishitani K (2004) **Genomic basis for cell-wall diversity in plants: A**

comparative approach to gene families in rice and Arabidopsis *Plant cell Physiology* **45**:

1111-1121.

Yuan Y, Teng Q, Lee C, Zhong R, Ye Z-H (2014) **Modification of the degree of 4-O-**

methylation of secondary cell wall glucuronoxylan *Plant Science* **219-220**: 42-50.

Zhang J, Xie M, Tuskan GA, Muchero W, Chen J-G (2018) **Recent advances in the**

transcriptional regulation of secondary cell wall biosynthesis in the woody plants

Frontiers in Plant Sciences **9**: 1535.

Zhao Y, Lin S, Qiu Z, Cao D, Wen J, Deng X, Wang X, Lin J, Li X (2015) **MicroRNA857 is**

involved in the regulation of secondary growth of vascular tissues in Arabidopsis *Plant*

physiology **169**: 2539-2552.

Zhong, R, Ye, Z-H (2014) **Complexity of the transcriptional network controlling**

secondary cell wall biosynthesis *Plant Science* **229**: 193-207.

Zhu S, Gu J, Yao J, Li Y, Zhang Z, Xia W, Wang Z, Gui X, Li L, Li D, Zhang H, Liu C (2022)

Liquid-liquid phase separation of RBGD2/4 is required for heat stress resistance in

Arabidopsis *Developmental Cell* **57**: 583-597.

1067

Acknowledgements

We thank Aurélie Le Ru for the histological staining and cell wall thickness measurements,

Hua Cassan-Wang for helping to the preparation of stem sections for staining, and Michele

Laudié for preparation of the mRNA libraries. This work was supported by the Centre

National de la Recherche Scientific (CNRS), and grants ANR-08-BLAN-0206 and 12-BSV6-

0010 from Agence National de la Recherche (ANR) to TL. This study was supported by the

"Ecole Universitaire de Recherche (EUR) TULIP-GS (ANR-18-EURE-0019), and the

EPIPLANT Groupement de Recherche (CNRS, France). The proteomic experiments

performed by LB and YC were partially supported by Agence Nationale de la Recherche

under projects ProFI (Proteomics French Infrastructure, ANR-10-INBS-08) and GRAL, a

program from the Chemistry Biology Health (CBH) Graduate School of University Grenoble

Alpes (ANR-17-EURE-0003). The funders had no role in study design, data collection and

analysis, decision to publish or preparation of the paper.

1081

Author contributions

1082

Study design: T.L., F.M., R.S., M.F., DP., and N.B.E.; mutant characterization and molecular

biology: A.K., D.P., C.P., R.M., and J.A.; RNA binding analysis: N.B.E.; MSIL co-

purification experiments, mass spectrometry and statistical analysis of co-IP data: D.G. and

P.H.; Total inflorescence stem proteomic and statistical analysis: L.B. and Y.C.;

Polysaccharide biochemical analysis: L.L-B. and R.S.; Lignin biochemical and FTIR analysis:

F.M.; immunohistochemical staining: Y.M.; MALDI-TOF analysis and statistical analysis:

R.S. and M.F.; writing: T.L., F.M., R.S., M.F., D.P., and N.B.E.; visualisation: T.L.;

supervision: T.L. and N.B.E.

1091

Competing interests

1092

1093 The authors declare no competing interests.

1094

1095

FIGURES

1096

Figure 1│ Musashi-like MSIL2/4 proteins redundantly control development in

Arabidopsis.

(**A**) Schematic representation of the Musashi-like (MSIL) protein family in *Arabidopsis*

thaliana. The invariant RNP1 and RNP2 motifs within the conserved RRM domains are

indicated. Numbers refer to amino acid identities between the Mouse Musashi/MSI RRM1/2

domains and the corresponding domains in Arabidopsis MSIL homologs. (**B**) Evolutionary

relationships between MSILs and related RNA-binding proteins. The scale bar indicates the

rate of evolutionary change expressed as number of amino acid substitutions per site. (**C**)

RNA-seq expression map of *MSIL* genes extracted from ARAPORT11. Shading is a $\log_2$

scale of transcripts per million (TPM). (**D**) General overview of the roots of 6-days-old wild

type Arabidopsis seedlings (Col-0) or transgenic seedlings expressing either a free GFP

protein or GFP-tagged versions of MSIL2/MSIL2G and MSIL4/MSIL4G. Scale bar, 10 μm.

(**E**) Photographs of representative rosettes of Col-0, *msil2/4*, *msil2/4-MSIL2*F1 and *msil2/4-*

*MSIL4*F1 plants. (**F**) Photographs of representative inflorescence stems of Col-0, *msil2/4*,

*msil2/4-MSIL2*F1 and *msil2/4-MSIL4*F1 plants. Abbreviations: Mus musculus Musashi2

(MmMsi2); human heterogeneous nuclear ribonucleoprotein A1 (hnRNPA1).

1113

Fig. 2│ RRM-dependent RNA-binding activity is essential for MSIL function *in planta*.

(**A**) Experimentally determined structures of the RRM1 domain of MmMSI2 (PDB ID: 6C8U)

and the RRM2 domain of MmMSI1 (PDB ID: 5x3y), and models of the RRM domains of

MSIL4 generated using the homology-modeling server SWISS-MODEL. The α-helices and

β-sheets of RRM domain as well as the phenylalanine residues that contact RNA bases are

highlighted. (**B**) Schematic representation of the mutations introduced in the RRM domains of

MSIL4. (**C**) Binding assays of MSIL4 RRM domains on ssRNA homopolymers *in* vitro.

Single-stranded polyA (pA), polyG (pG), polyC (pC) or polyU (pU) RNAs were subjected to

binding with the His-tagged recombinant RRM domains from MSIL4 (upper part) or its

mutated version (MSIL4^{RRM}, lower part). An anti-His antibody was used for detection. In.

represents the input fraction. (**D**) RRM-dependent RNA-binding activity is essential for

MSIL2/4 function in stem. Photographs of representative inflorescence stems of Col-0 and

msil2/4 mutants complemented or not with a WT version of MSIL4 (MSIL4G-8 and 3) or an

rrm mutant (MSIL4G^{RRM}-3 and 10).

1128

Fig. 3│ MSIL2/4 protein interactomes are enriched in proteins involved in 3'-UTR

binding and translation regulation. (**A**) The semi-volcano plots show the enrichment of

proteins co-purified with MSIL2F1 and MSIL4F1 as compared with control IPs. Y- and X-axis

display adjusted p-values and fold changes, respectively. The dashed lines indicate the

threshold above which proteins are significantly enriched/depleted (fold change > 2; adjP <

0.05). (**B**) Venn diagram showing the overlap between the MSIL2 and the MSIL4

interactomes. (**C**) PANTHER-based classification of the GO molecular function that are

overrepresented among the MSIL-interacting proteins. (**D**) Confocal monitoring of the

colocalization of MSIL2G, MSIL4G and PAB2R in root tips of 8-d-old seedlings were

monitored after 30 min of exposure to 38°C (heat stress) or 20°C for control treatment. (**E**)

Cycloheximide treatment inhibits the formation of MSILG and PAB2R cytosolic foci. The

transgenic plants expressing the stress granule marker PAB2R was used as a control. Root of

7-day-old seedlings expressing either MSIL4G, MSIL2G, or PAB2R were monitored after 1

hour of exposure to 38°C in the presence of DMSO control treatment (DMSO) or in the

presence of cycloheximide inhibitor (+CHX). Scale bar, 10 μm.

1144

Fig. 4│ MSIL proteins regulate the molecular architecture of SCW in Arabidopsis fibers.

(**A**) Expression profiles of the *MSIL2* and *MSIL4* genes in the inflorescence stem tissues as

retrieved from the Arabidopsis inflorescence stem tissue-specific transcriptome database[40]

(https://arabidopsis-stem.cos.uni-heidelberg.de/). The results of two replicates are shown. F,

fibers; X, xylem vessels; Cx Cambium (xylem side); Cp, Cambium (Phloem side); P, Phloem;

S, Starch sheath; E, Epidermis. (**B**) Cross-sections of Col-0 and *msil2/4* stems showing

vascular bundle (*xy*) and interfascicular fiber (*if*) cells stained with Toluidine blue.

Representative fiber and xylary areas analyzed in the panel 4C are outlined. (**C**)

Measurements of the SCW thickness of xylary fiber and interfascicular fiber cells. Values are

means (n>400)± SEM. Data were analyzed by unpaired Student's test. Asterisks indicate

significant differences relative to Col-0; ****p<0.0001. (**D**) Top: lignin autofluorescence

under ultraviolet (UV) using confocal microscopy in Col-0, *msil2/4* mutant, and

complemented *msil2/4-MSIL2*F1 and *msil2/4-MSIL4*F1 plant stem sections. xy: xylem fibers;

if: interfascicular fibers. Bottom: Phloroglucinol-HCl staining of Col-0, *msil2/4*, and

complemented *msil2/4-MSIL2*F1 and *msil2/4-MSIL4*F1 plant stem sections. Scale bar, 100 μm.

(**E**) Boxplots represent acetyl bromide lignin content, expressed as % of dry weight. Lignin

content was measured in the bottom section of mature stems (3 biological replicates per line).

Significant differences between Col-0, *msil2/4+MSIL2*F1, *msil2/4+MSIL4*F1, and *msil2/4*

plant lines are indicated by asterisks*, *P-value*< 0.01 according to Student's t test (n = 6 to 10).

(**F**) Confocal microscopy of Calcofluor-white staining of cross sections of Col-0 and *msil2/4*

mutant inflorescence stems. Scale bar, 100 μm. (**G**) Boxplots represent the glucose yield after

incubation of Col-0 and *msil2/4* lignocellulosic material with cellulases either without pre-

treatment (-), or after pre-treatment (+) with sodium hydroxide (NaOH).

1168

Fig. 5│The accumulation and activity of the glucuronoxylan decoration machinery are

altered in *msil2/4* mutant.

(**A**) MS-based quantitative comparison of Col-0 and *msil2/4* inflorescence stem proteomes.

Volcano plot displaying the differential abundance of proteins in Col-0 and *msil2/4* cells

analyzed by MS-based label-free quantitative proteomics. The volcano plot represents the -

log10 (limma p-value) on y axis plotted against the log2 (Fold Change msil2/4 vs Col-0) on x

axis for each quantified protein. Green and red dots represent proteins found significantly

enriched respectively in *msil2/4* and Col-0 Arabidopsis stems (log2(Fold Change) ≥ 1 and -

log10(p-value) ≥ 2.29, leading to a Benjamini-Hochberg FDR = 1.01 %). The up-regulated

proteins involved the glucuronoxylan biosynthetic pathways are indicated. (**B**) Gene Ontology

(GO) analysis of the proteins that are significantly up-regulated in the *msil2/4* mutant. The

gene ontology analysis of DEGs was performed using ShinyGO v0.76 software. Lollipop

diagrams provide information about GO fold enrichment, significance (FDR in log10), and

number of genes in each pathway. (**C**) Schematic model of glucuronoxylan substitution

patterns in Arabidopsis and the enzymes involved. IRX9/10/14, glycosyltransferases involved

in the synthesis of xylan (Xy) backbone. ESK1, eskimo1. GUX1, glucuronic acid substitution

of xylan1; GXM3, glucuronoxylan methyltransferase3 are involved in the glucuronoxylan

decoration. (**D**) Neutral monosaccharide composition of alcohol insoluble residue (AIR)

extracted from inflorescence stems of wild-type and *msil2/4* mutant plants that have been pre-

hydrolyzed or not with acid. Rha, rhamnose; Fuc, fucose; Ara, arabinose; Xyl, xylose; Man,

mannose; Gal, galactose; Glc, glucose. (**E**) Left: Immunofluorescence labeling of transverse

sections of Col-0 and *msil2/4* stems with the LM10 and LM11 antixylan antibodies.

Representative fiber and xylary regions analyzed by immunofluorescence are outlined. Xy,

xylem fibers; if, interfascicular fibers. Right: Quantification of the fluorescence was done

using ImageJ software and processed according to Supplementary file 5. Significant

differences are indicated by asterisks*, *P-value*< 0.01 according to Student's t test (n = 5 to 12)

1195

Fig. 6│MSILs restrain the degree of 4-O-methylation of glucuroxylan in Arabidopsis

inflorescence stem. (**A**) MALDI-TOF mass spectra of xylooligosaccharides generated by

xylanase digestion of xylan from Col-0, *msil2/4*, and complemented *msil2/4-MSIL2*F1 and

*msil2/4-MSIL4*F1 inflorescence stem materials. The ions at m/z 1705/1747 and 1719/1761

correspond to acetylated xylo-decapolysaccharides bearing a GlcA residu (GlcA-Xyl10-

Ac4//GlcA-Xyl10-Ac5) or a methylated GlcA residue (MeGlcA-Xyl10-Ac4//MeGlca-Xyl10-

Ac5). (**B**) Acidic monosaccharide composition of alcohol insoluble residue (AIR) extracted

from inflorescence stems of wild-type and *msil2/4* mutant plants that have been pre-

hydrolyzed or not with acid. GalA, galacturonic acid; GlcA, glucuronic acid. (**C**) Relative

changes in unmethylated/methylated GlcA decapolysaccharide ratio in Col-0 (black) and

msil2/4 mutant (red) as controlled by the addition of external spike-in control corresponding

to a pentaacetyl-chitopentaose. (**D**) GFP-based RNA-IP assays. Top panel (WB): western

blots performed using antiGFP or antiUGPase antibodies on protein fractions from inputs or

antiGFP immunoprecipitates from WT and the *msil2/4* mutant plants expressing the WT

(MSIL4G-3) or RRM mutant (MSIL4G^{RRM}-3) MSIL4-GFP fusions. Bottom panel (RT) :

Corresponding RT-PCR using *GXM1* or *GXM3* specific primers on RNA fractions.

1212

Fig. 7│ Model of MSIL-dependent control of glucuronoxylan methylation in

Arabidopsis and its consequence for SCW architecture. In the interfascicular fiber cells of

Col-0, MSIL2/4 restrain the translation of the glucuronoxylan biosynthesis enzymes,

including the rate-limiting GXM3 enzyme. This activity would keep the level of

glucuronoxylan methylation at an intermediate level, therefore providing a biochemical

environment that favors the interactions between the glucuronoxylan and lignin polymers. In

the interfascicular fiber cells of *msil2/4* mutant, the translation of the glucuronoxylan

biosynthesis machinery, including GXM3, is increased, leading to the deposition of an over-

methylated form of glucuronoxylan that would be less prone to interact with lignin and

establish SCW formation. In a non exclusive manner, MSIL2/4 could also have a positive role

in lignin synthesis, whose defect in *msil2/4* would lead to a decrease in lignin content. The

model was inspired from Grantham et al., 2017.

A

*MmMSI2*RRM1 *AtMSIL4*RRM1

*MmMSI1*RRM2 *AtMSIL4*RRM2

B

RRM1   RRM2
F F   F F   *MSIL4*
↓ ↓   ↓ ↓
D D   D D   *MSIL4^{RRM}*

C

MSIL4
kDa In. pA pG pC pU
55—
37—

55—
37—

In. pA pG pC pU
MSIL4^{RRM}

D

Col-0 / *MSIL4G-8*   Col-0 / *MSIL4G-3*   Col-0 / *MSIL4G^{RRM}_4*   Col-0 / *MSIL4G^{RRM}_10*
msil2/4   *msil2/4*   *msil2/4*   *msil2/4*

A

Volcano plot (Bait MSIL2F1,4IPs)

MSIL2F1

Volcano plot (Bait MSIL4F1,4IPs)

MSIL4F1

B

MSIL2F1
associated proteins

MSIL4F1
associated proteins

C

GO molecular function	Fold Enrichment	P value
Poly (A) binding	>100	$2.15^{E}-14$
mRNA 3'-UTR binding	>100	$5.81^{E}-11$
Single-stranded RNA binding	>100	$1.80^{E}-11$
Translation initiation factor activity	74.04	$2.77^{E}-07$
mRNA binding	24.56	$2.49^{E}-21$
RNA binding	16.04	$2.47^{E}-18$

D

MSIL4G   PAB2R   Merged

Control

Heat stress

Seedling roots

E

MSIL4G   PAB2R   MSIL2G   PAB2R

-CHX

+CHX

Seedling roots (38°C)

Interfascicular fiber cell (Col-0)

Interfascicular fiber cell (*msil2/4*)

AAA

MSIL

?

mRNA
(*lignin-related enzymes*)

mRNA
(*glucuronoxylan-related enzymes*)

Golgi apparatus

GXM3

GUX1

ESK1

IRX9
IRX10
IRX14

mexylan

Secretion

Lignin

meXylan

Cellulose microfibrils

AAA

MSIL

mRNA
(*lignin-related enzymes*)

mRNA
(*glucuronoxylan-related enzymes*)

Golgi apparatus

GXM3

GUX1

ESK1

IRX9
IRX10
IRX14

Mexylan

Secretion

Lignin

?

MeXylan

Cellulose microfibrils

Xylosyl   Glucuronosyl   MeGlucuronosyl   Acetyl   Electrostatic contact/linkages