# Mammals achieve common neural coverage of visual scenes using distinct sampling behaviors

Jason M. Samonds[1,*], Martin Szinte[2], Carrie Barr[1], Anna Montagnini[2], Guillaume S Masson [2] and Nicholas J. Priebe[1]

1. Center for Learning and Memory and The Institute for Neuroscience, The University of Texas at Austin, Austin, Texas, 78712

2. Institut de Neurosciences de la Timone (UMR 7289), Centre National de la Recherche Scientifique and Aix-Marseille Université, Marseille, France

*Corresponding author.
**Email:** samondjm@gmail.com

**Author Contributions:** J.M.S., M.S., A.M., G.S.M., and N.J.P. designed research; J.M.S., M.S., C.B. and A.M. performed research; J.M.S., M.S., C.B. and A.M. analyzed data; and J.M.S. wrote the original draft; J.M.S., M.S., A.M., G.S.M., and N.J.P. edited the paper.
**Competing Interest Statement:** No competing interests.
**Classification:** Biological Sciences: Neuroscience
**Keywords:** saccades, fixation, natural scenes, visual cortex

**This PDF file includes:**
>      Main Text
>      Figures 1 to 2

**Abstract**

Most vertebrates use saccadic eye movements to quickly change gaze orientation and sample different portions of the environment. Visual information is integrated across several fixations to construct a more complete perspective. In concert with this sampling strategy, neurons adapt to unchanging input to conserve energy and ensure that only information for novel fixations is processed. We demonstrate how adaptation recovery times and saccade properties interact, and thus shape spatiotemporal tradeoffs observed in the motor and visual systems of different species. These tradeoffs predict that in order to achieve similar visual coverage over time, animals with smaller receptive field sizes require faster saccade rates. We find comparable sampling of the visual environment by neuronal populations across mammals when integrating measurements of saccadic behavior with receptive field sizes and V1 neuronal density. We propose that these mammals share a common statistically driven strategy of maintaining coverage of their visual environment over time calibrated to their respective visual system characteristics.

**Significance Statement**

Mammals rapidly move their eyes to sample their visual environment over successive fixations, but they use different spatial and temporal strategies for this sampling. We demonstrate that these different strategies achieve similar neuronal receptive field coverage over time. Because mammals have distinct sensory receptive field sizes and neuronal densities for sampling and processing information, they require different eye movement strategies to encode natural scenes.

## Introduction

Saccadic eye movements have been demonstrated across vertebrates (1), but the size and frequency of these eye movements vary substantially (2). Saccadic eye movements are thought to overcome inhomogeneity in our sensory representation by providing receptive fields a unique and updated perspective for each fixation (Fig. 1a, b). The retina contains a differential density of photoreceptors (3-6) and occlusion by retinal vasculature (7). This inhomogeneity persists along the visual pathway and is related to the cortical magnification factor (8, 9), acuity (10-15), color sensitivity (16), and irregular ocular dominance domains (17). Additional inhomogeneity emerges within cortex, such as the irregular distribution of orientation (15) and disparity selectivity (18-20). By making eye movements to cover different regions of the scene, an internal representation may be constructed by integrating novel receptive field information over successive fixations (21).

Adaptation and saccades work together to provide novel receptive field inputs and conserve energy. All cells could respond equally over time, but that would be costly in terms of neurophysiological metabolic energy so neurons quickly adapt to unchanging visual input to conserve that energy (22). Eye movements, which change where a receptive field samples, could be continuous and uniform across the visual scene, but that would be costly. To conserve energy, the eyes move discretely and quickly at the minimum distance necessary to provide novel inputs to the visual system given natural scene statistics (2). This argument successfully accounts for the differences in distribution shapes of saccade sizes across mammals (2), but it is not clear if it also explains differences in saccade rates. Since sampling the visual environment depends on both space and time, we examined the differences in adaptation dynamics and saccade statistics across fixations across different mammals.

## Results

The adaptation properties of cortical neurons are remarkably distinct across mammals. For both mice and macaques, the responses to brief presentations (<0.5 s) within visual receptive fields decay within a couple hundred milliseconds, but macaque neurons recover their responsiveness very quickly (within a few hundred milliseconds, Fig. 1a, c, blue) (22-24) while mouse neurons

recover much slower (over several seconds, Fig. 1b, c, red) (25, 26). Cat neurons recover with a time constant somewhere in between these two values (27). Recovery time constants for these animals in these studies were measured for repeated presentations of the same stimuli. Saccadic eye movements that decorrelate input across fixations may decrease the effects of neural adaptation (2), but the adaptation that persists nonetheless requires time for recovery (24). Macaque neurons will still recover much faster than mouse neurons even with large enough saccades.

To quantify how different saccade rates influence adaptation, we constructed a model in which the dynamics of adaptation could be incorporated into the theoretical framework we previously used to explain the relationship between neural adaptation and saccade size (2). Responses ($r_m$) of a population of $M = 500$ neurons to $N = 500$ stimuli ($s_n$) following sequential saccades were multiplied by a divisive change in gain dependent on the response to the previous stimulus ($s_{n-1}$):

$$r_{m,adapted}(s_n) = \frac{k}{k+r_m(s_{n-1})}r_m(s_n) \tag{1}$$

Parameters of this function ($k$ and average responses) were chosen to approximately match observed adaptive changes in gain in macaques, mice, and cats, where the average gain never reaches zero for even the fastest repeat presentations (see Supplemental Fig. 1 and Methods for details) (23-27). We varied the time between stimuli to represent different saccade rates and model responses decayed over time ($t$) as an exponential decay function with a fixed time constant $\tau_{decay}$. The divisive gain based on the initial response after each saccade recovered over time based on a second exponential decay function:

$$r_{m,adapted}(t, s_n) = \frac{k}{k+e^{\frac{-t}{\tau_{recovery}}} r_m(1,s_{n-1})}r_m(1,s_n)\, e^{\frac{-t}{\tau_{decay}}} \tag{2}$$

We varied the time constant ($\tau_{recovery}$) of this function to cover a range that includes experimentally observed adaptation recovery time constants to brief presentations of stimuli in macaques and mice(23-26). To mimic the effects of saccades, we repeated this analysis but changed the stimulus. For some conditions, the stimulus changed randomly, thus providing a decorrelated input mimicking large saccades. For other conditions, the stimulus did not change,

mimicking small saccades. As saccade rate increases, neurons have less time to recover from adaptation and the overall responses decline (Fig. 1d). For both animals, large saccades reduced adaptation compared to small saccades yielding higher responses for faster saccade rates, reflecting a linkage between saccade size and rate (Fig. 1d, solid vs. dashed lines). To illustrate how the recovery time would relate to saccade rate, we plotted the saccade rate that would still maintain 75% of responsiveness against the recovery time constant (Fig. 1e). This simple model predicts a dramatic difference in saccade rate that depends on the adaptive dynamics in visual neurons, and captures the large differences in saccade rates observed in macaques (Fig. 1e, blue point; 3.8 saccades/s) (28) and mice (Fig. 1e, red point; 0.6 saccades/s) (2).

Our model suggests that there should be functional links between the spatiotemporal dynamics of the visual and motor systems. To characterize the spatiotemporal tradeoffs made by these systems, we first compared the saccade rates and sizes when presenting natural scenes to three different species known to have different visual system properties: humans, marmosets, and mice (Fig. 1f and Methods). As demonstrated previously (2), there are clear differences in saccade sizes between these species, with all animals exhibiting a skewed distribution in saccade sizes (Supplemental Fig. 2b). Marmosets have the smallest saccades and mice have the largest saccades. For all animals, saccade sizes were slightly larger for larger images (see also ref. (29)) (Supplemental Fig. 2b; bootstrapped, p < 0.001 for all comparisons). There are also clear differences in saccade rates between species (Fig. 1f). Marmosets have the highest saccade rates and mice have the lowest saccade rates. For all animals, saccade rates were slightly higher for larger images (see also ref. (29)) (Supplemental Fig. 2c; bootstrapped, p < 0.001 for all comparisons). Saccade behavior is similar between animals with only a shift in rates and sizes. Including previously reported data from other animals (macaques and cats) reveals an inverse relationship between saccade rate and saccade size across distinct mammalian species (Fig. 1f).

The decorrelation hypothesis predicts that saccade size differences between species are attributed to the properties of their respective visual systems (2). Human and macaque receptive fields are small relative to mouse receptive fields, whereas marmoset and cat receptive fields lie

between these sizes. Consequently, to achieve decorrelation and sample new information, mice need much larger saccades in order to move their larger receptive fields as compared to humans. Our model also suggests that saccade rates will be matched to differences in adaptation recovery time constants (Fig 1e). Since adaptation time constants in individual animals can vary depending on experimental conditions and data based on similar conditions is fairly limited across species, we rather considered preferred temporal frequency distributions across several species to provide a more comprehensive view of the relationship between spatial and temporal properties of their visual system neurons. Indeed, temporal frequency tuning has been measured in several animals under very similar experimental conditions (e.g., spike rate responses to sinusoidal luminance gratings; see Methods) in different laboratories (30-40). Similar to the saccade statistics, there is an inverse relationship between preferred temporal frequency and receptive field size across these animals (Fig. 1g). Both the motor and visual systems share spatiotemporal tradeoffs across species where those with smaller receptive fields can make smaller saccades, but require more frequent sampling and faster visual processing. Indeed, we find that low luminance contrast of natural images, which it is known to shift temporal frequency tuning lower (35, 41, 42), is associated with a corresponding decline in saccade rate (Supplemental Fig. 3).

The relationship we uncover between saccade size and rate may be from differences between species other than receptive field sizes. To test whether changes in receptive field size are linked to saccade rate, we reduced the spatial acuity of mice using monocular deprivation during the critical period. This manipulation shifted their saccades to larger sizes (Fig 1h, top; see ref. (2)) and significantly reduced their saccade rates compared to control mice (Fig 1h, bottom; bootstrapped, P < 0.001). This causal manipulation demonstrates that there is an inverse relationship between saccade size and rate. A similar change in saccade size and rate is observed in humans on a coarse scale with substantial reduced acuity due to macular degeneration or foveal occlusion (43, 44) and on a finer scale in subjects with myopia and amblyopia (45, 46).

Our modeling and empirical data in different species show that visual and motor systems share similar spatiotemporal tradeoffs to sample visual information. We reasoned that such a

tradeoff is constrained by a common computational goal: achieving equivalent decorrelated spatiotemporal coverage of their visual environment. The mouse with both large V1 receptive fields and eye movements would need to make eye movements less frequently compared to the human with both small receptive fields and eye movements in order to achieve similar updated coverage over time.

To examine coverage over time quantitatively, we applied a circular mask equal to the median V1 receptive field size for each animal to every fixation point (Fig. 2a, green circles). We then added up all of those masks (each equal to one for that area) across space and time for all images to generate cumulative spatial maps. Lastly, we divided those maps by the total of the intersaccadic interval times to get a percent coverage per second (Fig. 2a, right). Because receptive fields are based on distance squared, the mouse achieves much more coverage over time compared to marmosets and humans even with their slower saccade rates (Fig. 2b).

One notable exception to the general inverse relationship between saccade rate and size across species is observed for humans, who appear to have lower saccade rates than macaques although they have slightly smaller saccade sizes (Fig. 1f). In addition, if we assume that marmosets make larger saccades with their heads (28) with no decrease in saccade rate (see Supplemental videos 1-2 and Supplemental Figs 4-5), they would have higher saccade rates than macaques with larger saccade sizes. This suggests that saccade rates might also depend on species-specific factors other than saccade sizes and receptive field sizes.

We postulated that the main bottleneck constraining coverage over time is the receptive field size and density of the first cortical stage, the primary visual cortex. The animals we are examining have very different densities of V1 cortical neurons per degree of visual angle (dva), and therefore differences in densities of feature preferences (47) (Fig. 2c). The human has many more neurons for each receptive field location representing many more different visual features than the mouse and mathematically, larger networks can provide greater decorrelation (48). We can take our original coverage maps over time (Fig. 2b) and multiply by the two-dimensional (2D) V1 neuronal density for each animal (see Methods for details) to compare receptive field cortical coverage over time between species (Fig. 2d). Because marmosets often use head movements

for gaze changes and their eye movements, unlike humans, appear to be restricted to mostly ±10 dva (Fig. 2b, center; see also Supplemental videos 1-2 and Supplemental Figs 4-5 and ref (28)), we observe comparable coverage for smaller 20 dva images between humans and marmosets (Fig. 2d, center, light gray versus light cyan). Mice also use head movements for changes in gaze (49-51), but their eye movements at least cover ±20 dva (Fig. 2b, right) so we observe comparable coverage for 40-dva images between humans and mice (Fig. 2d, right, dark gray versus red). These comparisons suggest that saccades for these mammals are decorrelating similar numbers of neurons over time despite having very different saccade rates, saccade sizes, and receptive field sizes. With this sampling strategy, differences in saccade rates and sizes between animals are explained by corresponding differences in receptive field sizes and cortical densities with respect to visual space. If saccade rates and sizes are calibrated to achieve a certain level of visual coverage, this would also explain why saccade rates and sizes change with image size and decrease over viewing time (Supplemental Figs. 2 and 6; see also refs (2, 29)). As our model predicts, saccade rates are always higher in the condition where it is easier to achieve decorrelation. The oculomotor system indeed seems to favor larger versus smaller saccades, since it is easier and faster to make larger saccades (52-54).

**Discussion**

Our model makes some important assumptions. First, for the purposes of predicting saccade sizes, we assumed that the receptive field is static and only examined image differences across sequential fixations (2). In fact, saccades have complex dynamics over 10's of milliseconds and the eyes drift and oscillate continuously over 100's of milliseconds or seconds of fixation. These fast and slow oculomotor dynamics effectively whiten the visual inputs (55-57). This whitening is typically outside of the frequency spectrum of the V1 response preferences that we used to predict saccade sizes (Supplemental Fig. 7). Indeed, V1 responses following a saccade are very similar to those evoked by sequentially flashed stimuli (58). Nonetheless, saccades can produce an intra-saccadic flow of information that is sensed (59, 60) and can boost post-saccadic visual motion processing (61). Second, we assumed that the primary visual cortex was the main

processing bottleneck that constrains saccade sizes and rates. This choice was partly made because of the large number of comparable data available across multiple species. This assumption does not rule out that differences in receptive fields found earlier, later, or subcortically in the visual system between species also contribute to their oculomotor differences. Future work shall include additional processing that occurs during both saccades and fixation and how they influence both cortical and subcortical responses.

Yarbus illustrated that humans move their eyes deliberately towards objects and features of interest and the overall pattern of those movements can heavily depend on the task (62). Decades of research support that the primary purpose of saccadic eye movements is to move an area of interest into the high-spatial acuity portion of the visual field (62-66). Even afoveate animals, such as the mouse, use gaze changes to move objects of interest into the central portion of their upper visual field (50, 67) where binocularity and spatial features are represented better (14, 15, 19). Our hypothesis is complementary to foveation and does not attempt to predict saccade targets instantaneously. We use decorrelation as a measurement to quantify a statistical sampling strategy for natural scenes over a long time scale of several fixations to update an internal visual representation. It is important to point out that this sampling strategy can depend on and adapt to environmental and task demands as well. We see this when varying image contrast and size (Supplemental Figs. 3 and 6; see also ref. (29)), but previous studies have also shown predictable changes in coverage (size/rate) statistics using images with biased spatial frequency content (68), tasks with high acuity demands (69), or tasks with high cognitive demands (70, 71). Previous work examining transsaccadic integration also shows that information from past fixations influence perception during the current fixation. If that past information about a particular visual feature was more reliable than the current information, it has a stronger influence on perception (72-74). Additionally, a higher rate of saccades improves performance in detecting changes within a scene (75) highlighting the importance of maintaining sufficient coverage. Overall, our analyses illustrate that the oculomotor and visual systems of multiple mammals coordinate to sample the environment efficiently under a diverse range of processing constraints.

**Materials and Methods**

Adaptation model

For the model, we chose parameters for equation 2 that best matched the average response dynamics (decaying to 30% of the initial response) to brief sequential presentations reported for adaptation experiments in macaques and mice (23-27). The decay time constant $\tau_{decay}$ for all simulations was 60 ms and the recovery time constant $\tau_{recovery}$ varied between 100 and 900 ms. The initial unadapted firing rate was 50 spikes/s and the constant *k* for the divisive gain function was 30.

Human subjects

Twenty-four participants and staff members of the Institute des Neurosciences de la Timone took part in the different experiments. Five participants (22-29 years old, 2 females) took part in the free viewing tasks and 20 participants (19-54 years old, 10 females, 3 authors) took part in the fMRI experiments where we looked at their fixation data. Experiments were approved by the Ethics Committee of Aix-Marseille University and conducted in accordance with the Declaration of Helsinki. All participants gave written informed consent.

Mice

Experiments and procedures were performed on eight C57bl6 adult female and male mice (RRID: JAX: 00664), one female and two male PV-Cre mice (76), two female PV-Cre;Ai14 mice (76), and two female and four male PV-Cre;ChR2 mice (77). Mice were group housed and maintained on a 12-hour light/dark cycle under standard housing conditions. All procedures and care were performed in accordance with the guidelines of the Institutional Animal Care and Use Committees at the University of Texas at Austin.

Marmosets

Seven marmosets (1.5-4 years, 2 females) were obtained from TxBiomed and bred in-house and were group-housed. Food and water were provided ad libitum. All procedures and care were

performed in accordance with the guidelines of the Institutional Animal Care and Use Committees at the University of Texas at Austin.

Human saccades

For free viewing experiments, participants sat in a quiet and dimly illuminated room, with their head positioned on a chin and forehead rest. The experiments were controlled by a HP Z420 (Hewlett-Packard, Palo Alto, CA, USA) computer equipped with an Nvidia Quadro 600 video card (Nvidia, Santa Clara, CA, USA). Binocular eye position was recorded using an Eyelink 1000 tower mount (SR Research, Osgoode, ON, Canada) at a sampling rate of 2 kHz (1 kHz per eye). The experimental software controlling the display as well as the eye tracking was implemented in Matlab (The MathWorks, Natick, MA, USA), using the Psychophysics (78, 79) and Eyelink toolboxes (80). Stimuli were presented at a viewing distance of 60 cm on a 32-inch Display++ LCD monitor (Cambridge Research System, Rochester, UK) with a spatial resolution of 1920 by 1080 pixels and a vertical refresh rate of 120 Hz. Participants' gaze position was calibrated with a 13-point custom calibration sequence with sequences validated at the beginning of an experimental session as well as when necessary.

Participants took part in free viewing tasks composed each of 40 runs of eight trials each. These runs were completed in four experimental sessions (on different days) of about 60 min each (including breaks). Participants were instructed to freely explore a set of 80 images displayed at the center of a gray background screen. These pictures were presented on each trial for a duration of 30 seconds and separated each by two seconds of blank gray screen. Pictures were selected from the natural images from the McGill Calibrated Color Image Database (81). For the first 20 runs, participants randomly explored the screen with either the 80 full-sized selected images covering 40 by 32 dva of the visual field or the same 80 images center-cropped covering 20 by 16 dva of the visual field. For the next 20 runs, each participant explored 40 images selected randomly from the same set of 80 images, but with their luminance contrast scaled down uniformly at either 5 or 10% of the original contrast. These images covered 40 by 32 dva of the

visual field during the first 10 runs and 20 by 16 dva of the visual field in the last 10 runs. Data from images of both sizes were combined for contrast-related results.

For fixation data, participants laid down in a dimly illuminated Prisma 3T MRI scanner (Siemens Healthcare, Erlangen, Germany), with their head restrained within a 64-channel receiver coil array for an experiment unrelated to this study. The experiment was controlled by a Dell Precision 7820 computer (Dell, Round Rock, TX, USA) equipped with a Nvidia Quadro RTX 5000 video card (Nvidia, Santa Clara, CA, USA). The left eye gaze position for each subject was recorded using an Eyelink 1000 long-range mount (SR Research, Osgoode, ON, Canada) situated at the end of the bore, at a sampling rate of 1 kHz. The experimental software controlling the display as well as the eye tracking were identical to that used in the free viewing tasks. Stimuli were back projected on a projection screen situated at the end of the bore and viewed through a mirror at a viewing distance of 120 cm. Stimuli were presented with a PROPixx DLP LED projector (VPixx Technologies, Saint-Bruno, QC, Canada) with a spatial resolution of 1920 by 1080 pixels and vertical refresh rate of 120 Hz. The eye tracking calibration and validation was the same as the ones used in the free viewing tasks.

Participants who took part in this fixation task also participated in a longer session combining eye tracking and multi-modal MRI tasks (lasting about an hour). Here, we only report eye tracking results from the first 30 s of two runs of a "resting state" task lasting about 7 min each. Within each run, participants were instructed, while "letting their minds wonder", to keep their eyes on a white fixation mark (0.3-degree radius) presented at the screen center on a black background during 400 seconds (500 whole brain functional scans with a TR of 0.8 seconds).

For all experiments, saccades were detected as previously described for marmosets with a velocity threshold (2).

Marmoset saccades

Marmoset eye position was tracked as previously described using an EyeLink 1000 (SR Research, Osgoode, ON, Canada) camera to capture binocular eye position at sampling rate of 1 kHz (500 Hz for each eye) (2, 19) and saccades were detected with a velocity threshold (2).

Twenty-four small (20 by 16 dva) and 24 large (40 by 32 dva) images from the McGill Calibrated Color Image Database(81) were presented per session on a CRT monitor (85 Hz refresh rate) 50 cm away randomly interleaved them with unrelated 1-s fixation trials using Maestro software suite (https://sites.google.com/a/srscicomp.com/maestro/). In separate sessions, the images were presented with their luminance contrast scaled down uniformly at either 5 or 10% of the original contrast. Data from images of both sizes were combined for contrast-related results.

Marmosets had higher saccade rates than expected with respect to their temporal frequency relationship to the other animals (Fig. 3A-B). Previously, we found that marmosets have smaller saccade sizes than we predicted based on their receptive field sizes (2). We attributed this mismatch to marmosets normally using saccadic head movements to make rapid changes in gaze (28). To probe this idea further, head and eye position were measured in marmosets that were not head restrained using DeepLabCut (82) to track pupils and the center of the upper forehead in 400 by 400 pixel images collected at 60 frames per second (see Supplemental video 1). Calibration for these results was done on head restrained marmosets using the fixation methods described previously (28, 83) (see Supplemental video 2). Saccade statistics based on DeepLabCut of posted marmosets were similar to those measured with the EyeLink system (Supplemental Fig 2). For analysis of unposted marmosets, we only looked at continuous segments when the gaze was towards the monitor (within ±50 dva from the monitor center).

Marmosets clearly used combined head and eye movements for nearly all changes in gaze, which always produced larger saccades than with the eyes alone (Supplemental Fig 3A-C). We did not observe obvious differences in the saccade rates (Supplemental Fig 3D). Interestingly, marmosets also cock their heads frequently and rapidly, especially when presented with novel objects (84). This might increase their ability to decorrelate orientation tuned receptive field responses and allow them to sample visual information at a faster rate than with saccades alone (shift the curve in Fig. 1D to the right).

For comparison to our data, the two open circles (median size and rate for each subject) in Fig. 3A were reformatted from figures from a previous publication and the two marmosets were

viewing 44 by 34 dva images displayed for 20 s of natural images including humans, macaques, and marmosets (28).

Mouse saccades

For image size experiments, saccade statistics are from four C57bl6 and two PV-Cre;Ai14 female mice from a previously reported study(2). These mice were trained for a binocular discrimination task (19) and 8-12 small (40 by 32 dva) and 8-12 large (80 by 64 dva) images/per session were presented before or after 2-3 training sessions. Images from the McGill Calibrated Color Image Database (81) were presented onto a screen 22 cm away with a DLP LED projector. Binocular eye movements were tracked using custom software to analyze 250 by 250-pixel images collected at 20-30 frames per second with cameras aligned with the orbital axis of each eye. Since mice have a much higher contrast threshold and saturation point than primates, for image contrast experiments, we had them view natural images scaled down to 25 and 50% of the original contrast. This increase for mice compared to primates roughly corresponds to the ratio of increase in contrast thresholds for mice compared to humans (85). For three male and one female C57bl6 mice, images were scaled to modulate at a ratio of 0.25 or 0.5 of their original contrast modulation. Experiments were run in the same manner as the previous study except four images each of 25, 50, and 100% contrast were randomly interleaved for two sessions. Mice were first trained to discriminate black from white or crossed versus uncrossed disparities from dynamic random dot stimuli by licking left or right. Natural images were shown before or after a training session when mice reached consistent performance above 75% correct for any task. Eyes were tracked in the same manner as previously described except that DeepLabCut(82) was used to track pupil position. Saccades were detected as previously described with a velocity threshold and comparing detection for both eyes(2).

For monocular deprivation (MD) experiments(86), one eye in each mouse (two female and two male PV-Cre;ChR2) was sutured closed at P24 under anesthesia and sutures were removed at P33. Three age-matched PV-Cre mice (one female and two males) were used as

controls. Acuity was assessed and saccades for all mice viewing natural images were measured at P40 (2).

Macaque saccades

Macaque data in Fig. 3A were reformatted from figures from a previous publication (median size and rate for each subject) and the two macaques were viewing 44 by 34 dva images displayed for 20 s of natural images including humans, macaques, and marmosets(28). The median and 25th and 75th  percentile was also extracted from the aggregate distribution of both subjects.

Cat saccades

Cat saccade size statistics (median, 25th percentile quartile, and 75th percentile quartile of reported sizes) were reformatted from figures from previous publications where the cats were either free viewing in the dark (87) or viewing a 21-min natural video of unknown size (88). Cat saccade rate statistics (median, 25th percentile quartile, and 75th percentile quartile of intersaccadic intervals) were extracted from horizontal and vertical traces of unpublished data of a head-fixed cat watching a 30-s video of animals at the zoo (unknown size, but smaller than the 60 by 40 dva screen) provided by Theodore Weyand at Louisana State University School of Medicine. Saccades were detected as previously described for mice and marmosets with a velocity threshold. Dr. Weyand observed that cats made saccades more often for the movie compared to dots or a blank screen (unpublished observations).

Receptive field size and temporal frequency data

All receptive field size distributions were extrapolated from data from previously published studies that measured the areas or widths of the minimum response field from electrophysiological recordings for cats, marmosets, and macaques, and GCaMP fluorescence thresholded at half the maximum response for mice. Receptive field (RF) size distributions for cats(10-12), marmosets(9), and macaques(9, 13) were generated by combining receptive field size versus

eccentricity data fits with cortical magnification data fits, and for mice, we used the reported

distribution (89) (see reference (2) for more details).

All preferred temporal frequency distributions were aggregated from previously published

electrophysiology studies that were based on peak responses in tuning curves generated from

average spike rates measured from single neurons in anesthetized animals. For all studies,

responses for each temporal frequency were measured using drifting sinusoid gratings centered

on the receptive field of single neurons using gratings with the preferred orientation and spatial

frequency of the neuron. All spikes were measured extracellularly for single units. All studies used

single insulated tungsten electrodes, except a silicon microprobe with 16 recording sites was

used to record spikes in mice for one study (36). Only summary statistics from the microprobe

study were used to validate our aggregate statistics from other studies. For macaques, peak

temporal frequencies for 130 single neurons from ref. (31) were combined with fitted peak

temporal frequencies for 75 single neurons from ref. (32) to generate an aggregate distribution of

peak temporal frequencies. This distribution was consistent with summary statistics reported in

additional macaque studies using similar methods (34, 35, 38). For cats, peak temporal

frequencies for 36 single neurons from ref. (30) were combined with fitted peak temporal

frequencies for 72 single neurons from ref. (33) to generate an aggregate distribution of peak

temporal frequencies. For mice, fitted peak temporal frequencies for 192 single neurons from ref.

(37) were combined with fitted peak temporal frequencies for 69 single neurons from ref. (38) to

generate an aggregate distribution of peak temporal frequencies. This distribution was consistent

with summary statistics reported in an additional mouse study using similar methods (36). From

the complete aggregate distributions for each animal, we measured the median and 25th and

75th quartiles.

Cortical density

Cortical density per dva was estimated by taking the total number of V1 neurons and computing what the total 2D representation of neurons would be based on the surface area and depth of V1, and then dividing that by the total visual area in degrees. We estimated total numbers of neurons using the volume of V1 in $mm^3$ times a density of 150,000 neurons/$mm^3$ because this density has been a relatively consistent value reported in V1 for chimpanzees (90), macaques (91), marmosets (92), and mice (93). For humans, we used 1470 $mm^2$ for surface area (94) and 2.5 mm for depth (95) to get 211 neurons/$dva^2$. For marmosets, we used 200 $mm^2$ for surface area (96) and 1.1 mm for depth (97) to get 28.7 neurons/$dva^2$. For mice, we used 3 $mm^2$ for surface area (98) and 0.8 mm for depth (95) to get 0.43 neurons/$dva^2$.

Statistics and reproducibility

All statistical tests were nonparametric based on the median and error bars were based on bootstrap analysis of the median by resampled data 1000 times, allowing repeats, to produce surrogate datasets of the same size. The 160th and 840th samples were used for the standard error of the median for all results. For data we collected on humans, marmosets, and mice, we present the entire distribution of all subjects grouped or single subject medians as single solid data points, an outline connecting the standard error of the median of grouped data across rate and size distributions, and an outline connecting the 25th and 75th quartiles of grouped data across rate and size distributions. For receptive field, temporal frequency and saccade data published previously by other authors, we present the median of grouped data or medians of single subjects as open circles and an outline connecting the 25th and 75th quartiles of grouped data across y- and x-axis distributions. For statistical tests, bootstrapped data for one set of observations were used to find the percentile equal to the median of the compared set of observations. If the compared median was completely outside of the bootstrapped dataset, we reported that as $P < 0.001$.

**Data and code availability**

The data presented in the figures and the code used to generate the figures are available at

Figshare: https://figshare.com/projects/

Mammals_achieve_common_neural_coverage_of_visual_scenes_using_distinct_sampling_beha

viors/136912

**Acknowledgments**

We thank Allison Laudano for animal care and Jagruti Pattadkal for helping set up all marmoset

experiments. We thank Theodore Weyand for sharing cat data and helpful discussions. This

research was funded by ANR-20-NEUC-0002 (G.M.) and NIH-R01-NS120562 (N.P.).

**References**

1.  Land MF (1999) Motion and vision: why animals move their eyes. *J Comp Physiol A* 185(4):341-352.
2.  Samonds JM, Geisler WS, & Priebe NJ (2018) Natural image and receptive field statistics predict saccade sizes. *Nat Neurosci* 21(11):1591-1599.
3.  Goodchild AK, Ghosh KK, & Martin PR (1996) Comparison of photoreceptor spatial density and ganglion cell morphology in the retina of human, macaque monkey, cat, and the marmoset Callithrix jacchus. *J Comp Neurol* 366(1):55-75.
4.  Szel A*, et al.* (1992) Unique topographic separation of two spectral classes of cones in the mouse retina. *J Comp Neurol* 325(3):327-342.
5.  Jeon CJ, Strettoi E, & Masland RH (1998) The major cell populations of the mouse retina. *J Neurosci* 18(21):8936-8946.
6.  Rapaport DH & Stone J (1984) The area centralis of the retina in the cat and other mammals: focal point for function and development of the visual system. *Neuroscience* 11(2):289-301.
7.  Schiefer U*, et al.* (1999) Angioscotoma detection with fundus-oriented perimetry. A study with dark and bright stimuli of different sizes. *Vision Res* 39(10):1897-1909.
8.  Wassle H, Grunert U, Rohrenbeck J, & Boycott BB (1989) Cortical Magnification Factor and the Ganglion-Cell Density of the Primate Retina. *Nature* 341(6243):643-646.
9.  Chaplin TA, Yu HH, & Rosa MGP (2013) Representation of the visual field in the primary visual area of the marmoset monkey: Magnification factors, point-image size, and proportionality to retinal ganglion cell density. *J Comp Neurol* 521(5):1001-1019.
10. Albus K (1975) Quantitative Study of Projection Area of Central and Paracentral Visual-Field in Area-17 of Cat .2. Spatial Organization of Orientation Domain. *Exp Brain Res* 24(2):181-202.
11. Wilson JR & Sherman SM (1976) Receptive-Field Characteristics of Neurons in Cat Striate Cortex - Changes with Visual-Field Eccentricity. *J Neurophysiol* 39(3):512-533.
12. Tusa RJ, Palmer LA, & Rosenquist AC (1978) Retinotopic Organization of Area-17 (Striate Cortex) in Cat. *J Comp Neurol* 177(2):213-235.
13. Van Essen DC, Newsome WT, & Maunsell JHR (1984) The Visual-Field Representation in Striate Cortex of the Macaque Monkey - Asymmetries, Anisotropies, and Individual Variability. *Vision Res* 24(5):429-448.
14. van Beest EH*, et al.* (2021) Mouse visual cortex contains a region of enhanced spatial resolution. *Nat Commun* 12(1):4029.

15. Tan L, Ringach DL, & Trachtenberg JT (2022) The Development of Receptive Field Tuning Properties in Mouse Binocular Primary Visual Cortex. *J Neurosci* 42(17):3546-3556.
16. Rhim I, Coello-Reyes G, Ko HK, & Nauhaus I (2017) Maps of cone opsin input to mouse V1 and higher visual areas. *J Neurophysiol* 117(4):1674-1682.
17. Adams DL & Horton JC (2002) Shadows cast by retinal blood vessels mapped in primary visual cortex. *Science* 298(5593):572-576.
18. Sprague WW, Cooper EA, Tosic I, & Banks MS (2015) Stereopsis is adaptive for the natural environment. *Sci Adv* 1(4).
19. Samonds JM, Choi V, & Priebe NJ (2019) Mice Discriminate Stereoscopic Surfaces Without Fixating in Depth. *J Neurosci* 39(41):8024-8037.
20. La Chioma A, Bonhoeffer T, & Hubener M (2019) Area-Specific Mapping of Binocular Disparity across Mouse Visual Cortex. *Curr Biol* 29(17):2954-+.
21. Jonides J, Irwin DE, & Yantis S (1982) Integrating visual information from successive fixations. *Science* 215(4529):192-194.
22. Muller JR, Metha AB, Krauskopf J, & Lennie P (1999) Rapid adaptation in visual cortex to the structure of images. *Science* 285(5432):1405-1408.
23. Priebe NJ, Churchland MM, & Lisberger SG (2002) Constraints on the source of short-term motion adaptation in macaque area MT. I. the role of input and intrinsic mechanisms. *J Neurophysiol* 88(1):354-369.
24. Patterson CA, Wissig SC, & Kohn A (2013) Distinct effects of brief and prolonged adaptation on orientation tuning in primary visual cortex. *J Neurosci* 33(2):532-543.
25. Jin M, Beck JM, & Glickfeld LL (2019) Neuronal Adaptation Reveals a Suboptimal Decoding of Orientation Tuned Populations in the Mouse Visual Cortex. *J Neurosci* 39(20):3867-3881.
26. Jin M & Glickfeld LL (2020) Magnitude, time course, and specificity of rapid adaptation across mouse visual areas. *J Neurophysiol* 124(1):245-258.
27. Nelson SB (1991) Temporal interactions in the cat visual system. I. Orientation-selective suppression in the visual cortex. *J Neurosci* 11(2):344-356.
28. Mitchell JF, Reynolds JH, & Miller CT (2014) Active Vision in Marmosets: A Model System for Visual Neuroscience. *J Neurosci* 34(4):1183-1194.
29. Otero-Millan J, Macknik SL, Langston RE, & Martinez-Conde S (2013) An oculomotor continuum from exploration to fixation. *P Natl Acad Sci USA* 110(15):6175-6180.
30. Ikeda H & Wright MJ (1975) Spatial and Temporal Properties of 'Sustained' and 'Transient' Neurones in Area 17 of the Cat's Visual Cortex. *Exp Brain Res* 22:363-383.
31. Foster KH, Gaska JP, Nagler M, & Pollen DA (1985) Spatial and temporal frequency selectivity of neurones in visual cortical areas V1 and V2 of the macaque monkey. *J Physiol* 365:331-363.
32. Hawken MJ, Shapley RM, & Grosof DH (1996) Temporal-frequency selectivity in monkey visual cortex. *Vis Neurosci* 13(3):477-492.
33. Allison JD, Smith KR, & Bonds AB (2001) Temporal-frequency tuning of cross-orientation suppression in the cat striate cortex. *Vis Neurosci* 18(6):941-948.
34. Webb BS, Dhruv NT, Solomon SG, Tailby C, & Lennie P (2005) Early and late mechanisms of surround suppression in striate cortex of macaque. *J Neurosci* 25(50):11666-11675.
35. Priebe NJ, Lisberger SG, & Movshon JA (2006) Tuning for spatiotemporal frequency and speed in directionally selective neurons of macaque striate cortex. *J Neurosci* 26(11):2941-2950.
36. Niell CM & Stryker MP (2008) Highly selective receptive fields in mouse visual cortex. *J Neurosci* 28(30):7520-7536.
37. Gao EQ, DeAngelis GC, & Burkhalter A (2010) Parallel Input Channels to Mouse Primary Visual Cortex. *J Neurosci* 30(17):5912-5926.
38. Van den Bergh G, Zhang B, Arckens L, & Chino YM (2010) Receptive-field Properties of V1 and V2 Neurons in Mice and Macaque Monkeys. *J Comp Neurol* 518(11):2051-2070.

39. Yu HH*, et al.* (2010) Spatial and temporal frequency tuning in striate cortex: functional uniformity and specializations related to receptive field eccentricity. *Eur J Neurosci* 31(6):1043-1062.
40. Durand S*, et al.* (2016) A Comparison of Visual Response Properties in the Lateral Geniculate Nucleus and Primary Visual Cortex of Awake and Anesthetized Mice. *J Neurosci* 36(48):12144-12156.
41. Alitto HJ & Usrey WM (2004) Influence of contrast on orientation and temporal frequency tuning in ferret primary visual cortex. *J Neurophysiol* 91(6):2797-2808.
42. Camillo D, Ahmadlou M, & Heimel JA (2020) Contrast-Dependence of Temporal Frequency Tuning in Mouse V1. *Front Neurosci* 14:868.
43. Seiple W, Rosen RB, & Garcia PM (2013) Abnormal fixation in individuals with age-related macular degeneration when viewing an image of a face. *Optom Vis Sci* 90(1):45-56.
44. Kwon MY, Nandy AS, & Tjan BS (2013) Rapid and Persistent Adaptability of Human Oculomotor Control in Response to Simulated Central Vision Loss. *Curr Biol* 23(17):1663-1669.
45. Chen D, Otero-Millan J, Kumar P, Shaikh AG, & Ghasia FF (2018) Visual Search in Amblyopia: Abnormal Fixational Eye Movements and Suboptimal Sampling Strategies. *Invest Ophthalmol Vis Sci* 59(11):4506-4517.
46. Tang S*, et al.* (2019) Effects of visual blur on microsaccades during visual exploration. *J Eye Mov Res* 12(6).
47. Jang J, Song M, & Paik SB (2020) Retino-Cortical Mapping Ratio Predicts Columnar and Salt-and-Pepper Organization in Mammalian Visual Cortex. *Cell Rep* 30(10):3270-3279 e3273.
48. Fasoli D & Panzeri S (2019) Mathematical studies of the dynamics of finite-size binary neural networks: A review of recent progress. *Math Biosci Eng* 16(6):8025-8059.
49. Meyer AF, O'Keefe J, & Poort J (2020) Two Distinct Types of Eye-Head Coupling in Freely Moving Mice. *Curr Biol* 30(11):2116-2130 e2116.
50. Michaiel AM, Abe ET, & Niell CM (2020) Dynamics of gaze control during prey capture in freely moving mice. *Elife* 9.
51. Zahler SH, Taylor DE, Wong JY, Adams JM, & Feinberg EH (2021) Superior colliculus drives stimulus-evoked directionally biased saccades and attempted head movements in head-fixed mice. *Elife* 10.
52. Harwood MR, Madelain L, Krauzlis RJ, & Wallman J (2008) The spatial scale of attention strongly modulates saccade latencies. *J Neurophysiol* 99(4):1743-1757.
53. De Vries JP, Azadi R, & Harwood MR (2016) The saccadic size-latency phenomenon explored: Proximal target size is a determining factor in the saccade latency. *Vision Res* 129:87-97.
54. Poletti M, Intoy J, & Rucci M (2020) Accuracy and precision of small saccades. *Sci Rep* 10(1):16097.
55. Segal IY*, et al.* (2015) Decorrelation of retinal response to natural scenes by fixational eye movements. *P Natl Acad Sci USA* 112(10):3110-3115.
56. Kuang XT, Poletti M, Victor JD, & Rucci M (2012) Temporal Encoding of Spatial Information during Active Visual Fixation. *Curr Biol* 22(6):510-514.
57. Mostofi N*, et al.* (2020) Spatiotemporal Content of Saccade Transients. *Curr Biol* 30(20):3999-4008 e3992.
58. Parker PR*, et al.* (2022) A dynamic sequence of visual processing initiated by gaze shifts. *BioRxiv*.
59. Castet E & Masson GS (2000) Motion perception during saccadic eye movements. *Nat Neurosci* 3(2):177-183.
60. Castet E, Jeanjean S, & Masson GS (2002) Motion perception of saccade-induced retinal translation. *Proc Natl Acad Sci U S A* 99(23):15159-15163.
61. Miles FA, Kawano K, & Optican LM (1986) Short-latency ocular following responses of monkey. I. Dependence on temporospatial properties of visual input. *J Neurophysiol* 56(5):1321-1354.

62.     Yarbus AL (1967) *Eye Movements and Vision* (Springer US : Imprint: Springer, Boston, MA) pp XIII, 222 s.
63.     Koch C & Ullman S (1985) Shifts in Selective Visual-Attention - Towards the Underlying Neural Circuitry. *Hum Neurobiol* 4(4):219-227.
64.     Hayhoe M & Ballard D (2005) Eye movements in natural behavior. *Trends Cogn Sci* 9(4):188-194.
65.     Najemnik J & Geisler WS (2005) Optimal eye movement strategies in visual search. *Nature* 434(7031):387-391.
66.     Gegenfurtner KR (2016) The Interaction Between Vision and Eye Movements (dagger). *Perception* 45(12):1333-1357.
67.     Johnson KP*, et al.* (2021) Cell-type-specific binocular vision guides predation in mice. *Neuron* 109(9):1527-1539 e1524.
68.     Groner MT, Groner R, & von Muhlenen A (2008) The effect of spatial frequency content on parameters of eye movements. *Psychol Res-Psych Fo* 72(6):601-608.
69.     Intoy J & Rucci M (2020) Finely tuned eye movements enhance visual acuity. *Nat Commun* 11(1):795.
70.     Tatler BW, Baddeley RJ, & Vincent BT (2006) The long and the short of it: spatial statistics at fixation vary with saccade amplitude and task. *Vision Res* 46(12):1857-1862.
71.     Loh Z, Hall EH, Cronin D, & Henderson JM (2022) Working memory control predicts fixation duration in scene-viewing. *Psychol Res*.
72.     Niemeier M, Crawford JD, & Tweed DB (2003) Optimal transsaccadic integration explains distorted spatial perception. *Nature* 422(6927):76-80.
73.     Oostwoud Wijdenes L, Marshall L, & Bays PM (2015) Evidence for Optimal Integration of Visual Feature Representations across Saccades. *J Neurosci* 35(28):10146-10153.
74.     Ganmor E, Landy MS, & Simoncelli EP (2015) Near-optimal integration of orientation information across saccades. *J Vis* 15(16):8.
75.     Henderson JM & Hollingworth A (2003) Eye movements and visual memory: detecting changes to saccade targets in scenes. *Percept Psychophys* 65(1):58-71.
76.     Scholl B, Pattadkal JJ, Dilly GA, Priebe NJ, & Zemelman BV (2015) Local Integration Accounts for Weak Selectivity of Mouse Neocortical Parvalbumin Interneurons. *Neuron* 87(2):424-436.
77.     Choi V & Priebe NJ (2020) Interocular velocity cues elicit vergence eye movements in mice. *J Neurophysiol* 124(2):623-633.
78.     Pelli DG (1997) The VideoToolbox software for visual psychophysics: transforming numbers into movies. *Spat Vis* 10(4):437-442.
79.     Brainard DH (1997) The psychophysics toolbox. *Spatial Vision* 10(4):433-436.
80.     Cornelissen FW, Peters EM, & Palmer J (2002) The Eyelink Toolbox: eye tracking with MATLAB and the Psychophysics Toolbox. *Behav Res Methods Instrum Comput* 34(4):613-617.
81.     Olmos A & Kingdom FAA (2004) A biologically inspired algorithm for the recovery of shading and reflectance images. *Perception* 33(12):1463-1473.
82.     Mathis A*, et al.* (2018) DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat Neurosci* 21(9):1281-1289.
83.     Mitchell JF, Priebe NJ, & Miller CT (2015) Motion dependence of smooth pursuit eye movements in the marmoset. *J Neurophysiol* 113(10):3954-3960.
84.     Menzel CR & Menzel EW (1980) Head-Cocking and Visual Exploration in Marmosets (Sanguinus fuscicollis). *Behaviour* 75(3/4):219-234.
85.     Umino Y, Pasquale R, & Solessio E (2018) Visual Temporal Contrast Sensitivity in the Behaving Mouse Shares Fundamental Properties with Human Psychophysics. *Eneuro* 5(4).
86.     Scholl B, Pattadkal JJ, & Priebe NJ (2017) Binocular Disparity Selectivity Weakened after Monocular Deprivation in Mouse V1. *J Neurosci* 37(27):6517-6526.
87.     Lee D & Malpeli JG (1998) Effects of saccades on the activity of neurons in the cat lateral geniculate nucleus. *J Neurophysiol* 79(2):922-936.
88.     Kording KP, Kayser C, Betsch BY, & Konig P (2001) Non-contact eye-tracking on cats. *J Neurosci Meth* 110(1-2):103-111.

89.    Roth MM*, et al.* (2016) Thalamic nuclei convey diverse contextual information to layer 1 of visual cortex. *Nat Neurosci* 19(2):299-+.
90.    Miller DJ, Balaram P, Young NA, & Kaas JH (2014) Three counting methods agree on cell and neuron number in chimpanzee primary visual cortex. *Front Neuroanat* 8:36.
91.    O'Kusky J & Colonnier M (1982) A laminar analysis of the number of neurons, glia, and synapses in the adult cortex (area 17) of adult macaque monkeys. *J Comp Neurol* 210(3):278-290.
92.    Atapour N*, et al.* (2019) Neuronal Distribution Across the Cerebral Cortex of the Marmoset Monkey (Callithrix jacchus). *Cereb Cortex* 29(9):3836-3863.
93.    Keller D, Ero C, & Markram H (2018) Cell Densities in the Mouse Brain: A Systematic Review. *Front Neuroanat* 12:83.
94.    Dougherty RF*, et al.* (2003) Visual field representations and locations of visual areas V1/2/3 in human visual cortex. *J Vis* 3(10):586-598.
95.    DeFelipe J (2011) The evolution of the brain, the human nature of cortical circuits, and intellectual creativity. *Front Neuroanat* 5.
96.    Solomon SG & Rosa MG (2014) A simpler primate brain: the visual system of the marmoset monkey. *Front Neural Circuits* 8:96.
97.    Balaram P & Kaas JH (2014) Towards a unified scheme of cortical lamination for primary visual cortex across primates: insights from NeuN and VGLUT2 immunoreactivity. *Front Neuroanat* 8:81.
98.    Garrett ME, Nauhaus I, Marshel JH, & Callaway EM (2014) Topography and areal organization of mouse visual cortex. *J Neurosci* 34(37):12587-12600.

**Figures**

**Figure 1.** Saccade rates increase with decreasing saccade size matching faster recovery from adaptation for smaller receptive fields. a, b, Sequential fixations change receptive field inputs and differences in adaptation recovery time constants allow primates (a) to make more frequent saccades compared to mice (b). c, Top row: example macaque V1 responses to two repeated presentations of the same stimulus (28). Black represents neural activity to the first presentation and different shades of blue depict the neural activity after the second presentation in progressively more delayed conditions. Bottom row: example mouse V1 responses to a similar experiment[31]. See model examples in Supplemental Fig. 1. d, Magnitude of initial responses for different saccade rates for macaques and mice. For large saccades, the stimulus changed randomly after a saccade, while for small saccades, the stimulus stayed the same after a saccade. e, Saccade rates for a range of recovery time constants that maintain 75% of initial responses. f, Saccade rate versus size distributions for five species. Each solid point represents the medians of a single subject. Open circles of marmosets (33), macaques (33), and cats (86, 87) represent previously published data from other laboratories (see Methods). The large outline represents the central half of the distribution (between the 25th and 75th percentiles) and the small outline represents the standard error of the median for the distribution of all subjects. Due to the large number of samples, the standard error of the median outlines are smaller than even the data points. See example full distributions in Supplemental Fig. 2. g, Preferred temporal frequency versus receptive field size distributions from previously published data (see Methods) (8, 35-45). The open circle represents the median and the large outline represents the central half of the distribution (between the 25th and 75th percentiles). h, Saccade size increases (8) and saccade rate decreases in mice with monocular deprivation during the critical period (N = 4 mice: n = 1448 saccades) compared to control mice (N = 3 mice: n = 1586 saccades).

**Figure 2.** Fixations over time provide similar visual processing coverage across different mammals. a, Computing average receptive field coverage over time from saccades. b, Horizontal cross sections of receptive field coverage over time for each mammal. Lighter colors are data obtained when subjects were viewing small images. c, Relative brain sizes between each mammal. d, Horizontal coverage data from b multiplied by the average two-dimensional V1 cell density for each mammal. Human data are replotted with marmoset and mouse data to facilitate comparison.

**a** Primate

Sequential Fixations          Adaptation

Fast Recovery <150 ms

V1 Response

**b** Mouse

Sequential Fixations          Adaptation

Slow Recovery > 1s

V1 Response

**c**

Macaque

$\tau_{recovery}$ = 73 ms

100 ms

Mouse

100 ms

Response (spikes/s)

**d**

Initial Response at Fixation (spikes/s)

— Large Saccades
- - - Small Saccades

Macaque

Mouse

Maximum Adaptation

Saccade Rate (1/s)

**e**

Predicted Saccade Rate (1/s)

Macaque

Mouse

Recovery Time Constant (ms)

**f**

Motor System

Median Saccade Rate (1/s)

— Human
— Macaque
— Marmoset
— Cat
— Mouse

Saccade Size (degrees)

**g**

Visual System

Preferred Temporal Frequency (1/s)

Receptive Field Size (degrees)

**h**

Normalized Number of Saccades/Intervals

— Control Mice
— MD Mice

Saccade Size (degrees)

Normalized Number of Saccades/Intervals

Saccade Rate (1/s)