Genomic adaptation underlies nutrient-driven reprogramming in the malaria-causing parasite

Qian Li[1,4], Leonie Vetter[1], Cagla Sahin[1], Ákos Végvári[2], Ulf Ribacke[1], Mats Wahlgren[1], Ola Larsson[3], Sherwin Chun-Leung Chan[3,4*]

[1] Department of Microbiology, Tumor and Cell Biology (MTC), Karolinska Institutet, Stockholm, Sweden
[2] Division of Chemistry I, Department of Medical Biochemistry and Biophysics (MBB), Karolinska Institutet, Stockholm, Sweden
[3] Department of Oncology-Pathology, Science for Life Laboratories, Karolinska Institutet, Stockholm, Sweden
[4] These authors contributed equally

*Correspondence: Sherwin Chun-Leung Chan

Abstract (Summary)

Plasmodium falciparum has an AT-rich genome with highly biased codon and amino acid usage that is not balanced by the tRNA genes repertoire. To determine whether individual tRNA genes can be differentially expressed to compensate for the tRNA-codon imbalance, we characterized the tRNA profiles of the parasite. While we found that the parasite stably maintained a biased expression of different tRNAs throughout the intraerythrocytic cycle (IDC), the tRNA-codon imbalance remains. Specifically, the requirements for a limited tRNA subset in the codons of a gene negatively correlated with its mRNA level. A detailed analysis revealed that these observations implicate host adaptation during *P. falciparum* genome evolution, whereby housekeeping genes evolved to minimize dependence on exogenous amino acids, allowing these genes to mitigate the negative effects during acute nutrient deprivation. We found that these host-adapted features can derive a decentralized nutrient-dependent regulatory programme via sensing isoleucine as a proxy for nutrient stress, which causes a reduction in aminoacylated Ile-tRNA level. This, in turn, rapidly reprogrammes the global protein output and transcriptome in a targeted fashion. Our findings provide strong evidence showing that the primary sequences of proteins underlie gene expression programmes, suggesting new perspectives on protein evolution.

Introduction

Plasmodium falciparum is a protozoan that causes malaria, which imposes a significant global health burden. The parasite genome is AT-rich (81%), which is reflected by a high frequency of AT-rich codons and biased in amino acid (AA) usage[1,2]. The number of tRNA gene copies is usually positively correlated with the codon usage in unicellular organisms[3]. In *P. falciparum, however,* tRNA genes are nonredundant. This raises the question of how the parasite compensates for its codon bias during protein synthesis. Due to extensive posttranscriptional modifications and redundancy in most organisms, profiling tRNAs requires tailored sequencing protocols which include ARM-seq, hydro-tRNAseq and mim-tRNAseq[4-8]. These protocols allow simultaneous quantifications of tRNA levels, posttranscriptional modifications and aminoacylation ("charge") status[9]. In contrast to the commonly held belief that tRNAs are stably expressed house-keeping molecules, the application of these methods has revealed dynamic tRNA expression across tissues and cell lines[10-12]. Dynamic alterations in the levels of individual tRNAs may modulate protein synthesis by affecting translation elongation. Although rarely explored, this regulatory mechanism has been implicated in cell-fate decisions and pathological conditions[11,13,14]. Intriguingly, regulation of tRNA expression has been reported with changes in growth conditions and stress levels[15-17]. In *P. falciparum*, nutrient stress induces Asp-tRNA cleavage, which may promote gametocytogenesis[18], while MAF1-dependent inhibition of tRNA transcription is essential for mounting an effective response to AA starvation[19]. In addition to these examples, stress sensing and downstream responses in *P. falciparum* remain poorly understood, in part due to the absence of the conserved mTOR pathway[20], which integrates a multitude of intracellular and extracellular cues in other organisms[21]. Moreover, although AA and glucose starvation can be sensed in *P. falciparum* via the conserved GCN2-dependent phosphorylation of eIF2α and a genus-specific serine/threonine kinase, KIN, respectively[22-24], AA starvation is still effectively sensed in the absence of GCN2 or when only a nonphopsphorylatable eIF2α variant is expressed[23]. This suggests additional stress sensing mechanisms in *P. falciparum.* We therefore sought to understand how translation is regulated in *P. falciparum* under nutrient stress.

Results

P. falciparum maintains an imbalanced tRNA-codon profile throughout the IDC

The lack of correlation between codon usage and tRNA gene copy number in *P. falciparum* suggests a disparity in tRNA availability and demand (highlighted by the demand for Asn-GTT~14.41% vs Pro-CGG~0.09%)[25] that is not seen in *Toxoplasma gondii*, a related apicomplexan (Extended Data Fig. 1). To assess whether tRNA dynamics and/or expression may compensate for this low correlation, we applied an improved TGIRT-seq protocol to profile mature tRNAs (the 45 nuclear tRNAs are referred to as the anti-codon pool or tRNA supply) throughout the IDC (Ring, 8-12 hours post invasion (h.p.i.); Trophozoite, 24-28 h.p.i.; Schizont, 40-44 h.p.i.) with a high mapping ratio (Extended Data Fig. 2a-b)[8]. Approximately 80% of the RNA sequencing reads mapped to tRNA genes retained full 5' to 3' coverage (Extended Data Fig. 2c), with only a small fraction mapped to host tRNAs (from 8.7% in Ring to 0.85% in Schizont) (Extended Data Fig. 3a). These profiles showed substantial tRNA abundance variation among tRNA species (Fig. 1a). In particular, all Ser-tRNAs, which all carry an A-box element that diverges from the consensus sequence (Extended Data Fig. 3b)[27], appeared to be highly abundant (28.7-35.4% of the anti-codon pools). However, in contrast to the cyclic pattern of the coding transcripts[28], minimal differences in the anticodon pools were found across stages (Fig. 1a). Furthermore, the discrepancy in charged tRNA ratio throughout the IDC was minimal, so that a relatively high uncharged-to-charged tRNA ratio was maintained (Fig. 1b, Extended Data Fig. 4). We also quantified fluctuations in mRNA expression-weighted tRNA usage (referred to as the codon pool or tRNA demand). The results

revealed that despite stage-dependent transcriptions (Extended Data Fig. 3c), the codon pools
exhibited limited fluctuation (Fig. 1c, Extended Data Fig. 3d). As pools of both anticodons and
codons were relatively constant, the codon composition of a transcript determines its
accessibility to the required anticodon pool throughout the IDC. Surprisingly, the anticodon
pools correlated poorly with the codon pools, which allowed us to identify a subset of tRNAs
that were either under- or overrepresented (Fig. 1d). Since isoacceptors for a specific AA are
often found concertedly to be under- or overrepresented, this observation suggested that AA
composition, in addition to codon usage, may influence translation efficiency of a transcript.
140
tRNA modification profiles exhibit limited variations throughout the IDC
tRNA posttranscriptional modifications can directly and indirectly affect translation[16,29]. We
assessed whether these modifications can be dynamically modulated throughout the IDC
despite the stable anticodon pool. Some modifications frequently cause position-specific
mutations[6], and these mutation signatures allow *a priori* inference about the modification status
at reported tRNA-modification sites[5,6,30]. Using this strategy, we generated a tRNA-
modification atlas including m^1A on A9 and A58, m^1G on G9 and G37, $m^{2,2}G$ on G26, m^3C on
C32, and inosine on A34 (I34) in *P. falciparum* (Extended Data Fig. 5a,c & 6a). As a validation
for the inferences, m^1A, m^1G and m^3C were found to be sensitive to *E. coli* AlkB demethylase
treatment, as previously reported (Extended Data Fig. 6b)[5,6,8]. Interestingly, we observed A59
mutations in four Leu-tRNAs and mutations at the discriminator U73 base in all Thr- and Lys-
tRNAs (Extended Data Fig. 5a,c). Although identities of these signatures cannot be inferred, as
no relevant annotations have been deposited in MODOMICS[31], the A59 signatures were
sensitive to demethylase treatment, suggesting that A59 is methylated (Extended Data Fig. 6b).
While the same types of modification are often found on isoacceptors for a specific AA, some
isoacceptors exhibit distinct modification patterns among themselves (Extended Data Fig. 5a;
e.g. m^1G37 in Ile-tRNAs). Modifications on the anticodon loop (C32 and G37), which can
directly influence translation dynamics, generally exhibit lower stoichiometry (Extended Data
Fig. 5a)[16,32,33].
Furthermore, we performed bisulfite (BS) conversion to estimate the abundance of m^5C
(Extended Data Fig. 5b). Positions with heavily deposited m^5C included the conserved C38 of
Asp-GUC and the variable loops of most tRNAs. Clusters of m^5C were observed at low
stoichiometry on both sides of the T-stem of many tRNAs (Extended Data Fig. 5b,c).
Meanwhile, m^1G37 is the only modification detected on apicoplast tRNAs (Extended Data
Fig.7a). However, a frequent truncation at the wobble position of one Met-tRNA strongly
suggests K^2C34 modification, which would convert the codon specificity from AUG (Met) to
AUA (Ile) (Extended Data Fig. 7b-d)[34,35].
Notably, we only observed a modest global reduction of modifications in trophozoites
(Extended Data Fig. 6c). To clarify this finding in more detail, we specifically examined Tyr-
GUA. Because Tyr-GUA is the only tRNA that splices an 11-nt intron from pre-tRNA, the ratio
of unspliced-to-spliced Tyr-GUA reads was used to estimate nascent tRNA transcription. As
expected from their higher transcriptional activity, trophozoites exhibited higher levels of
unspliced Tyr-GUA than other stages (Extended Data Fig. 6d). Furthermore, the unspliced
reads harbored less abundant modifications at all sites, with m^1G37 completely absent
(Extended Data Fig. 6d). Therefore, the global reduction in tRNA modification abundance is
likely a result of the relatively high abundance of pre-tRNAs in trophozoites, suggesting the
lack of systemic modulation of these modifications during stage progression. Together, these
findings support the notion that the codon composition of a transcript can primarily affect
translation efficiency throughout the IDC.
180

Highly expressed genes minimize their dependence on AAs with lower usage frequency in host haemoglobin

To further explore the relationship between the codon and the anticodon pools, we sought to identify sources of variation in tRNA demand across genes. To this end, we used principal component analysis with data describing the relative demand for each tRNA across all transcripts (i.e. after normalizing the tRNA demand per transcript). The resulting PCA suggested that nine tRNAs deviated in principal component 1 (PC1; which explains 80.4% total variance; Fig. 2a) and thus showed the greatest frequency variation across transcripts (among the highest in combined contribution in PC1 and PC2). Notably, their corresponding codons account for a major fraction of the codon pool, but the supply of these tRNAs was underrepresented in the anticodon pool (Fig. 2b; i.e. they belong to the "underrepresented" tRNA group identified above). As the supply of these tRNAs was relatively restricted, variation in the transcripts' demand of them may affect translation efficiency. Interestingly, genes with the highest mRNA levels was associated with a lower demand for six of these tRNAs (Asn-GUU, Ile-AAU, Ile-UAU, Lys-UUU, Phe-GAA and Tyr-GUA) compared to genes with the lowest mRNA levels (Fig. 2c,d). Moreover, transcripts encoding highly abundant ribosomal proteins show reduced demand for these tRNAs, except for Lys-UUU (Fig. 2c). Finally, the demand for these tRNAs was negatively correlated with the gene expression throughout the IDC (Extended Data Fig. 8a,b). Accordingly, to bolster their translation, highly expressed genes may have evolved to avoid competing for these decoding-limiting tRNAs. Perplexingly, five of these tRNAs exclusively decode asparagine, isoleucine, phenylalanine and tyrosine. Concomitantly, the AA composition of proteins translated from these highly abundant transcripts is low in these AAs. Consistent with this finding, the frequencies of these AAs are negatively correlated with mRNA levels (Extended Data Fig. 8c), suggesting the possibility that the AA composition influences translational efficiency.

In the IDC, *P. falciparum* digests host haemoglobin (HB) to acquire most of the AAs needed for growth[36]. Human HB, however, lacks isoleucine (Ile), which the parasite takes up from the external milieu[23]. Intriguingly, the usage frequency of the two Ile-tRNAs was negatively correlated with the mRNA level, and among all AAs, Ile AA frequency showed the strongest negative correlation with mRNA level (Fig. 2c,e, Extended Data Fig. 8a-c). We therefore hypothesized that highly expressed genes in *P. falciparum* may have evolved to reduce their dependence on exogenous AA as a feature of host-adaptation, which allows their expression to be unperturbed during external nutrient deprivation. To test this hypothesis, we compared the AA demand in the codon pools to the AA composition of HB (considered a stable endogenous AA supplier). This revealed that the AAs showing the strongest negative correlation between demand and mRNA level (Tyr [Y], Ile [I] and Asn [N] in Extended Data Fig. 8c) were inadequately supplied by HB (Fig. 2f; Y, I and N are indicated in red and fall below the identity line). Furthermore, the AA composition encoded by genes with the highest mRNA levels showed higher Spearman correlation to the AA composition of HB compared to genes with the lowest mRNA levels (Fig. 2g), a pattern that was generalizable to all the coding genes (Fig. 2h). These analyses support that highly expressed transcripts encode an AA composition which corresponds well with the HB AA composition and the proteins encoded by these transcripts are likely to be more subsistent under nutrient stress, during which AAs supply will increasingly depends on AAs released from HB digestion. Accordingly, this intricate interplay between AA availability and demand may facilitate translation of mRNA subsets in response to exogenous nutrient stress by the parasite.

Isoleucine availability serves as a proxy for external nutrient stress

We next sought to study whether AA availability under stress conditions may selectively control protein synthesis. To this end, we first assessed the tRNA profile after acute depletion

of AA in the medium, which forced the parasites to acquire AAs only from HB digestion. Six
hours of AA depletion in late trophozoites (32-36 hpi) resulted in a reduction in the level of
charged Ile-tRNAs (Ile-AAU -53% and Ile-UAU -19%) (Fig. 3a), but did not induce significant
changes in the composition or modification of the anticodon pool (Extended Data Fig. 9a,b).
This decrease was milder than might be expected by the complete lack of exogenous Ile,
suggesting that Ile was recycled from the degradation of endogenous proteins. However, the
level of charged Ile-tRNAs was not reduced if elongation was also blockaded by halofuginone,
which is a prolyl tRNA synthetase inhibitor that abolishes aminoacylation of all Pro-tRNAs
(>85%) (Fig. 3a)[37]. This is consistent with depletion of charged Ile-tRNAs upon an unbalanced
demand and supply of Ile during translation elongation. Interestingly, the reduction of charged
Ile-AAU, which is the preferred Ile-tRNA used by highly expressed genes (Extended Data Fig.
9c), was more pronounced than that of Ile-UAU (Fig. 3a). The higher depletion rate of Ile-
AAU may imply that highly expressed genes were still actively, and might have been
selectively, translated during starvation. Accordingly, our data consolidate isoleucine as an
essential exogenous AA for *P. falciparum* development and that isoleucine availability may be
key to nutrient-sensing in this parasite[23,38].
247
Stage progression and parasite multiplication rates were sensitive to the isoleucine level
Next, we sought to examine biological processes that may be affected by Ile withdrawal (such
as that during deprivation of the exogenous AA supply). Delay in stage progression is
commonly observed upon prolonged AA starvation; however, the parasite can reportedly
recover and resume normal progression upon nutrient resupply, reminiscing a "hibernatory
state"[23,38]. We reaffirmed the stage-delay phenotype even after a short duration of AA depletion
(6 hours, Extended Data Fig. 10). Notably, we found that genes with upregulated mRNA
expression during the ring-to-trophozoite and the trophozoite-to-schizont transitions showed
higher average Ile-encoding content than genes with downregulated mRNA expression, while
the opposite pattern was observed when the schizont-to-ring transition was analysed (Fig. 3b,
Extended Data Fig. 11a,b). Furthermore, genes with mRNA levels that were the most variable
throughout the IDC encoded a higher Ile content than genes with mRNA expression showing
the least variation between stages (Extended Data Fig. 11c). The usage discrepancy, however,
are not specific for isoleucine, since leucine, which exhibits biochemical properties similar to
those of Ile, displays similar trends. Yet overall, this analysis provides a rationale for the stage
delay phenotype and the higher sensitivity to Ile withdrawal reported in pre-S phase compared
to post-S phase parasites[38].
Meanwhile, stress parasites consistently multiplied less during the immediate invasion cycle
than the unstressed parasites. We observed the differences despite the stressed parasites had
fully recovered to normal progression. Furthermore, Ile supplementation during the acute stress
period could restore the multiplication rate in a dose-dependent manner (Fig. 3c). Specifically,
a reduction in multiplication rate was observable only under starvation conditions in the late
trophozoite stage, but not in the early stage (8-12 hours) (Fig. 3c). These combined to suggest
trophozoites-specific biological processes were affected by acute AA depletion, which led to
reduced proliferation and failure to achieve full recovery.
To explore the underlying reasons for these effects, we performed a quantitative proteomic
analysis and found that 56 out of the 78 downregulated proteins (56%) in the starved late
trophozoites were merozoite-related proteins, which mostly associate to membrane and
organellar compartments, particularly the inner membrane complex (Fig. 3d, Supplementary
table 1). Interestingly, gene ontology (GO) analysis of the 5% genes with the highest Ile content
(Ile-rich, "IR") revealed enrichment of lipid metabolic pathways (Fig 3e). In contrast, genes
with the highest leucine content, which is also decoded by AT-rich codons, show no pathway
enrichment. Since lipid metabolic pathways are fundamental to membrane biogenesis needed

for merozoite production during schizogony, AA depletion, which abrogates the Ile supply, can
hamper the efficient translation of IR transcripts and eventually caused a reduction in
merozoites number and multiplication rate. Strikingly, >80% (43/53) of conserved IR genes
(defined as having >100 orthologues) encode an Ile content that is >3 standard deviations higher
than the mean Ile content in the respective orthologue groups (Fig. 3f)[39], suggesting that the
high Ile content encoded by IR genes is unlikely due to functional conservation in their proteins,
because if this is the cause, high Ile content would have been universally retained in their
orthologues, as is the case for genes encoding leucine-rich proteins (Fig. 3f; only 3/55 genes
have a leucine content >3 S.D. than the mean). The analysis provides a strong basis that the
high Ile content encoded by IR genes may have evolved uniquely in *P. falciparum* as an
adaptation.
292
Stress-mediated reprogramming is defined by isoleucine usage
To assess the impact of AA starvation on translation, we performed polysome-profiling, which
involves fractionation of mRNA on the basis of its ribosome-association. AA starvation led to
a reduction in mRNA associated with multiple ribosomes (polysomes) with a concomitant
increase in the 80S monosome peak (Fig. 4a). This profile is commonly observed in eukaryotes
under stress conditions and reflects the global suppression of translation initiation due to eIF2α
phosphorylation[40,41]. To determine how individual mRNAs are regulated upon AA-starvation,
we first performed RNA-sequencing to quantify polysome-associated mRNA. As 6h of AA
starvation led to a reduction in charged Ile-tRNAs, we compared Ile usage between the 247 and
152 transcripts that showed increased and decreased polysome-association, respectively,
following AA starvation (Fig. 4b; FDR<0.05 and $|\log_2$ fold-change$| > 1$). Indicative of increased
frequency of stalled Ile codon translation during elongation, transcripts with increased
polysome association showed a higher Ile demand than those with decreased polysome
association (Fig. 4c). Consistently, whereas changes in polysome-associated mRNA and
protein levels showed a substantial correlation for transcripts with decreased polysome
association (the larger subset with a $\log_2$ fold-change <0 for polysome-associated mRNA), the
corresponding correlation was lower for transcripts with increased polysome association (Fig.
4d). Therefore, increased ribosome stallings, but not translation efficiency, likely explains the
increase in polysome association.
The Kozak context of a start codon is a major determinant of the translation rate. Surprisingly,
the 247 transcripts with increased polysome association showed a Kozak contexts than the 152
with decreased polysome association (Fig. 4e, Extended Data Fig. 12a)[42]. Ribosome collisions
driven by excessive ribosome stalling can trigger the conserved No-Go-Decay (NGD) pathway,
leading to degradation of transcripts with collided ribosomes[43]. Therefore, transcripts with a
higher demand for Ile-tRNAs may have evolved to prevent ribosome collision by acquiring
lower translation initiation potential and thus escape NGD surveillance[44]. Hypothetically, this
would permit Ile-dependent stabilization of transcripts, which may then contribute to global
transcriptome reprogramming. To assess this possibility, we quantified total mRNA in samples
collected in parallel with the polysome-associated mRNA samples described above and
identified 555 and 353 genes with increased and decreased mRNA levels, respectively, after
AA starvation (Extended Data Fig. 12b). Because AA starvation delays stage progression,
starved parasites were expected to be less abundant in transcripts that are typically upregulated
in schizonts compared to trophozoites and vice versa. We thus categorized differentially
expressed genes as stage dependent and stage independent (Extended Data Fig. 12c). Changes
in expression of stage dependent genes are expected due to delayed stage progression, whereas
stage-independent changes would represent transcripts associated with the stress-
reprogramming. Among the stage-independent changes, higher Ile usage and a weaker initiation
context were identified in parasites with increased mRNA levels compared to those with

decreased mRNA levels. In contrast, no difference in Ile usage and only a minor difference in
initiation context were found when stage-dependent changes were compared among these
groups (Fig. 4f,g, Extended Data Fig. 12d). The shared features associated with polysome-
associated mRNA and stage-independent total mRNA changes suggest crosstalk between
polysome association and mRNA levels. Indeed, a stronger correlation was found between
changes in polysome-associated and total mRNA among mRNAs with increased polysome
association compared to those with decreased polysome association (Fig. 4h, Extended Data
Fig. 12e). This finding supports the idea that increased polysome association stabilizes
transcripts. Furthermore, many transcripts were not regulated as expected due to delayed stage
progression (Extended Data Fig. 12c), suggesting that these transcripts may have been
stabilized (i.e., they were expected to be downregulated but were unchanged) or destabilized
(i.e., they were expected to be upregulated but were unchanged) by reprogramming.
Accordingly, the stabilized transcripts also showed higher Ile usage and weaker initiation
context than the destabilized transcripts (Extended Data Fig. 12f,g), These features are
associated with increased polysome association and point to a stabilizing effect due to increased
polysome association. Interestingly, many transcripts encoding ribosomal and proteasomal
proteins were found in the subset of stage-independent downregulated genes (Extended Data
Fig. 12h). These changes may lead to reduced proteome turnover and potentially linked to the
"hibernatory state" reported during prolonged AA starvation[23]. In contrast, upregulated stage-
independent genes were enriched in several metabolic pathways (Extended Data Fig. 12i). As
upregulation may depend on transcript stability but not necessarily increased protein output,
these changes may prime a cell for rapid metabolic reactivation after nutrient stimulation.
Collectively, our data suggest that transcripts can be stabilized following ribosome stalling in
combination with averted ribosome collision, which would otherwise trigger NGD surveillance.
We used anota2seq to further explore the relationship between polysome-associated and total
mRNA. Anota2seq identifies changes in polysome-associated mRNA occurring independent of
changes in total mRNA (denoted "translation"); congruent changes in total and polysome-
associated mRNA ("abundance"); and changes in total mRNA not reflected by corresponding
changes in polysome-association ("buffering")[45]. Notably, anota2seq analysis revealed many
buffered transcripts (Extended Data Fig. 13a). Technically, buffered transcripts with increased
("buffered, mRNA up") or decreased ("buffered, mRNA down") total transcript levels
correspond to transcripts with relatively decreased or increased ribosome association,
respectively. Buffered transcripts therefore exhibit a discordance between ribosome association
and total mRNA levels and therefore would be expected to show differed RNA features
compared to those of the aforementioned transcripts with altered polysome association (Fig.
4b-d). Indeed, genes in the "buffered, mRNA down" subset have higher Ile usage than genes in
the "buffered, mRNA up" subset (Extended Data Fig. 13b), as well as having stronger Kozak
initiation context (Extended Data Fig. 13c). Tentatively, we reason that despite the enrichment
of ribosomes on Ile-codons, mRNAs that are reduced but buffered are destabilized due to the
use of strong initiation context, which lead to ribosome collision and NGD degradation.
As stalling at Ile codon is instrumental to the observed transcriptome reprogramming, we
believe that the expansion of homo-Ile repeats, which are potent stalling motifs, would have
been under purifying selection. Without functional constraint, AA repeats often appear at a
higher frequency than expected by chance due to replication slippage[46]. In contrast, di-Ile and
all successive expansions of Ile unit in *P. falciparum* genome occur at a frequency lower than
expected by chance, which is not the case in other hydrophobic AAs (Extended Data Fig. 14).
Therefore, occurrence of Ile repeats are likely under non-neutral selection, further asserting a
central role for Ile in stress-mediated reprogramming.
In summary, we proposed a mechanism in which the parasite senses nutrient stress through a
change in the charged-to-uncharged ratio of Ile-tRNAs. The stress-induced accumulation of

uncharged Ile-tRNAs not only induces eIF2α phosphorylation and repressed translation
initiation, but also allows targeted proteome and transcriptome reprogramming due to its
uniquely evolved genome (Fig. 4i).
384

Similar cases of adaptive evolution may be conserved in other organisms
Our analyses with *P. falciparum* reveal an evolutionary selection that influenced the protein
sequences on a genome-wide scale. This selection strongly implicates a feature of metabolic
adaptation to its HB dependence. We may therefore expect other *Plasmodium* spp., which also
digest HB, to show a similar evolutionary selection. However, the GC-rich genome of *P.vivax*,
for example, does not favor Ile codons and would apparently contradict the idea of a selection
that is primarily reliant on Ile. To investigate the generalizability of the adaptation, we
compared AA compositions of the components of glycolysis and the pentose phosphate
pathway (PPP) to the AA composition of HB in several organisms with different genome GC-
contents. Both pathways are highly conserved and use glucose-6-phosphate as input; however,
glycolysis generates ATP, while PPP supplies ribose-5-phosphates for nucleic acid synthesis[47].
Therefore, although these pathways compete for a common input, glycolysis is essential for
energy homeostasis, whereas PPP favours proliferation. Importantly, our results indicate that
glycolysis pathway components of all the investigated *Plasmodium* and *Babesia* spp., which
are RBC-resident parasites[36], show a higher correlation to host HB than the correlation shown
by the components of the PPP (Fig. 5). This finding supports an adaptation that leads to the
stronger harmonization of the AA usage in housekeeping glycolytic enzymes to the stable
nutrient source. It implies that synthesis of the glycolytic pathway maybe prioritized under
exogenous nutrient stress. Interestingly, despite being RBC-resident, *Theileria* spp. do not
exhibit this discrepancy (Fig. 5). Schizogy in *Theileria* exclusively takes place in nucleated
blood cells[48], but not in RBCs, which may have reduced the selection force needed to drive
similar evolution. Indeed, *Babesia* and *Theileria* share a high lineage relatedness and the
opposite observations between the two genera suggest that the selection is dynamic and depends
on the parasitism mode.
409

Discussion
P. falciparum genome presents a peculiar feature in which the extreme codon and AA usage
bias is not balanced by an expansion of tRNA genes. We studied this feature by simultaneously
followed the dynamics of codon and anticodon pools using tailored tRNA-seq. While detection
sensitivities may vary among tRNAs due to sequence-specific biases, such as differences in
posttranscriptional modifications, we found that the supply of anticodon correlates poorly to
the demand of the codon pool, reducing the availability of cognate tRNAs to certain codons.
Although our method enabled only an assessment of the relative compositions, but not the
absolute abundance, between the codon and the anticodon pools, our study concludes that *P.*
falciparum has evolved to exploit the translation elongation process as an important regulatory
mechanism. This is a sound strategy because the parasite encodes few regulators for
transcriptional control and, compared to other eukaryotes, has longer average coding sequences
to accentuate the relative importance of translation elongations in determining protein output
rate[1,49]. Indeed, our previous study also concluded that the excess usage of GU wobble codons
in the parasite genome serves to fine-tune but not to maximize global protein output[25].
In this study, we identified sweeping selection that shaped the evolution of *P. falciparum*
proteome, characterizing it as an adaptation to the lifestyle of parasitizing the RBCs. Proteins
with housekeeping function evolved to rely mainly on a stable nutrient supply provided by host
haemoglobins and minimize dependency on external nutrients. Therefore, translation of these
proteins can be buffered during acute nutrient deprivation. Phosphorylation of eIF2α is the only
characterized response pathway to AA starvation, however, the parasite can still mount an

effective response and recovered from AA starvation in the absence of phosphorylatable eIF2α
[23,24]. Therefore, an alternative stress response mechanism has long been expected. Our findings
suggest that the proteome architecture itself derives such an alternative mechanism through
which the parasite can make a decentralized, rapid, targeted and graded response by simply
sensing the isoleucine level, which is a proxy for AA starvation, resulting in a dose-dependent
autoregulation of its multiplication rate. These features bear remarkable resemblance to a
market economy in human society, where efficient resource allocation is based solely on
demand and supply.
Notably, isoleucine is the third-most encoded AA in *P. falciparum* genome; given that
isoleucine must be acquired exogenously, why has the genome not evolved to reduce its usage
altogether so as to minimize the exposure to nutrient deprivation? We believe this evolutionary
outcome confers two additional advantages. First, branched chain AAs, including isoleucine,
are potent activators of mTOR, which positively regulate haemoglobin synthesis in
erythroblasts[50,51]. Therefore, host nutrient status indirectly affects the internal nutrient supply
available for the parasite with the RBCs. By sensing the same cue that regulates haemoglobin
synthesis, the parasite can rapidly synchronize with the host nutrient status. Supporting this
hypothesis, isoleucine has been shown to be the only metabolite that synchronizes parasite
growth with the host feed-fast cycle *in vivo*[52]. Second, when the parasite was subjected to acute
AA depletion, we observed only a minor reduction in the charged Ile-tRNA level, indicating
that a continuous supply of isoleucine was maintained through recycling of endogenous
proteins. Since isoleucine usage is efficiently reallocated under nutrient stress condition, a
proteome with high isoleucine content can thus serve as a reservoir of isoleucine and enable the
parasite to endure prolonged starvation.
The high AT content of *P. falciparum* genome is maintained by a significant excess of G:C to
A:T transition, prompting the speculation that such mutational bias contributes to adaptive
evolution[53]. Coincidentally, AAs that are negatively associated with gene expression are
invariably decoded by AT-rich codons. A hypothesis suggesting that the substitution and
insertion of these AAs may be continuously selected is compelling, because these mutations
can successively increase the efficiency of resource reallocation during nutrient stress. This
evolutionary adaptation can account for the AT-biased mutational rate, at least within the
coding region of the genome. However, despite similar mode of parasitism, *P. vivax* harbors a
GC-rich genome and the emergence of similar isoleucine-sensing mechanism, although not
analyzed, is less favored. *P. vivax* shows a clear preference to invade reticulocytes, which
retains a substantial amount of host ribosomes; therefore, *P. vivax* can likely digest these
proteins to acquire isoleucine and is not a strict isoleucine auxotroph. Infections with *P. vivax*
also often result in lower parasitemia, as parasites compete for fewer nutrients in the host and
are less likely to cause severe nutrient stress. Together, these factors may weaken the selection
force that drives the adaptive evolution seen in *P. falciparum*.
A significant implication of our finding is the demonstration of a global selection sweep that
affected the primary sequences of many proteins but may not necessarily impart a functional
advantage. Although functional selection and neutral drift are believed to be the main drivers
of protein evolution[54,55], this study suggests that many non-functional missense mutations can
be under positive selection for their regulatory effect, which are otherwise manifested as neural
drift[55]. This adds new perspectives to the understanding of genome and protein evolution, such
as on the thesis of adaptive mutation bias and on the frequent occurrence of low complexity
regions in *P. falciparum* genome[2,46,54]. This finding also draws attention to the limitation of
studying translation dynamics by considering only codon usage bias, since in addition to codon-
identity, AA-identity can also be a prominent factor evolved to influence translation.
While NGD pathway has not been experimentally validated in *P. falciparum* and the genome
does not encode most of the homologues involved in the pathway[56], our findings implicate a

mechanism similar to NGD in *P. falciparum*. At present, only a highly divergent non-sense mediated decay pathway has been experimentally validated in *P. falciparum*[57]. Similarly, NGD may also be highly divergent, since the mechanism is poorly conserved outside metazoans. Further investigation on this will be needed.

Artemisinin resistance is closely related to an enhanced stress response mediated through pathways involved in proteostasis[58,59]. Furthermore, reduced haemoglobin digestion has been shown to be a resistance-associated trait[60], and parasites with this trait exhibits enhanced dependency on external nutrients to compensate for fitness loss[61,62]. Accordingly, the switch in nutrient acquisition from a balanced nutrient source to predominantly external source may influence global translation dynamics and contribute to the resistance mechanism. It warrants further study on this hypothesis.

References

Gardner, M. J. *et al.* Genome sequence of the human malaria parasite Plasmodium falciparum. *Nature* **419**, 498-511 (2002). https://doi.org:10.1038/nature01097

Singh, G. P. *et al.* Hyper-expansion of asparagines correlates with an abundance of proteins with prion-like domains in Plasmodium falciparum. *Mol Biochem Parasitol* **137**, 307-319 (2004). https://doi.org:10.1016/j.molbiopara.2004.05.016

Percudani, R., Pavesi, A. & Ottonello, S. Transfer RNA gene redundancy and translational selection in Saccharomyces cerevisiae. *J Mol Biol* **268**, 322-330 (1997). https://doi.org:10.1006/jmbi.1997.0942

Schwartz, S. & Motorin, Y. Next-generation sequencing technologies for detection of modified nucleotides in RNAs. *RNA Biol* **14**, 1124-1137 (2017). https://doi.org:10.1080/15476286.2016.1251543

Cozen, A. E. *et al.* ARM-seq: AlkB-facilitated RNA methylation sequencing reveals a complex landscape of modified tRNA fragments. *Nat Methods* **12**, 879-884 (2015). https://doi.org:10.1038/nmeth.3508

Zheng, G. *et al.* Efficient and quantitative high-throughput tRNA sequencing. *Nat Methods* **12**, 835-837 (2015). https://doi.org:10.1038/nmeth.3478

Gogakos, T. *et al.* Characterizing Expression and Processing of Precursor and Mature Human tRNAs by Hydro-tRNAseq and PAR-CLIP. *Cell Rep* **20**, 1463-1475 (2017). https://doi.org:10.1016/j.celrep.2017.07.029

Behrens, A., Rodschinka, G. & Nedialkova, D. D. High-resolution quantitative profiling of tRNA abundance and modification status in eukaryotes by mim-tRNAseq. *Mol Cell* **81**, 1802-1815.e1807 (2021). https://doi.org:10.1016/j.molcel.2021.01.028

Evans, M. E., Clark, W. C., Zheng, G. & Pan, T. Determination of tRNA aminoacylation levels by high-throughput sequencing. *Nucleic Acids Res* **45**, e133 (2017). https://doi.org:10.1093/nar/gkx514

Pinkard, O., McFarland, S., Sweet, T. & Coller, J. Quantitative tRNA-sequencing uncovers metazoan tissue-specific tRNA regulation. *Nat Commun* **11**, 4104 (2020). https://doi.org:10.1038/s41467-020-17879-x

Gingold, H. *et al.* A dual program for translation regulation in cellular proliferation and differentiation. *Cell* **158**, 1281-1292 (2014). https://doi.org:10.1016/j.cell.2014.08.011

Yu, P. *et al.* Dynamic landscapes of tRNA transcriptomes and translatomes in diverse mouse tissues. *Genomics Proteomics Bioinformatics* (2022). https://doi.org:10.1016/j.gpb.2022.07.006

529	13	Orioli, A. tRNA biology in the omics era: Stress signalling dynamics and cancer progression. *Bioessays* **39** (2017). https://doi.org:10.1002/bies.201600158
531	14	Knight, J. R. P. *et al.* Control of translation elongation in health and disease. *Dis Model Mech* **13** (2020). https://doi.org:10.1242/dmm.043208
533	15	Pang, Y. L., Abo, R., Levine, S. S. & Dedon, P. C. Diverse cell stresses induce unique patterns of tRNA up- and down-regulation: tRNA-seq for quantifying changes in tRNA copy number. *Nucleic Acids Res* **42**, e170 (2014). https://doi.org:10.1093/nar/gku945
536	16	Chan, C. T. *et al.* Highly Predictive Reprogramming of tRNA Modifications Is Linked to Selective Expression of Codon-Biased Genes. *Chem Res Toxicol* **28**, 978-988 (2015). https://doi.org:10.1021/acs.chemrestox.5b00004
539	17	Sørensen, M. A., Fehler, A. O. & Lo Svenningsen, S. Transfer RNA instability as a stress response in Escherichia coli: Rapid dynamics of the tRNA pool as a function of demand. *RNA Biol* **15**, 586-593 (2018). https://doi.org:10.1080/15476286.2017.1391440
542	18	Hammam, E. *et al.* Malaria Parasite Stress Tolerance Is Regulated by DNMT2-Mediated tRNA Cytosine Methylation. *mBio* **12**, e0255821 (2021). https://doi.org:10.1128/mBio.02558-21
545	19	McLean, K. J. & Jacobs-Lorena, M. Plasmodium falciparum Maf1 Confers Survival upon Amino Acid Starvation. *mBio* **8** (2017). https://doi.org:10.1128/mBio.02317-16
547	20	van Dam, T. J., Zwartkruis, F. J., Bos, J. L. & Snel, B. Evolution of the TOR pathway. *J Mol Evol* **73**, 209-220 (2011). https://doi.org:10.1007/s00239-011-9469-9
549	21	Sengupta, S., Peterson, T. R. & Sabatini, D. M. Regulation of the mTOR complex 1 pathway by nutrients, growth factors, and stress. *Mol Cell* **40**, 310-322 (2010). https://doi.org:10.1016/j.molcel.2010.09.026
552	22	Mancio-Silva, L. *et al.* Nutrient sensing modulates malaria parasite virulence. *Nature* **547**, 213-216 (2017). https://doi.org:10.1038/nature23009
554	23	Babbitt, S. E. *et al.* Plasmodium falciparum responds to amino acid starvation by entering into a hibernatory state. *Proc Natl Acad Sci U S A* **109**, E3278-3287 (2012). https://doi.org:10.1073/pnas.1209823109
557	24	Zhang, M. *et al.* PK4, a eukaryotic initiation factor 2α(eIF2α) kinase, is essential for the development of the erythrocytic cycle of Plasmodium. *Proc Natl Acad Sci U S A* **109**, 3956-3961 (2012). https://doi.org:10.1073/pnas.1121567109
560	25	Chan, S., Ch'ng, J. H., Wahlgren, M. & Thutkawkorapin, J. Frequent GU wobble pairings reduce translation efficiency in Plasmodium falciparum. *Sci Rep* **7**, 723 (2017). https://doi.org:10.1038/s41598-017-00801-9
563	26	Fuchs, R. T., Sun, Z., Zhuang, F. & Robb, G. B. Bias in ligation-based small RNA sequencing library construction is determined by adaptor and RNA structure. *PLoS One* **10**, e0126049 (2015). https://doi.org:10.1371/journal.pone.0126049
566	27	Marck, C. *et al.* The RNA polymerase III-dependent family of genes in hemiascomycetes: comparative RNomics, decoding strategies, transcription and evolutionary implications. *Nucleic Acids Res* **34**, 1816-1835 (2006). https://doi.org:10.1093/nar/gkl085
570	28	Bozdech, Z. *et al.* The transcriptome of the intraerythrocytic developmental cycle of Plasmodium falciparum. *PLoS Biol* **1**, E5 (2003). https://doi.org:10.1371/journal.pbio.0000005
573	29	Zaborske, J. M. *et al.* A nutrient-driven tRNA modification alters translational fidelity and genome-wide protein coding across an animal genus. *PLoS Biol* **12**, e1002015 (2014). https://doi.org:10.1371/journal.pbio.1002015

30 Ng, C. S. *et al.* tRNA epitranscriptomics and biased codon are linked to proteome
expression in. *Mol Syst Biol* **14**, e8009 (2018). https://doi.org:10.15252/msb.20178009
31 Boccaletto, P. *et al.* MODOMICS: a database of RNA modification pathways. 2017
update. *Nucleic Acids Res* **46**, D303-D307 (2018). https://doi.org:10.1093/nar/gkx1030
32 Masuda, I. *et al.* Loss of N^1-methylation of G37 in tRNA induces ribosome stalling and
reprograms gene expression. *Elife* **10** (2021). https://doi.org:10.7554/eLife.70619
33 Hou, Y. M., Masuda, I. & Gamper, H. Codon-Specific Translation by m^{1}G37 Methylation
of tRNA. *Front Genet* **9**, 713 (2018). https://doi.org:10.3389/fgene.2018.00713
34 Muramatsu, T. *et al.* Codon and amino-acid specificities of a transfer RNA are both
converted by a single post-transcriptional modification. *Nature* **336**, 179-181 (1988).
https://doi.org:10.1038/336179a0
35 Marck, C. & Grosjean, H. tRNomics: analysis of tRNA genes from 50 genomes of
Eukarya, Archaea, and Bacteria reveals anticodon-sparing strategies and domain-
specific features. *RNA* **8**, 1189-1232 (2002).
https://doi.org:10.1017/s1355838202022021
36 Sojka, D., Jalovecká, M. & Perner, J. Babesia, Theileria, Plasmodium and Hemoglobin.
Microorganisms **10** (2022). https://doi.org:10.3390/microorganisms10081651
37 Herman, J. D. *et al.* The cytoplasmic prolyl-tRNA synthetase of the malaria parasite is
a dual-stage target of febrifugine and its analogs. *Sci Transl Med* **7**, 288ra277 (2015).
https://doi.org:10.1126/scitranslmed.aaa3575
38 McLean, K. J. & Jacobs-Lorena, M. The response of Plasmodium falciparum to
isoleucine withdrawal is dependent on the stage of progression through the
intraerythrocytic cell cycle. *Malar J* **19**, 147 (2020). https://doi.org:10.1186/s12936-
020-03220-w
39 Altenhoff, A. M. *et al.* OMA orthology in 2021: website overhaul, conserved isoforms,
ancestral gene order and more. *Nucleic Acids Res* **49**, D373-D379 (2021).
https://doi.org:10.1093/nar/gkaa1007
40 Nagarajan, S. & Grewal, S. S. An investigation of nutrient-dependent mRNA translation
in Drosophila larvae. *Biol Open* **3**, 1020-1031 (2014).
https://doi.org:10.1242/bio.20149407
41 Gobet, C. *et al.* Robust landscapes of ribosome dwell times and aminoacyl-tRNAs in
response to nutrient stress in liver. *Proc Natl Acad Sci U S A* **117**, 9630-9641 (2020).
https://doi.org:10.1073/pnas.1918145117
42 Patakottu, B. R., Singh, P. K., Malhotra, P., Chauhan, V. S. & Patankar, S. In vivo analysis
of translation initiation sites in Plasmodium falciparum. *Mol Biol Rep* **39**, 2225-2232
(2012). https://doi.org:10.1007/s11033-011-0971-3
43 Ikeuchi, K., Izawa, T. & Inada, T. Recent Progress on the Molecular Mechanism of
Quality Controls Induced by Ribosome Stalling. *Front Genet* **9**, 743 (2018).
https://doi.org:10.3389/fgene.2018.00743
44 Simms, C. L., Yan, L. L. & Zaher, H. S. Ribosome Collision Is Critical for Quality Control
during No-Go Decay. *Mol Cell* **68**, 361-373.e365 (2017).
https://doi.org:10.1016/j.molcel.2017.08.019
45 Oertlin, C. *et al.* Generally applicable transcriptome-wide analysis of translation using
anota2seq. *Nucleic Acids Res* **47**, e70 (2019). https://doi.org:10.1093/nar/gkz223
46 Radó-Trilla, N. & Albà, M. Dissecting the role of low-complexity regions in the evolution
of vertebrate proteins. *BMC Evol Biol* **12**, 155 (2012). https://doi.org:10.1186/1471-
2148-12-155

623	47	Bertels, L. K., Fernández Murillo, L. & Heinisch, J. J. The Pentose Phosphate Pathway in
624		Yeasts-More Than a Poor Cousin of Glycolysis. *Biomolecules* **11** (2021).
625		https://doi.org:10.3390/biom11050725
626	48	Mans, B. J., Pienaar, R. & Latif, A. A. A review of Theileria diagnostics and epidemiology.
627		*Int J Parasitol Parasites Wildl* **4**, 104-118 (2015).
628		https://doi.org:10.1016/j.ijppaw.2014.12.006
629	49	Tiessen, A., Pérez-Rodríguez, P. & Delaye-Arredondo, L. J. Mathematical modeling and
630		comparison of protein size distribution in different plant, animal, fungal and microbial
631		species reveals a negative correlation between protein size and protein number, thus
632		providing insight into the evolution of proteomes. *BMC Res Notes* **5**, 85 (2012).
633		https://doi.org:10.1186/1756-0500-5-85
634	50	Chung, J. *et al.* The mTORC1/4E-BP pathway coordinates hemoglobin production with
635		L-leucine availability. *Sci Signal* **8**, ra34 (2015).
636		https://doi.org:10.1126/scisignal.aaa5903
637	51	Knight, Z. A., Schmidt, S. F., Birsoy, K., Tan, K. & Friedman, J. M. A critical role for
638		mTORC1 in erythropoiesis and anemia. *Elife* **3**, e01913 (2014).
639		https://doi.org:10.7554/eLife.01913
640	52	Kimberley, F. P. *et al.* Synchrony between daily rhythms of malaria parasites and hosts
641		is driven by an essential amino acid. *bioRxiv* (2021).
642		https://doi.org:10.1101/2020.08.24.264689
643	53	Hamilton, W. L. *et al.* Extreme mutation bias and high AT content in Plasmodium
644		falciparum. *Nucleic Acids Res* **45**, 1889-1901 (2017).
645		https://doi.org:10.1093/nar/gkw1259
646	54	Jayaraman, V., Toledo-Patiño, S., Noda-García, L. & Laurino, P. Mechanisms of protein
647		evolution. *Protein Sci* **31**, e4362 (2022). https://doi.org:10.1002/pro.4362
648	55	Svensson, E. I. & Berger, D. The Role of Mutation Bias in Adaptive Evolution. *Trends*
649		*Ecol Evol* **34**, 422-434 (2019). https://doi.org:10.1016/j.tree.2019.01.015
650	56	Pavlovic Djuranovic, S. *et al.* translational machinery condones polyadenosine repeats.
651		*Elife* **9** (2020). https://doi.org:10.7554/eLife.57799
652	57	McHugh , E., Bulloch , M. S., Batinovic , S., Sarna , D. K. & Ralph, S. A. A divergent
653		nonsense-mediated decay machinery in Plasmodium falciparum is inefficient and non-
654		essential. *BioRxiv* (2021). https://doi.org:10.1101/2021.04.14.439394
655	58	Zhang, M. *et al.* Inhibiting the Plasmodium eIF2α Kinase PK4 Prevents Artemisinin-
656		Induced Latency. *Cell Host Microbe* **22**, 766-776.e764 (2017).
657		https://doi.org:10.1016/j.chom.2017.11.005
658	59	Mok, S. *et al.* Drug resistance. Population transcriptomics of human malaria parasites
659		reveals the mechanism of artemisinin resistance. *Science* **347**, 431-435 (2015).
660		https://doi.org:10.1126/science.1260403
661	60	Birnbaum, J. *et al.* A Kelch13-defined endocytosis pathway mediates artemisinin
662		resistance in malaria parasites. *Science* **367**, 51-59 (2020).
663		https://doi.org:10.1126/science.aax4735
664	61	Bunditvorapoom, D. *et al.* Fitness Loss under Amino Acid Starvation in Artemisinin-
665		Resistant Plasmodium falciparum Isolates from Cambodia. *Sci Rep* **8**, 12622 (2018).
666		https://doi.org:10.1038/s41598-018-30593-5
667	62	Mesén-Ramírez, P. *et al.* The parasitophorous vacuole nutrient channel is critical for
668		drug access in malaria parasites and modulates the artemisinin resistance fitness cost.

669   *Cell   Host   Microbe*   **29**,   1774-1787.e1779   (2021).
670   https://doi.org/10.1016/j.chom.2021.11.002

671
672

Acknowledgement

We thank all members of the Larsson laboratory and the Ribacke laboratory for their constructive comments and feedback. We thank Pan Tao and Addgene for providing us the demethylase plasmids and E. Strandback, H.A. Korsah and T. Nyman in the Protein Science facility of Karolinska Institutet for assistance in synthesizing the recombinant proteins. Protein identification and quantification were carried out by the Proteomics Biomedicum core facility, Karolinska Institutet (https://ki.se/en/mbb/proteomics-biomedicum). The authors acknowledge the support of the Freiburg Galaxy Team: Person *X* and Björn Grüning, Bioinformatics, University of Freiburg (Germany) funded by the Collaborative Research Centre 992 Medical Epigenetics (DFG grant SFB 992/1 2012) and the German Federal Ministry of Education and Research BMBF grant 031 A538A de.NBI-RBC. This work was supported by a grant from Knut and Alice Wallenberg Foundation (2017.0055.) to M.W. and U.R, a grant from the Swedish Research Council (2021-03141) to S.CL.C. and M.W, a grant from Tore Nilsons Stiftelse För Medicinsk Forskning (2017-00532) to S.CL.C., S.CL.C. was also supported by a postdoctoral fellowship grant from Svenska Sällskapet för Medicinsk Forskning, Q.L. was supported by a scholarship from Chinese Scholarship Council and a doctoral (KID #2018-01037) grant from Karolinska Institutet.

Figure 1. Maintenance of an imbalanced codon and anticodon pools throughout the IDC.
a, The dynamics of the anticodon pool during the IDC. The fraction of each tRNA to the total aligned tRNA reads in each stage is shown. **b,** The dynamics of tRNA charging throughout the IDC. Charged tRNAs are represented by reads with 3' -CCA ends after periodate oxidation and beta-elimination reactions. **c,** The dynamics of the codon pools throughout the IDC. The codon usage frequency of the transcriptome in each stage was estimated by the codon usage of each expressed gene weighted by the expression level (transcript per million), and the fraction of codons that needed to be decoded by each tRNA is shown. **d,** Heatmap showing log$_2$-transformed averaged ratio of supply to demand (anti-codon/codon) for all tRNAs across the IDC. R: Ring; T: Trophozoite; S: Schizont. Error bars represent the standard deviation (SD) of biological replicates. For a & c, N=3 each for the Ring, Trophozoite and Schizont stages. For b, Ring, Trophozoite: N=3, Schizont: N=2. All apicoplast tRNAs are excluded.

Figure 2. tRNA usage delineates gene expression level and haemoglobin dependency.
a, Principal component analysis of tRNA usage among all the coding genes in the genome. The most variably used tRNAs among genes (cyan) were named and used for subsequent analyses. **b,** Plot showing the relationship of the usage of each tRNA in the codon pool (x-axis) and the $\log_2$-transformed ratio of supply to demand per tRNA (y-axis) in the trophozoite stage. Variable tRNAs (cyan) represent tRNAs named in (a). The line demarcates regions of underrepresentation (below the line) and overrepresentation (above the line). Pie charts show the combined usage of these tRNAs in the codon pool. **c,** tRNA usage in different gene categories. Genes were sorted by the averaged TPM values in the trophozoite stage and classified into three categories; Ribosomal: genes encoding ribosomal proteins (n=84), Top 5% genes: the 5% of the most highly expressed coding genes (n=258), Bottom 5% genes: the 5% of the least expressed coding genes (n=258). P-values of Mann-Whitney U-test between the top and bottom 5% genes are denoted by **** (p<0.0001); ** (p<0.01); ns (not significant, p>0.05). Glu-UUC shows an opposite trend with higher mean usage in the top 5% genes (red asterisk). **d,** Correlation of the gene expression levels in the trophozoite stage and the combined usage of Asn-GUU, Ile-AAU, Ile-UAU, Phe-GAA and Tyr-GUA tRNAs of genes. **e,** Correlation of the gene expression levels in the trophozoite stage and isoleucine usage of genes. **f,** Plot showing the relationship of AA usage frequency in the codon pool (x-axis) and in host haemoglobin (y-axis). AAs are defined as negatively associated with expression level if correlation between usage frequency in genes and gene expression level was < -0.2, not associated if -0.2 < ρ < 0.2, positively associated if ρ > 0.2. **g,** Correlation between the AA usage frequency in different gene categories and in host haemoglobin. **h,** The relationship

746   between the abundance of transcripts (x-axis) and the correlation between the AA usage
747   frequency of the transcripts and the AA composition of haemoglobin (y-axis). For all panels,
748   ρ: Spearman's rank correlation coefficients. Coding genes, n=5267.
749
750

751
Figure 3. Isoleucine availability is sensed as a proxy of nutrient status. a, Changes in
charged tRNA levels after 6 hours of treatment in late trophozoite cultures (CM: complete
medium, AA-: AA depletion, HF: halofuginone (70 nM) and AA depletion). All changes are
represented relative to the CM condition. Error bars represent the SD of N=3 biological
replicates. **b,** Box plot showing isoleucine (left) and leucine (right) usage in up- and
downregulated genes during stage progression. P-values of Mann-Whitney U-test are denoted
by **** (p<0.0001); ns (not significant, p>0.05). **c,** The multiplication of parasites after 6
hours of AA depletion in the presence of varying concentrations of isoleucine. The parasites
were starved from 34±2 hpi (red) or 8±2 hpi (cyan) for 6 hours and subsequently recovered by
culturing in complete medium until all parasites had completed reinvasion. The multiplication
rate was normalized to the complete medium (CM) group. Error bars represent the SD of N=3
biological replicates. P-values of the two-tailed unpaired t test are denoted by ****
(p<0.0001); ns (not significant, p>0.05). **d,** (Left) Volcano plot showing 1146 proteins
identified in all samples by LC-MS/MS after 6 hours of AA depletion. The dotted line
indicates the cut-off of the FDR-adjusted p-value=0.05. 56 of the 78 downregulated proteins
are associated with merozoite, and their reported localizations are indicated (right). **e,** Gene
Ontology (GO) analysis of the top 5% isoleucine-rich genes (n=265). **f,** The number of SDs
for the usage frequency in *P. falciparum* orthologue from the mean usage frequency of all
orthologous proteins for isoleucine (left) and leucine (right). High isoleucine: top 5% of
isoleucine-rich genes, high leucine: top 5% of leucine-rich genes, ribosomal protein:
ribosomal protein encoding genes. Only genes with >100 orthologues were considered as
functionally conserved and analyzed. Bracketed values = average orthologue counts. Red
lines indicate 0 and 3 SD.

775
Figure 4. Polysome profiling reveals targeted reprogramming after AA starvation.
a, (Below) Representative polysome traces of parasites growing in complete medium (CM) and
AA-depleted medium (AA-) on a 10-50% sucrose gradient. Fractions containing monosomes
and polysomes are indicated. (Upper) Immunoblotting of each fraction against RPS14. **b,**
Differential expression analysis of the genes in the polysome fractions after AA starvation. The
horizontal and vertical dotted lines show the cut-off of the FDR-adjusted p-value=0.05 and cut-
off of twofold changes, respectively. **c,** The differences in isoleucine and leucine usages
between genes with increased or decreased polysome association **d,** The Spearman correlation
of the changes in transcript abundance in the polysome fractions and the changes in protein
abundance. **e,** Kozak context of genes with decreased (DPA) or increased polysome association
(IPA) **f, g,** The differences in **(f)** isoleucine and leucine usage; and **(g)** the Kozak context
between genes up- and downregulated within (left) the stage independent and (right) the stage
dependent transcriptome changes. Kozak context - strong: Rnn<u>ATG</u>R; adequate:
Rnn<u>ATG</u>Y/Ynn<u>ATG</u>R; and weak: Ynn<u>ATG</u>Y (<u>ATG</u>= start codon. Mann-Whitney U test for
the AA usage analysis and Fisher's exact test for the Kozak context analysis, * (p<0.05); ns
(not significant, p>0.05). **h,** The Spearman correlation of the changes in transcript abundance
in the polysome fractions and the total transcriptomes. Only transcripts with an FDR<0.05 and
|log₂ fold-change| > 1 in polysome association are shown. **i,** Proposed model of a decentralized
nutrient-sensing mechanism. (Left) Nutrient deprivation due to host fasting reduces
haemoglobin synthesis in erythroblasts via a dampened mTOR activation by branched chain
AAs. Reduction in serum Ile levels below a threshold level causes an accumulation of
uncharged Ile-tRNAs and the selective decrease in translational output of the translatome. Lipid
metabolism genes are highly targeted due to generally high Ile usage, leading to lower
merozoite production and potentially reduced endocytosis of host haemoglobins. (Right) The
changes in ribosome loadings also reprogramme the transcriptome. Differential stabilization of
transcripts depends on the interplay between translation initiation and Ile-codon-induced

stalling. Reprogramming in conjunction with a global reduction in translation initiation by
eIF2α phosphorylation drives the cell into a "hibernation state" signified by reduced proteome
turnover but a transcriptome primed for rapid metabolic reactivation upon the cessation of
nutrient deprivation.
806

807
808
Figure 5. Reduced dependency of housekeeping genes on external nutrient sources is
conserved. Spearman correlation (Rho) between the AA compositions of the host haemoglobin
and that of the enzymes involved in glycolysis or the pentose phosphate pathway (PPP) in
different organisms. Correlations were calculated using *Homo sapiens* haemoglobin for *P.*
falciparum, *P. vivax*, *B. microti*, *T. brucei*, *T. gondii*. *H. sapiens* and *S. cerevisiae*; using *Bos*
Taurus haemoglobin for *B. bovis*, *T. parva*, *T.annulata* and *T. orientalis*; and using *Mus*
musculus haemoglobin for *P. berghei*. Non-parasitic multicellular (*H. sapiens*) and unicellular
(*S. cerevisiae*) organisms are shown for comparison. The red box represents organisms with
developmental stage within host RBCs. P-values of Mann-Whitney U-test are denoted by **
(p<0.01); * (p<0.05); ns (not significant, p>0.05).
819
820

Material Extension for

Genomic adaptation underlies nutrient-driven reprogramming in the malaria-causing parasite

Qian Li[1,4], Leonie Vetter[1], Cagla Sahin[1], Akos Vegvari[2], Ulf Ribacke[1], Mats Wahlgren[1], Ola Larsson[3], Sherwin Chun-Leung Chan[3,4*]

This file includes:
Materials and Methods
Extended Data Fig. 1 to 14
References

Materials and methods

Parasite Culturing

Plasmodium falciparum strain NF54 were cultured with human O+ erythrocytes in RPMI 1640 medium (Thermofisher Scientific, #42402-016) supplemented with 10% A^+ human serum, 50 µg/L gentamicin (Gibco) and 2mM L-glutamine (Hyclone). Cultures were maintained at 37°C in 5% O_2, 5% CO_2 gas mixture. Parasites were routinely synchronized by treating with 5% sorbitol solution (Sigma) for ring stage and by using percoll gradient (Cytiva) for late stage.

RNA isolation

For time series experiment, cultures were expanded to 30 ml at 3% hematocrit and ~8% parasitemia, 10 ml culture were collected for RNA extraction at each time point (8-12, 24-28 and 40-44 hpi).

For tRNA library

Small RNA (<200 nt) was isolated directly from the pellets of infected RBCs using the *mir*Vana miRNA isolation kit (Invitrogen) according to the manufacturer's instructions. 600 µl lysis buffer was added to 150 µl infected RBCs and the cell suspension was vortexed vigorously to completely lyse the cells. 1/10 volume of miRNA Homogenate Additive was added to the cell lysates and the samples were incubated 10min on ice. After addition of one v/v ratio of Acid-Phenol: Chloroform to the lysate, the samples were vortexed vigorously and centrifuged at 10,000 g for 5 min and the aqueous phase was transferred to a new tube. 1/3 volume of 100% ethanol was added to the aqueous phase and subsequently passed through the column. The flow through was mixed with 2/3 volume of 100% ethanol and passed through a second filter. Small RNAs were immobilized on the filter. Two consecutive washing were performed using miRNA wash solution 1 and wash solution 2/3, respectively. Small RNAs were eluted in 70µl pre-heated (95°C) nuclease-free water.

For mRNA library

Total RNA was extracted from 100 µl infected RBCs using Trizol reagent (Sigma). 1 ml Trizol was added to the cell pellet and pipetted up and down until complete homogenization of the lysate. The protocol was then carried out according to the manufacturer's instructions. Total RNAs and small RNAs were quantified by Qubit RNA high sensitivity kit (Invitrogen).

tRNA library construction

Periodate oxidation and β-elimination

RNA oxidation and β-elimination were performed as previously described[1]. 2 µg isolated small RNA was treated in 100 mM NaOAc/HOAc, pH 4.8 and freshly prepared 50 mM $NaIO_4$ at room temperature for 30 min. The reaction was quenched by incubating with 100 mM glucose for 5 min at room temperature. The treated RNA was purified first with RNA clean-up and concentration micro-elute kit (Norgen Biotek Corp.), and then by ethanol precipitation. For β-elimination, the purified RNA was treated in 60 mM sodium borate, pH 9.5 at 45°C for 90 min. Subsequently, the RNA was purified with RNA clean-up and concentration micro-elute kit followed by ethanol precipitation. Two short RNA standards with A-end and C-end were mixed in equimolar ratio and added to the purified RNA to a final concentration of 0.4 pmol/µg small RNA).

Two synthetic RNA standards:

A-ending: 5'-GCAGAUGGCUUCAAUUGCUAUUAAGGACCA

C-ending: 5'-GCAGAUGGCUUCAAUUGCUAUUAAGGACAC

96   *Bisulfite conversion*
97   Bisulfite conversion of RNA was performed to generate separate sequencing library. 1 μg of
98   isolated small RNA was used for the reaction using EZ RNA methylation kit (Zymo Research).
99
tRNA library generation
This was carried out as previously described with minor modifications[2,3]. For cDNA synthesis,
optimized R2R DNA (1:1 ratio of T-ending and G-ending oligos) was annealed to
complementary R2 RNA in 100 mM Tris-HCl, 0.5 mM EDTA, pH 7.5 at 82°C for 2 min, then
cooled down to 25°C with a ramp rate of 0.1°C /s[4]. A total of 100 ng tRNA (≈4 pmol) was
mixed with 4 pmol primer mixture and pre-incubated at room temperature for 30 min with low
salt buffer (50 mM Tris-HCl, pH7 .5, 75 mM KCl, 3 mM $MgCl_2$), 10 mM DDT and 500 nM
TGIRT-III enzyme (InGex, Inc). dNTPs mix were added to a final concentration of 1 mM to
initiate the reaction. The template switching reaction was performed at 42°C for 16 h.
The reactions were terminated by adding 1 μl of 5 M NaOH and incubated at 95°C for 3
min.The reactions were neutralized by addition of 1μl of 5M HCl. cDNAs were then purified
by MinElute Reaction Cleanup Kit (QIAGEN) and eluted in 10 μl nuclease free water. 10 μM
R1R DNA containing UMI sequences was adenylated with 2 μl of Mth RNA ligase (New
England Biolab) at 65°C for 1 h and subsequently inactivated at 85°C for 5 min. 2 μM of the
adenylated R1R DNA was ligated to the purified cDNAs at 65°C for 2 h with Thermostable 5'
AppDNA/RNA ligase (New England Biolab). After inactivation at 90°C for 3 min, the ligated
cDNAs were purified with MinElute Reaction Cleanup Kit (QIAGEN) and eluted in 23 μl
nuclease free water. PCR amplification for Illumina sequencing was performed using 2X
Phusion High-Fidelity PCR Master Mix (Thermofisher Scientific) for 12 PCR cycles (98°C 5s,
60°C 10s, 72°C 10s) using R1 multiplex PCR primer and barcoded primer. 1.3X Agencourt
AMPure XP beads (Beckman Coulter) were used to clean up the libraries and get rid of primer
dimers before Illumina sequencing. The quality and quantity of each library was analyzed on
the Bioanalyzer with a High Sensitivity DNA analysis Kit (Agilent technologies) and Collibri
Library Quantification Kit (Invitrogen).
R2 RNA:
5'-GAUCGGAAGAGCACACGUCUGAACUCCAGUCAC-3SpC3/
R2R DNA T-ending/ G-ending:
5'-GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT/G
R1R DNA:
5'-/5Phos/GATANNNNNNNGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT/3SpC3/
R1 multiplex PCR primer:
5'-AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCC
GATCT
Barcoded primer:
5'CAAGCAGAAGACGGCATACGAGAT*[BARCODE]*GTGACTGGAGTTCAGACGTGT
GCTCTTCCGATCT
136
137   **mRNA library construction**
138   100 ng total RNA was used to construct mRNA library using TruSeq Stranded mRNA kit
139   (Illumina). The protocol was carried out following manufacturer's instruction. The quality and
140   quantity of each library was analyzed on the Bioanalyzer with a High Sensitivity DNA analysis
141   Kit (Agilent technologies) and Collibri Library Quantification Kit (Invitrogen).
142
Recombinant synthesis of AlkB demethylases

pET30a-AlkB and pET30a-AlkB-D135S were gifts from Tao Pan (Addgene plasmid # 79050
and #79051). These plasmids contain wildtype *E.coli alkB* or D135S mutant sequences, both
with a deletion of 11 residues from the N-terminal[5]. Plasmids were transformed into *E.coli*
BL21(DE3) T1R pRARE2 and the cultures were grown in the presence of 50 μg/ml kanamycin
and 34 μg/ml chloramphenicol at 37°C until OD_{600} reached 2. Protein expression was then
induced with 0.5 mM IPTG at 18°C overnight. Cells were pelleted and resuspended in lysis
buffer (100 mM HEPES, 500 mM NaCl, 10% glycerol, 10 mM imidazole, 0.5 mM TCEP,
protease inhibitor cocktail (Roche), pH 8.0). The Lysate was sonicated by pulsed sonication
(4s/4s on/off cycle for 4 min, 80% amplitude) and then centrifuged and filtered through 0.45
μm filters. The soluble proteins were purified first using HisTrap HP (GE Healthcare) and then
by gel filtration on a HiLoad 16/60 Superdex 200 column (GE Healthcare) on an ÄKTA Xpress
system. Thrombin was added (1 unit/mg protein) to the purified recombinant proteins and
incubated overnight at 4°C. The cleavage tag was then moved by HisTrap HP. The proteins
were then concentrated in storage buffer (20 mM HEPES, 300 mM NaCl, 10% glycerol, 2 mM
TCEP, pH 7.5).
159
tRNA demethylation reaction
Demethylation assay was performed to validation the identities of the detected mutation
signatures. 1 μg small RNA was treated with 80 pmol WT and 160 pmol D135S AlkB in
demethylase reaction buffer (300 mM KCl, 50 mM MES buffer (pH 5.0), 2 mM $MgCl_2$, 50 μM
of $(NH_4)_2Fe(SO_4)_2·6H_2O$, 300 μM 2-ketoglutarate (2-KG), 2 mM L-ascorbic acid, 50 μg/ml
BSA). The reaction was incubated at room temperature for 2h and then quenched by adding 5
mM EDTA. The treated RNA was purified using RNA clean-up and concentration micro-elute
kit.
168
Library sequencing and analysis
tRNA library read preprocessing
All libraries were sequenced on an Illumina Nextseq 550 platform with a single-end read length
of 120 bp. Base calling and demultiplexing were performed with Generate FASTQ (v2.0.1).
Unique molecular identifiers (UMI) were extracted using UMI_tools (v1.1.2). Adapter
sequences were clipped from the reads using fastx_toolkit (v0.0.14) with a size filtering to
discard reads <15bp. These analysis was performed on the galaxy platform.
For our system, we expect that the low complexity tRNA transcriptome of the parasite would
generate the majority of reads sequenced in the libraries, with the few remaining reads
originated from the highly complex human tRNA transcriptome as well as from other non-
tRNA small RNA species. Therefore, a combined tRNA reference (including Plasmodium
nuclear and apicoplast tRNAs and all human tRNAs from GtRNAdb[6]) was first compiled and
curated using sequence data generated from a pilot sequencing trial. The curation includes re-
defining the exact 5' and 3' position of all tRNAs, addition of 3'-CCA to all tRNAs, changing
the reference from A to G for tRNAs that has inosine modification on position 34 and the
addition of a 5' G base on His-tRNA[7].
The sequence reads were then aligned to the curated tRNA reference using bowtie2 (v2.2.1)
with sensitive local mode[8]. The unaligned reads were then aligned to the 3D7 genome (v.
PlasmoDB-46) with sensitive local mode to identify mapping to other small RNA species. Two
addition steps were performed to assess the outcome of the mapping. First, the unaligned reads
from the genome mapping were again aligned to the tRNA reference with very sensitive local
mode, which should yield very few reads. Second, unaligned reads with 3'-CCA were
randomly assessed by blasting to 3D7 genome to ensure most reads with 3'-CCA were mapped
in prior operations. After alignment, reads with the same UMI and mapping coordinates were
de-duplicated using UMI_tools (v1.1.2).

tRNA composition was determined by the ratio of read counts per unique tRNA to the read
counts of total tRNAs found by htseq-count (v0.9.1) [9].
The determination for tRNA aminoacylation level is as follows: an aligned read is considered
CCA-ending(charged) if it aligns with no mismatches to the tRNA's 3' end. Consequently, if
the read aligns to the 3'end but only ends in -CC, it is considered uncharged. Aminoacylation
level were determined as the percentage of CCA-ending reads over the sum of the CCA-ending
and CC-ending reads, adjusted to the recovered ratio of the two spike-in RNA oligos.
Fractions of mutation signature were calculated as 1- (coverage of reference sequence at
position n/ total coverage at position n+1).
203
mRNA library read preprocessing
mRNA libraries were sequenced on an Illumina Nextseq 550 platform with a single-end read
length of 120 bp or 150bp (for the polysome profiles). Base calling was performed with
Generate FASTQ (v2.0.1). Sequence reads were first filtered with Trimmomatic (v0.38) using
the default setting[10]. The trimmed reads were then aligned to 3D7 genome (v. PlasmoDB-46)
with HISAT2 (v2.2.1) using default setting and read counts were obtained by htseq-count
(v0.9.1)[11].
211
Amino acid starvation assay
Parasite cultures were synchronized with two consecutive cycles of sorbitol and percoll
gradient treatment before the start of the assay. All starvation assays used parasites at 32-36
hpi maintained at 3% hematocrit and >5% parasitemia. The parasites were washed twice with
and resuspended in either AA-free RPMI (US biological, #R8999-04A) or complete RPMI (US
biological, #R8999-13) supplemented with 0.5% AlbumaxII and 20μg/ml hypoxanthine, pH
7.2. Parasites were incubated for 6 h and harvested immediately afterwards for the preparation
of tRNA library and quantitative proteomics analysis. 70 nM of halofuginone (Calbiochem) or
382 nM of L-isoleucine (Sigma) were also added where applicable.
221
Quantitative proteomics analysis
Sample preparation
Cell pellets were solubilized with 100 µL of 8M urea, 0.2% ProteaseMAX (Promega) and 5%
acetonitrile (ACN) in 100 mM Tris-HCl, pH 8.0 with 1 µL of 100x protease inhibitor cocktail
(Roche) during sonication in water bath for 10 min followed by probe sonicated with VibraCell
probe (Sonics & Materials, Inc.) for 20 s, with pulse 2/2, at 20% amplitude. Following
centrifugation at 10,000 g for 10 min at 4°C, the supernatant was transferred to a new tube and
the remaining pellet was supplemented with 50 µL of 1M urea in 100 mM Tris-HCl with 0.2%
ProteaseMAX and sonicated in water bath for 10 min. Protein concentration in the combine
supernatant was measured by BCA assay (Thermofisher Scientific). An aliquot of 25 µg
samples was transferred to a new tube and equalized with 50 mM Tris-HCl to a total volume
of 100 µL. Proteins were reduced with addition of 1 µL of 500 mM dithiothreitol (Sigma) at
25°C for 1 h and alkylated with 3 µL of 500 mM iodoacetamide for 1 h at room temperature in
the dark and quenched with 2 yL of dithiothreitol for 10 min. Proteolytic digestion was started
with 2 µL of 0.5 µg/µL Lys-C (Wako, Japan) at 37°C for 3 h with shake at 450 rmp and
supplemented with 300 µL of 100 mM Tris-HCl, pH 8.0 before adding 4 µL of 0.5 µg/µL
sequencing grade trypsin (Promega) and incubtaed over night at 37°C. The digestion was
stopped with 20 µL cc. formic acid, incubating the solutions at RT for 5 min. The sample was
cleaned on a C18 Hypersep plate with 40 µL bed volume (Thermofisher Scientific), dried using
a vacuum concentrator (Eppendorf). Biological samples were labeled with TMT-10plex
reagents in random order adding 200 µg TMT-reagent in 30 µL dry ACN to each digested
sample resolubilized in 70 µL of 50 mM triethylammonium bicarbonate (TEAB) and

incubating at room temperature (RT) for 2 h. The labeling reaction was stopped by adding 11
µL of 5% hydroxylamine and incubated at RT for 15 min before combining them in one vial.
246
Liquid Chromatography-Tandem Mass Spectrometry Data Acquisition
Peptides were reconstituted in solvent A and approximately, two µg samples injected on a 50
cm long EASY-Spray C18 column (Thermofisher Scientific) connected to an Ultimate 3000
nanoUPLC system (Thermofisher Scientific) using a 120 min long gradient: 4-26% of solvent
B (98% acetonitrile, 0.1% FA) in 120 min, 26-95% in 5 min, and 95% of solvent B for 5 min
at a flow rate of 300 nL/min. Mass spectra were acquired on a Q Exactive HF hybrid quadrupole
orbitrap mass spectrometer (Thermofisher Scientific) ranging from m/z 375 to 1700 at a
resolution of R=120,000 (at m/z 200) targeting 5×10^6 ions for maximum injection time of 80
ms, followed by data-dependent higher-energy collisional dissociation (HCD) fragmentations
of precursor ions with a charge state 2+ to 8+, using 45 s dynamic exclusion. The tandem mass
spectra of the top 18 precursor ions were acquired with a resolution of R=60,000, targeting
2×10^5 ions for maximum injection time of 54 ms, setting quadrupole isolation width to 1.4 Th
and normalized collision energy to 33%.
260
Data analysis
Acquired raw data files were analyzed using Proteome Discoverer v2.5 (Thermofisher
Scientific) with Sequest HT search engine against Plasmodium falciparum PF3D7 protein
database (SwissProt). A maximum of two missed cleavage sites were allowed for full tryptic
digestion, while setting the precursor and the fragment ion mass tolerance to 10 ppm and 0.02,
respectively. Carbamidomethylation of cysteine was specified as a fixed modification, while
TMT6plex on lysine and N-termini, oxidation on methionine as well as deamidation of
asparagine and glutamine were set as dynamic modifications. Initial search results were filtered
with 5% FDR using Percolator node in Proteome Discoverer. Quantification was based on the
TMT-reporter ion intensities. Differential quantitation analysis was performed by DESeq2[12].
271
Multiplication rate assay
Parasite cultures were synchronized with two consecutive cycles of sorbitol and percoll
gradient treatment before the start of the assay. Parasites at 8-12 hpi or 32-36 hpi were collected
for the assay. The parasites were washed twice in AA-free RPMI and then resuspended at 2%
hematocrit and 1% parasitemia in different conditions including in complete medium, AA-free
medium and AA-free medium supplied with varying concentrations of isoleucine (382 µM, 40
µM, 20 µM, 10 µM, 5 µM, 2.5 µM) in a 96-well plates. After 6h of incubation, the starved
cultures were re-supplemented with complete medium until all parasites had entered a new
invasion cycle. Microscopy assessment was used to ensure all parasites were either in ring or
trophozoite stage. Samples were then harvested for flow cytometry analysis.
282
Flow cytometry analysis
Samples were harvested for flow cytometry analysis at 0h, 6h, 24h and 48h after plate setup
for 6-10 hpi parasites and at 0h, 6h, 12h, 18h and 32h for 32-36hpi parasites. 20µl of sample
per timepoint were harvested and stained for 30 min at 37°C with Hoechst (Invitrogen) and
DHE (Sigma) at a final concentration of 20 µg/ml and 10 µg/ml respectively. Stained cells were
diluted in PBS to a final hematocrit of 0.2% prior to analysis. At least 3000 Hoechst+/DHE+
events/sample were acquired using a FACSVerse (BD Biosciences). Data was analyzed as
described previously using FlowJo (Version 10.6.2)[13].
291
Polysome fractionation

Polysome purification was adopted from a protocol previously reported but with major
modifications[14].
Cultures were expanded to 150 ml at 3% hematocrit and ~10% parasitemia and were
synchronized with two consecutive cycles of sorbitol and percoll gradient treatment before the
start of the assay. When the culture reached 32-36 hpi, infected RBCs were enriched with
magnetic separation using XS column (Miltenyi Biotec). After the enrichment, the cell pellet
was equally split and cultured either in AA-free RPMI or complete RPMI for 6h.
After the incubation, cycloheximide was added to a final concentration of 100 µg/ml to the
cultures. The cultures were briefly shaken and immediately incubated on ice for 10 min.
Afterwards, the cells were centrifuged and washed once with 10 ml ice-cold PBS supplemented
with 100 µg/ml cycloheximide. Equal volume of ice-cold 2x lysis buffer (800 mM potassium
acetate, 50 mM potassium Hepes, pH 7.2, 30 mM magnesium acetate, 10 µg/ml cycloheximide,
2 mM DTT, 2 mM AEBSF, 40 U/ml rRNasin (Promega), 1% sodium deoxycholate, 2% Igepal
CA-630) was added to the pellet and gently resuspended the pellet. The resuspended pellet was
incubated on ice for 5 min with a brief vortex every minute. The pellet was then centrifuged at
17000 g for 2 min to clear the lysate. 50 µl of the lysate was separated for total RNA extraction.
A 10% to 50% sucrose gradient solution in gradient buffer (20 mM HEPES pH 7.6, 100 mM
KCl, 5 mM $MgCl_2$) was prepared on a Gradient Station platform (Biocomp Instruments). 450
µl of the lysate was carefully layered on top of the sucrose gradient and centrifuged at 35000
rpm for 2h on a pre-chillled SW40 Ti rotor with the OptimaXL-100K ultracentrifuge system
(Beckman Coulter). After the centrifugation, the gradient was fractionated into 25 fractions on
a Gradient Station platform (Biocomp Instruments). RNA and proteins were extracted from the
fractions using Trizol reagent (Sigma) as per manufacturer's instruction and the RNA from the
polysomal fractions were pooled.
317
Immunoblotting
Western blot was performed with proteins purified from the polysome fractionation. Purified
proteins were solubilized in Laemmli buffer (Bio-Rad) with 2% β-mercaptoethanol and heated
at 95°C for 10 min. The proteins were resolved in a criterion TGX Stain-Free gel (Bio-Rad)
with Tris/Glycine/SDS running buffer (Bio-Rad). The resolved proteins were transferred to
nitrocellulose membrane in Tris/glycine buffer (Bio-Rad). The membrane was blocked with
5% skim milk, 2% BSA in PBS at 4°C overnight. The membrane was incubated with 1:2000
anti-RPS14 antibody (abcam, #ab174661) in 2% skim milk, 2% BSA in PBS for 2h at room
temperature. The membrane was washed thrice with 1x TBS, 0.1% Tween 20 and then
incubated for 30 min with 1:5000 HRP-conjugated anti-rabbit IgG (Amersham, #NA934).
Chemiluminescent signal was detected with a ChemiDoc imager (Bio-Rad) after developing
with WesternBright ECL kit (advansta).
330
331 **Orthologue analysis**
332 All orthologues to the isoleucine-rich genes of *P. falciparum* were obtained from OMA
333 orthology database using the OMA group fingerprint[15]. A cut-off of having at least 100
334 orthologues was applied to consider a *P. falciparum* protein as functionally conserved and used
335 for anaylsis.
336
Kozak Similarity score (KSS)
Calculation of KSS was inspired by a previous report[15]. Bit scores for each nucleotide from
position -3 to +4 of the annotation start codon was generated from all genes encoding ribosomal
protein and used as a reference. The KSS of a gene was calculated as the sum of all the bit
scores from these positions relative to the maximum possible score summed from these
positions.

Statistical analysis

Two tailed student's t-test, two tailed Mann-Whitney U test and Fisher's exact test were used for statistical analyses wherever specified. A p-value of <0.05 was used as the cut off for statistical significance. Multiple testings were corrected with Benjamini-Hochberg procedure, with a false discovery rate <0.05 used as the cut off for statistical significance.

393 Extended Data Fig. 1
394 **The codon usage bias in *P. falciparum* is not compensated by an expansion of tRNA genes.**
395 Correlations between tRNA gene copy numbers and codon usage in the genomes of *T. gondii*
396 and *P. falciparum*
397
398

400
Extended Data Fig. 2
An optimized workflow showing full-length cDNA library construction and data analysis
pipeline.
(a) Left panel: Schematic showing tRNA sequence library generation using TGIRT. A
DNA/RNA duplex with a T overhang that is complementary to the terminal adenosine of
mature tRNAs was used for the template-switching reaction. The 3' adapter was attached only
after the cDNA was synthesized. Right panel: Strategy for read processing of the tRNA
sequence libraries. **(b)** Bowtie2 alignment statistics for tRNA-sequencing data generated from
non-periodate (untreated, N=3) and periodate-treated (N=3) libraries. **(c)** Representative scaled
5'-to-3' sequence coverage plots spanning all nuclear tRNAs. The full-length tRNA was
divided into 25 bins with each bin representing 4% of the tRNA length, as shown on the x-axis.
The Y-axis value for each tRNA was normalized to the bin with the maximum coverage for
each respective tRNA. Left panel: samples without periodate treatment. Right panel: sample
treated with periodate.
415
416

417
Extended Data Fig. 3
Characteristics of the anticodon and codon pools.
(a) Origin of all reads mapped to tRNAs throughout the IDC. Error bars represent the standard
deviation of N=3 biological replicates. **(b)** Motif logo of the A-box in all *P. falciparum* tRNAs,
except Ser-tRNAs (upper), and of all Ser-tRNAs (lower). With a G-to-Y substitution at the
third position, the A-box of Ser-tRNAs diverges from the eukaryotic consensus (boxed). **(c)**
Principal component analysis of the coding transcriptomes throughout the IDC. **(d)** Coefficient
of variation of each anticodon (tRNA) and the respective codons it decodes. The abundance of
each anticodon and its decoded codons as a percentage of the total anticodon (size) and codon
pool (colour) are illustrated. Calculated from N=9, N=3 for the ring, the trophozoite and the
schizont stages. All apicoplast tRNAs are excluded.
429
430
431

Extended Data Fig. 4
Charge level of total tRNA throughout the IDC.
Charged ratio of tRNAs are represented by reads with 3'-CCA/-CC ends after periodate
oxidation and beta-elimination reactions. Error bars represent the standard deviation of
biological replicates; Ring and Trophozoite: N=3; Schizont: N=2. All apicoplast tRNAs are
excluded.
438
439
440

441
Extended Data Fig. 5
The mutation frequency measured by TGIRT-tRNAseq reflects modification
stoichiometry. Global heatmap of the average proportion of reads with **(a)** mutation signatures
and **(b)** m^{5}C methylation detected per position in all tRNAs at the ring stage. Numbers
represent the canonical position, e denotes positions in the variable loop, inferred identities of
the modifications that cause the mutations are shown (A$_9$, G$_9$, G$_{26}$, C$_{32}$, A$_{34}$, G$_{37}$ and A$_{58}$), and
identities of A$_{59}$ and U$_{73}$ cannot be inferred. **(c)** Schematic representation of the locations of
the modifications in relation to the secondary structure of tRNA. Blue represents the presence
of m^{5}C methylation with high (dark blue) or low (light blue) stoichiometry.

451
452

Extended Data Fig. 6
The dynamics of tRNA modifications throughout the IDC.
(a) Pairwise correlations of the mutation frequency at all detected sites between replicates. **(b)**
Changes in the average mutation frequencies after AlkB demethylase treatment (N=2).
Modifications of G26 and U73 were insensitive to treatment. **(c)** Heatmap showing changes in
the mutation frequency at each site during IDC. **(d)** Left panel: proportion of pre- and mature
Tyr-tRNA during IDC. Right panel: mutation frequencies of pre- and mature Tyr-tRNA at G26-
, G37- and A58-modified sites during IDC. Error bars in all panels represent the SD of N=3
biological replicates.
462
463

Extended Data Fig. 7

The characteristics of the apicoplast tRNA transcriptome.

(a) All G37 sites on apicoplast tRNAs are methylated. A37 (orange)/G37 (green) indicates the
nucleotide identity at position 37. (+) denotes the detection of mutation signature. (Dup.)
indicates duplicated genes. **(b)** Integrative Genomics Viewer (IGV) image showing the
truncation at the wobble position * in (lower) a potential Ile2-tRNA (PF3D7_API00600), and
no truncation was observed in (upper) the Met-tRNA (PF3D7_API05000). **(c)** An illustration
of all possible codon anticodon pairings predicted by the wobble hypothesis. The cognate
tRNA for the ATA codon was not found in the apicoplast genome. The number of tRNA genes
per anticodon (tRNA) and the number of codons used (frequency) by the apicoplast genome
are indicated. **(d)** tRNAscan-SE predicted an Ile2-tRNA.

475
476
477

478
Extended Data Fig. 8
Highly expressed genes minimize dependency on certain types of amino acids.
(a) Correlation between the usage frequency and gene expression level in the trophozoite
stage (LogTPM) based on a fitted linear trendline of nine tRNAs. Spearman's rank
correlation coefficients (ρ) are indicated. (n=5267)**. (b)** Spearman correlations between the
demand of tRNA usage frequency and gene expression level (LogTPM) throughout the IDC.
(c) Spearman correlation between amino acid usage frequency and gene expression level
(LogTPM) throughout the IDC. The LogTPM value for each stage is the average of N=3.
487
488
489

Extended Data Fig. 9

Limited changes in the abundance and modification levels of the anticodon pool after

acute amino acid depletion.

(a) Changes in tRNA composition and **(b)** mutation frequency at each position detected after

6 hours of treatment in trophozoites 32-36 h.p.i. (CM: complete medium; AA-: AA depletion;

and HF: halofuginone (70 nM) and AA depletion). Error bars represent the SD of N=3

biological replicates. **(c)** The relative isoacceptor usage of the 5% most highly expressed coding

genes (x-axis, n=258) and the 5% least expressed coding genes (y-axis, n=258). Relative

isoacceptor usage for each tRNA was calculated as (the fraction of all codons encoding a

specific amino acid that is decoded by an isoacceptor/the fraction of all isoacceptors for a

specific amino acid that are equally used). The most highly expressed genes preferred Ile-AAU,

but the least expressed genes preferred Ile-UAU.

Extended Data Fig. 10
Acute amino acid depletion causes delayed progression.
(a) FACS plots showing the fluorescence intensity of dihydroethidium (DHE) and Hoechst
DNA staining in parasites (left) at the beginning and (right) after 6 hours of AA depletion
(AA-) compared to parasites grown in complete medium (CM). Both DHE and Hoechst
staining were reduced in AA- parasites after 6 hours. **(b)** The geometric mean of the Hoechst
staining intensity in parasites (left) at the beginning and (right) after 6 hours of AA depletion
relative to that in CM parasites. Supplementation of AA- parasites with isoleucine rescued the
reduction in DNA content in a dose-dependent manner. Error bars represent the SD for N=3
biological replicates.

Extended Data Fig. 11

Stage progression is isoleucine sensitive.

(a) DESeq2 differential gene expression analyses of the transcriptomes during stage
progression from (left) the schizont-to-ring stage, (middle) the ring-to-trophozoite stage, and
(right) the trophozoite-to-schizont stage. The horizontal and vertical dotted lines show the cut-
off of the FDR-adjusted p-value=0.05 and cut-off of the twofold change, respectively. N=3
biological replicates. **(b)** Empirical CDF plot of isoleucine and leucine usage in up- (cyan) and
downregulated (red) genes during stage progression. There was an increase in isoleucine
sensitivity during the ring-to-schizont progression, which was reflected by higher isoleucine
usage for the upregulated genes but reduced sensitivity during the schizont-to-ring transition.
The p-values of Mann–Whitney U tests are indicated. **(c)** Isoleucine and leucine usage in the
10% of genes with the most variable expression (the 10th decile) and the 10% of genes with the
least variable expression (1st decile) throughout the IDC. Variability was defined by the
coefficient of variation in expression (CV of TPM) throughout the IDC.

Extended Data Fig. 12

Increased ribosome loading stabilizes transcripts and reprogrammes the transcriptome.
(a) Kozak similarity scores (KSS, using ribosomal genes as reference) in transcripts with
increased polysome association (IPA) or decreased polysome association (DPA) after amino
acid depletion. Transcripts with increased polysome association exhibited lower scores,
indicating an initiation context less similar to that of the highly translated ribosomal genes. The
p-value of the Mann–Whitney U test is indicated. **(b)** DESeq2 differential gene expression
analyses of the transcriptome after amino acid depletion. The horizontal and vertical dotted
lines show the cut-off of the FDR-adjusted p-value=0.05 and cut-off of twofold change,
respectively. N=3 biological replicates. **(c)** The strategy used to categorize stage-dependent and
stage-independent changes in the transcriptome. The transcriptome of the AA-starved parasites
(AA-) was expected to most closely resemble a trophozoite-like transcriptional profile due to
delayed stage progression (upper). Transcripts that were more abundant in trophozoites were
also expected to be upregulated in AA- parasites and vice versa, as illustrated in the Venn
diagram. **(d)** The KSS of up- and downregulated genes in the stage-dependent changes (left
panel) and stage-independent changes (right panel). Only stage-independent changes show a
disparity in KSS between the up- and downregulated genes similar to that in the polysome
fractions, supporting a positive feedback loop. The p-values of Mann–Whitney U tests are
indicated. **(e)** The correlation of the changes in transcript abundance in the polysome fractions
and those in the total transcriptome after amino acid depletion. The differences in **(f)** isoleucine
and leucine usage and **(g)** the Kozak context in transcripts that remained stable after amino acid
depletion but were expected to be downregulated (stabilized) or upregulated (destabilized)
according to the categorization shown in Panel (c). Gene Ontology analysis of **(h)** the
downregulated genes and **(i)** the upregulated genes in the stage-independent change group.
Only the top 20 enrichment terms are shown. Downregulated genes were enriched in
proteasomal pathways, suggesting a lower protein turnover. No pathway was significantly

enriched (FDR<0.05) with upregulated genes, but many genes were found to be involved in
various metabolic pathways, potentially priming cells for rapid metabolic reactivation.
566
567
568

Extended Data Fig. 13
Transcript stabilization depends on the interplay between translation initiation and Ile
codon-induced stalling.
(a) Anota2seq analysis of the polysome dataset using rlog normalization with default
parameters. Anota2seq delineates different regulatory modes. Buffered (mRNA down)
represents genes with a relative increase in ribosome loading but with no transcript-stabilizing
effect. N=3 biological replicates. **(b)** Isoleucine and leucine usage in "buffered mRNA up" and
"buffered mRNA down" transcripts. **(c)** (Left) KSS of "buffered mRNA up" and "buffered
mRNA down" transcripts. The p-values of the Mann–Whitney U tests are indicated. (Right)
The Kozak context of "buffered mRNA up" and "buffered mRNA down" transcripts. Kozak
context: strong, RnnATGR; adequate, RnnATGY/YnnATGR; and weak, YnnATGY (ATG=
start codon). The p-value of the Fisher's exact test is indicated.
581
582

Extended Data Fig. 14
The expansion of Ile homorepeats is under purifying selection.
(Upper) Heatmap showing the frequency of di-amino acid motifs relative to their expected
frequency. Values of (observed/expected frequency) were calculated for each di-amino acid
motif, and all values were Z-transformed to generate the heatmap. (Lower) The fraction of the
motifs retained in the successive additions of an extra residue. The dotted lines represent the
expected frequency based on the amino acid usage frequency in the genome. Four of the most
frequently used hydrophobic amino acids were analysed. Instead of being more likely to occur
due to replication slippage, both di-Ile and Ile homorepeat expansion were found to be less
frequent than expected.

Reference

1. Evans, M.E., Clark, W.C., Zheng, G., and Pan, T. (2017). Determination of tRNA aminoacylation levels by high-throughput sequencing. Nucleic Acids Res *45*, e133. 10.1093/nar/gkx514.

2. Zheng, G., Qin, Y., Clark, W.C., Dai, Q., Yi, C., He, C., Lambowitz, A.M., and Pan, T. (2015). Efficient and quantitative high-throughput tRNA sequencing. Nat Methods *12*, 835-837. 10.1038/nmeth.3478.

3. Behrens, A., Rodschinka, G., and Nedialkova, D.D. (2021). High-resolution quantitative profiling of tRNA abundance and modification status in eukaryotes by mim-tRNAseq. Mol Cell *81*, 1802-1815.e1807. 10.1016/j.molcel.2021.01.028.

4. Xu, H., Yao, J., Wu, D.C., and Lambowitz, A.M. (2019). Improved TGIRT-seq methods for comprehensive transcriptome profiling with decreased adapter dimer formation and bias correction. Sci Rep *9*, 7953. 10.1038/s41598-019-44457-z.

5. Cozen, A.E., Quartley, E., Holmes, A.D., Hrabeta-Robinson, E., Phizicky, E.M., and Lowe, T.M. (2015). ARM-seq: AlkB-facilitated RNA methylation sequencing reveals a complex landscape of modified tRNA fragments. Nat Methods *12*, 879-884. 10.1038/nmeth.3508.

6. Chan, P.P., and Lowe, T.M. (2016). GtRNAdb 2.0: an expanded database of transfer RNA genes identified in complete and draft genomes. Nucleic Acids Res *44*, D184-189. 10.1093/nar/gkv1309.

7. Cooley, L., Appel, B., and Söll, D. (1982). Post-transcriptional nucleotide addition is responsible for the formation of the 5' terminus of histidine tRNA. Proc Natl Acad Sci U S A *79*, 6475-6479. 10.1073/pnas.79.21.6475.

8. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat Methods *9*, 357-359. 10.1038/nmeth.1923.

9. Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work with high-throughput sequencing data. Bioinformatics *31*, 166-169. 10.1093/bioinformatics/btu638.

10. Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics *30*, 2114-2120. 10.1093/bioinformatics/btu170.

11. Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol *37*, 907-915. 10.1038/s41587-019-0201-4.

12. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550. 10.1186/s13059-014-0550-8.

13. Malleret, B., Claser, C., Ong, A.S., Suwanarusk, R., Sriprawat, K., Howland, S.W., Russell, B., Nosten, F., and Rénia, L. (2011). A rapid and robust tri-color flow cytometry assay for monitoring malaria parasite development. Sci Rep *1*, 118. 10.1038/srep00118.

14. Lacsina, J.R., LaMonte, G., Nicchitta, C.V., and Chi, J.T. (2011). Polysome profiling of the malaria parasite Plasmodium falciparum. Mol Biochem Parasitol *179*, 42-46. 10.1016/j.molbiopara.2011.05.003.

15. Altenhoff, A.M., Train, C.M., Gilbert, K.J., Mediratta, I., Mendes de Farias, T., Moi, D., Nevers, Y., Radoykova, H.S., Rossier, V., Warwick Vesztrocy, A., et al. (2021). OMA orthology in 2021: website overhaul, conserved isoforms, ancestral gene order and more. Nucleic Acids Res *49*, D373-D379. 10.1093/nar/gkaa1007.