Agrobacteria deploy two classes of His-Me finger superfamily nuclease effectors

exerting different antibacterial capacities against specific bacterial competitors

Mary Nia M. Santos[a,c,d,f], Katherine L. Pintor[a,1], Pei-Yu Hsieh[b,2], Yee-Wai Cheung[a], Li-

Kang Sung[a,c,d], Yu-Ling Shih[b] and Erh-Min Lai[a,c,e,*]

[a] Institute of Plant and Microbial Biology, Academia Sinica, Taipei 115201 Taiwan

[b] Institute of Biological Chemistry, Academia Sinica, Taipei 115201 Taiwan

[c] Molecular and Biological Agricultural Sciences Program, Taiwan International

Graduate Program, National Chung-Hsing University and Academia Sinica, Taipei,

Taiwan

[d] Graduate Institute of Biotechnology, National Chung-Hsing University, Taichung,

Taiwan

[e] Biotechnology Center, National Chung-Hsing University, Taichung, Taiwan

[f] Aquaculture Research and Development Division, Department of Agriculture -
National Fisheries Research and Development Institute (DA-NFRDI), Manila,
Philippines

Running title: His-Me finger T6SS nuclease toxins

*Corresponding author
Erh-Min Lai: emlai@gate.sinica.edu.tw
Institute of Plant and Microbial Biology, Academia Sinica
128 Sec. 2, Academia Road, Nankang, Taipei, Taiwan 115201

Current address:
[1]Department of Ecophysiology, Max Planck Institute for Terrestrial Microbiology,
Karl-von-Frisch-Str. 10, 35043 Marburg, Germany
[2]Department of Pharmacology, University of Cambridge, United Kingdom

ORCID
MNMS https://orcid.org/0000-0001-6544-0339
KLP https://orcid.org/0000-0001-9137-4726
PYH https://orcid.org/0000-0002-7263-1691
YWC https://orcid.org/0000-0002-1421-4419
LKS https://orcid.org/0000-0002-7078-8003
YLS https://orcid.org/0000-0002-8232-2569
EML https://orcid.org/0000-0003-3630-8683

ABSTRACT

Type VI secretion system (T6SS) assembles into a contractile nanomachine to inject

effectors across bacterial membranes for secretion. *Agrobacterium tumefaciens*

species complex is a group of soil inhabitants and phytopathogens that deploys T6SS

as an antibacterial weapon against bacterial competitors at both inter-species and

intra-species levels. *A. tumefaciens* strain 1D1609 genome encodes one main T6SS

gene cluster and four *vrgG* genes (i.e. *vgrGa-d*), each encoding a spike protein as an

effector carrier. Previous study reported that *vgrGa*-associated gene 2, named as *v2a,*

encodes a His-Me finger nuclease toxin (also named as HNH/ ENDO VII nuclease)

contributing to DNase-mediated antibacterial activity. However, the functions and roles

of other putative effectors remain unknown. In this study, we identified *vgrGc-*

associated gene 2 (*v2c*) that encodes another His-Me finger nuclease but with distinct

SHH motif differed from AHH motif of V2a. We demonstrated that the ectopic

expression of V2c caused growth inhibition, plasmid DNA degradation, and cell

elongation in *Escherichia coli*. The cognate immunity protein, V3c, neutralizes the

DNase activity and rescues phenotypes of the growth inhibition and cell elongation.

Ectopic expression of V2c DNase-inactive variants retains the cell elongation

phenotype while V2a induced cell elongation in a DNase-mediated manner. We also

showed that the amino acids of conserved SHH and HNH motifs are responsible for

the V2c DNase activity *in vivo* and *in vitro*. Notably, V2c also mediated the DNA

degradation and cell elongation of target cell in the context of interbacterial competition.

Importantly, V2a and V2c exhibit different capacities against different bacterial species

and function synergistically to exert stronger antibacterial activity against the soft rot

phytopathogen, *Dickeya dadantii*.

INTRODUCTION

The type VI secretion system (T6SS) is a nanomachine used by many Gram-negative

bacteria for antagonism or pathogenesis by injecting toxins into target bacterial or

eukaryotic cells [1-3]. T6SS is composed of a membrane complex in connection to a

baseplate, which is the docking site for polymerization of a tube surrounded by a

contractile sheath [4]. The tube is a puncturing device stacked of Hcp protein hexamer

tipped with the trimeric Valine-glycine repeat protein G (VgrG) spike and a PAAR

repeat-containing protein which sharpens the tip [5]. When the sheath contracts, the

puncturing device carrying effectors is propelled towards the target cell to deliver
effectors. After firing, the sheath disassembles and the subunits of the sheath are
recycled to build another machine [6].
75
In terms of delivery, the T6SS effectors can be classified as cargo or specialized
effectors. The cargo effectors interact non-covalently with Hcp, VgrG or PAAR while
in specialized effectors, the effector domain is covalently fused to Hcp, VgrG or PAAR
[7]. Functionally, most of the identified effectors mediate antibacterial activities
including nuclease, peptidoglycan hydrolase, lipase, phospholipase, NAD(P)+-
glycohydrolase and ADP ribosyltransferase by targeting conserved cellular structures
such as nucleic acids, peptidoglycan, and inner membrane as well as specific
metabolites or proteins [8, 9].
84
To date, the characterized T6SS nuclease effectors include members of HNH, N-Tox,
Tox-Rease, and PoNe superfamilies and were shown to target DNA substrates without
specificity [10-16]. Thus, expression of these T6SS nuclease effectors causes DNA
degradation *in vitro* and/or *in vivo*. Interestingly, the majority of T6SS DNase effectors
belong to HNH/ENDO VII nuclease superfamily, also named as His-Me finger
endonuclease superfamily consisting of 38 distinct families [17, 18]. These nuclease
superfamily proteins harbor three conserved His-Asp-His within a ~30-amino-acid
domain formed by a β1 hairpin followed by an α-helix to form a ββα-metal topology
for DNA binding and hydrolysis [17, 18]. Several T6SS effectors belonging to this
HNH/His-Me superfamily have been identified, including Tse7 of *Pseudomonas*
aeruginosa as Tox-GHH2 [15], Tse1 of *Aeromonas dhakensis* [13]*,* V2a of
Agrobacterium tumefaciens as Tox-AHH [11], and two Tox-SHH family effectors Tke4
of *Pseudomonas putida* [19] and Txe4 of fish pathogen *Pseudomonas plecoglossicida*
[20]. Amino acid substitution experiments on Tse7, V2a, and Tse1 have indicated that
this conserved catalytic site [A/G]HH is required for the nuclease activity and toxicity
[11, 13, 15]. However, the impact of SHH and HNH motifs of the Tox-SHH family
effectors in nuclease activity and T6SS toxicity have not been demonstrated.
102
A. tumefaciens is a Gram-negative Alphaproteobacteria belonging to the
Rhizobiaceae family. It is an economically important pathogen that causes crown gall
disease and a gene delivery tool with capability of transforming plant cells and fungi

[21]. T6SS main gene cluster is highly conserved in the genome of *A. tumefaciens*
genomospecies complex [22-24]. Besides the main cluster consisting of two divergent
operons, *imp* (<u>imp</u>aired in nitrogen fixation) encoding core structural and regulatory
components and *hcp* operon for effectors and effector delivery components (e.g. Hcp,
VgrG, PAAR). Most *A. tumefaciens* strains also encode additional *vgrG* operons
encoding effector-immunity (EI) gene pairs. *A. tumefaciens* strain 1D1609 encodes
four *vgrG* genes (*vgrGa, vgrGb, vgrGc, vgrGd*) with each *vgrG* module associated with
three to four associated genes, named as <u>*vgrG-associated*</u> gene <u>1-4</u> in cluster <u>*a/b/c/d*</u>
[11]. Each *vgrG* gene is genetically linked to a conserved chaperone/adaptor gene
and different known or putative EI pairs, including three effectors with a conserved N-
terminal PAAR-like DUF4150 domain but distinct C-terminal effector domains (V2a,
V2c, and V2d) [11] (**Figure S1**). For *vgrGb* module, V2b does not encode an effector
domain but downstream two genes encode a putative ADP-ribosylating enzyme (V3b)
and a Rhs-linked effector domain (V4b), followed by a putative immunity. Previous
study by deletion of single or multiple EI pairs suggested that *vgrGa-associated*
effector V2a, a DNase effector harboring C-terminal Tox-AHH domain, appears to be
the major antibacterial toxin but *vgrGd-associated* effector V2d also contributes to full
antibacterial activity for 1D1609 against *Escherichia coli* prey [11]. The biochemical
and biological functions of other putative effectors encoded in the other three orphan
vgrG modules remain uncharacterized.
126
In this study, we discovered that *vgrGc*-associated effector V2c is a Tox-SHH DNase
toxin and explored the rationale of having two T6SS DNase toxins in *A. tumefaciens*
strain 1D1609. We show that the expression of V2c caused growth inhibition, plasmid
DNA degradation and cell elongation, which the phenotypes are neutralized by co-
expression of the cognate immunity protein, V3c. We also show that the SHH motif is
responsible for the nuclease activity of V2c. Importantly, V2a AHH nuclease and V2c
SHH nuclease exert different antibacterial capacities against specific bacterial
competitors and function synergistically to exert stronger antibacterial activity against
the soft rot phytopathogen, *Dickeya dadantii*. Harboring two classes of His-Me finger
nuclease effectors with potential target-specific toxicity may grant 1D1609 with
versatile antibacterial weapons in facing different bacterial competitors in microbial
community.

RESULTS

V2c is a DNase belonging to the Tox-SHH clade of His-Me finger superfamily

In 1D1609, *vgrGa*-associated effector V2a appears to be the major antibacterial toxin

against *Escherichia coli* prey [11]. V2c is a putative PAAR (DUF4150)-linked

specialized effector (**Figure 1A**) but the presence or absence of *v2c* did not affect the

antibacterial activity of 1D1609 against *E. coli* prey [11]. Aside from the N-terminal

PAAR domain, V2c does not show any sequence similarity to V2a (**Figure S2**). BLAST

and CDD searches did not identify known or conserved domain at its C-terminal region.

However, multiple sequence alignment of V2c and V2c orthologs revealed conserved

SHH and HNH motifs at the C-terminal region (**Figure 1B**). These V2c orthologs all

harbor C-terminal Tox-SHH domain but their N-terminal region can be fused to other

known T6SS domains such as <u>F</u>ound in type s<u>IX</u> effector (FIX), DUF4150/PAAR-like

domain and Rhs core domain ([25], **Figure 1C).** Using the protein homology detection

program HHpred [26], we found that the C-terminal region of V2c encodes two parallel

β-strands connected by α-helix, which is signature structure in His-Me finger nuclease

[17]. Consistent with the result of secondary structure prediction, 3D structure

prediction using AlphaFold ([27, 28], **Figure 1D**) shows this domain consists of two

antiparallel β-strands connected with an Ω loop with a His (H384) at the C-terminus of

the first β-strand and followed by an α-helix, featuring a HNH motif of His-Me finger

endonuclease [17, 18]. These suggest V2c belongs to the Tox-SHH (Pfam PH15652)

clade of His-Me finger superfamily.

To determine the antibacterial and DNase activities of V2c, we expressed *v2c* using

an arabinose inducible promoter to determine if V2c is a bacterial toxin functioning as

a nuclease. Growth inhibition measured by optical density was observed when *v2c* is

expressed by arabinose induction in *Escherichia coli* and this can be neutralized by

co-expression of its putative cognate immunity *v3c* as compared to the vector control

(**Figure 2A**). Also, colony forming unit (CFU) count shows the reduced CFU of *v2c*-

expressed cells and higher CFU could be recovered with co-expression of *v3c* (**Figure**

2B). In addition, plasmid degradation assay showed that the expression of *v2c* upon

arabinose induction resulted in degradation of plasmid DNA in *E. coli* and this is

abolished when *v3c* is co-expressed (**Figure 2C**). The data indicated that V2c exhibits

DNase activity *in vivo*.

We further investigated whether the SHH and HNH motifs are involved in DNase

activity. His-Me finger superfamily has a strictly conserved His residue and catalytic

metal ion which are both essential for nucleic acid hydrolysis. Based on this premise,

we define a region spanning amino acid residues 380-409 as a His-Me finger domain

core region (CR). Based on the alignment, H384 is the catalytic His and H408 is for

metal ion binding while H383 may function as second residue for metal ion

coordination. To verify their roles in antibacterial and DNase activity, a series of

deletion as well as single and double amino acid substitution variants were generated.

Strikingly, no growth inhibition and plasmid degradation were observed in any of the

mutants expressing the CR deletion (V2c^{CR}), single (V2c^{H383A}, V2c^{H384A} and V2c^{H408A})

and double (V2c^{HAHA} substitution in both H383 and H384) variants (**Figure 2A, C, D**).

Higher CFU was also recovered from *E. coli* culture expressing these mutants alone

or with *v3c* (**Figure 2B**). To further confirm the DNase activity of V2c*, each of V2c and

its variants is fused with 6x-His and purified for *in vitro* DNase activity assay. The

results showed that purified V2c fraction is able to degrade DNA, but not the eluate

from *E. coli* expressing the vector (pET28a) (**Figure 2D**). Consistent with the *in vivo*

plasmid degradation assay, no activity could be detected from V2c^{H384A}*, V2c^{H408A}*, or

V2c^{HAHA} variants. However, purified V2c^{H383A} fraction exhibited DNase activity by

degrading the plasmid DNA substrate. Western blot analysis of the reaction samples

shows V2c and its variants used for the assay were at similar protein levels. Taken the

results of *in vivo* and *in vitro* DNA degradation assays, the conserved H384 and H408

in the HNH motif are crucial for V2c DNase activity whereas H383 in SHH motif

appears to be less critical for *in vitro* but may be required for *in vivo* nuclease activity.

Ectopic expression of V2c caused DNA degradation and cell elongation

We next examined the cell morphology and DNA integrity of *E. coli* cells expressing

v2c and the two variants (*v2c^{CR}* and *v2c^{HAHA}*) that are defective in DNase activity and

toxicity. Cell morphology was observed by phase contrast while bacterial cell

membrane and DNA were stained by FM-4-64 and Hoechst, respectively under

fluorescence microscope. Hoechst signals are significantly reduced when *v2c* is

expressed by arabinose induction in *E. coli* and this can be rescued by co-expression

of its cognate immunity *v3c* shown in the images and quantification (**Figure 3A, B**).

Hoechst signals are diminished only in *E. coli* cells producing *v2c* but not *v2c^{HAHA}*,

$v2c^{CR}$ and vector control, suggesting that the cellular DNA degradation is caused by

V2c DNase activity, which can be neutralized by its immunity protein V3c. In addition,

we also observed cell elongation in *E. coli* expressing *v2c* and co-expression of *v3c*

rescues the cell elongation phenotype (**Figure 3A, C**). However, *E. coli* cells

expressing $v2c^{CR}$ and $v2c^{HAHA}$ as well as other single amino acid substitution variants

($v2c^{H383A}$, $v2c^{H384A}$, $v2c^{H408A}$) do not abolish the elongated cell phenotype (**Figure 3A,**

C, Figure S3A, B).

Ectopic expression of V2a also caused cell elongation phenotype

Previous study showed that V2a is a Tox-AHH nuclease that serves as the major

antibacterial weapon in 1D1609 [11]. H385 and H386 are two of the predicted His

residues in AHH motif of V2a, corresponding to H383 and H384 in SHH motif of V2c.

The structure predicted by AlphaFold indicates V2a also features a His-Me finger,

having two α-helices with histidine residues (H430, H456) followed by the β-strands

(**Figure 4A**). Previous findings demonstrated that the expression of *v2a* but not

$v2a^{H385A}$ caused growth inhibition and plasmid degradation in *E. coli* [11]. To gather

further insight of V2a AHH motif in the His-Me finger domain, we evaluated the

growth and morphology of arabinose induced *v2a* mutants. Results show that the

expression of *v2a* caused reduction in Hoechst-stained DNA signals and resulted in

cell elongation (**Figure 4B, C, D**). The expression of $v2a^{H385A}$ has similar levels of

Hoechst-stained DNA signals and cell length as compared to the vector control.

These results suggest that the cell elongation phenotype observed were caused by

V2a DNase activity, whereas V2c shows DNase activity-independent cell elongation.

We also determined the toxicity of another putative effector V4b of 1D1609, a PAAR

(DUF4150)-linked specialized effector but harboring a Rhs domain with structural

similarity to the insecticidal toxin [11] (**Figure S4A**). Expression of *v4b* caused

growth inhibition in *E. coli,* in which the growth inhibition can be rescued by co-

expression of its putative cognate immunity *v5b* to a level similar to vector control

(**Figure S4B**). However, no DNA degradation and cell elongation could be observed

(**Figure S4C, D, E, F**). These results show that V4b is a bacterial toxin and V5b is

the cognate immunity protein and suggest that cell elongation phenotype may be

specific to toxicity of nuclease effectors.

V2c nuclease exhibits antibacterial activity at both intra-species and inter-species competition and functions synergistically with V2a against the soft rot phytopathogen, *Dickeya dadantii*

To correlate the DNA degradation and cell elongation phenotypes observed by ectopic expression of *v2c* in *E. coli* in a more biologically relevant context, we further performed interbacterial competition assay. We first selected *A. tumefaciens* strain C58, which possess incompatible EI pairs with 1D1609 and previously demonstrated to be susceptible to 1D1609 T6SS killing [22], for intra-species competition. Each of 1D1609 mutants lacking multiple EI pairs attackers was co-cultured with C58 strain harboring pRL-GFP(S65T) prey *in vitro* on AK agar [29] and the number of viable C58 prey cells (log$_{10}$ CFU/mL) were quantified. Similar to previous killing assay using *E. coli* prey [11], we detected ~ 2 log of reduced prey cell survival by 1D1609 WT attacker, as compared to the Δ*4EI* mutant with deletion of all four EI pairs (**Figure 5A**). Other double *(ΔEIbd)* or triple deletion mutants (Δ*EIabd*, Δ*EIbcd*) exhibited compromised but detectable antibacterial activity to C58, indicating that *v2a* or *v2c* alone is sufficient to exhibit antibacterial activity to C58.

Intrigued by the observation that *v2c* could exhibit antibacterial activity to *A. tumefaciens* C58 prey but no significant effect against *E. coli* prey [11], we reasoned that *E. coli* may not be physiologically relevant prey for *Agrobacterium* although *v2c* is able to cause toxicity when directly expressed in *E. coli* (**Figure 2A, B**). We then selected *Dickeya dadantii* as the prey cells for inter-species competition because it is a phytopathogen but also a close relative of *E. coli*, both belonging to *Enterobacteriaceae*. To avoid potential counterattack by *D. dadantii* T6SS [30], a T6SS-deficient mutant, Δ*imp* with deletion of *imp* operon, harboring pRL-GFP(S65T) (*D. dadantii* Δ*imp*-GFP(S65T)) was used as a prey. CFU counting of *D. dadantii* Δ*imp*-GFP(S65T) prey cells after co-culture show that ~ 1.5 log of reduced prey cell survival by 1D1609 WT attacker, as compared to the Δ*4EI* mutant (**Figure 5B**). The presence of *v2a* only (Δ*3EIbcd*) exhibited comparable antibacterial activity as the WT while *v2c* only (Δ*3EIabd*) showed weak but detectable antibacterial activity as compared to Δ*4EI,* indicating *v2c* itself is able to kill *D. dadantii* but with much weaker activity than *v2a*. Strikingly, 1D1609 expressing both *v2a* and *v2c* (Δ*2EIbd*) could result in stronger killing effects than WT. The use of pRL-GFP(S65T) with constitutive expression of GFP also allowed us to observe the *D. dadantii* Δ*imp*-GFP(S65T) prey cells under the

fluorescence microscope after co-culture. We observed very few cells expressing GFP

when competing with WT or Δ*3EIbcd* expressing *v2a* only*. The lysed and elongated

D. dadantii cells were also detected when co-cultured with 1D1609 with either strains

expressing *v2a* only (Δ*3EIbcd*), *v2c* only (Δ*3EIabd*), or both Δ*3EIbd* (**Figure 5C**). We

further quantified the cell length of GFP-expressing *D. dadantii* cells. The results show

that majority of cells were elongated after 18 hours of co-culture with 1D1609

expressing either *v2a* (Δ*3EIbcd*) or *v2c* (Δ*3EIabd*), or combination of both (Δ*3EIbd*),

as compared to those *D. dadantii* cells alone or co-incubated with Δ*4EI* (**Figure 5D**).

Taken together, these results suggest that V2c nuclease exhibits antibacterial activity

at both intra-species and inter-species competition but at different capacities. While

V2a and V2c exert similar antibacterial against *A. tumefaciens* C58, V2c only exhibits

weak antibacterial activity but functions synergistically with V2a when competing with

D. dadantii prey, resulting to elongated and lysed cells.

DISCUSSION

His-Me finger nucleases are a large and diverse superfamily of nucleases; however,

only a limited number of T6SS effectors have been identified as His-Me finger

nuclease. In the present study, we reported that *A. tumefaciens* strain 1D1609

encodes two His-Me finger superfamily DNases, V2a and V2c belonging to Tox-AHH

and Tox-SHH family respectively, exhibiting different capacities to specific bacterial

competitors.

His-Me finger nucleases are one-metal-ion-dependent nucleases, primarily dependent

on Mg^{2+} or in some cases dependent on Zn^{2+}. The only invariant residue in His-Me

endonuclease is the catalytic His located at the end of β1 strand (1st H in HNH motif,

H384 for V2c and H386 for V2a, **Figure 1D, 4A, S2)**, which functions as the general

base for hydrolysis of phosphodiester bonds. The other conserved His (last H in HNH

motif, H408 for V2c and H430 for V2a) and Asn/Asp/Glu/His located before the

invariant His (H383 for V2c and H384 for V2a) are involved in coordinating the single

metal ion. Indeed, our data show that H383, H384, and H408 of V2c and H385 and

H386 of V2a are required for exerting DNase activity *in vivo* (**Figure 2C,** [11]. However,

different from the loss of DNase activity *in vivo* and *in vitro* for both V2c[H384A] and

V2c^{H3408A}, V2c^{H383A} variant retains the DNase activity by an *in vitro* nuclease activity

assay (**Figure 2D**). This indicates the requirement of H383 in coordinating metal ion

binding for DNase activity may be subject to environmental factors.

His-Me finger encodes a small structural motif, so it is usually associated with other

domain architectures that could play additional functions. In addition, its three-element

structural motif is unable to provide enough of a hydrophobic core for stability [18].

Thus, many proteins containing the His-Me finger employ a variety of additional

structural elements or domains for stabilization and/or specificity. V2c and its orthologs

commonly found in *Rhizobiaceae* are not included in the reported 77 sequences with

conserved core regions of His-Me finger superfamily [18]. Our evidence further

suggests that V2c may represent a new branch of Tox-SHH toxin family (Pfam

PF15652), which includes proteins with additional domain architectures (**Figure 1C**).

These additional domain(s) are conserved regions found in many bacterial toxin

proteins and may facilitate the secretion of nuclease and affect its toxicity to the

competing cell. While both V2c and V2a harbor the conserved N-terminal DUF4150

PAAR-like domain, they share low sequence similarity in the C-terminal His-Me finger

domain (**Figure S2**), in which the protein structures predicted by AlphaFold reveal the

structural difference between V2c and V2a (**Figure 1C, 4A**). This discrepancy may

contribute to the difference on toxicity strength and the role of DNase on cell elongation.

Alternatively, V2a and V2c may have the same mechanism to bind and hydrolyze DNA

for toxicity and the difference in the phenotype is caused by another domain outside

the His-Me finger fold.

Cell elongation could still be observed in the *E. coli* strain carrying different DNase

defective *v2c* variants, even in the absence of arabinose (**Figure 3A, B, S2A**). Several

studies reported the leaky expression of T6SS toxin driven by arabinose-inducible

P$_{BAD}$ promoter resulted in toxicity in the absence of arabinose [16, 25]. Thus, we

speculated that the leaky expression of these *v2c* variants is sufficient to induce the

cell elongation phenotype. The data suggest that another domain outside the SHH

catalytic region in V2c is responsible for the cell elongation phenotype. This is different

from the nuclease activity-dependent filamentation observed previously [15, 31] and

V2a (**Figure 4**), suggesting that V2c-induced cell elongation is independent of

nuclease activity. The proposed biological role of bacterial filamentation is mainly

related to stress response to thrive for survival [32]. One of the conditions that lead to

filamentation is DNA damage. When the DNA is damaged, the SOS response is

induced and cells may delay the cell cycle and inhibit cell division until DNA repair and

replication are complete. Although V2c catalytic site variants lose activity for DNA

cleavage, they may still bind to DNA and induce cell elongation. Furthermore, cell

division arrest could be also caused by non-nuclease effectors. For example, a T6SS

ADP-ribosyltransferase effector called Tre1 in *Serratia proteamaculans* was reported

to inactivate bacterial cell division Tre1 by modifying arginine on FtsZ to block

polymerization [33]. Future work by domain dissection of V2c is needed to identify the

domain causing cell elongation.

While HNH/His-Me T6SS nucleases are the most prevalent nuclease toxins identified

in T6SS, most bacterial genomes only encode one type of HNH/His-Me nuclease

effector, including Tse7 of *P. aeruginosa* [15], Tse1 of *A. dhakensis* [13]*,* and Tke4 of

P. putida [19]. Besides *A. tumefaciens* 1D1609 encoding Tox-AHH (V2a) and Tox-

SHH (V2c) nucleases, the use of multiple HNH/His-Me nucleases as T6SS

antibacterial weapons have been identified in fish pathogen *P. plecoglossicida* [20]. *P.*

plecoglossicida T6SS-2 mediates interbacterial competition and encodes four putative

effectors, all contain C-terminal toxin domains belonging to HNH/His-Me superfamily

but distinct classes (Txe1 with a dipeptide HH motif, Txe2 as Tox-AHH, Txe3 with

HNHc, and Txe4 as Tox-SHH) [20]. Among them, Txe1, Txe2, and Txe4 exhibit *in vitro*

nuclease activity and toxicity when expressed in *E. coli*. However, only Txe1 and Txe4

contributes to interbacterial activity against *E. coli,* with Txe1 as the predominant toxin,

in which its C-terminal conserved dipeptide HH motif is required for nuclease activity

and toxicity to *E. coli.* Txe4 is also required for full interbacterial competition but the

role of SHH motif in DNase and toxicity has not been determined. Together with our

findings, it is possible that having multiple distinct or same classes of nuclease

effectors may provide versatile toxicity when competing with different preys, in which

some toxins may only exhibit activity against specific preys.

The strain 1D1609 is unique in *A. tumefaciens* species complex with multiple VgrG

spikes carrying different antibacterial effectors [11] . Our findings that *v2a* and *v2c*

exhibit different capacities against different preys support previous studies that some

T6SS toxins can only exert toxicity in specific bacterial species [23, 34]. From the

observation of the synergism of V2a and V2c against *D. dadantii,* harboring multiple

T6SS effectors in 1D1609 may provide an advantage for the strain to maintain a

growth competitiveness in different niches and environments.

MATERIALS AND METHODS

Bacterial strains and growth conditions

Bacterial strains and plasmids are listed in Table S1. *Escherichia coli* strains were

grown in Luria-Bertani (LB) medium at 37°C, *A. tumefaciens* strains in 523 medium at

25°C and *D. dadantii* 3937 in LB medium at 30°C. Antibiotics were added when

necessary: for *E. coli*, 30 µg/mL gentamicin and 250 µg/mL streptomycin; for *D.*

dadantii and *A. tumefaciens,* 50 µg/mL gentamicin.

Molecular cloning

Plasmid pJN105 was used for expression of toxin genes and pTrc200 plasmid was

used for expression of immunity gene. PCR was amplified with KAPA HiFi HotStart

DNA Polymerase (Roche, Switzerland) using genomic DNA of *A. tumefaciens* 1D1609.

The primers used in the study are listed in Table S2. Site directed mutagenesis was

done using either DpnI or Gibson Assembly (New England BioLabs, USA), ligated

using T4 ligase (Takara) and transformed into chemically competent *E. coli* DH10B.

Plasmid DNA was isolated using QIAprep miniprep kit (Qiagen, Germany). All

constructs were confirmed by sequencing. The in-frame deletion mutants in *A.*

tumefaciens were generated using a suicide plasmid via double crossing over by

electroporation or by conjugation as described [11].

Bioinformatic analysis

The orthologs of V2c were analyzed using amino acid sequences after the DUF4150

domain via BLASTP search. Multiple sequence alignment was performed using

MUSCLE or Multiple Sequence Comparison by Log- Expectation [35]. Full length

sequences of the V2c orthologs were analyzed to obtain domain architectures from

Pfam database [36]. The E-value threshold was set at 10^{-5}. HHpred [26] and Swiss

Model [37] were used for homology-based protein structure prediction. Protein

structure was predicted by AlphaFold [27, 28] from the UniProt database. Secondary

structure prediction was done using default settings in PSIPRED 4.0 Protein

Structure Prediction Server/ DMPfold 1.0 Fast Mode [38, 39].

Growth inhibition assay

Growth inhibition assay was performed as described previously [10]. In brief, overnight cultures of *E. coli* DH10B strain with empty vectors (pJN105 for effector gene and pTrc200 for immunity gene) or the derivatives were adjusted to OD_{600} of 0.1 in LB medium. 1 mM isopropyl β-D-1- thiogalactopyranoside (IPTG, Amresco) was used to induce the expression of the putative immunity protein. After 1 hour of IPTG induction, L-arabinose (Ara, Sigma-Aldrich) was added to 0.2% final concentration, to induce expression of the toxin. Cell growth was recorded every hour at OD_{600}. Empty vectors were used as controls. After 8 hours, cells were harvested to quantify the number of colony forming units ($\log_{10}$ CFU/mL) by automatic diluter and plater, easySpiral Dilute (Interscience, France).

Plasmid DNA degradation analysis in *E. coli* cells

In vivo plasmid DNA degradation analysis was performed as described previously [10]. In brief, overnight cultures of *E. coli* DH10B strain with empty vectors (pJN105 for effector gene and pTrc200 for immunity gene) or the derivatives grown in LB broth were harvested and adjusted to OD_{600} ~0.3 in LB medium. *E. coli* cultures were induced with 1 mM IPTG at 0 hour for *v3c* expressed from pTrc200 plasmid followed by L-arabinose (0.2%, final concentration) induction at 1 hour to induce *v2c* from pJN105 plasmid and cultured for 2 hours. Equal amounts of cells were used for plasmid DNA extraction and equal volume of extracted DNA was resolved in agarose gel followed by ethidium bromide staining and visualized using Gel Doc XR+ UV Gel Documentation Molecular Imager Universal Hood II (Bio-Rad, USA).

Cloning, expression and purification of V2c and mutants

For *in vitro DNase* activity assay, the effector gene *v2c* was constructed in pET28a plasmid and immunity gene *v3c* was constructed in pTrc200 plasmid. Site-directed mutagenesis was done by using overlap extension PCR. Plasmid for expressing effector protein, *v2c* or its variants, and plasmid for expressing immunity protein, *v3c*, were transformed into *E. coli* BL21(DE3) for IPTG-induced co-expression. Bacterial culture was incubated at 37 °C until OD_{600} reached 0.5, then 0.1 mM IPTG was added followed by 5-h expression at 37°C. Pellets were collected, sonicated, and expressed proteins were purified by using Ni Sepharose 6 Fast Flow histidine-tagged protein

purification resin (Cytiva, Germany). Briefly, cells were lysed in extraction buffer (50

mM Tris-Cl, pH7.5, 0.1 M NaCl, 20 mM imidazole) and proteins were eluted by using

elution buffer with up to 250 mM imidazole in 50 mM Tris-Cl, pH7.5, 0.1 M NaCl.

***In vitro* DNase activity assay**

A 10 µL reaction containing 200 ng of pTrc200HA plasmid DNA and 1× CutSmart

Buffer (New England BioLabs, USA) with partially purified V2c or its variants

normalized by the major V2c protein band was incubated at 37°C for 1 hour. The

digestion of DNA was determined by agarose gel electrophoresis and visualized by

SYBR Safe DNA gel stain (Invitrogen, USA). The reaction samples of V2c or its

variants were scaled up for Western blotting analysis. Anti-V2c antibody was

generated in rabbits against two V2c peptides, PFYWDFPNSQVGRD and

KRIAMLRGDPTRYD (indicated in Figure S1) by Yao-Hong Biotechnology Inc.

(Taiwan).

Western blot analysis

Western blot analysis was performed as described [40]. In general, proteins were

resolved by SDS-PAGE and transferred onto a PVDF membrane by using a transfer

apparatus (Bio-Rad, USA). The membrane was probed with primary antibody against

V2c (1:4000), Hcp (1:2500), TssB (1:4000) and GFP (1:1000), followed by incubation

with horseradish peroxidase-conjugated anti-rabbit secondary antibody (Chemichem)

(1:25000) and visualized with the ECL system (Perkin-Elmer Life Science, Boston,

USA).

Fluorescence microscopy and image analysis

E. coli DH10B carrying the empty vectors (pJN105 for effector gene and pTrc200 for

immunity gene) or the derivatives were grown overnight at 37°C. Overnight cultures

were sub-cultured with OD_{600} ~0.01. After 1 hour, L-arabinose (Ara) was added (0.2%

final concentration) to induction. Cultures were harvested after an hour and

resuspended with 30 µL PBS with 0.5 % Tween 20 (PBST). Three µL of cells was

placed on 2% agarose pad and imaged. To check the DNA degradation, the same

protocol was followed and live cells were stained with 2 µg/mL Hoechst (Life

Technologies, USA) and 1 µg/mL FM4-64FX, fixable analog of FM-4-64 (Invitrogen,

USA) for 5 mins and washed with PBS to remove the excess dye.

For co-culture of *D. dadantii* Δ*imp*-GFP(S65T) with various 1D1609 EI pair mutants,

strains were mixed at 1:10 ratio on AK agar, *Agrobacterium* kill-triggering (AK) medium

(3 g K_2HPO_4, 1 g NaH_2PO_4, 1 g NH_4Cl, 0.15 g KCl, 9.76 g MES in 900 mL ddH_2O, pH

5.5) [29] solidified by 2% (w/v) agar. Cells from 16 hour co-culture at 25°C were

concentrated to OD_{600} ~10 and spotted on 2% agarose pad for imaging.

Imaging was performed using an inverted fluorescent microscope (IX81, Olympus

Japan) with objective lens UPLSAPO 100XO (Olympus, Tokyo, Japan), DAPI (49000;

Chroma), GFP (49002; Chroma) and Cy3 (49005; Chroma) filter set and CCD camera

(ORCA-R2; Hamamatsu, Japan). The images were acquired using Olympus cellSens

Dimension software.

All the image analysis was done using Fiji analyze particles tool [41]. The micrograph

of FM-4-64 fluorescence was used to segment surfaces (i.e. individual bacterial cell)

with background subtraction and thresholding. Feret's diameter was used. The cell

length (μm) was determined from cumulative counts of three random frames per

sample. For quantification of cell degradation, the Hoechst intensity was normalized

using the formula intensity/unit area of cells, where intensity is major x minor x 3.14

(π).

Interbacterial competition assay

Bacterial competitions were carried out as described [22] with modifications. Briefly,

overnight culture of *A. tumefaciens* and *D. dadantii* 3937 Δ*imp* were mixed at a 10:1

attacker to prey ratio and spotted on Agrobacterium kill-triggering (AK) agar plates [29].

The prey *A. tumefaciens* C58 or *D. dadantii* 3937 Δ*imp* was transformed with pRL662-

GFP(S65T) for gentamycin selection or microscopy imaging. After 18 hour incubation

at 25°C, the co-cultured cells were collected, serially diluted and plated on LB agar

containing gentamicin to quantify surviving prey cells by counting CFU.

Statistical analysis

The statistical analyses were done using GraphPad Prism version 6.01 for Windows,

GraphPad Software, La Jolla California USA (www.graphpad.com). The number of

technical replicates and independent biological replicates, p values and statistical

tests performed are indicated in the figure legends. The mean of independent

replicates was compared using the following statistical tests: Student's t test, one-

way ANOVA followed by Tukey's multiple comparisons tests

(http://astatsa.com/OneWay_Anova_with_TukeyHSD/). The error bars indicate the

standard error of the mean (SEM).

ACKNOWLEDGEMENTS

The authors acknowledge Ching-Hong Yang for providing *Dickeya dadantii* 3937

strains, which were imported under the permit 106-B-003 issued by the Council of

Agriculture of Taiwan. We would like to thank the former and current Lai lab

members for their help and fruitful discussion throughout this study, as well as See-

Yeun Ting and Mao-Sen Liu for critically reading the manuscript and their valuable

comments. We also thank Venus Marie Gaela and Thomas Boudier for the guidance

and help in processing the images using Fiji and the Sanger DNA sequencing

service provided by Genomic Technology Core located at the Institute of Plant and

Microbial Biology, Academia Sinica. The funding was supported by grants National

Science and Technology Council (NSTC) of Taiwan (107-2311-B-001-019-MY3) and

Academia Sinica Investigator Award (AS-IA-107-L01) to EML, and NSTC grant (110-

2311-B001-011) to YLS. YWC was supported by the postdoctoral fellowship (NSTC

110-2811-B-001-504). The funders had no role in study design, data collection and

interpretation, or the decision to submit the work for publication.

AUTHOR CONTRIBUTION

518 Conceptualization: MNMS, EML; Investigation: MNMS, KP, PYH, YWC, LKS, YLS,

519 EML; Project Funding acquisition and administration: EML, YLS; Supervision: EML,

520 YLS; Writing of original draft: MNMS, EML; Writing – methodology, review & editing:

521 MNMS, KP, PYH, YWC, LKS, YLS, EML.

REFERENCES

1. Hood RD, Peterson SB, Mougous JD. From Striking Out to Striking Gold: Discovering
that Type VI Secretion Targets Bacteria. Cell Host Microbe. 2017;21(3):286-9. doi:
10.1016/j.chom.2017.02.001. PubMed PMID: 28279332.

2.   Coulthurst S. The Type VI secretion system: a versatile bacterial weapon.
Microbiology (Reading). 2019;165(5):503-15. Epub 2019/03/21. doi: 10.1099/mic.0.000789.
PubMed PMID: 30893029.

3.   Hachani A, Wood TE, Filloux A. Type VI secretion and anti-host effectors. Curr Opin
Microbiol. 2016;29:81-93. Epub 2016/01/02. doi: 10.1016/j.mib.2015.11.006. PubMed
PMID: 26722980.

4.   Wang J, Brodmann M, Basler M. Assembly and Subcellular Localization of Bacterial
Type VI Secretion Systems. Annu Rev Microbiol. 2019;73:621-38. Epub 2019/06/22. doi:
10.1146/annurev-micro-020518-115420. PubMed PMID: 31226022.

5.   Shneider MM, Buth SA, Ho BT, Basler M, Mekalanos JJ, Leiman PG. PAAR-repeat
proteins sharpen and diversify the type VI secretion system spike. Nature.
2013;500(7462):350-3. Epub 2013/08/09. doi: 10.1038/nature12453. PubMed PMID:
23925114; PubMed Central PMCID: PMC3792578.

6.   Basler M, Pilhofer M, Henderson GP, Jensen GJ, Mekalanos JJ. Type VI secretion
requires a dynamic contractile phage tail-like structure. Nature. 2012;483(7388):182-6. Epub
2012/03/01. doi: nature10846 [pii]

10.1038/nature10846. PubMed PMID: 22367545; PubMed Central PMCID: PMC3527127.

7.   Cianfanelli FR, Monlezun L, Coulthurst SJ. Aim, Load, Fire: The Type VI Secretion
System, a Bacterial Nanoweapon. Trends Microbiol. 2016;24(1):51-62. doi:
10.1016/j.tim.2015.10.005. PubMed PMID: 26549582.

8.   Lien YW, Lai EM. Type VI Secretion Effectors: Methodologies and Biology. Front Cell
Infect Microbiol. 2017;7:254. doi: 10.3389/fcimb.2017.00254. PubMed PMID: 28664151;
PubMed Central PMCID: PMC5471719.

9.   Cherrak Y, Flaugnatti N, Durand E, Journet L, Cascales E. Structure and Activity of the
Type VI Secretion System. Microbiol Spectr. 2019;7(4). Epub 2019/07/13. doi:
10.1128/microbiolspec.PSIB-0031-2019. PubMed PMID: 31298206.

10.   Ma LS, Hachani A, Lin JS, Filloux A, Lai EM. Agrobacterium tumefaciens deploys a
superfamily of type VI secretion DNase effectors as weapons for interbacterial competition
in planta. Cell Host Microbe. 2014;16(1):94-104. doi: 10.1016/j.chom.2014.06.002. PubMed
PMID: 24981331; PubMed Central PMCID: PMC4096383.

11.   Santos MNM, Cho ST, Wu CF, Chang CJ, Kuo CH, Lai EM. Redundancy and Specificity
of Type VI Secretion vgrG Loci in Antibacterial Activity of Agrobacterium tumefaciens
1D1609 Strain. Frontiers in Microbiology. 2020;10. doi: ARTN 3004

10.3389/fmicb.2019.03004. PubMed PMID: WOS:000510643200001.

12.   Ma J, Pan Z, Huang J, Sun M, Lu C, Yao H. The Hcp proteins fused with diverse
extended-toxin domains represent a novel pattern of antibacterial effectors in type VI
secretion systems. Virulence. 2017;8(7):1189-202. Epub 20170106. doi:
10.1080/21505594.2017.1279374. PubMed PMID: 28060574; PubMed Central PMCID:
PMCPMC5711352.

13.   Pei TT, Li H, Liang X, Wang ZH, Liu G, Wu LL, et al. Intramolecular chaperone-
mediated secretion of an Rhs effector toxin by a type VI secretion system. Nat Commun.

2020;11(1):1865. Epub 20200420. doi: 10.1038/s41467-020-15774-z. PubMed PMID:
32313027; PubMed Central PMCID: PMCPMC7170923.

14.   Fitzsimons TC, Lewis JM, Wright A, Kleifeld O, Schittenhelm RB, Powell D, et al.
Identification of Novel Acinetobacter baumannii Type VI Secretion System Antibacterial
Effector and Immunity Pairs. Infect Immun. 2018;86(8). Epub 20180723. doi:
10.1128/IAI.00297-18. PubMed PMID: 29735524; PubMed Central PMCID:
PMCPMC6056853.

15.   Pissaridou P, Allsopp LP, Wettstadt S, Howard SA, Mavridou DAI, Filloux A. The
Pseudomonas aeruginosa T6SS-VgrG1b spike is topped by a PAAR protein eliciting DNA
damage to bacterial competitors. Proc Natl Acad Sci U S A. 2018;115(49):12519-24. Epub
2018/11/21. doi: 10.1073/pnas.1814181115. PubMed PMID: 30455305; PubMed Central
PMCID: PMCPMC6298103.

16.   Fridman CM, Keppel K, Gerlic M, Bosis E, Salomon D. A comparative genomics
methodology reveals a widespread family of membrane-disrupting T6SS effectors. Nat
Commun. 2020;11(1):1085. Epub 20200227. doi: 10.1038/s41467-020-14951-4. PubMed
PMID: 32109231; PubMed Central PMCID: PMCPMC7046647.

17.   Wu CC, Lin JLJ, Yuan HS. Structures, Mechanisms, and Functions of His-Me Finger
Nucleases. Trends Biochem Sci. 2020;45(11):935-46. Epub 20200814. doi:
10.1016/j.tibs.2020.07.002. PubMed PMID: 32807610.

18.   Jablonska J, Matelska D, Steczkiewicz K, Ginalski K. Systematic classification of the
His-Me finger superfamily. Nucleic Acids Res. 2017;45(20):11479-94. doi:
10.1093/nar/gkx924. PubMed PMID: 29040665; PubMed Central PMCID: PMCPMC5714182.

19.   Bernal P, Allsopp LP, Filloux A, Llamas MA. The Pseudomonas putida T6SS is a plant
warden against phytopathogens. ISME J. 2017;11(4):972-87. Epub 2017/01/04. doi:
10.1038/ismej.2016.169. PubMed PMID: 28045455; PubMed Central PMCID:
PMCPMC5363822.

20.   Li Y, Yan X, Tao Z. Two Type VI Secretion DNase Effectors are Utilized for
Interbacterial Competition in the Fish Pathogen Pseudomonas plecoglossicida. Front
Microbiol. 2022;13:869278. Epub 20220406. doi: 10.3389/fmicb.2022.869278. PubMed
PMID: 35464968; PubMed Central PMCID: PMCPMC9020831.

21.   Hwang HH, Yu M, Lai EM. Agrobacterium-Mediated Plant Transformation: Biology
and Applications: American Society of Plant Biologists; 2017.

22.   Wu CF, Santos MNM, Cho ST, Chang HH, Tsai YM, Smith DA, et al. Plant-Pathogenic
Agrobacterium tumefaciens Strains Have Diverse Type VI Effector-Immunity Pairs and Vary
in In-Planta Competitiveness. Mol Plant Microbe In. 2019;32(8):961-71. doi: 10.1094/Mpmi-
01-19-0021-R. PubMed PMID: WOS:000477044200007.

23.   Wu CF, Weisberg AJ, Davis EW, 2nd, Chou L, Khan S, Lai EM, et al. Diversification of
the Type VI Secretion System in Agrobacteria. mBio. 2021;12(5):e0192721. Epub
2021/09/15. doi: 10.1128/mBio.01927-21. PubMed PMID: 34517758; PubMed Central
PMCID: PMCPMC8546570.

24.   Chou L, Lin YC, Haryono M, Santos MNM, Cho ST, Weisberg AJ, et al. Modular
evolution of secretion systems and virulence plasmids in a bacterial species complex. BMC

Biol. 2022;20(1):16. Epub 2022/01/14. doi: 10.1186/s12915-021-01221-y. PubMed PMID:
35022048; PubMed Central PMCID: PMCPMC8756689.

25.   Jana B, Fridman CM, Bosis E, Salomon D. A modular effector with a DNase domain
and a marker for T6SS substrates. Nat Commun. 2019;10(1):3595. Epub 20190809. doi:
10.1038/s41467-019-11546-6. PubMed PMID: 31399579; PubMed Central PMCID:
PMCPMC6688995.

26.   Soding J, Biegert A, Lupas AN. The HHpred interactive server for protein homology
detection and structure prediction. Nucleic Acids Res. 2005;33(Web Server issue):W244-8.
doi: 10.1093/nar/gki408. PubMed PMID: 15980461; PubMed Central PMCID:
PMCPMC1160169.

27.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly
accurate protein structure prediction with AlphaFold. Nature. 2021;596(7873):583-9. Epub
20210715. doi: 10.1038/s41586-021-03819-2. PubMed PMID: 34265844; PubMed Central
PMCID: PMCPMC8371605.

28.   Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold
Protein Structure Database: massively expanding the structural coverage of protein-
sequence space with high-accuracy models. Nucleic Acids Res. 2022;50(D1):D439-D44. doi:
10.1093/nar/gkab1061. PubMed PMID: 34791371; PubMed Central PMCID:
PMCPMC8728224.

29.   Yu M, Wang YC, Huang CJ, Ma LS, Lai EM. Agrobacterium tumefaciens Deploys a
Versatile Antibacterial Strategy to Increase its Competitiveness. J Bacteriol. 2020. Epub
2020/11/11. doi: 10.1128/JB.00490-20. PubMed PMID: 33168638.

30.   Koskiniemi S, Lamoureux JG, Nikolakakis KC, t'Kint de Roodenbeke C, Kaplan MD,
Low DA, et al. Rhs proteins from diverse bacteria mediate intercellular competition. Proc
Natl Acad Sci U S A. 2013;110(17):7032-7. Epub 2013/04/11. doi: 1300627110 [pii]

10.1073/pnas.1300627110. PubMed PMID: 23572593; PubMed Central PMCID:
PMC3637788.

31.   Yu M, Lai EM. Warfare between Host Immunity and Bacterial Weapons. Cell Host
Microbe. 2017;21(1):3-4. doi: 10.1016/j.chom.2016.12.012. PubMed PMID: 28081441.

32.   Justice SS, Hunstad DA, Cegelski L, Hultgren SJ. Morphological plasticity as a bacterial
survival strategy. Nature Reviews Microbiology. 2008;6(2):162-8. doi: 10.1038/nrmicro1820.
PubMed PMID: WOS:000252426700015.

33.   Ting SY, Bosch DE, Mangiameli SM, Radey MC, Huang S, Park YJ, et al. Bifunctional
Immunity Proteins Protect Bacteria against FtsZ-Targeting ADP-Ribosylating Toxins. Cell.
2018;175(5):1380-92 e14. Epub 20181018. doi: 10.1016/j.cell.2018.09.037. PubMed PMID:
30343895; PubMed Central PMCID: PMCPMC6239978.

34.   LaCourse KD, Peterson SB, Kulasekara HD, Radey MC, Kim J, Mougous JD. Conditional
toxicity and synergy drive diversity among antibacterial effectors. Nat Microbiol.
2018;3(4):440-6. Epub 2018/02/21. doi: 10.1038/s41564-018-0113-y. PubMed PMID:
29459733; PubMed Central PMCID: PMCPMC5876133.

35.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res. 2004;32(5):1792-7. doi: 10.1093/nar/gkh340. PubMed PMID:
15034147; PubMed Central PMCID: PMC390337.

36.   Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, et al.
Pfam: The protein families database in 2021. Nucleic Acids Res. 2021;49(D1):D412-D9. doi:
10.1093/nar/gkaa913. PubMed PMID: 33125078; PubMed Central PMCID:
PMCPMC7779014.

37.   Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, et al. SWISS-
MODEL: homology modelling of protein structures and complexes. Nucleic Acids Res.
2018;46(W1):W296-W303. doi: 10.1093/nar/gky427. PubMed PMID: 29788355; PubMed
Central PMCID: PMCPMC6030848.

38.   Greener JG, Kandathil SM, Jones DT. Deep learning extends de novo protein
modelling coverage of genomes using iteratively predicted structural constraints. Nat
Commun. 2019;10(1):3977. Epub 20190904. doi: 10.1038/s41467-019-11994-0. PubMed
PMID: 31484923; PubMed Central PMCID: PMCPMC6726615.

39.   Buchan DWA, Jones DT. The PSIPRED Protein Analysis Workbench: 20 years on.
Nucleic Acids Res. 2019;47(W1):W402-W7. doi: 10.1093/nar/gkz297. PubMed PMID:
31251384; PubMed Central PMCID: PMCPMC6602445.

40.   Lai EM, Kado CI. Processed VirB2 is the major subunit of the promiscuous pilus of
Agrobacterium tumefaciens. J Bacteriol. 1998;180(10):2711-7. Epub 1998/06/06. PubMed
PMID: 9573157; PubMed Central PMCID: PMC107224.

41.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82. Epub
20120628. doi: 10.1038/nmeth.2019. PubMed PMID: 22743772; PubMed Central PMCID:
PMCPMC3855844.

FIGURE LEGENDS

Figure 1. V2c is a PAAR-linked specialized effector containing C-terminal Tox-

SHH motif. (A) Schematic diagram of V2c protein showing N-terminal DUF4150

PAAR region and the catalytic region (amino acid 380-409) which covers the Tox-

SHH motif. **(B)** Partial sequence alignment of orthologs of V2c from BLASTP. The

strain name and accession number are indicated on the left. SHH motif is highlighted

in yellow, with H383 (indicated in red) as the metal ion coordinating residue, H384

and H408 (indicated in green) of HNH motif for catalysis and metal ion binding

respectively, are targeted for mutagenesis. The star corresponds to the specific

domains shown in Figure 1C: *DUF4150/PAAR-like, **DUF4150/PAAR-like + FIX-

like, ***FIX-like superfamily, **** PAAR + Rhs core domain. **(C)** Domain architectures

of V2c orthologs, Tox-SHH-containing proteins found in this study. Full length

sequences were analyzed and the domains were from Pfam database with E value

threshold set at 10^{-5}. **(D)** Cartoon model of V2c (residue 380-409). The structure of

V2c predicted by AlphaFold indicates V2c consisting a His-Me finger domain with

two antiparallel β-strands (yellow) connected with an Ω loop (light blue), a histidine

(H384) at the C-terminus of the first β-strand and followed by an α-helix (magenta).

The confidence score of this region is higher than 90, indicating high structure

accuracy.

Figure 2. V2c exhibits growth inhibition, DNase activity *in vivo* and *in vitro.* (A)

Growth inhibition analysis of expression of *v2c,* its variants, with or without *v3c* in *E.*

coli DH10B. *E. coli* cultures were induced with 1 mM IPTG at 0 hour for *v3c* expressed

from pTrc200 plasmid followed by L-arabinose (Ara) induction at 1 hour to induce *v2c*

from pJN105 plasmid. Cell growth was recorded every hour at OD_{600}. Strain

expressing empty vectors (vectors) were used as a control. **(B)** Number of colony

forming unit ($\log_{10}$ CFU/mL) quantified after growth inhibition assay at 8 hour. Data of

(A) and (B) represents mean ± SEM of four independent experiments (n=4), each

averaged with 3 technical repeats. Different letters above the bar indicate statistically

different groups of strains (P < 0.05) determined by Tukey's HSD test. **(C)** Plasmid

degradation assay performed by *E. coli* DH10B cells harboring pTrc200 and pJN105

plasmids (vector) or the derivatives induced with (+) or without (–) Ara for 2 hour.

Plasmid DNA was extracted and the degradation pattern was observed in agarose gel.

(D) *In vitro* DNase activity assay performed by using pTrc200HA plasmid DNA as DNA

substrate and incubated with purified V2c or its variants. Nuclease free water (ddH_2O),

protein elution buffer (elution buffer) and eluate from empty vector expression (pET28a)

served as negative controls while Benzonase nuclease served as a positive control.

The digestion of DNA substrate was determined by agarose gel electrophoresis. The

reaction samples of V2c or its variants were detected by western blotting using anti-

V2c antibody (α-V2c). Results of (C) and (D) are representative of three independent

experiments.

Figure 3. V2c Tox-SHH DNase exhibits cell elongation phenotype. (A) Images of

E. coli DH10B cells harboring vector(s) and the derivatives expressing *v2c* or its

variants in the presence or absence of its cognate immunity gene *v3c* with (+Ara) and

without (-Ara) Arabinose induction stained with FM-4-64 (red) and Hoechst stain (blue).

The micrographs from left to right are FM-4-64, Hoechst, phase contrast and a merged

image of FM-4-64 and Hoechst stains. Scale bar, 2 µm. Representative images of two

independent experiments are shown. **(B)** Normalized Hoechst intensity (intensity/unit

area of cells) and **(C)** Cell length (µm) as measured from the indicated number of cells

(n) per treatment. The graph shows a combined count from three frames of a

representative result of two independent experiments. The red line in (C) shows the

median with interquartile range. Statistics were performed with data mean ± SEM of

three frames. Different letters above the bar indicate statistically different groups of

strains ($P < 0.05$) determined by Tukey's HSD test.

Figure 4. V2a Tox-AHH DNase exhibits cell elongation phenotype. (A) Cartoon

model of V2a (residue 380-470). The structure of V2a predicted by AlphaFold indicates

V2a consisting of a His-Me finger domain with its signature antiparallel β-strands

(yellow) connected with an Ω loop (light blue) with a histidine (H386, orange) at the C-

terminus of the first β-strand. Two α-helices (pink) with histidine residues (H430, H456

magenta) followed the β-strands. N412 is found in the loop between the β-strands and

H430 is in the first α-helix (E429–G430). The predicted Local-Distance Difference Test

(pLDDTs) of the His-Me domains of V2a are from confident to very high, most of the

pLDDTs of the residues are > 90. **(B)** Morphological analysis of *E. coli* DH10B cells

harboring vector(s) or its derivatives expressing *v2a*, catalytic site (H385A) mutant

without (-Ara) or with (+Ara) arabinose induction. Cells were stained with FM-4-64 (red)

and Hoechst stain (blue). The micrographs from left to right are FM-4-64, Hoechst,

phase contrast and a merged image of the two fluorescent images. Scale bar, 2 µm.

(C) Cell length (µm) and **(D)** Hoechst intensity in different treatments as determined

from a combined count of three random frames of a representative result, the number

of cells (n) per sample is indicated. The graph shows a combined count from three

frames of a representative result of two independent experiments. The red line shows

the median with interquartile range. Statistics were performed with data mean ± SEM

of three frames. Different letters above the bar indicate statistically different groups of

strains ($P < 0.05$) determined by Tukey's HSD test.

Figure 5. V2a and V2c exhibit different toxicity against *A. tumefaciens* and *D.

dadantii*. (A) Intra-species competition between *A. tumefaciens* 1D1609 attacker and

prey C58. **(B), (C), (D)** Inter-species competition between *A. tumefaciens* 1D1609

attacker and prey *D. dadantii* 3937 Δ*imp.* The plasmid pRL-GFP(S65T) plasmid (Gm^R)

was transformed into prey cells, which were mixed at 1:10 ratio with each of the

attacker strains, 1D1609 WT or various EI mutant strains as indicated. After co-culture

on AK agar plates, cells was collected, serially diluted, and plated on LB agar plate

containing Gm for CFU counting of surviving prey cells **(B)**. Data are mean ± SEM (n

= 6 biological repeats from three independent experiments). For imaging, co-cultured

D. dadantii Δimp-GFP(S65T) and 1D1609 EI pair mutant were concentrated and

spotted on 2% agarose pad for fluorescence micrographs **(C)**. *D. dadantii Δimp*-

GFP(S65T) indicated as Dd only is included as a control. The inset in phase contrast

shows a magnified image of cells in the red box. White arrows point to *D. dadantii* cells.

Scale bar, 2 μm. **(D)** Cell length after competition between *A. tumefaciens* 1D1609 EI

mutants with *D. dadantii* 3937 *Δimp*-GFP(S65T). Cell length of GFP fluorescent *D.*

dadantii cells was measured using Fiji analyze particles tool. Cell length (μm) in

different treatments as determined from a combined count of at least 10 random

frames of two independent experiments. The red line shows the median with

interquartile range. Statistics were performed with data mean ± SEM of total counts.

Different letters above the bar indicate statistically different groups of strains (P < 0.05)

determined by Tukey's HSD test.

Figure S1. Genetic loci of *A. tumefaciens* 1D1609 T6SS gene cluster and *vgrG*

genetic modules. The four *vgrG* genes (*vgrGa, vgrGb, vgrGc, vgrGd*) and their

associated effector-immunity pairs are indicated on linear chromosome. The number

of *vgrG*-associated genes *(v)* represents the position to *vgrG* as defined previously

[11].

Figure S2. Sequence alignment of V2a and V2c. The conserved amino acid

residues are indicated as '∗'   (identical), ':' (similar) and '.' (slightly similar). The

DUF4150, PAAR-like domain is in green. The histidine residues in AHH and SHH

motifs in His-Me finger domain of V2a and V2c respectively, V2aH385/V2cH383 and

V2aH386/V2cH384, are in red. The histidine residues that predicted to responsible for

metal binding in His-Me finger domain, V2aH430/V2cH408, are in a bold blue font.

Two peptides used for generating antibodies were underlined.

Figure S3. Cell length analysis of *E. coli* DH10B cells expressing *v2c* catalytic
site variants. (A) Expression of *v2c* single catalytic site variants. **(B)** Expression of
catalytic site variants co-expressed with *v3c*. Cell length (μm) as measured from the
indicated number of cells (n) per sample. The graph shows a combined count from six
random frames of a representative result, the number of cells (n) per sample is
indicated. The graph shows a combined count from 6 random frames, 3 from each of
two independent experiments. The red line shows the median with interquartile range.
Statistics were performed with data mean ± SEM of 6 frames. Different letters above
the bar indicate statistically different group of strains (P < 0.05) determined by Tukey's
HSD test. The red line shows the median with interquartile range.

Figure S4. V4b exhibits growth inhibition, which could be neutralized by V5b.
(A) Schematic diagram of V4b. The Rhs domain is bordered by conserved motifs. The
CT is the C-terminal domain encoding a putative muramidase toxin. **(B)** *E. coli* growth
inhibition analysis. *E. coli* DH10B cultures were induced with 1 mM IPTG at 0 hour for
the expression of putative immunity gene *v5b* from pTrc200 plasmid followed by L-
arabinose (Ara) induction at 1 hour to induce putative *v4b* toxin gene from pJN105
plasmid. Cell growth was recorded every hour at OD_{600}. DH10B expressing empty
vectors were used as a control. Data are mean ± SD of three biological replicates.
Similar results were observed in three independent experiments. **(C)** Plasmid
degradation assay performed by using full length V4b (*v4b*^{FL}) and C-terminal domain
only (*v4b*^{CT}). *E. coli* cells with pTrc200 and pJN105 plasmid (vectors) or expressing
v4b and *v4b*+*v5b* were induced with (+) or without (-) arabinose (Ara) for 2 h. Plasmid
DNA was extracted and the degradation pattern was observed in agarose gel. Result
shown is a representative data from two independent experiments. **(D)** Images of *E.*
coli DH10B cells expressing *v4b* and *v4b* co-expressed with its cognate immunity gene
v5b (*v4b*+*v5b*) under condition with (+Ara) or without (-Ara) arabinose induction. *E.*
coli DH10B cells expressing empty vectors (vectors) under condition with (+Ara) and
without (-Ara) arabinose induction served as negative controls. Cells were stained with
FM-4-64 (red) and Hoechst stain (blue). The micrographs from left to right are FM-4-
64, Hoechst, phase contrast and a merged image of the two fluorescent micrographs.
Representative images of two independent experiments are shown. Scale bar, 2 μm.
(E) Cell length (μm) and **(F)** Hoechst intensity in cells analyzed in (D). The red line
shows the median with interquartile range. The graph shows a combined count from

3 random frames of a representative result, the number of cells (n) per sample is

indicated Statistics were performed with data mean ± SEM of three frames. Different

letters above the bar indicate statistically different groups of strains (P < 0.05)

determined by Tukey's HSD test.

Supplementary Table S1. Bacterial strain and plasmids used in this study

Supplementary Table S2. Primers used in this study**.**

Figure 1. V2c is a PAAR-linked specialized effector containing C-terminal Tox-SHH motif.

(A)

(B)

(C)

DUF4150/PAAR-like		Tox-SHH

DUF4150/PAAR-like	FIX-like superfamily	Tox-SHH

	FIX-like superfamily	Tox-SHH

PAAR	Rhs core domain	Tox-SHH

(D)

N388

H384

H408

Figure 2. V2c exhibits growth inhibition, DNase activity *in vivo* and *in vitro*.

Figure 3. V2c Tox-SHH DNase exhibits cell elongation phenotype.

Figure 4. V2a Tox-AHH DNase exhibits cell elongation phenotype.

Figure 5. V2a and V2c exhibit different toxicity against *A. tumefaciens* and *D. dadantii*.

Figure S1. Genetic loci of *A. tumefaciens* 1D1609 T6SS gene cluster and *vgrG* genetic modules.

Figure S2. Sequence alignment of V2a and V2c.

```
V2a    MSIPRDNYIGEPQYPGTWTTATPREGLRDIDEARIVSLAPDVCLTPVGSSVVPIPYPVVD
V2c    MSIPRDNYIGEPQYPGTWTTATPREGLRDIDEARIVSLAPDVCLTPVGSSVVPIPYPVVD
       ***********************************************************
V2a    FCGHDKNYTPSVRFTGKKAMVMRSFTTHVHGDAPGNRKGVKSGTVESICEPIGHADQVRA
V2c    FCGHDKNYTPSVRFTGKKAMVMRSFTTHVHGDAPGNRKGVKSGTVESICEPIGHADQVRA
       ***********************************************************
V2a    EGSHVIRHLDRFYMNSRNTQGEAIFVRSTQTYDPPKDDDPVRGSLRWQDGSDEGRVMSDA
V2c    EGSHVIRHLDRFYMNSRNTQGEAIFVRSTQTYDPPKDDDPVRGSLRWQDGSDEGRVMSDA
       ***********************************************************
V2a    SPEPFKEGEKYAYAGSAAAATTGAAQNSAQALRGGMPATRPTVPQTVPRGPGRVITPGPA
V2c    SPEPLIMGAQYAQAQTATIPQSGGV-----ARRGTYGRPTTTVDPRLLKGL-RFWSSARG
       ****:   * :** * :*: . :*..     * **   . .**   : .*   *. :.. .
V2a    RAP---QWYRTPPHRVVSDLA-KVG---------RLARVGIWAR-RLNVIGILSSVLIPD
V2c    AAEIMDMVQQISEDRAVQDSAGQLGLNLSDPTDVLVARAHAWSKYHIDASPFYWD--FPN
        *         . .  *.*.* * ::*       :**.  *:. .::.  . .  :*:
V2a    NGANLSDVVPRDDQERRIKEMFDEGVDAGHDSWEMRDWAMEELRRHRQRVREEPESRPQA
V2c    SQ------VGRDAATKAITSM--EAAAPG----TLGE-ALQGDRTQLEKLRTAIKNAVSA
       .       * **   . *..*  *.. .*     : : *::  * : :..:*   :.   .*
V2a    LPQSETVRVSDRDEYRKKCQVDRYGKMKSVCGQYGMQAHHIVPDWTLRYGAREEADKRIP
V2c    ATAAASVRVSG-----QPCAREALG----APGKYRGGAHFAT---ALPLGDNLDSHHTPA
        . : :****.     : *   : *    . *:*   **. .   :*   *  .:: .  .
V2a    NMPSLNDGMAICVIGNARVEQTEHWGGHLAD------KAIENAGRNSTPPFTAPLHKVVQ
V2c    NSASY--------------LPTWAGPAIQMLPEDHRATLSYGRKVTSPYISTQARLIN
       * .*                *.*  :      .*  . **: *.*: :.  .:::
V2a    ASAQAMISVRPDCAKEIMAALTTQYAGKNPFQLLRAK--QYPPLPEETISALKSGAVEGA
V2c    SG------------QFMKAINMDILDIKRIAMLRGDPTRYDAAIAQMMAYAECLKREGI
       :.            ::* *:. :   . : : :**.. .* .   : :: :.    **
V2a    TRKP
V2c    IK--
        .
```

Figure S3. Cell length analysis of _E. coli_ DH10B cells expressing _v2c_ catalytic site variants.

(A)

(B)

Figure S4. V4b exhibits growth inhibition, which could be neutralized by V5b.

Table S1. Bacterial strains and plasmids

Strain	Relevant characteristics	Reference/Source	EML No.
E. coli			
DH10B	Host of plasmid for cloning	Lab stock	EML455
S17-1	Host for conjugation	Lab stock	EML124
BL21 (DE3)	For protein expression	Lab stock	EML117
D. dadantii **3937**			
WT	Wild-type *Dickeya dadantii* 3937	Ching-Hong Yang, University of Wisconsin, Milwaukee	EML4911
Δ*imp*	T6SS inactive strain with deletion of *imp* operon	Ching-Hong Yang, University of Wisconsin, Milwaukee	EML4921
A. tumefaciens **1D1609**			
WT	Alfalfa isolate, virulent strain containing pTi1D1609	Clarence Kado, UC Davis	EML304
Δ*4* or Δ*vgrGabcd*	Deletion of *vgrGa*, *vgrGb*, *vgrGc* and *vgrGd*	Santos et al., 2020	EML4746
Δ2*EIbd*	Deletion of *vgrGb*- and *vgrGd*-associated effector-immunity pair	This study	EML5531
Δ3EIbcd #1	Deletion of *vgrGb*-, *vgrGc*- and *vgrGd*-associated effector - immunity pair using Δ2EIbc as background.	This study	EML5594

Δ3EIbcd #2	Deletion of *vgrGb*-, *vgrGc*- and *vgrGd*-associated effector - immunity pair using Δ2EIcd as background.	This study	EML5920
Δ3*EIabd*	Deletion of *vgrGa*-, *vgrGb*- and *vgrGd*-associated effector - immunity pair	Santos et al., 2020	EML5515
Δ4*EIabcd*	Deletion of *vgrGa*-, *vgrGb*- and *vgrGd*-associated effector - immunity pair and whole *vgrGc* cluster (Δ*vgrGc-v3*)	Santos et al., 2020	EML5540
Plasmids			
pJQ200KS	Gm^R, suicide plasmid containing GmR and sacB gene for selection of double crossover	Quandt and Hynes, 1993	EML121
pRL662	Gm^R, broad host range vector derived from pBBR1MCS-2	Vergunst et al., 2000	EML315
pTrc200	Sp^R, pVS1 origin, *lacIq, trc* promoter expression vector	Schmidt-Eisenlohr et al., 1999	EML904
pJN105	Gm^R, arabinose-inducible gene expression vector derived from pBBRMCS-1, *araC*-P$_{BAD}$	Newman and Fuqua, 1999	EML4220
pET28a	Km^R, Protein expression vector, N-terminal His-tagged	Lab stock	EML2485

pJN-*v2a*	Gm^R, pJN expressing *v2a*	Santos et al., 2020	EML5272
pJN-*v4b*^FL	Gm^R, pJN expressing the full-length *v4b*	This study	EML4758
pJN-*v4b*^CT	Gm^R, pJN expressing the C-terminal toxin domain of *v4b*	This study	EML4756
pJN-*v2c*	Gm^R, pJN expressing *v2c*	This study	EML5328
pTrc-*v5b*	Sp^R, pTrc200 expressing the immunity protein V5b	This study	EML4759
pTrc-*v3c*	Sp^R, pTrc200 expressing the immunity protein V3c	This study	EML5337
pJN-*v2a* (H385A)	Gm^R, pJN expressing *v2a* with amino acid substitution (H385A)	Santos et al., 2020	EML5298
pJN-*v2c* (H383A)	Gm^R, pJN expressing *v2c* with amino acid substitution (H383A)	This study	EML5925
pJN-*v2c* (H384A)	Gm^R, pJN expressing *v2c* with amino acid substitution (H384A)	This study	EML5926
pJN-*v2c* (H408A)	Gm^R, pJN expressing *v2c* with amino acid substitution (H408A)	This study	EML5927
pJN-*v2c* (HAHA)	Gm^R, pJN expressing *v2c* with amino acid substitution (H383AH384A)	This study	EML5922
pJN-*v2c* CR	Gm^R, pJN expressing *v2c* with deletion of the conserve region of Tox-SSH domain	This study	EML5589
pJQ200sk-*vag4vag5b*	Gm^R, plasmid to generate Δ3*EIbcd* #2	This study	EML5307

pJQ200sk-*vag2vag3d*	Gm^R, plasmid to generate Δ2*EIbd*, Δ3*EIbcd* #1 and Δ3*EIabd*	This study	EML5340
pRL662-GFP(S65T)	GFP from pRU1156 cloned into pRL662	Lab stock	EML3376
pET28a-*v2c*	pET28a expressing His-tagged V2c	This study	EML6050
pET28a-*v2c*HAHA	pET28a expressing His-tagged V2c H383AH384A	This study	EML6051
pET28a-*v2c*H408A	pET28a expressing His-tagged V2cH408A	This study	EML 6063
pET28a-*v2c*H383A	pET28a expressing His-tagged V2cH383A	This study	EML 6064
pET28a-*v2c*H384A	pET28a expressing His-tagged V2cH384A	This study	EML 6065

Table S2. Primers used in this study

Primer Name	Sequences	Purpose
V2c(H383A)	5'-CCG<u>GAATTC</u>AAGGAACAGCTATGAGCATCCCCCGCGAC-3' 5'- GACAATTTAGATAGCGCGCATACTCCTGCTAACTCT-3' 5'- AGAGTTAGCAGGAGTATGCGCGCTATCTAAATTGTC-3' 5'- CG<u>TCTAGA</u>TCACTTTATGATTCCTTCGCG-3'	mutagenesis
V2c(H384A)	5'-CCG<u>GAATTC</u>AAGGAACAGCTATGAGCATCCCCCGCGAC-3' 5'-GACAATTTAGATAGCCATGCGACTCCTGCTAACTCT-3' 5'-AGAGTTAGCAGGAGTCGCATGGCTATCTAAATTGTC-3' 5'- CG<u>TCTAGA</u>TCACTTTATGATTCCTTCGCG-3'	mutagenesis
V2c(H408A)	5'-CCG<u>GAATTC</u>AAGGAACAGCTATGAGCATCCCCCGCGAC-3' 5'- ATGCTTCCAGAAGACGCGCGAGCTACACTCTCATAT -3' 5'-TGAGAGTGTAGCTCGCGCGTCTTCTGGAAGCAT-3' 5'- CG<u>TCTAGA</u>TCACTTTATGATTCCTTCGCG-3'	mutagenesis
V2c(CR)	5'-GTTTTTTTGGGCTAGCGAAGGAACAGCTATGAGCATC-3' 5'-CCTTTCTTCCATATGAGAGTGTAGCATTGTCGCCAAGTGGGAGGGCCGTC-3' 5'-GACGGCCCTCCCACTTGGCGACAATGCTACACTCTCATATGGAAGAAAGG -3' 5'-GCGGTGGCGGCCGCTTCACTTTATGATTCCTTCGC-3'	mutagenesis
V2c(H383A H384A)	5'-CCG<u>GAATTC</u>AAGGAACAGCTATGAGCATCCCCCGCGAC-3' 5'-GACAATTTAGATAGCCATGCGACTCCTGCTAACTCT-3' 5'-AGAGTTAGCAGGAGTCGCATGGCTATCTAAATTGTC-3' 5'- CG<u>TCTAGA</u>TCACTTTATGATTCCTTCGCG-3'	mutagenesis
pJQ200KS-*v4bv5b*	5'-TT<u>TCTAGA</u>CATCTGTTGTTCGCGGCCGATG-3' 5'-GC<u>GGATCC</u>ATCACCCGCCATGTGAGGCCT-3' 5'-GC<u>GGATCC</u>TTCGCTGCCCTGTGACCGTTCT-3' 5'-GC<u>CCCGGGG</u>CAAGATGATTCCGGGAATTGAC-3'	deletion mutant
pJQ200KS-*v2dv3d*	5'-GC<u>TCTAGA</u>GAGCGCCGAAATCCGCTT-3' 5'-CG<u>GGATCC</u>GATGCTCATGCTGTCTCCCTCTC-3' 5'-CG<u>GGATCC</u>GCCAGAAAGCCATAGATTGTCG-3' 5'-TCCC<u>CCCGGGG</u>CATCACCACACTCTCGAGCCT-3'	deletion mutant

V4b	5'-CCGTCTAGAAAGGAACAGCTATGGCGGGTGAT-3' 5'-CGGAGCTCGTTTCGATACTCAGCATCGACCG-3'	overexpression
V5b	5'-TATAGGTACCCGAGAATAAGAACCGCGTTC-3' 5'-GCTCTAGAATTGGCGAGGCTACAGCG-3'	overexpression
V2c	5'-CCGGAATTCAAGGAACAGCTATGAGCATCCCCCGCGACAATTAT-3' 5'-CGTCTAGATCACTTTATGATTCCTTCGCGTTTAAGGC-3'	overexpression
V3c	5'-ATCCGAGCTCGTGACAGACAAGTCGGGAGCG-3' 5'-GCTCTAGACCTAGATCCATCGCTTACTGCCTC-3'	overexpression

pET28a-v2c	5'-AATCTAGAATGAGCATCCCCCGCGACAA-3'. 5'-AACTCGAGCTTTATGATTCCTTCGC-3'	Protein expression
pET28a-v2cHAHA	5'-GACAATTTAGATAGCGCGGCGACTCCTGCTAACTCT-3'. 5'-AGAGTTAGCAGGAGTCGCCGCGCTATCTAAATTGTC-3'	Protein expression
pET28a-v2c-H383A	5'-GACAATTTAGATAGCGCGCATACTCCTGCTAACTCT-3' 5'-AGAGTTAGCAGGAGTATGCGCGCTATCTAAATTGTC-3'	Protein expression
pET28a-v2c-H384A	5'-GACAATTTAGATAGCCATGCGACTCCTGCTAACTCT-3' 5'-AGAGTTAGCAGGAGTCGCATGGCTATCTAAATTGTC-3'	Protein expression
pET28a-v2c-H408A	5'-ATGCTTCCAGAAGACGCGCGAGCTACACTCTCATAT-3' 5'-TGAGAGTGTAGCTCGCGCGTCTTCTGGAAGCAT-3'	Protein expression

Restriction enzyme sites are underlined. UF (upstream forward), UR (upstream reverse), DF (downstream forward) and DR (downstream reverse).