Two classes of His-Me finger superfamily nuclease effectors exerting synergy for

Agrobacterium T6SS-mediated interbacterial competition

Mary Nia M. Santos[a1], Katherine L. Pintor[a2], Pei-Yu Hsieh[b3], Yee-Wai Cheung[a], Li-

Kang Sung[a], Yu-Ling Shih[b] and Erh-Min Lai[a*]

[a] Institute of Plant and Microbial Biology, Academia Sinica, Taipei 115201 Taiwan

[b] Institute of Biological Chemistry, Academia Sinica, Taipei 115201 Taiwan

Running title: His-Me finger T6SS nuclease toxins

*Corresponding author

Erh-Min Lai: emlai@gate.sinica.edu.tw

Institute of Plant and Microbial Biology, Academia Sinica

128 Sec. 2, Academia Road, Nankang, Taipei, Taiwan 115201

Current address:

[1]Aquaculture Research and Development Division, Department of Agriculture -

National Fisheries Research and Development Institute (DA-NFRDI), Manila,

Philippines

[2]Department of Ecophysiology, Max Planck Institute for Terrestrial Microbiology,

Karl-von-Frisch-Str. 10, 35043 Marburg, Germany

[3]Department of Pharmacology, University of Cambridge, United Kingdom

ORCID

MNMS https://orcid.org/0000-0001-6544-0339

KLP https://orcid.org/0000-0001-9137-4726

PYH https://orcid.org/0000-0002-7263-1691

YWC https://orcid.org/0000-0002-1421-4419

LKS https://orcid.org/0000-0002-7078-8003

YLS https://orcid.org/0000-0002-8232-2569

EML https://orcid.org/0000-0003-3630-8683

ABSTRACT

Type VI secretion system (T6SS) assembles into a contractile nanomachine to inject effectors across bacterial membranes for secretion. *Agrobacterium tumefaciens* species complex is a group of soil inhabitants and phytopathogens that deploys T6SS as an antibacterial weapon against bacterial competitors at both inter-species and intra-species levels. *A. tumefaciens* strain 1D1609 genome encodes four effector genes, in which all four are specialized effector harboring a conserved N-terminal PAAR-like DUF4150 domain but distinct C-terminal effector domains. Previous study reported that V2a is a His-Me finger nuclease toxin contributing to DNase-mediated antibacterial activity. However, it remains unknown about the functions and roles of other three effectors. In this study, we identified V2c is another His-Me finger nuclease encoded by 1D1609 genome but with distinct SHH motif differed from AHH motif of V2a. We demonstrated that the ectopic expression of V2c caused growth inhibition, plasmid DNA degradation, and cell elongation in *Escherichia coli*. The cognate immunity protein, V3c, neutralizes the DNase activity and rescues phenotypes of the growth inhibition and cell elongation. Ectopic expression of V2c DNase-inactive variants retain the cell elongation phenotype while V2a induced cell elongation in a DNase-mediated manner. We also showed that the amino acids of conserved SHH and HNH motifs are responsible for the V2c DNase activity *in vivo* and *in vitro*. Notably, V2c also mediated the DNA degradation and cell elongation of target cell in the context of interbacterial competition. Importantly, V2a and V2c function synergistically to exert stronger antibacterial activity against the soft rot phytopathogen, *Dickeya dadantii*.

INTRODUCTION

The type VI secretion system (T6SS) is a nanomachine used by many gram-negative bacteria for antagonism or pathogenesis by injecting toxins into target bacterial or eukaryotic cells. T6SS is composed of a membrane complex in connection to a baseplate, which is the docking site for polymerization of a tube surrounded by a contractile sheath (1). The tube is stacked of Hcp protein hexamer tipped with the trimeric VgrG puncturing device and a PAAR repeat-containing protein which sharpens the tip (2). When the sheath contracts, the tube is propelled towards the target cell to deliver effectors. After firing, the sheath disassemble and the subunits of the sheath are recycled to build another machine (3).

In terms of delivery, the T6SS effectors can be classified as cargo or specialized

effectors. The cargo effectors interact non-covalently with Hcp, VgrG or PAAR while

in specialized effectors, the effector domain is covalently fused to Hcp, VgrG or PAAR

(4). Functionally, most of the identified effectors mediate antibacterial activities

including nuclease, peptidoglycan hydrolase, lipase, phospholipase, NAD(P)+-

glycohydrolase and ADP ribosyltransferase by targeting conserved cellular structures

such as nucleic acids, peptidoglycan, and inner membrane as well as specific

metabolites or proteins (5).

Agrobacterium tumefaciens is a Gram-negative alphaproteobacteria belonging to the

Rhizobiaceae family. It is an economically important pathogen that causes crown gall

disease and a gene delivery tool with capability of transforming plant cells and fungi

(6). T6SS main gene cluster is highly conserved in the genome of *A. tumefaciens*

genomospecies complex (7, 8). Besides the main cluster consisting of two divergent

operons, *imp* (<u>imp</u>aired in nitrogen fixation) encoding core structural and regulatory

components and *hcp* operon for effectors and effector delivery components (e.g. *hcp,*

vgrG, paar), most *A. tumefaciens* strains also encode one to multiple *vgrG* operons

encoding effector-immunity (EI) gene pairs. *A. tumefaciens* strain 1D1609 encodes

largest number of *vgrG* (*vgrGa, vgrGb, vgrGc, vgrGd*) in *A. tumefaciens*

genomospecies complex with each module associated with an EI pair (9). Previous

study by deletion of single or multiple *vgrG*-associated effector-immunity (EI) pairs

suggested that *vgrGa*-associated effector V2a appears to be the major antibacterial

toxin but other effectors also contribute to full antibacterial activity for 1D1609 (9). V2a

is a PAAR (DUF4150)-linked specialized DNase effector harboring C-terminal Tox-

AHH domain. It is a member of the HNH/ENDO VII nuclease superfamily, which also

named as His-Me finger endonuclease superfamily (9, 12, 13). This study also

indicated there are three more putative EI pair encoded in the other three orphan *vgrG*

modules. However, the biochemical and biological functions of other three EI pairs

remain uncharacterized.

In this study, we discovered that *vgrGc*-associated effector V2c is another DNase toxin

and explored the rationale of having two T6SS DNase toxins in *A. tumefaciens* strain

1D1609. We show that V2c harbors a DNase toxin domain with a conserved SHH

motif representing a newly characterized T6SS bacterial toxin. SHH is one of the five

new clades identified in the His-Me finger nuclease superfamily (10). His-Me finger

nucleases are characterized by their strictly conserved histidine (His) (1st H in the HNH

motif) and a metal ion (Me) bound to conserved His (last H in the HNH motif) being

essential for nucleic acid hydrolysis (11). Here, we show that the expression of V2c

caused growth inhibition, plasmid DNA degradation and cell elongation, which the

phenotypes are neutralized by co-expression of the cognate immunity protein, V3c.

We also show that the SHH motif is responsible for the nuclease activity of V2c.

Importantly, V2a AHH nuclease and V2c SHH nuclease function synergistically to

exert stronger antibacterial activity against the soft rot phytopathogen, *Dickeya*

dadantii. V2c orthologs are also conserved in Rhizobiaceae family, suggesting the

importance of His-Me finger T6SS nuclease in this family for interbacterial competition.

RESULTS

V2c is a DNase belonging to the Tox-SHH clade of His-Me finger superfamily

Of the four identified effector-immunity (EI) pairs encoded on T6SS main and auxiliary

clusters in *A. tumefaciens* 1D1609, *v2c* (At1D1609_43220) resides in the auxiliary

cluster, upstream of its putative cognate immunity protein, *v3c* (At1D1609_43230) (9).

V2c is a PAAR (DUF4150)-linked specialized effector (**Figure 1A**) that similar to V2a,

the homolog encoded on the T6SS main cluster. Despite both V2a and V2c harboring

N-terminal PAAR domain (**Figure S4**), V2c does not show any sequence similarity to

V2a as well as other known T6SS effectors by BLAST and CDD searches. Multiple

sequence alignment of V2c and V2c orthologs revealed V2c containing conserved

SHH and HNH motifs at its C-terminal region (**Figure 1B**), in which these V2c

orthologs all harbor C-terminal Tox-SHH domain but their N-terminal region can be

fused to other known T6SS domains such as Found in type sIX effector (FIX) and FIX

RhsA AHH HNH-like (14, **Figure 1C).** Using the protein homology detection program

HHpred (15), we found that the C-terminal region of V2c encodes two parallel β-

strands connected by α-helix, which is signature structure in His-Me finger nuclease

(12). Consistent with the result of secondary structure prediction, structure prediction

using AlphaFold (**Figure 1D**) shows this domain consists of two antiparallel β-strands

connected with an Ω loop with a His (H384) at the C-terminus of the first β-strand and

followed by an α-helix, featuring a HNH motif of His-Me finger endonuclease (10, 11).

These suggest V2c belonging to the Tox-SHH (Pfam PH15652) clade of His-Me finger

superfamily.

To demonstrate the antibacterial and DNase activities of V2c, we expressed *v2c* using

an arabinose inducible promoter to determine if V2c is a bacterial toxin functioning as

a nuclease. Growth inhibition measured by optical density was observed when *v2c* is

expressed by arabinose induction in *Escherichia coli* and this can be neutralized by

co-expression of its putative cognate immunity *v3c* as compared to the vector control

(**Figure 2A**). Also, colony forming unit (CFU) count shows the reduced CFU of *v2c*-

expressed cells and higher CFU could be recovered with co-expression of *v3c* (**Figure**

2B)*.* In addition, plasmid degradation assay showed that the expression of *v2c* upon

arabinose induction resulted in degradation of plasmid DNA in *E. coli* and this is

abolished when *v3c* is co-expressed (**Figure 2C**). The data indicated that V2c exhibits

DNase activity *in vivo*.

We further investigated whether the SHH and HNH motif are involved in DNase activity.

His-Me finger superfamily has a strictly conserved His residue and catalytic metal ion

which are both essential for nucleic acid hydrolysis. Based on this premise, we define

a region that contains His residues as the putative catalytic site region as a His-Me

finger domain core region (CR), corresponding to the region spanning amino acid

residues 380-409. Based on the alignment, H384 is the catalytic His and H408 is for

metal ion binding while H383 may function as second residue for metal ion

coordination. To verify their roles in the His-Me finger domain, a series of deletion as

well as single and double amino acid substitution variants were generated. Strikingly,

no growth inhibition and plasmid degradation were observed in any of the mutants

expressing the CR deletion (V2c^{CR}), single (V2c^{H383A}, V2c^{H384A} and V2c^{H408A}) and

double (V2c^{HAHA,} substitution in both H383 and H384) variants (**Figure 2A, C, D**).

Higher CFU was also recovered from *E. coli* culture expressing these mutants alone

or with V3c (**Figure 2B**). To further confirm the DNase activity of V2c*,* each of V2c and

its variants is fused with 6x-His and purified for *in vitro* DNase activity assay. The

results showed that purified V2c fraction is able to degrade DNA, but not the eluate

from *E. coli* expressing the vector (pET28a) (**Figure 2D**). Consistent with the *in vivo*

plasmid degradation assay, no activity could be detected from V2c^{H384A}*,* V2c^{H408A}, or

V2c^{HAHA} variants. However, purified V2c^{H383A} fraction exhibited DNase activity by

degrading the plasmid DNA substrate. Western blot analysis of the reaction samples

shows V2c and its variants used for the assay were at similar protein levels. Taken the

results of *in vivo* and *in vitro* DNA degradation assays, the conserved H384 and H408

in the HNH motif are crucial for V2c DNase activity whereas H383 in SHH motif

appears to be less critical for *in vitro* but may be required for *in vivo* nuclease activity.

Ectopic expression of V2c caused DNA degradation and cell elongation

We next examined the cell morphology and DNA integrity of *E. coli* cells expressing

V2c and the two variants (V2c^{CR} and V2c^{H408A}) that are defective in DNase activity and

toxicity. Cell morphology was observed by phase contrast while bacterial cell

membrane and DNA were stained by FM4-64 and Hoechst, respectively under

fluorescence microscope. Hoechst signals are significantly reduced when V2c is

expressed by arabinose induction in *E. coli* and this can be rescued by co-expression

of its cognate immunity V3c shown in the images and quantification (**Figure 3A, B**).

Hoechst signals are diminished only in *E. coli* cells producing *v2c* but not *v2c*^{H383A}

v2c^{H384A}, *v2c*^{CR} and vector control, suggesting that the cellular DNA degradation is

caused by V2c DNase activity, which can be neutralized by its immunity protein V3c.

In addition, we also observed cell elongation in *E. coli* expressing *v2c* (**Figure 3A, C**).

Notably, co-expression of V3c rescues the cell elongation phenotype. However, *E. coli*

expressing *v2c*^{CR} and *v2c*^{H408A} does not abolish the elongated cell phenotype (**Figure**

3A, C).

Ectopic expression of V2a also caused cell elongation phenotype

Previous study showed that V2a is a Tox-AHH nuclease that serves as the major

antibacterial weapon in 1D1609 (9). H385 and H386 are two of the predicted His

residues in AHH motif of V2a, corresponding to H383 and H384 in SHH motif of V2c.

The structure predicted by AlphaFold indicates V2a also featuring a His-Me finger, in

which two α-helices with histidine residues (H430, H456) followed the β-strands

(**Figure 4A**). And, our previous findings demonstrated that expression of V2a but not

V2a^{H385A} caused growth inhibition and plasmid degradation in *E. coli* (9)*.* To gather

further insight of V2a AHH motif in the His-Me finger domain, we evaluated the

growth and morphology of arabinose induced V2a mutants. Results showed that

expression of V2a caused reduction in Hoechst-stained DNA signals and resulted in

cell elongation (**Figure 4B, C, D**). The expression of V2a^{H385A} has similar levels of

Hoechst-stained DNA signals and cell length as compared to the vector control.

These results suggest that the cell elongation phenotype observed were caused by

V2a DNase activity, whereas V2c shows DNase activity-independent cell elongation.

We also determined the toxicity of another putative effector V4b of 1D1609, a PAAR

(DUF4150)-linked specialized effector but harboring a Rhs domain with structural

similarity to the insecticidal toxin (9) (**Figure S1A**). Expression of *v4b* caused growth

inhibition in *E. coli,* in which the growth inhibition can be rescued by co-expression of

its putative cognate immunity *v5b* to a level similar to vectors control (**Figure S1B**).

However, no DNA degradation and cell elongation could be observed (**Figure S1C,**

D, E, F). These results suggest that cell elongation phenotype may be specific to

toxicity of nuclease effectors and V4b may also function as a bacterial toxin.

V2c and V2a nucleases function synergistically for antibacterial activity against

the soft rot phytopathogen, *Dickeya dadantii*

To correlate the DNA degradation and cell elongation phenotypes observed by ectopic

expression in *E. coli* in more biologically relevant context, we further performed

interbacterial competition assay using the phytopathogen, *Dickeya dadantii* 3937 as

the prey cells. Expression of TssB with a C-terminal fusion to green fluorescence

protein (TssB-GFP) has been shown to assemble into a functional TssBC sheath in *A.*

tumefaciens strain C58 (16). Here, we expressed 1D1609 TssB with a C-terminal

fusion to super-folder GFP (TssB-sfGFP) endogenously in both 1D1609 wild type

(WT::TssB-sfGFP) and a non-functional T6SS mutant lacking all four *vgrG* genes

(Δ*4*::TssB-sfGFP) (9). While a few GFP foci were also detected in Δ*4*::TssB-sfGFP,

the TssBC sheath structures and dynamics could only be observed in WT::TssB-

sfGFP cells (**Figure S2A**, **Supplementary video 1**), confirming TssBC sheath is not

formed in the mutant lacking all four *vgrG* genes. This is consistent with the lack of

Hcp secretion in Δ*4*::TssB-sfGFP although TssB-sfGFP fusion proteins are at similar

abundant in all strains (**Figure S2B**). Accordingly, some of *D. dadantii* prey cells (either

WT or Δ*imp,* a non-functional T6SS mutant) became elongated and were lysed after

competition with WT::TssB-sfGFP but not Δ*4*::TssB-sfGFP (**Figure 5A**).

Next, we constructed mutants lacking one or several EI pairs and used them to assess

the activity against *D. dadantii* Δ*imp* expressing GFP (*D. dadantii* Δ*imp*-GFP(S65T))

by microscopy and quantified the number of viable cells (log$_{10}$ CFU/mL) after co-

culture with *D. dadantii* Δ*imp*. The use of *D. dadantii* 3937 Δ*imp* mutant is to avoid

potential counterattack by *D. dadantii* T6SS (16). We observed very few cells

expressing GFP when competing with WT or Δ*3EIbcd* expressing V2a effector only*.*

The lysed and elongated *D. dadantii* cells were only detected when co-cultured with

1D1609 with either strains expressing V2a only (Δ*3EIbcd*), V2c only (Δ*3EIabc*), or both

Δ*3EIbd* (**Figure 5B**). No lysed and elongated cells were observed in Δ*4EI* and *D.*

dadantii only.

By counting CFU of prey cells after Interbacterial competition, the results also show

that V2a only (Δ*3EIbcd*) exhibited similar antibacterial activity as the WT against *D.*

dadantii (**Figure 5C**). In addition, 1D1609 expressing V2c only (Δ*3EIabd*) showed

weak but detectable antibacterial activity as compared to Δ*4EI,* indicating V2c itself is

able to detoxify *D. dadantii* but with weaker activity than V2a. Strikingly, 1D1609

expressing V2a and V2c only (Δ*2EIbd*) could result in stronger killing effects than WT.

We further quantified the cell length under the fluorescence microscope after

competition assay and results showed that *D. dadantii* cells were elongated after 18

hours of co-culture with either *v2a* and *v2c* effectors, or combination of both, as

compared to those *D. dadantii* cells alone or co-incubated with Δ*4EI* (**Figure 5D**).

Taken together, these results suggest that 1D1609 uses the two V2a and V2c DNase

effectors with synergistic antibacterial activity against *D. dadantii* resulting to elongated

and lysed cells.

DISCUSSION

Nucleases are considered as one of the most potent categories of T6SS effectors

because they poison the targets from the most essential in the origin of life, nucleic

acid. Although His-Me finger nucleases are a large and diverse superfamily of

nucleases, however, only a limited number of T6SS effectors have been identified as

His-Me finger nuclease. In the present study, we characterized a T6SS effector in *A.*

tumefaciens strain 1D1609, V2c, by its DNase activity. Results shows V2c caused

DNA degradation both *in vivo* and *in vitro* (**Figure 2C, 2D**) that confirmed V2c is a

DNase.　Moreover, our previous findings (9) suggest V2a and V2c that consisting of

Tox-AHH and Tox-SHH motifs respectively, belong to the His-Me finger superfamily.

The predicted structures (**Figure 1D, 4A**) show that both V2a and V2c contain the

signature structure of His-Me finger domain harboring the catalytic His located at the

C-terminus of their β1 strands (H384 for V2c and H386), which is predicted to be the

general base for hydrolysis of phosphodiester bonds, followed by α-helixes. With our

previous findings on V2a, these suggest both V2a and V2c are His-Me finger

nucleases.

Given that His-Me finger nucleases are one-metal-ion-dependent nucleases, primarily

dependent on Mg^{2+} or in some cases dependent on Zn^{2+}. H408 of V2c and H430 of

V2a are determined as the His residue for direct binding to metal ion whereas another

acidic or polar amino acid may also coordinate the metal ion, located before the

invariant His (first H in SHH or AHH motif), which could either be Asn/Asp/Glu or His

which preferentially bind to Mg^{2+} or Zn^{2+}, respectively (9). The latter is the case for

both V2a (AHH motif) and V2c (SHH motif). The two His (H383, H384) residues of V2c

SHH motif are both required for exerting DNase activity *in vivo,* consistent with the

requirement of H385 and H386 of V2a AHH motif for DNase activity (8). Different from

the loss of DNase activity *in vivo*, V2c[H383A] variant retains the DNase activity by an *in*

vitro nuclease activity assay (**Figure 2C, 2D**). This indicates the requirement of H383

in coordinating metal ion binding for DNase activity may be subject to environmental

factors. His-Me fingers approach DNA from the minor groove and clamp the phosphate

backbone via the α-helix and β1 strand. Apart from having similar DNA-binding modes,

His-Me finger nucleases also share a conserved catalytic mechanism for hydrolyzing

nucleic acids. They all cleave P-O3' bonds to generate cleaved products containing

5'-phosphate and 3'-OH ends (22). If both V2a and V2c have the same mechanism to

bind and hydrolyze DNA, then the other domains outside the His-Me finger fold may

explain the difference in the phenotype.

His-Me finger encodes a small structural motif, so it is usually associated with other

domain architectures that could play additional functions. In addition, its three-element

structural motif is unable to provide enough of a hydrophobic core for stability (10).

Thus, many proteins containing the His-Me finger employ a variety of additional

structural elements or domains for stabilization and/or specificity. As V2c and its

orthologs commonly found in Rhizobiaceae are not included in the 77 sequences with

conserved core regions of His-Me finger superfamily reported by Jablonska et al (10),

suggesting they are SHH motif variants belonging to SHH clade of His-Me finger

nuclease. Our evidence also suggests that V2c may represent a new branch of Tox-

SHH toxin family (Pfam PF15652), which includes proteins with additional domain

architectures (**Figure 1**). This includes PAAR-domain containing protein, <u>F</u>ound in

type s<u>IX</u> effector (FIX) superfamily, and FIX RhsA AHH-HNH-like. Most of these are

conserved regions found in many bacterial toxin proteins. Moreover, these additional

domain(s) may facilitate the secretion of nuclease and affect its toxicity to the

competing cell. While both V2c and V2a harbor the conserved N-terminal DUF4150

PAAR-like domain, they share low sequence similarity in the C-terminal region

including the AHH or SHH motifs containing His-Me finger domain that may contribute

to the difference on toxicity strength and the role of DNase on cell elongation (**Figure**

S4). The protein structures predicted by AlphaFold reveal the structural difference

between V2c and V2a (**Figure 1C, S1B**). Moreover, His-Me finger motif is nearly a

stand-alone domain and separate from the DNA sequence-specific binding domains

(22, 23). Collectively, we discovered *A. tumefaciens* strain 1D1609 is equipped with

two T6SS nuclease effectors, V2a and V2c, that belong to His-Me finger superfamily.

The findings expand the variety of T6SS His-Me finger nucleases on top of those have

been functionally characterized (9, 17, 18, 19).

Cell elongation caused by T6SS-mediated nuclease effector have been reported (17,

24). Besides nuclease effectors, cell division arrest is also caused by non-nuclease

effectors. For example, a T6SS ADP-ribosyltransferase effector called Tre1 in *Serratia*

proteamaculans was reported to inactivate bacterial cell division Tre1 by modifying

arginine on FtsZ to block polymerization (25). We also observed the elongation

phenotype in ectopic expression of *v2c* in *E. coli* and in *D. dadantii* during interbacterial

competition. The proposed biological role of bacterial filamentation is mainly related to

stress response to thrive for survival (26). One of the conditions that lead to

filamentation is DNA damage. When the DNA is damaged, the SOS response is

induced and cells may delay the cell cycle and inhibit cell division until DNA repair and

replication are complete. In this study, while both V2a and V2c cause DNA degradation,

the mechanism of induced cell elongation seems to be different. For V2a*,* the cell

elongation phenotype is unobservable in DNase inactive mutant by disrupting H385,

the metal ion coordinating residue (**Figure 4**). This contrasts with V2c, wherein the

DNase activity is independent of cell elongation phenotype. Cell elongation could still

be observed when different DNase defective V2c mutant proteins were expressed.

(**Figure 3, Figure S3A**). In addition, co-expressing of V3c can restore V2c-induced

cell elongation back to normal cell length, but not to all the catalytic variants (**Figure**

3C, S3B). The data suggest that another domain outside the SHH catalytic region in

V2c is responsible for the cell elongation phenotype. Different from the nuclease

activity-dependent filamentation observed previously (17, 24) and V2a, V2c-induced

cell elongation independent of nuclease activity may represent a new mechanism of

stress response that yet to be elucidated. Future work by domain dissection is needed

to identify the domain that causing cell elongation.

To date, a few T6SS effectors were reported to contain SHH motif, including Tke4 of

Pseudomonas putida (20) and Txe4 of fish pathogen *Pseudomonas plecoglossicida*

strain XSDHY-P that exerting DNase-mediated toxicity when expressed in *E. coli* (21).

However, it remains unclear whether SHH and HNH motifs of these T6SS His-Me

finger nucleases are directly involved in interbacterial competition. Here, we

demonstrate the DNase activities related to both SHH and HNH motifs are critical

during interbacterial competition against the soft rot phytopathogen *D. dadantii*. We

also discovered V2a and V2c exert synergistic antibacterial activity against *D. dadantii*.

The strain 1D1609 is unique in *A. tumefaciens* species complex with multiple VgrG

spikes carrying different antibacterial effectors (9). Several studies showed that some

T6SS toxins can only exert toxicity in specific bacterial species (27, 28). For 1D1609,

V2c-mediated antibacterial activity was observed when against *D. dadantii* was acted

as a prey, whereas it was not observed when *E. coli* was used as a prey (9). On the

other hand, V2a appears to be the major antibacterial weapon for interbacterial

competition against either *E. coli* or *D. dadantii*. From the observation of the synergism

of V2a and V2c against *D. dadantii,* harboring multiple T6SS effectors in 1D1609 is an

advantage for the strain to maintain a growth competitiveness in different niches and

environments.

In conclusion, in the present study we identified V2c is SHH motif bearing His-Me

finger nuclease, which its DNase activity can be neutralize by its cognate immunity

protein, V3c. We also found that V2c mediated the DNA degradation and cell

elongation of target cell independently.   Importantly, V2a and V2c function

synergistically to exert stronger antibacterial activity against the soft rot phytopathogen,

D. dadantii 3937.

MATERIALS AND METHODS

Bacterial strains and growth conditions

Bacterial strains and plasmids are listed in Table S1. *Escherichia coli* strains were

grown in LB medium at 37°C, *A. tumefaciens* strains in 523 medium at 25°C and *D.*

dadantii 3937 in LB medium at 30°C. Antibiotics were added when necessary: for *E.*

coli, 30 µg/mL gentamicin and 250 µg/mL streptomycin; for *D. dadantii* 50 µg/mL

gentamicin.

Molecular cloning

Plasmid pJN105 was used for expression of toxin genes and pTrc200 plasmid was

used for expression of immunity gene. PCR was amplified with KAPA HiFi HotStart

DNA Polymerase (Roche) using genomic DNA of *A. tumefaciens* 1D1609. The C-

terminal fusion of *tssB* with superfolder GFP (sfGFP) was constructed in pJQ200ks to

replace the endogenous *tssB* in WT and Δ*4*Δ*tssB* background. The primers used in

the study are listed in Table S2. Site directed mutagenesis was done using either DpnI

or Gibson Assembly (New England BioLabs), ligated using T4 ligase (Takara) and

transformed into chemically competent *E. coli* DH10B. Plasmid DNA was isolated

using QIAprep miniprep kit (Qiagen). All constructs were confirmed by sequencing.

The in-frame deletion mutants in *A. tumefaciens* were generated using a suicide

plasmid via double crossing over by electroporation or by conjugation as described (9).

Bioinformatic analysis

The orthologs of V2c were analyzed using amino acid sequences after the DUF4150

domain via BLASTP search. Multiple sequence alignment was performed using

MUSCLE or Multiple Sequence Comparison by Log- Expectation (29). Full length

sequences of the V2c orthologs were analyzed to obtain domain architectures from

Pfam database (30). The E-value threshold was set at 10^{-5}. HHpred and Swiss

Model were used for homology-based protein structure prediction while the

availability of protein structure predicted by AlphaFold was from the UniProt

database. Secondary structure prediction was done using default settings in

PSIPRED 4.0 Protein Structure Prediction Server/ DMPfold 1.0 Fast Mode.

Growth inhibition assay

Growth inhibition assay was performed as described previously (31). In brief, overnight

cultures of *E. coli* DH10B strain with vectors were adjusted to OD_{600}. 1 mM isopropyl

β-D-1- thiogalactopyranoside (IPTG, Amresco) was used to induce the expression of

the putative immunity protein. L-Arabinose (Ara, Sigma-Aldrich) was added to 0.2%

final concentration, to induce expression of the toxin. Cell growth was recorded every

hour at OD_{600}. Empty vectors were used as controls. The number of colony forming

units (log_{10} CFU/mL) was quantified in each sample grown on agar plates.

Plasmid DNA degradation analysis in *E. coli* cells

In vivo plasmid DNA degradation analysis was performed as described previously (31).

In brief, overnight cultures of *E. coli* DH10B strain with empty vectors or expressing

the effectors and mutants were harvested and adjusted to OD_{600} ~0.3. L-Arabinose

(0.2%, final concentration) was added to induce toxin expression and cultured for 2 h.

Equal amounts of cells were used for plasmid DNA extraction and equal volume of

extracted DNA was resolved in agarose gel followed by ethidium bromide staining and

visualized using Gel Doc XR+ UV Gel Documentation Molecular Imager Universal

Hood II (Bio-Rad).

Cloning, expression and purification of V2c and mutants

The effector gene *v2c* was constructed in pET28a plasmid and immunity gene *v3c*

was constructed in pTrc200 plasmid. Site directed mutagenesis were done by using

overlap extension PCR. Plasmid for expressing effector protein, V2c WT or its variants,

and plasmid for expressing immunity protein, V3c, were transformed into *E. coli*

BL21(DE3) for IPTG-induced co-expression. Bacterial culture was incubated at 37 °C

until OD_{600} reached 0.5, then 0.1 mM IPTG was added followed by 5-h expression at

37 °C. Pellets were collected, sonicated, and expressed proteins were purified by

using Ni Sepharose 6 Fast Flow histidine-tagged protein purification resin (Cytiva).

Briefly, cells were lysed in extraction buffer (50 mM Tris-Cl, pH7.5, 0.1 M NaCl, 20 mM

imidazole) and proteins were eluted by using elution buffer with up to 250 mM

imidazole in 50 mM Tris-Cl, pH7.5, 0.1 M NaCl.

In vitro **DNase activity assay**

A 10 µL reaction containing 200 ng of pTrc200HA plasmid DNA and 1× CutSmart

Buffer (NEB) with partially purified V2c or its variants normalized by the major V2c

protein band was incubated at 37°C for 1 hour. The digestion of DNA was

determined by agarose gel electrophoresis and visualized by SYBR Safe DNA gel

stain (Invitrogen). The reaction samples of V2c or its variants were scaled up for

Western blotting analysis as described (31). Anti-V2c antibody was generated in

rabbits against two V2c peptides, PFYWDFPNSQVGRD and KRIAMLRGDPTRYD

(indicated in Figure S1).

Western blot analysis

Western blot analysis was performed as described (ref). In general, proteins were

resolved by SDS-PAGE and transferred onto a PVDF membrane by using a transfer

apparatus (Bio-Rad). The membrane was probed with primary antibody against V2c

(1:4000), Hcp (1:2500), TssB (1:4000) and GFP (1:1000), followed by incubation with

horseradish peroxidase-conjugated anti-rabbit secondary antibody (Chemichem)

(1:25000) and visualized with the ECL system (Perkin-Elmer Life Science, Boston,

USA).

Fluorescence microscopy and image analysis

E. coli DH10B carrying the effector-expressing plasmids were grown overnight at 37°C.

Overnight cultures were sub-cultured with OD_{600} ~0.01. After 1 h, L-arabinose (Ara)

was added (0.2% final concentration) to induce toxin expression. Cultures were

harvested after an hour and resuspended with 30 µL PBS with 0.5 % Tween 20 (PBST).

Three µL of cells was placed on 2% agarose pad and imaged. To check the DNA

degradation, the same protocol was followed and live cells were stained with 2 µg/mL

Hoechst (Life Technologies, USA) and 1 µg/mL FM4-64FX (Invitrogen) for 5 mins and

washed with PBS to remove the excess dye.

For co-culture of *D. dadantii* Δ*imp*-GFP(S65T) with various 1D1609 EI pair mutants,

strains were mixed at 1:10 ratio on AK agar. This is an optimized growth medium,

named as *Agrobacterium* Kill-triggering, *Agrobacterium* kill-triggering (AK) medium (3

g K_2HPO_4, 1 g NaH_2PO_4, 1 g NH_4Cl, 0.15 g KCl, 9.76 g MES in 900 mL ddH_2O, pH 5.5)

(24) and solidified by 2% (w/v) agar. Cells from 16 h co-culture were concentrated to

OD ~10 and spotted on 2% agarose pad.

Imaging was performed using an inverted fluorescent microscope (IX81, Olympus
Japan) with objective lens UPLSAPO 100XO (Olympus, Tokyo, Japan), DAPI (49000;
Chroma), GFP (49002; Chroma) and Cy3 (49005; Chroma) filter set and CCD camera
(ORCA-R2; Hamamatsu, Japan). The images were acquired using Olympus cellSens
Dimension software.

All the image analysis was done using Fiji analyze particles tool (32). The micrograph
of FM4-64 fluorescence was used to segment surfaces (i.e. individual bacterial cell)
with background subtraction and thresholding. Feret's diameter was used. The cell
length (μm) was determined from cumulative counts of three random frames per
sample. For quantification of cell degradation, the Hoechst intensity was normalized
using the formula intensity/unit area of cells, where intensity is major x minor x 3.14
(π).

Interbacterial competition assay

Interbacterial competition assay
Bacterial competitions were carried out as described before (7). Briefly, overnight
culture of *A. tumefaciens* and *D. dadantii* Δ*imp* were mixed at a 10:1 attacker to target
ratio and spotted on AK agar plates. The target strain harbor pRL662-GFP (S65T).
After 18 h incubation at 25°C, the co-cultured cells were collected, serially diluted and
plated on LB agar containing gentamicin to quantify surviving *D. dadantii* Δ*imp* by
counting CFU.

Statistical analysis
The statistical analyses were done using GraphPad Prism version 6.01 for Windows,
GraphPad Software, La Jolla California USA (www.graphpad.com). The number of
technical replicates and independent biological replicates, p values and statistical
tests performed are indicated in the figure legends. The mean of independent
replicates was compared using the following statistical tests: Student's t test, one-
way ANOVA followed by Tukey's multiple comparisons tests
(http://astatsa.com/OneWay_Anova_with_TukeyHSD/). The error bars indicate the
standard error of the mean (SEM).

ACKNOWLEDGEMENTS

The authors acknowledge Ching-Hong Yang for providing *Dickeya dadantii* 3937

strains, which were imported under the permit 106-B-003 issued by the Council of

Agriculture of Taiwan. We would like to thank the former and current Lai lab

members for their help and fruitful discussion throughout this study, as well as See-

Yeun Ting and Mao-Sen Liu for critically reading the manuscript and their valuable

comments. We also thank Venus Marie Gaela and Thomas Boudier for the guidance

and help in processing the images using Fiji and the Sanger DNA sequencing

service provided by Genomic Technology Core located at the Institute of Plant and

Microbial Biology, Academia Sinica. The funding was supported by grants National

Science and Technology Council (NSTC) of Taiwan (107-2311-B-001-019-MY3) and

Academia Sinica Investigator Award (AS-IA-107-L01) to EML, and NSTC grant (110-

2311-B001-011) to YLS. YWC was supported by the postdoctoral fellowship (NSTC

110-2811-B-001-504). The funders had no role in study design, data collection and

interpretation, or the decision to submit the work for publication.

AUTHOR CONTRIBUTION

Conceptualization: MNMS, EML; Investigation: MNMS, KP, PYH, YWC, LKS, YLS,

EML; Project Funding acquisition and administration: EML, YLS; Supervision: EML,

YLS; Writing of original draft: MNMS, EML; Writing – methodology, review & editing:

MNMS, KP, PYH, YWC, LKS, YLS, EML.

FIGURE LEGENDS

Fig 1. V2c is a PAAR-linked specialized effector containing C-terminal Tox-SHH motif. (A) Schematic diagram of V2c protein showing N-terminal DUF4150 PAAR region and the catalytic region (amino acid 380-409) which covers the Tox-SHH motif. **(B)** Partial sequence alignment of orthologs of V2c from BLASTP. The strain name and accession number are indicated on the left. SHH motif is highlighted in yellow, H383 in red as the metal ion coordinating residue, H384 and H408 of HNH motif for catalysis and metal ion binding, respectively shown in green targeted for mutagenesis are indicated. **(C)** Domain architectures of V2c orthologs, Tox-SHH-containing proteins found in this study. Full length sequences were analyzed and the domains were from Pfam database with E value threshold set at 10^{-5}. **(D)** Cartoon model of V2c (residue 380 – 409). The structure of V2c predicted by AlphaFold indicates V2c consisting a His-Me finger domain with two antiparallel β-strands (yellow) connected with an Ω loop (light blue), a histidine (H384) at the C-terminus of the first β-strand and followed by an α-helix (magenta). The confidence score of this region is higher than 90, indicating high structure accuracy.

Fig 2. V2c exhibits DNase activity *in vivo* and *in vitro*. (A) Growth inhibition analysis of expression of *v2c,* its variants, with or without *v3c* encoding putative immunity protein in *E. coli* DH10B. *E. coli* cultures were induced with 1 mM IPTG at 0 h for *v3c* expressed from pTrc200 plasmid followed by L-arabinose (Ara) induction at 1 h to induce *v2c* from pJN105 plasmid. Cell growth was recorded every hour at OD_{600}. Empty vectors (vectors) were used as control. Data are mean ± SEM from four independent experiments. **(B)** Number of colony forming unit ($\log_{10}$ CFU/mL) quantified after growth inhibition assay (8 h) in each sample grown on agar plates. Data represents mean ± SEM of four independent experiments. Different letters above the bar indicate statistically different groups of strains (P < 0.05) determined by Tukey's HSD test. **(C)** Plasmid degradation assay performed by *E. coli* DH10B cells harboring pTrc200 and pJN105 plasmid (vector) or derivatives expressing *v2c* or its variants with pTrc200, *v2c* or its variants with pTrc200 expressing *v3c* were induced with (+) or without (–) Ara for 2 h. Plasmid DNA was extracted and the degradation pattern was observed in agarose gel. Result shown is a representative data from three

independent experiments. **(D)** *In vitro* DNase activity assay performed by using

pTrc200HA plasmid DNA as DNA substrate and incubated with purified V2c or its

variants. Nuclease free water (ddH$_2$O), protein elution buffer (elution buffer) and eluate

from empty vector expression (pET28a) served as negative controls while Benzonase

nuclease served as a positive control. The digestion of DNA substrate was determined

by agarose gel electrophoresis. The reaction samples of V2c or its variants were

detected by western blotting using anti-V2c antibody (α-V2c).

Fig 3. V2c-dependent cellular phenotypes. (A) Images of *E. coli* DH10B cells

harboring vector(s) and its derivatives expressing *v2c* or its variants in the presence

or absence of its cognate immunity gene *v3c* with (+Ara) and without (-Ara) Arabinose

induction stained with FM4-64 (red) and Hoechst stain (blue). The micrographs from

left to right are FM4-64, Hoechst, phase contrast and a merged image of FM-4-64 and

Hoechst stains. Scale bar, 2 μm. Two representative images from two independent

experiments are shown. **(B)** Normalized Hoechst intensity (intensity/unit area of cells)

in different treatments and **(C)** Cell length (μm) as measured from the indicated

number of cells (n) per treatment. The graph shows a combined count from 12 frames,

with three from each of four independent experiments. The red line in (C) shows the

median with interquartile range. Data represents mean ± SEM of four independent

experiments. Different letters above the bar indicate statistically different groups of

strains (P < 0.05) determined by Tukey's HSD test.

Fig 4. V2a Tox-AHH DNase exhibits cell elongation phenotype. (A) Cartoon model

of V2a (residue 380 – 470). The structure of V2a predicted by AlphaFold indicates V2a

consisting a His-Me finger domain with its signature antiparallel β-strands (yellow)

connected with an Ω loop (light blue) with a histidine (H386, orange) at the C-terminus

of the first β-strand. Two α-helices (pink) with histidine residues (H430, H456 magenta)

followed the β-strands. N412 is found in the loop between the β-strands and H430 is

in the first α-helix (E429–G430). The predicted Local-Distance Difference Test

(pLDDTs) of the His-Me domains of V2a are from confident to very high, most of the

pLDDTs of the residues are > 90. **(B)** Morphological analysis of *E. coli* DH10B cells

harboring vector(s) and its derivatives expressing *v2a*, catalytic site (H385A) mutant

without (-Ara) and with (+Ara) arabinose induction. Cells were stained with FM4-64

(red) and Hoechst stain (blue). The micrographs from left to right are FM4-64, Hoechst,

phase contrast and a merged image of the two fluorescent images. Scale bar, 2 μm.

(C) Cell length (μm) and **(D)** Hoechst intensity in different treatments as determined

from a combined count of three random frames, the number of cells (n) per sample is

indicated. The graph shows a representative result of three independent experiments.

The red line shows the median with interquartile range Different letters above the bar

indicate statistically different groups of strains (P < 0.05) determined by Tukey's HSD

test.

Fig 5. Interbacterial competition with *D. dadantii*. (A) Fluorescence micrographs of

1D1609 WT::TssB-sfGFP and Δ*4*::TssB-sfGFP co-cultures against *D. dadantii* 3937

WT and Δ*imp* mutant. A representative area selected from several random fields is

shown. **(B)** Fluorescence micrographs of *D. dadantii* Δ*imp*-GFP(S65T) and 1D1609 EI

pair mutants' co-cultures, at 1:10 ratio, observed after 16h on AK agar. *D. dadantii*

Δ*imp*-GFP(S65T) indicated as Dd only is included as a control. The inset in phase

contrast shows a magnified image of cells in the red box. White arrows point to *D.*

dadantii cells. Scale bar, 2 μm. **(C)** Survival of target strains *D. dadantii* 3937 Δ*imp*-

GFP(S65T) harboring pRL662 plasmid (Gm^R) mixed at 1:10 ratio with the attacker

1D1609 EI mutant strains on the x axis. Surviving *D. dadantii* 3937 were quantified by

CFU. The attacker and target strains were spotted on AK agar plates. After 18h, the

co-culture was collected, serially diluted and spotted on LB agar plate containing Gm

to count the CFU. Data are mean ± SEM (n = 6 biological repeats from three

independent experiments). **(D)** Cell length after competition between *A. tumefaciens*

1D1609 EI mutants with *D. dadantii* 3937 Δ*imp*-GFP(S65T). Cell length of *D. dadantii*

was measured using Fiji analyze particles tool. Data are mean ± SEM (n=6 biological

repeats from three independent experiments).

Fig S1. (A) Schematic diagram of V4b. The Rhs domain is bordered by conserved

motifs. The CT is the C-terminal domain that encodes the putative toxin. **(B)** *E. coli*

growth inhibition analysis in *vgrGb*-associated EI pair. *E. coli* cultures were induced

with 1 mM IPTG at 0 h for the expression of putative immunity protein expressed on

pTrc200 plasmid followed by L-arabinose (Ara) induction at 1 h to induce putative toxin

gene from pJN105 plasmid. Cell growth was recorded every hour at OD_{600}. Empty

vector was used as control. Data are mean ± SEM of three independent experiments.

(C) Plasmid degradation assay performed by using full length V4b (*v4b*^{FL}) and C-

terminal domain only (*v4b*^{CT}). *E. coli* cells with pTrc200 and pJN105 plasmid (vector)

or expressing *v4b* and *v4b+v5b* were induced with (+) or without (-) arabinose (Ara)

for 2 h. Plasmid DNA was extracted and the degradation pattern was observed in

agarose gel. Result shown is a representative data from two independent experiments.

(D) Images of *E. coli* DH10B cells expressing *v4b* and *v4b* co-expressed with its

cognate immunity gene *v5b* (*v4b+v5b*) under condition with (+Ara) and without (-Ara)

arabinose induction. *E. coli* DH10B cells expressing empty vector (vector) under

condition with (+Ara) and without (-Ara) arabinose induction served as negative control.

Cells were stained with FM4-64 (red) and Hoechst stain (blue). The micrographs from

left to right are FM4-64, Hoechst, phase contrast and a merged image of the two

fluorescent micrographs. Two representative images from independent experiments

are shown. Scale bar, 2 μm. **(E)** Cell length (μm) as measured from the indicated

number of cells (n) per sample. The graph shows a combined count from 3 random

frames as a representative of two independent experiments. The red line shows the

median with interquartile range. **(F)** Hoechst intensity in cells analyzed in (D). Result

shown is a representative data of two independent experiments. Different letters above

the bar indicate statistically different groups of strains ($P < 0.05$) determined by

Tukey's HSD test.

Fig S2. (A) Representative fluorescence micrographs of *A. tumefaciens* 1D1609

(1D1609WT::TssB-sfGFP) and quadruple *vgrG* deletion mutant- (1D1609Δ4::TssB-

sfGFP) expressing TssB-sfGFP from the native *tssB* locus (Movie S1). No sheath

assembly was observed in 1D1609Δ4::TssB-sfGFP during four min time-lapse

observation period. Scale bar: 2 μm. **(B)** Western blot analysis of 1D1609WT::TssB-

sfGFP and 1D1609Δ4::TssB-sfGFP. The strains were grown in AB-MES (pH5.5). The

cellular and secreted (S) fractions were subjected to immunoblotting probed with

antibodies as indicated. Molecular weight markers (in kDa) are indicated on the left,

TssB is about 19 kDa while GFP is 27 kDa, resulting 46-kDa TssB-GFP fusion protein.

The T6SS inactive mutant *Δ4ΔtssB* served as a negative control. Similar results were

obtained from two independent experiments.

Fig S3. Cell length (μm) of **(A)** *v2c* single catalytic site mutants and **(B)** catalytic

variants co-expressed with *v3c*. Cell length (μm) as measured from the indicated

number of cells (n) per sample. The graph shows a combined count from six random

frames. Data represents mean ± SEM of four independent experiments. Different

letters above the bar indicate statistically different group of strains (P < 0.05)

determined by Tukey's HSD test. The red line shows the median with interquartile

range.

Fig S4. Sequence alignment of V2a and V2c. The conserved amino acid residues are

indicated as '*' (identical), ':' (similar) and '.' (slightly similar). The DUF4150, PAAR-

like domain is in green. The histidine residues in AHH and SHH motifs in His-Me finger

domain of V2a and V2c respectively, V2aH385/V2cH383 and V2aH386/V2cH384, are

in red. The histidine residues that predicted to responsible for metal binding in His-Me

finger domain, V2aH430/V2cH408, are in a bold blue font. Two peptides used for

generating antibodies were underlined.

SUPPLEMENTARY VIDEO

Video S1. Time-lapse recording of *A. tumefaciens* 1D1609 cells expressing TssB-

sfGFP from the native *tssB* locus (WT::TssB-sfGFP) showing the full cycle of T6SS

assembly: sheath initiation, extension, contraction and disassembly, visualized in less

than ~75 s. Up to three visible structures were observed in a single cell and the sheath

spreads throughout the width and length of the cell. T6SS machine cycle from one

location in the cell to another location.

Supplementary Table S1. Bacterial strain and plasmids used in this study

Supplementary Table S2. Primers used in this study

REFERENCES

1. Brackmann M, Nazarov S, Wang J, Basler M. Using Force to Punch Holes: Mechanics of Contractile Nanomachines. Trends in Cell Biology. 2017;27(9):623-32.

2. Shneider MM, Buth SA, Ho BT, Basler M, Mekalanos JJ, Leiman PG. PAAR-repeat proteins sharpen and diversify the type VI secretion system spike. Nature. 2013;500(7462):350-3.

3. Basler M, Pilhofer M, Henderson GP, Jensen GJ, Mekalanos JJ. Type VI secretion requires a dynamic contractile phage tail-like structure. Nature. 2012;483(7388):182-6.

4. Cianfanelli FR, Monlezun L, Coulthurst SJ. Aim, Load, Fire: The Type VI Secretion System, a Bacterial Nanoweapon. Trends in Microbiology. 2016;24(1):51-62.

5. Coulthurst S. The Type VI secretion system: a versatile bacterial weapon. Microbiology. 2019;165(5):503-15.

6. Hwang HH, Yu M, Lai EM. *Agrobacterium*-mediated plant transformation: biology and applications. Arabidopsis Book. 2017;15:e0186.

7. Wu CF, Santos MNM, Cho ST, Chang HH, Tsai YM, Smith DA, Kuo CH, Chang JH, Lai EM. Plant pathogenic *Agrobacterium tumefaciens* strains have diverse type VI effector-immunity pairs and vary in in planta competitiveness. Molecular Plant-Microbe Interactions. 2019; 32(8).

8. Lin C, Lin YC, Haryono M, Santos MNM, Cho ST, Weisberg A, Wu CF, Chang J, Lai EM, Kuo CH. Modular evolution of secretion systems and virulence plasmids in a bacterial species complex. BMC Biology. 2022; 20:16.

9. Santos MNM, Cho ST, Wu CF, Chang CJ, Kuo CH, Lai EM. Redundancy and Specificity of Type VI Secretion *vgrG* Loci in Antibacterial Activity of *Agrobacterium tumefaciens* 1D1609 Strain. Frontiers in Microbiology. 2019;10(3004):3004.

10. Jablonska J, Matelska D, Steczkiewicz K, Ginalski K. Systematic classification of the His-Me finger superfamily. Nucleic Acids Research. 2017;45(20):11479-94.

11. Zhang D, de Souza RF, Anantharaman V, Iyer LM, Aravind L. Polymorphic toxin systems: Comprehensive characterization of trafficking modes, processing, mechanisms of action, immunity and ecology using comparative genomics. Biology Direct. 2012;7(1):18.

12. Wu CC, Lin LJ and Yuan HS. Structures, Mechanisms, and Functions of His-Me Finger Nucleases. Trends in Biochemical Sciences. 2020;45(11):935-946.

13. Zhang D, Iyer LM, Aravind L. A novel immunity system for bacterial nucleic acid
degrading toxins and its recruitment in various eukaryotic and DNA viral systems. Nucleic
Acids Research. 2011;39(11):4532-52.

14. Jana B, Fridman CM, Bosis E, Salomon D. 2019. A modular effector with a DNase
domain and a marker for T6SS substrates. Nature Communications 10:3595 (2019).

15. Söding J, Biegert A, Lupas AN. The HHpred interactive server for protein homology
detection and structure prediction. Nucleic Acids Research. 2005;33:W244-W8.

16. Wu CF, Lien YW, Bondage D, Lin JS, Pilhofer M, Shih YL, Chang JH, Lai EM. Effector
loading onto the VgrG carrier activates type VI secretion system assembly. EMBO Reports.
2019; 5:e47961.

17. Pissaridou P, Allsopp LP, Wettstadt S, Howard SA, Mavridou DAI, Filloux A. The
Pseudomonas aeruginosa T6SS-VgrG1b spike is topped by a PAAR protein eliciting DNA
damage to bacterial competitors. Proceedings of the National Academy of Sciences of the
United States of America. 2018;115(49):12519-24.

18. Pei T-T, Li H, Liang X, Wang Z-H, Liu G, Wu L-L, et al. Intramolecular chaperone-mediated
secretion of an Rhs effector toxin by a type VI secretion system. Nature Communications.
2020;11(1):1865.

19. Fridman CM, Jana B, Ben-Yaakov R, Bosis E, Salomon D. A DNase Type VI Secretion
System Effector Requires Its MIX Domain for Secretion. Microbiology Spectrum.
2022;13:e0246522.

20. Bernal P, Allsopp LP, Filloux A, Llamas MA. The *Pseudomonas putida* T6SS is a plant
warden against phytopathogens. ISME Journal. 2017;11(4):972-987.

21. Li Y, Yan X, Tao Z. Two Type VI Secretion DNase Effectors are Utilized for Interbacterial
Competition in the Fish Pathogen *Pseudomonas plecoglossicida*. Frontiers in Microbiology.
2022;13:869278.

22. Wu C-C, Lin JLJ, Yuan HS. Structures, Mechanisms, and Functions of His-Me Finger
Nucleases. Trends in Biochemical Sciences. 2020;45(11):935-46.

23. Galburt EA, Chevalier B, Tang W, Jurica MS, Flick KE, Monnat RJ, et al. A novel
endonuclease mechanism directly visualized for I-PpoI. Nature Structural Biology.
1999;6(12):1096-9.

24.   Yu M, Wang YC, Huang CJ, Ma LS, Lai EM. *Agrobacterium tumefaciens* deploys a
versatile antibacterial strategy to increase its competitiveness. Journal of Bacteriology
2021;203(3):e00490-20.

25.   Ting S-Y, Bosch DE, Mangiameli SM, Radey MC, Huang S, Park Y-J, et al. Bifunctional
Immunity Proteins Protect Bacteria against FtsZ-Targeting ADP-Ribosylating Toxins. Cell.
2018;175(5):1380-92.e14.

26.   Justice SS, Hunstad DA, Cegelski L, Hultgren SJ. Morphological plasticity as a bacterial
survival strategy. Nature Reviews Microbiology. 2008;6(2):162-8.

27.   LaCourse KD, Peterson SB, Kulasekara HD, Radey MC, Kim JY, Mougous JD. Conditional
toxicity and synergy drive diversity among antibacterial effectors. Nature Microbiology
2018;3:440–446.

28.   Wu CF, Weisberg AJ, Davis EW 2nd, Chou L, Khan S, Lai EM, Kuo CH, Chang JH.
Diversification of the Type VI Secretion System in Agrobacteria. mBio.
2021;26;12(5):e0192721.

29. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res. 2004;32(5):1792-7.

30. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar Gustavo A, Sonnhammer ELL, et
al. Pfam: The protein families database in 2021. Nucleic Acids Research. 2021;49(D1):D412-
D9.

31. Ma L-S, Hachani A, Lin J-S, Filloux A, Lai E-M. *Agrobacterium tumefaciens* deploys a
superfamily of type VI secretion DNase effectors as weapons for interbacterial competition in
planta. Cell Host Microbe. 2014;16(1):94-104.

32. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
source platform for biological-image analysis. Nature Methods. 2012;9(7):676-82.

Fig 1. Domain and sequence analysis of V2c.

(A)

(B)

Agrobacterium tumefaciens 1D1609	AVH44366.1	-NLDSHHTPANS---A--SYLPTWAG-----PAIQMLPEDHR	409
Silvanigrella aquatica 1915309#	WP_148696535.1	-GLDSHHMPAKS---AYEGSKMNPDGEAKQGPAIKMEPKDHR	268
S. enterica serovar Enteritidis IVB 176/82*	HAU6875067.1	-GLDSHHMPAKD---S--SPLSHDDG-----PAIHMDPLDHR	149
S. enterica serovar Enteritidis SARB17*	ELO80288.1	-GLDSHHMPAKD---S--SPLSHDDG-----PAIHMDPLDHR	150
Ralstonia solanacearum SEPPX05**	ARU21619.1	-GKDSHHMPADD---V--SPLARNGG-----PAIQMEPKDHK	221
Ralstonia solanacearum 305**	WP_087451447.1	-GKDSHHMPADD---V--SPLARNGG-----PAIQMEPKDHK	367
Paraburkholderia phosphatilytica 2282883**	WP_124800779.1	-GKDSHHMPADD---A--SPLPKKDG-----PAIQMDPADHH	217
Caballeronia mineralivorans 2010198**	WP_053064266.1	-GKDSHHMPADD---V--SPLKRNDG-----PAIQMEPDDHA	323
Burkholderia vietnamiensis 60552**	WP_196483818.1	-GKDSHHMPADD---V--SPLKRSDG-----PAIQMDPDDHA	227
Burkholderia vietnamiensis 60552**	WP_155626574.1	-GKDSHHMPADD---V--SPLKRSDG-----PAIQMDPDDHA	243
Burkholderia cenocepacia 95486**	WP_200039262.1	-GKDSHHMPADD---V--SPLKRSDG-----PAIQMDPDDHA	243
Burkholderia anthina 179879**	WP_155634575.1	-GKDSHHMPADD---V--SPLKRSDG-----PAIQMDPDDHA	227
Burkholderia cenocepacia 95486**	WP_124550222.1	-GKDSHHMPADD---V--SPLKRSDG-----PAIQMDPDDHA	243
Massilia armeniaca 2072590**	WP_181259249.1	-GKDSHHLPADD---V--SPLKRSDG-----PAIQMDPEDHK	236
Herbaspirillum rubrisubalbicans M1**	ALU88908.1	-GKDSHHMPADN---V--SPLKSNEG-----PAIQMDPLDHK	244
Herbaspirillum rubrisubalbicans M1**	ALU90982.1	-GKDSHHMPADD---V--SPLKRNEG-----PAIQMDPLDHK	244

(C)

(D)

Fig 2. V2c exhibits DNase activity *in vivo* and *in vitro.*

Fig 3. V2c-dependent cellular phenotypes.

Fig 4. V2a also exhibits cell elongation phenotype.

Fig 5. Interbacterial competition with *D. dadantii*.

(A)

(B)

(C)

(D)

Figure S1.

Figure S2

(A)

(B)

(A)

(B)

Figure S4.

```
V2a    MSIPRDNYIGEPQYPGTWTTATPREGLRDIDEARIVSLAPDVCLTPVGSSVVPIPYPVVD
V2c    MSIPRDNYIGEPQYPGTWTTATPREGLRDIDEARIVSLAPDVCLTPVGSSVVPIPYPVVD
       ************************************************************
V2a    FCGHDKNYTPSVRFTGKKAMVMRSFTTHVHGDAPGNRKGVKSGTVESICEPIGHADQVRA
V2c    FCGHDKNYTPSVRFTGKKAMVMRSFTTHVHGDAPGNRKGVKSGTVESICEPIGHADQVRA
       ************************************************************
V2a    EGSHVIRHLDRFYMNSRNTQGEAIFVRSTQTYDPPKDDDPVRGSLRWQDGSDEGRVMSDA
V2c    EGSHVIRHLDRFYMNSRNTQGEAIFVRSTQTYDPPKDDDPVRGSLRWQDGSDEGRVMSDA
       ************************************************************
V2a    SPEPFKEGEKYAYAGSAAAATTGAAQNSAQALRGGMPATRPTVPQTVPRGPGRVITPGPA
V2c    SPEPLIMGAQYAQAQTATIPQSGGV-----ARRGTYGRPTTTVDPRLLKGL-RFWSSARG
       ****:   *  :**  *  :*:   . :*..      *  **    . .**    :  *   *. :... .
V2a    RAP---QWYRTPPHRVVSDLA-KVG---------RLARVGIWAR-RLNVIGILSSVLIPD
V2c    AAEIMDMVQQISEDRAVQDSAGQLGLNLSDPTDVLVARAHAWSKYHIDASPFYWD--FPN
        *         . .  *.*.* *  ::*          :**.   *:.  .::.   :   .  :*:
V2a    NGANLSDVVPRDDQERRIKEMFDEGVDAGHDSWEMRDWAMEELRRHRQRVREEPESRPQA
V2c    SQ------VGRDAATKAITSM--EAAAPG----TLGE-ALQGDRTQLEKLRTAIKNAVSA
       .        * **   . *..*  *..  .* 　  : : *:: 　 * : :.:* 　:.   .*
V2a    LPQSETVRVSDRDEYRKKCQVDRYGKMKSVCGQYGMQAHHIVPDWTLRYGAREEADKRIP
V2c    ATAAASVRVSG-----QPCAREALG----APGKYRGGAHFAT---ALPLGDNLDSHHTPA
       . : :*****. 　  : * : *  　 . *:* 　 **. . 　 :* 　 * . :: : .
V2a    NMPSLNDGMAICVIGNARVEQTEHWGGHLAD------KAIENAGRNSTPPFTAPLHKVVQ
V2c    NSASY---------------LPTWAGPAIQMLPEDHRATLSYGRKVTSPYISTQARLIN
       * .*                    *.*   :        .*  .  **: *.*: :.   .:::
V2a    ASAQAMISVRPDCAKEIMAALTTQYAGKNPFQLLRAK--QYPPLPEETISALKSGAVEGA
V2c    SG------------QFMKAINMDILDIKRIAMLRGDPTRYDAAIAQMMAYAECLKREGI
       :.            ::* *:. : 　 . : : :**.. 　.* . 　 : :: 　 :.    **
V2a    TRKP
V2c    IK--
        .
```

Supplementary Video 1.

Table S1. Bacterial strains and plasmids

Strain	Relevant characteristics	Reference/Source	EML No.
E. coli			
DH10B	Host of plasmid for cloning	Lab stock	EML455
S17-1	Host for conjugation	Lab stock	EML124
BL21 (DE3)	For protein expression	Lab stock	EML117
D. dadantii **3937**			
WT	Wild-type *Dickeya dadantii* 3937	Ching-Hong Yang, University of Wisconsin, Milwaukee	EML4911
Δ*imp*	T6SS inactive strain with deletion of *imp* operon	Ching-Hong Yang, University of Wisconsin, Milwaukee	EML4921
A. tumefaciens **1D1609**			
WT	Alfalfa isolate, virulent strain containing pTi1D1609	Clarence Kado, UC Davis	EML304
Δ*4* or Δ*vgrGabcd*	Deletion of *vgrGa*, *vgrGb*, *vgrGc* and *vgrGd*	Santos et al., 2020	EML4746
Δ2*EIbd*	Deletion of *vgrGb*- and *vgrGd*-associated effector-immunity pair	This study	EML5531
Δ3EIbcd #1	Deletion of *vgrGb*-, *vgrGc*- and *vgrGd*-associated effector - immunity pair	This study	EML5594

Δ3EIbcd #2	Deletion of *vgrGb*-, *vgrGc*- and *vgrGd*-associated effector - immunity pair	This study	EML5920
Δ3*EIabd*	Deletion of *vgrGa*-, *vgrGb*- and *vgrGd*-associated effector - immunity pair	Santos et al., 2020	EML5515
Δ4*EIabcd*	Deletion of *vgrGa*-, *vgrGb*- and *vgrGd*-associated effector - immunity pair and whole *vgrGc* cluster (Δ*vgrGc-v3*)	Santos et al., 2020	EML5540
1D1609 WT::TssB-sfGFP	*tssB*-sfGFP integrated into WT chromosome to replace the endogenous *tssB*	This study	EML5559
Δ4::TssB-sfGFP	*tssB*-sfGFP integrated into Δ*4* chromosome to replace the endogenous *tssB*	This study	EML5567

Plasmids			
pJQ200KS	Gm^R, suicide plasmid containing GmR and sacB gene for selection of double crossover	Quandt and Hynes, 1993	EML121
pRL662	Gm^R, broad host range vector derived from pBBR1MCS-2	Vergunst et al., 2000	EML315
pTrc200	Sp^R, pVS1 origin, *lacIq, trc* promoter expression vector	Schmidt-Eisenlohr et al., 1999	EML904

pJN105	Gm^R, arabinose-inducible gene expression vector derived from pBBRMCS-1, *araC*-P$_{BAD}$	Newman and Fuqua, 1999	EML4220
pET28a	Km^R, Protein expression vector, N-terminal His-tagged	Lab stock	EML2485
pJN-*v2a*	Gm^R, pJN expressing *v2a*	Santos et al., 2020	EML5272
pJN-*v4b*^{FL}	Gm^R, pJN expressing the full-length *v4b*	This study	EML4758
pJN-*v4b*^{CT}	Gm^R, pJN expressing the C-terminal toxin domain of *v4b*	This study	EML4756
pJN-*v2c*	Gm^R, pJN expressing *v2c*	This study	EML5328
pTrc-*v5b*	Sp^R, pTrc200 expressing the immunity protein V5b	This study	EML4759
pTrc-*v3c*	Sp^R, pTrc200 expressing the immunity protein V3c	This study	EML5337
pJN-*v2a* (H385A)	Gm^R, pJN expressing *v2a* with amino acid substitution (H385A)	Santos et al., 2020	EML5298
pJN-*v2c* (H383A)	Gm^R, pJN expressing *v2c* with amino acid substitution (H383A)	This study	EML5925
pJN-*v2c* (H384A)	Gm^R, pJN expressing *v2c* with amino acid substitution (H384A)	This study	EML5926
pJN-*v2c* (H408A)	Gm^R, pJN expressing *v2c* with amino acid substitution (H408A)	This study	EML5927
pJN-*v2c* (HAHA)	Gm^R, pJN expressing *v2c* with amino acid substitution (H383AH384A)	This study	EML5922

pJN-*v2c* CR	Gm^R, pJN expressing *v2c* with deletion of the conserve region of Tox-SSH domain	This study	EML5589
pJQ200sk-*vag4vag5b*	Gm^R, plasmid to generate Δ3*EIbcd* #2	This study	EML5307
pJQ200sk-*vag2vag3d*	Gm^R, plasmid to generate Δ2*EIbd*, Δ3*EIbcd* #1 and Δ3*EIabd*	This study	EML5340
pRL662-GFP(S65T)	GFP from pRU1156 cloned into pRL662	Lab stock	EML3376
pJQ200sk-tssBsfGFP	C-terminal fusion of TssB with superfolder GFP (sfGFP) in pJQ200ks to replace the endogenous *tssB* in WT and Δ*4* background	This study	EML5557
pET28a-*v2c*	pET28a expressing His-tagged V2c	This study	EML6050
pET28a-*v2c*HAHA	pET28a expressing His-tagged V2c H383AH384A	This study	EML6051
pET28a-*v2c*H408A	pET28a expressing His-tagged V2cH408A	This study	EML 6063
pET28a-*v2c*H383A	pET28a expressing His-tagged V2cH383A	This study	EML 6064
pET28a-*v2c*H384A	pET28a expressing His-tagged V2cH384A	This study	EML 6065

Table S2. Primers used in this study

Primer Name	Sequences	Purpose
V2c(H383A)	5'-CCG<u>GAATTC</u>AAGGAACAGCTATGAGCATCCCCCGCGAC-3' 5'- GACAATTTAGATAGCGCGCATACTCCTGCTAACTCT-3' 5'- AGAGTTAGCAGGAGTATGCGCGCTATCTAAATTGTC-3' 5'- CG<u>TCTAGA</u>TCACTTTATGATTCCTTCGCG-3'	mutagenesis
V2c(H384A)	5'-CCG<u>GAATTC</u>AAGGAACAGCTATGAGCATCCCCCGCGAC-3' 5'-GACAATTTAGATAGCCATGCGACTCCTGCTAACTCT-3' 5'-AGAGTTAGCAGGAGTCGCATGGCTATCTAAATTGTC-3' 5'- CG<u>TCTAGA</u>TCACTTTATGATTCCTTCGCG-3'	mutagenesis
V2c(H408A)	5'-CCG<u>GAATTC</u>AAGGAACAGCTATGAGCATCCCCCGCGAC-3' 5'- ATGCTTCCAGAAGACGCGCGAGCTACACTCTCATAT -3' 5'-TGAGAGTGTAGCTCGCGCGTCTTCTGGAAGCAT-3' 5'- CG<u>TCTAGA</u>TCACTTTATGATTCCTTCGCG-3'	mutagenesis
V2c(CR)	5'-GTTTTTTTGGGCTAGCGAAGGAACAGCTATGAGCATC-3' 5'-CCTTTCTTCCATATGAGAGTGTAGCATTGTCGCCAAGTGGGAGGGCCGTC-3' 5'-GACGGCCCTCCCACTTGGCGACAATGCTACACTCTCATATGGAAGAAAGG -3' 5'-GCGGTGGCGGCCGCTTCACTTTATGATTCCTTCGC-3'	mutagenesis
V2c(H383A H384A)	5'-CCG<u>GAATTC</u>AAGGAACAGCTATGAGCATCCCCCGCGAC-3' 5'-GACAATTTAGATAGCCATGCGACTCCTGCTAACTCT-3' 5'-AGAGTTAGCAGGAGTCGCATGGCTATCTAAATTGTC-3' 5'- CG<u>TCTAGA</u>TCACTTTATGATTCCTTCGCG-3'	mutagenesis
pJQ200KS- *v4bv5b*	5'-TT<u>TCTAGA</u>CATCTGTTGTTCGCGGCCGATG-3' 5'-GC<u>GGATCC</u>ATCACCCGCCATGTGAGGCCT-3' 5'-GC<u>GGATCC</u>TTCGCTGCCCTGTGACCGTTCT-3' 5'-GC<u>CCCGGGG</u>CAAGATGATTCCGGGAATTGAC-3'	deletion mutant
pJQ200KS- v2dv3d	5'-GC<u>TCTAGA</u>GAGCGCCGAAATCCGCTT-3' 5'-CG<u>GGATCC</u>GATGCTCATGCTGTCTCCCTCTC-3' 5'-CGG<u>GATCC</u>GCCAGAAAGCCATAGATTGTCG-3' 5'-TCCC<u>CCCGGGG</u>CATCACCACACTCTCGAGCCT-3'	deletion mutant

V4b	5'-CCGTCTAGAAAGGAACAGCTATGGCGGGTGAT-3' 5'-CGGAGCTCGTTTCGATACTCAGCATCGACCG-3'	overexpression
V5b	5'-TATAGGTACCCGAGAATAAGAACCGCGTTC-3' 5'-GCTCTAGAATTGGCGAGGCTACAGCG-3'	overexpression
V2c	5'-CCGGAATTCAAGGAACAGCTATGAGCATCCCCCGCGACAATTAT-3' 5'-CGTCTAGATCACTTTATGATTCCTTCGCGTTTAAGGC-3'	overexpression
V3c	5'-ATCCGAGCTCGTGACAGACAAGTCGGGAGCG-3' 5'-GCTCTAGACCTAGATCCATCGCTTACTGCCTC-3'	overexpression
tssB-sfGFP	UF 5'-TTTCTAGACGAGCTTCGACGCCATCATGGCGG-3' UR 5'-TTTGCTCGCGCCGTTCTTTTCCTT-3' sfGFP F 5'-CAAGGAAAAGAACGGCGCGAGCAAAATGCGTAAAGGCGAAGAGCTGTTCA-3' sfGFP R 5'-GGGAGGTTTTCCTCCGCACAGGCGATCATTTGTACAGTTCATCCATACCA-3' DF 5'-TGGTAGGGATGAACTGTACAAATGATCGCCTGTGCGGAGGAAAACCTCCC-3' DR 5'-TCCCCCCGGGCAGCTGTACGTCATCGAAGGAA-3'	Chromosomal replacement
pET28a-v2c	5'-AATCTAGAATGAGCATCCCCCGCGACAA-3'. 5'-AACTCGAGCTTTATGATTCCTTCGC-3'	Protein expression
pET28a-v2cHAHA	5'-GACAATTTAGATAGCGCGGCGACTCCTGCTAACTCT-3'. 5'-AGAGTTAGCAGGAGTCGCCGCGCTATCTAAATTGTC-3'	Protein expression
pET28a-v2c-H383A	5'-GACAATTTAGATAGCGCGCATACTCCTGCTAACTCT-3' 5'-AGAGTTAGCAGGAGTATGCGCGCTATCTAAATTGTC-3'	Protein expression
pET28a-v2c-H384A	5'-GACAATTTAGATAGCCATGCGACTCCTGCTAACTCT-3' 5'-AGAGTTAGCAGGAGTCGCATGGCTATCTAAATTGTC-3'	Protein expression
pET28a-v2c-H408A	5'-ATGCTTCCAGAAGACGCGCGAGCTACACTCTCATAT-3' 5'-TGAGAGTGTAGCTCGCGCGTCTTCTGGAAGCAT-3'	Protein expression

Restriction enzyme sites are underlined. UF (upstream forward), UR (upstream reverse), DF (downstream forward) and DR (downstream reverse).