Title: Plasma cells in human pancreatic ductal adenocarcinoma secrete antibodies to self-

antigens

Authors: Min Yao[1]†, Jonathan Preall[1], Johannes Yeh[1], Darryl Pappin[1], Paolo Cifani[1], Yixin Zhao[1],

Sophia Shen[2], Philip Moresco[1,3,4], Brian He[1], Hardik Patel[1], Amber Habowski [1], Daniel A. King[5], Kara

Raphael[5], Arvind Rishi[5], Divyesh Sejpal[5]‡, Matthew Weiss[5], David Tuveson[1], Douglas Fearon[1,6*]

Affiliations:

[1]Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724.

[2]Cold Spring Harbor High School, Cold Spring Harbor, NY 11724.

[3]Graduate Program in Genetics, Stony Brook University, Stony Brook, NY 11794.

[4]Medical Scientist Training Program, Stony Brook University Renaissance School of Medicine, Stony

Brook University, Stony Brook, NY 11794.

[5]North Shore University Hospital, Manhasset, NY 11030.

[6]Weill Cornell Medicine, New York, NY 10065.

*Corresponding author. Email: dfearon@cshl.edu

†Present addresses: Tri-Institutional Therapeutics Discovery Institute, New York, NY 10021.

‡ Present addresses: Digestive Disease Institute, CommonSpirit Health Southwest Division and Dignity

Health Medical Group, Phoenix, NY 85012.

One Sentence Summary: We have identified that plasma cells in human pancreatic ductal
adenocarcinoma prominently secrete antibodies recognizing intracellular unmutated self-antigens.

Abstract:

Intratumoral B cell responses are associated with more favorable clinical outcomes in human pancreatic

ductal adenocarcinoma (PDAC). However, the antigens driving these B cell responses are largely

unknown. We sought to discover these antigens by using single-cell RNA sequencing (scRNA-Seq) and

immunoglobulin (Ig) sequencing of tumor-infiltrating immune cells from seven primary PDAC samples.

We identified activated T and B cell responses and evidence of germinal center reactions. Ig sequencing

identified plasma cell (PC) clones expressing isotype-switched and hyper-mutated Igs, suggesting the

occurrence of T cell-dependent B cell responses. We assessed the reactivity of 41 recombinant antibodies

that represented the products of 235 PCs and 12 B cells toward multiple cell lines and PDAC tissues, and

observed frequent staining of intracellular self-antigens. Three of these antigens were identified: the

filamentous actin (F-actin), the nucleic protein, RUVBL2, and the mitochondrial protein, HSPD1.

Antibody titers to F-actin and HSPD1 were elevated in the plasma of PDAC patients, and also detectable

in healthy donors. Thus, PCs in PDAC produce auto-antibodies reacting with intracellular self-antigens,

which may result from promotion of pre-existing, autoreactive B cell responses. These observations

indicate that the chronic inflammatory microenvironment of PDAC can support the adaptive immune

response.

Main Text:

INTRODUCTION

Microsatellite-stable PDACs do not respond to immunotherapy, have few T cells in cancer cell nests, and

have been thought to minimally stimulate the adaptive immune system (*1*). Recently, however, it has been

observed that PDAC patients whose tumors contained tertiary lymphoid structures (TLS) with germinal

centers, memory B cells, and memory CD4+ T cells had improved long-term survival (*2-4*), suggesting

the occurrence of clinically relevant, ongoing anti-PDAC immune responses. The possibility of these

immune responses is supported by a study involving patients with PDAC and colorectal cancer in which

one-week continuous administration of an inhibitor to the chemokine receptor, CXCR4, revealed on-

going anti-tumor immune responses (*5*).

The antigens driving these intratumoral immune reactions in human PDAC are unknown. Although

PDAC has a low mutational frequency compared to other cancers (*1*), attention has been directed to T cell

clones from PDAC patients with specificity to neoantigens arising from mutations that predicted cross-

reactive microbial epitopes (*6*). The intratumoral immune response may also be directed towards

germline-encoded antigens, a concept that has been supported by the finding that immunization of mice

with induced pluripotent stem cells confers protection against several tumor models (*7, 8*). Thus, defining

the range of antigens that are driving the intratumoral immune response in PDAC may expand our

knowledge of the interaction between this cancer and the immune system.

B cell activation is triggered by antigen interaction with membrane immunoglobulin (Ig). This

interaction may lead to B cell activation, antigen-presentation to CD4 T cells, and the formation of

germinal center reactions. The latter will lead to B cell clonal expansion, heavy (H) chain isotype

switching, H chain and light (L) chain variable region somatic hypermutation (SHM), and ultimately

differentiation into antibody-secreting PCs. Therefore, antibodies derived from activated B cell responses

may help identify those antigens that are driving intratumoral immune responses. In the current report, we

have used scRNA-Seq of PCs and B cells from primary PDAC specimens to identify their paired H and L

chains which enabled screening of antibody reactivity.

RESULTS

Identification of B and T cell responses by scRNA-Seq in human PDAC

We performed scRNA-Seq of CD45+ immune cells from seven untreated primary PDAC samples,

including four microsatellite-stable surgically resected tumors and three fine-needle aspiration (FNA)

biopsies (Table S1). scRNA-Seq identified a total of 26,702 immune cells and 327 non-immune cells. The

major immune cells included T cells (66%), myeloid cells (19%), PCs (7.4%) and B cells (3.6%), with T

and myeloid cells being the most frequent cell types in each sample (Fig. 1A-B and Fig. S1A). In the

CD8+ T cells, features of T cell activation were prominent, as evidenced by frequent expression of the

effector genes, *PRF1* (46% of CD8 T cells) and *GZMB* (31%), as well as the cytokines, *IFNG* (13%) and

TNF (11%). Subsets of *CD8+* T cells also expressed the inhibitory receptors, *PD-1* (14%) and *CTLA4*

(3%) (Fig. 1C and Fig. S1B), which also identify effector cells. In the *CD4+* T cell population, a Th1

response was apparent, as assessed by the expression of the Th1 lineage marker, *TBX21* (13% of CD4 T

cells), and the Th1 cytokines, *TNF* (18%) and *IFNG* (6%). *FOXP3+ CD4+* regulatory T cells represented

a relatively abundant CD4 T cell population (22%), and characteristically co-expressed *IL2RA*, *PD-1* and

CTLA4. *CD4+* T follicular helper T cells (Tfh cells) were identified by their expression of *BCL6* (2%),

IL-21 (1%), and the chemokine receptor, *CXCR5* (2%) (Fig. 1C and Fig. S1C-D). In summary, the PDAC

tumors contained activated CD8 and CD4 T cells, including Tfh cells of presumed germinal center origin.

The B cell cluster (n=963 cells) was defined by the expression of the pan B cell markers, *CD19* and

CD20. 35% of B cells expressed the naïve H chain isotype, *IgD*, and B cells that expressed isotype

switched *IgG1* (19%) were also present (Fig. 1C). Additional evidence for an active intratumoral B cell

response was indicated by the presence of germinal center B cells, as assessed by their expression of

BCL6 (4%) and *AICDA* (1%), the latter of which mediates SHM and isotype switching (Fig. 1C and Fig.

S1E). PCs (n=1984) were identified by the makers, *J chain*, *PRDM1*, and *CD138*, and high expression of

Ig H and L chains (Fig. 1C). Most PCs (95% of PCs) no longer expressed *CD19*, indicating they were

relatively mature PCs rather than plasmablasts. Thus, PDAC tumors contained activated B cells and

terminally differentiated PCs.

We further obtained histopathology images from the four resected PADC samples which we have

performed scRNA-Seq. We have observed presence of multiple TLS-like structures in each tumor

sample, with TLSs typically located in the tumor border (Fig. 1D and Fig.S2). This observation supports

that the B and T cell response revealed by scRNA-Seq may take place inside those TLSs.

Igs from PCs in PDAC featuring isotype switching, somatic hypermutation and clonal expansion

A total of 615 PCs with paired H and L chains were detected in six PDAC samples (Fig. 2A and Fig.

S3A). A minority of PCs (0-17%) expressed the *IgM* isotype, while the majority had undergone class

switching to *IgG1*, *IgG2*, *IgA2*, or *IgA1* isotypes (Fig. 2B and Fig. S3B). Analyzing SHM in the V regions

revealed an average of 23 and 15 mutations in the H and L chains, respectively (Fig. 2C). These rates of

SHM are comparable to those in antibodies induced by viral infections (*9*). Expanded PC clones were

identified in four of the six PDAC samples, with clone sizes ranging from 2 to 46. Strikingly, in patients

19 and 20, more than half of the Ig-paired PCs were products of clonal expansion (Fig. 2D). Analysis of

the evolution of V region SHM sequences revealed that most PCs originated from single expanded

nodules, although additional lineage evolutions were present (Fig. 2E and Fig. S3C), consistent with

typical germinal center reactions. Fewer B cells with paired Igs were sequenced (n=473), likely due to

lower Ig expression, which yielded only four small, expanded clones (clone size 2-3). Therefore, PCs in

PDAC likely arose from germinal center reactions in the PDAC stroma.

Antibodies from PCs binding to intracellular self-antigens

Forty-one antibodies with paired H and L chains from the six PDAC patients were selected for

recombinant antibody synthesis. These antibodies included the mostly expanded PC cell clones, as well as

some single-clone PC and B cells, and represented the products of 235 PCs and 12 B cells (Fig. 3A and

Table S2). Immunofluorescent staining of human PDAC cell lines and tumors was used to screen the

reactivity of the recombinant antibodies. Twenty-five of the 41 antibodies showed positive binding to

PDAC cell lines, with the expanded clones showing more frequent binding (Fig. 3A). Recombinant

antibodies reacted with antigens in all sub-cellular locations, including cytoplasmic (e.g. 8-3 and 15-7),

cytoplasmic-enriched (e.g. 19-1), both cytoplasmic and nuclear (e.g. 19-4), and nuclear-enriched

reactivities (e.g. 19-3 and 20-1) (Fig. 3B-C and Fig. S4). We did not identify any antibodies staining cell

surface antigens. Antibodies derived from a single patient's PDAC tumor could exhibit diverse staining

patterns (for example, 19-1 and 19-3), indicating the occurrence of an adaptive immune response to

multiple intracellular antigens in the same patient. All the reactive antibodies could stain multiple PDAC

cell lines with a similar staining pattern, as well as non-tumor human pancreatic ductal cell line, HPDE,

and human fibroblasts (Fig. 3D and Fig. S5). When staining PDAC tumors from which antibody

sequences were derived, the antibodies bound both to cancer cells and stromal cells (Fig. 3E). Thus, non-

mutated, non-cancer cell-specific intracellular antigens drive common humoral immune responses in

human PDAC.

Identification of antigens driving PC cell response in PDAC

Three PDAC antigens were identified by using antibody-mediated immunoprecipitation from MiaPaca2

cell lysates, followed by mass spectrometry analysis. Antibody 8-3, which was derived from the most

expanded PC clone (clone size 20) from patient 8, recognized a cortical structure in MiaPaca2 cells (Fig.

3C). This antibody immunoprecipitated a major protein band of approximately 45 kDa, which was

identified as ACTIN by mass spectrometry, as well as the ACTIN associated proteins, MYH10 and

MYH9 (Fig. 4A). Antibody 8-3 co-localized with filamentous actin (F-actin) in the cell cortex, and its

antigen re-localized to perinuclei foci after treatment with the actin destabilizing drug, cytochalasin D

(Fig. 4B). Knockdown of ACTIN with siRNA reduced staining by antibody 8-3, but not staining by a

MYH10-specific antibody (Fig. S6A), indicating that ACTIN was the target of antibody 8-3. This

antibody was confirmed to bind to polymerized F-actin, but not monomeric G-actin, using in vitro actin

polymerization and depolymerization assays (Fig. 4C). We developed an F-actin specific ELISA and

measured the EC_{50} of antibody 8-3 for F-actin to be 9.1 nM, a relatively high affinity/avidity that was

probably a result of SHM (Fig. 4D and Fig. 2E). Sequencing the actin genes (ACTB and ACTG1) from

the tumor organoid derived from patient 8, the source of the PCs encoding antibody 8-3, showed no

mutations in the protein-coding regions, confirming the self-antigen nature.

Similarly, antibody 19-3, the second most expanded PC clone in patient 19 (clone size 23),

immunoprecipitated a protein complex of RUVBL1 and RUVBL2, as determined by mass spectrometry.

Knockdown of RUVBL2 reduced antibody 19-3 binding in the MiaPaca2 cell line (Fig. 4F and Fig. S6B).

This antibody recognized recombinant RUVBL2 but not RUVBL1 protein, with an EC_{50} of 0.2 uM (Fig.

4G and Fig. S6C), confirming that antibody 19-3 recognizes RUVBL2.

Antibody 15-7, which was derived from a single PC clone in the tumor from patient 15, co-stained with

antibodies specific for the mitochondrial markers, COX4I1 and HSPA9 (Fig. S6D). This antibody

recognized a mitochondrial-enriched protein band with an apparent molecular weight of 60 kDa (Fig.

4H), which was identified as heat-shock protein D1 (HSPD1) by mass spectrometry. HSPD1 was

confirmed to be the antigen of antibody 15-7 by immunofluorescent co-localization in MiaPaca2 cells,

siRNA knockdown, and recombinant protein binding, with an EC_{50} of 7.9 nM (Fig. 4I-J and Fig. S6D).

In summary, our identification of three self-antigens, including antigens for two antibodies from highly

expanded PC clones, indicates that widely expressed, intracellular self-antigens may be targets of humoral

immune responses in human PDAC.

The frequent occurrence of antibody responses to F-actin and HSPD1 in PDAC patients

We collected plasma samples from 59 PDAC patients and 61 healthy donors, and measured IgG titers to

F-actin, RUVBL2, and HSPD1, respectively (Table S3). Demographics, such as gender, age, and race, did

not significantly affect antibody concentration. Interestingly, plasma from normal donors contained

detectable levels of IgG responses to these three antigens. In PDAC, the IgG F-actin and HSPD1 antibody

titers were significantly higher than those in normal individuals (Fig. 4K and Fig. S7A). IgG F-actin

antibody titers were significantly reduced in patients who had received neoadjuvant therapy, but this was

not observed for either HSPD1 or RUVBL2 antibody titers (Fig. S7B), suggesting that B cell responses

for different antigens may have different sensitivities to cytotoxic chemotherapy. These antibody

responses did not correlate with other pathological features such as tumor stage, size or grade. Thus,

PDAC patients likely promote auto-antibody responses which pre-exist in healthy donors.

DISCUSSION

PDAC is generally considered to be one of the more immunologically "silent" carcinomas and to be resistant to cancer immunotherapy for this reason. Reports, however, of TLSs in PDAC (*2-4*), of neoantigens with homology to infectious disease-derived peptides in long-term survivors of PDAC (*6*), and of signs of improved anti-PDAC immunity in metastatic lesions of patients with PDAC after inhibition of CXCR4 (*5*), all point to a need for a deeper understanding of the relationship between this cancer and the immune system. The antigens driving the B cell response are more readily identified by the Ig product of the B cells and PCs. Thus, we chose to interrogate the Ig products of intratumoral B/PCs in PDAC.

Our scRNA-Seq analysis of CD45+ immune cells in primary PDAC samples indicated active intratumoral immune reactions, such as the presence of effector CD8 T cells expressing *PRF1*, *GZMB*, and *IFNG*, and effector CD4 T cells expressing *TNF* and *IFNG*. We were able to confirm the four resected PDAC tumor samples as being microsatellite-stable, representing the majority of PDAC (*10*). The presence of effector T cells in our cohort is consistent with previous studies of human and mouse autochthonous PDAC immune microenvironments in which activated, effector T cells were found (*11*). In addition, scRNA-Seq identified CD4 T cells with the Tfh phenotype by their transcriptional expression of *BCL6* and *CXCR5*, and germinal center B cells expressing *BCL6*. The presence of a B cell response is further supported by Ig sequencing that identified isotypically switched, somatically hypermutated and expanded PC clones. The presence of TLS in PDAC stroma from the paired histology images further support the integrated B and T cell responses. It is interesting that these indicators of an active adaptive immune reaction are occurring despite the presence of large proportion of FOXP3-expressing regulatory T cells.

There are a few previous studies exploring the antigen reactivity of B cells and PCs in other tumors (*12, 13*), but our study is the first to apply this approach to PDAC. While we were unsuccessful in our initial intent to discover tumor-specific B cell antigens, the finding of Igs from PCs frequently binding to widely expressed, non-mutated intracellular self-antigens in human PDAC is intriguing. Our screening in

PDAC identified most antibodies as recognizing intracellular antigens that were shared by multiple

PDAC cell lines, normal epithelial cells and fibroblasts. This conclusion was confirmed by the

identification of three self-antigens, F-actin, RUVBL2 and HSPD1, as targets of these humoral immune

responses. All the seven PDAC patients enrolled for this study have no records of auto-immune disease or

pancreatitis. In this context, it is worth noting that 20% of naïve B cells in human peripheral blood are

reported to be autoreactive (*14*), which is consistent with our detection of autoantibody responses to those

antigens in healthy donors. Self-reactive B cell responses are likely promoted in PDAC because of

continual exposure of intracellular self-antigens through cell death and chronic inflammation, reminiscent

of the antibody responses in autoimmune diseases. Autoantibody responses to F-actin, RUVBL2 and

HSPD1 have been reported in various autoimmune conditions (*15-17*) and cancers (*18, 19*). Previously,

immunization with mouse syngeneic ES cells reduce mouse pancreatic tumor progression, indicating self-

antigens likely drive such immune response (*7, 8*). Additionally, a recent study reported that the self-

protein matrix metallopeptidase 14 was a major auto-antigen in human ovarian cancer, and some Igs were

likely originated from germline encoded auto-reactive B cell (*13*). Thus, the B cell response to self-

antigens may be a common feature of cancer.

Our analysis of plasma titer to F-actin, RUVBL2 and HSPD1 did reveal that PDAC patients have

significantly higher titers to F-actin and HSPD1 as compared to a group of healthy donors. Our analysis

did not reveal a significant correlation with tumor-associated pathological factors, although this may be

partially due to our relatively small cohort of patients. Our studies can be potentially strengthened by

analyzing serial tumor biopsies and plasma obtained during the course of tumor progression or response

to therapy such as CXCR4 inhibition, thus providing more dynamic pictures of B cell response and

antibody titer changes.

One open question from this study is whether the B cell response to intracellular self-antigens affects

PDAC progression. Given the generally positive correlation of a B cell response with a favorable outcome

in PDAC (*2-4*), we speculate that those autoreactive B cell responses may indirectly have an anti-tumor

role. While antibodies binding intracellular antigens cannot directly target live tumor cells, these antigens

can be exposed to antibody binding during cancer cell death, such as necrosis, and potentially induce

inflammation through complement activation and Fc receptor binding. Moreover, we expect that B cells

binding such self-antigens will internalize and engage in antigen presentation to CD4+ T cells. F-actin has

been reported to promote antigen presentation through the F-actin receptor Clec9A on dendritic cells in

breast cancer model (20). Further understanding the B cell response in PDAC may potentially provide

opportunities for modulation of cancer immunotherapy. In any event, the finding of T cell and B cell

responses in the PDAC tumors indicates that an adaptive immune response is occurring in the tumor

microenvironment, which may be exploited for tumor control.

MATERIALS AND METHODS

Study design

This study is designed to study B cell response and identify B cell targeted antigens in human PDACs.

This study consists of :1) ScRNA-Req of immune cells infiltrating human PDAC and Igs of B and

plasma cells; 2) Synthesizing of recombinant antibodies from Ig sequencing; 3) Screening of recombinant

antibodies reactivities toward PDAC cell lines and tumor tissues; 4) Identification of antigens. This

pipeline was used to study four resected and three FNA naïve-treated PDAC samples. The antibodies

synthesizing and screening aim to profile the most expanded plasma cells clones in PDAC to get a better

picture of B cell reactivity in human PDAC.

Human samples

All human samples were obtained with written consent and IRB approval. Four resected PDAC

specimens and three FNA (two passes each) specimens were obtained from Northwell Health. Archived

plasma samples from PDAC patients (n=59) were received from Northwell Health. Plasma samples from

healthy donors (n=61) were obtained from volunteer donors or purchased from commercial resources

(Innovative Research and BioChemed Services). Microsatellite instability status of resected PDAC

samples were determined by immunohistochemistry staining of MLH1, MSH2, MSH6 and PMS2,

performed and interpreted by pathologists from Northwell Health. Histopathologic hematoxylin and eosin

images were obtained from Northwell pathology department acquired by Leica Aperio slide scanner. All

patient samples were de-identified.

Cell lines and culture

264   Human PDAC cell lines MiaPaca2, Suit2, Panc-1, and foreskin fibroblast (BJ-hTert) were obtained from

265   Cold Spring Harbor Laboratory cell validation center, and had been validated by STR profiling. The

266   normal human pancreatic ductal epithelial cell line (HPDE) was obtained from Dr. Krainer lab in CSHL,

267   originally purchased from Kerafast. All cell lines had been tested mycoplasma free and cultured in the

268   conditions recommended by ATCC or the manufacturer.

Tumor samples preparation and Single-cell sequencing

A small portion of the resection tumor was snap-frozen for histology. The remaining human PDAC tumor

samples were digested into single cells using collagenases and about 10% of single cells digest was used

for tumor organoid culture as described (*21*). Live immune cells were sorted by $EpCAM^-$ /$CD45^+$

(Biolegend, # clone 2D1 and 9C4) on a Sony SH800 cell sorter. About 10,000 immune cells were

barcoded with 10X Genomics Chromium single cell 5' kit, and both gene expression and Ig libraries were

prepared and sequenced according to the 10X Genomics manual.

Sequencing data analysis

The scRNA-Seq and Ig reads were aligned and quantified using the Cell Ranger pipeline (10X

Genomics). Dead cells and cells with the number of genes expressed less than 200 were also filtered out.

ScRNA-Seq Data was normalized, logarithm transformed and scaled. Doublets were removed using a R

package 'Scrublet'. Principle component analysis (PCA) was run with 50 components using the top 4000

genes of each sample. The nearest neighbor algorithm was run with 50 PCAs and clustered in UMAP

projections using Leiden clustering. All additional analyses were performed using Loupe Browser (10X

Genomics) and the Python toolkit Scanpy. Immune cells clusters were defined using the set of markers

and UMAP clusters: T cells ($CD3^+$, TCR^+, $NCAM^-$), NK cells ($CD3^-$, $NCAM^+$), B cells ($CD20^+$, $CD79A^+$,

$IgKC^{low}$), PCs ($JChain^{hi}$, $IgKC^{hi}$), myeloid cells ($CD14^+$, $CSF1R^+$), mast cells ($CPA3^+$, $GATA2^+$),

fibroblasts (FAP^+, $ACTA2^+$, $PDGFRA1^+$), epithelial cells ($CK19^+$, $CK18^+$, $CK5^+$, $CK14^+$).

Ig data analysis

PCs and B cells clusters defined by scRNA-Seq were used to select cell type specific Igs analysis. In one

patient (Pt-15), gene expression sequencing data was not available, the PCs were defined by high

expression of H chain (UMI>=100). B or PC specific Igs were exported from the Loupe V(D)J Browser

(10X Genomics) and analyzed for isotypes and clonotypes. For somatic hypermutation analysis, V region

sequences were analyzed by Igblast (NCBI) and visualized by Graph Prism. For lineage tree analysis, Ig

sequences were first analyzed using R packages Change-O and alakazam (22), and the Ig lineages were

built using a R package PHYLIP. Finally, Lineage trees were visualized using a R package igraph.

Recombinant Antibodies production

The selected antibody V(D)J sequences were synthesized from IDT, and cloned into pFUSEss H and L

vectors (Invivogen) using NEB HIFI assembly kit (NEB, #5520s). The heavy chains were cloned into

human IgG1 or customer-designed human IgG4-His vector. A 6x His tag with a Gly-Ser-Gly linker was

inserted before the stop codon of human IgG4 constant region, synthesized and replaced the IgG1 in

pFUSE-CHIg-hG1 vector. The light chain was cloned into a vector containing IgKC, IgLC2, or

costumed-made IgLC1 vector, to match the original light chain. Sequences confirmed antibody vectors

were mixed at H:L 1:1.5 ratio, and transiently transfected into 293T cells, cultured in 10% IgG reduced

serum. Culture supernatants were collected four days later and antibodies were purified using the Protein

G column (Thermo Fisher, #89956). The antibodies were further concentrated and the buffer was

exchanged into PBS using Amicon 50K centrifugal filter device (Sigma, #UFC805096). The antibody

concentration was quantified by NanoDrop (Thermo Fisher) and human IgG ELISA kit (Mabtech, #3850-

1H-6).

Antibody screening using immunofluorescent staining

Cells were seeded in the 24-well glass-bottom cell culture plate (Chemglass, #CLS-1812-024), and used

2-3 days later at 50-80% confluency. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton-

X100, blocked with 3% FBS, and stained with 10 ug/ml human query antibodies diluted in PBST

containing 1% FBS overnight at 4 °C. Cells were then washed three times with PBST, and stained with

fluorescent conjugated secondary anti-human IgG antibody (Thermo Fisher, #A21090 and #A-11013) and

DNA dye DAPI. In human tumor section staining, anti-His secondary antibody (Biolegend, #362607) was

used to detect IgG4-His primary antibodies. Other antibodies and dyes used were: Phalloidin (Thermo

Fisher, #A12379), MYH10 (Atlas Antibodies, #HPA047541), cytochalasin D (Thermo Fisher,

#PHZ1063), ACTIN (Cell Signaling Technology, #3700S), RUVBL2 (Atlas Antibodies, #HPA067966),

RUVBL1 (Atlas Antibodies, # HPA019947), HSPD1 (Atlas Antibody, #HPA050025), HSPA9 (Atlas

Antibody, #HPA000898), COX4I1 (Atlas antibody, #HPA002485), KRT19 (Abcam, #ab203445), human

IgG isotype controls (Biolegend, #403502 and #403702). The samples were imaged in Leica SP8

confocal microscope under 40x magnification. Antibody screening images were acquired using the same

confocal setting for isotype control staining, and the gain of query antibodies was reduced if images were

saturated in the setting. Images were analyzed using Leica LAS X or Image J software.

Immunoprecipitation and mass spectrometry

Antibodies were covalently crosslinked to magnetic beads using the Thermo Fisher Dynabeads Antibody

Coupling Kit. The total cell lysate was prepared from MiaPaca2 cell line using TNET lysis buffer (50 mM

Tris-HCl, pH 7.5, 150 mM NaCl, 5 mM EDTA and 1% Triton X-100). Dynabeads containing 2-5 ug

crosslinked antibodies or isotype control were first blocked with 1%BSA, then incubated with up to 1mg

cell lysate overnight in 4C in TNET buffer. The beads were separated using a magnetic rack, washed

three times with TNET buffer (or with increased NaCl concentration), and eluted with 2X SDS under

boiling. The immunoprecipitation elute was running in gradient SDS-PAGE gel and stained with a silver

staining kit (Thermo Fisher, #24600).

Distinct gel bands were excised, de-stained, reduced with 3 mM TCEP and alkylated with 10 mM

CEMTS, and then digested with trypsin. Eluting peptides were ionized and transferred into an Exploris

Orbitrap mass spectrometer (Thermo Fisher). Spectral data were searched against the human database and

a database of common contaminants. M-oxidation and N/Q-deamidation were set as variable

modifications. Peptide-spectral matches were filtered to maintain FDR <1% using the Percolator.

siRNA knockdown

Cells were transfected with control or siRNA at 20-60uM, cultured for three days, and proceeded for

immunofluorescent staining. siRNA against human ACTB (#SASI_Hs01_00204238,

#SASI_Hs01_00204239), MYH10 (#SASI_Hs01_00072460, #SASI_Hs02_00340636), MYH9

(#SASI_Hs01_00197338, #SASI_Hs01_00197339), RUVBL2 (GATGATTGAGTCCCTGACCAA,

GAAGATGTGGAGATGAGTGAG), control (GGATGTAAGTGGGAAAGTGGA) were purchased

from Sigma. siRNA targeting HSPD1 (L-010600-00-0005) was purchased from Horizon Discovery.

Actin polymerization or de-polymerization and dot-blot assay

Human non-muscle actin (Cytoskeleton, # APHL99) was polymerized or de-polymerized according to the

manufacturer's protocol. Briefly, actin was first diluted into 0.4 mg/ml in G-buffer (5 mM Tris-HCl, 0.2

mM CaCl$_2$, 0.2 mM ATP), and kept in ice for 1h. For polymerization, 10% polymerization buffer (500

mM KCl, 20 mM MgCl2) with 1 mM final ATP and 5uM phalloidin (Simga, #P2141) was added to the

actin and incubated at room temperature for 1h. For de-polymerization, 5 uM latrunculin B (Sigma,

#428020) was added into the actin with G-buffer, and incubated in ice for 1h. Actin samples were further

diluted into G-buffer or polymerization buffer supplemented with phalloidin or Latrunculin B, and serial

dilutions of F-actin or G-actin were blotted on nitrocellulose membrane using Bio-Dot microfiltration

apparatus (Bio-Rad) according to the manufacturer's instructions. The membrane was furthered blocked

and incubated with human IgG4 isotype control, Ab 8-3, mouse anti-human Actin antibody (Thermo

Fisher, # MA1-140) at 1 ug/ml for 1h, washed and detected with HRP conjugated anti-human IgG

(Thermo Fisher, #A18811) or anti-mouse IgG (Biolegend, #405306) secondary antibody, and developed

by ECL substrate.

ELISA

Recombinant proteins RUVBL2 (Creative Biomart, #RUVBL2-30950), RUVBL1 (Novus, #NBP1-

50845), HSPD1 (Origene, #TP760396), or BSA control were coated in PBS with 0.25 ug per well in 96-

well ELISA plate (Corning) overnight in 4C. The plate was blocked with 1% BSA, washed with PBST,

incubated with query antibodies or isotype control at 1 ug/ml diluted in 0.1% BSA in PBST for 1h,

detected with HRP conjugated secondary antibody (1ug/ml), and developed by TMB substrate

(Biolegend, #421101) for 5-10 min. For F-actin ELISA, polymerized F-actin was diluted in

polymerization buffer supplemented with 5uM phalloidin and 1mM ATP, and coated in the ELISA plate

for 1h at room temperature, followed by antibody binding and development.

For human plasma studies, the antigens were coated as above, then blocked with 5% milk in PBS.

PDAC or healthy donor plasma was diluted 1/100 and 1/1000 in 0.5% milk in PBST, and each sample

was run in duplicates. Positive controls are using antibodies 8-3, 19-3 or 15-7. Background signal was

obtained using the higher signal from secondary antibody only and plate coated with BSA. All samples to

the same antigen were analyzed at the same time to avoid batch differences. Each ELISA analysis was

repeated twice.

Mitochondria cell fraction and western blot

For crude mitochondria fraction, the cells were first lysed using low salt buffer (0.25 M sucrose, 20 mM

HEPES-KOH pH 7.5, 10 mM KCl, 1.5 mM MgCl2, 1 mM EDTA, 1 mM EGTA, 1 mM DTT)

supplemented with proteinase/phosphatase inhibitors, homologized with a Teflon-glass homogenizer, and

then centrifuged at 700 g for 10 min in 4C to collect the cytosol fraction in the supernatant. The cytosol

fraction was further centrifuged at 10,000 g for 15 min to collect the mitochondria fraction in the pellet.

Same amounts of SDS-reduced cytosol or mitochondria fractions were run in western blot, and detected

with antibody 15-7, HSPA9 and GAPDH (Cell Signaling Technology, #97166). A separate sample gel

was silver stained and cut around 60kd for mass spectrometry analysis.

Histopathology image analysis

The scanned resected PDAC histopathologic images were opened with ImageScope (Leica), and

lymphocytes aggregates with bigger than 0.01 mm^2 size were counted and exported for size

measurement using image J.

Statistical analysis

Statistical analysis was performed using Graphpad (Prism). The plasma IgG data was first tested for

normality distribution in Graphpad. Mann-Whitney t-test was used for a two-group comparison of non-

normal distribution data, Kruskal-Wallis ANOVA test was used for comparisons of three or more groups

of non-normal distribution data. Fisher's test was used for demographic factors comparison, and

Pearson's correlation was used for correlation studies. P<0.05 was used for statistical significance.

List of Supplementary Materials

Fig. S1 Characterization of primary PDAC immune cells microenvironment by scRNA-Seq

Fig. S2. Presence of TLS like structures in PDAC.

Fig. S3 Additional Ig sequences evolution trees from PCs in PDAC

Fig. S4 Antibodies staining atlas on MiaPaca2 cell line.

Fig. S5 Examples of antibodies staining in PDAC and non-cancer cell lines

Fig. S6 Characterization of antibodies 8-3, 19-3 and 15-7

Fig. S7 Comparison of plasma IgG titers to F-actin, RUVBL2 and HSPD1 between healthy donors and

PDAC patients.

Table S1 Human PDAC samples information for scRNA-Seq studies.

Table S2 List of antibodies synthesized and screening results.

Table S3 Human samples information for plasma studies.

References and Notes

1.   A. S. Bear, R. H. Vonderheide, M. H. O'Hara, Challenges and Opportunities for Pancreatic Cancer
Immunotherapy. *Cancer Cell* **38**, 788-802 (2020).
2.   N. Hiraoka, Y. Ino, R. Yamazaki-Itoh, Y. Kanai, T. Kosuge, K. Shimada, Intratumoral tertiary
lymphoid organ is a favourable prognosticator in patients with pancreatic cancer. *Br J Cancer*
112, 1782-1790 (2015).
3.   G. F. Castino, N. Cortese, G. Capretti, S. Serio, G. Di Caro, R. Mineri, E. Magrini, F. Grizzi, P.
Cappello, F. Novelli, P. Spaggiari, M. Roncalli, C. Ridolfi, F. Gavazzi, A. Zerbi, P. Allavena, F.
Marchesi, Spatial distribution of B cells predicts prognosis in human pancreatic adenocarcinoma.
Oncoimmunology **5**, e1085147 (2016).
4.   J. G. A, V. Rajamanickam, C. Bui, B. Bernard, J. Pucilowska, C. Ballesteros-Merino, M. Schmidt, K.
McCarty, M. Philips, B. Piening, C. Dubay, T. Medler, P. Newell, P. Hansen, E. Tran, E. Tang, C.
Bifulco, M. Crittenden, M. Gough, K. H. Young, Germinal center reactions in tertiary lymphoid
structures associate with neoantigen burden, humoral immunity and long-term survivorship in
pancreatic cancer. *Oncoimmunology* **10**, 1900635 (2021).
5.   D. Biasci, M. Smoragiewicz, C. M. Connell, Z. Wang, Y. Gao, J. E. D. Thaventhiran, B. Basu, L.
Magiera, T. I. Johnson, L. Bax, A. Gopinathan, C. Isherwood, F. A. Gallagher, M. Pawula, I.
Hudecova, D. Gale, N. Rosenfeld, P. Barmpounakis, E. C. Popa, R. Brais, E. Godfrey, F. Mir, F. M.
Richards, D. T. Fearon, T. Janowitz, D. I. Jodrell, CXCR4 inhibition in human pancreatic and

colorectal cancers induces an integrated immune response. *Proc Natl Acad Sci U S A* **117**, 28960-
28970 (2020).
6. V. P. Balachandran, M. Luksza, J. N. Zhao, V. Makarov, J. A. Moral, R. Remark, B. Herbst, G.
Askan, U. Bhanot, Y. Senbabaoglu, D. K. Wells, C. I. O. Cary, O. Grbovic-Huezo, M. Attiyeh, B.
Medina, J. Zhang, J. Loo, J. Saglimbeni, M. Abu-Akeel, R. Zappasodi, N. Riaz, M. Smoragiewicz, Z.
L. Kelley, O. Basturk, I. Australian Pancreatic Cancer Genome, R. Garvan Institute of Medical, H.
Prince of Wales, H. Royal North Shore, G. University of, H. St Vincent's, Q. B. M. R. Institute, C. f.
C. R. University of Melbourne, I. f. M. B. University of Queensland, H. Bankstown, H. Liverpool, C.
O. B. L. Royal Prince Alfred Hospital, H. Westmead, H. Fremantle, H. St John of God, H. Royal
Adelaide, C. Flinders Medical, P. Envoi, H. Princess Alexandria, H. Austin, I. Johns Hopkins
Medical, A. R.-N. C. f. A. R. o. Cancer, M. Gonen, A. J. Levine, P. J. Allen, D. T. Fearon, M. Merad,
S. Gnjatic, C. A. Iacobuzio-Donahue, J. D. Wolchok, R. P. DeMatteo, T. A. Chan, B. D. Greenbaum,
T. Merghoub, S. D. Leach, Identification of unique neoantigen qualities in long-term survivors of
pancreatic cancer. *Nature* **551**, 512-516 (2017).
7. N. G. Kooreman, Y. Kim, P. E. de Almeida, V. Termglinchan, S. Diecke, N. Y. Shao, T. T. Wei, H. Yi,
D. Dey, R. Nelakanti, T. P. Brouwer, D. T. Paik, I. Sagiv-Barfi, A. Han, P. H. A. Quax, J. F. Hamming,
R. Levy, M. M. Davis, J. C. Wu, Autologous iPSC-Based Vaccines Elicit Anti-tumor Responses In
Vivo. *Cell Stem Cell* **22**, 501-513 e507 (2018).
8. X. Ouyang, Y. Liu, Y. Zhou, J. Guo, T. T. Wei, C. Liu, B. Lee, B. Chen, A. Zhang, K. M. Casey, L.
Wang, N. G. Kooreman, A. Habtezion, E. G. Engleman, J. C. Wu, Antitumor effects of iPSC-based
cancer vaccine in pancreatic cancer. *Stem Cell Reports* **16**, 1468-1477 (2021).
9. R. Di Niro, L. Mesin, N. Y. Zheng, J. Stamnaes, M. Morrissey, J. H. Lee, M. Huang, R. Iversen, M. F.
du Pre, S. W. Qiao, K. E. Lundin, P. C. Wilson, L. M. Sollid, High abundance of plasma cells
secreting transglutaminase 2-specific IgA autoantibodies with limited somatic hypermutation in
celiac disease intestinal lesions. *Nat Med* **18**, 441-445 (2012).
10. C. Luchini, L. A. A. Brosens, L. D. Wood, D. Chatterjee, J. I. Shin, C. Sciammarella, G. Fiadone, G.
Malleo, R. Salvia, V. Kryklyva, M. L. Piredda, L. Cheng, R. T. Lawlor, V. Adsay, A. Scarpa,
Comprehensive characterisation of pancreatic ductal adenocarcinoma with microsatellite
instability: histology, molecular pathology and clinical implications. *Gut* **70**, 148-156 (2021).
11. E. Elyada, M. Bolisetty, P. Laise, W. F. Flynn, E. T. Courtois, R. A. Burkhart, J. A. Teinor, P. Belleau,
G. Biffi, M. S. Lucito, S. Sivajothi, T. D. Armstrong, D. D. Engle, K. H. Yu, Y. Hao, C. L. Wolfgang, Y.
Park, J. Preall, E. M. Jaffee, A. Califano, P. Robson, D. A. Tuveson, Cross-Species Single-Cell
Analysis of Pancreatic Ductal Adenocarcinoma Reveals Antigen-Presenting Cancer-Associated
Fibroblasts. *Cancer Discov* **9**, 1102-1123 (2019).
12. J. DeFalco, M. Harbell, A. Manning-Bog, G. Baia, A. Scholz, B. Millare, M. Sumi, D. Zhang, F. Chu,
C. Dowd, P. Zuno-Mitchell, D. Kim, Y. Leung, S. Jiang, X. Tang, K. S. Williamson, X. Chen, S. M.
Carroll, G. Espiritu Santo, N. Haaser, N. Nguyen, E. Giladi, D. Minor, Y. C. Tan, J. B. Sokolove, L.
Steinman, T. A. Serafini, G. Cavet, N. M. Greenberg, J. Glanville, W. Volkmuth, D. E. Emerling, W.
H. Robinson, Non-progressing cancer patients have persistent B cell responses expressing
shared antibody paratopes that target public tumor antigens. *Clin Immunol* **187**, 37-45 (2018).
13. R. D. Mazor, N. Nathan, A. Gilboa, L. Stoler-Barak, L. Moss, I. Solomonov, A. Hanuna, Y. Divinsky,
M. D. Shmueli, H. Hezroni, I. Zaretsky, M. Mor, O. Golani, G. Sabah, A. Jakobson-Setton, N.
Yanichkin, M. Feinmesser, D. Tsoref, L. Salman, E. Yeoshoua, E. Peretz, I. Erlich, N. M. Cohen, J.
M. Gershoni, N. Freund, Y. Merbl, G. Yaari, R. Eitan, I. Sagi, Z. Shulman, Tumor-reactive
antibodies evolve from non-binding and autoreactive precursors. *Cell* **185**, 1208-1222 e1221
(2022).
14. H. Wardemann, S. Yurasov, A. Schaefer, J. W. Young, E. Meffre, M. C. Nussenzweig, Predominant
autoantibody production by early human B cell precursors. *Science* **301**, 1374-1377 (2003).

15. A. Granito, L. Muratori, P. Muratori, G. Pappas, M. Guidi, F. Cassani, U. Volta, A. Ferri, M. Lenzi,
F. B. Bianchi, Antibodies to filamentous actin (F-actin) in type 1 autoimmune hepatitis. *J Clin*
Pathol **59**, 280-284 (2006).
16. K. Kaji, N. Fertig, T. A. Medsger, Jr., T. Satoh, K. Hoshino, Y. Hamaguchi, M. Hasegawa, M. Lucas,
A. Schnure, F. Ogawa, S. Sato, K. Takehara, M. Fujimoto, M. Kuwana, Autoantibodies to RuvBL1
and RuvBL2: a novel systemic sclerosis-related antibody associated with diffuse cutaneous and
skeletal muscle involvement. *Arthritis Care Res (Hoboken)* **66**, 575-584 (2014).
17. G. Schett, Q. Xu, A. Amberger, R. Van der Zee, H. Recheis, J. Willeit, G. Wick, Autoantibodies
against heat shock protein 60 mediate endothelial cytotoxicity. *J Clin Invest* **96**, 2569-2577
(1995).
18. M. H. Hansen, H. Nielsen, H. J. Ditzel, The tumor-infiltrating B cell response in medullary breast
cancer is oligoclonal and directed against the autoantigen actin exposed on the surface of
apoptotic cancer cells. *Proc Natl Acad Sci U S A* **98**, 12659-12664 (2001).
19. Y. He, Y. Wu, Z. Mou, W. Li, L. Zou, T. Fu, A. Zhang, D. Xiang, H. Xiao, X. Wang, Proteomics-based
identification of HSP60 as a tumor-associated antigen in colorectal cancer. *Proteomics Clin Appl*
1, 336-342 (2007).
20. E. Giampazolias, O. Schulz, K. H. J. Lim, N. C. Rogers, P. Chakravarty, N. Srinivasan, O. Gordon, A.
Cardoso, M. D. Buck, E. Z. Poirier, J. Canton, S. Zelenay, S. Sammicheli, N. Moncaut, S. Varsani-
Brown, I. Rosewell, C. Reis e Sousa, Secreted gelsolin inhibits DNGR-1-dependent cross-
presentation and cancer immunity. *Cell* **184**, 4016-4031 e4022 (2021).
21. S. F. Boj, C. I. Hwang, L. A. Baker, Chio, II, D. D. Engle, V. Corbo, M. Jager, M. Ponz-Sarvise, H.
Tiriac, M. S. Spector, A. Gracanin, T. Oni, K. H. Yu, R. van Boxtel, M. Huch, K. D. Rivera, J. P.
Wilson, M. E. Feigin, D. Ohlund, A. Handly-Santana, C. M. Ardito-Abraham, M. Ludwig, E. Elyada,
B. Alagesan, G. Biffi, G. N. Yordanov, B. Delcuze, B. Creighton, K. Wright, Y. Park, F. H. Morsink, I.
Q. Molenaar, I. H. Borel Rinkes, E. Cuppen, Y. Hao, Y. Jin, I. J. Nijman, C. Iacobuzio-Donahue, S. D.
Leach, D. J. Pappin, M. Hammell, D. S. Klimstra, O. Basturk, R. H. Hruban, G. J. Offerhaus, R. G.
Vries, H. Clevers, D. A. Tuveson, Organoid models of human and mouse ductal pancreatic
cancer. *Cell* **160**, 324-338 (2015).
22. N. T. Gupta, J. A. Vander Heiden, M. Uduman, D. Gadala-Maria, G. Yaari, S. H. Kleinstein,
Change-O: a toolkit for analyzing large-scale B cell immunoglobulin repertoire sequencing data.
Bioinformatics **31**, 3356-3358 (2015).
503
504

Acknowledgments:

We thank Sharon Fox and Cristina Valente and other members from Northwell Health Biospecimen

Repository in helping coordinate the human sample collections. We thank Ledong Wan from Dr. Krainer

lab at CSHL in providing the HPDE cell line. We thank Dr. Semir Bayez of CSHL for his thoughtful

input in this project. We thank members of Fearon lab for their thoughtful discussions of this project and

help review of this manuscript. This project was funded by Lustgarten Foundation and Simons

Foundation awarded to Dr. Douglas Fearon.

Funding:

Lustgarten Foundation for Pancreatic Research (DF)

Simons Foundation (DF)

Author contributions:

Conceptualization: MY, DF

Methodology: MY, JP, JY, DP, PC, YZ, BH, HP, AH, DK, KP, AR, DS, MW, DT

Investigation: MY, SS, PM,

Visualization: MY, JP, YZ, BH

Funding acquisition: DF

Project administration: MY

Supervision: DF

Writing – original draft: MY, DF

Writing – review & editing: YZ, PM, KP, AR

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: The sequencing data from this study will be deposited into the NCBI

database. Public codes/software was used for the data analysis as described in the paper. The codes for

sequencing data analysis and experimental material can be accessed through communication with the

corresponding authors.

 Figures

Fig.1. Identification of active T and B cell responses in PDAC by scRNA-Seq. (**A**) UMAP clusters are

shown of immune cells isolated from six primary PDAC. (**B**) The cell numbers of different immune cell

types identified in individual patient are shown. (**C**) The expression of selected genes in T cell, B cell and

PC clusters are shown. The intensity of the colors represents the normalized log2 transformation of UMI.

(**D**) Examples of TLS like structures (indicated by arrowhead and labelled with 1,2 and 3) in patient-19.

The zoom in examples of TLS were shown in the right. Scale bar represents 600um.

550
Fig.2. Identification of isotype switched and somatically hypermutated Igs by sequencing of PCs

from PDAC. (A) Numbers are shown of PCs in which paired H and L chains were sequenced in each

patient. The distribution of Ig isotypes (**B**) and the number of somatic mutations in V regions (**C**) in PCs

from individual patients are determined. (**D**) The PCs with paired H and L chains and their clonotype

distribution are shown for each patient. PCs within expanded clones are labelled with colored doughnuts,

with colors indicating isotypes. Each slice represents one clone, with clone size proportional to slice size.

Single clone PCs are pooled and labeled with gray color. (**E**) Examples are presented of antibody lineage

evolution within expanded PC clones for individual patients. Clone size is indicated by the size of the

node (not scaled) and labeled by numbers on the right. The number of somatic mutations in the combined

V(D)J regions is shown in parenthesis.

Fig.3. Recognition of intracellular self-antigens by recombinant antibodies from expanded PCs of

PDAC tumors. (**A**) Summary is presented of the results of staining various cell lines by 41 recombinant

antibodies based on scRNA-Seq of PCs and B cells from six PDAC tumors, distributed according to clone

sizes. (**B-C**) The distribution and examples are shown of four distinct antibody staining patterns:

cytoplasmic (Cyto, e.g. 8-3 and 15-7), cytoplasmic enriched (Cyto enriched. e.g. 19-1), evenly distributed

in cytoplasmic and nuclei (Cyto/nuclei even, e.g. 19-1), and nuclei enriched (e.g. 19-3 and 20-1).

Examples of positive antibody staining (red) in MiaPaca2 cell line, co-stained with DAPI (blue). Original

antibody isotype and clone size (cs) are indicated. (**D**) The recombinant antibodies, 8-3 and 19-3 (red),

respectively, were used for staining different PDAC and non-cancer cell lines, along with nuclear staining

with DAPI (blue). (**E**) Antibodies 8-3 and 19-3, respectively, were used to stain the PDAC tumors from

which their PCs were derived. Staining was also done with anti-KRT19 antibody to demonstrate the

cancer cells. Scale bar in C, 25um, and in D and E, 50um.

Fig.4. Identification of F-actin, RUVBL2 and HSPD1 as antigens in PDAC. (A)

Proteins that were immunoprecipitated by antibody 8-3 from lysates of MiaPaca2 cells were

separated by SDS-PAGE and visualized by silver staining. Two bands were identified by

mass spectrometry (red arrowheads). (**B**) MiaPaca2 cells were co-stained with 8-3,

phalloidin, and an antibody specific for MYH10 after 30min treatment of the cells with

DMSO or cytochalasin D. (**C**) Dot-blot assay is shown the reaction with F-actin or G-actin

of 8-3, isotype control IgG, and a commercial actin specific antibody. (**D**) The binding of

incremental concentrations of 8-3 to F-actin was measured by ELISA. (**E**) Proteins that

were immunoprecipitated by recombinant antibody 19-3 from lysates of MiaPaca2 cells

were separated by SDS-PAGE and visualized by silver staining. The proteins, RUVBL1

and RUVBL2, were identified by mass spectrometry. (**F**) MiaPaca2 cell were co-stained

with 19-3 and an antibody specific for RUVBL2. (**G**) The binding of incremental

concentrations of 19-3 to RUVBL2 was measured by ELISA. (**H**) Western blot analysis of

recombinant antibody 15-7 detection of cytoplasmic proteins (Cyto) and mitochondria-

enriched proteins (Mito) is shown. (**I**) Immunofluorescent staining of MiaPaca2 cells by 15-

7 and an antibody specific for HSPD1 is shown. (**J**) The binding of incremental

concentrations of 15-7 to HSPD1 was measured by ELISA. (**K**) The serum IgG titers of

normal individuals and PDAC patients to F-actin (sera diluted 1:100), to RUVBL2 (sera

diluted 1:1000), and to HSPD1(sera diluted 1:1000), respectively, were measured by

ELISA. The non-parametric t-test was used for comparison. Scale bars in **B**, **F** and **I** are

50um.

Title: Plasma cells in human pancreatic ductal adenocarcinoma secrete antibodies to self-

antigens

Authors: Min Yao[1]†, Jonathan Preall[1], Johannes Yeh[1], Darryl Pappin[1], Paolo Cifani[1], Yixin Zhao[1],

Sophia Shen[2], Philip Moresco[1,3,4], Brian He[1], Hardik Patel[1], Amber Habowski [1], Daniel A. King[5], Kara

Raphael[5], Arvind Rishi[5], Divyesh Sejpal[5]‡, Matthew Weiss[5], David Tuveson[1], Douglas Fearon[1,6*]

630
631 **Supplementary materials**

Supplementary figures

Fig. S1. Characterization of primary PDAC immune cells microenvironment by scRNA-Seq. (A)

Heatmap of the top five most upgraded genes in each cell cluster defined in Figure 1A. Color scale is log2

fold changes. (**B-E**) The percentage of cells expressing selected gene markers in the clusters of CD8 T

(**B**), CD4 (**C, D**) and B cells (**E**).

665   **Fig. S2. Presence of TLS like structures in PDAC. (A)** Examples of TLS like structures in patient 8, 19

666   and 20 (TLS indicated by arrowhead and labelled by numbers). The zoom in images were shown in the

667   right. Scale bar is shown in figures. (**B**) Quantification and size distribution of the TLS in the four

668   resected PDAC samples. Only TLS with size bigger than 0.01 mm^2 were analyzed.

Fig. S3. Additional Ig sequences evolution trees from PCs in PDAC. Additional examples of antibody

lineage evolution among top expanded PC clones screened in this study. Clone size is indicated by the

673   size of the node (not scaled) and labeled by numbers on the right. The clone size includes both paired and

674   unpaired chains. Number of somatic mutations in combined V(D)J regions is shown in parenthesis.

Fig. S4. Antibodies staining atlas on MiaPaca2 cell line. MiaPaca2 cells were stained with each

antibody (red) and co-stained with DAPI (blue). Secondary antibody only (2nd Ab) and isotype IgG were

used as control. Antibodies with positive staining are labeled green, and antibodies with negative staining

are labeled white. All images were acquired using the same confocal setting for isotype control staining,

and the gain on the query antibody was reduced if the images were saturated. Scale bar is 50um.

Fig. S5. Examples of antibodies staining in PDAC and non-cancer cell lines. Cells were stained with

each antibody (red) and co-stained with DAPI (blue). Scale bar is 50um.

Fig. S6. Characterization of antibodies 8-3, 19-3 and 15-7. (**A**) 8-3 staining after siRNA knockdown of

ACTB in MiaPaca2 cells, co-stained with phalloidin, MYH10 antibody and DAPI. (**B**) Measurement of

19-3 binding to recombinant RUVBL1 and RUBL2 proteins by ELISA. Positive controls using

polyclonal rabbit antibodies anti-RUVBL1 (a-RUVBL1) or anti-RUVBL2 (a-RUVBL2). (**C**) 19-3

staining after siRNA knockdown of *RUVBL2* in MiaPaca2 cells, co-stained with anti-RUVBL2 rabbit

antibody and DAPI. (**D**) Co-staining of 15-7 with mitochondria markers COX4I1, HSPA9 and HSPD1 in

MiaPaca2 cells. Scale bars in A, C and D are 50um.

Fig. S7. Comparison of plasma IgG titers to F-actin, RUVBL2 and HSPD1 between healthy donors and PDAC patients. (**A**) Comparison of IgG titer to F-actin, RUVBL2 and HSPD1 of plasma samples diluted at 1/100 or 1/1000 by ELISA. Background signals from secondary antibody only are shown in a dashed line. (**B**) Comparison of IgG response to F-actin, RUVBL2 and HSPD1 between naïve treated (naive) and neoadjuvant treated (NAT) PDAC patients. The non-parametric t-test is used for comparisons.

715 **Table S1. Human PDAC samples information for scRNA-Seq studies.**

Patient ID	Pt-8	Pt-17	Pt-19	Pt-20	Pt-13	Pt-14	Pt-15
Diagnosis	PDAC	PDAC	PDAC	PDAC	PDAC	PDAC	PDAC
Tumor collection method	resection	resection	resection	resection	FNA	FNA	FNA
Age	58	75	71	67	40	89	53
Gender	Male	Male	Female	Female	Female	Male	Female
Race	White	White	African-American	Asian	Latino/Hispanic	White	African American
Neoadjuvant therapy	No	No	No	No	No	No	No
TNM stage	pT2N2Mx	pT2N1Mx	pT2N1	pT2N0	NA	NA	NA
Tumor size (cm)	2.2*1.5*1	2.7*2.6*2.5	2.7*2.3*2.2	2.2*1.7*0.8	NA	NA	NA
MSI status	MSS	MSS	MSS	MSS	NA	NA	NA
Tumor organoid available	Yes	No	No	Yes	Yes	Yes	No
Tumor section available	Yes	Yes	Yes	Yes	No	No	No
Gene Expressing Sequencing	Yes	Yes	Yes	Yes	Yes	Yes	Failed
Ig sequencing	Yes	Yes (low B and PC cells)	Yes	Yes	Yes	Yes	Yes

716 Abbreviations: FNA, fine needle aspiration; MSI, microsatellite instable; MSS, microsatellite stable; NA,

717 not available/appliable

719   **Table S2. List of antibodies synthesized and screening results.**

No	Patient	Ab-name	Clone size	Cell type	Isotype	Light chain	Binding reactivity	Antigen location	Antigen identity	Synthesized Antibody format
1	Pt-8	8-1	4	PC	IgG1	IgK	Negative	NA		IgG4-His
2	Pt-8	8-2	3	PC	IgA1	IgK	Positive	nuclei enriched	Unknown	IgG4-His
3	Pt-8	8-3	20	PC	IgG2	IgLC3	Positive	cyto	F-actin	IgG4-His
4	Pt-19	19-1	46	PC	IgG1	IgK	Positive	cyto enriched	Unknown	IgG4-His
5	Pt-19	19-3	23	PC	IgG1	IgK	Positive	nuclei enriched	RUVBL2	IgG4-His
6	Pt-19	19-4	3	PC	IgA2	IgK	Positive	cyto/nuclei even	Unknown	IgG4-His
7	Pt-20	20-1	25	PC	IgG2	IgLC2	positive	nuclei enriched	Unknown	IgG4-His
8	Pt-20	20-2	6	PC	IgG2	IgLC1	positive	nuclei enriched	Unknown	IgG4-His
9	Pt-20	20-3	26	PC	IgG2	IgLC1	positive	nuclei enriched	Unknown	IgG4-His
10	Pt-20	20-4	25	PC	IgG2	IgK	positive	nuclei enriched	Unknown	IgG4-His
11	Pt-20	20-5	14	PC	IgG2	IgK	positive	cyto	Unknown	IgG4-His
12	Pt-13	13-1	1	PC	IgG1	IgK	Negative	NA		IgG1
13	Pt-13	13-2	1	PC	IgG1	IgLC2	Negative	NA		IgG1
14	Pt-13	13-3	1	PC	IgA1	IgK	Negative	NA		IgG1
15	Pt-13	13-4	1	PC	IgG1	IgK	Negative	NA		IgG1
16	Pt-13	13-5	1	PC	IgA1	IgK	Negative	NA		IgG1
17	Pt-13	13-6	1	GC-B	IgG1	IgK	Negative	NA		IgG1
18	Pt-13	13-7	1	GC-B	IgG3	IgLC2	Positive	cyto	Unknown	IgG1
19	Pt-13	13-8	2	B	IgA1	IgLC2	Negative	NA		IgG1
20	Pt-14	14-1	4	PC	IgA2	IgK	Positive	cyto/nuclei even	Unknown	IgG1
21	Pt-14	14-2	4	PC	IgA2	IgK	Positive	cyto/nuclei even	Unknown	IgG1
22	Pt-14	14-3	3	PC	IgA2	IgK	Positive	nuclei enriched	Unknown	IgG1
23	Pt-14	14-4	2	PC	IgM	IgK	Negative	NA		IgG1
24	Pt-14	14-5	2	PC	IgA1	IgK	Positive	cyto enriched	Unknown	IgG1
25	Pt-14	14-6	2	PC	IgA2	IgK	Negative	NA		IgG1
26	Pt-14	14-7	2	PC	IgA2	IgK	Positive	nuclei enriched	Unknown	IgG1
27	Pt-14	14-8	2	PC	IgA2	IgLC3	Negative	NA		IgG1
28	Pt-14	14-9	1	PC	IgM	IgK	Negative	NA		IgG1
29	Pt-14	14-10	1	PC	IgM	IgLC2	Positive	nuclei enriched	Unknown	IgG1

30	Pt-14	14-11	1	PC	IgA2	IgLC1	Positive	nuclei enriched	Unknown	IgG1
31	Pt-14	14-12	1	PC	IgG1	IgK	Positive	nuclei enriched	Unknown	IgG1
32	Pt-14	14-13	1	PC	IgA2	IgK	Positive	cyto/nuclei even	Unknown	IgG1
33	Pt-14	14-14	1	PC	IgA2	IgK	Positive	cyto	Unknown	IgG1
34	Pt-14	14-17	1	PC	IgG3	IgK	Positve	cyto enriched	Unknown	IgG1
35	Pt-15	15-1	3	B	IgA2	IgK	Positive	cyto/nuclei even	Unknown	IgG1
36	Pt-15	15-2	3	B	IgA1	IgK	Positive	cyto/nuclei even	Unknown	IgG1
37	Pt-15	15-3	2	PC	IgM	IgKC	Negative	NA		IgG1
38	Pt-15	15-4	2	PC	IgA1	IgK	Negative	NA		IgG1
39	Pt-15	15-5	2	PC	IgA2	IgK	Negative	NA		IgG1
40	Pt-15	15-6	2	B	IgA2	IgLC1	Negative	NA		IgG1
41	Pt-15	15-7	1	PC	IgG1	IgK	Positive	cyto	HSPD1	IgG1

721   Abbreviations: PC-plasma cells, B- B cells, GC-B: Germinal center B cells, cyto-cytoplasmic

722   Light orange color indicates the three antibodies that have antigens identified.

737 **Table S3. Human samples information for plasma studies.**

		Normal (n=61)		PDAC (n=59)		Comparison
		No.	Ratio	No.	Ratio	p value
Gender	Male	22	36%	22	37%	n.s.
	Female	39	64%	32	54%	
	Unknown	0	0%	5	8%	
Age	Median (range)	36	17-66	69.5	49-89	<0.0001
	Unknown	0		5		
Race	Asian	7	11%	5	8%	<0.001
	African American	13	21%	7	12%	
	Caucasian	8	13%	40	68%	
	Hispanic	33	54%	2	3%	
	NA	0	0%	5	8%	
Treatment	Neoadjuvant	-		16	27%	
	No pre-treatment	-		41	69%	
	NA	-		2	3%	
T stage	T1	-		4	7%	
	T2	-		26	44%	
	T3	-		13	22%	
	T4	-		1	2%	
	NA	-		15	25%	
N stage	N0	-		15	25%	
	N1	-		17	29%	
	N2	-		12	20%	
	NA	-		15	25%	
M stage	Mx	-		54	92%	
	M1	-		5	8%	

746