Widespread readthrough events in plants reveal unprecedented plasticity of stop codons

Yuqian Zhang[1,2,6], Hehuan Li[1,6], Yanting Shen[3,6], Shunxi Wang[1], Lei Tian[1], Haoqiang Yin[1], Jiawei Shi[1], Anqi Xing[4], Jinghua Zhang[1], Usman Ali[1], Abdul Sami[1], Xueyan Chen[1], Chenxuan Gao[1], Yangtao Zhao[1], Yajing Lyu[1], Xiaoxu Wang[1], Yanhui Chen[1], Zhixi Tian[3,5,*], Shu-Biao Wu[2,*], Liuji Wu[1,2,*]

[1] National Key Laboratory of Wheat and Maize Crop Science, Collaborative Innovation Center of Henan Grain Crops, College of Agronomy, Henan Agricultural University, Zhengzhou, China.

[2] School of Environmental and Rural Science, University of New England, Armidale, NSW, Australia.

[3] State Key Laboratory of Plant Cell and Chromosome Engineering, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China.

[4] Department of Plant and Environmental Sciences, Clemson University, Clemson, SC, USA.

[5] College of Advanced Agriculture Sciences, University of Chinese Academy of Sciences, Beijing, China.

[6] These authors contributed equally.

*Corresponding authors

Zhixi Tian (zxtian@genetics.ac.cn), Shu-Biao Wu (shubiao.wu@une.edu.au), and Liuji Wu (wlj200120@163.com)

Abstract

Stop codon readthrough (SCR), the decoding of a stop codon as a sense codon by the ribosome, has important biological implications but remains largely uncharacterized in plants. Here, we identified 1,009 SCR events in two monocots (maize, rice) and two dicots (soybean, *Arabidopsis*) using a proteogenomic strategy with 80 customized databases. SCR transcripts were mostly significantly shorter and had fewer components than non-SCR transcripts in two monocot plants, although these differences were not as significant in the dicots. Mass spectrometry evidence revealed that all three stop codons involved in SCR events could be recoded as 20 standard amino acids, some of which were also supported by suppressor transfer RNA analysis. In addition, we observed multiple functional signals in the C-terminal extensions of 34 maize SCR proteins, and characterized the structural and subcellular localization changes in the extended protein of BASIC TRANSCRIPTION FACTOR 3. Overall, our study not only demonstrates that SCR events are widespread in plants but also reveals the unprecedented recoding plasticity of stop codons, which provides important new insights into the flexibility of genetic decoding.

Introduction

The rules of the genetic code were long considered universal dogma in all organisms after the standard genetic code was deciphered decades ago. However, exceptions to the rules have been increasingly observed in different species over the years, substantially altering the perception of the decoding rules from being static to dynamic[1,2]. Living organisms are now widely understood to employ various modes to expand the coding potential of their genome, such as alternative translation initiation[3], ribosome frameshifting[4], and translational bypassing[5]. Stop codon readthrough (SCR) is another important means of recoding that is classified as a non-canonical translational event[6,7]. SCR occurs when the ribosome reads a stop codon (UGA, UAG, or UAA) as a sense codon rather than the stop signal with the translation terminating at a later in-frame stop codon[7,8]. Such an event can generate a C-terminally extended protein isoform which may acquire novel biological functions, since the amino acid sequence appended at the C-terminus possibly alters the activity[9], stability[10], and/or subcellular localization[2,11] of the protein. Indeed, SCR proteins have been implicated in a variety of biological processes, such as the regulation of intercellular cytoplasm flow[12], neuronal integrity and reproductive health[13], cellular redox homeostasis[2,14], and angiogenesis[15,16].

Initially discovered in viruses as a strategy to expand protein-coding potential[17-19], SCR was subsequently identified in other organisms. In plant pathogens, *Phosphoglycerate kinase* (*PGK*) in *Ustilago maydis* and *Glyceraldehyde-3-phosphate dehydrogenase* (*GAPDH*) in *Aspergillus nidulans* are typical functional readthrough genes conserved in fungal species[2]. In the yeast *Saccharomyces cerevisiae*, the [PSI+] strain forms prion-like polymers of the abnormal eukaryotic release factor 3 (eRF3), which is implicated in translation termination and peptide release, resulting in a high level of readthrough across many genes[20], including *Phosphodiesterase 2* (*PDE2*), *Interacting with mpp10p 3* (*IMP3*), and *Bypass of stop codon 4* (*BSC4*)[10,21,22]. In *Drosophila melanogaster*, readthrough is estimated to affect over 300 genes[23,24], such as *kelch* (*kel*)[12,25], *Synapsin* (*Syn*)[26], *headcase* (*hdc*)[27], and *traffic jam* (*tj*)[13]. In

mammals, *Hemoglobin beta chain* (*HBB*)[28], *Myelin protein zero* (*MPZ*)[29], *Vascular endothelial growth factor-A* (*VEGF-A*)[15], *Mitochondrial carrier 2* (*MTCH2*)[30], *Argonaute 1* (*AGO1*)[31], *Opioid related nociceptin receptor 1* (*OPRL1*)[32], *Mitogen-activated protein kinase 10* (*MAPK10*)[32], *Aquaporin 4* (*AQP4*)[33], *Lactate dehydrogenase B* (*LDHB*)[34], *Malate dehydrogenase 1* (*MDH1*)[34], *Vitamin D receptor* (*VDR*)[35], and *Adenosylmethionine decarboxylase 1* (*AMD1*)[36,37] have been confirmed to undergo readthrough. However, SCR events in plants remain largely uncharacterized, with only one study reporting 144 readthrough genes in *Arabidopsis* detected with ribosome profiling[38]. Therefore, further exploration of SCR events with a more robust analysis is warranted in a wide range of plant species.

It is crucial to identify the amino acids decoded at the stop codon sites, as they can strongly affect the functionality of readthrough protein isoforms[11,21,39]. Previous studies have shown that UGA could be translated to arginine, asparagine, cysteine, glutamine, glycine, serine, threonine, tryptophan, and tyrosine[30,31,39-41]; UAA to alanine, glutamate, glutamine, lysine, serine, tryptophan, and tyrosine[39,42-45]; and UAG to alanine, arginine, aspartate, glutamate, glutamine, glycine, leucine, lysine, serine, threonine, tryptophan, tyrosine, and valine[30,39,46,47]. However, knowledge about the amino acids translated at stop codon sites in SCR events remains unknown in Plantae. In addition, the potential for stop codons to be recoded as amino acids other than those reported above has not been investigated yet.

Here, we employed a mass spectrometry (MS)-based proteogenomic strategy to study SCR events in plants and to identify the amino acids at the readthrough stop codon sites. We detected a total of 1,009 SCR events in the monocots maize and rice, and dicots soybean and *Arabidopsis*, and characterized their features. Interestingly, we discovered that the three stop codons could be recoded as all 20 standard amino acids, which reveals the unprecedented recoding plasticity of stop codons and the dynamic nature of genetic decoding. We also delineated the first comprehensive landscape of SCR events in plants at the protein level, uncovering a collection of previously unknown readthrough proteins to further explore their biological functions.

Results

A proteogenomic strategy to identify SCR events in plants

The SCR event is a means of alternative translation by redefining UGA, UAG, or UAA to specify amino acids, thus extending the translation at the C-terminus. The most definitive evidence for SCR occurrence is the direct detection of translation at the stop codon sites. However, SCR events are non-canonical and cannot be identified by a typical proteomic strategy. Therefore, we implemented a proteogenomic strategy integrating MS-based proteomics with 20 customized databases to identify SCR events in maize at the protein level (Fig. 1). First, total protein was extracted from maize leaves, followed by digestion, fractionation, and liquid chromatography-tandem mass spectrometry (LC-MS/MS) analysis to obtain MS/MS datasets of peptides for subsequent identification. We then constructed 20 customized databases with in-house scripts based on the assumption that the annotated stop codons of genes were readable as sense codons and could be decoded as one of the 20 standard amino acids. Each of the 20 customized databases therefore contained the *in silico*-translated protein isoforms whose annotated stop codon sites were replaced by one of the 20 standard amino acids, and the protein sequences were extended to the subsequent in-frame stop codon.

To capture SCR events in maize, we used the obtained MS/MS data to search against the 20 SCR customized databases with MaxQuant search engine. This strategy identified 708 unique "through" peptides spanning the stop codon site. For further quality control of MS data, we next applied more stringent criteria for "through" peptides to identify SCR events with high confidence (see Methods). The screening steps resulted in much more reliable "through" peptides, which corresponded to 367 SCR events involving 1,063 proteins (Supplementary Table 1). All identified peptides for SCR proteins are listed in Supplementary Table 2. Collectively, these results not only demonstrate the abundance of readthrough events naturally occurring in maize, but also uncover a previously unknown source of protein isoforms produced by SCR.

Verifying SCR events in maize

To verify the SCR events in maize identified above, we retrieved the available

ribosome profiling (Ribo-seq) datasets of 56 maize samples from NCBI to analyze the

SCR events at the mRNA level. Four Ribo-seq datasets with higher quality were

finally retained after further screening (see Methods). We then analyzed the three-

nucleotide (3-nt) periodicity of the Ribo-seq reads (30-mers) in the coding sequences

(CDSs), putative extensions, and distal 3′ untranslated regions (UTRs). The results

showed that the putative extensions exhibited a visible hallmark of 3-nt periodicity (P

< 0.05, Fig. 2a), i.e., the reading frame 0 occurred significantly higher than the other

two, which was similar to that of the CDSs (Fig. 2a). By contrast, such difference was

not observed in the distal 3′ UTRs (Fig. 2a). In addition, the average ribosome

footprint density (in RPKM) of putative extensions was significantly higher than that

of the distal 3′ UTRs in the four datasets ($P < 0.001$, Fig. 2b). These results suggest

that readthrough is a significant contributor to the ribosome footprint density in

putative extensions. After applying the conditions described in the Methods section

for the four Ribo-seq datasets to verify SCR, we found that 55 out of the 367 SCR

events identified by the proteogenomic approach were also corroborated by Ribo-seq

analysis (Fig. 2c and Supplementary Table 3).

We selected two SCR representative genes, *ZmH3C2* and *ZmCEQORH*, which

exerted functional roles in distinct biological processes related to chromatin assembly

and stress response, respectively, for further analysis. For *ZmH3C2*, which encodes

HISTONE H3.2, MS analysis identified the readthrough peptide AAAAGRPDRPWR

with its second alanine (A) translated at the UAG site (Fig. 2d and Supplementary Fig.

1a). The readthrough of this gene was also evidenced by ribosome footprints within

the extended region downstream of the annotated stop codon (Fig. 2e). For

ZmCEQORH,   encoding   CHLOROPLAST   ENVELOPE   QUINONE

OXIDOREDUCTASE,   MS   analysis   identified   the   "through"   peptide

AQEVWAKSIDGHATGKIVVEMGGAAEETR,   in   which   the   antepenultimate

glutamate (E) was incorporated at the UGA site (Fig. 2h and Supplementary Fig. 1b).

We also observed ribosome footprints beyond the annotated stop codon of the
corresponding transcript (Fig. 2i).

In addition, we performed *in vitro* translation experiments using wheat germ
extract to further demonstrate the readthrough of the above two genes. The results
showed that *ZmH3C2* and *ZmCEQORH* had significantly higher readthrough
activities compared to the SCR-negative gene *Zm00001d009138* ($P < 0.001$), as
demonstrated by the detection of luciferase activities. The SCR rates of *ZmH3C2* and
ZmCEQORH were 12% and 5%, respectively (Fig. 2f, j). It is known that the rate of
the basal translation error is lower than 0.1%[8], thus the results indicate that *ZmH3C2*
and *ZmCEQORH* underwent the programmed SCR events. Further MS analysis of the
readthrough products identified the "through" peptides AAAAGRPDRPWR and
GGAAEETR for the two genes, where the UAG and UGA were recoded as alanine
and glutamate (Fig. 2g, k), respectively. Taken together, our Ribo-seq analysis, *in vitro*
translation experiments, and MS analysis validated the SCR events identified using
proteogenomics in maize, and demonstrated the feasibility of the proteogenomic
method for large-scale identification of SCR events in plants.

Characteristics of SCR events in maize

To understand the characteristics of SCR events in maize, we first analyzed the stop
codon preferences. All three stop codons were read through among the 367 SCR
events: UGA was involved in 174 events (47.4%), followed by UAG in 102 events
(27.8%) and UAA in the remaining 91 events (24.8%) (Fig. 3a). We found that UGA
was the most frequent next in-frame stop codon following these readthrough stop
codons, accounting for 76 cases after the readthrough UGA, 46 cases after the
readthrough UAG, and 40 cases after the readthrough UAA (Fig. 3a). The genome-
wide distribution pattern showed that these SCR events were distributed
heterogeneously across each chromosome of maize, and many of the SCR events were
closer to the telomeres (Fig. 3b). In addition, a collection of 379 hotspot regions were
obtained, 244 of which were SCR hotspot regions while 135 were non-SCR hotspot
regions (Fig. 3b). These results revealed the wide distribution of SCR events in the

maize genome.

Furthermore, we compared the length and complexity of SCR and non-SCR

transcripts. Interestingly, we observed that the SCR primary transcripts were

significantly shorter than the non-SCR transcripts (Fig. 3c). The average lengths of

the elements from the primary transcripts, including coding sequences (CDSs), introns,

and 3′ UTRs, were also markedly shorter in SCR events than those in non-SCR events

(Fig. 3c). The 5′ UTR was also shorter in readthrough transcripts, albeit not

statistically significant. Consistent with the length of readthrough transcripts, their

corresponding proteins had notably shorter lengths with fewer amino acids

(Supplementary Fig. 2a) and lower molecular weight (Supplementary Fig. 2b).

Moreover, SCR primary transcripts had significantly fewer exons (Fig. 3d) and splice

variants (Fig. 3e) than those of non-SCR primary transcripts.

Earlier studies have shown that *cis*-acting elements such as nucleotides

immediately surrounding the readthrough stop codon and motifs downstream the stop

codon can influence readthrough[48-50]. We thus examined the local mRNA context of

SCR transcripts. An analysis of nucleotide frequencies in positions adjacent to the

readthrough stop codons showed a bias of A>U>C>G at the -2 position, A>G>C>U at

the -1 position, A>G>U>C at the +4 position, and U>C>G>A at the +5 position

(Supplementary Fig. 3a). The most frequent tetranucleotide pattern (stop codon plus

the following nucleotide) was UGA-U (Supplementary Fig. 3b). Extended regions

downstream of the annotated stop codons in the readthrough transcripts were enriched

with featured sequence motifs (Supplementary Dataset 1). Three motif examples,

ZNF263, ASCL1, and ETV6 are shown in Supplementary Fig. 3c. These motifs

enriched in extended regions likely exert effects on the particular facets of SCR by

interacting with specific proteins. In addition, extended regions of SCR transcripts

harbored remarkably lower free energy than their analogs of non-SCR transcripts

(Supplementary Fig. 3d), suggesting more stable secondary structures in the

extensions of the readthrough transcripts.

Stop codons are recoded as 20 amino acids in maize SCR events

Since translation of the stop codon is crucial for SCR events, we examined which

amino acids were translated at every readthrough stop codon site based on the MS

data. Quite unexpectedly, we found that, with the exception of UAA which was

recoded to 19 amino acids (except serine), UGA and UAG were translated to all 20

standard amino acids in maize SCR events (Fig. 4a). Among all the amino acids,

glutamate was the most recoded amino acid for UGA (in 17 SCR events, 8.9%),

methionine for UAG (in 17 cases, 15.6%), and aspartate for UAA (in 11 cases, 11.6%).

Previous literature has reported several amino acids that were able to be

translated at stop codon sites in different species (Fig. 4b and Supplementary Table 5),

prompting us to explore the reasons behind the recoding plasticity of the stop codons

in our results. Suppressor transfer RNA s (tRNAs) are specific tRNA species forming

classical base pairing with the stop codon, which can lead to the translation[24]. We thus

conducted a genome-wide search for such potential suppressor tRNAs capable of

reading stop codons in maize by running tRNAscan-SE[51] and ARAGORN[52]. As a

result, we acquired a collection of UGA-, UAG-, and UAA-recognizing tRNAs.

Multiple sequence alignments of the predicted suppressor tRNAs against the NCBI

Nucleotide (NT) database resulted in different groups of suppressor tRNAs well

aligned with the canonical tRNAs of a range of amino acids (Supplementary Table 6).

We identified six classes of UGA suppressor tRNAs (Supplementary Tables 5 and 6):

$tRNA^{Met}$(UCA), $tRNA^{Lys}$(UCA), $tRNA^{Val}$(UCA), $tRNA^{Gly}$(UCA), $tRNA^{Ile}$(UCA), and

$tRNA^{Thr}$(UCA). The predicted secondary structures of the first two UGA suppressor

tRNAs are shown in Fig. 4c I, II, and their nucleotide sequence alignments are

presented in Supplementary Fig. 4a, b. We also detected six classes of UAG

suppressor tRNAs (Supplementary Tables 5 and 6): $tRNA^{Glu}$(CUA) (Fig. 4c III and

Supplementary Fig. 5a), $tRNA^{Trp}$(CUA) (Fig. 4c IV and Supplementary Fig. 5b),

$tRNA^{Arg}$(CUA), $tRNA^{Asp}$(CUA), $tRNA^{Gly}$(CUA), and $tRNA^{Lys}$(CUA). Furthermore,

we identified eight classes of UAA suppressor tRNAs (Supplementary Tables 5 and 6):

$tRNA^{Lys}$(UUA) (Fig. 4c V and Supplementary Fig. 6a), $tRNA^{His}$(UUA) (Fig. 4c VI

and Supplementary Fig. 6b), $tRNA^{Ala}$(UUA), $tRNA^{Asn}$(UUA), $tRNA^{Gly}$(UUA),

tRNA^{Phe}(UUA), tRNA^{Met}(UUA), and tRNA^{Thr}(UUA). Collectively, the suppressor tRNAs identified in maize provide potential reasons for the translation of at least some amino acids at stop codon sites.

SCR events and recoding plasticity of stop codons in other plants

In light of the abundant SCR events identified in maize, we questioned if SCR also occurred in other plant species. We thus followed the same proteogenomic method with the respective 20 custom databases to explore SCR events in rice, soybean, and *Arabidopsis*. Indeed, we detected abundant SCR events in these species with similar features to those in maize. We identified 198 SCR events in rice (Supplementary Table 7), 179 SCR events in soybean (Supplementary Table 9), and 265 SCR events in *Arabidopsis* (Supplementary Table 11), corresponding to 269, 385, and 468 proteins, respectively.

In rice, we observed that the stop codon usage SCR events was in the order of UGA > UAA > UAG, and the next in-frame stop codon following the readthrough stop codons also displayed a preference for UGA (Fig. 5a). For the dicots soybean and *Arabidopsis*, the stop codon usages were the same in the order of UGA > UAA > UAG in SCR events, while the most frequent next in-frame stop codon after the readthrough stop codons was UGA for soybean and UAA for *Arabidopsis* (Fig. 5b, c). Genome-wide distribution analysis showed that SCR events were distributed across all chromosomes in their respective plant species (Fig. 5d, e, f). We also analyzed the features of SCR and non-SCR transcripts in the three plants. In rice, the length of SCR primary transcripts and transcript elements including CDSs, introns and the 5′ UTR were shorter than those of non-SCRs ($P < 0.05$, Supplementary Fig. 7a, b). In addition, the SCR primary transcripts tended to possess fewer exons (Supplementary Fig. 7c) and had significantly fewer splice variants ($P < 0.001$, Supplementary Fig. 7d) than non-SCR transcripts. In the dicots soybean and *Arabidopsis*, however, the SCR transcript features were not apparent as compared to those in monocots. SCR transcripts in soybean had significantly shorter UTR lengths and fewer exons than the non-SCR transcripts ($P < 0.05$, Supplementary Fig. 8b, c) but no significance for

other feature comparison (Supplementary Fig. 8a, b, d). Similarly, SCR transcripts

from *Arabidopsis* showed no significant different from the non-SCR counterparts

except fewer splice variants (Supplementary Fig. 9a, b, c, d).

Analysis of the recoded amino acids at stop codon sites revealed that UGA

exhibited great plasticity to be recoded as the 20 standard amino acids in all the three

plant species (Fig. 5g, h, i). For UAG, it was recoded to 18 amino acids (except serine

and tryptophan) in rice (Fig. 5g), 17 amino acids (except cystine, aspartate, and

phenylalanine) in soybean (Fig. 5h), and 18 amino acids (except aspartate and

arginine) in *Arabidopsis* (Fig. 5i). Similarly, UAA was translated to 17, 16, and 20

amino acids in rice, soybean, and *Arabidopsis*, respectively (Fig. 5g, h, i). These

results further uncover the high recoding plasticity of stop codons in plant SCR events.

C-terminal extensions in maize SCR events contain multiple functional signals

To examine the biological implications of SCR events, we searched the functional

signals in all C-terminal extensions of readthrough proteins in maize. As a result,

multiple functional signals were detected in 34 extensions, including the

transmembrane domains in 28 extensions, nuclear localization signals in two

extensions, and peroxisomal targeting signals (PTS1) in four extensions (Fig. 6a and

Supplementary Table 13). Among them, we selected BASIC TRANSCRIPTION

FACTOR 3 (ZmBTF3), which was predicted to possess a transmembrane domain in

its C-terminal extension (54 amino acids), for further analysis. First, we conducted the

in vitro translation experiment to validate the readthrough of this gene. We observed

high readthrough activity of *ZmBTF3* with an SCR rate of 15% (Fig. 6b), which

revealed that *ZmBTF3* was read through and generated a novel isoform (hereafter

termed ZmBTF3x). Next, we confirmed the expression of the fusion protein

ZmBTF3x-Fluc by immunoblotting using an anti-Fluc antibody (Fig. 6c). The results

showed that the molecular weight of the readthrough protein ZmBTF3x-Fluc was the

same as its positive control ZmBTF3x^{ACA}-Fluc and higher than that of ZmBTF3-Fluc

(Fig. 6c). By contrast, no visible band of readthrough products expressed from the

SCR-negative control *Zm00001d009138-TAG-ISR-Fluc* was observed in the immunoblot (Fig. 6c). The fusion protein ZmBTF3x-Fluc was then subjected to mass spectrometric analysis, resulting in the detection of different types of peptides labeled as "before", "through", and "after", which further validated the production of the readthrough protein isoform ZmBTF3x (Fig. 6d, top). The spectrum for "through" peptide "EPEEKKESTFCCV" revealed that the stop codon UAA was translated as threonine (T) (Fig. 6d, bottom).

The single transmembrane domain predicted in the C-terminus of ZmBTF3x was found to contain 20 amino acid residues crossing the membrane (Fig. 6e). The predicted 3D structures of ZmBTF3 and ZmBTF3x revealed an α-helix appended at the ZmBTF3x C-terminus that was not present in ZmBTF3 (Fig. 6f). Furthermore, the α-helix with a high pLDDT score and low PAE score corresponded exactly to the region of the predicted transmembrane domain (Fig. 6f and Supplementary Fig. 10a, b). To examine the biological significance of the C-terminal extension derived from SCR of *ZmBTF3*, we constructed *35S:ZmBTF3-GFP* and *35S:ZmBTF3x-GFP* vectors and individually transfected them in maize protoplasts for subcellular localization analysis using confocal microscopy. The results showed that ZmBTF3 localized to the nucleus and cytosol (Fig. 6g), consistent with the previous study reporting that *Arabidopsis* BTF3 acted as the transcription factor and the β-subunit of nascent polypeptide-associated complex (NAC)[53]. By contrast, the readthrough protein ZmBTF3x localized to the plasma membrane and cytosol (Fig. 6g), confirming the localization change between the isoforms ZmBTF3 and ZmBTF3x. In summary, these results indicate that SCR can lead to the structural and subcellular localization change of the readthrough protein isoform, which may confer new biological functions.

Discussion

SCR, classified as the recoding of stop codons, is a regulatory mode of translation that could produce multiple protein isoforms from the same transcript[1,54]. SCR has been reported in a variety of organisms with important biological implications. Recently, Sahoo *et al*. have identified mRNAs of 144 genes that exhibited SCR in *Arabidopsis*

by analyzing Ribo-seq datasets[38]. However, a systematic study of SCR events in

plants at the protein level is still lacking. In the current study, we employed a high-

resolution MS-based proteogenomic strategy to analyze SCR events in different plants

and to identify the amino acids translated at stop codon sites. We established 20

customized databases for each of the four plant species with one of the 20 amino acids

defined at annotated stop codon sites in each database, which enabled detection of any

potential SCR events with all possible amino acids translated at stop codon sites.

Using this strategy, we detected 1,009 SCR events corresponding to a collection of

2,185 proteins in four plant species.

Our systematic analysis of SCR events revealed their similar characteristics

among the four plant species. Firstly, SCR events occur in all chromosomes of the

four species, and SCR events are prone to be dispersively distributed in the

chromosome, except in maize where the higher distribution of SCR was closer to

telomeres. Secondly, consistent with previous studies[6,23], UGA is the most preferential

stop codon for readthrough, and it also turned out to be the most preferred next in-

frame stop codon in SCR events of these species. In addition, SCR transcripts were

mostly significantly shorter and had fewer exons and splice variants than non-SCR

transcripts in two monocot plants, although these were not as significant as that in the

dicots. However, in other eukaryotes like *D. melanogaster* and *Anopheles gambiae*,

SCR transcripts were found to be much longer than the non-SCR transcripts, possibly

because the occurrence of SCR requires more regulatory elements embedded in their

longer transcripts[23,24]. Our findings suggest that the regulation of SCR in plants might

differ from that in animals.

The stop codons are able to be recoded as all 20 standard amino acids in the four

plant species in our study, which has not been reported elsewhere. This result suggests

a greater recoding potential of stop codons underestimated in the past and reveals a

more flexible mode of decoding. Since tRNA is the adapter molecule in decoding a

codon to an amino acid, we explored the reasons for the plasticity of stop codons by

tRNA analysis. Suppressor tRNAs, of which the anti-codon could fully form

canonical base-pairing with stop codons, have been reported to contribute to the translation of stop codons[24,55,56]. Here, we detected an array of suppressor tRNAs corresponding to different amino acids in maize, which may be the underlying reasons for the translation of six amino acids at UGA, six amino acids at UAG, and eight amino acids at UAA sites. In addition to suppressor tRNAs, near-cognate tRNAs of stop codons are also an important reason for the translation due to the noncanonical base pairing of one position between a stop codon and the anticodon of the tRNA[41,44]. The amino acids carried by these near-cognate tRNAs include arginine, serine, tryptophan, cysteine, glycine, and leucine for UGA; lysine, glutamate, glutamine, tryptophan, serine, leucine, and tyrosine for UAG; and lysine, glutamate, glutamine, tyrosine, serine, and leucine for UAA[57,58]. However, we did not find suppressor tRNAs for the rest small portion of the 20 amino acids recoded at stop codon sites in our study, nor the tRNAs of these amino acids were near-cognate tRNAs (e.g., the case of UAG recoded as alanine and UGA recoded as glutamate in Fig. 2). Indeed, these amino acids are not exclusively observed in this study, as previous studies have also reported some of their translation at stop codon sites[15,30,39,46]. These cases suggest that our present understanding of stop codon reading by suppressor or near-cognate tRNAs is still incomplete. Considering the sophisticated modifications and regulation of the decoding adaptor tRNAs and multiple factors that influence SCR events[8,59], the mechanism underlying SCR associated with different tRNAs warrants further study.

In summary, we implemented a proteogenomic strategy and identified abundant SCR events across the genomes of four plant species. Stop codons in these SCR events could be recoded as all 20 standard amino acids. We demonstrated that SCR events are widespread in plant species and uncovered the unprecedented plasticity of stop codons. Our findings provide evidence that genetic decoding is far more flexible than has been recognized. We also identified a large volume of novel protein isoforms derived from SCR, which not only expands the plant proteome but also lays a foundation for further exploring the functionality of SCR products.

Methods

Sample preparation

Maize (*Zea mays,* B73 inbred line), rice (*Oryza sativa* L. ssp. *japonica* cv. Nipponbare), *Arabidopsis thaliana* (Columbia-0), and soybean (*Glycine max* L., ZH13) plants were grown in a temperature- and light-controlled greenhouse. The growth conditions for maize were a 15-h light/9-h dark photoperiod and a 28/25 °C day/night air temperature, and samples were collected at the three-leaf stage. For rice, the growth conditions were a 12-h light/12-h dark photoperiod and a 28/25 °C day/night air temperature, and samples were collected on day 56. *Arabidopsis* plants were grown in a greenhouse under a 16-h light/8-h dark photoperiod and a 22/21 °C day/night air temperature, and samples were collected on day 21. Soybean plants were grown in a greenhouse under a 16-h light/8-h dark photoperiod and a 22/21 °C day/night air temperature, until the trifoliate leaves were fully expanded. Three replicates of leaf samples for each of the plant species were harvested, frozen in liquid nitrogen, and stored at –80 °C until analysis.

Protein extraction

Total proteins from each sample were extracted as previously reported[60]. Briefly, leaf tissues (1 g) for each plant species were pulverized in liquid nitrogen, and the powder was precipitated in 10% (w/v) trichloroacetic acid/acetone solution containing 65 mM dithiothreitol (DTT) at –20 °C for 1 h. Then, the solution was centrifuged at 10,000×g at 4 °C for 45 min and the supernatant was removed. The precipitate was then vacuum-dried and solubilized in 1/10 volumes of SDT buffer (4% [w/v] SDS, 100 mM DTT, and 150 mM Tris-HCl, pH 8.0). After a 3-min incubation in boiling water, the suspensions were ultrasonicated (80 w, ultrasonic 10 s/time, every 15 s, repeated 10 times) and incubated at 100 °C for 3 min. The extract was centrifuged at 12,000×g at 25 °C for 10 min, and the total protein contents were quantified using BCA Protein Assay Reagent (Promega Corporation). The supernatants were stored at –80 °C until subsequent analysis.

Protein digestion and peptide fractionation

The total proteins (250 μg for each sample) were digested by FASP procedure using

trypsin (Promega Corporation) as described previously[60]. Then, the digested peptide filtrate was fractionated by strong cation exchange (SCX) chromatography using an AKTA Purifier system (GE Healthcare). The dried peptide mixture was reconstituted and acidified with 2 ml buffer A (10 mM KH_2PO_4 in 25%[v/v] of ACN, pH = 2.7) and loaded onto a PolySULFOETHYL 4.6×100 mm column (5 µm, 200 Å, PolyLC Inc). The peptides were eluted at a flow rate of 1 ml/min buffer B (500 mM KCl, 10 mM KH_2PO_4 in 25% of ACN, pH 2.7) in a gradient of 0-10% (v/v) for 2 min, 10-20% (v/v) for 25 min, 20-45% (v/v) for 5 min, and 50-100% (v/v) for 5 min. The elution was monitored by absorbance at 214 nm, and fractions were collected every 1 min. The 30 collected fractions were combined into 15 pools and desalted on C18 Cartridges (Empore SPE Cartridges C18 standard density, bed I.D. 7 mm, volume 3 ml, Sigma-Aldrich). Each fraction was concentrated by vacuum centrifugation and reconstituted in 40 µl of 0.1% (v/v) trifluoroacetic acid. All samples were stored at –80 °C before LC-MS/MS analysis.

Liquid chromatography-tandem mass spectrometry analysis

The samples were analyzed on a Q-Exactive HF-X mass spectrometer coupled to the Easy nLC 1200 (Thermo Fisher Scientific). Peptide (2 µg) was loaded onto a 25 cm, 75 µm inner diameter C18-reversed phase column packed with 3 µm resin (EASY-Column Capillary Columns, Thermo Fisher Scientific) in buffer A (2% [v/v] acetonitrile and 0.1% [v/v] formic acid) and separated with a linear gradient of buffer B (80% [v/v] acetonitrile and 0.1% formic acid) at a flow rate of 250 nl/min controlled by IntelliFlow technology over 90 min. MS data were acquired using a data-dependent top 10 method dynamically choosing the most abundant precursor ions from the survey scan (m/z 300-1,800) for higher-energy collision dissociation fragmentation (HCD). Determination of the target value was based on predictive Automatic Gain Control (pAGC). The dynamic exclusion duration was 25 s. Survey scans were acquired at a resolution of 70,000 (m/z 200), and the resolution for HCD spectra was set to 17,500 (m/z 200). Normalized collision energy was 30 eV, and the underfill ratio, which specifies the minimum percentage of the target value likely to

be reached at maximum fill time, was defined as 0.1%. The instrument was run with

peptide recognition mode enabled. MS experiments for each sample were conducted

in triplicate for each sample.

Customized database construction

The complete genome sequences of the four plants were downloaded from Ensembl

Plants (ftp://ftp.ensemblgenomes.org/pub/plants/release-41/fasta/zea_mays/dna/) for

maize, MSU Rice Genome Annotation Project

(http://rice.uga.edu/pub/data/Eukaryotic_Projects/o_sativa/annotation_dbs/pseudomol

ecules/version_7.0/all.dir/) for rice, the National Genomics Data Center

(ftp://download.big.ac.cn/gwh/Plants/Glycine_max_Gmax_ZH13_v2.0_GWHAAEV

00000000.1/GWHAAEV00000000.1.genome.fasta.gz) for soybean, and TAIR 10

(ftp://ftp.ensemblgenomes.org/pub/plants/release-

28/fasta/Arabidopsis_thaliana/dna/Arabidopsis_thaliana.

TAIR10.28.dna.toplevel.fa.gz) for *Arabidopsis* in FASTA format. The tailored

databases for SCR event identification were generated from the computational

translation of plant genome sequences to amino acid sequences by EMBOSS 6.6.0, as

previously described with modifications[46]. *In silico* translation followed the principle

that the annotated stop codon (UGA, UAG, or UAA) of a given gene was considered

a translatable codon to be designated as the symbol "X", representing any one of the

20 standard amino acids, and the elongation terminated at the next in-frame stop

codon in the 3′ UTR. The *in silico* translation was conducted by in-house scripts based

on Shell under a Linux environment to produce 20 customized databases for each

plant species. Each database has been in conjunction with a public plant protein

database for every plant species as follows: Ensembl Plants

(http://ftp.ebi.ac.uk/ensemblgenomes/pub/release-41/plants/fasta/zea_mays/pep/Zea_

mays.B73_RefGen_v4.pep.all.fa.gz) for maize, MSU Rice Genome Annotation

Project protein database (http://rice.uga.edu/pub/data/Eukaryotic_Projects/o_sativa

/annotation_dbs/pseudomolecules/version_7.0/all.dir/all.pep) for rice, the National

Genomics Data Center

(ftp://download.big.ac.cn/gwh/Plants/Glycine_max_Gmax_ZH13_v2.0_GWHAAEV 00000000.1/GWHAAEV00000000.1.Protein.faa.gz) for soybean, and Ensembl Plants (ftp://ftp.ensemblgenomes.org/pub/plants/release-51/fasta/Arabidopsis_thaliana/pep/Arabidopsis_thaliana.TAIR10.pep.all.fa.gz) for *Arabidopsis*. The conjunction of the public protein database ensured the searching precision and calculation of the false discovery rate (FDR) in the subsequent searching stage.

Identification of SCR events by MaxQuant

The raw data were analyzed using MaxQuant software (v.1.6.1.10)[61]. The spectra were searched against the 20 customized databases for each of the four plant species. The initial search was set at a precursor mass window of 6 ppm. The search followed an enzymatic cleavage rule of trypsin and allowed a maximum of two missed cleavage sites and a mass tolerance of 20 ppm for fragment ions. Carbamidomethylation of cysteines was defined as fixed modification, while methionine oxidation was defined as variable modification for database searching. The cutoff of global FDR for peptide and protein identification was set to 0.01. Protein quantification was calculated by normalized spectral protein intensity.

All matched peptides were then relocated to the corresponding positions of protein sequences in the reference databases and mapped to their genomic locations. Every stop codon site was given an ID for genomic position recognition. The non-redundant peptides uniquely mapped to the regions spanning the annotated stop codon were labeled as "through", the peptides mapped to the coding sequence (CDS) region were labeled as "before", and the peptides mapped to the 3′ UTR were labeled as "after". We further implemented more stringent criteria to determine the "through" peptides so as to identify SCR events. Only the unique "through" peptides that met the following conditions were retained: (1) presented in at least two of the three replicates in MS experiment; (2) at least two of the quality scores were higher than 20. Finally, proteins sharing one stop codon site and containing a unique "through" peptide were merged into one group and the group was defined as one SCR event.

Ribosome profiling analysis

We obtained ribosome profiling (Ribo-seq) datasets of 56 maize samples from previous reports for SCR event analyzing[62-66]. The Ribo-seq data was analyzed according to the methods described by Dunn et al. (2013)[11] and Sapkota et al. (2019)[6] with modifications. Briefly, the ribosomal sequencing data were quality-controlled using fastax (v.36.06) or cutadapt (v.1.13)[62] to remove low-quality bases, adaptor sequences, and random primers. Reads of 25-35 nucleotides (nts) were retained by seqkit (v.0.5.4) for subsequent analysis. The remaining reads were aligned to the maize genome by STAR (v.2.7.3a)[67]. Ribosome-protected footprint alignments were mapped to their estimated P-sites as follows: 12 nt were cut from each end of the alignment, resulting in a fragment of $n \times$ nt in length ($n = l$-12×2). Each genomic position covered by a nucleotide remaining in the truncated alignment was then incremented by 1/n (count number). Then, the reads were mapped to the following positions: 12 nts after the start codon, 15 nts before and 9 nts after the first stop codon, and 15 nts after the second stop codon. This resulted in four regions remaining in the transcripts: the 5′ UTR, CDS, readthrough region, and the distal 3′ UTR.

We then analyzed the length distribution of Ribo-seq reads in each dataset. Four Ribo-seq datasets in which the read length was mainly distributed in 30-nt (30-mer) and prepared with RNase I (little cutting bias) were finally retained for further SCR validation. For analysis of 3-nt periodicity (phasing) of footprints, R package ribosomeProfilingQC (v.1.10.0) was used and only 30-nt RPFs were counted, as shown in maize with good-phased footprint population[62]. Reads were then counted in all putative extended regions. Mean value of normalized RPKM (reads per kilobase per million mapped reads) was calculated in R (v.3.2.5) in the four datasets. The ribosome footprint density of putative extensions and the distal 3′ UTRs were calculated in RPKM. Only extensions with RPKM > 1, read coverage > 10%, and no overlap with any annotated 5′ UTRs, CDSs, and other non-coding RNAs were retained and regarded as readthrough. Full data of samples that exhibited readthrough in Ribo-seq analysis for each of the SCR events are shown in Supplementary Table 3.

Distribution of readthrough events at the genomic level

The genome-wide density of SCR and non-SCR events in maize chromosomes was plotted using a sliding window of 20 Mb with 2-Mb steps according to the annotated maize genome in Ensembl (https://plants.ensembl.org/index.html). We used "hotspot region" to reflect the distribution density of SCR and non-SCR events. Hotspot regions were defined as ≥ 5 SCR events for SCR hotspots and ≥ 335 for non-SCR events in a window size of 20 Mb. The chromosomal distributions of rice SCR and non-SCR events were drawn using a window size of 4 Mb with a step of 400 kb based on the annotated rice genome from MSU (http://rice.uga.edu/). The same distribution pattern was plotted with a window size of 10 Mb with a 1-Mb step based on the ZH13 reference genome (https://ngdc.cncb.ac.cn/gwh/Assembly/652/show) for soybean. For *Arabidopsis*, the SCR and non-SCR genomic localizations were obtained by a 1.2-Mb sliding window with a step size of 100 kb according to the annotated *Arabidopsis* genome in TAIR (https://www.Arabidopsis.org/). Hotspot regions were defined as ≥3 for SCR events and ≥178 for non-SCR events in a window size of 4 Mb for rice, ≥2 for SCR events and ≥253 for non-SCR events in a window size of 10 Mb for soybean, and ≥4 for SCR events and ≥157 for non-SCR events in a window size of 1.2 Mb for *Arabidopsis*.

Motif and minimum free energy analysis in readthrough regions of SCR transcripts in maize

Transcription factor-binding motifs in readthrough regions of maize SCR transcripts were identified using the MEME-ChIP webserver (v.5.4.1)[68], and the annotations of the obtained motifs were searched against the available databases for comparison using Tomtom (v.5.0.5) with default parameters[69].

The minimum free energy of readthrough regions for maize SCR transcripts and the comparable length between the first and downstream second stop codons for non-readthrough transcripts were calculated using ViennaRNA Package (v.2.0) as described[70].

Identification and alignment of suppressor tRNAs in maize

The complete maize genome sequences were downloaded from the Ensembl database (ftp://ftp.ensemblgenomes.org/pub/plants/release-41/fasta/zea_mays/dna/). A genome-wide search for suppressor tRNA genes was conducted using tRNAscan-SE (v.2.0)[51] with default settings and ARAGORN[52] with both default settings and 0.95 default settings as a lower scoring threshold. This analysis generated a collection of suppressor/suppressor-like tRNAs recognizing the three stop codons from each search. Comparison of the canonical tRNAs from different species with these predicted suppressor tRNAs was performed using BLASTn (v.2.5.0+) against the NCBI GenBank nucleotide (NCBI NT) database as previously described[55,56]. The potential identity of the suppressor tRNAs corresponding to different amino acids could be predicted from the high similarity of the sequence alignment between the suppressor tRNAs and the canonical tRNAs of the standard amino acids.

Motif and structure prediction of SCR proteins in maize

C-terminal extensions at least 20 amino acids in length in maize SCR events were scanned for transmembrane helices using TMHMM (v.2.0)[71] with default settings and for nuclear localization signals using cNLS mapper[72]. C-terminal extensions at least 12 amino acids in length were searched for peroxisomal targeting signals using PTS1 Predictor[73] and prenylation signals using PrePS[74]. The 3D structures of ZmBTF3 and ZmBTF3x were predicted using Alphafold[75].

Plasmid construction

Luciferase constructs for the luminescence-based SCR assay were generated in the reconstructive pFN19K T7 SP6 flexi vector (Promega Corporation). The coding sequences of the tested genes, *ZmH3C2*, *ZmCEQORH*, *ZmBTF3* and *Zm00001d009138*, along with their canonical stop codons and inter-stop codon regions (ISRs, the extension regions) were cloned upstream of the coding sequence of the firefly luciferase gene (*Fluc*) without ATG between *EcoRI* and *KpnI* sites. A linker sequence (5′ GGCGGCTCCGGCGGCTCCCTCGTGCTCGGG 3′) was added upstream of the Fluc sequence to avoid potential distortions between the tested protein and Fluc in the fusion protein[15,31,38]. Primer sequences are listed in Supplementary

Table 15.

For subcellular localization analysis, the ZmBTF3 and ZmBTF3x coding

sequences were introduced into the pAN580 vector containing the 35S promoter and

the coding sequence for GFP. GFP was fused to the C-terminus of ZmBTF3 and

ZmBTF3x. Primer sequences are listed in Supplementary Table 15.

***In vitro* translation for SCR Assay**

For SCR assay, we used *in vitro* translation system based on Wheat Germ Extract

(Promega Corporation), which has been used for SCR validation as reported

previously[31,38]. The constructed plasmid DNA including the test gene and *Fluc* (no

ATG) was linearized using BamH I or XbaI enzyme, and 2 μg of the linearized DNA

was transcribed *in vitro* using T7 RNA polymerase (Promega Corporation). The

obtained RNA was then purified using GeneJET kit (Thermo Fisher Scientific). The

concentration and purity of the RNA were measured using Nanodrop 2000 (Thermo

Fisher Scientific). A total of 2 μg purified RNA was translated *in vitro* using the

Wheat Germ Extract at 25 °C for 2 h according to the manufacturer′s instructions.

Luciferase activity was then measured at room temperature (below 25 °C) using the

Luciferase Assay System (Promega Corporation) in a Glomax luminometer (Promega

Corporation) with a 10-s measurement read for luciferase activity.

Immunoblotting and mass spectrometric analysis

The proteins extracted from the *in vitro* protein expression system were separated by

10% (w/v) SDS-polyacrylamide gel electrophoresis (SDS-PAGE). One PAGE gel was

stained with Coomassie Brilliant Blue (CBB), while another was transferred to a

polyvinylidene difluoride membrane (0.22 μm, Millipore). Proteins were detected by

immunoblotting with anti-Fluc antibodies (1:10,000, ab185924, Abcam). The regions

of interest in the CBB staining gel were excised manually according to the

immunoblotting band, and then destained with 30% (v/v) ACN in 100 mM NH_4HCO_3

and dried in a vacuum centrifuge. The extracted proteins were reduced with

dithiothreitol (DTT) (10 mM DTT in 100 mM NH_4HCO_3) for 30 min at 56 °C, then

alkylated with iodoacetamide (200 mM IAA in 100 mM NH_4HCO_3) in the dark at

room temperature for 30 min. Gel pieces were briefly rinsed with 100 mM NH_4HCO_3

and ACN, respectively. The gel pieces were then divided equally into three parts and

separately digested overnight in 12.5 ng/μl trypsin, overnight in chymotrypsin in 25

mM NH_4HCO_3, and for 30 min in 12.5 ng/μl pepsin in PH3 HCl. The peptides were

extracted using 60% (v/v) ACN in 0.1% (v/v) trifluoroacetic acid (TFA) three times,

and the extracts were pooled and dried completely by a vacuum centrifuge.

LC-MS/MS was performed on a Q-Exactive HF-X mass spectrometer coupled to

Easy nLC 1200 (Thermo Fisher Scientific). The peptide mixture was loaded onto a

C18-reversed phase column (15 cm, 75 μm inner diameter) packed in-house with RP-

C18 (5 μm resin, Thermo Fisher Scientific) in buffer A (0.1% [v/v] formic acid in

HPLC-grade water) and separated with a linear gradient of buffer B (0.1% formic acid

in 84% [v/v] ACN) at a flow rate of 250 nl/min controlled by IntelliFlow technology

over 60 min. MS data were acquired using a data-dependent top-10 method

dynamically choosing the most abundant precursor ions from the survey scan (m/z

300-1,800) for HCD fragmentation. The target value was determined based on pAGC.

The dynamic exclusion duration was 20 s. Survey scans were acquired at a resolution

of 70,000 (m/z 200) and the resolution for HCD spectra was set to 17,500 (m/z 200).

Normalized collision energy was 27 eV and the underfill ratio, which specifies the

minimum percentage of the target value likely to be reached at maximum fill time,

was defined as 0.1%.

The MS data were then separately searched against the target protein sequences

(readthrough protein sequences of *H3C2*, *ZmCEQORH*, and *ZmBTF3*) using

MaxQuant software (v.1.6.1.10)[61]. An initial search was set with a precursor mass

window of 6 ppm. The search followed an enzymatic cleavage rule of "unspecified"

and a mass tolerance of 20 ppm for fragment ions. Carbamidomethylation of cysteines

was defined as fixed modification, while methionine oxidation was defined as

variable modification for database searching. The cutoff of the global FDR for peptide

and protein identification was set to 0.01.

Subcellular localization of ZmBTF3-GFP and ZmBTF3x-GFP fusions.

The GFP fusion plasmids were transformed into maize mesophyll protoplasts by

polyethylene glycol-mediated transformation. Protoplast preparation and transfection

were conducted as previously described[76]. A plasma membrane-localized marker

35S:ZmPDCK7-mCherry and a nucleus-localized marker *35S:AtHY5-mCherry* were

used for colocalization analyses[77,78]. Fluorescence signals were observed on a

confocal laser scanning microscope (LSM800, Zeiss).

Data availability

655 All data generated or analyzed in this study are included in this article or the

656 Supplementary Information and Tables. The raw datasets of proteomics have been

657 deposited at the ProteomeXchange under the accession ID PXD035958 and

658 PXD038762.

Code availability

660 Custom codes are available for research purposes from the corresponding authors

661 upon request.

References

1. Baranov, P. V., Atkins, J. F. & Yordanova, M. M. Augmented genetic decoding: global, local and temporal alterations of decoding processes and codon meaning. *Nat. Rev. Genet.* **16**, 517-529 (2015).

2. Freitag, J., Ast, J. & Bolker, M. Cryptic peroxisomal targeting via alternative splicing and stop codon read-through in fungi. *Nature* **485**, 522-525 (2012).

3. Bazykin, G. A. & Kochetov, A. V. Alternative translation start sites are conserved in eukaryotic genomes. *Nucleic Acids Res.* **39**, 567-577 (2011).

4. Jacks, T. & Varmus, H. E. Expression of the Rous sarcoma virus *pol* gene by ribosomal frameshifting. *Science* **230**, 1237-1242 (1985).

5. Lang, B. F. et al. Massive programmed translational jumping in mitochondria. *Proc. Natl Acad. Sci. USA* **111**, 5926-5931 (2014).

6. Sapkota, D. et al. Cell-type-specific profiling of alternative translation Identifies regulated protein isoform variation in the mouse brain. *Cell Rep.* **26**, 594-607.e597 (2019).

7. Rodnina, M. V. et al. Translational recoding: canonical translation mechanisms reinterpreted. *Nucleic Acids Res.* **48**, 1056-1067 (2020).

8. Palma, M. & Lejeune, F. Deciphering the molecular mechanism of stop codon readthrough. *Biol. Rev. Camb. Philos. Soc.* **96**, 310-329 (2021).

9. Torabi, N. & Kruglyak, L. Genetic basis of hidden phenotypic variation revealed by increased translational readthrough in yeast. *PLoS Genet.* **8**, e1002546 (2012).

10. Namy, O., Duchateau-Nguyen, G. & Rousset, J. P. Translational readthrough of the *PDE2* stop codon modulates cAMP levels in *Saccharomyces cerevisiae*. *Mol. Microbiol.* **43**, 641-652 (2002).

11. Dunn, J. G., Foo, C. K., Belletier, N. G., Gavis, E. R. & Weissman, J. S. Ribosome profiling reveals pervasive and regulated stop codon readthrough in *Drosophila melanogaster*. *eLife* **2**, e01179 (2013).

12. Xue, F. & Cooley, L. *kelch* encodes a component of intercellular bridges in

Drosophila egg chambers. *Cell* **72**, 681-693 (1993).

13. Karki, P. et al. Tissue-specific regulation of translational readthrough tunes
functions of the traffic jam transcription factor. *Nucleic Acids Res.* (2021).

14. Schueren, F. et al. Peroxisomal lactate dehydrogenase is generated by
translational readthrough in mammals. *eLife* **3**, e03640 (2014).

15. Eswarappa, S. M. et al. Programmed translational readthrough generates
antiangiogenic VEGF-Ax. *Cell* **157**, 1605-1618 (2014).

16. Xin, H., Zhong, C., Nudleman, E. & Ferrara, N. Evidence for pro-angiogenic
functions of VEGF-Ax. *Cell* **167**, 275-284.e276 (2016).

17. Weiner, A. M. & Weber, K. Natural read-through at the UGA termination
signal of Q-beta coat protein cistron. *Nat. New Biol.* **234**, 206-209 (1971).

18. Pelham, H. R. Leaky UAG termination codon in tobacco mosaic virus RNA.
Nature **272**, 469-471 (1978).

19. Yoshinaka, Y., Katoh, I., Copeland, T. D. & Oroszlan, S. Murine leukemia
virus protease is encoded by the *gag-pol* gene and is synthesized through
suppression of an amber termination codon. *Proc. Natl Acad. Sci. USA* **82**,
1618-1622 (1985).

20. True, H. L. & Lindquist, S. L. A yeast prion provides a mechanism for genetic
variation and phenotypic diversity. *Nature* **407**, 477-483 (2000).

21. Namy, O. et al. Identification of stop codon readthrough genes in
Saccharomyces cerevisiae. Nucleic Acids Res. **31**, 2289-2296 (2003).

22. Baudin-Baillieu, A. et al. Genome-wide translational changes induced by the
prion [PSI+]. *Cell Rep.* **8**, 439-448 (2014).

23. Jungreis, I. et al. Evidence of abundant stop codon readthrough in *Drosophila*
and other metazoa. *Genome Res.* **21**, 2096-2113 (2011).

24. Jungreis, I. et al. Evolutionary dynamics of abundant stop codon readthrough.
Mol. Biol. Evol. **33**, 3108-3132 (2016).

25. Hudson, A. M. et al. Tissue-specific dynamic codon redefinition in *Drosophila*.
Proc. Natl Acad. Sci. USA **118** (2021).

26.   Klagges, B. R. et al. Invertebrate synapsins: a single gene codes for several
721   isoforms in Drosophila. *J Neurosci.* **16**, 3154-3165 (1996).

722   27.   Steneberg, P. & Samakovlis, C. A novel stop codon readthrough mechanism
723   produces functional Headcase protein in *Drosophila* trachea. *EMBO Rep* 2,
724   593-597 (2001).

725   28.   Geller, A. I. & Rich, A. A UGA termination suppression tRNATrp active in
726   rabbit reticulocytes. *Nature* **283**, 41-46 (1980).

727   29.   Yamaguchi, Y. et al. L-MPZ, a novel isoform of myelin P0, is produced by
728   stop codon readthrough. *J. Biol. Chem.* **287**, 17765-17776 (2012).

729   30.   Manjunath, L. E. et al. Stop codon read-through of mammalian *MTCH2*
730   leading to an unstable isoform regulates mitochondrial membrane potential. *J.*
731   *Biol. Chem.* **295**, 17009-17026 (2020).

732   31.   Singh, A. et al. Let-7a-regulated translational readthrough of mammalian
733   *AGO1* generates a microRNA pathway inhibitor. *EMBO J.* **38**, e100727 (2019).

734   32.   Loughran, G. et al. Evidence of efficient stop codon readthrough in four
735   mammalian genes. *Nucleic Acids Res.* **42**, 8928-8938 (2014).

736   33.   De Bellis, M. et al. Translational readthrough generates new astrocyte AQP4
737   isoforms that modulate supramolecular clustering, glial endfeet localization,
738   and water transport. *Glia* **65**, 790-803 (2017).

739   34.   Stiebler, A. C. et al. Ribosomal readthrough at a short UGA stop codon context
740   triggers dual localization of metabolic enzymes in Fungi and animals. *PLoS*
741   *Genet.* **10**, e1004685 (2014).

742   35.   Loughran, G. et al. Stop codon readthrough generates a C-terminally extended
743   variant of the human vitamin D receptor with reduced calcitriol response. *J.*
744   *Biol. Chem.* **293**, 4434-4444 (2018).

745   36.   Yordanova, M. M. et al. *AMD1* mRNA employs ribosome stalling as a
746   mechanism for molecular memory formation. *Nature* **553**, 356-360 (2018).

747   37.   Yordanova, M. M., Loughran, G., Atkins, J. F. & Baranov, P. V. Stop codon
748   readthrough contexts influence reporter expression differentially depending on

the presence of an IRES. *Wellcome Open Res.* **5**, 221 (2020).

38.   Sahoo, S. et al. Identification and functional characterization of mRNAs that

exhibit stop codon readthrough in *Arabidopsis thaliana*. *J. Biol. Chem.* **298**,

102173 (2022).

39.   Blanchet, S., Cornu, D., Argentini, M. & Namy, O. New insights into the

incorporation of natural suppressor tRNAs at stop codons in *Saccharomyces*

cerevisiae. *Nucleic Acids Res.* **42**, 10061-10072 (2014).

40.   Feng, Y. X., Copeland, T. D., Oroszlan, S., Rein, A. & Levin, J. G.

Identification of amino acids inserted during suppression of UAA and UGA

termination codons at the *gag-pol* junction of Moloney murine leukemia virus.

Proc. Natl Acad. Sci. USA **87**, 8860-8863 (1990).

41.   Roy, B., Leszyk, J. D., Mangus, D. A. & Jacobson, A. Nonsense suppression

by near-cognate tRNAs employs alternative base pairing at codon positions 1

and 3. *Proc. Natl Acad. Sci. USA* **112**, 3038-3043 (2015).

42.   Strigini, P. & Brickman, E. Analysis of specific misreading in *Escherichia coli*.

J. Mol. Biol. **75**, 659-672 (1973).

43.   Sánchez-Silva, R., Villalobo, E., Morin, L. & Torres, A. A new noncanonical

nuclear genetic code: translation of UAA into glutamate. *Curr. Biol.* **13**, 442-

447 (2003).

44.   Blanchet, S. et al. Deciphering the reading of the genetic code by near-cognate

tRNA. *Proc. Natl Acad. Sci. USA* **115**, 3018-3023 (2018).

45.   Choi, S. et al. Proteogenomic Approach to UTR Peptide Identification. *J*

Proteome Res. **19**, 212-220 (2019).

46.   Aerni, H. R., Shifman, M. A., Rogulina, S., O'Donoghue, P. & Rinehart, J.

Revealing the amino acid composition of proteins within an expanded genetic

code. *Nucleic Acids Res.* **43**, e8 (2015).

47.   Valle, R. P., Morch, M. D. & Haenni, A. L. Novel amber suppressor tRNAs of

mammalian origin. *EMBO J.* **6**, 3049-3055 (1987).

48.   Namy, O., Hatin, I. & Rousset, J. P. Impact of the six nucleotides downstream

of the stop codon on translation termination. *EMBO Rep.* **2**, 787-793 (2001).

49.   Tork, S., Hatin, I., Rousset, J. P. & Fabret, C. The major 5' determinant in stop

codon read-through involves two adjacent adenines. *Nucleic Acids Res.* **32**,

415-421 (2004).

50.   Mangkalaphiban, K. et al. Transcriptome-wide investigation of stop codon

readthrough in *Saccharomyces cerevisiae. PLoS Genet.* **17**, e1009538 (2021).

51.   Chan, P. P., Lin, B. Y., Mak, A. J. & Lowe, T. M. tRNAscan-SE 2.0: improved

detection and functional classification of transfer RNA genes. *Nucleic Acids*

Res. **49**, 9077-9096 (2021).

52.   Laslett, D. & Canback, B. ARAGORN, a program to detect tRNA genes and

tmRNA genes in nucleotide sequences. *Nucleic Acids Res.* **32**, 11-16 (2004).

53.   Ding, Y. et al. OST1-mediated BTF3L phosphorylation positively regulates

CBFs during plant cold responses. *EMBO J.* **37**, *e98228* (2018).

54.   Atkins, J. F. & Baranov, P. V. The distinction between recoding and codon

reassignment. *Genetics* **185**, 1535-1536 (2010).

55.   Ivanova, N. N. et al. Stop codon reassignments in the wild. *Science* **344**, 909-

913 (2014).

56.   Swart, E. C., Serra, V., Petroni, G. & Nowacki, M. Genetic codes with no

dedicated stop codon: context-dependent translation termination. *Cell* **166**,

691-702 (2016).

57.   Beier, H. & Grimm, M. Misreading of termination codons in eukaryotes by

natural nonsense suppressor tRNAs. *Nucleic Acids Res.* **29**, 4767-4782 (2001).

58.   Dabrowski, M., Bukowy-Bieryllo, Z. & Zietkiewicz, E. Translational

readthrough potential of natural termination codons in eucaryotes--The impact

of RNA sequence. *RNA Biol.* **12**, 950-958 (2015).

59.   Schimmel, P. The emerging complexity of the tRNA world: mammalian tRNAs

beyond protein synthesis. *Nat. Rev. Mol. Cell Biol.* **19**, 45-58 (2018).

60.   Wiśniewski, J. R., Zougman, A., Nagaraj, N. & Mann, M. Universal sample

preparation method for proteome analysis. *Nat. Methods* **6**, 359-362 (2009).

61. Tyanova, S., Temu, T. & Cox, J. The MaxQuant computational platform for
mass spectrometry-based shotgun proteomics. *Nat. Protoc.* **11**, 2301-2319
(2016).

62. Lei, L. et al. Ribosome profiling reveals dynamic translational landscape in
maize seedlings under drought stress. *Plant J.* **84**, 1206-1218 (2015).

63. Chotewutmontri, P. & Barkan, A. Dynamics of chloroplast translation during
chloroplast differentiation in maize. *PLoS Genet.* **12**, e1006106 (2016).

64. Chotewutmontri, P. & Barkan, A. Multilevel effects of light on ribosome
dynamics in chloroplasts program genome-wide and psbA-specific changes in
translation. *PLoS Genet.* **14**, e1007555 (2018).

65. Zoschke, R., Chotewutmontri, P. & Barkan, A. Translation and co-translational
membrane engagement of plastid-encoded chlorophyll-binding proteins are
not influenced by chlorophyll availability in maize. *Front. Plant. Sci.* **8**, 385
(2017).

66. Jiang, J. et al. An RNA chaperone-like protein plays critical roles in
chloroplast mRNA stability and translation in *Arabidopsis* and maize. *Plant*
Cell **31**, 1308-1327 (2019).

67. Dobin, A. et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**,
15-21 (2013).

68. Machanick, P. & Bailey, T. L. MEME-ChIP: motif analysis of large DNA
datasets. *Bioinformatics* **27**, 1696-1697 (2011).

69. Gupta, S., Stamatoyannopoulos, J. A., Bailey, T. L. & Noble, W. S.
Quantifying similarity between motifs. *Genome Biol.* **8**, R24 (2007).

70. Lorenz, R. et al. ViennaRNA Package 2.0. *Algorithms Mol. Biol.* **6**, 26 (2011).

71. Krogh, A., Larsson, B., von Heijne, G. & Sonnhammer, E. L. Predicting
transmembrane protein topology with a hidden Markov model: application to
complete genomes. *J. Mol. Biol.* **305**, 567-580 (2001).

72. Kosugi, S., Hasebe, M., Tomita, M. & Yanagawa, H. Systematic identification
of cell cycle-dependent yeast nucleocytoplasmic shuttling proteins by

prediction of composite motifs. *Proc. Natl Acad. Sci. USA* **106**, 10171-10176

(2009).

73.   Neuberger, G., Maurer-Stroh, S., Eisenhaber, B., Hartig, A. & Eisenhaber, F.

Prediction of peroxisomal targeting signal 1 containing proteins from amino

acid sequence. *J. Mol. Biol.* **328**, 581-592 (2003).

74.   Maurer-Stroh, S. & Eisenhaber, F. Refinement and prediction of protein

prenylation motifs. *Genome Biol.* **6**, R55 (2005).

75.   Jumper, J. et al. Highly accurate protein structure prediction with AlphaFold.

Nature **596**, 583-589 (2021).

76.   Yoo, S. D., Cho, Y. H. & Sheen, J. *Arabidopsis* mesophyll protoplasts: a

versatile cell system for transient gene expression analysis. *Nat. Protoc.* **2**,

1565-1572 (2007).

77.   Zhao, Y. et al. The calcium-dependent protein kinase ZmCDPK7 functions in

heat-stress tolerance in maize. *J. Integr. Plant. Biol.* **63**, 510-527 (2021).

78.   Li, Y. et al. Plasma membrane-nucleo-cytoplasmic coordination of a receptor-

like cytoplasmic kinase promotes EDS1-dependent plant immunity. *Nat.*

Plants **8**, 802-816 (2022).

Acknowledgments

This study was supported by the National Natural Science Foundation of China (no. 32172073, U22A20474, 31872872), the National Key Research and Development Program of China (2022YFD1201802), and the Seed-Industrialized Development Program in Shandong Province (no. 2021LZGC003). We are grateful to Steven P. Briggs and David Backhouse for their constructive comments and suggestions. We also thank Dr. Anguo Sun and Dr. Yanwen Xiang for the technical assistance and Yingying Sha and Mengxia Niu for their assistance with the microscopic observations.

Author Contributions

L.W., S.-B.W., and Z.T. designed the project. Y. Zhang, Y.S., L.T., H.Y., A.X., J.Z., U.A., A.S., X.C. and Y.C. were involved in bioinformatics analysis. Y. Zhang, H.L., S.W. and J.S. conducted the experiments. C.G., Y. Zhao, Y. L. and X.W. collected the materials for mass spectrometry detection and subsequent experiments. Y. Zhang, H.L. and Y.S. wrote the manuscript. L.W., S.-B.W., and Z.T. contributed substantially to revisions. All authors read and approved the manuscript.

Competing interests

The authors declare no competing interests.

Figures

Fig. 1 | Proteogenomic workflow for identifying plant stop codon readthrough (SCR) events. The strategy consists of two procedures: shotgun proteomics experiment (green box) and customized database construction (pink boxes). Total protein was extracted from plant leaves and subjected to digestion, fragmentation, and LC-MS/MS analysis for identification of spectra datasets. For customized database construction, the plant genome sequences were downloaded from public databases and subjected to *in silico* translation. The annotated stop codon sites were replaced with "X", representing one of the 20 standard amino acids during translation each time. The translation process was assigned to terminate at the next in-frame stop codon, producing 20 customized databases containing theoretical readthrough proteins with the annotated stop sites corresponding to the 20 amino acids. The distal 3′ UTR represents the portion of the 3′ UTR downstream of the putative readthrough extension. MS/MS spectra datasets were searched against the 20 customized databases merged with the public plant protein database by the MaxQuant. We defined the MS/MS-identified peptides that exclusively span the annotated stop codon sites as "through" peptides, and classified proteins sharing a unique stop codon site and containing a "through" peptide as one SCR event.

Fig. 2 | Verifying SCR events in maize. a Three-nucleotide periodicity of 30-mers in the CDSs, putative extensions, and the distal 3′ UTRs. Data are mean ± s.d. Different letters indicate significant differences at $P < 0.05$ by one-way ANOVA with Tukey's multiple comparisons test. Green, light green, and gray bars represent frame 0, 1, and 2, respectively. **b** Distribution of ribosome footprint density in putative extensions and the distal 3′ UTRs in the four Ribo-seq datasets. Distal 3′ UTR was defined as the region 40 codons after the next stop-codon or the region from the next stop-codon to the third stop codon within 40 codons. Box edges represent the 0.25 and 0.75 quantiles. The bold lines represent the median and whiskers represent 1.5×IQR. Statistical significance was determined using a two-sided Student's *t*-test,

P < 0.001. **c** The proportion of SCR events confirmed by ribosome profiling among those identified by proteogenomics. **d-k** Validation of two readthrough examples, *ZmH3C2* (**d-g**) and *ZmCEQORH* (**h-k**). **d, h** The complete protein sequences extended to the next in-frame stop codon of the two genes showing the underlined peptides identified by mass spectrometry analysis, and the readthrough peptides are in bold. The amino acids translated at the stop codon sites in readthrough peptides are highlighted in red, which are alanine (A) at UAG (**d**) and glutamate (E) at UGA (**h**). **e, i** Ribo-seq analysis revealing clear footprint signals in the C-terminal extensions of the two transcripts. Dashed lines represent the annotated stop codon and the next in-frame stop codon. **f, j** *In vitro* translation experiment showing readthrough activities of *ZmH3C2* and *ZmCEQORH*. Plasmids containing test cDNA-stop codon-ISR-*Fluc* were firstly transcribed *in vitro*, followed by *in vitro* translation using wheat germ extract. Luciferase activities were used to represent the readthrough activities. Constructs with the stop codon mutated to a sense codon were used as a positive control of the translation to measure the efficiency of SCR. The non-SCR gene *Zm00001d009138* with its TAG and ISR sequence fused with *Fluc* (*Zm00001d009138-TAG-ISR-Fluc*) was used as the SCR-negative control, as it showed no SCR sign in either proteogenomics or Ribo-seq analysis. All samples were tested in biological triplicates. Data are mean ± s.e.m (n=3). Asterisks indicate statistical significance by two-sided Student's *t*-test (P < 0.001). ISR, inter-stop codon region (extension region). **g, k** MS/MS spectra of the identified readthrough peptides AAAAGRPDRPWR for *ZmH3C2* (**g**) and GGAAEETR for *ZmCEQORH* (**k**) readthrough products expressed from (**f**) and (**j**), respectively.

Fig. 3 | Characteristics of SCR events in maize. a Stop codon usage and the next stop codon

preference in SCR events. Stop codon 1, readthrough stop codon; Stop codon 2, the next in-

frame stop codon. **b** Genome-wide distribution of SCR (pink) and non-SCR (green) events in

maize. For each chromosome, the distribution contains three columns. Left: SCR (pink) and

non-SCR (green) events across the chromosomes. Black dots are centromeres. Middle:

distribution patterns of SCR (pink) and non-SCR (green) events are shown by curves, shown

as a 20-Mb sliding window with a 2-Mb step based on the B73 reference genome. Right:

distribution of hotspot regions for SCR (pink) and non-SCR (green) events shown with

asterisks. We defined hotspot regions as ≥5 SCR events and ≥335 non-SCR events in a 20-Mb

window. **c** Violin plot describing the length distribution of primary transcripts and transcript

elements in SCR and non-SCR events. The length of the primary transcript is the sum of

CDSs, introns, and UTRs. CDS length represents the length of the open reading frame (ORF).

Internal boxplots show the median values (white dots in the center) and the interquartile range

(IQR; 0.25 and 0.75 quantiles) of the length distribution. **d** Number of exons in SCR and non-

SCR primary transcripts. **e** Number of splice variants in SCR and non-SCR primary

transcripts. For all boxplots, box edges represent the 0.25 and 0.75 quantiles. The bold lines

represent the median values and the whiskers represent 1.5×IQR. ***$P < 0.001$ (two-sided

Student's *t*-test).

Fig. 4 | Stop codons are recoded as 20 amino acids in maize SCR events. a Frequency of amino acids specified by the three stop codons. The x-axis represents the 20 standard amino acids. The y-axis is the number of SCR events. **b** Previously reported amino acids translated at UGA, UAG, and UAA in different species. Blue, virus; orange, bacteria; light blue, fungi; yellow, animal; green, algae. **c** Examples of UGA-, UAG-, and UAA suppressor tRNAs identified by tRNAscan-SE and ARAGORN corresponding to amino acids shown at UGA, UAG, and UAA. The predicted secondary structures of the tRNAs are shown (I-VI): tRNA^Met (UCA) and tRNA^Lys (UCA) recognize UGA, tRNA^Glu (CUA) and tRNA^Trp (CUA) recognize UAG, and tRNA^Lys (UUA) and tRNA^His (UUA) recognize UAA.

960 three stop codons in rice (**g**), soybean (**h**), and *Arabidopsis* (**i**) SCR events. The x-axis

961 represents the 20 standard amino acids. The y-axis is the number of SCR events.

Fig. 6 | C-terminal extensions in maize SCR events contain multiple functional signals. a Multiple functional signals predicted in the extensions of SCR events. **b** Readthrough validation of *ZmBTF3* using the *in vitro* translation experiment. The plasmid containing *ZmBTF3-TAA-ISR-Fluc* was firstly transcribed *in vitro*, followed by *in vitro* translation using wheat germ extract. *ZmBTF3* with TAA mutated to ACA was used as the positive control of the translation to measure the efficiency of SCR. The *Zm00001d009138-TAG-ISR-Fluc* was used as the SCR-negative control. Luciferase activity reflects the readthrough activity. All experiments were performed in triplicates. Data are mean ± s.e.m (n=3). Statistical difference was calculated using a two-sided Student's *t*-test, ***$P < 0.001$. ISR, inter-stop codon region. **c** SDS-PAGE and immunoblot analysis of the ZmBTF3x-Fluc (ZmBTF3-TAA-ISR-Fluc), ZmBTF3x^ACA-Fluc (ZmBTF3-ACA-ISR-Fluc), and ZmBTF3-Fluc fusion proteins. Top:

Coomassie Brilliant Blue (CBB)-stained gel of the expressed products derived from (**b**).

Bottom: Immunoblot of the ZmBTF3x-Fluc, ZmBTF3x[ACA]-Fluc, and ZmBTF3-Fluc fusion

proteins using an anti-Fluc antibody. The ZmBTF3x[ACA]-Fluc was used as the positive control.

The Zm00001d009138-TAG-ISR-Fluc was used as the negative control. ISR, inter-stop codon

region. **d** MS/MS analysis of the digested peptides of the ZmBTF3x-Fluc fusion protein

excised from the CBB gel in (**c**). The sequences of MS-detected peptides, labeled as "before",

"through", and "after", are shown on the top. An MS/MS spectrum of the "through" peptide

EPEEKKESTFCCV is shown on the bottom. UAA was recoded as threonine (T) shown in red.

e Transmembrane domain prediction in the C-terminus of ZmBTF3x using the TMHMM

server. The blue, red, and magenta lines represent residues 1-201 inside the membrane, 202-

221 as a transmembrane helix, and 222-223 outside the membrane, respectively. The amino

acid sequence from the annotated stop codon to the next stop codon of BTF3x is shown on the

top. **f** AlphaFold structural models for ZmBTF3 and its readthrough protein ZmBTF3x. Green:

ZmBTF3; dark blue: parent protein region of the readthrough protein ZmBTF3x; magenta: C-

terminal extension region of ZmBTF3x. The predicted α-helix is shown at the C-terminus. **g**

Subcellular localization of ZmBTF3 and ZmBTF3x. Constructs encoding C-terminal GFP

fusions of ZmBTF3 or ZmBTF3x were transfected into maize protoplasts for localization

assays under a confocal microscope. GFP-vector transfected protoplasts were used as the

control. The experiments were performed three times with similar results. HY5-mCherry was

the nucleus marker and PDCK7-mCherry was the plasma membrane marker. Scale bars, 5 μm.

Supplementary Information, Tables and Dataset have been submitted

as separate files.

Supplementary Information contains Supplementary Figures 1-10 and

References for Supplementary Table 5

Supplementary Table 1. SCR events identified in maize by proteogenomics.

Supplementary Table 2. Peptides identified by mass spectrometry for SCR and

parent proteins in maize.

Supplementary Table 3. SCR events detected in maize by both proteogenomics

and ribosome profiling.

Supplementary Table 4. Non-SCR events identified in maize by proteogenomics.

Supplementary Table 5. Standard amino acids translated by stop codons in the

literature and their translation evidence in maize. I, UGA; II, UAG; III, UAA.

UGA-, UAG-, and UAA suppressor tRNAs were predicted by tRNAscan-SE or

ARAGORN.

Supplementary Table 6. Predicted suppressor tRNAs in maize aligned with

tRNAs in the NCBI NT database.

Supplementary Table 7. SCR events identified in rice by proteogenomics.

Supplementary Table 8. Non-SCR events identified in rice by proteogenomics.

Supplementary Table 9. SCR events identified in soybean by proteogenomics.

Supplementary Table 10. Non-SCR events identified in soybean by

proteogenomics.

Supplementary Table 11. SCR events identified in *Arabidopsis* by proteogenomics.

Supplementary Table 12. Non-SCR events identified in *Arabidopsis* by

proteogenomics.

Supplementary Table 13. Functional signals predicted in C-terminal extensions

of maize SCR events.

Supplementary Table 14. Primers used in this study.

1022

1023 **Supplementary Dataset 1: Motifs predicted by Tomtom in the C-terminal**

1024 **extensions of SCR transcripts in maize.**