1 ***Metabolic dynamics of the coral-algal symbiosis from fertilization to settlement identify***
2 ***critical coral energetic vulnerabilities***
3
4 **Authors and Author Affiliations**
5

6 Ariana S. Huffmyer[1,2,6]*, Kevin H. Wong[3], Danielle M. Becker[2], Emma Strand[4], Tali Mass[5], Hollie
7 M. Putnam[2]
8
[1] School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA, USA
[2] Department of Biological Sciences, University of Rhode Island, Kingston, RI, USA
[3] Rosenstiel School of Marine and Atmospheric Science, Department of Marine Biology and
Ecology, University of Miami, Miami, FL, USA
[4] Gloucester Marine Genomics Institute, Gloucester, MA, USA
[5]Department of Marine Biology, The Leon H. Charney School of Marine Sciences, University of
Haifa, Mt. Carmel, Haifa, Israel
[6] Lead contact ashuffmyer@gmail.com
17
18 * Correspondence: Ariana S. Huffmyer, ashuffmyer@gmail.com

19
20 **Summary**
21

Climate change accelerates coral reef decline and jeopardizes recruitment essential for
ecosystem recovery. Adult corals rely on a vital nutritional exchange with their symbiotic algae
(Symbiodiniaceae), but the dynamics and sensitivity of this reliance from fertilization to
recruitment are understudied. We investigated the physiological, metabolomic, and
transcriptomic changes across 13 developmental stages of *Montipora capitata*, a coral in
Hawaiʻi that inherits symbionts from parent to egg. We found that embryonic development
depends on maternally provisioned mRNAs and lipids, with a rapid shift to symbiont-derived
nutrition in swimming larvae. Symbiont density and photosynthesis peak once swimming to fuel
pelagic larval dispersal. In contrast, respiratory demand increases significantly during
metamorphosis, settlement, and calcification, reflecting this energy-intensive morphological
reorganization. Symbiont ontogenetic proliferation is driven by symbiont ammonium assimilation
with little evidence of nitrogen metabolism in the coral host. As development progresses, the
host enhances nitrogen sequestration, regulating symbiont populations, and ensuring the
transfer of fixed carbon to support metamorphosis, with both metabolomic and transcriptomic
indicators of increased carbohydrate availability. Although algal symbiont communities remained
stable, bacterial communities shifted with ontogeny, associated with holobiont metabolic
reorganization. Our study reveals extensive metabolic changes during development,
increasingly reliant on symbiont nutrition. Metamorphosis and settlement emerge as the most
critical periods of energetic vulnerability to projected climate scenarios that destabilize
symbiosis. This highly detailed elucidation of symbiotic nutritional exchange relative to sensitive
early life stages provides essential knowledge for understanding and forecasting nutritional
symbiosis integration, and specifically, coral survival and recruitment in a future of climate
change.

Introduction

Symbiotic relationships between animals and microbial organisms enable survival in nutritionally challenging environments [1–4]. The function and structure of one of the most ecologically and economically important ecosystems on Earth, coral reefs, is driven by the productivity of marine symbioses. The nutritional symbiosis between reef-building corals and endosymbiotic algae in the family Symbiodiniaceae [5] can provide >100% of the coral's daily metabolic demands through translocated organic carbon generated through symbiont photosynthesis [6,7]. In return, metabolic waste from the coral host, including nitrogen and inorganic carbon, is available to Symbiodiniaceae to support their cellular metabolism and population growth [8,9]. The energy-enhancing nutrient exchange between Symbiodiniaceae and corals, as well as their diverse bacterial and archaeal microbiome [10], enables coral biomineralization and thus the formation of the 3D skeletal structure that provides a habitat for high biodiversity in coral reef ecosystems [11] and their resulting invaluable goods and services for humans [12]. However, the delicate coral-Symbiodiniaceae symbiosis is susceptible to destabilization by environmental stress. Rising sea surface temperatures and acute marine heat waves cause coral bleaching, or the dysbiosis and loss of Symbiodiniaceae cells and/or photosynthetic pigments, and reduced translocation of carbon surplus to the coral [13], which can, and often does, culminate in coral starvation and mass mortality [14]. Increasingly, the stability of the nutritional relationship between corals, their algal endosymbionts, and symbiotic microbes is critical for the continued survival of coral reef ecosystems and the maintenance of essential ecosystem services humans rely on.

Globally, half of the live coral cover on reefs has been lost since the 1950s [15], significantly increasing the importance of coral reproduction and recruitment to replenish and maintain reef ecosystems. Corals have complex and biphasic life cycles, in which sessile adults produce gametes that are internally or externally fertilized and develop into swimming larvae that then undergo metamorphosis into the sessile polyp recruit stage and grow into mature colonies attached to the benthos [16]. Survivorship from fertilization to recruitment represents the primary bottleneck for marine larvae [17], due to environmental impacts on fertilization, larval dispersal and behavior, predation, substrate suitability, and post-settlement survival [16]. Key among the myriad challenges facing these sensitive life stages is the energetic requirement for successful development, metamorphosis, and calcification [18]. To buffer the energetic challenges of living in oligotrophic waters, corals formed associations with Symbiodiniaceae ~250 mya [19]. This symbiotic nutritional exchange is initiated either immediately through vertical transmission (i.e., symbionts are provisioned to the eggs), or within days to weeks of release and/or settlement through the process of horizontal transmission (i.e., symbionts are acquired from the environment) [20]. Initiation of symbiosis is a topic of intense study [21], but despite its increasing importance to reef persistence and recovery [22], the contribution of symbiont-derived nutrition to

support energy demands of early life history are not well characterized. Broadly, an improved
understanding of the development of symbiotic relationships in foundational marine
invertebrates, like corals, provides critical insight on how environmental change may impact
nutritional symbioses that are widespread across organisms and ecosystems.
Coral embryos and larvae initially metabolize limited endogenous lipid reserves [23],
supporting the paradigm that parental provisioning is the primary energetic fuel for early life
stages [24–26]. For example, high-performance thin-layer chromatography lipid analysis in
aposymbiotic life stages of corals (i.e., larvae) indicates a decline in wax ester abundance as
larvae age and a strong reliance on parentally provisioned fatty acids for energetic fuel [26].
Additionally, lipid usage in *Acropora tenuis* larvae without symbionts is highest within optimum
thermal temperatures, suggesting that lipid stores are an important energy source for larval
development and that dispersal potential will shift as ocean temperatures increase [25]. However,
in symbiotic larvae, the demand for energy fueled by lipid reserves has the potential to be
substantially supplemented by photosynthetically derived carbon translocation. Radiolabel
tracing in symbiotic *Pocillopora damicornis* larvae demonstrated that ~70% of photosynthetically
fixed carbon is translocated to the host [27]. Further, the use of lipid reserves slows under light
conditions in symbiotic *P. damicornis* and *Montipora digitata* larvae with lipids accounting for 38-
41% of metabolic energy in the light compared to 84-90% of energy in the dark [28]. Similarly,
symbiotic *A. tenuis* larvae exhibited decreased lipid consumption compared to those that did not
have symbionts, increasing dispersal potential and the ability of larvae to settle in favorable
conditions [29]. In contrast, high resolution local scale translocation using nanoscale secondary
ion mass spectrometry (NanoSIMS) imaging combined with stable isotope in symbiotic *P.*
damicornis larvae identified that the energetic contribution of photosynthates was systematically
lower across tissues in larvae compared to the adult stage [30]. These contrasting studies
highlight the need to clarify the dynamics of nutritional exchange across early life history.
Recent evidence using stable isotope metabolomic tracing confirms that
photosynthetically fixed carbon in the form of a major photosynthate, glucose, can be
translocated to symbiotic larvae at approximately 7 days old (165 hours post fertilization) in
Montipora capitata [31]. Under thermal stress, *M. capitata* larvae increased nitrogen assimilation
and sequestration to maintain glucose translocation, demonstrating that symbiotic relationships
require metabolic investment to maintain [31]. This previous research demonstrates that symbiotic
relationships can contribute to nutritional requirements in coral larvae and have the potential to
affect performance in coral larvae. However, it remains unclear at which point along
development symbiotic nutritional exchange begins and the timing of transitions from reliance
on lipids to active metabolism of photosynthates. Considering the need to increase our
understanding of the sensitive coral symbiosis and energetic demands of recruitment, we
examined physiological, molecular, and metabolomic data to characterize ontogenetic shifts in
central metabolic pathways in early life stages of the vertically transmitting coral, *M. capitata*
(Fig 1). Considering the energetic consequences of nutritional symbioses across taxa, here we
examined an archetypal symbiotic nutritional model of reef-building corals to characterize: (1)
symbiotic nutritional exchange relative to energetic demands across development,
metamorphosis and settlement; and (2) mechanisms and dynamics of the transition from
reliance on parentally provisioned lipid reserves to active utilization of symbiont nutrition.
132

Results

A. Ontogenetic shifts in physiology, metabolic rates, and endosymbiont photosynthesis

We tracked size, respiratory demand, symbiont density, and photosynthesis to characterize shifts in metabolic production and demand across development (65-231 hours post fertilization, hpf) (Fig 1AB). There were significant changes in individual size across development (Kruskal-Wallis, $\chi^2(9,562)=235.86$, P<0.001; Fig 2A). Embryo size decreased 41% as development progressed from egg (1 hpf) through larval stages as corals went through size contraction associated with the transition from the late gastrula (65 hpf) to planula (93 hpf) stages (Fig 2A). Size then increased, reaching a maximum at 231 hpf (142% growth relative to egg size; Fig 2A) in larvae and metamorphosed polyps.

Size-normalized respiration rates also changed across ontogeny (Kruskal-Wallis, $\chi^2(5,91)=55.68$, P<0.001), decreasing steadily from 65 hpf embryos to the lowest observed value at 231 hpf in larvae (Fig 2B). Respiratory demand then increased sharply in metamorphosed polyps, at 183-231 hpf, with rates significantly higher than swimming larvae of the same age (Fig 2B). The largest difference was seen at 183 hpf with respiration rates 83% higher than swimming larvae of the same age (Fig 2B). Similarly, size-normalized gross photosynthesis rates exhibited a significant decrease across development (Kruskal-Wallis, $\chi^2(5,91)=34.98$, P<0.001), with the lowest photosynthesis rates at 231 hpf in both swimming larvae and metamorphosed polyps (Fig 2C). Comparison of the ratio of oxygen production to oxygen demand was calculated as P:R ratio, where values >1 indicates a surplus of oxygen produced. We found that P:R ratios changed significantly across development (Kruskal-Wallis, $\chi^2(5,91)=38.91$, P<0.001), characterized by an initial increase from embryos at 65 hpf to larvae at 183 hpf, peaking at 2.71 ± 0.18 (mean ± standard error of mean; Fig 2D). After this point in development, P:R ratios decreased significantly (Fig 2D), reaching a minimum (1.19 ± 0.16) in metamorphosed polyps (183-231 hpf), driven by elevated rates of respiratory demand at this stage (Fig 2B).

Endosymbiont populations increased across development when calculated as total cells per individual (ANOVA, SS=1.02e8, F(9,30)=25.06, P<0.001), peaking in the oldest life stages characterized (larvae and metamorphosed polyps at 231 hpf) (Fig S1). Between the earliest larval stage and the latest metamorphosed polyp stage, there was a 138% increase in cell densities per individual (post hoc P<0.001). Because the size of life stages changed across development and cells per individual correlated positively with size (Spearman's correlation, S=4876, rho=0.543, P<0.001), we further calculated cells per unit volume (cells mm^3). There was a significant change in size-normalized cell density between time points (ANOVA, SS=5.29e10, F(9,30)=21.28, P<0.001) (Fig 2E). Size-normalized cell densities were highest between 65-93 hpf (post hoc P<0.05). Densities at 65 hpf were 300% higher than density at the previous stage of 38 hpf and population density decreased by 43% from 93 hpf to 163 hpf (Fig 2E). There was a trend for increased cell densities within attached recruit stages (183, 231, and 255 hpf), but this was not statistically significant (Kruskal-Wallis, $\chi^2(2)=5.56$, *P*=0.062; Fig 2F). Due to different units of size normalization, we cannot directly compare symbiont densities between attached recruits and earlier stages.

B. Symbiotic and microbial community composition across development

Our samples contained a mixture of *Cladocopium* sp. (ITS2 Profile
C31.C17d.C21.C21ac.C31a; [32]) (56.0 ± 0.42%) and *Durusdinium glynnii* (ITS2 Profile
D1.D4.D6.D1ab.D17d; [33]) (44.0 ± 0.42%). Symbiont community similarity at the level of Defining
Intragenomic Variants (DIVs [34]) significantly changed across development (PERMANOVA,
SS=0.09, DF=9, R^2=0.50, F=2.97, P=0.001; Fig S2AB) but there was no change in alpha
diversity (Shannon and Inverse Shannon metrics, P>0.05 for all metrics). We calculated the
ratio of relative abundance of *Cladocopium* sp. and *Durusdinium glynnii* (C:D ratio) across
development. We found a significant effect of time point on C:D ratios (one-way ANOVA,
SS=0.38, DF=9, F=3.06, P=0.011) (Fig S2C); however, there was only a significant difference in
C:D ratios found between eggs (1 hpf) and one larval (231 hpf stage (increased C:D ratio; post
hoc P=0.025; Fig S2C) with no other significant pairwise differences (P>0.05; Fig S2C). Relative
abundance of C31 was elevated at 231 hpf in larvae compared to eggs at 1 hpf, specifically (Fig
S2B).
We further investigated bacterial community composition in *M. capitata* early life stages.
However, due to lower than anticipated sequencing depth in our samples, we constrained our
analysis at the taxonomic level of genus, when available, or family and above. Bacterial
communities varied significantly across development (PERMANOVA, SS=4.879, R^2=0.643,
F(7,17)=4.378, P=0.001) (Fig S3A), and alpha diversity peaked during metamorphosis and
recruitment (one-way ANOVA; Shannon: SS=20.247, F(7,17)=22.31, P<0.001; Inverse
Simpson: SS=1.69e4, F(7,17)=17.46, P<0.001) (Fig S3B). Specifically, we found that
Alteromonadaceae (*Alteromonas* sp.), *Vibrio* sp., and Pseudoalteromonadaceae
(*Psychrosphaera* sp.) peaked early in development followed by Oceanospiriliaceae
(unclassified) increasing in larval stages (Fig S3CD). In larvae and recruits, the relative
abundance of Rhodobacteraceae and Flavobacteriaceae also increased (Fig S3CD).

C. Ontogenetic shifts in gene expression

There were strong shifts in gene expression across development (PERMANOVA;
SS=3.19e5, R^2=0.75, F(9,28)=9.36, P=0.001), as characterized from all genes that were
expressed and passed filtering thresholds (*n* = 11,475 genes). Gene expression displayed three
clear ontogenetic groupings in early (eggs at 1 hpf through embryos at 38 hpf), mid (embryos at
65 hpf through larvae at 163 hpf) and late developmental stages (larvae at 183 hpf through
attached recruits at 255 hpf) (Fig 3A). 66% of the variance in gene expression was explained by
the separation of early developmental stages (PC1; Fig 3A) with 16% of the variance explained
by the separation between mid and late developmental groupings (PC2; Fig 3A). Of the 11,475
genes expressed, 9,181 were detected as differentially expressed (DEG) through likelihood ratio
tests (Fig S4). Expression patterns of the DEGs across development (i.e., clusters) were then
detected through hierarchical clustering analyses with 64 total clusters identified that contained
at least 15 genes (Fig S5; Table S2). 9,072 out of the 9,181 DEGs (98.8% of DEGs) were
assigned to a cluster. These clusters all fell within one of three main patterns with peak
expression in either early (1-38 hpf), mid (65-163 hpf), or late (183-255 hpf) developmental

groupings (Table S2) and were combined for analyses of each of the three developmental
patterns (Fig 3B). In total, 2,679 genes were identified as peaking in expression in early
development, 3,192 peaking in mid-development, and 3,201 peaking in late development (Fig
3B).
There were 386 GO terms enriched in early development (eggs at 1 hpf through
embryos at 38 hpf) that included functions involved in embryonic development including cell
growth, cell cycle, and DNA and RNA processing cell cycle, nucleic acid metabolic process,
mRNA processing, chromosome organization, RNA processing and localization, signal
transduction, cytoskeleton organization, developmental process, cell division, and methylation
and demethylation (Fig S6). Early development gene expression was dominated by GO terms
involved in development and not by functions related to energetics or metabolism (Table S3).
In mid-development (embryos at 65 hpf through larvae at 163 hpf), there were 27
enriched GO terms (Table S4). These terms included terms related to morphological
development and swimming behavior including cilium movement and cilium-dependent cell
motility (Fig 4A). A full list of GO terms enriched and statistical results in mid-development are
provided in Table S4. There were also terms related to energy and central metabolism including
ATP metabolic processes, organonitrogen compound biosynthetic process, oxidative
phosphorylation, translation and protein targeting (Fig 4A).
Gene expression in late development (183-255 hpf including larvae, metamorphosed
polyps, and recruits) was also enriched for terms involved in morphological development
including muscle tissue and organ development, morphogenesis of a branching structure, and
cell differentiation including stem cell differentiation and neuron fate commitment (Fig 4B). A full
list of GO terms enriched and statistical results in late development are provided in Table S5.
There was also enrichment of GO terms related to intracellular chemical and ion homeostasis,
including sequestration of iron ions, intracellular cation homeostasis, and inorganic ion
homeostasis (Fig 4B). In addition, there was enrichment of terms related to ion transport,
detoxification of inorganic compounds, and response to oxygen-containing compounds (Fig 4B).
There were several enriched terms directly related to glucose and carbohydrate availability and
metabolism. These terms included glucose and carbohydrate intracellular homeostasis,
response to glucose, hexose, and carbohydrates, and cellular response to glucose, hexose, and
carbohydrate stimuli (Fig 4B). Central metabolism terms including organic acid, oxoacid,
carboxylic acid, and NADP metabolic processes were enriched, as was organic and carboxylic
acid transport and lipid localization (Fig 4B).
254
255   ***D. Holobiont metabolomic responses across ontogeny***
256
There were significant shifts in holobiont metabolomic responses across ontogeny
characterized as changes in metabolite pool size from 183 identified compounds
(PERMANOVA; SS=5361.80, R^2=0.64, F(12,46)=5.13, P=0.001). Unsupervised PCA
visualizations showed shifts in metabolomic responses across time points (Fig 5A) with 34.7%
of variance on PC1 describing shifts across developmental time and 12.09% of variance on PC2
primarily describing separation of attached recruits from other life stages (Fig 5A). Due to the
significant shifts in metabolites seen in unsupervised analyses, we proceeded with supervised
PLS-DA analysis (Fig 5B) and variables of importance on projection (VIP) identification to

determine metabolites that significantly contribute to variation across time points. PLS-DA
analyses showed that 40% of variance was explained on X-variate 1, describing shifts across
developmental time, with X-variate 2 explaining 12% of variance driven primarily by separation
of eggs and attached recruits from other life stages (Fig 5B).

We identified 82 VIP metabolites with VIP values ≥ 1 that significantly contribute to
separation between time points (45% of all metabolites; Table S6; Fig S7; Fig 5C). Four
Montiporic Acids displaying the highest VIP values (ranging from 1.82 to 1.84), all peaking in
attached recruits (Table S6; Fig 5D). Pool sizes of VIP metabolites changed across ontogeny,
with groups of metabolites peaking during distinct developmental periods (Fig 5D; Table S6).
Specifically, there were 23 metabolites peaking in early development in egg, embryo, and larval
stages from 1-163 hpf. These metabolites include purine and pyrimidine nucleotide sugars and
amino acids (e.g., FAD). There were 10 metabolites also peaking in early development in
embryos and larvae from 5-163 hpf, including ATP and ADP, carbohydrates (e.g., glucose-6-
phosphate), and amino acids (e.g., glutathione, glutamate). In mid-development, there were 6
metabolites with elevated pool size, including organic phosphoric acids (e.g., carbamoyl
phosphate), fatty acid esters (e.g., carnitines), and indole. As development progressed into
larval, polyp, and recruit stages between 93-255 hpf, metabolites including amino acids (e.g.,
betaine, glutamine), dicarboxylic acids, phenylpropanoic acids, and organosulfonic acids
increased in concentration. In attached recruit stages (183-255 hpf), there was a peak in several
carbohydrates (e.g., cellobiose, sucrose, fructose), pyruvate), and amino acids. Interestingly, a
set of metabolites peaked in both eggs (1 hpf) and attached recruits (183-255 hpf) including
glucose, dipeptides (arginine-alanine, arginine-glutamine), and ribonucleosides.

To examine metabolic pathways that shift across development, we conducted
enrichment analyses of metabolite classes (Table S7) and KEGG pathways (Table S8) of all
VIP metabolites. At the metabolite class level, there was significant enrichment of pyrimidine
ribonucleosides ($p<0.001$) and nucleotides ($p=0.007$) as well as purine ribonucleotides
($p<0.001$), and purine ($p=0.005$) and pyrimidine nucleotide sugars ($p=0.007$) in metabolites that
discriminate between developmental stages (Fig 5E). Tryptamines ($p=0.019$), dicarboxylic acids
($p=0.034$), pterins ($p=0.034$) and their derivatives were also enriched in VIP metabolites along
with carbohydrates and conjugates ($p<0.001$) and amino acids, peptides, and analogous
($p<0.001$) (Fig 5E). KEGG pathways involved in carbohydrate metabolism including starch and
sucrose ($p=0.001$), galactose ($p=0.009$), fructose and mannose ($p=0.036$), and riboflavin
metabolism ($p=0.049$) were enriched in VIP metabolites along with glyoxylate and dicarboxylate
metabolism (Fig 5F). Nitrogen ($p=0.019$) and amino acid metabolism (e.g., phenylalanine
($p=0.022$), valine, leucine, and isoleucine ($p=0.022$), and phenylalanine, tyrosine, and
tryptophan metabolism ($p=0.049$)) were also enriched in metabolites that changed across
development (Fig 5F).

Finally, we examined changes in pool size of four VIP metabolites of interest related to
symbiotic nutritional exchange including glucose (a primary photosynthate) [21,35–37], glutamine
and glutamate (metabolites involved in ammonium assimilation) [9,31,38,39], and arginine-glutamine
(a dipeptide involved in nitrogen sequestration) [31,40]. Glucose concentrations varied significantly
across development (ANOVA, SS=0.46, $F(12,34)=4.16$, $P<0.001$), peaking in attached recruits
at 231 hpf with lowest levels see in metamorphosed polyps at 231 hpf (post hoc $P<0.05$; Fig
6A). Glutamine exhibited significant increases across development (ANOVA, SS=3.40,

F(12,34)=35.01, P<0.001). Glutamine concentrations were lowest in eggs and reached a peak
in attached recruits at 183-255 hpf (post hoc P<0.05; Fig 6B). Glutamate exhibited the opposite
pattern across development, decreasing significantly (ANOVA, SS=0.54, F(12,34)=7.96,
P<0.001) from eggs and embryos to attached recruits (Fig 6C). Arginine-glutamine
concentrations also varied significantly (ANOVA, SS=2.17, F(12,34)=17.50, P<0.001) with a
peak in eggs, larvae (231 hpf), and attached recruits (183-255 hpf) (Fig 6D).
315
316   **Discussion**
317
The results of this study provide an extensive organismal and multi-omic view of
metabolism and symbiotic interactions across multiple early life history stages of a vertically
transmitting coral, advancing understanding of metabolic shifts during development in a
foundational nutritional symbiosis. Critically, the results of our study detail mechanisms by which
corals undergo extensive metabolic changes during development and that both symbiont-
derived and stored energy reserves are important to support energetic demands (Fig 7). We
found that although nutritional exchange may begin in larval stages in this species [31], a strong
metabolic response to carbohydrate photosynthates was not seen until after metamorphosis
(Fig 7). In addition, we found that lipid metabolism was important in early development with
clear evidence of increased importance of symbiont-derived nutrition after metamorphosis (Fig
7). Collectively, our integrative approach clarifies and deepens understanding of ontogenetic
metabolic shifts and highlights the large molecular and metabolic reorganization that occurs
during coral early life history. Because the metabolism of symbiont-derived nutrition increases
during sensitive post-settlement stages, symbiotic disruption under climate change driven
environmental stress has the potential to reduce energetic capacity and, consequently,
successful coral reef recruitment and persistence.
334
335   *A.* *Early developmental energy demands are supported by parentally provisioned*
336   *reserves*
337
Stored parental reserves are an important source of nutrition in early stages of coral
development. Coral offspring are lecithotrophic (i.e., feed on maternally provisioned egg yolk)
and under normal environmental conditions are provisioned with stored energy reserves
sufficient to complete development until symbiotic nutrition and/or heterotrophic feeding provide
necessary energy for growth and survival [23,26,28,41,42]. In early development, embryos undergo a
reorganization of tissues and cellular structures, gastrulation and differentiation of germ layers,
and rapid cell proliferation [43–45]. These processes require maternal and paternal transcripts,
structural molecules, and metabolic building blocks (e.g., glycogen, amino acids, lipids) to
support cell functions that are provisioned to differing extents to the eggs from the parents [46–48].
The total amount and rate at which these lipids are metabolized can determine larval pelagic
duration and energy available to meet the demands of development, metamorphosis,
settlement, and survival [49–51].
We found evidence of metabolism of parentally provisioned lipids in early embryos (1-38
hpf). Specifically, there were increased concentrations of intermediates and cofactors involved
in lipid and stored energy metabolism in early development (e.g., acetyl CoA, lysine,

pantothenate). Glyoxylate metabolism was one of the enriched metabolic pathways that changed across development. The glyoxylate pathway generates acetyl CoA from lipid stores when carbohydrate availability is limited and may facilitate stored reserve metabolism during stress and bleaching [52–55]. Furthermore, there were increased concentrations of cofactors involved in fatty acid and central metabolism including pantothenate, riboflavin, and lysine and precursors of glycerol storage (e.g., UDP-Glucose). Our results support that metabolism of stored lipids contribute to the energetic needs of early embryonic development.

Lipid metabolism also played a role in supporting development throughout larval stages. During larval stages, there were increases in carnitine metabolites (i.e., O–Decanoyl–L–carnitine and L–Octanoylcarnitine), which transport fatty acids during lipid metabolism, suggesting that larvae were continuing to metabolize lipid stores, as demonstrated in previous research [24–26]. However, there was no enrichment of lipid metabolism processes found in our gene expression analysis and in combination with our observation of decreased concentrations of lipid metabolism cofactors (e.g., acetyl CoA, lysine, pantothenate, riboflavin) supports the finding that stored lipid metabolism decreases in contribution as development progresses. Parental provisioning is critical for early embryonic development in our study and underscores the importance of characterizing how environmental stressors may compromise gamete provisioning, especially during reproductive events following mass bleaching events [48,56].

B. Shifts in nitrogen metabolism across ontogeny support nutritional requirements and symbiotic relationships

Nitrogen cycling is a central function in the coral-algal symbiosis and is a driver of symbiotic stability and productivity [39,57–59]. We found that nitrogen metabolism changed significantly across development to support nutritional demands and maintain a stable symbiotic relationship. Nitrogen availability can be modulated by both the host and symbiont through assimilation of free ammonium by each partner through the glutamine synthetase-glutamine oxoglutarate aminotransferase/glutamate synthase pathways (GS-GOGAT) [8,38,60–62]. Through these pathways, ammonium is assimilated into amino acids through sequential action of glutamine synthetase (GS; produces glutamine from glutamate and ammonium) and glutamine oxoglutarate aminotransferase/glutamate synthase (GOGAT; produces two molecules of glutamate from glutamine and α-ketoglutarate), which generates amino acids that can be used for a wide variety of cellular functions and metabolism [38,40,63].

Signatures of increased ammonium metabolism in our study first appear in mid-developmental stages with significant shifts in the concentration of glutamine, glutamate, and arginine-glutamine dipeptides in larvae. Our metabolomic results revealed a consistent increase in glutamine pools with concurrent decrease in glutamate pools across development, indicating activity of the glutamine synthase pathway to assimilate ammonium in the holobiont through production of glutamine [38,64]. This was likely driven by nitrogen uptake in the algal symbiont community, because ammonium assimilation by the symbiont can account for 14-23 times greater ammonium assimilation than host assimilation in adult corals [38] and there was no enrichment of nitrogen metabolism functions in host gene expression. Importantly, we observed increased symbiont densities between 65-93 hpf in larvae as well as increased photosynthesis

between 93-183 hpf, corresponding with this period of elevated ammonium assimilation by the algal symbiont in larval stages and further suggesting symbiont proliferation is a nitrogen-limited response [65].

Other mechanisms of nitrogen mechanisms, such as production arginine-glutamine dipeptides have been hypothesized to play a role in nitrogen sequestration by the coral host in adult corals [40] and in *Montipora capitata* larvae [31]. Interestingly, there were three distinct peaks in arginine-glutamine dipeptide concentrations across development - first in eggs (1 hpf), then in larvae (231 hpf), and finally in attached recruits (183-255 hf). The high production of arginine-glutamine in our study during larval and recruit stages suggests that the coral host is actively sequestering nitrogen to reduce nitrogen availability to the symbiont, as demonstrated previously in *M. capitata* larvae [31]. Nitrogen limitation of the symbiont population reduces the rate of population growth and can favor the translocation of excess photosynthates from the symbiont to the host [2,8,39,57,58,60,66]. Recent research in the coral *Stylophora pistillata*, the sea anemone *Exaiptasia diaphana*, and jellyfish *Cassiopea andromeda* have demonstrated that passive feedback loops between carbon and nitrogen availability are crucial for a stable symbiosis and disruption in these feedback loops can lead to dysbiosis [39,58,59].

We suggest that alongside ammonium assimilation via the GS pathway, the coral host performs nitrogen sequestration via the production of arginine-glutamine dipeptides during early development to stabilize symbiont growth and translocation. We found that symbiont population densities decreased by 43% after 93 hpf and were stable from 163 hpf onward. Similarly, photosynthesis rates increased a few hours later at 183 hpf and remained stable further into development. This stabilization in cell densities and photosynthetic rates supports the interpretation of regulation of nitrogen availability to symbiont populations by the host [8,66,67]. Alongside metabolomic evidence of nitrogen metabolism, we documented increased gene expression enrichment for host response to organonitrogen compounds and monatomic ion transport, which could indicate increased transport of ions such as ammonium and/or bicarbonate [68–70]. Also increased in mid-development, were carbamoyl phosphate concentrations which are involved in the urea cycle to remove nitrogen through the production of urea [31,40,71]. In a previous study in *M. capitata* larvae, the host used ammonium assimilation and nitrogen sequestration via the urea cycle and dipeptide production as a mechanism to limit nitrogen availability to the symbiont and maintain stable symbiont populations under thermal stress [31]. We suggest that these mechanisms continue throughout metamorphosis and settlement to maintain stable symbiotic relationships and control algal symbiont population growth.

In early development, however, arginine-glutamine dipeptides may play a different role by being used to meet nutritional requirements. Arginine-glutamine was elevated in eggs along with enriched gene expression of nitrogen metabolism, suggesting that parentally provisioned dipeptides may play different roles depending on developmental stage and may provide a source of nitrogen for developing embryos. Arginine-glutamine is more soluble than either amino acid individually [72] and may therefore be a stable mechanism for providing a source of nitrogen to gametes in corals. Specifically, the catabolism of arginine-glutamine may provide a source of glutamine for catabolic glutamate dehydrogenase activity that can generate a-ketoglutarate for metabolism through the tricarboxylic acid cycle [39]. Recent work has demonstrated that heterotrophically-sourced nitrogen is indeed packaged into coral gametes [73]

and we suggest that dipeptides may be one form of provisioned nitrogen. Future work should
examine the role of dipeptides in supporting the nutritional needs of developing embryos and
how the role of this metabolite changes across development.
Throughout this developmental time series, our samples contained a mixture of
Cladocopium sp. and *Durusdinium* sp., which is characteristic of this species in Kāneʻohe Bay,
Hawaiʻi [74,75]. Stability in Symbiodiniaceae communities suggests that the nutritional and
metabolic changes documented in this study are not driven by changes in symbiont identity and
thus function. However, as environmental conditions continue to change, the thermal tolerance
advantages of hosting *D. glynnii* previously documented in adult *M. capitata* [76,77] could manifest
in a shift to a greater abundance of *D. glynnii* than *Cladocopium sp.* provisioned to offspring [78].
Interestingly, *Acropora tumida* recruits (approx. 1.5 months old) exhibit higher infection rates of
Durusdinium symbionts under elevated temperature, but with no clear fitness advantages 31
days after infection [79]. Since *D. glynnii* translocate lower amounts of photosynthates than
Cladocopium sp. in *M. capitata* [80], this could result in detrimental impacts on nutritional
availability for offspring in the future and tradeoffs that can result in reduced growth and
reproduction at later stages and should be examined in additional work.
Bacterial communities play important roles in nutritional cycling [57,81,82] and acquisition of
micronutrients [83], and can act as nutritional mediators between host and symbiont [84,85]. For
example, larvae of the brooding coral *Pocillopora acuta* transfer bacterially derived nitrogen to
their symbiont populations [84]. Shifts in bacterial taxa indicate potential changes in microbial
nutritional relationships in early life history. We found that Alteromonadaceae (*Alteromonas* sp.),
Pseudoalteromonadaceae (*Psychrosphaera* sp.) were highest in relative abundance in early
development, with Oceanospiriliaceae peaking in larvae and Rhodobacteraceae and
Flavobacteriaceae highest in late development. These taxa play important roles in carbon [86],
nitrogen [84], sulfur (e.g., DMSP/DMS [87]), and micronutrient cycling [83] along with the production of
antimicrobial compounds [88]. Further, taxa in Rhodobacteraceae are associated with symbiont
communities and may signal a shift in the microbial communities associated with symbionts
across development [89]. Therefore, we hypothesize that our observations of an increase in
bacterial diversity could confer increased nutritional functional diversity and capacity for nutrient
cycling as corals age [90,91]. Although our limited sequencing depth of microbial communities
prevented us from making detailed functional conclusions on shifts in bacterial communities, the
increases in microbial diversity across ontogeny are compelling and are deserving of future
research to understand the functional and nutritional role of bacterial associations during coral
development. To test this hypothesis, quantitative measures of nutrient cycling between
bacterial communities and the coral host should be completed in future studies with approaches
such as metatranscriptomics, DNA-stable isotope probing (e.g., DNA-SIP), as well as
metabolomic stable isotope probing and microscopy techniques (e.g., NanoSIMS) applied to
early life stages [92].
479
480 ### C. Photosynthate metabolism becomes increasingly important after metamorphosis
481
Progressing through metamorphosis and settlement is an energetically demanding
process for marine taxa [93–95] due to rearrangement of the body plan [96] and initiation of
attachment and biomineralization [69,97]. The high energetic demand of metamorphosed polyps as

compared to swimming larvae at the same age, indicates *M. capitata* larvae underwent high
energetic stress during metamorphosis. This is also evidenced by a sharp depletion in glucose
pools in metamorphosed polyps and enrichment of genes involved in oxidative phosphorylation
and ATP metabolism during this period. Thus, after metamorphosis, it is critical for newly settled
corals to obtain sufficient nutrition to recover from energetic stress and meet demands of
calcification and growth.

Metabolism of glucose, a major photosynthate [21,35,36], increased after metamorphosis.
Importantly, glucose was elevated in attached recruit stages following a depletion of glucose
during metamorphosis, suggesting that glucose was heavily used as a source of nutrition during
metamorphosis, with pools recovering to previous concentrations after settlement. Metabolomic
analyses also revealed depletion of a glucose intermediate, glucose-6-phosphate, and
increased production of pyruvate, a glycolysis product in late developmental stages.
Additionally, there were decreases in NAD+ and NADP+, important cofactors for glycolysis and
central metabolism [98]. Together, these shifts in glycolysis metabolites and cofactors indicate
increased glycolytic metabolism after metamorphosis, as more pyruvate is produced from high
turnover of intermediate metabolites. Strong enrichment of carbohydrate, glucose, and hexose
metabolism, stimulus, and homeostasis in gene expression of the host provide evidence that
central carbon metabolism is occurring in the host, that glucose was indeed translocated from
the algal symbiont, and photosynthate metabolism plays an important role in supporting
nutritional requirements after settlement.

Carbon produced through photosynthesis is the primary source of carbon packaged into
coral gametes [48]. In early development, we also found that glucose supported developmental
energy demands, but that this was likely not a result of active translocation from the symbiont.
The earliest developmental stages in *M. capitata* are unlikely to experience active translocation
from their symbionts during the first few hours of development, even though symbionts are
present. This is because spawning occurs at approximately 21:45 with fertilization complete by
approximately 23:00 and therefore, the first 8 hours of development are completed in the dark
with no potential for photosynthetic activity. Therefore, glucose pools in the eggs were likely
parentally provisioned as a source of nutrition to support early development. As climate change
intensifies and coral reproduction occurs under higher temperatures, symbiotic nutritional
exchange is increasingly vulnerable in both adult and larval corals. A current paradigm of the
field is that energetic contribution of symbiont nutrition to metabolic demands of swimming coral
larvae is negligible [30]. Previous work examining the influence of symbiotic nutritional exchange
on larval energy requirements has largely occurred in brooded larvae in vertically transmitting
species (e.g., [27,28,30]), with ample lipid reserves [26,28,99], thus the relative contribution of symbiotic
nutrition to energy needs may be less than that of smaller, spawned larvae. Here, we propose
that prior research based on vertically transmitting brooding corals has underestimated the
energetic contribution of symbiont nutrition to metabolic demands in spawning species. Our
observation that post-settlement stages utilize photosynthates to support energy demand, and
previous observations that spawned *Acropora* larvae use lipid reserves at reduced rates if they
have symbionts [29], suggest that symbiotic nutrition is impactful for pelagic larval dispersal
potential during early development for vertically transmitting spawning species. Further, the
contribution of symbiont nutrition must be considered in the context of energetic demands. For
example, [30] suggested that the contribution of symbiotic nutrition to larval energy demands is

negligible compared to adult conspecific in *Pocillopora damicornis.* However, adult corals are expending energy for expensive processes such as calcification and reproduction [100–102], while larvae require energy for dispersal and metamorphosis [26,95,103], and therefore, while symbiotic nutritional exchange may be at lower rates in larvae, this does not discount that it may be energetically meaningful for life stage specific energy demands. Comparative studies that clarify the influence of symbiont nutrition on energetic demands across life history strategies in the context of life stage specific energetic budgets will be essential for understanding trajectories of reef recruitment under climate change. Specifically, future work conducting direct tracing of translocation using stable isotope methods across multiple developmental stages would further clarify the contribution of photosynthates to energy demands during vulnerable life history transitions.

The benefits of photosynthate translocation and increase in nutritional exchange are, however, not without dangers from temperature and irradiance driven production of reactive oxygen species (ROS) as photosynthesis byproducts [104–106]. In our study, we observed that *M. capitata* larvae had enriched gene expression including detoxification and response to oxygen-containing compounds during late development. These pathways are involved with stress and ROS response to photosynthesis byproducts and are central components of metabolism in symbiotic corals [21,70,107,108]. Elevated expression of cellular stress response genes supports that early life stages require activation of cellular stress management to neutralize oxidative stress from symbiont metabolic activity [104–106]. Observed response to ROS, sequestration of trace metals, and increased carbohydrate production by the symbiont during late developmental stages suggest that nutritional exchange is active during *M. capitata* development and that the host is actively investing in the management and enhancement of symbiont photosynthesis. We found enrichment of iron ion sequestration in late developmental stages, which is a critical trace metal that affects symbiont population productivity and symbiotic stability [83,109]. In addition, there was increased production of storage carbohydrates including sucrose, fructose, raffinose, and cellobiose in late developmental stages, which can be fixed by the symbiont and translocated to the host [110]. Previous work demonstrating that pools of starches and carbohydrates decrease when the symbiosis is compromised under bleaching conditions [111] further suggests that increased carbohydrate production in attached recruits in our study is a product of photosynthetic carbon fixation. Indeed, the swimming larvae of *M. capitata* have the capacity to invest in symbiotic regulation [31] and stress response. Yet, in a future of increasing heat wave intensity and frequency [112], further increasing the energetic costs of cellular stress management to maintain symbiotic relationships will further hinder settlement and developmental success [106,113].

D. Conclusions

In this study, we documented large-scale ontogenetic shifts in metabolism and symbiotic relationships in a vertically transmitting species, *Montipora capitata*, in Hawai□i. We found that parentally provisioned reserves support early embryonic development. During larval stages, we documented increases in symbiont population and photosynthetic capacity, with evidence that this is driven by symbiont ammonium assimilation. Following this initial symbiont proliferation, the coral host employed nutrient cycling to enhance symbiont productivity and stabilize growth

via nitrogen limitation [31,58,59] to support high energetic costs of settlement and metamorphosis.
Active symbiotic nutritional exchange and metabolism of photosynthates increased strongly
after settlement, highlighting their value to the holobiont success at the most vulnerable coral life
stages. The results of this study demonstrate that the symbiotic relationship is maturing during
early life stages, and that symbiotic interactions and nutritional exchange play important roles in
shaping metabolic responses across development. Nutritional symbioses are ubiquitous in
nature, and it is critical to understand how symbiotic relationships form in foundational
organisms with complex early life histories. As environmental stress continues to increase on
reefs, it is critical to conduct additional research to understand how symbiotic stress may impact
the energetic vulnerabilities and dispersal potential of vertically transmitting species.
583

Resource availability

Lead contact

Requests for further information and resources should be directed to and will be fulfilled by the
lead contact, Ariana S. Huffmyer (ashuffmyer@gmail.com).

Materials availability

This study did not generate new unique reagents.

Data and code availability

Data and scripts are openly available on GitHub as a static release at
https://github.com/AHuffmyer/EarlyLifeHistory_Energetics/releases/tag/v2.1.1 and Open
Science Framework (OSF) at DOI 10.17605/OSF.IO/KMU7H [114]. Raw sequence files (TagSeq,
ITS2, and 16S) are stored at the NCBI Sequence Read Archive under project PRJNA900235.
Raw and processed metabolomic data are archived at OSF at DOI 10.17605/OSF.IO/KMU7H
[114].

Acknowledgements

This research was supported by the National Science Foundation Ocean Sciences Postdoctoral
Fellowship to ASH (award no. 2205966), US-Israeli Binational Science Foundation to HMP and
TM (award no. BSF 2016-321), National Science Foundation Rules of Life-Epigenetics (award
no. EF-1921465), and was supported by a gift of the Washington Research Foundation to the
University of Washington eScience Institute. As guests, we recognize and give thanks for the
land and water resources of the ʻāina and the traditional owners of the land, kānaka ʻōiwi,
both past and present, as well as future generations, on which this experimental work was
conducted in the Kāneʻohe Ahupuaʻa and the islands of Hawaiʻi. We are grateful for the
logistical support provided by the Coral Resilience Lab and the Hawaiʻi Institute of Marine
Biology. We acknowledge Margaret Schedl for RNA extraction, Craig Nelson and Shayle
Matsuda for discussions on 16S bacterial analyses, Eric Chiles and Xiaoyang Su for

metabolomics discussions, and members of the Putnam Lab (University of Rhode Island) and Roberts Lab (University of Washington) for feedback on earlier versions of this manuscript. We acknowledge use of the computational resources of the URI Center for Computational Research for this work.

Author contributions

Conceptualization: ASH, HMP; Data curation: ASH; Formal analysis: ASH; Funding acquisition: ASH, HMP, TM; Investigation: ASH, KHW, DMB, ES; Methodology: ASH, KHW, HMP; Project administration: ASH, HMP; Resources: HMP; Supervision: ASH, HMP; Validation: ASH; Visualization: ASH, HMP; Writing - original draft: ASH; Writing - review & editing: ASH, KHW, DMB, ES, TM, HMP

Author contributions

The authors declare no competing interests.

STAR Methods

   A. Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, peptides, and recombinant proteins		
DNA/RNA Shield	Zymo Research	Cat #R1100-250
Phusion High-Fidelity PCR Master Mix with HF Buffer	New England Biolabs	Cat #M0531S
Critical commercial assays		
Quick DNA/RNA Miniprep Plus Kit	Zymo Research	Cat #D7003
Invitrogen Qubit RNA High Sensitivity Assay Kit	ThermoFisher Scientific	Cat #Q32852
KAPA Biosystems Illumina Kit	Roche Sequencing	Cat #KK4824
Illumina Nextera XT Index Kit	Illumina	Cat #FC-131-1024
Deposited data		

Montipora capitata reference genome assembly (version 3)	Stephens et al. [124]	http://cyanophora.rutgers.edu/montipora
Datasets and R code for running analyses and figures	This paper; GitHub repository	https://github.com/AHuffmyer/EarlyLifeHistory_Energetics/releases/tag/v2.1.1
Raw sequence data (TagSeq, ITS2, 16S)	This paper; NCBI repository	BioProject PRJNA900235
Raw metabolomics data	This paper; Open Science Framework repository [114]	https://osf.io/kmu7h/
Silva 16S database release 138.1	Quast et al. [154]	https://www.arb-silva.de/documentation/release-1381/
mothur RDP training dataset version 9	Schloss et al. [153]	https://mothur.org/wiki/rdp_reference_files/
Oligonucleotides		
ITSintfor2 (forward) and ITS2-reverse (reverse) ITS2 primers	Camp et al. [132]	https://doi.org/10.1007/s00338-019-01889-3
515F (forward) and 806RB (reverse) 16S rRNA V4 primers	Apprill et al. [133]	https://doi.org/10.3354/ame01753
Software and algorithms		
FastQC	Andrews [121]	https://www.bioinformatics.babraham.ac.uk/projects/fastqc/
MultiQC	Ewels et al. [122]	https://github.com/MultiQC/MultiQC
FastP	Chen et al. [123]	https://github.com/OpenGene/fastp
HISAT2	Kim et al. [125]	https://daehwankimlab.github.io/hisat2/download/
StringTie2	Kovaka et al. [126]	https://github.com/skovaka/stringtie2
GFFcompare	Pertea and Pertea [127]	https://github.com/gpertea/gffcompare

MAVEN	Melamud et al. [130]	https://github.com/eugenemel/maven
Fiji	Schindelin et al. [139]	https://imagej.net/software/fiji/downloads
ImageJ	Rueden et al. [138]	https://imagej.net/ij/download.html
R Statistical Programming v4.2.2	The R Project for Statistical Computing [136]	https://cran.r-project.org
LoLinR R package	Olito et al. [134]	https://github.com/colin-olito/LoLinR
FSA R package	Ogle et al. [137]	https://cran.r-project.org/web/packages/FSA/index.html
emmeans R package	Lenth et al. [141]	https://cran.r-project.org/web/packages/emmeans/index.html
vegan R package	Okansen et al. [142]	https://cran.r-project.org/web/packages/vegan/index.html
car R package	Fox and Weisberg [135]	https://cran.r-project.org/web/packages/car/index.html
DEGreport R package	Pantano [144]	https://www.bioconductor.org/packages/release/bioc/html/DEGreport.html
goseq R package	Young et al. [145]	https://bioconductor.org/packages/release/bioc/html/goseq.html
rrvgo R package	Sayols [147]	https://www.bioconductor.org/packages/release/bioc/html/rrvgo.html
org.Ce.eg.db R package	Carlson [146]	https://doi.org/doi:10.18129/B9.bioc.org.Ce.eg.db
mixOmics R package	Rohart et al. [148]	https://bioconductor.org/packages/release/bioc/html/mixOmics.html
RVAideMemoire R package	Hervé [149]	https://cran.r-project.org/web/packages/RV

		AideMemoire/index.html
MetaboAnalyst v5.0	Pang et al. [151]	https://www.metaboanalyst.ca/
SymPortal	Hume et al. [34]	https://symportal.org/
phyloseq R package	McMurdie and Holmes [152]	https://www.bioconductor.org/packages/release/bioc/html/phyloseq.html
mothur	Schloss et al. [153]	https://mothur.org/
VSEARCH	Rognes et al. [155]	https://github.com/torognes/vsearch

638
639
640   **B. Experimental model and subject details**
641
642   **i. Gamete collection and developmental rearing**
643

Montipora capitata is a hermaphroditic broadcast spawning, vertically transmitting coral
species [115,116] that is a dominant reef-builder in Kāne☐ohe Bay, O☐ahu, Hawai☐i where this
study was conducted (Fig 1B.1). During the July 2020 spawning period, *M. capitata* egg and
sperm bundles were collected on the night (21:00) of the new moon by scooping spawned
bundles from the surface of the water in Kāne☐ohe Bay under the State of Hawai'i Division of
Aquatic Resources Special Activities Permit 2019-60. Egg sperm bundles were then transported
to the laboratory at the Hawai☐i Institute of Marine Biology (HIMB) and approximately 3-5 mL of
bundles were added to 50 mL falcon tubes filled with 0.2 µm filtered seawater (FSW) and
allowed to separate and fertilize for 1 h at 27.5°C (Fig 1B.2). Following fertilization, embryos
were washed with 1 µm FSW to remove sperm and then added to 20 L conical rearing vessels
(N = 30) following methods in [117]. Conicals were supplied with 1 µm filtered seawater (FSW) at
ambient temperature at 27.5°C and 50-100 µmol photons m^{-2} s^{-1} (photosynthetic active
radiation; PAR). Throughout development, embryos were gently stirred, and debris, lipid slicks,
and dead material were removed every 3-6 hrs. Embryos were reared in conical vessels
through development to the swimming larval stage (Fig 1B.3). At approximately 165 hours post-
fertilization, larvae were transferred in 53 µm mesh-bottom plastic bins (N = 8; ~ 15 cm x 30 cm)
containing aragonite settlement plugs (circular surface; surface area 2.85 cm^2) conditioned for 3
days in seawater to outdoor flow-through seawater tables for settlement following methods in
[117]. Settlement was conducted at ambient temperature (~28°C) and light of 50-100 µmol m^{-2} s^{-1}
µmol photons m^{-2} s^{-1} on a 12h:12h light:dark cycle achieved by covering bins with two layers of
shade cloth. Larvae were allowed to settle in these conditions for 48 h, at which point we
examined larvae for metamorphosis. Two settlement states were sampled in this study – larvae
that metamorphosed and attached to the surface of aragonite plugs (Fig 1B.4) and those that
remained in the water metamorphosed without attachment (i.e., metamorphosed polyps). We
collected samples at 13 time points across the early life history of *M. capitata* including fertilized

eggs (1 hour post-fertilization, hpf), embryos (cleavage at 5 hpf, late gastrula stage at 38 and 65
hpf), larvae (93, 163, 183, 231 hpf), metamorphosed polyps (83 and 213 hpf), and attached
recruits (183, 231, and 255 hpf) (Fig 1A) for molecular, metabolomic, physiological, and
Symbiodiniaceae and bacterial amplicon sequencing assays. Sample size for all response
variables are presented in Table S1.

C. Method details

i. Respirometry

Respiration (oxygen consumption) and photosynthesis (oxygen production) rates were
measured using an Oxy-10 Sensor Dish Reader (PreSens Precision Sensing) in combination
with a 24-well microplate with spot oxygen sensors (80 µL; Loligo Systems) at six stages:
embryos (65 hpf), larvae (93, 183, and 231 hpf) and metamorphosed polyps (183 and 231 hpf)
(Fig 1A). Six individuals were added to each 80 µL microplate well filled with 1 µm FSW
(continuously aerated for 30 min prior to measurements) and sealed with glass coverslips.
Sensor spots were allowed to condition for 15 min in 1 µm FSW prior to measurements. Two
wells of each plate were filled with 1 µm FSW as blanks. Larvae were exposed to 10 min of dark
acclimation prior to measurements. Oxygen production ("photosynthesis") was measured as
oxygen concentration ($\mu mol\ L^{-1}$) first under light conditions for 15-20 min at ~ 500 $\mu mol\ m^{-2}\ s^{-1}$
PAR using LED aquarium lighting (AquaIllumination Hydra) with oxygen measurements
collected every 15 sec. Measurements under light were conducted first to standardize light
conditions prior to measurement of respiration due to light-enhanced dark respiration (LEDR)
effects resulting from stimulation of respiration through photosynthesis [118]. The lights were then
turned off and LEDR oxygen consumption ("respiration") was measured for an additional 15-20
min in the dark. Light levels during photosynthesis measurements are within the saturating
irradiance for adults of this species [119], but PI curves were not conducted for the early life
stages. Two replicate runs were conducted for embryo and larval time points (93 and 163 hpf, *n*
= 10 samples per run, *n* = 6 runs). At time points where both larvae and metamorphosed polyps
were analyzed (183 and 231 hpf), replicate runs were conducted in which larval and
metamorphosed polyps were included in each run and randomly assigned to wells (*n* = 5
samples per time point per run, *n* = 7 runs).

ii. Size and symbiont density

At each time point (Fig 1A), pools of approx. 100 individuals were collected from
replicate conicals (*n* = 4 conicals or bins), fixed in 4% paraformaldehyde for 24-36 h then stored
in 70% ethanol at 4°C (*n* = 4 samples per time point; approx. 100 individuals per sample) for
size and symbiont density measurements. Samples were digitally photographed using an
OMAX Digital Microscope Camera (Model A35180U3; Kitchener, ON, CA) on a OMAX
dissecting light microscope (Kitchener, ON, CA) with an OMAX 0.01 mm stage micrometer
(Model A36CALM1; Kitchener, ON, CA) for calibration in each photograph. Symbiont cell
density at each time point (Fig 1A) was measured by conducting replicate cell counts (*n* = 6-8
counts per sample, *n*= 4 samples per time point) of homogenized (Pro Scientific Bio-Gen

PRO200 Homogenizer; Oxford, CT, USA) tissue using a hemocytometer (Hausser Scientific
Improved Neubauer Hemocytometer; Horsham, PA, USA).
715
716   **iii. Gene expression in the coral host**
717
From all time points, except for attached recruits that were snap-frozen in liquid nitrogen,
pools of 50-200 individuals were preserved in DNA/RNA Shield (Zymo Research, Irvine, CA,
USA, Catalog #R1100-250). RNA and DNA extractions were performed with the Quick
DNA/RNA Miniprep Plus Kit (Zymo Research, Irvine, CA, USA, Catalog #D7003) according to
the manufacturer's instructions (see detailed protocol in data repository [114]) and stored at -80°C.
RNA quantity was determined using Invitrogen Qubit™ RNA High Sensitivity Assay Kit
(ThermoFisher, Waltham, MA, USA, Catalog #Q32852) and quality was assessed using 4200
TapeStation System for ribosomal bands (Agilent Technologies, Santa Clara, CA, USA). Purity
was assessed using a NanoDrop™ 8000 Spectrophotometer (ThermoFisher, Waltham, MA,
USA). We sequenced RNA using TagSeq, which is a 3' short transcript method that is a cost-
effective alternative to RNAseq and allows for accurate estimates of transcript abundance [120].
Libraries were prepared for sequencing on one lane of Illumina NovaSeq 6000 SR100 with
standard coverage of 3-5 million 100 bp single end reads (University of Texas Austin, Genomic
Sequencing and Analysis Facility, Austin, TX, USA). Quality of raw sequences was assessed
using *FastQC* [121] and visualized with *MultiQC* [122]. Raw reads were trimmed to remove poly-A
tails and adapters and sequences with low quality (phred quality <30) and low complexity
(threshold = 50%) using *FastP* [123]. Reads were then aligned to the *Montipora capitata* reference
genome assembly Version 3 [124] accessed at (http://cyanophora.rutgers.edu/montipora/; data
repository [114]) and assembled for quantification (exon only assembly) using *HISAT2* [125] in
combination with *StringTie2* [126], and compared to the reference using *GFFcompare* [127]. Finally,
a gene count matrix was generated using *prepDE* [126].
739
740   **iv. Metabolomics**
741
Samples for metabolomic analysis were snap-frozen in liquid nitrogen and stored at -
80°C (Fig 1A). Metabolites were extracted from the coral holobiont (homogenized tissue
containing host tissue and intracellular symbionts) using 40:40:20 methanol:acetonitrile:water
with 0.5%[v/v] formic acid according to [128] and identified and quantified at the Rutgers Cancer
Institute of New Jersey Metabolomics (New Brunswick, NJ, USA) facility using hydrophilic
interaction liquid chromatography separation and liquid chromatography-mass spectrometry
(LC-MS) quantification. Metabolite quantification and identification was performed at Rutgers
Cancer Institute of New Jersey Metabolomics facility. Hydrophilic interaction liquid
chromatography (HILIC) separation was performed on a Vanquish Horizon UHPLC system
(Thermo Fisher Scientific, Waltham, MA) with XBridge BEH Amide column
(150 mm × 2.1 mm, 2.5 µm particle size, Waters, Milford, MA) using a gradient of solvent A
(95%:5% H_2O:acetonitrile with 20 mM acetic acid, 40 mM ammonium hydroxide, pH 9.4), and
solvent B (20%:80% H_2O:acetonitrile with 20 mM acetic acid, 40 mM ammonium hydroxide, pH
9.4). The gradient was 0 min, 100% B; 3 min, 100% B; 3.2 min, 90% B; 6.2 min, 90% B; 6.5
min, 80% B; 10.5 min, 80% B; 10.7 min, 70% B; 13.5 min, 70% B; 13.7 min, 45% B; 16 min,

45% B; 16.5 min, 100% B and 22 min, 100% B [129]. The flow rate was 300 μL/min. Injection
volume was 5 μL and column temperature was 25 °C. The autosampler temperature was set to
4°C and the injection volume was 5μL. Full scan mass spectrometry analysis was performed on
a Thermo Q Exactive PLUS with a HESI source which was set to a spray voltage of -2.7kV
under negative mode and 3.5kV under positive mode. The sheath, auxiliary, and sweep gas
flow rates of 40, 10, and 2 (arbitrary unit) respectively. The capillary temperature was set to
300°C and aux gas heater was 360°C. The S-lens RF level was 45. The m/z scan range was
set to 72 to 1000m/z under both positive and negative ionization mode. The AGC target was set
to 3e6, and the maximum IT was 200ms. The resolution was set to 70,000. Full scan data was
processed with a targeted data pipeline using MAVEN software package [130]. The compound
identification was assessed using accurate mass and retention time matched to the metabolite
standards from the in-house library. Prior to running the samples, the LC-MS system was
evaluated for performance readiness by running a commercially available standard mixture and
an in-house standard mixture to assess the mass accuracy, signal intensities and the retention
time consistency. All known metabolites in the mixture were detected within 5 ppm mass
accuracy. Method blank samples matching the composition of the extraction solvent are used in
every sample batch to assess background signals and ensure there was no carryover from one
run to the next. In addition, the sample queue was randomized with respect to sample treatment
to eliminate the potential for batch effects.
776
777 **v. ITS2 symbiont amplicon sequencing**
778
Internal Transcribed Spacer 2 (ITS2) rRNA gene amplicon sequencing was conducted to
identify Symbiodiniaceae communities across time points (Fig 1A). A two-step PCR protocol
was used for sequencing according to a modification of the Illumina 16S Metagenomic
Sequencing Library Preparation protocol (see protocol in OSF data repository [114]) to amplify the
ITS2 region [131] of Symbiodiniaceae nuclear ribosomal DNA. A first round of PCR was performed
to amplify the ITS2 region of Symbiodiniaceae nuclear ribosomal DNA using ITSintfor2 and
ITS2-reverse primers [132]. Duplicate PCRs were performed on each sample using 4 ng of input
DNA in 1uL, 12.5 μL of 1X Phusion HiFi Mastermix (New England Biolabs, Ipswich, MA, Catalog
#Cat #M0531S), 0.5 μL each forward and reverse 10 μM primer and 10.5 μL of Ultra-Pure water
for a total reaction volume of 25 μL. Negative controls (no template DNA) were performed with
Ultra-Pure water instead of DNA. PCR thermocycler protocol included initial denaturation at
95°C for 3 min, 35 cycles of 95°C for 30s, 52°C for 30s, and 72°C for 30s, followed by final
extension at 72°C for 2 min. Duplicate PCR products were pooled by sample and run on a 2%
agarose gel to confirm successful amplification.
25 μL of each sample was then submitted to the RI-INBRE Molecular Informatics Core at
the University of Rhode Island (Kingston, RI, USA) for the index PCR and sequencing. PCR
products from the first round of PCR were cleaned with NucleoMag beads (Takara Bio, San
Jose, CA, USA) and then visualized by agarose gel electrophoresis. A second round of PCR (50
ng of template DNA, 8 cycles) was performed to attach Nextera XT indices and adapters using
the Illumina Nextera XT® Index Kit (Illumina, San Diego, CA) and Phusion HiFi PCR master mix
(New England Biolabs, Ipswich, MA, USA, Catalog #Cat #M0531S). PCR products from the
second PCR were cleaned with NucleoMag beads and analyzed by agarose gel

electrophoresis. Selected samples were confirmed on an Agilent BioAnalyzer DNA1000 chip (Santa Clara, CA, USA). Quantification was performed on all samples prior to pooling using a Qubit fluorometer (Invitrogen, Carlsbad, CA, USA), and the final pooled library was quantified by qPCR in a Roche LightCycler480 with the KAPA Biosystems Illumina Kit (Roche Sequencing, Indianapolis, IN, USA). Samples were sequenced on an Illumina MiSeq platform (Illumina, San Diego, CA, USA) using 2x300 bp paired-end sequencing at the RI-INBRE Molecular Informatics Core (University of Rhode Island, Kingston, RI, USA).

vi. 16S rRNA amplicon sequencing

To identify bacterial communities, we conducted 16S rRNA amplicon sequencing across development (Fig 1A). Extracted DNA was prepared using a two-step PCR protocol using published primers 515F (F) and 806RB (R) for the 16S rRNA V4 variable region [133]. To identify microbial communities, 12 ng per sample of extracted DNA was used in a two-step polymerase chain reaction (PCR) protocol using 515F and 806RB primers for the 16S rRNA V4 variable region [133]): 515F with Illumina adapter overhang marked in italics (5'-*TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG* GTG CCA GCM GCC GCG GTA A-3') and 806RB with Illumina adapter overhang marked in italics (5'-*GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA G*GG ACT ACN VGG GTW TCT AAT-3'). Samples were amplified in triplicate (4 ng per replicate) with the following mix for each reaction (25 µL total): 12.5 µL 1X Phusion HiFi Mastermix (New England Biolabs, Ipswich, MA, USA, Catalog #Cat #M0531S), 0.5 µL of each primer (10 µM concentration), 10.5 µL of Ultra-Pure water, and 1 µL of DNA input. All samples were amplified with the following thermocycler conditions: initial denaturation for 2 minutes at 95°C, 35 cycles of 20 seconds at 95°C, 15 seconds at 57°C, and 5 minutes at 72°C, followed by 10 minutes at 72°C. Post-PCR, triplicates were pooled, and amplification was confirmed with a 2% agarose gel. 25 µL of PCR product per sample was then sent for index PCR and sequencing at the RI-INBRE Molecular Informatics Core at the University of Rhode Island (Kingston, RI, USA). PCR products were cleaned with NucleoMag beads (Takara Bio, San Jose, CA, USA) and visualized with agarose gel electrophoresis. 50 ng of cleaned DNA per sample was used in a 8-cycle PCR protocol to attach Nextera XT indices and adapters using the Illumina Nextera XT® Index Kit (Illumina, San Diego, CA, USA, Catalog #FC-131-1024) and Phusion HiFi PCR master mix (New England Biolabs, Ipswich, MA, USA, Catalog #Cat #M0531S). PCR products were then cleaned again with NucleoMag beads and visualized with agarose gel electrophoresis. Quality of selected samples was assessed on an Agilent BioAnalyzer DNA1000 chip (Santa Clara, CA, USA). Quantity was measured for all samples using a Qubit fluorometer (Invitrogen, Carlsbad, CA, USA) prior to pooling. The final pooled library was quantified using qPCR in a Roche LightCycler480 with the KAPA Biosystems Illumina Kit (Roche Sequencing, Indianapolis, IN, USA). Samples were then sequenced on an Illumina MiSeq (Illumina, San Diego, CA, USA) for 2x300 bp paired-end sequencing (University of Rhode Island, Kingston, RI, USA)

D. Quantification and statistical analyses

i. Respirometry

Photosynthesis and respiration rates were extracted using localized linear regressions
(LoLinR [134]) and normalized to average size of each respective life stage to calculate size-
normalized respiration and photosynthesis rates (nmol O_2 mm^{-3} min^{-1}).
Photosynthesis:respiration ratios (P:R) were calculated as gross photosynthesis divided by
respiration. Assumptions of normality and homogeneity of variance for these and following tests
were assessed by quantile-quantile plots and Levene's tests in the *car* package, respectively [135].
Due to violations in normality and homogeneity of variance, size-normalized respiration and
gross photosynthesis rates and P:R ratios were analyzed using Kruskal Wallis rank sum tests in
R Statistical Programming v4.2.2 [136]. Post hoc tests were performed using Dunn tests with
Bonferroni P-value adjustment in the *FSA* package [137].

ii. Size and symbiont density

Size was determined as individual volume by measuring the length and width of the
longitudinal and transverse axes of randomly selected individuals from each image (n = 15
individuals per time point, n = 4 samples per time point) in Fiji/ImageJ [138,139]. Eggs and
embryos (1, 5, and 65 hpf) were approximated as the volume of a sphere, V = $\frac{4}{3}\pi a^3$, where *a* is
the width. Larvae (93-231 hpf) were approximated as a prolate spheroid, V = $\frac{4}{3}\pi a^2 b$, where *a* is
$\frac{1}{2}$ width and *b* is $\frac{1}{2}$ length [140]. Metamorphosed polyps were approximated as an oblate spheroid,
V = $\frac{4}{3}\pi ab^2$, where *a* is $\frac{1}{2}$ width and *b* is $\frac{1}{2}$ length. We did not statistically compare the size of
individual attached recruits on plugs because recruits often settled in aggregation, preventing
measurement of individual size. Total surface area of attached recruit tissue on each plug was
measured for normalization of symbiont cell density. We tested for differences in size between
time points using a non-parametric Kruskal Wallis test. Post hoc tests were performed using
Dunn tests with Bonferroni P-value adjustment in the *FSA* package [137].
Symbiont cell counts were normalized to the number of individuals in the sample
measured by counting all individuals in each sample (egg through metamorphosed polyps; cells
individual^{-1}). Cell densities were also normalized to the mean size of each life stage (eggs
through metamorphosed polyps; cells mm^{-3}), or total surface area of tissue sampled (attached
recruits; cells mm^{-2}). We tested for correlations between size and cells per individual using a
Spearman correlation test. Symbiont densities per individual and size-normalized densities were
analyzed using a one-way ANOVA test with time point as the main effect, from the stages of
eggs to metamorphosed polyps. Post hoc tests were performed using estimated marginal
means in the *emmeans* package [141]. The differences in symbiont cell densities between time
points for attached recruits (symbiont cells mm^{-2}) were analyzed separately using a non-
parametric Kruskal Wallis test.

iii. Gene expression in the coral host

Gene expression across development was first visualized using PCA analyses and
tested for differences between stages using a permutational analysis of variance
(PERMANOVA) in the *vegan* package with variance stabilized transformed gene counts [142].

Raw gene counts analyzed were analyzed using a likelihood ratio test (LRT) approach to
identify differentially expressed genes in the *DESeq2* package [143] with a full model design
including time point as the main effect and the reduced model removing the effect of time point
(reduced=~1). The LRT approach detects genes that significantly change across the
developmental time series by testing whether the main effect of developmental time point
(removed in the reduced model) explains a significant amount of variance. Differentially
expressed genes (DEGs) were identified as genes with false discovery rate (FDR) adjusted P-
value < 0.05.
Patterns of expression of DEGs across development were then visualized using the
DEGreport package [144], which uses hierarchical clustering to identify groups, or clusters, of
DEGs that share expression patterns (minimum cluster size = 15). Clusters were assigned to a
developmental pattern based on the developmental window during which expression peaked:
early development (1-38 hpf), mid development (65-163 hpf) and late development (183-231
hpf). Functional enrichment of genes in each developmental pattern (DEGs peaking in early,
mid, or late development) was conducted by analyzing the list of DEGs against all genes
expressed and passed filtering using gene ontology (GO) enrichment analysis with length bias
adjustment in *goseq* [145] and published functional annotation accompanying the *M. capitata*
reference genome Version 3 (http://cyanophora.rutgers.edu/montipora/) [124]; data repository [114]).
Significantly enriched GO terms (Biological Process ontology) were selected as those with false
discovery rate adjusted P-value < 0.05. Redundant GO terms were reduced by grouping with
semantic similarity (threshold = 0.7, *Caenorhabditis elegans*; [146]) in the *rrvgo* package [147].
909
910   **iv. Metabolomics**
911
Ion counts for metabolites detected at both positive and negative polarities in each
sample were center normalized to the respective sample median ion count to account for
variability in sample concentrations (n = 183 total detected and annotated metabolites). For
metabolites that were detected at both positive and negative polarities, we selected the
normalized ion counts from the polarity with the strongest detection (i.e., greatest normalized ion
count) for each metabolite. Following normalization and polarity selection, normalized ion counts
were then log_{10} transformed. We examined holobiont metabolomic profiles across development
by first examining unsupervised PCA visualizations and PERMANOVA tests in the vegan
package [142]. Metabolomic profiles were then analyzed using supervised partial least squares
discriminant analysis (PLS-DA) followed by identification of metabolite variables of importance
in projection (VIP) in the mixOmics [148] and RVAideMemoire [149] packages. The VIP score
(calculated as the sum of squared correlations between the PLS-DA components and the
individual variable) indicates a metabolite's importance in discriminating between time points in
the PLS-DA model. Metabolites with a VIP score greater than or equal to 1 (VIP ≥ 1) are
statistically important in discriminating variation due to developmental time point [150]. The list of
significant VIP metabolites was then analyzed for compound class and KEGG pathway
enrichment in the MetaboAnalyst 5.0 web interface [151]. The effect of time point on pool size of
specific VIP metabolites of interest (including glucose, glutamine, glutamate, and arginine-
glutamine) was tested using one-way ANOVA tests with post hoc comparisons conducted using
estimated marginal means tests in the emmeans package in R [141].

v. ITS2 symbiont amplicon sequencing

Sequences were quality controlled and subsampled to even sequencing depth of 10,551
(the minimum sequencing depth in our samples) and analyzed with the SymPortal workflow [34]
locally to identify Symbiodiniaceae Defining Intragenomic Variants (DIVs) that determine ITS2
type profiles. Relative abundance was calculated and visualized for DIVs at >1% relative
abundance. Alpha diversity of DIVs was calculated using Shannon and Inverse Simpson indices
and analyzed using one-way ANOVA with time point as the main effect. Shifts in
Symbiodiniaceae communities (based on Bray-Curtis dissimilarity) were visualized across
ontogeny using a non-metric multidimensional scaling analysis (NMDS) in the *phyloseq*
package [152]. Changes in the relative abundance of major ITS2 profiles and the ratio of major
profiles were analyzed using one-way ANOVA with time point as the main effect. Post hoc tests
were conducted in the *emmeans* package [141].

vi. 16S rRNA amplicon sequencing

We conducted data analysis using the *mothur* analytical pipeline (v1.46.1) [153]. Contigs
were assembled using trimoverlap=TRUE and primers were removed. Contigs were then
filtered to remove sequences with ambiguous calls and sequences <200 bp and >350 bp length.
Unique sequences were identified and aligned to the V4 region of the Silva database (release
138.1 [154]) using the *mothur*-formatted version of the RDP training set (v.9;
https://mothur.org/wiki/rdp_reference_files/). Sequences that did not align within the region of
interest and those with >8 polymers (maxhomop=8) were removed. Sequences were then pre-
clustered with nucleotide differences of 1 (diffs=1). Chimeras were removed using VSEARCH
[155] in *mothur* (https://github.com/torognes/vsearch). Remaining sequences were then classified
to the Silva database (release 138.1) using the *mothur*-formatted version of the RDP training set
(v.9) removing sequences classified as chloroplast, mitochondria, Archaea, Eukaryote, or
unknown domain. 13% of sequences passed these QC and filtering parameters due to a high
proportion of sequences that did not align to the Silva database, which may have been due to
unintentional amplification of host sequences. Remaining sequences were clustered to
operational taxonomic units (OTU; cutoff=0.03) and subsampled to 900 sequences to obtain
even sampling depth. Due to low bacterial sequence reads, 13 out of 39 samples were removed
after subsampling. The removed samples included egg (1 hpf), embryo (5 hpf), and half of the
samples at 38 and 65 hpf stages, indicating low bacterial sequences generated from the earliest
life stages in the time series, which corresponded to samples with low DNA concentration during
extraction. These samples were not included in further analysis. Bacterial OTU counts were
analyzed in R using the *phyloseq* package [152]. Shifts in bacterial communities (calculated using
Bray-Curtis dissimilarity) across ontogeny were visualized using a non-metric multidimensional
scaling analysis (NMDS) and analyzed using PERMANOVA tests. Taxa abundance was
calculated as percent relative abundance and filtered to those present at >0.5% for visualization
at the genus level. Alpha diversity was calculated using Shannon and Inverse Simpson indices.
Changes in alpha diversity were analyzed using a one-way ANOVA with time point as the main
effect.

976
977 **Supplemental information titles and legends**
978
979 **Fig S1. Symbiont cell density per individual from egg to metamorphosed polyp stages.**
980 Data are represented as mean ± standard error. Life stage is indicated on the x-axis as hours
981 post fertilization (hpf). Color corresponds to major life history grouping (eggs=brown,
982 embryos=yellow, larvae=cyan, metamorphosed polyps=green, attached recruits=pink).
983
Fig S2. Symbiodiniaceae ITS2 analysis. (A) NMDS plot of Symbiodiniaceae communities
across development (eggs=brown, embryos=yellow, larvae=cyan, metamorphosed
polyps=green, attached recruits=pink). Shape indicates life stage. (B) Heatmap of ITS2 DIV
relative abundance of taxa abundant at >1% relative abundance across life stages. Red =
relative abundance of 1; Blue = relative abundance of 0. (C) Ratio of *Cladocopium* sp. to
Durusdinium glynii relative abundance across life stages. Error bars indicate standard error of
mean. Shared letters indicate groups are not significantly different. Groups with red letters are
significantly different at post hoc p<0.05.
992
Fig S3. Bacterial community 16S analysis. (A) NMDS plot of bacterial communities across
development (eggs=brown, embryos=yellow, larvae=cyan, metamorphosed polyps=green,
attached recruits=pink). Shape indicates life stage. (B) Alpha diversity metrics of bacterial
communities calculated as Shannon (top) and Inverse Simpson (bottom) metrics. (C) Heatmap
of bacterial relative abundance across life stages at the phylum level. (D) Heatmap of bacterial
relative abundance across life stages at the family:genus level in taxa at greater than 0.5%
relative abundance. Red colors indicate higher relative abundance.
1000
1001 **Fig S4. Differentially expressed genes across development.** Heatmap of expression (z-
1002 score) of differentially expressed genes (9,181 total DEGs; y-axis) across time points (x-axis)
1003 identified by likelihood ratio tests (FDR P-value < 0.05). Within the heatmap, orange indicates
1004 increased expression with purple indicating reduced expression relative to the average of that
1005 gene's expression in all samples. Column annotation colors indicate life stages.
1006
Fig S5. Expression patterns of gene clusters across development. Panels show detected
clusters (minimum number of genes = 15) of genes that share expression patterns across life
stages. Expression displayed as z-score of variance stabilized transformed gene counts. Dotted
lines indicate 38 hpf and 163 hpf that separate early, mid, and late developmental stages.
Clusters were assigned to developmental patterns based on the time at which expression
peaked: early development (1-38 hpf), mid development (65-163 hpf) and late development
(183-231 hpf). Red lines indicate expression for individual genes with black lines indicating the
cluster mean expression across development.
1015
Fig S6. Functional enrichment of genes in early development. Lollipop plot of functional
enrichment of genes peaking in early development (1-38 hpf). Plot shows significant (FDR
P<0.05) gene ontology (GO) terms (Biological Process; BP). x-axis and dot size indicate % hits
(percentage of the number of DEGs in the respective GO category compared to total number of

genes included in that category) with color indicating P-value. Parent term BP is indicated in
panels on the right and individual GO terms on the left.
1022
1023 **Fig S7. Metabolite variables of importance in project (VIP) scores for all metabolites.** Red
1024 dashed line indicates VIP = 1, above which metabolites are statistically important in
1025 discriminating between life stage groups. VIP scores calculated for components 1 and 2 in the
1026 partial least squares discriminant model.
1027
1028 **Table S1. Sample size of response variables across developmental stages.** Time point
1029 identified by life stage group and hours post-fertilization (hpf). Shaded boxes indicate that the
1030 response was not measured at the respective time point.
1031
1032 **Table S2. Gene expression patterns across development.** Number of genes identified in
1033 each gene expression cluster and the assignment of each cluster to developmental patterns.
1034 Developmental patterns describe clusters of genes that peaked in expression in early
1035 development (1-38 hpf), mid development (65-163 hpf), and late development (183-231 hpf)
1036 periods.
1037
Table S3. Functional enrichment of genes in early development. Significantly enriched
(adjusted P-value < 0.05) Biological Process (BP) gene ontology (GO) terms in *Montipora*
capitata early developmental stages (1-38 hpf). Functional enrichment analyses performed in
the goseq package in R with redundant terms reduced by grouping with semantic similarity in
the rrvgo package in R to generate parent terms. Percent Hits calculated as the number of
differentially expressed genes in the GO category divided by the total number of genes from the
background set in the GO category.
1045
Table S4. Functional enrichment of genes in mid-development. Significantly enriched
(adjusted P-value < 0.05) Biological Process (BP) gene ontology (GO) terms in *Montipora*
capitata mid-developmental stages (65-163 hpf). Functional enrichment analyses performed in
the goseq package in R with redundant terms reduced by grouping with semantic similarity in
the rrvgo package in R to generate parent terms. Percent Hits calculated as the number of
differentially expressed genes in the GO category divided by the total number of genes from the
background set in the GO category.
1053
Table S5. Functional enrichment of genes in late development. Significantly enriched
(adjusted P-value < 0.05) Biological Process (BP) gene ontology (GO) terms in *Montipora*
capitata late developmental stages (183-231 hpf). Functional enrichment analyses performed in
the goseq package in R with redundant terms reduced by grouping with semantic similarity in
the rrvgo package in R to generate parent terms. Percent Hits calculated as the number of
differentially expressed genes in the GO category divided by the total number of genes from the
background set in the GO category.
1061
1062 **Table S6. Metabolites that significantly vary across development.** Variable importance in
1063 projection (VIP) values for metabolites that significantly (VIP ≥ 1) contribute to discrimination

1064 between life stages as determined by partial least squares discriminant analyses. Pathway or
1065 function indicates metabolic function of each metabolite. Ontogenetic peak indicates
1066 developmental periods in which metabolite pool size was highest.
1067
1068 **Table S7. Class enrichment of metabolites that vary across development.** Enrichment of
1069 metabolite compound classes (sub-class) of VIP metabolites conducted using Metaboanalyst
1070 v5.0 web interface. Bold indicates classes significantly enriched at FDR p-value < 0.05.
1071
1072 **Table S8. KEGG enrichment of metabolites that vary across development.** Enrichment of
1073 KEGG pathways of VIP metabolites conducted using Metaboanalyst v5.0 web interface. Bold
1074 indicates pathways significantly enriched at FDR p-value < 0.05.
1075
1076
1077
References

1.   Muscatine, L., and Porter, J.W. (1977). Reef Corals: Mutualistic Symbioses Adapted to
Nutrient-Poor Environments. Bioscience *27*, 454–460. https://doi.org/10.2307/1297526.

2.   Yellowlees, D., Rees, T.A.V., and Leggat, W. (2008). Metabolic interactions between algal
symbionts and invertebrate hosts. Plant Cell Environ. *31*, 679–694.
https://doi.org/10.1111/j.1365-3040.2008.01802.x.

3.   van der Heijden, M.G.A., Dombrowski, N., and Schlaeppi, K. (2017). Continuum of root-
fungal symbioses for plant nutrition. Proc. Natl. Acad. Sci. U. S. A. *114*, 11574–11576.
https://doi.org/10.1073/pnas.1716329114.

4.   Dubilier, N., Bergin, C., and Lott, C. (2008). Symbiotic diversity in marine animals: the art of
harnessing chemosynthesis. Nat. Rev. Microbiol. *6*, 725–740.
https://doi.org/10.1038/nrmicro1992

5.   LaJeunesse, T.C., Parkinson, J.E., Gabrielson, P.W., Jeong, H.J., Reimer, J.D., Voolstra,
C.R., and Santos, S.R. (2018). Systematic revision of Symbiodiniaceae highlights the
antiquity and diversity of coral endosymbionts. Curr. Biol. *28*, 2570–2580.
https://doi.org/10.1016/j.cub.2018.07.008.

6.   Muscatine, L., Mccloskey, L.R., and Marian, R.E. (1981). Estimating the daily contribution
of carbon from zooxanthellae to coral animal respiration. Limnol. Oceanogr. *26*, 601–611.
https://doi.org/10.4319/lo.1981.26.4.0601.

7.   Falkowski, P.G., Dubinsky, Z., Muscatine, L., and Porter, J.W. (2006). Light and the
bioenergetics of a symbiotic coral. Bioscience *34*, 705–709.
https://doi.org/10.2307/1309663.

8.   Falkowski, P.G., Dubinsky, Z., Muscatine, L., and McCloskey, L. (1993). Population control
in symbiotic corals - Ammonium ions and organic materials maintain the density of
zooxanthellae. Bioscience *43*, 606–611. https://doi.org/10.2307/1312147.

9.   Roberty, S., Béraud, E., Grover, R., and Ferrier-Pagès, C. (2020). Coral Productivity Is Co-
Limited by Bicarbonate and Ammonium Availability. Microorganisms *8,* 640.

https://doi.org/10.3390/microorganisms8050640.

10. Bourne, D.G., Morrow, K.M., and Webster, N.S. (2016). Insights into the Coral Microbiome:
Underpinning the Health and Resilience of Reef Ecosystems. Annu. Rev. Microbiol. *70*,
317–340. https://doi.org/10.1146/annurev-micro-102215-095440.

11. Connolly, S.R., Bellwood, D.R., and Hughes, T.P. (2003). Indo-pacific biodiversity of coral
reefs: Deviations from a mid-domain model. Ecology *84*, 2178–2190.
https://doi.org/10.1890/02-0254.

12. Woodhead, A.J., Hicks, C.C., Norström, A.V., Williams, G.J., and Graham, N.A.J. (2019).
Coral reef ecosystem services in the Anthropocene. Funct. Ecol. *33*, 1023–1034.
https://doi.org/10.1111/1365-2435.13331.

13. Weis, V.M. (2008). Cellular mechanisms of Cnidarian bleaching: stress causes the collapse
of symbiosis. J. Exp. Biol. *211*, 3059–3066. https://doi.org/10.1242/jeb.009597.

14. Sully, S., Burkepile, D.E., Donovan, M.K., Hodgson, G., and van Woesik, R. (2019). A
global analysis of coral bleaching over the past two decades. Nat. Commun. *10*, 1264.
https://doi.org/10.1038/s41467-019-09238-2.

15. Eddy, T.D., Lam, V.W.Y., Reygondeau, G., Cisneros-Montemayor, A.M., Greer, K.,
Palomares, M.L.D., Bruno, J.F., Ota, Y., and Cheung, W.W.L. (2021). Global decline in
capacity of coral reefs to provide ecosystem services. One Earth *4*, 1278–1285.
https://doi.org/10.1016/j.oneear.2021.08.016.

16. Ritson-Williams, R., Arnold, S.N., Fogarty, N.D., Steneck, R.S., Vermeij, M.J.A., and Paul,
V.J. (2009). New perspectives on ecological mechanisms affecting coral recruitment on
reefs. Smithson. Contrib. Mar. Sci. *38*, 437–457.
https://doi.org/10.5479/si.01960768.38.437.

17. Pineda, J. (2000). Linking larval settlement to larval transport: assumptions, potentials, and
pitfalls. Oceanography of the eastern Pacific *1*, 84–105.

18. Marshall, D.J., Pettersen, A.K., Bode, M., and White, C.R. (2020). Developmental cost
theory predicts thermal environment and vulnerability to global warming. Nat. Ecol. Evol. *4*,
406–411. https://doi.org/10.1038/s41559-020-1114-9.

19. Drake, J.L., Mass, T., Stolarski, J., Von Euw, S., van de Schootbrugge, B., and Falkowski,
P.G. (2020). How corals made rocks through the ages. Glob. Chang. Biol. *26*, 31–53.
https://doi.org/10.1111/gcb.14912.

20. Baird, A.H., Guest, J.R., and Willis, B.L. (2009). Systematic and Biogeographical Patterns
in the Reproductive Biology of Scleractinian Corals. Annu. Rev. Ecol. Evol. Syst. *40*, 551–
571. https://doi.org/10.1146/annurev.ecolsys.110308.120220.

21. Rosset, S.L., Oakley, C.A., Ferrier-Pagès, C., Suggett, D.J., Weis, V.M., and Davy, S.K.
(2020). The Molecular Language of the Cnidarian–Dinoflagellate Symbiosis. Trends
Microbiol. *29*, 320-333. https://doi.org/10.1016/j.tim.2020.08.005.

22. Edmunds, P.J. (2023). Coral recruitment: patterns and processes determining the dynamics
of coral populations. Biol. Rev. Camb. Philos. Soc. *98*, 1862-1886.

1144   https://doi.org/10.1111/brv.12987.

1145 23. Harii, S., Nadaoka, K., Yamamoto, M., and Iwao, K. (2007). Temporal changes in
1146   settlement, lipid content and lipid composition of larvae of the spawning hermatypic coral
1147   Acropora tenuis. Mar. Ecol. Prog. Ser. *346*, 89–96. https://doi.org/10.3354/meps07114.

1148 24. Rivest, E.B., Chen, C.-S., Fan, T.-Y., Li, H.-H., and Hofmann, G.E. (2017). Lipid
1149   consumption in coral larvae differs among sites: a consideration of environmental history in
1150   a global ocean change scenario. Proc. R. Soc. Lond. Ser. B. Biol. Sci. *284*, 20162825.
1151   https://doi.org/10.1098/rspb.2016.2825.

1152 25. Graham, E.M., Baird, A.H., Connolly, S.R., Sewell, M.A., and Willis, B.L. (2016).
1153   Uncoupling temperature-dependent mortality from lipid depletion for scleractinian coral
1154   larvae. Coral Reefs *36*, 97-104. https://doi.org/10.1007/s00338-016-1501-5.

1155 26. Figueiredo, J., Baird, A.H., Cohen, M.F., Flot, J.-F., Kamiki, T., Meziane, T., Tsuchiya, M.,
1156   and Yamasaki, H. (2012). Ontogenetic change in the lipid and fatty acid composition of
1157   scleractinian coral larvae. Coral Reefs *31*, 613–619. https://doi.org/10.1007/s00338-012-
1158   0874-3.

1159 27. Gaither, M.R., and Rowan, R. (2010). Zooxanthellar symbiosis in planula larvae of the coral
1160   *Pocillopora damicornis*. J. Exp. Mar. Bio. Ecol. *386*, 45–53.
1161   https://doi.org/10.1016/j.jembe.2010.02.003.

1162 28. Harii, S., Yamamoto, M., and Hoegh-Guldberg, O. (2010). The relative contribution of
1163   dinoflagellate photosynthesis and stored lipids to the survivorship of symbiotic larvae of the
1164   reef-building corals. Mar. Biol. *157*, 1215–1224. https://doi.org/10.1007/s00227-010-1401-0.

1165 29. Hazraty-Kari, S., Masaya, M., Kawachi, M., and Harii, S. (2022). The early acquisition of
1166   symbiotic algae benefits larval survival and juvenile growth in the coral *Acropora tenuis*. J.
1167   Exp. Zool. A. Ecol. Integr. Physiol. *337*, 559–565. https://doi.org/10.1002/jez.2589.

1168 30. Kopp, C., Domart-Coulon, I., Barthelemy, D., and Meibom, A. (2016). Nutritional input from
1169   dinoflagellate symbionts in reef-building corals is minimal during planula larval life stage.
1170   Sci Adv *2*, e1500681. https://doi.org/10.1126/sciadv.1500681.

1171 31. Huffmyer, A., Ashey, J., Strand, E., Chiles, E., Su, X., and Putnam, H. (2024). Coral larvae
1172   increase nitrogen assimilation to stabilize algal symbiosis and combat bleaching under
1173   increased temperature. PLoS Biol. *22*, e3002875.
1174   https://doi.org/10.1371/journal.pbio.3002875.

1175 32. LaJeunesse, T.C., and Thornhill, D.J. (2011). Improved resolution of reef-coral
1176   endosymbiont (Symbiodinium) species diversity, ecology, and evolution through psbA non-
1177   coding region genotyping. PLoS One *6*, e29013.
1178   https://doi.org/10.1371/journal.pone.0029013.

1179 33. Wham, D.C., Ning, G., and LaJeunesse, T.C. (2017). *Symbiodinium glynnii* sp. nov., a
1180   species of stress-tolerant symbiotic dinoflagellates from pocilloporid and montiporid corals
1181   in the Pacific Ocean. Phycologia *56*, 396–409. https://doi.org/10.2216/16-86.1.

1182 34. Hume, B.C.C., Smith, E.G., Ziegler, M., Warrington, H.J.M., Burt, J.A., LaJeunesse, T.C.,
1183   Wiedenmann, J., and Voolstra, C.R. (2019). SymPortal: A novel analytical framework and

platform for coral algal symbiont next-generation sequencing ITS2 profiling. Mol. Ecol.
Resour. *19*, 1063–1080. https://doi.org/10.1111/1755-0998.13004.

35. Burriesci, M.S., Raab, T.K., and Pringle, J.R. (2012). Evidence that glucose is the major
transferred metabolite in dinoflagellate-cnidarian symbiosis. J. Exp. Biol. *215*, 3467–3477.
https://doi.org/10.1242/jeb.070946.

36. Gordon, B.R., and Leggat, W. (2010). Symbiodinium—Invertebrate Symbioses and the
Role of Metabolomics. Mar. Drugs *8*, 2546–2568. https://doi.org/10.3390/md810546.

37. Hillyer, K.E., Dias, D., Lutz, A., Roessner, U., and Davy, S.K. (2018). 13C metabolomics
reveals widespread change in carbon fate during coral bleaching. Metabolomics *14*, 12.
https://doi.org/10.1007/s11306-017-1306-8

38. Pernice, M., Meibom, A., Van Den Heuvel, A., Kopp, C., Domart-Coulon, I., Hoegh-
Guldberg, O., and Dove, S. (2012). A single-cell view of ammonium assimilation in coral–
dinoflagellate symbiosis. ISME J. *6*, 1314–1324.
https://doi.org/10.1038/ismej.2011.196https://doi.org/10.1038/ismej.2011.196.

39. Rädecker, N., Pogoreutz, C., Gegner, H.M., Cárdenas, A., Roth, F., Bougoure, J.,
Guagliardo, P., Wild, C., Pernice, M., Raina, J.-B., et al. (2021). Heat stress destabilizes
symbiotic nutrient cycling in corals. Proc. Natl. Acad. Sci. U. S. A. *118*, e2022653118.
https://doi.org/10.1073/pnas.2022653118.

40. Chiles, E.N., Huffmyer, A.S., Drury, C., Putnam, H.M., Bhattacharya, D., and Su, X. (2022).
Stable isotope tracing reveals compartmentalized nitrogen assimilation in scleractinian
corals. Front. Mar. Sci., *9*, 1035523. https://doi.org/10.3389/fmars.2022.1035523

41. Padilla-Gamiño, J.L., Bidigare, R.R., Barshis, D.J., Alamaru, A., Hédouin, L., Hernández-
Pech, X., Kandel, F., Leon Soon, S., Roth, M.S., Rodrigues, L.J., et al. (2013). Are all eggs
created equal? A case study from the Hawaiian reef-building coral *Montipora capitata*.
Coral Reefs *32*, 137–152. https://doi.org/10.1007/s00338-012-0957-1.

42. Buttari, F., Tsai, S., Wen, Z.-H., Cheng, J.-O., and Lin, C. (2024). Lipid profiling during
embryogenesis of coral *Galaxea fascicularis*. Mar. Biol. *171*, 199.
https://doi.org/10.1007/s00227-024-04521-3.

43. Okubo, N., Mezaki, T., Nozawa, Y., Nakano, Y., Lien, Y.-T., Fukami, H., Hayward, D.C.,
and Ball, E.E. (2013). Comparative embryology of eleven species of stony corals
(Scleractinia). PLoS One *8*, e84115. https://doi.org/10.1371/journal.pone.0084115.

44. Reyes-Bermudez, A., Villar-Briones, A., Ramirez-Portilla, C., Hidaka, M., and Mikheyev,
A.S. (2016). Developmental Progression in the Coral *Acropora digitifera* Is Controlled by
Differential Expression of Distinct Regulatory Gene Networks. Genome Biol. Evol. *8*, 851–
870. https://doi.org/10.1093/gbe/evw042.

45. Heather, Q.M., and Mark, Q.M. (2007). Embryonic development in two species of
scleractinian coral embryos: *Symbiodinium* localization and mode of gastrulation. Evol.
Dev. *9*, 355–367. https://doi.org/10.1111/j.1525-142X.2007.00173.x.

46. Fadlallah, Y.H. (1983). Coral Reefs Sexual Reproduction , Development and Larval Biology
in Scleractinian Corals A Review. Coral Reefs *2*, 129–150.

https://doi.org/10.1007/BF00336720.

47. Ghosh, S., Körte, A., Serafini, G., Yadav, V., and Rodenfels, J. (2022). Developmental energetics: Energy expenditure, budgets and metabolism during animal embryogenesis. Semin. Cell Dev. Biol. *138*, 83-93. https://doi.org/10.1016/j.semcdb.2022.03.009.

48. Rodrigues, L.J., and Padilla-Gamiño, J.L. (2022). Trophic provisioning and parental trade-offs lead to successful reproductive performance in corals after a bleaching event. Sci. Rep. *12*, 18702. https://doi.org/10.1038/s41598-022-21998-4.

49. Graham, E.M., Baird, A.H., Connolly, S.R., Sewell, M.A., and Willis, B.L. (2013). Rapid declines in metabolism explain extended coral larval longevity. Coral Reefs *32*, 539–549. https://doi.org/10.1007/s00338-012-0999-4.

50. Christiansen, F.B., and Fenchel, T.M. (1979). Evolution of marine invertebrate reproductive patterns. Theor. Popul. Biol. *16*, 267–282. https://doi.org/10.1016/0040-5809(79)90017-0.

51. Hadfield, M., and Strathmann, M. (1996). Variability, flexibility and plasticity in life histories of marine invertebrates. Oceanol. Acta *19*, 323–334.

52. Axworthy, J.B., Timmins-Schiffman, E., Brown, T., Rodrigues, L.J., Nunn, B.L., and Padilla-Gamiño, J.L. (2022). Shotgun Proteomics Identifies Active Metabolic Pathways in Bleached Coral Tissue and Intraskeletal Compartments. Front. Mar. Sci. *9*, 797517.

53. DeSalvo, M.K., Sunagawa, S., Voolstra, C.R., and Medina, M. (2010). Transcriptomic responses to heat stress and bleaching in the elkhorn coral *Acropora palmata.* Mar. Ecol. Prog. Ser. *402*, 97–113.

54. Kenkel, C.D., Meyer, E., and Matz, M.V. (2013). Gene expression under chronic heat stress in populations of the mustard hill coral (*Porites astreoides*) from different thermal environments. Mol. Ecol. *22*, 4322–4334. https://doi.org/10.1111/mec.12390.

55. Petrou, K., Nunn, B.L., Padula, M.P., Miller, D.J., and Nielsen, D.A. (2021). Broad scale proteomic analysis of heat-destabilised symbiosis in the hard coral *Acropora millepora*. Sci. Rep. *11*, 19061. https://doi.org/10.1038/s41598-021-98548-x.

56. Leinbach, S.E., Speare, K.E., Rossin, A.M., Holstein, D.M., and Strader, M.E. (2021). Energetic and reproductive costs of coral recovery in divergent bleaching responses. Sci. Rep. *11*, 23546. https://doi.org/10.1038/s41598-021-02807-w.

57. Radecker, N., Pogoreutz, C., Voolstra, C.R., Wiedenmann, J., and Wild, C. (2015). Nitrogen cycling in corals: The key to understanding holobiont functioning? Trends Microbiol. *23*, 490–497. https://doi.org/10.1016/j.tim.2015.03.008.

58. Rädecker, N., Escrig, S., Spangenberg, J.E., Voolstra, C.R., and Meibom, A. (2023). Coupled carbon and nitrogen cycling regulates the cnidarian–algal symbiosis. Nat. Commun. *14*, 6948. https://doi.org/10.1038/s41467-023-42579-7.

59. Cui, G., Mi, J., Moret, A., Menzies, J., Zhong, H., Li, A., Hung, S.-H., Al-Babili, S., and Aranda, M. (2023). A carbon-nitrogen negative feedback loop underlies the repeated evolution of cnidarian–Symbiodiniaceae symbioses. Nat. Commun. *14*, 6949. https://doi.org/10.1038/s41467-023-42582-y.

60. Cui, G., Liew, Y.J., Li, Y., Kharbatia, N., Zahran, N.I., Emwas, A.-H., Eguiluz, V.M., and
Aranda, M. (2019). Host-dependent nitrogen recycling as a mechanism of symbiont control
in Aiptasia. PLoS Genet. *15*, e1008189. https://doi.org/10.1371/journal.pgen.1008189.

61. Rahav, O., Dubinsky, Z., Achituv, Y., Falkowski, P.G., and Smith, D.C. (1989). Ammonium
metabolism in the zooxanthellate coral, *Stylophora Pistillata*. Proc. R. Soc. Lond. Ser. B,
Biol. Sci. *236*, 325–337. https://doi.org/10.1098/rspb.1989.0026.

62. Morris, L.A., Voolstra, C.R., Quigley, K.M., Bourne, D.G., and Bay, L.K. (2019). Nutrient
Availability and Metabolism Affect the Stability of Coral–Symbiodiniaceae Symbioses.
Trends Microbiol. *27*, 678–689. https://doi.org/10.1016/j.tim.2019.03.004.

63. Roberts, J.M., Fixter, L.M., and Davies, P.S. (2001). Ammonium metabolism in the
symbiotic sea anemone *Anemonia viridis*. Hydrobiologia *461*, 25–35.
https://doi.org/10.1023/a:1012752828587.

64. Su, Y., Zhou, Z., and Yu, X. (2018). Possible roles of glutamine synthetase in responding to
environmental changes in a scleractinian coral. Mol. Biol. Rep. *45*, 2115–2124.
https://doi.org/10.1007/s11033-018-4369-3.

65. Cui, G., Liew, Y.J., Konciute, M.K., Zhan, Y., Hung, S.-H., Thistle, J., Gastoldi, L., Schmidt-
Roach, S., Dekker, J., and Aranda, M. (2022). Nutritional control regulates symbiont
proliferation and life history in coral-dinoflagellate symbiosis. BMC Biol. *20*, 103.
https://doi.org/10.1186/s12915-022-01306-2.

66. Krueger, T., Horwitz, N., Bodin, J., Giovani, M.-E., Escrig, S., Fine, M., and Meibom, A.
(2020). Intracellular competition for nitrogen controls dinoflagellate population density in
corals. Proc. Biol. Sci. *287*, 20200049. https://doi.org/10.1098/rspb.2020.0049.

67. Xiang, T., Lehnert, E., Jinkerson, R.E., Clowez, S., Kim, R.G., DeNofrio, J.C., Pringle, J.R.,
and Grossman, A.R. (2020). Symbiont population control by host-symbiont metabolic
interaction in Symbiodiniaceae-cnidarian associations. Nat. Commun. *11*, 108.
https://doi.org/10.1038/s41467-019-13963-z.

68. Zoccola, D., Ganot, P., Bertucci, A., Caminiti-Segonds, N., Techer, N., Voolstra, C.R.,
Aranda, M., Tambutté, E., Allemand, D., Casey, J.R., et al. (2015). Bicarbonate
transporters in corals point towards a key step in the evolution of cnidarian calcification. Sci.
Rep. *5*, 9983. https://doi.org/10.1038/srep09983.

69. Tambutté, S., Holcomb, M., Ferrier-Pagès, C., Reynaud, S., Tambutté, É., Zoccola, D., and
Allemand, D. (2011). Coral biomineralization: From the gene to the environment. J. Exp.
Mar. Bio. Ecol. *408*, 58–78. https://doi.org/10.1016/j.jembe.2011.07.026.

70. Levy, S., Elek, A., Grau-Bové, X., Menéndez-Bravo, S., Iglesias, M., Tanay, A., Mass, T.,
and Sebé-Pedrós, A. (2021). A stony coral cell atlas illuminates the molecular and cellular
basis of coral symbiosis, calcification, and immunity. Cell *184*, 2973–2987.e18.
https://doi.org/10.1016/j.cell.2021.04.005.

71. Matthews, J.L., Oakley, C.A., Lutz, A., Hillyer, K.E., Roessner, U., Grossman, A.R., Weis,
V.M., and Davy, S.K. (2018). Partner switching and metabolic flux in a model cnidarian-
dinoflagellate symbiosis. Proc. Biol. Sci. *285,* 20182336.
https://doi.org/10.1098/rspb.2018.2336.

72. Fürst, P. (2001). New developments in glutamine delivery. J. Nutr. *131*, 2562S – 8S.
https://doi.org/10.1093/jn/131.9.2562S.

73. Jaffe, M.D., Padilla-Gamiño, J.L., Nunn, B.L., and Rodrigues, L.J. (2023). Coral trophic
pathways impact the allocation of carbon and nitrogen for egg development after bleaching.
Front. Ecol. Evol. *11,* 1251220. https://doi.org/10.3389/fevo.2023.1251220.

74. Cunning, R., Ritson-Williams, R., and Gates, R.D. (2016). Patterns of bleaching and
recovery of *Montipora capitata* in Kane'ohe Bay, Hawai'i, USA. Mar. Ecol. Prog. Ser. *551*,
131–139. https://doi.org/10.3354/meps11733.

75. Wall, C.B., Kaluhiokalani, M., Popp, B.N., Donahue, M.J., and Gates, R.D. (2020).
Divergent symbiont communities determine the physiology and nutrition of a reef coral
across a light-availability gradient. ISME J. *14*, 945–958. https://doi.org/10.1038/s41396-
019-0570-1.

76. Roach, T.N.F., Dilworth, J., Christian Martin, H., Daniel Jones, A., Quinn, R., and Drury, C.
(2021). Metabolomic signatures of coral bleaching history. Nat. Ecol. Evol. *5*, 495–503.
https://doi.org/10.1101/2020.05.10.087072.

77. Dilworth, J., Caruso, C., Kahkejian, V.A., Baker, A.C., and Drury, C. (2021). Host genotype
and stable differences in algal symbiont communities explain patterns of thermal stress
response of *Montipora capitata* following thermal pre-exposure and across multiple
bleaching events. Coral Reefs *40*, 151–163. https://doi.org/10.1007/s00338-020-02024-3.

78. Padilla-Gamiño, J.L., Pochon, X., Bird, C., Concepcion, G.T., and Gates, R.D. (2012). From
parent to gamete: vertical transmission of *Symbiodinium* (Dinophyceae) ITS2 sequence
assemblages in the reef building coral *Montipora capitata*. PLoS One *7*, e38440.
https://doi.org/10.1371/journal.pone.0038440.

79. Ng, T.Y., Chui, A.P.Y., Tsang, R.H.L., Yeung, C.W., and Ang, P. (2024). Hong Kong coral
recruits do not gain advantage by being flexible in establishing symbiosis under simulated
climate change conditions. J. Coast. Res. *113*, 936–940. https://doi.org/10.2112/jcr-si113-
183.1.

80. Allen-Waller, L., and Barott, K.L. (2023). Symbiotic dinoflagellates divert energy away from
mutualism during coral bleaching recovery. Symbiosis *89*, 173-186.
https://doi.org/10.1007/s13199-023-00901-3.

81. Glaze, T.D., Erler, D.V., and Siljanen, H.M.P. (2022). Microbially facilitated nitrogen cycling
in tropical corals. ISME J. *16*, 68–77. https://doi.org/10.1038/s41396-021-01038-1.

82. Peixoto, R.S., Rosado, P.M., Leite, D.C. de A., Rosado, A.S., and Bourne, D.G. (2017).
Beneficial Microorganisms for Corals (BMC): Proposed Mechanisms for Coral Health and
Resilience. Front. Microbiol. *8*, 341. https://doi.org/10.3389/fmicb.2017.00341.

83. Reich, H.G., Camp, E.F., Roger, L.M., and Putnam, H.M. (2022). The trace metal economy
of the coral holobiont: supplies, demands and exchanges. Biol. Rev. Camb. Philos. Soc. *98*,
623-642. https://doi.org/10.1111/brv.12922.

84. Ceh, J., Kilburn, M.R., Cliff, J.B., Raina, J.B., Van Keulen, M., and Bourne, D.G. (2013).
Nutrient cycling in early coral life stages: *Pocillopora damicornis* larvae provide their algal

symbiont (*Symbiodinium*) with nitrogen acquired from bacterial associates. Ecol. Evol. *3*,
2393–2400. https://doi.org/10.1002/ece3.642.

85. Matthews, J.L., Raina, J.-B., Kahlke, T., Seymour, J.R., van Oppen, M.J.H., and Suggett,
D.J. (2020). Symbiodiniaceae-bacteria interactions: rethinking metabolite exchange in reef-
building corals as multi-partner metabolic networks. Environ. Microbiol. *22*, 1675–1687.
https://doi.org/10.1111/1462-2920.14918.

86. Kimes, N.E., Van Nostrand, J.D., Weil, E., Zhou, J., and Morris, P.J. (2010). Microbial
functional structure of *Montastraea faveolata*, an important Caribbean reef-building coral,
differs between healthy and yellow-band diseased colonies. Environ. Microbiol. *12*, 541–
556. https://doi.org/10.1111/j.1462-2920.2009.02113.x.

87. Raina, J.-B., Dinsdale, E.A., Willis, B.L., and Bourne, D.G. (2010). Do the organic sulfur
compounds DMSP and DMS drive coral microbial associations? Trends Microbiol. *18*, 101–
108. https://doi.org/10.1016/j.tim.2009.12.002.

88. Raina, J.-B., Tapiolas, D., Motti, C.A., Foret, S., Seemann, T., Tebben, J., Willis, B.L., and
Bourne, D.G. (2016). Isolation of an antimicrobial compound produced by bacteria
associated with reef-building corals. PeerJ *4*, e2275. https://doi.org/10.7717/peerj.2275.

89. Matthews, J.L., Khalil, A., Siboni, N., Bougoure, J., Guagliardo, P., Kuzhiumparambil, U.,
DeMaere, M., Le Reun, N.M., Seymour, J.R., Suggett, D.J., et al. (2023). Coral
endosymbiont growth is enhanced by metabolic interactions with bacteria. Nat. Commun.
14, 6864. https://doi.org/10.1038/s41467-023-42663-y.

90. Cárdenas, A., Raina, J.-B., Pogoreutz, C., Rädecker, N., Bougoure, J., Guagliardo, P.,
Pernice, M., and Voolstra, C.R. (2022). Greater functional diversity and redundancy of coral
endolithic microbiomes align with lower coral bleaching susceptibility. ISME J. *16*, 2406–
2420. https://doi.org/10.1038/s41396-022-01283-y.

91. Sogin, E.M., Putnam, H.M., Nelson, C.E., Anderson, P., and Gates, R.D. (2017).
Correspondence of coral holobiont metabolome with symbiotic bacteria, archaea and
Symbiodinium communities. Environ. Microbiol. Rep. *9*, 310–315.
https://doi.org/10.1111/1758-2229.12541.

92. Engelberts, J.P., Robbins, S.J., Damjanovic, K., and Webster, N.S. (2021). Integrating
novel tools to elucidate the metabolic basis of microbial symbiosis in reef holobionts. Mar.
Biol. *168*, 175. https://doi.org/10.1007/s00227-021-03952-6.

93. Mos, B., and Dworjanyn, S.A. (2016). Early metamorphosis is costly and avoided by young,
but physiologically competent, marine larvae. Mar. Ecol. Prog. Ser. *559*, 117–129.
https://doi.org/10.3354/meps11914.

94. Pechenik, J.A. (2006). Larval experience and latent effects-metamorphosis is not a new
beginning. Integr. Comp. Biol. *46*, 323–333. https://doi.org/10.1093/icb/icj028.

95. Edmunds, P.J., Cumbo, V.R., and Fan, T.-Y. (2013). Metabolic costs of larval settlement
and metamorphosis in the coral *Seriatopora caliendrum* under ambient and elevated pCO2.
J. Exp. Mar. Bio. Ecol. *443*, 33–38. https://doi.org/10.1016/j.jembe.2013.02.032.

96. Grasso, L.C., Negri, A.P., Fôret, S., Saint, R., Hayward, D.C., Miller, D.J., and Ball, E.E.

(2011). The biology of coral metamorphosis: molecular responses of larvae to inducers of
settlement and metamorphosis. Dev. Biol. *353*, 411–419.
https://doi.org/10.1016/j.ydbio.2011.02.010.

97. Gilis, M., Meibom, A., Domart-Coulon, I., Grauby, O., Stolarski, J., and Baronnet, A. (2014).
Biomineralization in newly settled recruits of the scleractinian coral *Pocillopora damicornis*.
J. Morphol. *275*, 1349–1365. https://doi.org/10.1002/jmor.20307.

98. Amjad, S., Nisar, S., Bhat, A.A., Shah, A.R., Frenneaux, M.P., Fakhro, K., Haris, M.,
Reddy, R., Patay, Z., Baur, J., et al. (2021). Role of NAD+ in regulating cellular and
metabolic signaling pathways. Mol Metab *49*, 101195.
https://doi.org/10.1016/j.molmet.2021.101195.

99. Rivest, E.B., and Hofmann, G.E. (2015). Effects of temperature and pCO2 on lipid use and
biological parameters of planulae of *Pocillopora damicornis*. J. Exp. Mar. Bio. Ecol. *473*,
43–52. https://doi.org/10.1016/j.jembe.2015.07.015

100.Leuzinger, S., Willis, B.L., and Anthony, K.R.N. (2012). Energy allocation in a reef coral
under varying resource availability. Mar. Biol. *159*, 177–186.
https://doi.org/10.1007/s00227-011-1797-1.

101.Neder, M., Saar, R., Malik, A., Antler, G., and Mass, T. (2022). New Insights on the Diurnal
Mechanism of Calcification in the Stony Coral, *Stylophora pistillata*. Front. Mar. Sci. *8*,
745171. https://doi.org/10.3389/fmars.2021.745171

102.Edmunds, P.J. (2012). Effect of pCO2 on the growth, respiration, and photophysiology of
massive *Porites* spp. in Moorea, French Polynesia. Mar. Biol. *159*, 2149–2160.
https://doi.org/10.1007/s00227-012-2001-y

103.Richmond, R.H. (1987). Energetics, competency, and long-distance dispersal of planula
larvae of the coral *Pocillopora damicornis.* Mar. Biol. *93*, 527–533.
https://doi.org/10.1007/BF00392790.

104.Nesa, B., Baird, A.H., Harii, S., Yakovleva, I., and Hidaka, M. (2012). Algal symbionts
increase DNA damage in coral planulae exposed to sunlight. Zool. Stud. *51*, 6.

105.Baird, A.H., Yakovleva, I.M., Harii, S., Sinniger, F., and Hidaka, M. (2021). Environmental
constraints on the mode of symbiont transmission in corals. J. Exp. Mar. Bio. Ecol. *538*,
151499. https://doi.org/10.1016/j.jembe.2020.151499.

106.Kitchen, S.A., Jiang, D., Harii, S., Satoh, N., Weis, V.M., and Shinzato, C. (2022). Coral
larvae suppress heat stress response during the onset of symbiosis decreasing their odds
of survival. Mol. Ecol. *31*, 5813–5830. https://doi.org/10.1111/mec.16708.

107.Matthews, J.L., Crowder, C.M., Oakley, C.A., Lutz, A., Roessner, U., Meyer, E., Grossman,
A.R., Weis, V.M., and Davy, S.K. (2017). Optimal nutrient exchange and immune
responses operate in partner specificity in the cnidarian-dinoflagellate symbiosis. Proc.
Natl. Acad. Sci. U. S. A. *114*, 13194–13199. https://doi.org/10.1073/pnas.1710733114.

108.Kitchen, S.A., and Weis, V.M. (2017). The sphingosine rheostat is involved in the cnidarian
heat stress response but not necessarily in bleaching. J. Exp. Biol. *220*, 1709–1720.
https://doi.org/10.1242/jeb.153858.

109. Ferrier-Pagès, C., Sauzéat, L., and Balter, V. (2018). Coral bleaching is linked to the
capacity of the animal host to supply essential metals to the symbionts. Glob. Chang. Biol.
24, 3145–3157. https://doi.org/10.1111/gcb.14141

110. Hillyer, K.E., Dias, D.A., Lutz, A., Roessner, U., and Davy, S.K. (2017). Mapping carbon
fate during bleaching in a model cnidarian symbiosis: the application of 13C metabolomics.
New Phytol. *214*, 1551–1562. https://doi.org/10.1111/nph.14515

111. Hillyer, K.E., Dias, D.A., Lutz, A., Wilkinson, S.P., Roessner, U., and Davy, S.K. (2017).
Metabolite profiling of symbiont and host during thermal stress and bleaching in the coral
Acropora aspera. Coral Reefs *36*, 105–118. https://doi.org/10.1007/s00338-016-1508-y

112. Lee, H., Calvin, K., Dasgupta, D., Krinner, G., Mukherji, A., Thorne, P., Trisos, C., Romero,
J., Aldunce, P., Barret, K., et al. (2023). IPCC, 2023: Climate Change 2023: Synthesis
Report, Summary for Policymakers. Contribution of Working Groups I, II and III to the Sixth
Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing
Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland. (Intergovernmental Panel
on Climate Change (IPCC)), p. 42.

113. Hartmann, A.C., Marhaver, K.L., Klueter, A., Lovci, M.T., Closek, C.J., Diaz, E.,
Chamberland, V.F., Archer, F.I., Deheyn, D.D., Vermeij, M.J.A., et al. (2019). Acquisition of
obligate mutualist symbionts during the larval stage is not beneficial for a coral host. Mol.
Ecol. *28*, 141–155. https://doi.org/10.1111/mec.14967

114. Huffmyer, A.S. (2023). Cross-scale and multi-omic integration reveals transitions to
symbiotic nutritional integration across coral ontogeny. Preprint at Open Science
Framework, https://doi.org/10.17605/OSF.IO/KMU7H
https://doi.org/10.17605/OSF.IO/KMU7H.

115. Padilla-Gamiño, J.L., and Gates, R.D. (2012). Spawning dynamics in the Hawaiian reef-
building coral *Montipora capitata*. Mar. Ecol. Prog. Ser. *449*, 145–160.
https://doi.org/10.3354/meps09530.

116. Padilla-Gamiño, J.L., Hédouin, L., Waller, R.G., Smith, D., Truong, W., and Gates, R.D.
(2014). Sedimentation and the reproductive biology of the Hawaiian reef-building coral
Montipora capitata. Biol. Bull. *226*, 8–18. https://doi.org/10.1086/bblv226n1p8

117. Hancock, J.R., Barrows, A.R., Roome, T.C., Huffmyer, A.S., Matsuda, S.B., Munk, N.J.,
Rahnke, S.A., Drury, C. (2021). Coral husbandry for ocean futures: leveraging abiotic
factors to increase survivorship, growth, and resilience in juvenile *Montipora capitata*. Mar.
Ecol. *657*, 123-133. https://doi.org/10.3354/meps13534

118. Edmunds, P.J., and Davies, P.S. (1988). Post-illumination stimulation of respiration rate in
the coral Porites porites. Coral Reefs *7*, 7–9. https://doi.org/10.1007/BF00301975

119. Gibbin, E.M., Putnam, H.M., Gates, R.D., Nitschke, M.R., and Davy, S.K. (2015). Species-
specific differences in thermal tolerance may define susceptibility to intracellular acidosis in
reef corals. Mar. Biol. *162*, 717–723. https://doi.org/10.1007/s00227-015-2617-9.

120. Lohman, B.K., Weber, J.N., and Bolnick, D.I. (2016). Evaluation of TagSeq, a reliable low-
cost alternative for RNAseq. Mol. Ecol. Resour. *16*, 1315–1321.
https://doi.org/10.1111/1755-0998.12529.

121. Andrews, S. (2010). FastQC: A Quality Control Tool for High Throughput Sequence Data.
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

122. Ewels, P., Magnusson, M., Lundin, S., and Käller, M. (2016). MultiQC: summarize analysis
results for multiple tools and samples in a single report. Bioinformatics *32*, 3047–3048.
https://doi.org/10.1093/bioinformatics/btw354

123. Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ
preprocessor. Bioinformatics *34*, i884–i890. https://doi.org/10.1093/bioinformatics/bty560

124. Stephens, T.G., Lee, J., Jeong, Y., Yoon, H.S., Putnam, H.M., Majerová, E., and
Bhattacharya, D. (2022). High-quality genome assembles from key Hawaiian coral species.
Gigascience *11*, giac098. https://doi.org/10.1093/gigascience/giac098.

125. Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based
genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat. Biotechnol. *37*,
907–915. https://doi.org/10.1038/s41587-019-0201-4

126. Kovaka, S., Zimin, A.V., Pertea, G.M., Razaghi, R., Salzberg, S.L., and Pertea, M. (2019).
Transcriptome assembly from long-read RNA-seq alignments with StringTie2. Genome
Biol. *20*, 278. https://doi.org/10.1186/s13059-019-1910-1

127. Pertea, G., and Pertea, M. (2020). GFF Utilities: GffRead and GffCompare. F1000Res. *9*,
304. https://doi.org/10.12688/f1000research.23297.2

128. Williams, A., Chiles, E.N., Conetta, D., Pathmanathan, J.S., Cleves, P.A., Putnam, H.M.,
Su, X., and Bhattacharya, D. (2021). Metabolomic shifts associated with heat stress in coral
holobionts. Sci Adv *7,* eabd4210. https://doi.org/10.1126/sciadv.abd4210.

129. Su, X., Chiles, E., Maimouni, S., Wondisford, F.E., Zong, W.-X., and Song, C. (2020). In-
Source CID Ramping and Covariant Ion Analysis of Hydrophilic Interaction
Chromatography Metabolomics. Anal. Chem. *92*, 4829–4837.
https://doi.org/10.1021/acs.analchem.9b04181.

130. Melamud, E., Vastag, L., and Rabinowitz, J.D. (2010). Metabolomic analysis and
visualization engine for LC-MS data. Anal. Chem. *82*, 9818–9826.
https://doi.org/10.1021/ac1021166

131. Lajeunesse, T.C., and Trench, R.K. (2000). Biogeography of two species of *Symbiodinium*
(Freudenthal) inhabiting the intertidal sea anemone *Anthopleura elegantissima* (Brandt).
Biol. Bull. *199*, 126–134. https://doi.org/10.2307/1542872.

132. Camp, E.F., Suggett, D.J., Pogoreutz, C., Nitschke, M.R., Houlbreque, F., Hume, B.C.C.,
Gardner, S.G., Zampighi, M., Rodolfo-Metalpa, R., and Voolstra, C.R. (2020). Corals exhibit
distinct patterns of microbial reorganisation to thrive in an extreme inshore environment.
Coral Reefs *39*, 701–716. https://doi.org/10.1007/s00338-019-01889-3.

133. Apprill, A., McNally, S., Parsons, R., and Weber, L. (2015). Minor revision to V4 region SSU
rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquat.
Microb. Ecol. *75*, 129–137. https://doi.org/10.3354/ame01753.

134. Olito, C., White, C.R., Marshall, D.J., and Barneche, D.R. (2017). Estimating monotonic

rates from biological data using local linear regression. J. Exp. Biol. *220*, 759–764.
https://doi.org/10.1242/jeb.148775.

135. Fox, J., and Weisberg, S. (2018). An R Companion to Applied Regression Second. (SAGE
Publications).

136. R Core Team (2022). R: A language and environment for statistical computing (R
Foundation for Statistical Computing). https://www.r-project.org/

137. Ogle, D.H., Doll, J.C., Wheeler, A.P., and Dinno, A. (2025). FSA: Simple Fisheries Stock
Assessment Methods. R package version 0.9.6.
https://doi.org/10.32614/CRAN.package.FSA

138. Rueden, C.T., Schindelin, J., Hiner, M.C., DeZonia, B.E., Walter, A.E., Arena, E.T., and
Eliceiri, K.W. (2017). ImageJ2: ImageJ for the next generation of scientific image data.
BMC Bioinformatics *18*, 529. https://doi.org/10.1186/s12859-017-1934-z

139. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: An open-source
platform for biological-image analysis. Nat. Methods *9*, 676–682.
https://doi.org/10.1038/nmeth.2019

140. Isomura, N., and Nishihira, M. (2001). Size variation of planulae and its effect on the
lifetime of planulae in three pocilloporid corals. Coral Reefs *20*, 309–315.
https://doi.org/10.1007/s003380100180.

141. Lenth, R. (2024). emmeans: Estimated Marginal Means, aka Least-Squares Means. R
package version 1.10.2. https://doi.org/10.32614/CRAN.package.emmeans

142. Okansen, J., and et al. (2017). vegan: Community Ecology Package. R package version
2.6-6.1. https://doi.org/10.32614/CRAN.package.vegan

143. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.
https://doi.org/10.1186/s13059-014-0550-8

144. Pantano, L. (2024). DEGreport: Report of DEG analysis. R package version 1.40.0.
https://doi.org/doi:10.18129/B9.bioc.DEGreport

145. Young, M.D., Wakefield, M.J., Smyth, G.K., and Oshlack, A. (2010). Gene ontology
analysis for RNA-seq: accounting for selection bias. Genome Biol. *11*, R14.
https://doi.org/10.1186/gb-2010-11-2-r14.

146. Carlson, M. (2019). *org.Ce.eg.db: Genome wide annotation for Worm*. R package version
3.19.1. https://doi.org/doi:10.18129/B9.bioc.org.Ce.eg.db

147. Sayols, S. (2020). rrvgo: a Bioconductor package to reduce and visualize Gene Ontology
terms. microPubl. Biol. https://doi.org/10.17912/micropub.biology.000811

148. Rohart, F., Gautier, B., Singh, A., and Lê Cao, K.-A. (2017). mixOmics: An R package for
'omics feature selection and multiple data integration. PLoS Comput. Biol. *13*, e1005752.
https://doi.org/10.1371/journal.pcbi.1005752.

149. Hervé, M. (2020). RVAideMemoire: Testing and Plotting Procedures for Biostatistics. R
package version 0.9-83-7. https://doi.org/doi:10.18129/B9.bioc.rrvgo

150. Galindo-Prieto, B., Eriksson, L., and Trygg, J. (2014). Variable influence on projection (VIP)
for orthogonal projections to latent structures (OPLS). J. Chemom. *28*, 623–632.
https://doi.org/10.1002/cem.2627.

151. Pang, Z., Chong, J., Zhou, G., de Lima Morais, D.A., Chang, L., Barrette, M., Gauthier, C.,
Jacques, P.-É., Li, S., and Xia, J. (2021). MetaboAnalyst 5.0: narrowing the gap between
raw spectra and functional insights. Nucleic Acids Res. *49*, W388–W396.
https://doi.org/10.1093/nar/gkab382.

152. McMurdie, P.J., and Holmes, S. (2013). phyloseq: an R package for reproducible
interactive analysis and graphics of microbiome census data. PLoS One *8*, e61217.
https://doi.org/10.1371/journal.pone.0061217.

153. Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B.,
Lesniewski, R.A., Oakley, B.B., Parks, D.H., Robinson, C.J., et al. (2009). Introducing
mothur: open-source, platform-independent, community-supported software for describing
and comparing microbial communities. Appl. Environ. Microbiol. *75*, 7537–7541.

154. Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and
Glöckner, F.O. (2013). The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. Nucleic Acids Res. *41*, D590–D596.
https://doi.org/10.1093/nar/gks1219.

155. Rognes, T., Fluori, T., Nichols, B., Quince, C., and Mahé, F. (2016). VSEARCH: a versatile
open source tool for metagenomics. PeerJ, e2584. https://doi.org/10.7717/peerj.2584

1564

1565   **Figures**

1566

1568   **Fig 1. Characterization of *Montipora capitata* coral physiology and symbiosis across**
1569   **development.** (A) Variables measured across *Montipora capitata* development. Filled squares
1570   indicate the response variable was sampled at the respective time point; white squares indicate

the response was not measured. Color corresponds to major life history grouping (eggs=brown, embryos=yellow, larvae=cyan, metamorphosed polyps=green, attached recruits=pink). Hpf indicates hours post fertilization. (B) Photographs of 1 - *M. capitata* colony; 2 - Conical larval rearing chambers; 3 - Planula larvae, note pigmentation due to presence of symbiont cells, scale bar represents 1 mm; 4 - Attached recruits on settlement plug, scale bar represents 1 mm.

Fig 2. Metabolic rates and physiological responses change across *Montipora capitata* development. (A) Size (mm³) of each time point across development. (B) Size-normalized light enhanced dark respiration (R; oxygen consumed; µmol O_2 mm⁻³ min⁻¹). (C) Gross photosynthesis (oxygen produced; µmol O_2 mm⁻³ min⁻¹). (D) Gross photosynthesis : respiration ration (P:R). Dotted line indicates P:R of 1:1. (E) Symbiont cell density (cells per mm³) across development for egg through metamorphosed polyp stages. (F) Symbiont cell density (cells per mm²) in attached recruits. Data are represented as mean ± standard error for all responses. Life stage is indicated on the x-axis as hours post fertilization (hpf; eggs=brown, embryos=yellow, larvae=cyan, metamorphosed polyps=green, attached recruits=pink). P-value of Kruskal Wallis tests (KW) or analysis of variance tests (ANOVA) displayed in each panel. Letters indicate significance of post hoc comparisons with shared letters indicating no significant difference.

1595
Fig 3. Shifts in gene expression across coral early life history stages. (A) Principal
components analysis (PCA) of gene expression across *M. capitata* ontogeny (hours post-
fertilization and life stage indicated by shape) in eggs (brown), embryos (yellow), larvae (cyan),
metamorphosed polyps (green) and attached recruits (pink). Developmental patterns in gene
expression are labeled as early (1-38 hpf), mid (65-163 hpf) and late (183-255 hpf) stages. (B)
Gene expression (z-score) of DEGs peaking in early (1-38 hpf), mid (65-163 hpf) or late (183-
255 hpf) developmental time points (x-axis). The number of DEGs within each developmental
pattern are displayed above each plot. Gray lines indicate expression across time points for
individual genes with black lines indicating the mean of all DEGs in the respective group. Dotted
vertical lines indicate 38 hpf and 163 hpf separating early, mid, and late developmental phases.
1606

1607
Fig 4. Functional enrichment of differentially expressed genes in mid and late
development. Lollipop plots of the enrichment of significant (adjusted P<0.05) gene ontology
(GO) terms (Biological Process; BP) in (A) mid and (B) late development stages. x-axis and dot
size indicate % hits (percentage of the number of DEGs in the respective GO category
compared to total number of genes included in that category) with color indicating P-value.
Parent term is indicated in panels on the right and individual GO terms on the left.
1614

Fig 5. Metabolomic transitions across *Montipora capitata* ontogeny. (A) Unsupervised principal components analysis (PCA) of multivariate metabolite concentrations (normalized pool size) across *M. capitata* ontogeny (time point indicated by shape) in eggs (brown), embryos (yellow), larvae (cyan), metamorphosed polyps (green) and attached recruits (pink). (B) Supervised partial least squares discriminant analysis (PLSDA) of multivariate metabolite

concentrations. (C) Variable importance in projection values for metabolites that are statistically
significant in discriminating between time points (VIP ≥1). (D) Heatmap of expression (z-score)
of VIP metabolites (82 total VIPs) across time points (x-axis). Orange indicates increased
concentration of a metabolite relative to the mean of all samples, with purple indicating reduced
concentration. Annotations indicate ontogenetic patterns as developmental periods in which
metabolite concentration peaks. (E) Sub-class metabolite enrichment analysis conducted in
MetaboAnalyst showing significantly (P<0.05) enriched classes in VIP metabolites. X-axis
indicates enrichment ratio (number of VIP metabolites in the respective pathway / number of
expected metabolites in the respective pathway) with color indicating significance of adjusted P-
values. (F) KEGG pathway enrichment analysis conducted in MetaboAnalyst showing
significantly (P<0.05) enriched pathways in VIP metabolites.
1632

1633
Fig 6. Changes in key metabolite concentrations related to nutritional exchange and
nitrogen cycling across *Montipora capitata* development. Pool size (log-transformed
median-normalized) of four metabolites of interest: (A) glucose, (B) glutamine, (C) glutamate,
and (D) arginine-glutamine. Data are represented as mean ± standard error for all metabolites.
Life stage is indicated on the x-axis as hours post fertilization (hpf) (eggs=brown,
embryos=yellow, larvae=cyan, metamorphosed polyps=green, attached recruits=pink). Letters
indicate significance of post hoc comparisons with shared letters indicating no significant
difference.
1642

1643

Fig 7. Developmental shifts in metabolism across *Montipora capitata* ontogeny observed in this study. Thicker bars indicate qualitative representation of increased relative contribution of each metabolic pathway or energy source to development.