Early acting inbreeding depression can evolve as an inbreeding avoidance mechanism.

Yaniv Brandvain[1,✉], **Lia Thomson**[1,2], **and Tanja Pyhäjärvi**[3]

[1]Department of Plant & Microbial Biology. University of Minnesota - Twin Cities. St Paul, MN
[2]Department of Biology. Texas Tech University - Twin Cities. Lubbock, TX.
[3]Department of Forest Sciences. University of Helsinki. Helsinki, Finland.

Despite the potential for mechanical, developmental and/or chemical mechanisms to prevent self-fertilization, some amount of incidental self-fertilization is inevitable in some predominantly outcrossing species. In such cases, inbreeding (by self-fertilization or biparental inbreeding) may severely compromise individual fitness by exposing the recessive genetic load, which is otherwise hidden in the heterozygous state. Unquestionably, much of this load is an inevitable consequence of the mutational process and the masking of deleterious recessive alleles in predominantly outcrossing populations. However, we show that when back-up embryos efficiently compensate for inviable embryos lost early in development, natural selection can favor highly deleterious recessive variants that functionally induce self-sacrificial death of inbred embryos. Our work provides numerous testable hypotheses, and suggests that the distinction between late self-incompatibility and early acting inbreeding depression may be somewhat artificial in some taxa.

Inbreeding depression | Self Incompatibility | Altruism | Reproductive compensation | Polyembryony

Correspondence: *ybrandva@umn.edu*

Introduction

A recessive genetic load, caused by rare, recessive, and highly deleterious mutations (1, 2) as well as a deviation from phenotypic optima under multi-locus stabilizing selection (3), is inevitable in predominantly outcrossing populations. As such, numerous mechanisms have adaptively evolved to avoid inbreeding across plants, animals and fungi (4, 5). For example, many plant species have a chemically-mediated Self-Incompatibility (SI) system which rejects self-pollen. The mechanisms and timing of SI are diverse (6) (SI is often achieved by rejecting pollen from plants with a matching genotype at the so called S-locus), but the best-characterized cases involve an interaction between the diploid maternal genotype and the pollen's haploid (gametophytic SI) or pollen-parent's diploid (sporophytic SI) genotype at the S-locus well before fertilization.

In other "self-incompatible" taxa, self-pollen can effectively germinate and grow down self stigmas and in some cases can enter and even fertilize the ovule before ovular development arrests (7–9). These cases, are collectively known as "Late-Acting SI" (10). "Post-zygotic" SI – the scenario in which self-incompatibility is realized after fertilization – is the most extreme form of Late-Acting SI, and seems to be particularly common in the Bignoniaceae (11).

Claims of Late Acting SI, and post-zygotic SI in particular, have attracted much attention and criticism both because this mechanism of incompatibility is unusual (12), and because it is difficult to distinguish late-acting SI from early-acting inbreeding depression (6, 13). According to Charlesworth (13), the key distinction between these mechanisms is that SI is "*a mechanism operated by the maternal parent plant and controlled by the genotypes of that plant and its pollen donor*" and that early-acting inbreeding depression is "*a process acting in the progeny zygote, determined by its own genotype.*"

Substantial effort has gone into distinguishing between early acting inbreeding depression and late-acting SI (reviewed in 12, 14), with evidence supporting late-acting SI in some taxa (e.g. 15–18) and early inbreeding depression (e.g. 19, 20) in others. While these studies are quite interesting, we point out that with high reproductive compensation the distinction between early acting inbreeding depression and late acting SI is not necessarily evolutionarily relevant – that is, if there is both substantial inbreeding depression after seed development, and if embryos lost to early acting inbreeding depression can be compensated by viable embryos, early acting inbreeding can adaptively evolve as a mechanism to minimize reproductive investment in low fitness selfed seed. That is, an adaptive early expressed recessive load which functions to trade in inbred for outbred seeds plays the same functional and strategic role as traditional self-incompatibility, regardless of specifics of the genetic control of the phenomenon. We therefore argue that all forms of SI can be considered as a special case of this sort of altruistic act, and as such the distinction between late-acting SI and early-acting inbreeding depression is not meaningful if there is high reproductive compensation. Such altruistic early death has been previously considered in relation to the reinforcement of postzygotic isolating mechanisms (21, 22), the evolution of recessive lethality of a selfish transmission distorter that makes homozygous males sterile (23), and earlier onset of of genetic diseases that would guarantee low fitness (24).

After introducing the plausibility of this argument, we show that, depending on the mutation rate, ongoing evolution of this early-acting "altrusitic" adaptive load will lead to pat-

terns of variation that are difficult to distinguish from standard models of mutation selection balance. This second result potentially explains why no signal of this load was observed in a recent study of the evolution of polyembryony and its consequences (25). As such, while self-sacrifice may favor an elevated genetic load, Occam's razor argues for a null interpretation of early embryonic death as a simple consequence of standard mutation selection balance, rather than a panglossian (26) interpretation of embryonic death as an altruistic act . We therefore conclude with a series of specific predictions which allow for distinguishing between an adaptive recessive load and the load expected under standard mutation-selection balance.

Methods and models

Throughout, we assume that there is an opportunity for reproductive compensation ("that is a dead embryo can be replaced by a potentially viable one" (27)). For simplicity, we model reproductive compensation by providing each seed with a single backup - such that if the first 'primary' seed is inviable, it can be replaced by another, which may or may not be viable. This roughly follows the case of polyembryony in pines which we modeled previously (25), with the distinction that in this model maternal half-sibs do not necessarily inherit the same maternal allele. For tractability, we further assume that inbreeding depression is expressed after seed development (such that initially all embryos live to become full seeds), and that this inbreeding depression is both unpurgeable (i.e. the selfing rate is too low relative to the mutation rate to allow natural selection to decrease the recessive load) and lethal. Such an "unpurgeable load" evolved when correlations in homozygosity across unlinked loci induced by inbreeding exposes many recessive deleterious mutations at once (28).

Deterministic Models Without Mutation. To evaluate if an early-expressed recessive load can adaptively evolve, we first assume that all primary embryos survive, and that following dispersal seeds die if they are inbred and survive otherwise. We then introduce a highly recessive (with dominance coefficient, h) allele, A, which drastically decreases (by selection coefficient, s) the probability that an embryo survives. Primary embryos killed by exposing the effects of the A allele early in development can be replaced by a backup embryo, which cannot be replaced if it dies. Primary embryos that do not express this mutation will survive until later in development, when they cannot be replaced by a backup if they are killed by their inbreeding depression. We assume that the resource transfer from primary to backup embryos is perfect – so there is no maternal cost to losing their primary embryo (unless the backup embryo is either inviable or has a lower expected fitness than the primary embryo). We then explore the invasion of this allele under differing selection and dominance coefficients, and find the equilibrium frequency of such an allele in the absence of mutation. Throughout, we assume an infinite population size, but make no assumptions about Hardy-Weinberg

Equilibrium due to the recessivity an strong selection on the embryo-expressed allele.

Stochastic Model With Mutation.

Simulation framework: To complement our deterministic model showing that recessive mutations decreasing embryo survival could increase in frequency by selection, we developed genome-scale, individual-based stochastic simulations to examine the genomic signatures of this process. Specifically, we ran non-Wright-Fisher forward simulations in SLiM 3.0 (29) in which early-acting deleterious mutations recurrently arise (at rate, μ) at any location in the genome. We followed these simulations until populations achieved mutation-selection balance. SLiM implements an "infinite sites" mutational model but follows a finite number of genomic locations at which (potentially multiple) mutations can arise, with recombination only occurring between these finite sites. We specify a genome of one hundred thousand sites, such that e.g. a mutation rate of 0.5×10^{-6} corresponds to an expectation of one mutation per diploid genome per generation. These sites are uniformly spread over ten chromosomes, each of which is one Morgan in length with a uniform recombination map.

Evolving an altruistic load: Our SLiM model matches the essence our analytical framework for the adaptive evolution of an altruistic load. Each generation, 10,000 survivors were randomly sampled with replacement to act as mothers. Each mother produced two maternal siblings, each of which could be the product of selfing or outcrossing based on a set selfing rate, α. Next, early-acting deleterious mutations were expressed. If both siblings survived, one was randomly selected to be the unnecessary backup sibling and was discarded. If one or zero siblings survived, no discarding was necessary. Finally, all surviving embryos (at most one per sibling-pair) experienced viability selection in which survival depended on whether they were the product of selfing or not. The fitness of inbred offspring was a constant that we varied across simulations.

A control: As a control, we developed a second model where the early expressed load was not favored by natural selection. As in the first model, two maternal siblings were produced from 10,000 mothers every generation. Embryos were then affected by their early-acting deleterious mutations. Unlike the first model, replacement did not occur between siblings; instead occurring between random pairs. If both embryos in the pair survived, one was randomly selected and died, this removes the opportunity for altruistic selection to favor an early-expressed genetic load. We finally instituted viability selection on surviving embryos as described above.

Fig. 1. The critical selfing threshold required for the invasion of a self-sacrificial allele as a function of its selection and dominance coefficients. This model assumes that all selfed seed dies and unpurgeable inbreeding depression. The colored text reflects the approximation that a self-sacrificial allele increases when rare if $crit.self > 6h$. This approximation is exact as s approaches zero, but $crit.self$ must exceed $6h$ as s increases (e.g. with $s = 1$ $crit.self$ equals 0.155, 0.321, and 0.499 as opposed to 0.15, 0.30, and 0.45 which h is 0.025, 0.05, and 0.075, respectively.)

Results

Selection can favor recessive deleterious alleles.

Invasion of a self-sacrificial mutation: To evaluate invasion conditions, we assume the frequency of the self-sacrificial allele, p_A is very low, such that homozygotes are rare i.e. only generated by selfing. As such, we consider groups of two siblings from aa and aA mothers. Each offspring is the product of selfing or outcrossing with probabilities α and $1-\alpha$, respectively. Outcrossed ovules are pollinated by A or a bearing pollen, with probabilities p_A and $1-p_A$, respectively.

At invasion (i.e. $\lim_{p_A \to 0}$) aa mothers (which occur in frequency $1-2p_A$) produce aa offspring by selfing with probability α and produce aa and Aa offspring by outcrossing with probabilities $(1-p_A)(1-\alpha)$, and $(p_A)(1-\alpha)$, respectively. The probability that the primary embryo in an aa maternal family is replaced by a backup embryo is ψ_{aa}, and equals the product of probability that an embryo is fertilized by A pollen and dies in early development i.e. $h s p_A (1-\alpha)$.

At invasion, Aa mothers (which occur in frequency $2p_A$) produce aa, Aa, and AA offspring by selfing with probability $\frac{\alpha}{4}$, $\frac{\alpha}{2}$ and $\frac{\alpha}{4}$, respectively, and produce aa Aa, and AA offspring by outcrossing with probabilities $(1-p_A)\frac{(1-\alpha)}{2}, \frac{1}{2}$, and $p_A\frac{(1-\alpha)}{2}$, respectively. The probability that the primary embryo in an Aa maternal family is replaced by a backup embryo is ψ_{Aa}, and equals $\frac{s}{2}(h+\alpha/2+p_A(1-\alpha))$.

At invasion, the population mean fitness is $(1-\alpha)(\psi_{AA}) = (1-\alpha)$, and the mean fitness of rare Aa maternal families, $\overline{w}_{\text{Aa:fam}}$ is

$$\overline{w}_{\text{Aa:fam},\lim p \to 0} = (1-hs)(1-\alpha)(1+\frac{s}{4}(h+\frac{\alpha}{2}))$$

Fig. 2. The equilibrium frequency, p^*, of a self-sacrificial (partially) recessive allele for which lost embryos can be compensated, under a variety of selfing rates and dominance coefficients.

Assuming full recessivity (i.e. $h = 0$), $\overline{w}_{\text{Aa:fam}}$ at invasion becomes $\frac{1}{8}(1-\alpha)(8+s\alpha)$, which is always greater than mean population fitness at invasion. Because the fitness of Aa families exceeds that of aa families, it is likely that a fully recessive allele that diminishes embryo fitness and allows this embryo to be replaced will be favored by natural selection. More specifically, the expected increase in frequency of a rare recessive allele that acts in this manner is $p_A(1-\alpha)(s\alpha)/(4\overline{w})$. We note that the underlying logic to our argument holds even if selfed embryos do not all die – assuming that all outbred seed survive, that there is no biparental inbreeding, that the load cannot be purged, and a recessive allele decreases embryo fitness invades when the probability that a selfed offspring survives is less than $\frac{s(1-\alpha)}{4-s\alpha}$.

Furthermore, complete recessivity is not required for the deterministic spread of this self-sacrificial allele. A rare partially recessive allele which harms early embryos that can be replaced by siblings is expected to increase in frequency when the primary selfing rate exceeds a critical selfing level, which equals

$$\text{crit.self} = \frac{2h(3+hs)}{1-hs}$$

Figure 1 shows that at first approximation, crit.self is linear with its recessivity (h), such that a rare self-sacrificial A allele will spread when $\alpha > 6h$. That is, a greater probability of generating a low fitness selfed seed (i.e. a higher primary selfing rate) is required to favor the evolution of a self-sacrificial allele that can decrease fitness as a heterozygote.

Equilibrium frequency of a self-sacrificial mutation: As the self-sacrificial allele increases in frequency, it begins to be exposed commonly in the allozygous, rather than the autozygous, state. As such, as A increases in frequency, it becomes a less reliable indicator of recent inbreeding. To demonstrate this phenomenon, we apply Bayes' theorem to show that the probability that an embryo was the product of selfing, given that it is homozygous for A equals $\frac{p_\alpha/2}{\alpha/2+p_A(1-\alpha)}$ under the assumption of complete recessivity and lethality of the

Fig. 3. The mean frequency of early acting, recessive lethal ($\delta = 1$, A), and sublethal ($\delta = 0.75$, B) alleles, over a combination of selfing rates (σ), mutation rates (μ), selection (s) and dominance (h) coefficients. Blue lines display results for cases in which primary embryos are replaced by siblings (our model), while blue lines display cases in which primary embryos are replaced by a non-sibling (a control).

self-sacrificial allele. This results in a pattern of frequency dependent selection in which the self-sacrificial allele can be favored when rare but selected against when common, resulting in a stable intermediate frequency, p_A^*. Because our model contains both family structure and departures from Hardy-Weinberg Equilibrium, analytically deriving p_A^* is quite challenging.

Rather than deriving p_A^*, we iterate the genotype frequency recursion equations until genotype frequencies cease to change (i.e. after 1000 generations and the sum of the change in absolute difference in genotype frequencies across generations is less than 10^{-14}). The equilibrium frequency of a self sacrificial is highest for strongly recessive alleles in populations that self frequently and is lower for more less recessive alleles in more outcrossing populations (Figure 2).

A genome-wide altruistic load. We cannot easily intuit how this force will scale up to the genome scale. Is this deterministic selective benefit of self-sacrifice strong enough to overcome the sampling process of random genetic drift? If so, when will the benefit of replacing oneself with a viable half sibling leave an observable pattern on the genetic load, such that it can be separated from the standard process of mutation selection balance? What should the frequency spectrum of such self-sacrificial alleles look like – should it be marked by an excess of relatively rare or relatively common variants? Answers to such questions are essential for our ability to test the adaptive load hypothesis in natural populations.

Figure 3 compares the mean frequency deleterious reces-sive alleles per individual for early-expressive recessive ($h = 0$) alleles across three selfing rates ($\sigma = 0.1$, $\sigma = 0.2$, and $\sigma = 0.3$, x-axis) two mutation rates ($\mu = 5x10^{-8}$, or

$\mu = 5x10^{-6}$, rows) to such alleles, two levels of inbreeding depression, three selection coefficients ($s = 0.2$, $s = 0.6$, and $s = 1.0$, columns), and two levels of inbreeding depression ($\delta = 1$ Fig. 3A or $\delta = 0.75$, Fig. 3B). Although we present results only for mean allele frequency below, we find qualitatively similar results for different summaries of the mutational load (e.g. the mean number of mutations per individual, Fig S1). Consistent with our single-locus results that mean mutation frequency only weekly depends on the selection coefficient (Fig. 3).

With lethal inbreeding depression and low mutation rates, there is a clear elevation in the frequency of early-expressed deleterious recessive mutations with reproductive compen-sation as compared to the control (Top panel of Fig. 3A), consistent with expectations from our analytical theory. This pattern intensifies as the selfing rate increases, presumably because a higher selfing rate increases the necessity of removing selfed embryos. These patterns disappear as the mutation rate increases (Bottom panel of Figure 3A), presumably because the mutational input supplied under these higher mutation rates outpaces what is required to ensure the death of selfed embryos and rather acts to increase the allozygous expression of these alleles. This explanation is consistent with the increased number of such mutation per individual with this higher mutation rate (Fig S1) and their lower frequency when segregating.

With sublethal inbreeding depression ($\delta = 0.75$, 3B), we find a similar elevation in the frequency of these early expressed deleterious recessives with reproductive assurance as compared to controls. However, in this case, the frequency decreases, rather than increases with the selfing rate. This difference in the relationship between selfing rate and early expressed recessive load with lethal versus sublethal inbreeding could reflect the fact that with a sublethal load

there is some cost two having inviable selfed embryos.

Discussion

We show that early expressed recessive alleles can adaptively invade a partially selfing population with high inbreeding depression and reproductive compensation. Such a scenario is likely found in polyembryonic gymnosperms which lack a self-incompatibility mechanism, cannot avoid geitonogamous selfing, and display high inbreeding depression (30). In fact, a recent study of the evolution of polyembryony (25) found a subtle shift towards an excess of common alleles that decrease embryonic fitness when there is reproductive compensation and embryos do not compete.

Sakai (31, 32), recently found that simulations with reproductive compensation plus both early and late acting deleterious recessive mutations allowed for both mutational types to segregate at higher frequencies than they would on their own. We interpret this "two-stage effect" as an increase in the early expressed load favored by the mechanisms proposed above, followed by the weakening of selection against late expressed recessive load accompanying the masking effect of the deficit of homozygosity in surviving seeds.

Precedent for the adaptive evolution of self-sacrificial alleles. Ours is not the first study to examine the possibility of the evolution of self sacrificial alleles. For example, cytoplasmic male killers can adaptively spread in maternally transmitted organelles in taxa with reproductive compensation and/or sibling competition (33), and more broadly males are thought to be "cursed" by mitochondrial alleles that can harm male fitness at a benefit to female fitness (34, 35). Similarly, maternal effect dominant effect alleles (MEDEA) that epigenetically program the arrest of embryos that do not inherit this allele are also favored by natural selection (36). These scenarios are somewhat similar to the adaptive load modelled above in that these alleles induce self harm based on contextual information of their situation to increase their inclusive fitness. However, contrary to the case of the adaptive load modelled here, both cytoplasmic male killers and maternal effect dominant embryonic arrest alleles increase their inclusive fitness at the expense of unlinked loci – setting up a genetic conflict between these selfish alleles and the rest of the genome. By contrast, in our model of the adaptive load, the self-sacrificial allele increases the inclusive fitness of alleles at unlinked loci (at least for the maternally-derived genome). As such, rather than representing a genomic conflict, ours is a model of cooperation, and is consistent with Hamilton's (24) idea that "infant mortality may evolve when the early death of one infant makes more likely the creation or survival of a close relative" (see also (37).) The closest published analogue to our model of the adaptive load is Charlesworth's (23) model for the adaptive evolution of recessive lethal alleles linked to a driving t-haplotype, which would otherwise lead to sterility when homozygous.

The efficacy of an altruistic load in preventing inbreeding. The efficacy of an adaptive recessive load in preventing self-fertilization depends on the genetic architecture of this load. For example, our SLiM model with a selfing rate of ten percent, lethal inbreeding, and complete recessivity and lethality of self-sacrificial alleles, found about five such alleles per individual. Assuming these loci are unlinked, we expect an index of self-fertilization of $\frac{1}{2}^5 = 0.96875$ for this parameter combination – a value well-below the traditional cutoff of 0.80 for labelling a species as self-incompatible. However, the location, size, and selection coefficient of the mutational target for loci potentially underlying such early acting inbreeding depression are unknown, so we have no firm predictions for the expected index of self-incompatibility under a model of an altruistic recessive load.

Implications for the evolution of Self Incompatibility. Although we model the altruistic evolution of an early-expressed genetic load (rather than the spiteful load represented by cytoplasmic male killing and MEDEA), this evolutionary model is not without conflict. Here, the altruistic load clearly provides a greater patrilineal than matrilineal fitness cost, because an incidentally sacrificed embryo can be replaced by another from the same mother, while this pollen grain is replaced by one from a randomly chosen father. Although not directly modelled here, this conflict between maternally and paternally derived genes could favor the differential expression of the early embryonic load depending on parent of origin (especially once the deleterious mutation is at high frequency), much like conflict over maternal resource provisioning favors parent-of-origin expression under the kinship theory of genomic imprinting (38). This view of the early adaptive load as a conflict between parental genomes over a self-sacrificial act provides a potential connection between self-incompatibility and the early expressed recessive load. In fact, this view is quite similar to Uyenoyama's models that show that the differing costs and benefits of self-incompatibility alleles to pollen and style can favor pollen with novel SI specificities (39, 40).

Testable predictions.

> Adaptation is a special and onerous concept that should not be used unnecessarily.
> — George C. Williams. 1966.
> *Adaptation and Natural Selection* (41).

Our model predicts that inadvertent selfing in outcrossing species with reproductive compensation can favor the adaptive evolution of early expressed recessive deleterious variation. However, our genome-scale simulations point to a more parsimonious explanation for an early expressed load. That is, when the mutation rate is sufficiently high, the early expressed load that evolves by standard mutation-selection balance can equal or exceed the load favored by the adaptive

process presented here, obviating the need for an adaptive load. Given that a recessive load can be efficiently generated by the process of mutation rather than selection we urge for careful tests of the ideas developed here before accepting the adaptive load hypothesis. We present some testable predictions below.

Taxa with reproductive compensation should have a greater recessive embryonic load than taxa without reproductive compensation. The benefit of early embryonic death is that this death can be compensated by redirected resources that would go to this embryo to a current or future sibling. As such, we predict the load of early expressed recessive deleterious variants to increase in taxa with more reproductive compensation. However, while this observation is consistent with our adaptive load model, it is also consistent with the more parsimonious explanation that reproductive compensation relaxes selection against an early-expressed load (27, 42), counteracting the purging process in inbred populations (43).

Early acting inbreeding depression should increase with the primary selfing rate (up to a point). The greater the risk of inadvertent selfing, the greater the benefit of adaptive early-acting inbreeding depression. As such, we predict a larger early expressed recessive load in species with a higher selfing risk than those that effectively prevent the risk of self pollination. In such comparisons, the post-embryonic load – which should be unaffected by the process modelled above – could serve as a useful control. We note however, that we assumed that inbreeding depression was extreme and could not evolve. Such an unpurgeable load can arise when selfing is rare and recessive mutations are common (28), and has been observed in self-incompatible populations (e.g. 44), however both reproductive compensation and higher selfing rates make the recessive load easier to purge (27). Because we do not model the co-evolution of mating system and the early genetic load, our predictions are only valid for taxa with an near lethal inbreeding depression such that the recessive load cannot be purged by selfing.

Recessive loci underlying late-acting inbreeding depression should show signatures of balancing selection.. When the recessive load evolves adaptively (rather than under mutation selection balance) we see that recessive alleles are maintained under balancing selection and are on average older than those maintained under mutation selection balance. Thus, if the early expressed load is an adaptation, we predict a signature of balancing selection around loci underlying it – i.e. under a model of mutation-selection balance recessive deleterious alleles should be relatively young, while under a model of an altruistic load such alleles should be relatively old.

Examination of the embryo lethality in conifers. At first glance, polyembryony in gymnosperms seems to provide an ideal testing ground for these ideas, as conifers cannot avoid inbreeding, have early embryo but have relatively high levels of inbreeding depression. However the detailed study of early embryo death in conifers reveals the complex biological reality underpinning the abstractions represented as simple model parameters. Williams (45) has argued that embryo death in pines occurs by two processes (1) Numerous "Embryo Viability Loci" which are expressed throughout development and which potentially allow for embryo replacement, and (2) The Embryo Lethal System – a coordinated process of embryo death which occurs after a dominant embryo has been established with limited opportunity for replacement of a dead embryo. One potential interpretation of these two forms of embryo lethality is that an altruistic recessive load can evolve via the expression of "embryo viability loci", and that the embryo lethal system is an evolved mechanism to suppress the non-dominant embryo if both embryos are viable. However, Williams (45) further argues that most experimental studies comparing embryo filling and realized selfing rates in controlled crosses with self and non-self pollen fail to find evidence for the idea that polyembryony allows for the effective removal of selfed seeds (we note that this heterodox interpretation differs from the meta-analytical result that gymnosperms have a lower realized selfing rate than expected given their primary selfing rates (46)). Thus, evidence for an early expressed adaptive load in a potential model system for the ideas presented here are not decisive.

Summary. We show that when mothers can compensate for inviable embryos, a self-sacrificial early expressed recessive load can adaptively evolve by natural selection. Our model shows that the line between "early acting inbreeding depression" and "late-acting self-incompatibility" might not be as conceptually clear-cut as previous authors have insisted. As such, our model potentially refutes the orthodox idea that "Abortion of inbred embryos is not an evolved trait for ensuring outcrossing, but is considered an early expression of genetic load, a lethal 'price' a species pays for maintaining genetic variability" (19) – and the price paid is the that of incomplete reproductive compensation. However, our model also shows that such a mechanism only meaningfully operates when such mutations are rare as selection does not favor such mutations when they arise so frequently that a substantial recessive load would evolve in the absence of selection – as such, the key evolutionary question is not to distinguish between early-acting inbreeding depression and late acting self-incompatibility, but rather to distinguish between adaptive and non-adaptive mechanisms that prevent the production of selfed seed.

ACKNOWLEDGEMENTS
We would like to acknowledge members of the Brandvain lab for comments and the Academy of Finland for supporting our project 'Genetic determinants of embryo destiny in a polyembryonic seed – mortality, inbreeding depression and sibling rivalry in a pine nutshell'.

Bibliography

1. A S Kondrashov. Deleterious mutations as an evolutionary factor. ii. facultative apomixis and selfing. *Genetics*, 111(3):635–653, Nov 1985. ISSN 0016-6731 (Print); 0016-6731 (Linking). doi: 10.1093/genetics/111.3.635.
2. Deborah Charlesworth and John H Willis. The genetics of inbreeding depression. *Nat Rev Genet*, 10(11):783–796, Nov 2009. ISSN 1471-0064 (Electronic); 1471-0056 (Linking). doi: 10.1038/nrg2664.

3. R Lande. The influence of the mating system on the maintenance of genetic variability in polygenic characters. *Genetics*, 86(2 Pt. 1):485–498, Jun 1977. ISSN 0016-6731 (Print); 0016-6731 (Linking).

4. Darwin CR. On the various contrivances by which british and foreign orchids are fertilised by insects, and on the good effects of intercrossing, 1862.

5. Anne Pusey and Marisa Wolf. Inbreeding avoidance in animals. *Trends in Ecology & Evolution*, 11(5):201–206, 1996. ISSN 0169-5347. doi: https://doi.org/10.1016/0169-5347(96)10028-8.

6. SCH Barrett. The evolution, maintenance and loss of self-incompatibility system. In J. Lovett-Doust and L. Lovett-Doust, editors, *Plant reproductive ecology: patterns and strategies*, pages 98–123. Oxford University Press, Oxford, 1988.

7. R. A. Brink and D. C. Cooper. Somatoplastic sterility in medicago sativa. *Science*, 90(2345): 545–546, 1939. ISSN 00368075, 10959203.

8. D. C. Cooper and R. A. Brink. Partial self-incompatibility and the collapse of fertile ovules as factors affecting seed formation in alfalfa. *Journal of Agricultural Research*, 60:453–472, 1940.

9. Nelson S. Bittencourt Jr, Peter E. Gibbs, and João Semir. Histological Study of Post-pollination Events in Spathodea campanulata Beauv. (Bignoniaceae), a Species with Late-acting Self-incompatibility. *Annals of Botany*, 91(7):827–834, 06 2003. ISSN 0305-7364. doi: 10.1093/aob/mcg088.

10. Steven R. Seavey and Kamaljit S. Bawa. Late-acting self-incompatibility in angiosperms. *Botanical Review*, 52(2):195–219, 1986. ISSN 00068101, 18749372.

11. PETER E. Gibbs and MARTA B. Bianchi. Does late-acting self-incompatibility (lsi) show family clustering? two more species of bignoniaceae with lsi: *Dolichandra cynanchoides* and *Tabebuia nodosa*. *Annals of Botany*, 84(4):449–457, 1999. ISSN 0305-7364. doi: https://doi.org/10.1006/anbo.1999.0933.

12. Peter E. Gibbs. Late-acting self-incompatibility – the pariah breeding system in flowering plants. *New Phytologist*, 203(3):717–734, 2014. doi: https://doi.org/10.1111/nph.12874.

13. D Charlesworth. Distribution of dioecy and self-incompatibility in angiosperms. In P.J. Greenwood, P.H. Harvey, and M. Slatkin, editors, *Evolution: Essays in honour of John Maynard Smith*, pages 237–268. Cambridge University Press, Cambridge, 1988.

14. Tammy L Sage, Robert I Bertin, and Elizabeth G Williams. Ovarian and other late-acting self-incompatibility systems. In Elizabeth G Williams, Adrienne E Clarke, and R Bruce Knox, editors, *Genetic control of self-incompatibility and reproductive development in flowering plants*, chapter 7, pages 116–140. Kluwer Academic Publishers, 1994.

15. Tammy L Sage, Mary V Price, and Nickolas M Waser. Self-sterility in ipomopsis aggregata (polemoniaceae) is due to prezygotic ovule degeneration. *American Journal of Botany*, 93 (2):254–262, 2006. doi: https://doi.org/10.3732/ajb.93.2.254.

16. S R Lipow and R Wyatt. Single gene control of postzygotic self-incompatibility in poke milkweed, asclepias exaltata l. *Genetics*, 154(2):893–907, Feb 2000. ISSN 0016-6731 (Print); 0016-6731 (Linking). doi: 10.1093/genetics/154.2.893.

17. Marta B Bianchi, Thomas R Meagher, and Peter E Gibbs. Do s genes or deleterious recessives control late-acting self-incompatibility in Handroanthus heptaphyllus (Bignoniaceae)? A diallel study with four full-sib progeny arrays. *Annals of Botany*, 127(6):723–736, 02 2021. ISSN 0305-7364. doi: 10.1093/aob/mcab031.

18. Nelson Sabino Bittencourt Júnior. Evidence for post-zygotic self-incompatibility in handroanthus impetiginosus (bignoniaceae). *Plant Reproduction*, 30(2):69–79, 2017. doi: 10.1007/s00497-017-0300-7.

19. Stephen L. Krebs and James F. Hancock. Embryonic genetic load in the highbush blueberry, *Vaccinium corymbosum* (ericaceae). *American Journal of Botany*, 78(10):1427–1437, 1991. doi: https://doi.org/10.1002/j.1537-2197.1991.tb12609.x.

20. Nic Lughadha, SJ Owen, and PJ Rudall. Preferential outcrossing in gomidesia (myrtaceae) is maintained by a post-zygotic mechanism. In S.J. Owens and P. J. Rudall, editors, *Reproductive Biology in systematics, conservation and economic botany*, pages 363–379. Kew Royal Botanic Gardens, London, 1998.

21. Norman A Johnson and M J Wade. Conditions for soft selection favoring the evolution of hybrid inviability. *Journal of Theoretical Biology*, 176(4):493 – 499, 1995. ISSN 0022-5193. doi: https://doi.org/10.1006/jtbi.1995.0215.

22. Jerry A Coyne. The evolutionary origin of hybrid inviability. *Evolution*, 28(3):505–506, Sep 1974. doi: 10.1111/j.1558-5646.1974.tb00781.x.

23. Brian Charlesworth. The evolution of lethals in the *t* - haplotype system of the mouse. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 258(1352):101–107, 1994. doi: 10.1098/rspb.1994.0149.

24. W.D. Hamilton. The moulding of senescence by natural selection. *Journal of Theoretical Biology*, 12(1):12–45, 1966. ISSN 0022-5193. doi: https://doi.org/10.1016/0022-5193(66)90184-6.

25. Yaniv J Brandvain, Alexander J Harkness, and Tanja Pyhäjärvi. Reproductive compensation and embryo screening drive the evolution of polyembryony. *bioRxiv*, 2021. doi: 10.1101/2020.11.17.387340.

26. S. J. Gould and R. C. Lewontin. The spandrels of san marco and the panglossian paradigm: a critique of the adaptationist programme. *Proceedings of the Royal Society of London. Series B. Biological Sciences*, 205(1161):581–598, 1979. doi: 10.1098/rspb.1979.0086.

27. Emmanuelle Porcher and Russell Lande. Reproductive compensation in the evolution of plant mating systems. *The New Phytologist*, 166(2):673–684, 2005. ISSN 0028646X, 14698137.

28. Russell Lande, Douglas W Schemske, and Stewart T Schultz. High inbreeding depression, selective interference among loci, and the threshold selfing rate for purging recessive lethal mutations. *Evolution*, 48(4):965–978, August 1994.

29. Benjamin C Haller and Philipp W Messer. SLiM 3: Forward Genetic Simulations Beyond the Wright–Fisher Model. *Molecular Biology and Evolution*, 36(3):632–637, 01 2019. ISSN 0737-4038. doi: 10.1093/molbev/msy228.

30. Veikko Koski. On self pollination, genetic load and subsequent inbreeding in some conifers. *Communicationes Instituti Forestalis Fenniae*, 78:1–42, 1973.

31. Satoki Sakai. Maintenance of high inbreeding depression in selfing populations: Two-stage effect of early- and late-acting mutations. *Journal of Theoretical Biology*, 502:110307, 2020. ISSN 0022-5193. doi: https://doi.org/10.1016/j.jtbi.2020.110307.

32. Satoki Sakai. Why are deleterious mutations maintained in selfing populations? an analysis of the effects of early- and late-acting mutations by a two-locus two-allele model. *Journal of Theoretical Biology*, 533:110956, 2022. ISSN 0022-5193. doi: https://doi.org/10.1016/j.jtbi.2021.110956.

33. Laurence D. Hurst. The incidences and evolution of cytoplasmic male killers. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 244(1310):91–99, 1991. doi: 10.1098/rspb.1991.0056.

34. S. A. Frank and L. D. Hurst. Mitochondria and male disease. *Nature*, 383(6597):224–224, 1996. doi: 10.1038/383224a0.

35. Neil J. Gemmell, Victoria J. Metcalf, and Fred W. Allendorf. Mother's curse: the effect of mtdna on individual fitness and population viability. *Trends in Ecology & Evolution*, 19(5): 238–244, 2021/12/14 2004.

36. M J Wade and R W Beeman. The population dynamics of maternal-effect selfish genes. *Genetics*, 138(4):1309–1314, 12 1994. ISSN 1943-2631. doi: 10.1093/genetics/138.4.1309.

37. IAN M. HASTINGS. Models of human genetic disease: how biased are the standard formulae? *Genetical Research*, 75(1):107–114, 2000. doi: 10.1017/S0016672399003961.

38. David Haig. The kinship theory of genomic imprinting. *Annual Review of Ecology and Systematics*, 31(1):9–32, 2000. doi: 10.1146/annurev.ecolsys.31.1.9.

39. M K Uyenoyama and E Newbigin. Evolutionary dynamics of dual-specificity self-incompatibility alleles. *The Plant cell*, 12(3):310–315, 03 2000. doi: 10.1105/tpc.12.3.310.

40. M K Uyenoyama, Y Zhang, and E Newbigin. On the origin of self-incompatibility haplotypes: transition through self-compatible intermediates. *Genetics*, 157(4):1805–1817, 04 2001. doi: 10.1093/genetics/157.4.1805.

41. GC Williams. *Adaptation and natural selection: A critique of some current evolutionary thought*. Princeton University Press, Princeton, NJ, 1966.

42. I M Hastings. Reproductive compensation and human genetic disease. *Genetics Research*, 77(3):277–283, Jun 2001. doi: 10.1017/s0016672301004992.

43. A D J Overall, M. Ahmad, and R A Nichols. The effect of reproductive compensation on recessive disorders within consanguineous human populations. *Heredity*, 88(6):474–479, 2002.

44. Sarah J. Baldwin and Daniel J. Schoen. Inbreeding depression is difficult to purge in self-incompatible populations of leavenworthia alabamica. *New Phytologist*, 224(3):1330–1338, 2019. doi: https://doi.org/10.1111/nph.15963.

45. Claire G Williams. Re-thinking the embryo lethal system within the pinaceae. *Botany*, 85 (7):667–677, 2007.

46. Brian C. Husband and Douglas W. Schemske. Evolution of the magnitude and timing of inbreeding depression in plants. *Evolution*, 50(1):54–70, 1996. doi: https://doi.org/10.1111/j.1558-5646.1996.tb04472.x.

Appendix

Mating Table, Single locus model. Here we show all possible offspring genotype, its adult and embryo fitness, and its frequency in each maternally family after early embryo selection. Maternal families occur in frequencies of their genotype after selection. The relative fitness of a selfed plant equals $1-\delta$, which we assume to equal zero throughout the analytical portion of the manuscript.

The values, ψ_{aa}, ψ_{Aa}, and ψ_{AA} denote the probability that an embryo is replaced by another (i.e. compensated) in aa, Aa, and AA families, respectively. The value, replace in maternal family, k equals $\text{replace}_k = \sum\sum \text{freq}_{\text{Geno:i, of type: j, in Fam:k}} \times$ embryo fitness$_i$ summed across all primary embryos for each maternal family. These values are: $\psi_{aa} = phs(1-\alpha)$, $\psi_{Aa} = s(2h+2p(1-\alpha)+\alpha)/4$, and $\psi_{AA} = s(p+(1-p)(h(1-\alpha)+\alpha))$.

maternal fam	genotype	type	adult fitness	primary	embryo fitness	freq in fam
aa	aa	Self	$1-\delta$	Primary	1	α
aa	Aa	Self	$1-\delta$	Primary	$1-hs$	0
aa	AA	Self	$1-\delta$	Primary	$1-s$	0
aa	aa	Out	1	Primary	1	$(1-p)(1-\alpha)$
aa	Aa	Out	1	Primary	$1-hs$	$p(1-\alpha)$
aa	AA	Out	1	Primary	$1-s$	0
aa	aa	Self	$1-\delta$	Backup	1	$\psi_{aa}\alpha$
aa	Aa	Self	$1-\delta$	Backup	$1-hs$	0
aa	AA	Self	$1-\delta$	Backup	$1-s$	0
aa	aa	Out	1	Backup	1	$\psi_{aa}(1-\alpha)(1-p)$
aa	Aa	Out	1	Backup	$1-hs$	$\psi_{aa}(1-\alpha)p$
aa	AA	Out	1	Backup	$1-s$	0
Aa	aa	Self	$1-\delta$	Primary	1	$\alpha/4$
Aa	Aa	Self	$1-\delta$	Primary	$1-hs$	$\alpha/2$
Aa	AA	Self	$1-\delta$	Primary	$1-s$	$\alpha/4$
Aa	aa	Out	1	Primary	1	$(1-\alpha)(1-p)/2$
Aa	Aa	Out	1	Primary	$1-hs$	$(1-\alpha)/2$
Aa	AA	Out	1	Primary	$1-s$	$(1-\alpha)(p/2)$
Aa	aa	Self	$1-\delta$	Backup	1	$\psi_{Aa}(\alpha/4)$
Aa	Aa	Self	$1-\delta$	Backup	$1-hs$	$\psi_{Aa}(\alpha/2)$
Aa	AA	Self	$1-\delta$	Backup	$1-s$	$\psi_{Aa}(\alpha/4)$
Aa	aa	Out	1	Backup	1	$\psi_{Aa}(1-\alpha)(1-p)/2$
Aa	Aa	Out	1	Backup	$1-hs$	$\psi_{Aa}(1-\alpha)/2$
Aa	AA	Out	1	Backup	$1-s$	$\psi_{Aa}(1-\alpha)p/2$
AA	aa	Self	$1-\delta$	Primary	1	0
AA	Aa	Self	$1-\delta$	Primary	$1-hs$	0
AA	AA	Self	$1-\delta$	Primary	$1-s$	α
AA	aa	Out	1	Primary	1	0
AA	Aa	Out	1	Primary	$1-hs$	$(1-p)(1-\alpha)$
AA	AA	Out	1	Primary	$1-s$	$p(1-\alpha)$
AA	aa	Self	$1-\delta$	Backup	1	0
AA	Aa	Self	$1-\delta$	Backup	$1-hs$	0
AA	AA	Self	$1-\delta$	Backup	$1-s$	$\psi_{AA}\alpha$
AA	aa	Out	1	Backup	1	0
AA	Aa	Out	1	Backup	$1-hs$	$\psi_{AA}(1-\alpha)(1-p)$
AA	AA	Out	1	Backup	$1-s$	$\psi_{AA}(1-\alpha)p$

Selection, Single locus model. . The frequency of each genotype after selection is the product of its frequency in a maternal family, the maternal family's frequency, its embryonic and adult fitness, weighted by mean fitness. The results of these calculations are somewhat long and tedious and are therefore presented in the supplemental Mathematica file.
By appropriately summing and weighting the content of the table above, and assuming complete inbreeding depression ($\delta = 1$), we find mean fitness equals

$$\overline{w} = (1-\sigma)\left(p_{\text{aa}}(\psi_{\text{aa}}+1)(1-hps) + p_{\text{Aa}}(\psi_{\text{Aa}}+1)(1-\frac{s(h+p)}{2}) + p_{\text{AA}}(\psi_{\text{AA}}+1)(1-hqs-ps)\right)$$

693 Where σ is the selfing rate, p and q are frequencies of alleles a (wild type), and A (self-sacrificial), respectively, and subscripted
694 ψ is the probability that a primary embryo in the subscripted maternal genotype is replaced.
695 From the table above, we find that the frequencies of each genotype in the next generation are as follows:

$$p'_{aa} = \frac{(1-p)(1-\alpha)(2p_{aa}(\psi_{aa}+1)+p_{pAa}(\psi_{Aa}+1))}{2\overline{w}}$$

$$p'_{Aa} = \frac{((1-\alpha)(1-hs))(2p p_{aa}(\psi_{aa}+1)+2(1-p)p_{pAA}(\psi_{AA}+1)+p_{Aa}(\psi_{Aa}+1))}{2\overline{w}}$$

$$p'_{AA} = \frac{p(1-s)(1-\alpha)\left(p+\frac{p_{pAa}\psi_{Aa}}{2}+p_{AA}\psi_{AA}\right)}{\overline{w}}$$

Fig S1: The mean number of early acting, recessive lethal ($\delta = 1$, A), and sublethal ($\delta = 0.75$, B) alleles per individual, over a combination of selfing rates (σ), mutation rates (μ), selection (s)and dominance (h) coefficients. Blue lines display results for cases in which primary embryos are replaced by siblings (our model), while blue lines display cases in which primary embryos are replaced by a non-sibling (a control).