**Title: Gene conversion limits the cost of asexuality in somatically immortal worms**

Authors: Simon Kershenbaum*, Manuel Jara-Espejo, Ashleigh S. Griffin, *Aziz Aboobaker

Author affiliations: Department of Biology, University of Oxford, Oxford, United Kingdom,

OX1 3SZ

*Corresponding authors: simon.kershenbaum@new.ox.ac.uk ;

aziz.aboobaker@biology.ox.ac.uk

Keywords: Asexuality; Muller's ratchet; Gene conversion; Planarians

## Abstract

Most multicellular organisms reproduce sexually despite the costs associated with sexuality. This has been explained as the result of selection favouring the ability to recombine the genome. The lack of recombination in asexual species constrains their adaptability and leads to the accumulation of deleterious mutations, ultimately increasing their risk of extinction. Nonetheless, successful asexual life histories persist among multicellular organisms, and explanatory mechanisms which may help limit the cost of asexuality remain enigmatic. In search of these mechanisms, we looked at that the molecular evolutionary changes in the sexual and obligately asexual strains of the planarian flatworm, *Schmidtea mediterranea*. We find that the accumulation of deleterious mutations in highly conserved genes is largely avoided in the asexual strain. We find evidence that this is achieved by somatic gene conversion in stem cells allowing for the restoration of beneficial alleles and purification of deleterious mutations. Taken together, our analysis identifies gene conversion as a mechanism which may contribute to the maintenance of asexuality in an obligately fissiparous metazoan.

## Introduction

Most multicellular life reproduces sexually, while obligate asexuality is exceedingly rare (Judson and Normark, 1996). This pattern persists despite the high costs of sex, including the "two-fold" cost of sex (Gibson et al., 2017), the breaking up of beneficial haplotypes (Charlesworth and Barton, 1996), and the broad costs associated with securing a mate (Lehtonen et al., 2012). The maintenance of sexuality in the face of such costs suggests that

asexuality, the alternative reproductive strategy, is somehow fundamentally limited and is

outcompeted by sexual strategies in the long term (Lynch et al., 1993). However, some

ancient asexual organisms exist, which raises the question of how they avoid the limitations

of asexuality (Bast et al., 2018; Butlin et al., 1998; Judson and Normark, 1996; Lázaro et al.,

2011; Wilson et al., 2018).

The potential disadvantages of asexual reproduction are driven by the tight linkage between

alleles in an asexual genome (Box. 1) (Comeron et al., 2008; Felsenstein, 1974; Hill and

Robertson, 1966). This linkage means that selection cannot act on individual alleles or allele

combinations, but instead must act on the genomes of individuals as a whole - a problem

known as "The Hill-Robertson effect" (Comeron et al., 2008; de Visser and Elena, 2007; Hill

and Robertson, 1966). This reduces the adaptability of asexual populations, which has been

shown to be particularly problematic in the context of changing environments and biotic

competition (Hamilton et al., 1990; Lively and Morran, 2014; Vergara et al., 2014).

Additionally, the inability to remove deleterious mutations from the next generation through

recombination results in the accumulation of deleterious mutations over time in a process

called "Muller's ratchet" (Lynch et al., 1993; Muller). This occurs since even the fittest

individual in an asexual population contains some deleterious mutations, and as this fittest

genome reaches fixation, so do its deleterious mutations, leading to a ratchet-like process of

mutation accumulation (Lynch et al., 1993; Muller). Taken together, the reduced ability to

unlink alleles in asexual genomes is thought to drive Muller's ratchet and the Hill-Robertson

effect, both suggested as explanations for why obligate asexual reproduction is rare

(Felsenstein, 1974).

**Box. 1**: The limitations of obligate asexual reproduction stem from the tight linkage of alleles in

the genomes that lack recombination. Muller's ratchet describes how this lack of

recombination leads to the accumulation of deleterious mutations over time. This is because

even the fittest asexual genome will have some deleterious mutations that will be passed on to

the next generation, while in sexual reproduction, genomes that only contain beneficial alleles

can theoretically be generated and selected for. Hill-Robertson interference is a broad term

describing the inefficient selection that occurs when alleles are tightly linked due to the lack of

recombination. This means that deleterious mutations can interfere with the selection for

beneficial mutations (and vice versa), while in sexual populations these mutations can be

selected upon independently. Clonal interference describes the inefficacy resulting from the

fact that two beneficial mutations in different lineages will compete with one another. Since

asexual lineages don't recombine with one another, only the fittest variant will persist, while in

sexual populations a genome containing both variants can be formed.

However, some obligate asexual species persist over evolutionary time (Barraclough et al.,

2007; Bast et al., 2018; Lázaro et al., 2011; Schön et al., 2009), suggesting that they may

have evolved mechanisms to limit the impact of long-term asexuality. Such ancient asexuals

include bdelloid rotifers, oribatid mites, darwinulid ostracods and more, all of which have

likely been asexual for millions of generations (Brandt et al., 2017; Brandt et al., 2021; Flot et

al., 2013; Mark Welch and Meselson, 2000; Schön et al., 2003; Schön et al., 2009). To date,

there is no consensus regarding the methods by which they avoid the costs associated with

asexuality. One suggested mechanism is gene conversion (Box. 2): a form of mitotic

recombination in which one allele overwrites another (Butlin et al., 1998; Flot et al., 2013;

Schön et al., 2009). While deleterious and beneficial alleles are equally likely to get

overwritten in the process of gene conversion, selection may be able to act to remove the

deleterious gene conversion events and drive the spread of beneficial alleles (Box. 2).

Ancient asexuals such as bdelloid rotifers (Flot et al., 2013) and darwinulid ostracods (Schön

et al., 2009) exhibit excess homozygosity, which has been proposed to be the result of gene

conversion, leading the authors to suggest a role for gene conversion in the long term survival

of these asexuals. However, evidence from other species suggests that gene conversion may

not be sufficient to limit mutation accumulation, since young asexual lineages of Daphnia

show relatively high rates of gene conversion while still showing a significant accumulation

of deleterious mutations (Omilian et al., 2006; Tucker et al., 2013). In neither case has it been

shown that selection acts upon gene conversion events to remove deleterious mutations and

fix beneficial alleles in the population. Taken together, there is currently no consensus about

the mechanisms that limit the costs of asexuality in ancient asexuals, and we need to assess

more groups of long-term asexuals with diverse modes of asexual reproduction to better

understand how ancient asexuals might overcome the limitations of asexuality.

**Box. 2**: Gene conversion as a method to remove deleterious mutations. Gene conversion can

limit the progression of Muller's ratchet by removing deleterious mutations from some of the

progeny. A heterozygous diploid is depicted by homologous regions of red and blue

chromosomes. The red chromosome contains a region with a deleterious mutation. The

progeny containing a gene conversion event which removes the deleterious mutation in the

red chromosome has the highest fitness (left) and can be selected for, while gene conversion

events leading to a homozygous deleterious mutation (right) will be selected against. The

most deleterious genome (a/a) is indicated by high transparency, followed by a/A, and A/A

respectively.

The planarian flatworm *S. mediterranea* is a highly regenerative animal that has both a sexual

and an asexual strain (Tan et al., 2012). The sexual strain is a cross fertilizing hermaphrodite

and the obligate asexual strain lacks reproductive organs, reproducing solely by fission

followed by whole body regeneration (Newmark et al., 2008; Saberi et al., 2016) (Fig. 1).

The asexual  strain is evidently able to thrive, despite the limitations of asexual reproduction,

providing us with an opportunity to examine potential mechanisms for dealing with the

limitations of asexuality (Lázaro et al., 2011; Sahu et al., 2017; Tan et al., 2012). The

fissiparous nature of asexual planarian reproduction is based on whole body regeneration that

is mediated by somatic, immortal, pluripotent adult stem cells called neoblasts (Wagner et al.,

2011). Continuous BrdU labelling suggests that these neoblasts have a generation time of 24-

72 hours as they are constantly in the cell cycle, suggesting that since the origin of the

asexual lineage there have been at least tens to hundreds of millions of asexual neoblast

generations (Fields and Levin, 2018; Newmark and Sánchez Alvarado, 2000), dependent

upon when the asexual strain arose (Lázaro et al., 2011). Such a high number of asexual

generations is equivalent to some of the previously studied ancient asexuals but represents an

understudied mode of asexual reproduction in metazoans. This fissiparous lifestyle and the

presence of a closely related sexual strain which allows for the direct genetic comparison of

sexual and asexual strains, makes *S. mediterranea* a valuable species for understanding how

such asexual lineages might persist over evolutionary time.

In this study, we investigate how asexual planarians may limit the costs of asexuality.

Specifically, we test whether planarian flatworms are able to escape the costs of asexuality

through gene conversion. We find that asexual planarians limit the impact of Muller's ratchet,

and that they show GC-biased excess homozygosity indicative of gene conversion.

Furthermore, we find that mutations that are predicted to be more deleterious are less

homozygous, suggesting that gene conversion events can be acted upon by selection to fix the

beneficial allele in the population and remove deleterious mutations. Thus, we find evidence

that gene conversion mediated removal of deleterious mutations may allow asexual

planarians to limit Muller's ratchet and persist despite their obligate asexuality.

**Results**

**Population structure of laboratory *S. mediterranea***

To better understand how laboratory populations of S. mediterranea are related to one

another, we used 169 asexual RNAseq libraries from 7 studies and 46 RNAseq libraries from

3 studies and evaluated their population structure (SI1) (Currie and Pearson, 2013; Davies et

al., 2017; Duncan et al., 2015; Khan and Newmark, 2022; Scimone et al., 2014; Tu et al.,

2015; van Wolfswinkel et al., 2014; Vogg et al., 2014; Zhu et al., 2015). We found that all

asexual samples clustered together independently from the sexual samples (Fig. 1A). Asexual

populations also largely clustered by BioProject, suggesting that populations are evolving

independently to a certain extent in each laboratory (Fig. 1B). To ensure that no one

population would be biasing the results, for example by being abnormally homozygous, we

checked for outliers in mean allele frequency and proportion of homozygosity across all

BioProjects (Fig. S1). We found no outliers in the asexual population (Fig. S1A-C). One

sexual population(Davies et al., 2017), had a higher proportion of rare alleles compared to the

other sexual populations (Fig. S1D). To ensure that our results are not biased by this

population, all further analyses comparing asexual to sexual allele frequencies were also done

separately against each sexual population. Most of the variants isolated were specific to the

asexual lineage (806,712 variants, which amounts to 67.1% of the variants found in this

study), and only 7.7% were shared between the sexual and asexual populations (Fig. 1C, Fig.

S1E). This abundance of asexual variants is expected since variants were called against the

available sexual genome (Grohme et al., 2018). In both sexual and asexual populations, the

majority of variants were non-coding, followed by synonymous mutations, non-synonymous

mutations, and indels respectively (Fig. 1D, Fig. S1F). In summary, our analysis suggests that

asexual populations form subpopulations across different laboratories, and are distinct from

the sexual population, with over 800,000 variants that are unique to the asexual population.

Figure 1 - Description of population structure and variants. A) PCA of all libraries in study;

B) PCA of asexual libraries in study coloured by BioProject ID; C) Venn Diagram showing

the percentage of shared and unique variants between sexual and asexual populations. D)

Barplot showing the numbers of different types of variants in sexual and asexual populations.

**Limited impact of Muller's ratchet in asexual planarians**

We began by looking for evidence of deleterious mutation accumulation in the obligate

asexual strain by using dN/dS analyses (SI2-3). We found that across the entire set of protein

coding genes (Neiro et al., 2022) (SI4), the median asexual dN/dS value is 3.6% higher than

that found in the sexual strain (Wilcoxon rank sum; $N_{Asexual} = 16,445$, $N_{Sexual} = 10,700$, $W =$

91,647,379, $P < 0.001$, Fig.2A). High dN/dS values can indicate both deleterious mutation

accumulation and positive selection, thus we assessed dN/dS values in genes that are unlikely

to be under positive selection. Specifically, we compared the dN/dS values of exons

belonging to highly conserved single copy BUSCO genes (SI5) (Manni et al., 2021), and

found no significant difference between the strains (Wilcoxon signed rank; $N_{pairs}$ = 160, V =

7,069, P = 0.284, Fig. 2B). However, when we compared the dN/dS values of exons under

significant negative selection in both strains (SI6), we found that the asexual dN/dS values

were 27% higher than in the sexual strain (Wilcoxon signed rank; Npairs = 41, V = 665, p-

value = 0.002, Fig. 2C). This is in contrast to some previous studies that found higher impacts

of Muller's ratchet in asexual lineages (dN/dS values ~200%-~3,000% higher than in their

sexual counterparts) (Barraclough et al., 2007; Bast et al., 2018; Henry et al., 2012; Johnson

and Howard, 2007; Neiman et al., 2010; Paland and Lynch, 2006). Additionally, we isolated

genes that contain exons under positive selection in the asexual population (Fig. 2D, SI7-8).

These included multiple genes related to GLIPR1-like genes (see methods), which may

suggest that these genes evolved new functions in the asexual population. Taken together,

while asexual planarians may be subject to Muller's ratchet, the accumulation of deleterious

mutations is somehow limited, and is not present in highly conserved single copy genes. This

requires a mechanistic explanation in the absence of sexual recombination.

**Figure 2** - dN/dS analysis in sexual and asexual populations. A) Violin plot showing the

dN/dS values of all exons for both sexual and asexual populations. Sexual mean dN/dS =

0.444, asexual mean dN/dS = 0.46 ; B) dN/dS of all BUSCO exons that contained both

synonymous and nonsynonymous variants in both sexual and asexual population. Sexual

mean dN/dS = 0.318, asexual mean dN/dS = 0.325; C) dN/dS of exons under negative

selection in both sexual and asexual populations. Sexual mean dN/dS = 0.084, asexual mean

dN/dS = 0.107; D) Volcano plot showing exons under positive selection in the asexual

population.

**Loss of heterozygosity mechanisms in asexuals may limit Muller's ratchet**

Mechanisms limiting Muller's ratchet in asexual planarians may involve somatic

recombination mechanisms, such as gene conversion, that in combination with selection can

remove deleterious mutations (Box. 2). A signature of such a process would be

homogenisation of the genome, leading to low polymorphism and high homozygosity. We

compared allele frequencies of synonymous variants in asexual and sexual populations and

found that asexual allele frequency is significantly higher than in the sexual population for

both shared ancestral variants and lineage specific variants suggesting that there is lower

polymorphism in the asexual population (Wilcoxon signed rank$_{ancestral}$; $N_{pairs}$ = 9,206, V =

40,369,398, $P_{ancestral}$ < 0.001, Fig. 3A; Wilcoxon rank sum$_{lineage-specific}$, $N_{asexual}$ = 30,463, $N_{sexual}$

= 13,428, W = 338,332,851, $P_{lineage-specific}$ < 0.001, Fig. 3B). Additionally, in asexuals, the

allele frequencies of ancestral variants were 45% higher than lineage specific variants, while

in the sexual population the allele frequencies of ancestral variants were only 4% higher than

lineage specific variants (Wilcoxon rank sum; $N_{asexual}$ = 309, $N_{sexual}$ = 305, W = 82,102, P <

0.001, Fig. 3C). The increased allele frequency in ancestral variants suggests a process of

homogenisation over time, which is more effective in the asexual genome.

High allele frequencies in asexuals may be explained by the clonal nature of asexual

planarians, so we tested whether they also show excess homozygosity which would be

indicative of a homogenising process that cannot be accounted for by clonality. In the

absence of recombination, half of the ancestral variants shared between the sexual and

asexual strains should be homozygous in the asexual strain (see methods). However, we find

that the mean proportion of homozygosity in asexual ancestral variants is 0.82, suggesting

that some homogenising process is taking place over time (Wilcoxon signed rank; $N_{asexual-}$

$_{ancestral\ =}$ 9,206, V = 34,205,206, P < 0.001, Fig. 3D).

We also tested whether there is excess homozygosity in the uniquely asexual variants. For

asexual specific variants, the expected proportion of homozygosity depends on the proportion

of variants that formed by mutation after the sexual-asexual split, versus those that were

inherited from their sexual ancestors and subsequently lost from the sexual population. At the

extremes, if all of them formed after the sexual-asexual split we would expect almost none of

them to be homozygous, and if all of them were inherited from their sexual ancestors we

would expect a third of them to be homozygous (see methods). We find that the mean

proportion of homozygosity in asexual specific variants is 0.32 (Fig. 3D), which in the

absence of recombination would suggest that the vast majority (97%) of asexual specific

variants were inherited from sexual ancestors and then lost from the sexual population. This

would correspond to only 914 de-novo synonymous mutations in the asexual population since

the sexual-asexual split, which is unlikely given that the asexual population split from the

sexuals hundreds of thousands to millions of years ago (Lázaro et al., 2011). Thus, we

conclude that some form of recombination acts as a homogenising force.

However, we wanted to explicitly test whether variant inheritance from a sexual ancestor

followed by subsequent loss in the sexual population can explain the homozygosity levels in

uniquely asexual variants. To this end we assessed the proportion of homozygosity in

uniquely asexual variants that are fixed within the asexual population. These represent the

ancestral asexual variants of our population, and so are likely to be older than other variants

that are only found in a subset of samples. We found that the mean proportion of

homozygosity in ancestral asexual specific variants was 0.58, suggesting that there is excess

homozygosity even if all these asexual specific ancestral variants were originally inherited

from sexual ancestors rather than forming de-novo in the asexual lineage (Wilcoxon signed

rank; $N_{\text{ancestral asexual specific}} = 2,275$, $V = 2,121,916$, $P < 0.001$, Fig. 3E). Additionally, ancestral

variants shared between the sexual and asexual population are more homozygous than

asexual specific variants (Wilcoxon rank sum; $N_{\text{asexual-ancestral}} = 9,206$, $N_{\text{Asexual specific}} = 30,463$,

$W = 211,577,670$, $P < 0.001$, Fig. 3D), suggesting that more ancient variants are more likely

to be homozygous. We also found that within asexual specific variants, those found in more

samples were more likely to be homozygous (Binomial GLM, $N = 30,463$, $P < 0.001$, Fig3E).

Taken together, this suggests that some homogenising process might be taking place in the

asexual genome, and that its impact is larger on the more ancient variants compared to those

with a more recent origin. These results indicate a genome wide mechanism in driving the

loss of heterozygosity in asexual planarians that could potentially limit mutation

accumulation in this asexual lineage.

**Figure 3** - Excess homozygosity in the asexual population. Red dots indicate means. A)

Allele frequencies of ancestral variants shared by sexual and asexual populations. Mean

sexual allele frequency = 0.32. Mean asexual allele frequency = 0.9; B) Allele frequencies of

lineage specific. Mean sexual allele frequency = 0.31. Mean asexual allele frequency = 0.64;

C) Violin plot showing the ratio of average allele frequencies per scaffold of ancestral vs

lineage specific variants. Mean sexual ratio = 1.038. Mean asexual ratio= 1.446; D) Boxplot

showing the proportion of homozygous libraries per variant in ancestral (shared between the

sexual and asexual populations) and lineage specific variants in the asexual population.

Dashed blue lines indicate the expected proportion of homozygosity in the absence of

recombination; E) Binomial GLM showing that the proportion of homozygosity in asexual

lineage specific variants increases with the number of libraries containing the variant. When

all libraries contain the variant the mean proportion of homozygosity is 0.58. The dashed line

indicates the expected homozygosity assuming that all asexual specific variants were

inherited from sexual ancestors.

**GC-biased gene conversion in asexual planarians**

We hypothesised that since gene conversion has a GC bias (Chen et al., 2007), homozygous

variants should display this signature if they were indeed generated by gene conversion. We

find that homozygous variants are GC enriched compared to heterozygous variants, with 53%

of homozygous variants being composed of GC nucleotides compared to 35% in

heterozygous variants (Chi-sqr = 1,328.6, df = 1, $N_{HOM}$ = 17,487, $N_{HET}$ = 22,517, P < 0.001,

Fig. 4A), suggesting a GC bias in the mechanism driving high asexual homozygosity.

Additionally, we find that homozygosity is a significant predictor of GC-AT identity, with

more homozygous variants being more likely to have a GC identity (Binomial GLM, N =

44,669, P < 0.001, Fig. 4B). Furthermore, we show that AT to GC mutations are 1.5 times

more homozygous than GC to AT mutations, with mean homozygosity proportions of 0.57

and 0.38 respectively (Wilcoxon rank sum, $N_{AT \to GC}$ = 17,984, $N_{GC \to AT}$ = 21,483 W =

232,985,919, P < 0.001, Fig. 4C). This GC bias is not driven by AT to GC mutations across

the transcriptome generally (Fig. S3). Finally, we find that variants with higher allele

frequency are more likely to have a GC identity, and that this bias is stronger in the asexual

strain compared to the sexual (binomial GLM, P < 0.001, Fig. 4D). Taken together, this

suggests that GC biased gene conversion takes place in the asexual planarian, leading to a

loss of heterozygosity which could limit the impact of Muller's ratchet (Box. 2).

**Figure 4** - GC bias in homozygous variants. A) Bar plot showing that the proportion of GC is

higher in homozygous variants compared to heterozygous variants. Black line indicates the

50% line; B) Binomial GLM showing that variants with a higher proportion of homozygosity

are more likely to be of GC identity in asexual planarians; C) Violin plot showing that AT to

GC mutations are more homozygous than GC to AT mutations. Black dots indicate medians

and black cross bars indicate means; D) Binomial GLM showing that variants with higher

allele frequencies are more likely to be of GC identity in both sexuals and asexual, but in

asexuals the effect is larger.

**Selection on gene conversion events removes deleterious mutations**

Since gene conversion has the potential to generate homozygotes for both beneficial and

deleterious variants, selection must be able to remove deleterious gene conversion events for

gene conversion to be a viable solution to Muller's ratchet. We find that mutations that are

likely to be more deleterious due to their effect on predicted proteins (see methods), are less

likely to be homozygous, suggesting that deleterious gene conversion events are selected

against. We find that coding SNPs have a mean homozygosity proportion of 0.41 and are

more homozygous than coding indels which have a mean homozygosity proportion of 0.25

(Wilcoxon rank sum; $N_{indels} = 883$, $N_{SNPs} = 64,220$, $W = 23,118,081$, $P < 0.001$, Fig. 5A).

Additionally, coding SNPs have higher allele frequencies than coding indels, with mean

allele frequencies of 0.69 and 0.36 respectively (Wilcoxon rank sum; $N_{indels} = 883$, $N_{SNPs} =$

64,220, $W = 11,652,733$, $P < 0.001$, Fig. 5B). These patterns could be driven by other factors

such as error rates in each category of mutation, so we also compared the homozygosity and

allele frequencies of mutations within each category. We found that the homozygosity and

allele frequencies were higher in synonymous vs nonsynonymous SNPs (Wilcoxon rank sum;

$N_{Synonymous} = 39,669$, $N_{Nonsynonymous}$ 24,471=, $W_{Hom} = 462,855,666$, $P_{Hom} < 0.001$, $W_{AF} =$

452,432,400, $P_{AF} < 0.001$, Fig5C-D), non-framshifting indels vs framshiting indels (Wilcoxon

rank sum; $N_{Non\text{-}frameshifting} = 154$, $N_{Frameshifting} = 664$, $W_{Hom} = 44,439$, $P_{Hom} < 0.001$, $W_{AF} =$

23,168, $P_{AF} < 0.001$, Fig5C-D), and Nonsynonymous SNPs vs nonsense SNPs (Wilcoxon

rank sum; $N_{Nonsynonymous} = 24,471$, $N_{Nonsense} = 65$, $W_{Hom} = 613,538$, $P_{Hom} < 0.001$, $W_{AF} =$

494097, $P_{AF} < 0.001$ Fig. 5C-D). This suggests that mutations more likely to be deleterious

are less likely to be homozygous and are rarer in the population. However, in contrast to this

pattern, mutations in noncoding regions were altogether more heterozygous and had lower

allele frequencies than those in coding regions (Wilcoxon rank sum; $N_{Noncoding} = 21,908$,

$N_{Coding} = 74,724$, $W_{Hom} = 891,850,288$, $P_{Hom} < 0.001$, $W_{AF} = 860,143,632$, $P_{AF} < 0.001$, Fig.

S4) That being said, the majority of noncoding mutations are found within repetitive regions

(Fig. S4) which were shown in previous studies to have high mutation and erroneous

mapping rates that may also explain their abnormally high levels of heterozygosity (Nishant

et al., 2009; Patil and Vijay, 2021). We propose that when gene conversion events lead to a

homozygous deleterious mutation in a cell, the homozygous deleterious clone is eliminated

by intra-organismal and inter-organismal selection, leading to the lower observed

homozygosity in variants that are predicted to be more deleterious. The separation of alleles

via gene conversion followed by subsequent selection against unfit stem cell clones may help

limit the impact of both Muller's ratchet and the Hill-Robertson effect.

**Figure 5** - Predicted deleterious mutations are less likely to be homozygous. Dots indicate

medians, crossbars indicate means. A) Mean proportions of homozygosity are lower in

coding indels than coding SNPs; B) Mean allele frequencies are lower in coding indels than

coding SNPs; C) Frameshifting indels have lower homozygosity than non-frameshifting

indels (mean proportion homozygous of 0.215 and 0.335 respectively). Nonsense variants

have lower homozygosity than nonsynonymous variants (mean proportion homozygous of

0.15 and 0.38 respectively). Nonsynonymous variants have lower homozygosity than

synonymous variants (mean proportion homozygous of 0.38 and 0.43 respectively); D)

Frameshifting indels have lower allele frequencies than non-frameshifting indels (mean allele

frequencies of 0.3 and 0.57 respectively). Nonsense variants have lower allele frequencies

than nonsynonymous variants (mean allele frequencies of 0.46 and 0.66 respectively).

Nonsynonymous variants have lower allele frequencies than synonymous variants (mean

allele frequencies of 0.66 and 0.7 respectively).

**Discussion**

Mechanisms explaining how some asexual species persist despite the potential costs of

asexuality have remained elusive. Our results support a role for gene conversion in unlinking

loci in the asexual genome of *S. mediteranea*. Combined with selection against deleterious

homozygotes this allows these asexual planarians to limit the impact of Muller's ratchet and

Hill-Robertson effects, potentially explaining how this obligate asexual can persist over

evolutionary time.

Our results suggest that the impact of Muller's ratchet is relatively limited in asexual

planarians. This is surprising since it has been shown both empirically and by simulation that

Muller's ratchet should drive the rapid accumulation of deleterious mutations even in recent

asexual lineages (Loewe and Lamatsch, 2008; Lynch and Gabriel, 1990; Neiman et al., 2010;

Tucker et al., 2013). For example, *Daphnia pulex*, an asexual lineage that is estimated to be

only around 1,000 years old, is already showing signs of deleterious mutation accumulation

(Tucker et al., 2013). Additionally, *Potamopyrgus antipodarum*, a freshwater snail in which

the sexual form coexists with a recent obligate asexual form, shows higher rates of

deleterious mutations in the asexual strain (Johnson and Howard, 2007; Neiman et al., 2010).

This suggests that Muller's ratchet can act rapidly to drive mutation accumulation in

asexuals. Thus, one might expect asexual planarians to show clear signs of Muller's ratchet,

given that the asexual stem cells that make up their soma have been asexual for at least tens

of millions of generations. However, they seem to only show an 27% increase of dN/dS in

exons under negative selection, and this may be driven by the higher number of variants in

the asexual dataset, increasing the statistical power for calling exons under negative selection

and including those with higher dN/dS ratios. Furthermore, there is no evidence for elevated

dN/dS ratios in highly conserved BUSCO genes, where nonsynonymous mutations are likely

to be deleterious.

This lack of high mutation accumulation in asexual planarians could explain how they have

persisted as asexuals and is similar to the patterns seen in some other long term asexual

lineages (Brandt et al., 2017; Swanstrom et al., 2011). For example, asexual oribatid mites

have been shown to have lower dN/dS ratios compared to their sexual counterparts,

suggesting that, in this case, purifying selection might be stronger in the asexual species

(Brandt et al., 2017). In bdelloid rotifers nuclear genes do not show signs of Mullers ratchet

(Birky et al., 2005; Welch and Meselson, 2001). Others have shown an accumulation of

deleterious mutations in bdelloid mitochondrial genes (Barraclough et al., 2007), but have

later suggested that this may be due to other factors which when taken into account suggest

no signs of Muller's ratchet in the mitochondrial genes of bdelloid rotifers (Swanstrom et al.,

2011). The lack of strong impacts of Muller's ratchet on some long term asexuals suggests

that they may possess mechanisms to limit the accumulation of deleterious mutations.

Some authors have previously suggested gene conversion as a mechanism to limit Muller's

ratchet in asexual lineages (Flot et al., 2013; Loewe and Lamatsch, 2008; Schön et al., 2009).

The idea is that deleterious alleles can be overwritten by fit alleles, and that selection will

favour these events, ultimately removing deleterious mutations from the population (Box. 2).

However, beneficial alleles can equally be overwritten by deleterious alleles in the gene

conversion process, and so selection must be able to act against these events to effectively

counteract Muller's ratchet. This might explain why some asexuals show high rates of gene

conversion without limiting Mullers ratchet (Omilian et al., 2006; Tucker et al., 2013).

Furthermore, for gene conversion to stop the progression of Mullers ratchet, it may need to

occur at a rate higher than the rate of point mutations. Otherwise, mutations may accumulate

at a rate faster than gene conversion can remove them. This might be the case in some

asexuals such as *Timema* stick insects, which show lower rates of gene conversion compared

to their sexual relatives, and dN/dS values 3.6-13.4 times higher compared to their sexual

counterparts (Bast et al., 2018; Henry et al., 2012). Despite these caveats, gene conversion

remains a potential mechanism for limiting the cost of asexuality. This is in part because it is

a byproduct of DNA damage repair mechanisms that are available to diploid (or higher

ploidy) organisms (Chen et al., 2007) and so, it does not need to evolve *de novo* as a solution

to asexuality. Supporting this are the shared signatures of gene conversion between bdelloid

rotifers (Flot et al., 2013), darwinulid ostracods (Butlin et al., 1998), and now

*S.mediterranea*.

While gene conversion may be important in both fissiparous and parthenogenic asexuality,

we hypothesise that it might be significantly more effective in fissiparous asexuality. This is

due to the variation within the soma being heritable in fissiparous organisms, increasing the

amount of variation subject to selection (Howe et al., 2022). This is particularly relevant in

*S.mediterranea* whose immortal stem cells are extremely abundant and represent a

considerable proportion of the soma (van Wolfswinkel et al., 2014). These high numbers of

stem cells are important since each stem cell will experience unique gene conversion events,

and with a large enough population of stem cells it is likely that some of them will overwrite

deleterious alleles with their beneficial counterparts. These stem cells may then outcompete

other stem cells within the worm and spread, potentially reaching fixation in the worm.

Selection may then act between individual worms, selecting for those that have purged

deleterious mutations, ultimately removing deleterious mutations from the population. Thus,

fissiparous organisms may be more efficient than parthenogenetic asexuals at removing

deleterious mutations from the population since gene conversion within their soma can also

be selected upon to remove deleterious mutations.

However, it is important to point out that there is always the possibility that asexual

planarians have rare cryptic sex that we are simply not aware of, a fundamental difficulty

when researching the impacts of asexuality. For example, the brine shrimp, *Artemia*

*parthenogenetica*, was once thought to be asexual but was later shown to undergo rare sex in

a mass crossing experiment (Boyer et al., 2021). Additionally, even the presumed ancient

asexual bdelloid rotifer *Macrotrachella quadricornifera* has had its asexuality contested with

evidence supporting allele sharing between different isolates (Laine et al., 2022). However,

similar patterns can potentially be explained without sex by ancestral duplications followed

by independent allele losses (Wilson et al., 2024). Either way, it is entirely possible that in the

case of *S. mediterrannea* rare cryptic sex might occur and might explain how they avoid the

accumulation of deleterious mutations. However, it would not explain the strong signatures of

gene conversion, which is not as clear in the genome of sexual planarians (Fig. 3C, Fig. 4D).

Finally, even if rare sex occurs, gene conversion may still be useful as a mechanism to deal

with the cost of highly reduced levels of sexual recombination.

In summary, obligate asexual reproduction in multicellular organisms is rare and thought to

be an evolutionarily ephemeral phenomenon due to the fundamental limitations of asexuality.

When we examine *S. mediterrannea*, which adopts this strategy successfully, we are able to

assess how it limits the costs of obligate asexuality. Our results suggest that it likely uses

asexual recombination in the form of gene conversion, to unlink loci and thus limit the impact

of Muller's ratchet and Hill-Robertson interference. While such mechanisms may be available

to most asexual lineages, the opportunity for selection to act on a population of somatic stem

cells may be key in allowing asexual planarians to persist.

## Materials and Methods

### Collecting available RNAseq data

RNAseq data was collected from previous studies with a total of 169 libraries from 7 studies

for the asexual and 46 libraries from 3 different sexual populations (SI1) (Currie and Pearson,

2013; Davies et al., 2017; Duncan et al., 2015; Khan and Newmark, 2022; Scimone et al.,

2014; Tu et al., 2015; van Wolfswinkel et al., 2014; Vogg et al., 2014; Zhu et al., 2015).

When choosing the studies, we only used those that used illumina sequencing and had clear

annotations of the strains used. The Oxford sexual *S. mediterranea* population data was

acquired by extracting RNA from whole, mature sexual worms using a standard Trizol

extraction (Liu and Rink, 2018). Extracted RNA was then sequenced with Ilumina

sequencing of 150bp paired end reads. Raw reads were deposited at the SRA repository

(XXXX/XXXX).

### Variant calling and filtration

RNAseq data was then trimmed and quality filtered with trimmomatic and FASTQC using

default settings (Bolger et al., 2014). STAR was then used to map the RNAseq reads to the

dd_Smes_g4 genome (Dobin et al., 2013; Grohme et al., 2018). For variant calling we used

the GATK  4.2.2.0 software following the best practice (Van der Auwera et al., 2013). We

used the suggested parameters for variant filtration (SI9), except that we allowed for variants

that had no reads supporting the reference (these are usually filtered out since some of the

quality measures require a comparison of reads supporting the reference and alternative

alleles). This provided us with a list of variants in each population (SI10).

**Defining Orthologues**

We ran Orthofinder using a proteome from the Neiro et al annotation to isolate the

orthologues of each predicted gene (Emms and Kelly, 2019; Neiro et al., 2022). To reduce

redundancy, we first ran CD-hits to collapse all peptides with 99% identity to a single

transcript ID.

**dN/dS analysis**

We used the dNdScv R package for dN/dS analysis and annotation of mutation impact in

coding regions (Martincorena et al., 2017). dNdScv uses an expansion on the Poisson

framework to obtain dN/dS values and a full tri-nucleotide model to avoid systematic bias

due to sequence context (Martincorena et al., 2017). To get the list of genes for dN/dS

analysis we used the list included in the OrthoFinder output, and further filtered for the

longest transcript per gene. We then ran dNdScv with no covariates and no limits on the

number of variants per CDS. When comparing dN/dS values between the strains in genes

under purifying selection, we used the local model from dNdScv, since the overdispersion

value for the sexual dataset was lower than 1. To compare dN/dS values of BUSCO gene

exons between the strains, we isolated single copy BUSCO genes in planarians using BUSCO

(metazoan gene list). We then isolated BUSCO exons that had both synonymous and

nonsynonymous mutations within them (to allow for meaningful dN/dS assessment using the

dNdScv local model) and compared their dNdS values provided by dNdScv. We also

compared dN/dS ratios in exons that were under negative selection in both strains. These

were defined by having a qmis_loc < 0.05 and wmis_loc < 1 in the dN/dS output (SI3-SI4,

SI6). Finally, we used the global model to isolate genes under positive selection in asexual

planarians (SI7-SI8), and top blast hits were taken from planmine (Rozanski et al., 2019).

**Analysis of zygosity and allele frequency**

To assess allele frequency and homozygosity levels accurately, we only used information

about variants from samples that had a minimum depth of 10 at the site (SI11). To achieve

this, we split the VCF into individual VCFs (one per sample), and then filtered out any

variants that were not covered by at least 10 reads. Allele frequency in each sample was

calculated using VCFtools (0.1.16) freq function (Danecek et al., 2011). This resulted in each

variant obtaining a 0, 0.5 or 1 value for each sample, with the numbers corresponding to

homozygous reference, heterozygous, and homozygous alternative respectively. We could

then use this information to calculate allele frequency in the population for each variant (total

number of alternative alleles in the population / total number of alleles in the population). We

also calculated the proportion of libraries that were homozygous for each variant (number of

libraries homozygous for alternative variant / number of libraries containing the variant).

Note that assuming a complete lack of recombination in asexuals, the measure of

homozygosity from a group of individuals should be very similar to the measure of

homozygosity within a single individual.

**Expected values for mean proportion of homozygosity**

To test whether there is excess homozygosity in the asexual strain, we compared the observed

and expected proportions of homozygosity in the asexual population. When testing for excess

homozygosity, or GC bias in homozygous variants (Fig. 4), only synonymous mutations were

used to limit the impact of selection. Assuming there is no recombination, the expected mean

proportion of homozygosity for ancestral variants (those shared between sexual and asexual

strains) is 0.5. This is because for a variant to be ancestral, it must have been inherited from a

sexual worm as at least one of the alleles in the founding asexual population. Thus, whether

an ancestral variant is homozygous depends only on whether the variants was also inherited

as the second allele, meaning that half of the variants are expected to be homozygous. The

expected proportion of homozygous asexual specific variants cannot be precisely known

because it depends on how many of them were formed by mutation after the sexual-asexual

split, and how many of them were inherited from a sexual ancestor and subsequently lost

from the sexual population. However, we can place lower and upper limits for the expected

homozygosity in the two extremes. The first case, which defines the lower limit, is a scenario

in which all asexual specific variants arose by mutation after the sexual-asexual split. In the

absence of recombination, we would expect this to lead to no homozygous variants (since the

de-novo mutations should be heterozygous). In the second case, which sets the upper limit,

all asexual specific variants were inherited from a sexual ancestor and so the proportion of

homozygous variants depends on Mendelian principles. This means that there are 3 possible

ways that the alternative alleles could have been inherited, alt/alt, alt/ref, and ref/alt, meaning

that if all 3 options are equally likely, we would expect one third of all asexual specific

variants to be homozygous (Fig. S4).

**Assessing evidence for effective selection on gene conversion events**

To assess whether there is effective selection to remove deleterious gene conversion events,

we wanted to test whether more harmful mutations are less likely to be homozygous. Since

we cannot know the precise impact of each mutation on fitness, we broadly ranked variants

based on the amount of change the mutation is likely to have on the protein sequence. In our

analysis we assumed that coding indels are more deleterious on average than coding SNPs,

frameshifting indels are more deleterious than non-frameshifting indels, nonsynonymous

mutations are more deleterious than synonymous mutations, and nonsense mutations are

more deleterious than nonsynonymous mutations.

It would also be reasonable to assume that that, on average, noncoding mutations are less

deleterious than coding mutations. However, noncoding regions are often rich in repeats

which can complicate the assessment of homozygosity and allele frequency. We wanted to

test if this was indeed the case in our data, so we ran RepeatMasker 4.1.2-p1 on the

dd_Smes_g4 genome which gave us a gff3 file containing all the repetitive regions in the

genome. We then used bedtools (Quinlan and Hall, 2010) intersect to test the proportion of

coding vs noncoding variants that are found within repetitive regions.

**Supplementary Data:**

SI1 – List of all libraries used in study (Sexual RNAseq from Oxford is unpublished).

SI2 - Asexual dN/dS local model.

SI3 – Sexual dN/dS local model.

SI4 – List of all exons used in study.

SI5 – List of all BUSCO exons with synonymous and nonsynonymous mutations in both

sexual and asexual strains.

SI6 – List of all exons under negative selection in both sexual and asexual strains

SI7 – Asexual dN/dS global model.

SI8 - List of exons under positive selection in asexual strain using the global dNdScv model.

SI9 – variant filtration criteria.

S10 – List of variants

S11 – List of variants where homozygosity and allele frequency information only comes from

libraries with at least a depth of 10 at the variant site.

**Acknowledgements**

We would like to thank Tim Barraclough, Chris Wilson, and Tymoteusz Pieszko for their

helpful comments on the manuscript.

**Data availability**

Supplementary information and data can be found at:

https://github.com/SimonOX-Gene/Smed_GeneConversion

Additional information is available upon request.

**Bibliography**

**Barraclough, T. G., Fontaneto, D., Ricci, C. and Herniou, E. A.** (2007). Evidence for Inefficient
Selection Against Deleterious Mutations in Cytochrome Oxidase I of Asexual Bdelloid
Rotifers. *Molecular Biology and Evolution* **24**, 1952–1962.

**Bast, J., Parker, D. J., Dumas, Z., Jalvingh, K. M., Tran Van, P., Jaron, K. S., Figuet, E., Brandt,**
**A., Galtier, N. and Schwander, T.** (2018). Consequences of Asexuality in Natural
Populations: Insights from Stick Insects. *Molecular Biology and Evolution* **35**, 1668–
1677.

**Birky, C. W., Wolf, C., Maughan, H., Herbertson, L. and Henry, E.** (2005). Speciation and
Selection without Sex. *Hydrobiologia* **546**, 29–45.

**Bolger, A. M., Lohse, M. and Usadel, B.** (2014). Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120.

**Boyer, L., Jabbour-Zahab, R., Mosna, M., Haag, C. R. and Lenormand, T.** (2021). Not so
clonal asexuals: Unraveling the secret sex life of Artemia parthenogenetica. *Evol Lett* **5**,
164–174.

**Brandt, A., Schaefer, I., Glanz, J., Schwander, T., Maraun, M., Scheu, S. and Bast, J.** (2017).
Effective purifying selection in ancient asexual oribatid mites. *Nat Commun* **8**, 873.

**Brandt, A., Tran Van, P., Bluhm, C., Anselmetti, Y., Dumas, Z., Figuet, E., François, C. M.,**
**Galtier, N., Heimburger, B., Jaron, K. S., et al.** (2021). Haplotype divergence supports
long-term asexuality in the oribatid mite Oppiella nova. *Proceedings of the National*
*Academy of Sciences* **118**, e2101485118.

**Butlin, R., Schön, I. and Martens, K.** (1998). Asexual reproduction in nonmarine ostracods.
*Heredity* **81**, 473–480.

**Charlesworth, B. and Barton, N. H.** (1996). Recombination load associated with selection for
increased recombination. *Genetics Research* **67**, 27–41.

**Chen, J.-M., Cooper, D. N., Chuzhanova, N., Férec, C. and Patrinos, G. P.** (2007). Gene
conversion: mechanisms, evolution and human disease. *g e n e ti c s* 14.

**Comeron, J. M., Williford, A. and Kliman, R. M.** (2008). The Hill–Robertson effect: evolutionary
consequences of weak selection and linkage in finite populations. *Heredity* **100**, 19–31.

**Currie, K. W. and Pearson, B. J.** (2013). Transcription factors lhx1/5-1 and pitx are required for
the maintenance and regeneration of serotonergic neurons in planarians. *Development*
**140**, 3577–3588.

**Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker,**
**R. E., Lunter, G., Marth, G. T., Sherry, S. T., et al.** (2011). The variant call format and
VCFtools. *Bioinformatics* **27**, 2156–2158.

**Davies, E. L., Lei, K., Seidel, C. W., Kroesen, A. E., McKinney, S. A., Guo, L., Robb, S. M.,**
**Ross, E. J., Gotting, K. and Alvarado, A. S.** (2017). Embryonic origin of adult stem cells
required for tissue homeostasis and regeneration. *eLife* **6**, e21052.

**de Visser, J. A. G. M. and Elena, S. F.** (2007). The evolution of sex: empirical insights into the
roles of epistasis and drift. *Nat Rev Genet* **8**, 139–149.

**Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,**
**M. and Gingeras, T. R.** (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*
**29**, 15–21.

**Duncan, E. M., Chitsazan, A. D., Seidel, C. W. and Alvarado, A. S.** (2015). Set1 and MLL1/2
target distinct sets of functionally different genomic loci in vivo. *Cell Rep* **13**, 2741–2755.

**Emms, D. M. and Kelly, S.** (2019). OrthoFinder: phylogenetic orthology inference for
comparative genomics. *Genome Biology* **20**, 238.

630 **Felsenstein, J.** (1974). THE EVOLUTIONARY ADVANTAGE OF RECOMBINATION. *Genetics* **78**,
631   737–756.

632 **Fields, C. and Levin, M.** (2018). Are Planaria Individuals? What Regenerative Biology is Telling
633   Us About the Nature of Multicellularity. *Evol Biol* **45**, 237–247.

634 **Flot, J.-F., Hespeels, B., Li, X., Noel, B., Arkhipova, I., Danchin, E. G. J., Hejnol, A.,**
635   **Henrissat, B., Koszul, R., Aury, J.-M., et al.** (2013). Genomic evidence for ameiotic
636   evolution in the bdelloid rotifer Adineta vaga. *Nature* **500**, 453–457.

637 **Gibson, A. K., Delph, L. F. and Lively, C. M.** (2017). The two-fold cost of sex: Experimental
638   evidence from a natural system. *Evol Lett* **1**, 6–15.

639 **Grohme, M. A., Schloissnig, S., Rozanski, A., Pippel, M., Young, G. R., Winkler, S., Brandl,**
640   **H., Henry, I., Dahl, A., Powell, S., et al.** (2018). The genome of Schmidtea
641   mediterranea and the evolution of core cellular mechanisms. *Nature* **554**, 56–61.

642 **Hamilton, W. D., Axelrod, R. and Tanese, R.** (1990). Sexual reproduction as an adaptation to
643   resist parasites (a review). *Proceedings of the National Academy of Sciences* **87**, 3566–
644   3573.

645 **Henry, L., Schwander, T. and Crespi, B. J.** (2012). Deleterious Mutation Accumulation in
646   Asexual Timema Stick Insects. *Molecular Biology and Evolution* **29**, 401–408.

647 **Hill, W. G. and Robertson, A.** (1966). The effect of linkage on limits to artificial selection.
648   *Genetics Research* **8**, 269–294.

649 **Howe, J., Rink, J. C., Wang, B. and Griffin, A. S.** (2022). Multicellularity in animals: The
650   potential for within-organism conflict. *Proc. Natl. Acad. Sci. U.S.A.* **119**, e2120457119.

651 **Johnson, S. G. and Howard, R. S.** (2007). Contrasting Patterns of Synonymous and
652   Nonsynonymous Sequence Evolution in Asexual and Sexual Freshwater Snail Lineages.
653   *Evolution* **61**, 2728–2735.

654 **Judson, O. P. and Normark, B. B.** (1996). Ancient asexual scandals. *Trends in Ecology &*
655   *Evolution* **11**, 41–46.

656 **Khan, U. W. and Newmark, P. A.** (2022). Somatic regulation of female germ cell regeneration
657   and development in planarians. *Cell Reports* **38**, 110525.

658 **Laine, V. N., Sackton, T. B. and Meselson, M.** (2022). Genomic signature of sexual
659   reproduction in the bdelloid rotifer *Macrotrachella quadricornifera*. *Genetics* **220**,
660   iyab221.

661 **Lázaro, E. M., Harrath, A. H., Stocchino, G. A., Pala, M., Baguñà, J. and Riutort, M.** (2011).
662   Schmidtea mediterraneaphylogeography: an old species surviving on a few
663   Mediterranean islands? *BMC Evolutionary Biology* **11**, 274.

664 **Lehtonen, J., Jennions, M. D. and Kokko, H.** (2012). The many costs of sex. *Trends in Ecology &*
665   *Evolution* **27**, 172–178.

666    **Liu, S.-Y. and Rink, J. C.** (2018). Total RNA Isolation from Planarian Tissues. In *Planarian*
667           *Regeneration: Methods and Protocols* (ed. Rink, J. C.), pp. 259–265. New York, NY:
668           Springer.

669    **Lively, C. M. and Morran, L. T.** (2014). The ecology of sexual reproduction. *Journal of*
670           *Evolutionary Biology* **27**, 1292–1303.

671    **Loewe, L. and Lamatsch, D. K.** (2008). Quantifying the threat of extinction from Muller's
672           ratchet in the diploid Amazon molly (Poecilia formosa). *BMC Evolutionary Biology* **8**, 88.

673    **Lynch, M. and Gabriel, W.** (1990). Mutation Load and the Survival of Small Populations.
674           *Evolution* **44**, 1725–1737.

675    **Lynch, M., Bürger, R., Butcher, D. and Gabriel, W.** (1993). The Mutational Meltdown in
676           Asexual Populations. *Journal of Heredity* **84**, 339–344.

677    **Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A. and Zdobnov, E. M.** (2021). BUSCO
678           Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic
679           Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology*
680           *and Evolution* **38**, 4647–4654.

681    **Mark Welch, D. B. and Meselson, M.** (2000). Evidence for the Evolution of Bdelloid Rotifers
682           Without Sexual Reproduction or Genetic Exchange. *Science* **288**, 1211–1215.

683    **Martincorena, I., Raine, K. M., Gerstung, M., Dawson, K. J., Haase, K., Loo, P. V., Davies, H.,**
684           **Stratton, M. R. and Campbell, P. J.** (2017). Universal Patterns of Selection in Cancer
685           and Somatic Tissues. *Cell* **171**, 1029-1041.e21.

686    **Muller, H. J.** THE RELATION OF RECOMBINATION TO MUTATIONAL ADVANCE. 8.

687    **Neiman, M., Hehman, G., Miller, J. T., Logsdon, J. M. and Taylor, D. R.** (2010). Accelerated
688           Mutation Accumulation in Asexual Lineages of a Freshwater Snail. *Mol Biol Evol* **27**,
689           954–963.

690    **Neiro, J., Sridhar, D., Dattani, A. and Aboobaker, A.** (2022). Identification of putative
691           enhancer-like elements predicts regulatory networks active in planarian adult stem
692           cells. *eLife* **11**, e79675.

693    **Newmark, P. A. and Sánchez Alvarado, A.** (2000). Bromodeoxyuridine Specifically Labels the
694           Regenerative Stem Cells of Planarians. *Developmental Biology* **220**, 142–153.

695    **Newmark, P. A., Wang, Y. and Chong, T.** (2008). Germ Cell Specification and Regeneration in
696           Planarians. *Cold Spring Harb Symp Quant Biol* **73**, 573–581.

697    **Nishant, K. T., Singh, N. D. and Alani, E.** (2009). Genomic mutation rates: What high-
698           throughput methods can tell us. *Bioessays* **31**, 912–920.

699    **Omilian, A. R., Cristescu, M. E. A., Dudycha, J. L. and Lynch, M.** (2006). Ameiotic
700           recombination in asexual lineages of Daphnia. *Proceedings of the National Academy of*
701           *Sciences* **103**, 18638–18643.

702    **Paland, S. and Lynch, M.** (2006). Transitions to Asexuality Result in Excess Amino Acid
703           Substitutions. *Science* **311**, 990–992.

**Patil, A. B. and Vijay, N.** (2021). Repetitive genomic regions and the inference of demographic
history. *Heredity (Edinb)* **127**, 151–166.

**Quinlan, A. R. and Hall, I. M.** (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* **26**, 841–842.

**Rozanski, A., Moon, H., Brandl, H., Martín-Durán, J. M., Grohme, M. A., Hüttner, K.,**
**Bartscherer, K., Henry, I. and Rink, J. C.** (2019). PlanMine 3.0—improvements to a
mineable resource of flatworm biology and biodiversity. *Nucleic Acids Research* **47**,
D812–D820.

**Saberi, A., Jamal, A., Beets, I., Schoofs, L. and Newmark, P. A.** (2016). GPCRs Direct
Germline Development and Somatic Gonad Function in Planarians. *PLoS Biol* **14**,
e1002457.

**Sahu, S., Dattani, A. and Aboobaker, A. A.** (2017). Secrets from immortal worms: What can we
learn about biological ageing from the planarian model system? *Seminars in Cell &*
*Developmental Biology* **70**, 108–121.

**Schön, I., Martens, K., Van Doninck, K. and Butlin, R. K.** (2003). Evolution in the slow lane:
molecular rates of evolution in sexual and asexual ostracods (Crustacea: Ostracoda):
MOLECULAR EVOLUTION IN SEXUAL and ASEXUAL OSTRACODS. *Biological Journal of*
*the Linnean Society* **79**, 93–100.

**Schön, I., Rossetti, G. and Martens, K.** (2009). Darwinulid Ostracods: Ancient Asexual
Scandals or Scandalous Gossip? In *Lost Sex: The Evolutionary Biology of*
*Parthenogenesis* (ed. Schön, I.), Martens, K.), and Dijk, P.), pp. 217–240. Dordrecht:
Springer Netherlands.

**Scimone, M. L., Kravarik, K. M., Lapan, S. W. and Reddien, P. W.** (2014). Neoblast
Specialization in Regeneration of the Planarian Schmidtea mediterranea. *Stem Cell*
*Reports* **3**, 339–352.

**Swanstrom, J., Chen, K., Castillo, K., Barraclough, T. G. and Fontaneto, D.** (2011). Testing for
evidence of inefficient selection in bdelloid rotifers: do sample size and habitat
differences matter? *Hydrobiologia* **662**, 19–25.

**Tan, T. C. J., Rahman, R., Jaber-Hijazi, F., Felix, D. A., Chen, C., Louis, E. J. and Aboobaker,**
**A.** (2012). Telomere maintenance and telomerase activity are differentially regulated in
asexual and sexual worms. *Proceedings of the National Academy of Sciences* **109**,
4209–4214.

**Tu, K. C., Cheng, L.-C., TK Vu, H., Lange, J. J., McKinney, S. A., Seidel, C. W. and Sánchez**
**Alvarado, A.** (2015). Egr-5 is a post-mitotic regulator of planarian epidermal
differentiation. *eLife* **4**, e10501.

**Tucker, A. E., Ackerman, M. S., Eads, B. D., Xu, S. and Lynch, M.** (2013). Population-genomic
insights into the evolutionary origin and fate of obligately asexual Daphnia pulex. *Proc*
*Natl Acad Sci U S A* **110**, 15740–15745.

**Van der Auwera, G. A., Carneiro, M. O., Hartl, C., Poplin, R., del Angel, G., Levy-Moonshine,**
**A., Jordan, T., Shakir, K., Roazen, D., Thibault, J., et al.** (2013). From FastQ Data to

High-Confidence Variant Calls: The Genome Analysis Toolkit Best Practices Pipeline.
*Current Protocols in Bioinformatics* **43**, 11.10.1-11.10.33.

**van Wolfswinkel, J. C., Wagner, D. E. and Reddien, P. W.** (2014). Single-cell analysis reveals
functionally distinct classes within the planarian stem cell compartment. *Cell Stem Cell*
**15**, 326–339.

**Vergara, D., Jokela, J., Lively, C. M. and Bronstein, E. J. L.** (2014). Infection Dynamics in
Coexisting Sexual and Asexual Host Populations: Support for the Red Queen
Hypothesis. *The American Naturalist* **184**, S22–S30.

**Vogg, M. C., Owlarn, S., Pérez Rico, Y. A., Xie, J., Suzuki, Y., Gentile, L., Wu, W. and**
**Bartscherer, K.** (2014). Stem cell-dependent formation of a functional anterior
regeneration pole in planarians requires Zic and Forkhead transcription factors.
*Developmental Biology* **390**, 136–148.

**Wagner, D. E., Wang, I. E. and Reddien, P. W.** (2011). Clonogenic Neoblasts Are Pluripotent
Adult Stem Cells That Underlie Planarian Regeneration. *Science* **332**, 811–816.

**Welch, D. B. M. and Meselson, M. S.** (2001). Rates of nucleotide substitution in sexual and
anciently asexual rotifers. *Proceedings of the National Academy of Sciences* **98**, 6720–
6724.

**Wilson, C. G., Nowell, R. W. and Barraclough, T. G.** (2018). Cross-Contamination Explains
"Inter and Intraspecific Horizontal Genetic Transfers" between Asexual Bdelloid
Rotifers. *Current Biology* **28**, 2436-2444.e14.

**Wilson, C. G., Pieszko, T., Nowell, R. W. and Barraclough, T. G.** (2024). Recombination in
bdelloid rotifer genomes: asexuality, transfer and stress. *Trends in Genetics* **40**, 422–
436.

**Zhu, S. J., Hallows, S. E., Currie, K. W., Xu, C. and Pearson, B. J.** (2015). A mex3 homolog is
required for differentiation during planarian stem cell lineage development. *eLife* **4**,
e07025.

770