Evaluation of In2Care® mosquito stations for suppression of the Australian backyard

mosquito, *Aedes notoscriptus*

Véronique Paris[1], Nicholas Bell[1], Thomas L. Schmidt[1], Nancy M. Endersby-Harshman[1], Ary A.

Hoffmann[1,2]

[1]School of BioSciences, Bio21 Institute, University of Melbourne, Parkville, Victoria 3010, Australia

[2]Department of Chemistry and Bioscience, Aalborg University, Aalborg 9220, Denmark

Keywords: *Aedes notoscriptus*, In2Care®, pyriproxyfen, *Beauveria bassiana*

ABSTRACT

Aedes notoscriptus is a container-breeding mosquito endemic to Australia that vectors

arboviruses and is suspected to transmit *Mycobacterium ulcerans*, the cause of Buruli ulcer.

Localised control measures for this species are limited by its high dispersal capability. Here we

evaluated the effectiveness of the In2Care® system in controlling *Ae. notoscriptus*. The In2Care®

system suppresses mosquito populations via the entomopathogenic fungus, *Beauveria*

bassiana, and the insect growth regulator pyriproxyfen, the latter of which is autodisseminated

among larval habitats by contaminated mosquitoes. A field trial was conducted using 110

In2Care® stations in a 50,000 m^2 target area and the results were compared to four control

areas that did not receive the In2Care® treatment. Efficacy was evaluated by comparing egg

counts and measuring larvicidal impact in surrounding breeding sites. Laboratory experiments

were conducted to validate the effect of *B. bassiana* on adult survival. Field results showed that

six weeks after the In2Care® stations were deployed, treatment site ovitraps contained 43%

fewer eggs than control site ovitraps, and 33% fewer eggs after ten weeks, indicating that the

In2Care® system was able to reduce the population of *Ae. notoscriptus*. Population reduction

remained evident for up to three weeks after In2Care® stations removal. Treatment site

ovitraps had significantly fewer *Ae. notoscriptus* eclosing than control site ovitraps, confirming

the pyriproxyfen autodissemination feature of the stations, and an average reduction of 50% in

adult eclosion was achieved. Adult mosquitoes had a significantly shorter lifespan after

exposure to *B. bassiana.* The strategy has potential for suppressing *Ae. notoscriptus*

populations.

1   INTRODUCTION

Mosquitoes of the genus *Aedes* continue to pose a significant threat to human and animal

health. While much research has been focused on the yellow fever mosquito (*Ae. aegypti*) and

the tiger mosquito (*Ae. albopictus*), other container-breeding and human-biting *Aedes* species

have received less attention, limiting information about effective control measures. One such

species is *Ae. notoscriptus*, a container-breeding mosquito that is broadly distributed in its

native range in Australia and has become invasive in other regions, including the Torres Strait

Islands, New Zealand, Papua New Guinea, New Caledonia, Indonesia (Dobrotworsky 1965, Lee

et al. 1987, Sunahara and Mogi 2004), and the United States (specifically California) (Metzger et

al. 2021). *Aedes notoscriptus* is a vector of human arboviruses (Doggett and Russell 1997), is

suspected to play a significant role in the transmission of *Mycobacterium ulcerans* (the

bacterium responsible for Buruli ulcer (BU)) (Wallace et al. 2017), and is also a primary vector of

dog heartworm (Russell 1985). *Aedes notoscriptus* has greater dispersal capacity than other

container-breeding *Aedes* (Watson et al. 2000, Trewin et al. 2019, Paris et al. 2023), making

localised control efforts more challenging. In a recent pilot study in Victoria, Australia, gravid

traps were applied to reduce the population size of *Ae. notoscriptus* on the Mornington

Peninsula, with the objective to further explore the species' role in the transmission of Buruli

ulcer (https://www.health.vic.gov.au/infectious-diseases/beating-buruli-in-victoria; Mee *et al.*

unpublished). However, no measurable reduction in the number of *Ae. notoscriptus* eggs was

observed, likely due to the species' capacity for long distance dispersal, which is evident from

genetic data (Paris et al. 2023). In addition, the species breeds in many cryptic breeding sites

(Trewin et al. 2019) in properties that cannot be easily accessed to apply larvicides.

One potential solution for controlling pests with high dispersal rates in an environment with an

abundance of cryptic breeding sites is through autodissemination of insecticides by the target

pest, a method that has been implemented in both agriculture (Ignoffo 1999) and public health

settings (Cook et al. 2009). An example of this approach is the In2Care® system, an outdoor

contamination station designed to reduce local mosquito populations and thereby reduce the

risk of disease transmission (https://www.in2care.org). The system lures gravid container

breeding mosquitoes into the station and contaminates them with the insect juvenile hormone

analogue and insect growth regulator, pyriproxyfen, which affects larvae and pupae (Unlu et al.

2020). Contaminated mosquitoes spread the insecticide to other breeding sites so that

mosquito larvae are not only controlled inside the station, but also in other (often cryptic)

breeding sites in the vicinity.

Pyriproxyfen is combined with the entomopathogenic fungus *Beauveria bassiana*.

Entomopathogenic fungi are necrotrophic parasites (i.e. kill their host in order to support their

own growth and complete their life cycle), potentially useful for biocontrol (Lacey et al. 2015).

Beauveria bassiana infects a broad range of hosts (Ferron 1978) such as ants (Broome et al.

1976), termites (Culliney and Grace 2000), agricultural pests such as grasshoppers (Bidochka

and Khachatourians 1991) and arthropod vectors of human diseases, including mosquitoes

(Howard et al. 2010, Bukhari et al. 2011). Infected mosquitoes usually die of the infection after

a few days, reducing the length of time available to transmit disease and removing them from

the breeding population. The efficacy of the In2Care® system has previously been validated by

experiments in some cases for *Ae. aegypti* and *Ae. albopictus* (Buckner et al. 2017, 2021, Khater

et al. 2022), but not for other *Aedes* species.

In this study, we aimed to evaluate the potential of the In2Care® system in controlling a

population of *Ae. notoscriptus* in Victoria, Australia. We conducted a field trial using 110

In2Care® stations in a designated area and compared temporal changes in mosquito numbers

to those in four control areas that did not receive the In2Care® station treatment. The efficacy

of the In2Care® system was evaluated by comparing egg counts between all sites, as well as by

measuring the larvicidal impact achieved in surrounding mosquito breeding sites due to

pyriproxyfen autodissemination. We also conducted laboratory experiments to validate the

effect of *B. bassiana* on adult mosquito survival. The results provide insights into the

effectiveness of the In2Care® system in suppressing populations of *Ae. notoscriptus* and suggest

a promising new approach for controlling this species in locations where it has become invasive.

2 MATERIAL AND METHODS

2.1 Field experiments

2.1.1 Permits

This work was performed under APVMA Small-scale Trial Permit PER7250. Trials conducted to generate data relating to efficacy, residues, crop or animal safety or other scientific information outside the confines of a research facility where the size of the trial annually does not exceed the following: a total of 5 hectares nationally, with a maximum of 1 hectare in any one jurisdiction in the case of any food and/or fibre field crop (https://permits.apvma.gov.au/PER7250.PDF). The active ingredient In2Mix® sachets were imported under biosecurity permit 0006570560 from the Australian Government Department of Agriculture, Fisheries and Forestry. Permit conditions meant that the trial was limited to one treatment site, but we had access to several control sites.

2.1.2 Study sites

We assigned one treatment site and four control sites for this study: Eynesbury (In2Care® treatment) (Figure 1), Darley (C1), Aintree (C2), Diggers Rest (C3) and Wyndham Vale (C4). All sites were located north-west of Melbourne in Victoria, Australia and comprised ~80 residential houses over a total ~50,000 m^2 area. Previous surveillance has shown that all sites have established populations of *Ae. notoscriptus*. All sites represented residential areas with 1-2 storey houses on allotments of 572.42 m^2 to 958.83 m^2.

2.1.3 Mosquito monitoring

Mosquito population density surveillance was conducted using egg counts at control and In2Care® treatment from the 20th of January 2022 until the 21st of April 2022. A total of 150 ovitraps (30 per site) were used, each consisting of a 500 mL black plastic bucket half-filled with water and containing alfalfa pellets to attract gravid *Ae. notoscriptus*. A 15 cm long strip of red felt extending into the water provided a consistent oviposition substrate. Egg counts were taken at weekly intervals for two weeks prior to the deployment of In2Care® stations. After the deployment of In2Care® stations, eggs were collected fortnightly, followed by weekly

collections for two weeks after the intervention concluded. The same researcher performed all

egg counts under a microscope to ensure consistency. To prevent the ovitraps from becoming

functional breeding sites, all larvae and pupae present in the ovitraps were discarded at each

collection.

2.1.4 In2Care® station deployment, monitoring, and servicing

We placed In2Care® stations in volunteer properties, as well as on nature strips and other

public land, with a density of approximately one station per 450 m^2 close to the manufacturer's

recommendation (one station per 400 m^2, https://www.In2Care.org). This resulted in a total

deployment of 110 stations as shown in Figure 1. The stations were strategically located in

shaded or semi-shaded areas that were expected to attract *Ae. notoscriptus*, such as areas with

vegetation or proximity to other breeding grounds. During the station deployment process, we

filled the water reservoir with approximately 4.7 litres of tap water before attaching the netting

containing the biocides to the floater and placing it onto the water surface. We aimed for a

homogeneous distribution of the biocides by shaking each In2Mix® sachet containing the

biocide powder, odour tablets, and netting well. The remaining biocide powder and odour

tablets were placed into the water reservoir before the lid was locked into place.

Figure 1 In2Care® station and ovitrap placement in the treatment site: Stations were placed at a density of one station per 450 m^2, resulting in 110 stations deployed, represented by the In2Care® station symbol. Ovitraps (n=30) were placed on public land to monitor egg number before, during and after the intervention, shown by the bucket symbol (mean distance of ovitraps to stations = ~158 m).

Regular monitoring of the In2Care® stations was conducted on a fortnightly basis. The monitoring included checking for signs of disturbance, ensuring that the water level was sufficient, (i.e., filled more than halfway) and that the netting was still dry and intact. The presence of larvae and live and/or dead pupae in the stations was also recorded in order to validate the station's attractiveness to gravid female mosquitoes and to ensure that no mosquitoes developed into adults inside the stations. When recording larval and pupal counts, we did not discriminate between *Ae. notoscriptus* and other species that might have been

141 present in the stations. After four weeks, the biocides and odour tablets in the stations were
142 refreshed with In2Mix® as previously described.

2.2   Laboratory experiments

2.2.1   Mosquito rearing

All laboratory experiments in this study used mosquitoes from laboratory colonies of
Ae. notoscriptus and *Ae. aegypti*. These colonies were established from field collections made in
Cairns, Australia in 2019 for *Ae. aegypti*, and Brisbane, Australia in 2014 for *Ae. notoscriptus*.
The colonies were maintained in a temperature-controlled insectary at 26°C ± 1°C, with a 12-
hour photoperiod, as per the protocol outlined by Ross et al. (2017) for rearing of *Ae. aegypti*.
The mosquitoes were provided with constant access to a 10% sucrose solution, and were
housed in 27 cm^3 BugDorm-1® cages (MegaView Science Co., Ltd., Taichung City, Taiwan). Adult
females were blood-fed on a human volunteer's arm, once per generation, to initiate egg-laying
(with ethics approval from The University of Melbourne 0723847). The colonies were
maintained in replicate cages, each containing approximately 500 mosquitoes.

2.2.2   Laboratory validation of pyriproxyfen autodissemination

To determine whether pyriproxyfen was being disseminated from the In2Care® stations to
other breeding sites, we collected fortnightly water samples from 15 of the ovitraps monitoring
the treatment site. The traps were selected randomly after excluding ovitraps that did not
contain any eggs or larvae which ensured selected ovitraps had been visited by mosquitoes. A
total of 200 mL of water per trap was collected in a sealable plastic container which was then
placed into an individual zip-lock bag to avoid cross-contamination between samples. We also
collected 200 mL of water from five randomly selected In2Care® stations to serve as positive
controls. Finally, five 200 mL water samples were taken from randomly chosen ovitraps from
each control site as negative controls (20 samples total).

To measure residual intervention effects after the In2Care® stations were removed from the
treatment site, we collected additional water samples from ovitraps in week 12 and 13 of the

trial. We collected ten water samples from In2Care® stations in week 10 to act as positive
controls for those additional samples.

All collected water samples were strained through fine mesh to remove any organisms or
organic material that might have been collected with the water. We used individual pieces of
mesh for each sample to avoid cross contamination between samples. We then added 25 L3
laboratory-reared *Ae. notoscriptus* larvae into each water sample and added two TetraMin®
tropical fish food tables (Tetra, Melle, Germany). The containers were placed into a 26°C
climate-controlled cabinet with a 12:12 light:dark cycle. All containers were checked daily to
ensure larvae had sufficient food, and any mosquito eclosing was recorded. We used the
number of free pupal exuviae as a proxy for successful adult eclosing.

2.2.3 Laboratory validation of *B. bassiana* efficacy against *Ae. notoscriptus*

To validate the adulticidal effect of *B. bassiana* on adult *Ae. notoscriptus* and *Ae. aegypti,* we
performed survival assays in the laboratory. All mosquitoes used in the experiment were adults
seven days post-eclosion and were allowed to mate. We exposed 25 females to a fresh In2Mix®
netting (containing approximately 0.2 g of In2Mix® powder with 74.03% pyriproxyfen and 10%
B. bassiana spores) using the WHO standard forced exposure bioassay (The World Health
Organisation 2006). Eight replicates of 25 females were prepared for the treatment and the
control groups. We provided each replicate group with 10% sucrose solution. Individuals that
died within one hour post experiment set up were excluded from all analyses. Sucrose solution
was refreshed as needed. We scored survival daily until all individuals in each treatment cohort
died. Individuals were scored as dead if no movement occurred after being manipulated with
forceps. To test whether mosquitoes were still able to blood feed and produce eggs post
infection, we blood fed groups on day two post exposure. We recorded the proportion of
mosquitoes that successfully blood fed for each group and collected and counted eggs on day
four and five of the experiment. We repeated the same setup with In2Mix® nets collected from
In2Care® stations that were deployed in the field for four weeks.

We calculated the germination rate of spores used to infect the treatment groups to validate
the spore quality and to ensure consistent spore qualities across replicates. Germination rate

was also used to compare the quality of fresh In2Mix® with that of four-week-old nettings. To

estimate germination rate, we inoculated three Sabouraud Dextrose Agar (SDA) plates for each

used netting by rubbing a sterile cotton tip on the netting and transferring the spore sample

onto the plates. The plates were then sealed with Parafilm® "M" Laboratory Film (Amcor

Flexibles North America) and stored at 26°C for 24 hours. After incubation we identified

germinated and not germinated spores (300 spores total) on three randomly chosen areas of

each plate using a light microscope. The germination rate was calculated using the following

formula: [number of germinated spores] ÷ [total number of spores counted (germinated + not-

germinated)] x 100. We then averaged the calculated germination rate across the three

replicate plates.

2.3   Statistical analyses

All statistical analyses were performed in R Studio v1.14 (R Core Team 2021). We used one

sample t-tests to determine whether the average numbers of eggs per trap in the treatment

site differed from the control sites.

Contingency tests were used to test if the number of weeks since station deployment

influenced the number of larvae (alive) and pupae (alive and dead) in the In2Care® stations.

We tested whether the water samples collected from the treatment site ovitraps, the control

sites ovitraps and the In2Care® stations differed in the percentage of mosquitoes that

successfully eclosed by using Kruskal-Wallis tests (a Kologorov-Smirnov test revealed that the

data were not normally distributed). We ran a Mantel test to determine whether the

percentage of eclosed mosquitoes was correlated with the distance from the nearest In2Care®

station.

To investigate whether mosquitoes infected with *B. bassiana* had a shorter lifespan than

control groups, whether there were differences in survival between mosquitoes tested on fresh

or older nettings, and whether there were differences in survival between *Ae. notoscriptus* and

Ae. aegypti, we performed a Cox regression survival analysis using the 'survival' package in R

(Therneau and T. Lumley 2015). The variables 'species', 'treatment' and 'netting' were used as

explanatory variables, while the 'replicate' was added as a random factor.

We ran independent sample t-tests to investigate differences in the proportion of mosquitoes

blood feeding between control groups and groups infected with *B. bassiana.*

3. RESULTS

3.1 Mosquito surveillance

The results of our mosquito population density surveillance, which was performed by counting

eggs in ovitraps, revealed that the number of eggs was not obviously linked to rainfall events,

but did gradually decrease as conditions became colder (Figure 2). The average number of *Ae.*

notoscriptus eggs was higher in the control sites than the average value in the treatment site at

weeks 8, 10, 12 and 13 of the trial (week 8 (week 6 post station deployment): t = 7.39, df = 3, p

= 0.003; week 10 (week 8 post station deployment): t = 7.03, df = 3, p = 0.003; week 12 (week

10 post station deployment): t = 8.4, df = 3, p = 0.002; week 13 (week 11 post station

deployment): t = 3.49, df = 3, p = 0.019).

Figure 2 Egg counts and weather data during the In2Care® station trial: (A) *Weekly mean*

minimum and maximum temperature are shown as red solid and blue double dashed lines

respectively. Mean cumulative rainfall is represented by the light blue area plot. Weather data

were obtained from the Bureau of Meteorology (bom.gov.au) Melbourne Airport Observation

Station. **(B)** *Weekly mean egg counts in the treatment site are shown by the solid red line.*

Dashed green, brown, purple, and blue lines represent egg counts in individual control sites. Egg

counts are averaged from n=30 ovitraps per site. Black error bars show standard errors per

 collection and site. The In2Care® station symbols indicate the timing of station set and removal.

The In2Care® floater symbol indicates maintenance of stations in week 6 of the trial (4 weeks

post-deployment).

Two weeks after the In2Care® stations were deployed, larvae were found in 53.64% (95% CI:

46.77% - 60.51%) of stations, indicating that most stations were visited. After four weeks

67.15% (95% CI: 58.22%, 76.32%) of stations were infested and infestation rates increased to

87.27% (95% CI: 80.91% - 91.28%) after week 6 since deployment, and 63.63% (95% CI: 53.60%

- 71.68%) after week 8 (Figure 3). Over the entire monitoring period, only two stations were

never visited. A contingency test showed that the weeks since initial deployment of the

In2Care® stations significantly influences the number of stations that were positive for larvae

(X^2 = 9.67, df = 3, p = 0.022).

Figure 3 Number of In2Care® stations that presented live larvae since In2Care® stations were deployed: *Number of larvae are represented by different shades of grey, with lighter shades indicating fewer larvae and darker shades more larvae. n = 110 In2Care® stations per week. Fresh water and In2Mix® were added at 4 weeks.*

The numbers of live and dead pupae changed significantly across time (Contingency test: alive: X^2 = 121.68, df = 3, P < 0.001; dead: X^2 = 177.82, df = 3, P < 0.001). No In2Care® stations showed any pupae after being in the field for two weeks. After four weeks since In2Care® stations were deployed, 15.45% (95% CI: 14.96% - 15.95) of stations had live pupae in them. This increased to 81.82% (95% CI: 75.0% - 88.6%) after six weeks and 50% (95% CI: 40.43% - 59.57%) after eight

weeks since deployment. Dead pupae were observed in In2Care® stations after six (48.18%

(95% CI: 38.4% - 57.9%.) and eight weeks (89.09% (95% CI: 85.04% - 93.14%) of deployment

(Figure 4).

Figure 4 Number of In2Care® stations that presented live pupae (A) and dead pupae (B) since

In2Care® stations were deployed: *Number of pupae are represented by different shades of*

grey, with lighter shades indicating fewer pupae and darker shades more pupae. n = 110

In2Care® stations per week.

3.2 Autodissemination

Significantly more mosquitoes successfully eclosed from pupae if water samples were collected

in the control sites when compared with the samples collected from ovitraps in the In2Care®

treatment site (Kruskal-Wallis tests, X^2 = -41.043, df = 2, p < 0.001) and from samples directly

taken from the In2Care® stations (X^2= 82.926, df = 2, p < 0.001). Eclosion of mosquitoes from

pupae in water samples collected from ovitraps from the treatment site were also significantly

higher compared to samples collected from In2Care® stations and showed also a significant

difference (X^2= 41.925, df = 2, p < 0.001) (Figure 5). This indicates that pyriproxyfen was

autodisseminated from the In2Care® stations to the ovitraps. Baseline adult eclosing rates in
the control site samples was on average 92%, compared to 42% in the intervention site.

283 *Figure 5 Percentage of eclosed mosquitoes from water samples collected during the In2Care®*

284 *field trial.* *Eclosed mosquitoes from all collections combined. The red box represents water*

285 *samples that were collected from ovitraps in the In2Care® treatment site (n=60). The green box*

286 *represents water samples collected from control sites (n=64) and the blue box shows water*

287 *samples collected from field In2Care® stations (n=20).*

Prolonged autodissemination effects were observed. Water samples collected from ovitraps

two weeks after the In2Care® stations were removed from the treatment site showed that

Ae. notoscriptus larvae had a significantly reduced percentage of eclosed adults compared to

samples collected from control sites (Kruskal Wallis test: X^2= 15.83, df = 2, p < 0.001). Positive

control samples (from In2Care® stations) showed 100% eclosion inhibition, which was

significantly higher than the average 55% adult emergence inhibition in the treatment site

ovitraps for week 12 and 13 combined (Kruskal Wallis test: X^2= 26.380, df = 2, p < 0.001) (Figure

6), which also reflects the larvicidal effect of the In2Care® stations.

We also tested whether the results differed between the week 12 and week 13 water

collections to determine how long the residual effect of pyriproxyfen lasted in ovitraps. We

found a significant difference in eclosed adult between the treatment site samples and samples

from control sites (Kruskal Wallis test: X^2 = 8.71, df = 2, p = 0.030) when collected two weeks

after the removal of the In2Care® stations (week 12). The reductions in adult emergence

inhibition were not significantly different in samples collected three weeks after station

removal (Kruskal Wallis test: X^2 = 7.92, df = 2, p = 0.260) although eclosion in controls was lower

at this time (Figure 6). Baseline adult eclosing rates in the control site samples was on average

87%, compared to 55% in the intervention site.

■

Figure 6 Percentage of eclosed mosquitoes from water samples collected after the In2Care®
field trial. (Above) Both collections combined for week 12 and 13 (2 and 3 weeks after removal
of In2Care® Stations from the treatment site). Red boxes represent water samples that were
collected from ovitraps in the In2Care® treatment site (n=30). Green boxes represent water

samples collected from control sites (n=40) and blue boxes show water samples collected as

positive control from field In2Care® stations in week 10 (n=10). Error bars indicate standard

error.

(Below) Collections two and three weeks after removal of In2Care® Stations from the treatment

site separately (In2Care® treatment site n=15; control sites n=20; In2care® stations n=5). Error

bars indicate standard error.

There was no correlation between the proximity of the ovitraps to the closest In2Care® station

and to all In2Care® stations (mean distance = ~158 m; longest distance = ~307 m; shortest

distance = ~4 m) and the percent of successful eclosing of mosquitoes from collected water

samples (Mantel test: r = -0.05, p = 0.983; Figure S1)

3.3 Efficacy of *B. bassiana* on *Ae. notoscriptus* and *Ae. aegypti*

The results of the Cox-regression analysis indicated that there was a significant effect of the

species tested (*Ae. notoscriptus* or *Ae. aegypti)* (z = -3.53, df = 3, p = 0.001), the age of the

In2Care® netting (fresh or 4-weeks-old) (z = -59.49, df = 3, p < 0.001), and the treatment

(infected or control) (z = 48.15, df = 3, p < 0.001) on survival. On average, survival of *Ae. aegypti*

was 9% lower compared to *Ae. notoscriptus* (HR = 0.91 (0.87 − 0.96), p < 0.001). Additionally,

both species had a four-times higher chance of dying if infected with In2Care® netting

compared to uninfected control groups (HR = 4 (3.8 - 4.1), p < 0.001). Furthermore, if either

species was infected using fresh In2Care® nettings, average survival was reduced by 88%

compared to older nettings (HR = 0.12 (0.11 − 0.13), p < 0.001) (see Figure 7).

Figure 7 Survival of adult Aedes notoscriptus **and** Aedes aegypti **after exposure to fresh**

In2Care® netting (above) and nettings that were in the field for four weeks (below). *Solid lines*

represent the survival of the treatment groups with Ae. notoscriptus *in magenta and* Ae. aegypti

in red. Dashed lines show the survival of control groups with Ae. notoscriptus *in blue and* Ae.

aegypti *in green. Data shows averaged proportion of survival from n = 8 replicates per group*

and species. Error bars indicate standard error.

We calculated an average *Beauveria* spore germination rate of 93.41% (95% CI: 91.18% - 97.79)

for the fresh In2Mix® samples with no replicate being lower than 90%. The four-week-old

nettings presented an average germination rate of 53.74% (95% CI: 49.92% - 57.71%).

Figure 8 Proportion of mosquitoes that blood feed after* B. bassiana *infection. (A) Fresh

In2Care® netting (B) 4-weeks-old In2Care® netting. *Blue boxes indicate the proportion that*

blood feed of the treatment groups, red boxes indicate the proportion that blood feed in the

control groups. Data is averaged from n = 8 replicates per group and species. Error bars show

standard error.

We did not find a significant difference in the proportion of females that blood fed, between

the infected and control groups (t-test: fresh In2Care® mix: *Ae. notoscriptus:* t = -0.03, df = 7, p

= 0.98; *Ae. aegypti:* t = -0.01, df = 7, p = 0.41; 4-weeks old In2Care® mix: *Ae. notoscriptus:* t =

0.88, df = 7, p = 0.79; *Ae. aegypti:* t = 0-.34, df = 7, p = 0.74) (Figure 8). We did not observe any

oviposition by any of the infected groups, while control groups laid eggs.

4. DISCUSSION

Our experiments demonstrated the efficacy of the In2Care® system for control of the container breeding mosquito *Ae. notoscriptus*. We found a reduction in egg numbers in the treatment site after In2Care® stations had been deployed for six weeks with the effect persisting until three weeks after station removal. The autodissemination results indicate that the In2Care® stations were visited by mosquitoes which then successfully contaminated surrounding ovitraps with pyriproxyfen and that the disseminated larvicide persisted at lethal levels for at least two weeks after the In2Care® stations were removed. Additionally, we found that the fungal pathogen *B. bassiana*, which is incorporated in the In2Care® system, significantly decreased the lifespan of mosquitoes in a laboratory setting.

The In2Care® stations were successful in attracting mosquitoes as shown by the high proportion of stations that housed larvae (Figure 3). After two weeks of deployment, more than half of the stations contained larvae, which continued to increase over following weeks, before dropping somewhat to 63% after eight weeks. This decrease may reflect a reduction in the local *Ae. notoscriptus* population, particularly given that traps had been serviced at the 4-week mark when attraction levels should have been reset. A lower number of new eggs laid in the stations, as well as continued development of existing larvae to the pupal stage, would also have contributed to a decrease in larval numbers. This is further supported by an increase in the number of pupae in the stations after four weeks (Figure 4). The reduction in live pupae is caused by pyriproxyfen-induced mortality, which was observed in about half of the stations at six weeks post-deployment and 87% of stations after eight weeks of deployment. Pupal mortality might have started earlier than detected as larvae in the stations might have fed on dead pupae. These findings also suggest that the In2Care® stations function as an 'egg dump' that contributes to the overall reduction of the mosquito population, as eggs laid in the stations do not mature into adult mosquitoes.

Mosquito population sizes naturally fluctuate with rainfall and temperature (Bomblies 2012), with rainfall triggering egg hatching and higher temperatures accelerating development. We found that the general trend of egg counts in all sites followed these weather parameters

(Figure 2). However, after six weeks of the In2Care® stations being in place, we observed a

significant reduction in egg counts in the treatment site (Figure 2). The decrease in egg numbers

at the treatment site cannot be attributed to weather factors, as all control sites displayed

increasing egg numbers, while the treatment site showed a decline (Figure 2).

Our experiments on water collected from ovitraps at the treatment site confirm the effective

autodissemination of pyriproxyfen. The results reveal a significant average decrease of about

50% in eclosing of adults in water samples from the treatment site compared to control sites

(Figure 5). This finding suggests that mosquitoes carried pyriproxyfen from the In2Care®

stations to ovitraps and likely to other surrounding breeding sites. Female *Aedes* mosquitoes

tend to distribute their eggs across multiple breeding sites instead of laying them in one place,

which helps reduce the risk of losing all offspring if the chosen breeding site proves to be

unfavourable (Colton et al. 2003). This strategy of 'skip oviposition' favours the

autodissemination of pyriproxyfen, as it increases the number of breeding sites that

mosquitoes use. The autodisseminated pyriproxyfen likely contributed to the decrease in adult

mosquitoes in the treatment site, with a cumulative effect over multiple generations.

Importantly, this positive impact persisted for two weeks even after the removal of the

In2Care® stations from the treatment site (Figure 6).

The potential for mosquitoes to distribute pyriproxyfen to other parts of the ecosystem where

it would influence other insect systems should be considered before widespread application of

the technology is implemented, as some lab studies have shown pyriproxyfen transfer by male

mosquitoes (artificially dosed with high quantities of pyriproxyfen) to bees during nectar

feeding (Kancharlapalli et al. 2021). In2Care® Stations, however, specifically target gravid

female *Aedes* mosquitoes that only disseminate pyriproxyfen to water bodies during egg-laying,

and these will be less likely to visit flowers for sugar feeding at that stage. Moreover,

pyriproxyfen has already been registered and approved globally (including Australia) for

widescale agricultural applications given its acceptable risk for non-target organisms and

pollinators. If studies confirm these effects are found to be minimal or less damaging than the

effects of broadscale mosquito control methods, then implementation of the In2Care® system

will be of benefit in reducing the vectoring capacity of *Ae. notoscriptus* and other container

breeding mosquitoes.

We validated the adulticide effect of *B. bassiana* on *Ae. notoscriptus* and compared them to

Ae. aegypti, using both fresh In2Mix® and nettings that had been in the field for four weeks.

Both species had a 75% higher chance of reduced survival if infected with In2Mix®-treated

netting compared to control groups, and using fresh In2Care® nettings significantly reduced the

average survival rate of both species by 88% compared to older nettings (Figure 7). These

results show that older nettings still infect and kill adult *Aedes* mosquitoes, though their

lifespan increases by approximately three to six days, which indicated that the nettings should

be refreshed after four to six weeks, as recommended by the manufacturer. Our laboratory

investigations suggest that *B. bassiana* used in the In2Care® technology has the potential to be

an effective tool in controlling mosquito populations and has most likely contributed to the

positive results in our field intervention.

Our results are consistent with studies investigating the effect of the In2Care® system on

Ae. aegypti, Ae. albopictus and *Culex quinquefasciatus* (Buckner *et al.* 2017; Buckner *et al.* 2021;

Khater *et al.* 2022; Su et al. 2021) and suggest that the In2Care® stations successfully kill larvae

of container breeding mosquitoes by spreading pyriproxyfen to surrounding breeding sites. The

In2Care® system, in conjunction with source reduction and other integrated vector

management tools, has potential to provide sustained vector control and reduce transmission

of *Aedes*-borne arboviruses, parasites, and mechanically transmitted bacteria. Benefits of the

In2Care® system to public health may extend beyond the reduction of mosquito vectors.

Broadscale insecticide application may be obviated in an In2Care®-treated area, thereby

reducing public exposure as well as off-target environmental impacts.

Though the results present a promising demonstration of the potential for the In2Care® system

for *Ae. notoscriptus* control, there may be some challenges in implementing the approach in

other areas. In our trial, the treatment and control sites were relatively isolated, reducing the

potential for migration of mosquitoes back into the treated area. Implementation of the

strategy in less isolated areas may require the treatment of larger areas. Our study took place

in a relatively new developed area, where most houses are occupied and there are few empty

blocks or construction sites, making it easier to place traps. We did identify a few mosquito

breeding sources (mostly bird baths or drip trays of pot plants), but the area is generally well

maintained, reducing the likely incidence of cryptic breeding sites. In areas with a low

occupation rate of houses, there may be an increased risk of overflowing gutters and containers

in backyards that can act as breeding sites. It is worth noting that in a study utilising the

In2Care® system conducted in Hawaii (Brisco et al. 2023), a failure to find a reduction in

mosquito numbers was partly attributed to numerous cryptic breeding sites in properties that

did not participate in the intervention (in addition to an inadequate number of In2Care®

stations being deployed), and source reduction measures may be required in some contexts.

5. CONCLUSION

The In2Care® system was found to be effective in controlling a population of the container

breeding mosquito *Ae. notoscriptus* in Victoria, Australia. The system demonstrated a reduction

in egg numbers in the treatment site after six weeks of deployment. The autodissemination

results indicate that the In2Care® stations were visited by mosquitoes which then successfully

contaminated ovitraps with pyriproxyfen, and the larvicidal effect persisted for at least three

weeks after the In2Care® stations were removed. The experiments validated the adulticide

effect of the fungal pathogen *B. bassiana* on *Ae. notoscriptus* and *Ae. aegypti* and suggest that

it has the potential to be an effective tool in controlling mosquito populations.

6. ACKNOWLEDGEMENTS

Authors thank the Melton Council, especially Lawrie Conole and residents for permission and

assistance. Jessica Home, Sophie Collier, Matthew Whitney, and Sarah Jaboor are thanked for

assistance with fieldwork and lab assays. We thank Tim Möhlmann and Marit Farenhorst of

In2Care Trading B.V., and Steve Broadbent of Ensystex for assistance in planning the

intervention, lab experiments, provision of In2Care® stations and In2Mix® refills used in this

study and for feedback on the manuscript.

7. FUNDING

The study was funded by the National Health and Medical Research Council Partnership Project grant 1196396 'Stopping Buruli ulcer in Victoria'. VP was financially supported by the Australian Government Research Training Program Scholarship.

8. REFERENCES

Bidochka, M. J., and G. G. Khachatourians. 1991. The implication of metabolic acids produced by *Beauveria bassiana* in pathogenesis of the migratory grasshopper, *Melanoplus sanguinipes*. J. Invertebr. Pathol. 58.

Bomblies, A. 2012. Modeling the role of rainfall patterns in seasonal malaria transmission. Clim. Change. 112.

Brisco, K., C. Jacobson, S. Seok, X. Wang, Y. Lee, O. Arkabi, and A. J. Cornel. 2023. Field Evaluation of In2Care mosquito traps to control *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) in Hawai'i Island. J. Med. Entomol. 1–9.

Broome, J. R., P. P. Sikorowski, and B. R. Norment. 1976. A mechanism of pathogenicity of *Beauveria bassiana* on larvae of the imported fire ant, *Solenopsis richteri*. J. Invertebr. Pathol. 28.

Buckner, E. A., K. F. Williams, A. L. Marsicano, M. D. Latham, and C. R. Lesser. 2017. Evaluating the vector control potential of the In2Care® mosquito trap against *Aedes aegypti* and *Aedes albopictus* under semifield conditions in Manatee County, Florida. J. Am. Mosq. Control Assoc. 33.

Buckner, E. A., K. F. Williams, S. Ramirez, C. Darrisaw, J. M. Carrillo, M. D. Latham, and C. R. Lesser. 2021. A field efficacy evaluation of In2Care mosquito traps in comparison with routine integrated vector management at reducing *Aedes aegypti*. J. Am. Mosq. Control Assoc. 37.

Bukhari, T., W. Takken, and C. J. M. Koenraadt. 2011. Development of *Metarhizium anisopliae*

and *Beauveria bassiana* formulations for control of malaria mosquito larvae. Parasites and

Vectors. 4.

Colton, Y. M., D. D. Chadee, and D. W. Severson. 2003. Natural skip oviposition of the

mosquito *Aedes aegypti* indicated by codominant genetic markers. Med. Vet. Entomol. 17.

Cook, A., B. Devine, C. Rodriguez, D. Roser, S. Khan, N. McGuinness, N. Ashbolt, and P.

Weinstein. 2009. Assessing the public health impacts of recycled water use. Interim report

1. Gov. West. Aust. Dep. Water, West. Aust. Premier's Water Found.

Culliney, T. W., and J. K. Grace. 2000. Prospects for the biological control of subterranean

termites (Isoptera: Rhinotermitidae), with special reference to *Coptotermes formosanus*.

Bull. Entomol. Res.

Dobrotworsky, N. V. 1965. The mosquitoes of Victoria (Diptera, Culicidae). Melbourne

University Press, Carlton, Australia.

Doggett, S. L., and R. C. Russell. 1997. *Aedes notoscriptus* can transmit inland and coastal

isolates of Ross River and Barmah Forest viruses from New South Wales. Arbovirus Res

Aust. 7: 79–81.

Ferron, P. 1978. Biological control of insect pests by entomogenous fungi. Annu. Rev. Entomol.

23.

Howard, A. F. V., R. N'Guessan, C. J. M. Koenraadt, A. Asidi, M. Farenhorst, M. Akogbéto, M.

B. Thomas, B. G. J. Knols, and W. Takken. 2010. The entomopathogenic fungus *Beauveria*

bassiana reduces instantaneous blood feeding in wild multi-insecticide-resistant *Culex*

quinquefasciatus mosquitoes in Benin, West Africa. Parasites and Vectors. 3.

Ignoffo, C. M. 1999. The first viral pesticide: Past, present, and future. J. Ind. Microbiol.

Biotechnol. 22.

Kancharlapalli, S. J., C. J. Crabtree, K. Surowiec, S. D. Longing, and C. L. Brelsfoard. 2021.

Indirect transfer of pyriproxyfen to European honeybees via an autodissemination

approach. PLoS Negl. Trop. Dis. 15.

Khater, E., D. Autry, M. Gaines, and R. Xue. 2022. Field evaluations of autocidal gravid ovitraps

and In2Care traps against *Aedes* mosquitoes in Saint Augustine, northeastern Florida. J.

Florida Mosq. Control Assoc. 69.

Lacey, L. A., D. Grzywacz, D. I. Shapiro-Ilan, R. Frutos, M. Brownbridge, and M. S. Goettel.

2015. Insect pathogens as biological control agents: Back to the future. J. Invertebr. Pathol.

132.

Lee, D. J., M. M. Hicks, M. Griffiths, M. L. Debenham, J. H. Bryan, R. C. Russel, and M. Geary.

1987. The Culicidae of the Australasian Region⊡: Genus *Anopheles* (Anopheles, Cellia),

Monogr. Ser. Entomol. Monogr. Australian Government Publishing Service.

Metzger, M. E., J. W. Wekesa, S. Kluh, K. K. Fujioka, R. Saviskas, A. Arugay, N. McConnell, K.

Nguyen, L. Krueger, G. M. Hacker, R. Hu, and V. L. Kramer. 2021. Detection and

establishment of *Aedes notoscriptus* (Diptera: Culicidae) mosquitoes in Southern

California, United States. J. Med. Entomol.

World Health Organisation. 2006. Guidelines for testing mosquito adulticides for Indoor

Residual Spraying and treatment of mosquito nets. Who/Cds/Ntd/Whopes/Gcdpp/2006.3.

Paris, V., R. V. Rane, P. T. Mee, S. E. Lynch, A. A. Hoffmann, and T. L. Schmidt. 2023. Urban

population structure and dispersal of an Australian mosquito (*Aedes notoscriptus*) involved

in disease transmission. Heredity. 130: 99–108.

R Core Team. 2021. R core team (2021). R A Lang. Environ. Stat. Comput. R Found. Stat.

Comput. Vienna, Austria. URL http//www. R-project. org.

Russell, R. C. 1985. Report of a field study on mosquito (Diptera: Culicidae) vectors of dog

heartworm, *Dirofilaria immitis* Leidy (Spirurida:Onchocercidae) near Sydney, N.S.W., and

the implications for veterinary and public health concern. Aust. J. Zool. 33.

Su, T., P. Mullens, J. Thieme, A. Melgoza, R. Real, and M. Q. Brown. 2021. Deployment and

fact analysis of the In2Care mosquito trap, a novel tool for controlling invasive *Aedes*

species. J. Am. Mosq. Control Assoc. 36.

Sunahara, T., and M. Mogi. 2004. Searching clusters of community composition along multiple

spatial scales: a case study on aquatic invertebrate communities in bamboo stumps in

West Timor. Popul. Ecol. 2004 462. 46: 149–158.

Therneau, T. M., and T. Lumley. 2015. Package ' survival .' R Top. Doc.

Trewin, B., D. E. Pagendam, J. M. Darbro, Q. Health, and G. J. Devine. 2019. Urban landscape

features influence the movement and distribution of the Australian container-inhabiting

mosquito vectors *Aedes aegypti* (Diptera: Culicidae) and *Aedes notoscriptus* (Diptera:

Culicidae). J. Med. Entomol. 57.2: 443-453.

Unlu, I., I. Rochlin, D. S. Suman, Y. Wang, K. Chandel, and R. Gaugler. 2020. Large-scale

operational pyriproxyfen autodissemination deployment to suppress the immature asian

tiger mosquito (Diptera: Culicidae) Populations. J. Med. Entomol. 57.

Wallace, J. R., K. M. Mangas, J. L. Porter, R. Marcsisin, S. J. Pidot, B. Howden, T. F. Omansen,

W. Zeng, J. K. Axford, P. D. R. Johnson, and T. P. Stinear. 2017. *Mycobacterium ulcerans*

low infectious dose and mechanical transmission support insect bites and puncturing

injuries in the spread of Buruli ulcer. PLoS Negl. Trop. Dis. 11: 1–16.

Watson, T. M., A. Saul, and B. H. Kay. 2000. *Aedes notoscriptus* (Diptera: Culicidae) survival and

dispersal estimated by mark-release-recapture in Brisbane, Queensland, Australia. J. Med.

Entomol. 37: 380–4.

561