Multiple dynamic interactions from basal ganglia direct and indirect pathways mediate action selection

Hao Li[1], and Xin Jin[1,2,3,4*]

[1] Molecular Neurobiology Laboratory, The Salk Institute for Biological Studies, 10010 North Torrey Pines Road, La Jolla, CA 92037, USA

[2] Center for Motor Control and Disease, Key Laboratory of Brain Functional Genomics, East China Normal University, 3663 North Zhongshan Road, Shanghai 200062, China

[3] NYU–ECNU Institute of Brain and Cognitive Science, New York University Shanghai, 3663 North Zhongshan Road, Shanghai 200062, China

[4] Lead Contact

*Correspondence to: xjin@bio.ecnu.edu.cn

Summary

     The basal ganglia are known to be essential for action selection. However, the functional role of basal ganglia direct and indirect pathways in action selection remains unresolved. Here by employing cell-type-specific neuronal recording and manipulation in mice trained in a choice task, we demonstrate that multiple dynamic interactions from the direct and indirect pathways control the action selection. While the direct pathway regulates the behavioral choice in a linear manner, the indirect pathway exerts a nonlinear inverted-U-shaped control over action selection, depending on the inputs and the network state. We propose a new center (direct) - surround (indirect) - context (indirect) "Triple-control" functional model of basal ganglia, which can replicate the physiological and behavioral experimental observations that cannot be simply explained by either the traditional "Go/No-go" or more recent "Co-activation" model. These findings have important implications on understanding the basal ganglia circuitry and action selection in health and disease.

In Brief

Using behavior analysis, in vivo electrophysiology, optogenetics and computational modeling in mice, Li and Jin unveiled the neuronal dynamics of basal ganglia direct and indirect pathways underlying action selection, and proposed a new "Triple-control" functional model of basal ganglia.

Highlights

• Outputs of opponent SNr subpopulations determine the action selection

• Striatal direct/indirect pathways exhibit distinct physiology and function during action selection

• Cell ablation and optogenetic inhibition of indirect pathway exert opposite behavioral effects

• A new "Triple-control" functional model of basal ganglia pathways is proposed

Introduction

Selecting the proper actions is essential for organism's survival and reproduction in the ever-changing environment (Gallistel, 1980). Numerous studies have implicated that the basal ganglia, a series of interconnected subcortical nuclei including the striatum and substantia nigra, play a primary role in action selection (Graybiel, 1998; Hikosaka et al., 1998; Jin and Costa, 2015; Mink, 2003; Redgrave et al., 1999). Indeed, a wide range of neurological and psychiatric disorders associated with the dysfunctional basal ganglia circuitry, including Parkinson's disease (Benecke et al., 1987), Huntington's disease (Phillips et al., 1995), Obsessive-compulsive disorder (OCD) (Graybiel and Rauch, 2000), are characterized by major deficits in action selection and movement control. Anatomically, commands for motor control are processed by basal ganglia through two major pathways, termed direct and indirect pathway, originating from striatal D1- and D2-expressing spiny projection neurons (D1-/D2-SPNs), respectively (Albin et al., 1989; DeLong, 1990). These two pathways collectively modulate substantia nigra pars reticulata (SNr) activity and the basal ganglia output, thus influence behavioral decisions. There are currently two major types of thinking on how the basal ganglia pathways work. An early classic theory has suggested that the basal ganglia direct and indirect pathways oppose each other to facilitate and inhibit action, respectively (the "Go/No-go" model) (Albin et al., 1989; DeLong, 1990; Kravitz et al., 2010). In contrast, a recent theory has proposed that direct pathway selects the desired action, while the indirect pathway inhibits other competing actions in order to highlight the targeted choice (the "Co-activation" model) (Cui et al., 2013; Hikosaka et al., 2000; Mink, 1996).

The two theories have essentially agreed upon the function of direct pathway being the positive driving force for initiating or facilitating the desired actions. Yet, the ideas about the indirect pathway function are largely controversial as either impeding the desired action in the "Go/No-go" model or inhibiting the competing actions in the "Co-activation" model. While the precise neuroanatomy on how the D2-SPNs control SNr through indirect pathway has yet to be mapped out at single-cell level to differentiate the two hypotheses, either theory has found its supports from behavioral and physiological observations. For instance, it has been found that stimulation of striatal direct and indirect pathways can bidirectionally regulates locomotion (Durieux et al., 2012; Kravitz et al., 2010), consistent with the traditional 'Go/No-go' model. On the other hand, in vivo electrophysiological and imaging experiments revealed that the striatal direct and indirect pathways are both activated during action initiation (Barbera et al., 2016; Cui et al., 2013; Geddes et al., 2018; Isomura et al., 2013; Jin et al., 2014; Klaus and Plenz, 2016; Markowitz et al., 2018; Nonomura et al., 2018), as the 'Co-activation' model predicted. Furthermore, physiological and optogenetic studies concerning complex behavior such as learned action sequences have further complicated the issue, and unveiled various neuronal subpopulations in both pathways are activated during the initiation, termination and switching of actions (Geddes et al., 2018; Jin and Costa, 2015; Jin et al., 2014; Tecuapetla et al., 2016). So far, how exactly the basal ganglia direct and indirect pathways work together to control action selection has been controversial and inconclusive, and the underlying circuit mechanism remains largely unclear (Calabresi et al., 2014).

Here we trained mice to perform an operant action selection task where they were required to select one out of two actions to achieve reward, based on self-monitored time intervals (Howard et al., 2017). By employing in vivo neuronal recording, we found that the net

output of two opponent SNr neuron populations is predictive of the behavioral choices. Through

identifying striatal pathway-specific neuronal activity with optogenetic tagging, we found that

there are neuronal subpopulations in either direct or indirect pathway who are activated during

selecting one action and suppressed during another. Optogenetic inhibition, as well as selective

ablation of direct pathway impairs action selection, and optogenetic excitation of direct pathway

enhances current choice, confirming a role of direct pathway in facilitating desired actions.

Furthermore, optogenetic inhibition of indirect pathway improves action selection and excitation

of indirect pathway impairs behavioral choices, as predicted by the 'Go/No-go' model. However,

selective ablation of indirect pathway impairs action selection, opposite from the behavioral

effect of optogenetic inhibition and at odds with the 'Go/No-go' model, but in consistent with the

prediction from the 'Co-activation' model. To resolve these contradictions, we propose a new

center (direct) - surround (indirect) - context (indirect) "Triple-control" functional model of basal

ganglia pathways, in which there are two interacting indirect pathway subcircuits exerting

opposite controls over the basal ganglia output. The new model can reproduce the neuronal and

behavioral experimental results that cannot be simply explained by either the "Go/No-go" or the

"Co-activation" model. Further systematic analyses from this new model suggested that the

direct and indirect pathways modulate behavioral outputs in a linear and nonlinear manner,

respectively. Notably, in the new 'Triple-control' model, the direct and indirect pathways can

work together to dynamically control action selection and operate in a manner similar to 'Go/No-

go' or 'Co-activation' model, depending on the activity level and the network state. These results

revise our current understanding on how the basal ganglia control actions, and have important

implications for a wide range of movement and psychiatric diseases where the dynamic balance

between the two pathways is compromised (Albin et al., 1989; Benecke et al., 1987; Calabresi et

al., 2014; DeLong, 1990; Graybiel and Rauch, 2000; Mink, 1996; Phillips et al., 1995).

Results

Opponent SNr activities underlie action selection

To address the role of basal ganglia in action selection, we trained mice in a recently

developed 2-8 s task in which they are required to choose the left versus right action based on

self-monitored time intervals (Howard et al., 2017). Specifically, mice were put into an operant

chamber with both left and right levers extended (Figure 1A, see Methods). For a given trial,

both levers retract at trial initiation, and after either 2 s or 8 s (50% for each, randomly

interleaved), both levers extend. The mouse has to judge the interval between lever retraction and

extension as 2 s vs. 8 s and make a corresponding action choice by pressing the left vs. right

lever, respectively (Figure 1A). The first lever press after lever extension was registered as the

mouse's choice. The correct choice leads to sucrose delivery (10 µl) as reward, and any lever

presses beyond the first press after lever extension yield no outcome. A new trial starts at lever

retraction again after a random inter-trial-interval (ITI, 30 s on average; Figure 1A). Across 14

consecutive days of training, mice (n = 10) significantly increased the correct rate of choice from

chance level to more than 90% (Figure 1B). In addition, the animals gradually shortened the

choice latency and demonstrated a strong preference toward the left lever due to its association

with the shorter waiting time (Figure S1A, B). As a result, during the longer-waiting 8 s trials the

mouse initially moved toward the left lever, then crossing the midpoint between left and right

levers at around 4 s, and stayed around the right lever afterwards (Howard et al., 2017) (Figure

1C; Supplemental Video 1). Note that, the mouse showed no stereotyped movement trajectories

during the incorrect trials (Figure 1C). This emerged stereotyped movement trajectory in the 8-s

trials thus provided us a unique opportunity for investigating the neural mechanisms underlying

the internally-driven, dynamic action selection process.

The substantia nigra pars reticulata (SNr) is one of the major output nuclei of basal

ganglia (Albin et al., 1989; DeLong, 1990; Hikosaka et al., 2000; Mink, 1996). To investigate

how the basal ganglia contribute to the dynamic process of action selection, we began by

recording the SNr neuronal activity in mice trained in the 2-8 s task (Figure 1D, Figure S1C, see

Methods). It was found that a large proportion (211/261, 80.8%; recorded from n = 9 mice) of

SNr neurons changed firing rate significantly during the correct 8-s trials as mice dynamically

shifted the internal action selection from the left to the right (Figure 1E). The Z-score of the task-

related neuronal firing rate, reflecting the firing activity changes related to baseline, was defined

as Firing Rate Index (FRI, see Methods). We focus on the data analyses in the 8-s trials since the

first 2-s of 8-s trials consists of the identical behavioral and neuronal profiles of the 2-s trials due

to the task design (Figure 1C, Figure S1D-F). The task-related SNr neurons were categorized

into four subtypes based on the dynamics of FRI in the correct 8-s trials: Type 1 - monotonic

decrease (Figure 1E, F, 102/211, 48.3%), Type 2 - monotonic increase (Figure 1E, G, 56/211,

26.5%), Type 3 - transient phasic increase (Figure 1E, H, 25/211, 11.9%) and Type 4 - transient

phasic decrease (Figure 1E, I, 28/211, 13.3%). These types of neuronal dynamics in SNr only

appeared in the correct but not the incorrect trials (Figure 1F-I), nor on the day 1 of task training

(Figure S1G-K), suggesting a tight correlation between the SNr neuronal dynamics and the

behavioral performance. Specifically, the Type 1 and Type 2, but not the Type 3 and Type 4

neurons, exhibit firing changes co-varying with the action selection and these two types together

consist in around 80% of all task-related SNr neuron population (Figure 1J).

It has been suggested that SNr suppresses movements through the inhibition of

downstream motor nuclei and releases action via disinhibition (Hikosaka et al., 2000; Mink,

1996). We thus ask whether the opponent neuronal dynamics in Type 1 and Type 2 SNr

subpopulations mediate the dynamic shift of choice, by suppressing the competing selection of

right vs. left action, respectively. Indeed, the SNr net output by subtracting Type 2 and Type 1

SNr neuronal dynamics (Figure 1K) is highly reminiscent of the animal's stereotyped movement

trajectory during choice (Figure 1C). To further determine the relationship between the SNr net

output and action selection, we tested the behavioral choice of the 2-8 s trained mice in a series

of non-rewarded probe trials with novel intervals of 2.5, 3.2, 4, 5 and 6.3 s (see Methods).

Consistent with what reported before (Howard et al., 2017), the probability of mice selecting the

action associated with the long duration (8 s) gradually increases along with the time intervals of

probe trials (Figure 1L). The resulting psychometric curve thus represents the animal's real-time

action selection process during the 8-s trials. Further comparison between the psychometric

curve and the SNr net output revealed a strong linear correlation (Figure 1M), indicating that the

SNr net output faithfully predicts momentary behavioral choice. Together, these results suggest

that mice can learn to dynamically shift their choice based on internally-monitored time, and the

opponent neuronal activities in SNr correlate with the action selection.

SNr neuronal dynamics reflect action selection but not simply time or value

In the 2-8 s task, the passage of time and expectation of reward both change

simultaneously with the animal's internal choice. One may argue that the Type 1 and Type 2

neuronal dynamics observed in SNr during the 8-s trials might reflect the passage of time or

value of expected reward rather than action selection. To differentiate these possibilities and

specify the functional role of SNr activity, we presented mice previously trained in the 2-8 s task

with random probe trials of 16-s interval (Figure 2A). In these 16-s probe trials which they have

never experienced before during training, the animals sometimes wait on the right side and press

the right lever, or shift back to the left side and press the left lever when the levers are extended

at 16 s (Figure 2B, C). This arbitrary choice situation in the 16-s probe trials thus provides a

special window to determine the functional relationship between SNr activity and behavioral

choice. If the Type 1 and Type 2 SNr subpopulations encode information about time passage or

expectation value, their neuronal activities would continue changing monotonically between 8

and 16 s. In contrast, if the Type 1 and Type 2 SNr subpopulations encode action selection, their

neuronal activities would predict the behavioral choice and differentiate between the right vs. left

action selection. Indeed, it was found that when the firing activity of Type 1 SNr neurons

maintained below baseline from 8 to 16 s, the mice tended to select the right lever later (Figure

2D). However, when the firing activity reversed the decreasing tendency to increase, the mice

chose the left lever instead (Figure 2D). A similar relationship between the neuronal activity and

behavior choice was also evident in Type 2 SNr neurons, albeit with opposite dynamics (Figure

2E). This is especially evident in the subtraction of Type 2 and Type 1 SNr neuronal dynamics,

in which the SNr net output is strongly correlated with and predictive of behavioral choice

(Figure 2F). These results thus suggested that the neuronal activities in SNr likely encode the

ongoing action selection but not simply reflect time passage or reward value.

To further confirm this point, we recorded the firing activity from the same SNr neurons

during both the 2-8 s control task (Figure 2G, 2s-left and 8s-right) and a modified version of 2-8

s task in which the contingency between action and interval is reversed (Figure 2J, 2s-right and

8s-left) on the same day (see Methods). It was found that the mice performed at around 80%

correct in both tasks on the same day (Figure 2H, K, Figure S2A). Accordingly, the movement

trajectories of the same mice in 8-s trials were reversed from left-then-right in the control task

(Figure 2I) to right-then-left in the reversed task (Figure 2L). The left-lever preference during the

ITI in the control task was also switched to right-lever preference in the reversed 2-8 s task

(Figure S2B). Notably, the passage of time and expected value as well as other environmental

factors are all identical in both versions of task, except that the animal's choice is now reversed

from right to left for the 8-s trials (Figure 2H, I vs. Figure 2K, L). If Type 1 or Type 2 SNr

neurons encode time or value, either neuronal population will exhibit the same neuronal

dynamics in 8-s trials for both versions of task. On the other hand, if Type 1 and Type 2 SNr

neurons encode action selection, their neuronal dynamics will reverse in the reversed version of

2-8 s task compared to the standard version. In fact, the Type 1 SNr neurons which showed

monotonic decreasing dynamics in the control 2-8 s task (Figure 2M) reversed their neuronal

dynamics to a monotonic increase in the reversed 2-8 s task (Figure 2P), consistent with the

behavioral choice. The same reversal of neuronal dynamics was also observed in Type 2 SNr

neurons in the reversed version of standard task (Figure 2N, Q). The SNr net output by

subtracting Type 2 and Type 1 SNr neuronal dynamics, which was tightly correlated with the

action selection in the standard 2-8 s task (Figure 2O), is reversed and now predictive of the new

behavioral choice in the reversed 2-8 s task (Figure 2R). Notably, Type 3 and Type 4 SNr

neurons exhibiting transient change when mice switching between choices maintained the same

neuronal dynamics in both tasks (Figure S2C-F). Together, these results therefore demonstrate

that the output of basal ganglia reflects the dynamic action selection rather than simply time or

value.

Distinct striatal direct vs. indirect pathway activity during action selection

The basal ganglia output is largely controlled by two major neural pathways, called 'direct' and 'indirect' pathway, originating from D1- vs. D2-expressing spiny projection neurons (D1- vs. D2-SPNs) in the striatum, respectively (Albin et al., 1989; DeLong, 1990; Hikosaka et al., 2000; Mink, 1996). We then decided to determine the neuronal dynamics in the striatum, specifically the neuronal activity in the direct and indirect pathways during action selection. We employed in vivo extracellular electrophysiology to record the neuronal activity in the dorsal striatum when mice perform the 2-8 s task, and classified putative SPNs based on the spike waveforms and firing properties (Geddes et al., 2018; Jin and Costa, 2010; Jin et al., 2014). Among all the SPNs recorded from the trained mice (n = 24), 341 out of 409 SPNs (83.4%) were defined as task-related neurons for showing significant firing changes during the 2-s and 8-s lever retraction period (Figure 3A, Figure S3A). Similar to the various types of neuronal dynamics observed in SNr, task-related SPNs showed Type 1 (Figure 3B, monotonic decrease, 159/341, 46.6%), Type 2 (Figure 3C, monotonic increase, 103/341, 30.2%), Type 3 (Figure 3D, transient phasic increase, 49/341, 14.4%) and Type 4 (Figure 3E, transient phasic decrease, 30/341, 8.8%) activity profiles during the correct 8-s trials (Figure 3A, Figure S3A). These neural dynamics were largely absent in SPNs on day 1 of training (Figure S3B-F). These results indicate that the striatum, as one of the major input nuclei of basal ganglia, demonstrates the four types of neuronal dynamics similar with SNr during the dynamic process of action selection.

To further determine the neuronal activity in the direct and indirect pathways during action selection, we utilized an optogenetics-aided photo-tagging method (Geddes et al., 2018; Howard et al., 2017; Jin and Costa, 2010; Jin et al., 2014; Lima et al., 2009) to record and identify striatal D1- vs. D2-SPNs in freely behaving mice. Channelrhodopsin-2 (ChR2) was

selectively expressed in D1- or D2-SPNs by injecting AAV-FLEX-ChR2 in the dorsal striatum

of D1- and A2a-Cre mice, respectively (Geddes et al., 2018; Jin et al., 2014). In the end of each

behavioral session with recording, optogenetic stimulation via an optic fiber attached to the

electrode array was delivered to identify D1- vs. D2-SPNs through photo-tagging (Figure 3F,

Figure S3G-J) (Geddes et al., 2018; Jin et al., 2014). Only those neurons exhibiting a very short

latency ($\leq$ 6 ms) to light stimulation (Figure 3G-I) and showing identical spike waveforms (R $\geq$

0.95, Pearson correlation coefficient) between behavior and light-evoked response (Figure 3J, K)

were identified as Cre-positive thus D1- or D2-SPNs (Geddes et al., 2018; Jin et al., 2014).

Within all positively identified D1-SPNs (n = 92 from 6 mice) and D2-SPNs (n = 95 from 6

mice), 74 out of 92 (80.4%) D1-SPNs and 79 out of 95 (83.1%) D2-SPNs showed a significant

change in firing rate during the correct 8-s trials. In addition, all four types of neuronal dynamics

during action selection were found in both D1-SPNs (Figure 3L, M) and D2-SPNs (Figure 3N,

O), as observed in SNr. The Type 1 and Type 2 neuronal dynamics showing monotonic firing

change (Figure 3M, O) were the predominant task-related subpopulations within either D1-

(Figure 3L) or D2-SPNs (Figure 3N). Notably, the striatal D1-SPNs consist of significantly more

Type 1 than Type 2 neurons (Figure 3L), while D2-SPNs show a similar proportion between the

two Types (Figure 3N). These data thus suggest while neurons in both the striatal direct and

indirect pathways encode information related to behavioral choice, the two pathways might

reflect and contribute to distinct aspects of action selection.

Ablation of striatal direct vs. indirect pathway differently impaired action selection

Given the action-selection-related neuronal dynamics observed in striatum, we next asked

whether the neural activity in striatum is necessary for learning and execution of action selection,

and furthermore, what is the functional difference between the direct and indirect pathways. It

has been reported that the NMDA receptors on striatal SPNs are critical for sequence learning

(Geddes et al., 2018; Jin and Costa, 2010) and action selection (Howard et al., 2017). To further

identify the functional role of NMDA receptors on D1- vs. D2-SPNs for action selection, we

employed a genetic strategy to specifically delete NMDA receptors from D1- vs. D2-SPNs by

crossing mice carrying a floxed NMDAR1 (NR1) allele with a dorsal striatum-dominant D1-cre

line (Gong et al., 2007) and A2a-cre line (Geddes et al., 2018; Jin et al., 2014), respectively

(referred to as D1-NR1 KO and D2-NR1 KO mice, respectively; see Methods). Both the D1-

NR1 KO and D2-NR1 KO mice are significantly impaired in learning the 2-8 s task compared to

their littermate controls (Figure 4A, B), suggesting that NMDA receptors on either D1- or D2-

SPNs are critical for learning of proper action selection. In the end of two-week training, when

given the probe trials with various intervals across 2 to 8 s, it was found that D1-NR1 KO mice

showed a systematic bias toward the lever associated with short interval and made deficient

behavioral choice only in long interval trials (Figure 4C). In contrast, D2-NR1 KO mice showed

impaired action selection across various probe trials of both short and long intervals (Figure 4D).

These data suggest that while NMDA receptors on both D1- and D2-SPNs are required for action

learning, the deletion of NMDA receptors in direct and indirect pathways impairs action

selection in a different manner.

We then asked whether that neural activity in dorsal striatum is necessary for the proper

execution of action selection after learning. We first conduct striatal inactivation in trained

wildtype mice by bilateral intra-striatal infusion of muscimol (Figure 4E, see Methods). Striatal

muscimol infusion significantly reduced the animal's overall performance in comparison with

the pre- and post-saline injection controls (Figure 4F). When tested with probe trials, the

psychometric curve indicated that the inactivation of striatum impairs action selection for the

long trials (Figure 4G). These data thus suggested that the striatal neural activity is critical for

appropriate execution of learned action selection.

To further elucidate the functional role of specific striatal pathways in action selection,

we next employed a viral approach to bilaterally express diphtheria toxin receptors (AAV-

FLEX-DTR-eGFP) in the dorsal striatum of trained D1- and A2a-Cre mice, followed by

diphtheria toxin (DT) injections to selectively ablate D1- or D2-SPNs (Geddes et al., 2018)

(Figure 4H, I, N; see Methods). Ablation of either D1- or D2-SPNs significantly impaired action

selection and reduced the correct rate of choice (Figure 4J, O). Notably, the psychometric curve

revealed that D1-SPNs ablation mice showed a selective impairment of choice in long interval

trials (Figure 4K). In contrast, mice with D2-SPNs ablation exhibited choice deficits in both long

and short trials (Figure 4P). Consistent with the D1- and A2a-NR1 KO data, these results suggest

that the direct and indirect pathways are both needed yet play distinct roles in action selection.

The classic 'Go/No-go' model of basal ganglia suggests the direct and indirect pathways

work antagonistically to release and inhibit action, respectively (Albin et al., 1989; DeLong,

1990; Kravitz et al., 2010). On the other hand, more recent 'Co-activation' model of basal

ganglia proposes that direct pathway initiates the selected action and at the same time, the

indirect pathway inhibits the competing actions (Cui et al., 2013; Hikosaka et al., 2000; Mink,

1996). For visualization purpose, we diagram 'Go/No-go' and 'Co-activation' models as center-

surround receptive field with D1-SPNs as the center and D2-SPNs as the surround (Figure 4S;

Figure S4A, D). While the direct pathway plays the similar role in both models (Figure S4B, E),

the function of indirect pathway differs dramatically (Figure 4S). Lesion of the indirect pathway

thus leads to contrast predictions on action selection from the two models (Figure S4C, F).

Specifically, ablation of D2-SPNs would facilitate the action being selected through removing

inhibition according to the Go/No-go model (Figure S4C) (Albin et al., 1989; DeLong, 1990;

Kravitz et al., 2010), while blockage of indirect pathway would impair the action selection due to

disinhibition of competing actions according to the Co-activation model (Figure S4F) (Cui et al.,

2013; Hikosaka et al., 2000; Mink, 1996). Although our D1-SPNs ablation experiment indicates

that direct pathway is required for action selection as suggested in both models (Figure 4J, K),

the D2-SPNs ablation result favorably supports the Co-activation model over the Go/No-go

model (Figure 4O, P, Figure S4F). In fact, close inspection of the movement trajectories of D1-

SPNs lesioned mice in the 8-s trials showed that compared to control mice (Figure 4L), they tend

to stick on the left side more often with impaired right choice when lever extension at 8s (Figure

4M). In contrast, D2-SPNs lesioned mice demonstrated overall rather random movement

trajectories, and the stereotyped left-then-right movement sequences were largely disrupted in

comparison with the controls (Figure 4Q, R). These observations are mostly consistent with the

idea of indirect pathway inhibiting competing actions in the Co-activation model (Figure 4S) and

lesion of indirect pathway disrupts action selection for both the short and long trials (Figure 4P-

R). Together, these data suggest that ablation of direct and indirect pathways both impair choice

behavior but in a distinct manner due to their different roles in action selection.

Optogenetic manipulation of D1- vs. D2-SPNs distinctly regulates action selection

To further determine the specific function of direct vs. indirect pathway in action

selection, we employed optogenetics to alter the D1- and D2-SPNs activity in vivo with high

temporal precision and investigated its effects on the ongoing action selection process. Both the

classic 'Go/No-go' (Albin et al., 1989; DeLong, 1990; Kravitz et al., 2010) and more recent 'Co-

activation' (Cui et al., 2013; Hikosaka et al., 2000; Mink, 1996) models predict that activation of

the direct pathway enhances the action selection (Figure S5A, E, I, K, O, Q), while inhibition of

direct pathway reduces the correct choice (Figure S5B, F, L, R). To experimentally validate the

models' predictions, AAV-FLEX-ChR2 was injected into the dorsal striatum of D1- or A2a-Cre

mice and optic fibers were implanted bilaterally for in vivo optogenetic stimulation (Figure 5A,

Figure S3K, L; see Methods) (Geddes et al., 2018; Jin et al., 2014). After mice learned the 2-8s

task, 1-s pulse of constant light (wave length 473 nm) was delivered right before lever extension

in randomly chosen 50% of 2-s and 50% of 8-s trials (Figure 5B, C, see Methods). The correct

rate of optogenetic stimulation trials is compared with the non-stimulation trials of the same

animal as a within-subject design. We observed no significant change on the correct rate in 2-s

trials, whereas the correct rate was significant increased by optogenetic stimulation in 8-s trials

(Figure 5D), indicating a facilitation effect on action selection by stimulating the D1-SPNs. We

then sought to determine the effect of suppressing D1-SPN activity on action selection by viral

expression of Halorhodopsin (AAV5-EF1a-DIO-eNpHR3.0-eYFP) in the dorsal striatum of D1-

cre mice (Gradinaru et al., 2010). As expected, inhibiting D1-SPNs right before lever extension

in trained mice reduced the correct rates in 8-s but not 2-s trials (Figure 5C, E), opposite to D1-

SPN stimulation effects. These experimental data with bidirectional optogenetic manipulation

suggest that the D1-SPN activity is positively correlated with the choice performance, in

consistent with the hypothesis of direct pathway facilitating the action selected in both the

Go/No-go and Co-activation models (Figure S5K, L, Q, R).

Nevertheless, the two models have distinct views on the function of indirect pathway.

While the classic 'Go/No-go' model suggests that the indirect pathway inhibits the selected

action (Albin et al., 1989; DeLong, 1990; Kravitz et al., 2010), the 'Co-activation' model

hypothesizes that the indirect pathway inhibits the competing actions instead (Cui et al., 2013;

Hikosaka et al., 2000; Mink, 1996). These models thus provide contrast predictions about the

effect of activation of the indirect pathway on action selection, being decreased correct rate

based on the Go/No-go model (Figure S5M) and increased correct rate from the Co-activation

model (Figure S5S), respectively. We thus decided to test the distinct predictions from the two

models by optogenetic manipulation of indirect pathway during action selection in the 2-8s task.

ChR2 or Halorhodopsin (eNpHR3.0) was expressed in the dorsal striatum of A2a-cre mice for

bilaterally optogenetic activation or inhibition during behavior (Figure 5A, Figure S3L; see

Methods). Notably, optogenetic excitation of D2-SPNs for 1s right before lever extension

decreased the correct rate in both 2-s and 8-s trials (Figure 5F). In contrast, transient optogenetic

inhibition of D2-SPNs before behavioral choice increased correct rates for both 2-s and 8-s trials

(Figure 5G). These data suggest that opposite to the D1-SPN manipulation, optogenetic

stimulation of D2-SPNs impairs action selection, while inhibition of D2-SPNs facilitates

behavioral choice. These optogenetic results further unveil the distinct roles of direct vs. indirect

pathway in action selection, and are in line with the predictions from the Go/No-go (Figure 5H,

Figure S5M, N) but not the Co-activation model (Figure S5S, T).

A 'Triple-control' model of basal ganglia circuit for action selection

Our DT lesion experiments found that ablation of indirect pathway impairs action

selection (Figure 4O, P), as predicted from the Co-activation but not Go/No-go model (Figure

4S), while the optogenetic results suggested that inhibition of D2-SPNs enhances behavioral

choice (Figure 5G), a result in favor of the Go/No-go rather than Co-activation model (Figure

5H). We wonder whether these seemly discrepant effects are attributed to a more complex circuit

mechanism involving in the indirect pathway different from either the Go/No-go or Co-

activation model. To systematically investigate the cell type- and pathway-specific mechanisms

underlying action selection, we firstly add Go/No-go and Co-activation models together to

examine the whether the resulted combination model could explain the experimental

observations (Figure S4G). The lesion of D1-SPNs in the combination model indeed selectively

impaired choice in long interval trials (Figure S4H). However, the effect of D2-SPNs ablation in

the combination model was neutralized due to the opposing contributions from Go/No-go and

Co-activation models respectively (Figure S4I). Based on these simulation results, none of the

Go/No-go, Co-activation and combination models was able to fully capture the underlying

mechanism of basal ganglia in action selection. Inspired by the data in current experiments, we

decided to build a new computational model of the cortico-basal ganglia circuitry based on the

realistic neuroanatomy (Aoki et al., 2019; Mailly et al., 2003; Schmidt and Berke, 2017; Taverna

et al., 2008) and empirical neuronal physiology during action selection (Figures 1-3).

Different from the dual control of action by direct vs. indirect pathway in either the

Go/No-go or Co-activation model (Figure S4), our new model adds an additional layer of control

derived from the indirect pathway, thus called 'Triple-control' model for action selection.

Specifically, the new model consists of one direct pathway and two indirect pathways defined as

D2-SPN #1 and D2-SPN #2 two subpopulations, corresponding to the Co-activation and Go/No-

go functional modules, respectively (Figure 6A, B). In addition, the indirect pathway D2-SPNs

in the Co-activation module inhibits the indirect pathway D2-SPNs in the Go/No-go module

through the well-known D2-SPN collaterals with the properties of short-term depression in the

striatum (Gustafson et al., 2006; Schmidt and Berke, 2017; Taverna et al., 2008; Tecuapetla et al.,

2007) (Figure 6A; see Methods), providing asymmetric modulation to D2-SPN subgroups and

promoting Co-activation module as the dominant functional module at rest. In this 'Triple-

control' basal ganglia model, striatal D1- and D2-SPNs associated with left and right actions

receive excitatory inputs from corresponding cortical inputs (Figure 6A) to generate Type 1 and

Type 2 neuronal dynamics (Figure S6A-D) (Lo and Wang, 2006). The D1- and D2-SPNs then

regulate the SNr neuronal dynamics through the direct and indirect pathways, respectively

(Figure S6) (Albin et al., 1989; DeLong, 1990; Hikosaka et al., 2000; Mink, 1996). The net SNr

output (Figure S6F, I), which controls the downstream brainstem and thalamic circuits necessary

for action selection (Hikosaka, 2007; Lo and Wang, 2006; Redgrave et al., 1999), will determine

the final behavioral choice (Figure S6G, J). Our computational simulations showed that this

'Triple-control' network model could faithfully recapitulate the neuronal activity across the basal

ganglia circuitry and predict the behavioral choice (Figure S6).

To dissect the functional role of direct vs. indirect pathway in action selection, we

simulate the cell ablation experiments and examine the behavioral output in the 'Triple-control'

basal ganglia model. Ablation of D1-SPNs in the network model (Figure S6E) modulates both

Type 1 and Type 2 SNr dynamics but in different magnitude due to the biased striatal input to

SNr left output and mutual inhibition between SNr left vs. right outputs (Figure S6F; see

Methods). As a result, the lesion causes a downward shift in the net SNr output, especially

evident at the late section of 8 s (Figure S6G). This change in net SNr output predicts a

behavioral bias towards left choice as seen in the psychometric curve (Figure 6C), in consistent

with experimental results in mice with D1-SPNs ablation (Figure 4K). In contrast, ablation of

D2-SPNs in the network model (Figure S6H), by removing the indirect pathways of both the

Go/No-go and Co-activation modules, alters Type 1 and Type 2 SNr dynamics (Figure S6I) and

change the net SNr output dramatically around 2s as well as 8s (Figure S6J). The model thus

predicts behavioral choice deficits for both short and long trials during D2-SPNs ablation (Figure

6D), consistent with experimental observations (Figure 4P). Together, these data suggest that our

new 'Triple-control' basal ganglia model, based on realistic neuroanatomy and empirical

neuronal physiology, can perform action selection similar to the behavior of mice, and

successfully replicate the pathway-specific lesion effects on choice.

We further simulate the neuronal and behavioral effects of optogenetic manipulation of

D1- and D2-SPNs in the cortico-basal ganglia model. Consistent with the experimental results

(Figure 5D, F), optogenetic stimulation of D1-SPNs facilitates the ongoing choice (Figure 6E),

while optogenetic inhibition of D1-SPNs suppresses ongoing choice in the model (Figure 6G). In

addition, optogenetic stimulation of D2-SPNs impairs the ongoing choice and causes switching

(Figure 6F), while optogenetic inhibition of D2-SPNs facilitates ongoing choice, due to the now

dominant Go/No-go module mediated by the short-term depression of D2 collaterals in the

model (Figure 6H). Consistent with the experimental observations, the optogenetic inhibition

effect is opposite from the D2-SPNs cell ablation in the model (Figure 6D).

We next investigate how the striatum influence SNr outputs. Since the collateral

projection with STD in D2-SPNs is the key in our 'Triple-control' model to switch between

Go/No-go and Co-activation modules, we first built a motif of indirect pathway with two D2-

SPNs subgroups defined as D2-SPN #1 and D2-SPN #2 (Figure S7A). We tested this indirect

pathway motif with monotonic neural dynamics observed in experiments meanwhile simulating

the optogenetic activation at 1s and 7s (Figure S7D-I). The SNr therefore received more

activation at 1s than at 7s (Figure S7J, K), suggesting that the D2-SPNs with short-term

depression in collateral inhibition modulates SNr activities in a firing rate-dependent manner.

We next sought to test the model's predictions and experimentally investigate the

distinctions in modulating SNr activities between the direct and indirect pathways during action

selection. In order to manipulate D1- or D2-SPNs and monitor SNr responses at the same time,

we simultaneously implanted optogenetic fibers and recording array into striatum and SNr

respectively on a single mouse (Figure S7L). While the mice performing the 2-8s task,

optogenetic stimulation was delivered to activate either D1- or D2-SPNs. It was found that

optogenetic activation of D1- or D2-SPNs caused both inhibition and excitation in Type 1 and

Type 2 SNr neurons (Figure S7M-O). To further compare SNr activities responding to striatal

activation at different time points during the lever retraction period, for a given trial, we

activated D1-SPNs (or D2-SPNs) either at 1s or 7s after the lever retraction (Figure S7P-R). For

direct pathway, the change of FRI in SNr activities caused by activation of D1-SPNs showed no

significant difference between 1s and 7s (Figure S7P, S). For indirect pathway, activating D2-

SPNs at 1s caused smaller activation of FRI than at 7s in Type 1 SNr neurons (Figure S7Q),

whereas for Type 2 SNr neurons, activating D2-SPNs at 1s induced bigger FRI increase at 1s

than at 7s (Figure S7R). Overall, activating D2-SPNs tended to bias the firing rate downward at

1s but upward at 7s in Type 1 SNr neurons, which was counteractive to the decreasing tendency

of Type 1 SNr neuron (Figure S7T). In contrast, Type 2 SNr neurons showed higher FRI increase

and smaller decrease in response to activating D2-SPNs at 1s than at 7s, which was opposing to

the increasing dynamics of Type 2 SNr neurons (Figure S7T). This firing rate-dependent

modulation on SNr activities through indirect pathway is consistent with the computational

simulation (Figure S7J, K; Figure S8). Therefore, the underlying D2-SPNs collaterals might

indeed be a key mechanism contributing to the modulation of SNr activity and action selection in

vivo, as simulated in the 'Triple-control' model.

Taken together, our new 'Triple-control' basal ganglia model, based on realistic

neuroanatomy and empirical neurophysiology, successfully reproduces both the lesion and

optogenetic data we collected during the animal experiments. It could thus potentially provide

essential insights into the circuit mechanism of basal ganglia underlying action selection.

Linear and nonlinear control of action selection by direct vs. indirect pathway

To gain an overall picture of how basal ganglia control action selection, we run through

the model with a wide continuous range of manipulation to mimic the effects from lesion to

optogenetic inhibition and optogenetic activation (Figure 7A, B, E, F). The simulations of cell

ablation and bidirectional optogenetic manipulations of D1-SPNs activity in the model reveal no

significant effects at 2-s trials (Figure 7C), but a linear relationship between the neuronal activity

in direct pathway and the behavioral performance of choice in 8 s trials (Figure 7D), as observed

in animal experiments. It thus further confirms that direct pathway selects action and facilitates

ongoing choice, consistent with the predictions from both the classic Go/No-go and recent Co-

activation models (Figure 7I, Figure S9A-D) (Albin et al., 1989; DeLong, 1990; Hikosaka et al.,

2000; Mink, 1996).

In contrast, manipulations of D2-SPNs activity from cell ablation to optogenetic

inhibition and then optogenetic stimulation in the model demonstrate an inverted-U-shaped

nonlinear relationship between the neuronal activity in indirect pathway and action selection, for

both 2-s and 8-s trials (Figure 7G, H). Detailed analyses reveal that D2-SPNs ablation removes

both the Co-activation and Go/No-go module in the indirect pathway and leaves SNr activity

dictated by D1-SPN inputs. However, due to the inhibition from Co-activation to Go/No-go

module in the indirect pathway via D2-SPN collaterals and short-term plasticity of these

synapses (Figure S7A-C; see Methods), optogenetic manipulation of D2-SPNs differentially

affects the D2-SPN subpopulations groups and promotes Go/No-go module to dominate the

basal ganglia network (Figure S7C-S). This dynamic switch of dominance between Co-activation

and Go/No-go modules on the basal ganglia network gives rise to a nonlinear relationship

between D2-SPNs manipulation and the behavioral outcome (Figure 7J).

Note that when the same inputs were applied to the Go/No-go or Co-activation model

alone, the behavioral performance in either model exhibits linear negative (Figure S9E) or

positive correlation (Figure S9F) with D2-SPNs activity, respectively. Both our experimental and

modeling results thus indicate that different from either the Go/No-go or Co-activation model,

the indirect pathway regulates action selection in a nonlinear manner, depending on the state of

the network and D2-SPNs activity level. Besides collaterals within D2-SPNs, other collateral

connections, for example connections between D1-SPNs or connections between D1- and D2-

SPNs, could also contribute to the regulation of action selection (Taverna et al., 2008). We tested

our 'Triple-control' model with adding additional collateral connections as D1→D1 (Figure

S10A-C), D1→D2 (Figure S10D-F) and D2→D1 (Figure S10G-I), respectively. It was found

while these additional collaterals further quantitatively regulate action selection, the general

principle of linear vs. nonlinear modulation of action selection by direct and indirect pathways

still qualitatively hold (Figure S10). Interestingly, our current 'Triple-control' model can also

replicate the behavioral effects of optogenetic manipulation of nigrostriatal dopamine on

behavioral choice (Howard et al., 2017), and further unveils an inverted-U-shaped relationship

between striatal dopamine concentration change and action selection (Figure S11). Together,

these results suggest that there are multiple levels of interactions from D1- and D2-SPNs to

dynamically control SNr output, and the basal ganglia direct and indirect pathways distinctly

control action selection in a linear and nonlinear manner, respectively.

Discussion

Here, by using an internally-driven 2-8s action selection task in mice, we investigated the

function of basal ganglia direct and indirect pathways in mediating dynamic action selection. We

found that the neuronal activities in SNr, the major output of basal ganglia, directly reflect animals'

internal action selection process, other than simply time or value. It was also observed that the

striatum, the main input of basal ganglia, shares the similar action selection-related neuronal

dynamics with SNr and is needed for both learning and execution of proper action selection.

Furthermore, the striatal direct and indirect pathways exhibit distinct neuronal activity and during

manipulation, they have different functional effects on controlling action selection. Notably, the

experimental observations on the physiology and function of direct and indirect pathways cannot be

simply explained by either the traditional 'Go/No-go' model or the more recent 'Co-activation'

model. We proposed a new 'Triple-control' functional model of basal ganglia, suggesting a critical

role of dynamic interactions between different neuronal subpopulations within the indirect pathway

for controlling basal ganglia output and behavior. In the model, a 'center (direct pathway) – surround

(indirect pathway) – context (indirect pathway)' three layers of structure exerts dynamic control of

action selection, depending on the input level and network state. This new model respects the realistic

neuroanatomy, and can recapitulate and explain the essential in vivo electrophysiological and

behavioral findings. It also provides a new perspective on understanding many behavioral phenomena

involving in dopamine and basal ganglia circuitry in health and disease.

Our current 2-8s action selection task offers a unique opportunity to observe the animal's

internal switch from one choice to another and monitor the underlying neuronal dynamics

correspondingly. We observed two major types of monotonically-changing SNr neuronal dynamics

during the internal choice switching, presumably one type associated with selecting one action and

another with selecting the competing action, respectively. The classic view on SNr activity is that it

tonically inhibits the downstream motor nuclei and releases action via disinhibition (Albin et al.,

1989; DeLong, 1990; Hikosaka and Wurtz, 1983; Wurtz and Hikosaka, 1986). The increased

response in SNr, however, could potentially inhibit the competing actions or the movements toward

the opposing direction through projections to the contralateral brain regions like superior colliculus

(Jiang et al., 2003). Here, we found that two subpopulations in SNr showed opposite monotonic

firing change during the left-then-right choice, and notably, their neuronal dynamics switched when

the animals performed the reversed version of task which requires a right-then-left choice. It thus

suggests that these SNr neurons are indeed associated with different action options during choice

behavior, and actively adjust their firing rates to facilitate respective action selection. Given the

opposite neuronal dynamics and functionally antagonistic nature of these two groups of SNr neurons,

we defined the net output of basal ganglia by the subtracting the neuronal activity between the two

SNr subpopulations and correlated it with the behavioral choice. We found that the basal ganglia net

output exhibited a tight correlation with the psychometric curve of behavioral choice, and faithfully

represented a neural basis for the dynamic action selection process.

As one of the major input nuclei of basal ganglia, striatum influences SNr activity

through direct/indirect pathways and undisputedly, plays an essential role in action selection

(Ding and Gold, 2012; Geddes et al., 2018; Jin et al., 2014; Lauwereyns et al., 2002; Tai et al.,

2012). By genetic manipulation and pharmacological inactivation, we showed that striatum is

indispensable for both learning action selection and the proper performance of learned behavioral

choice. The recording of striatal neuronal activity during action selection further revealed that

striatal spiny projection neurons share the similar types of neuronal dynamics as SNr. Through

optogenetic-tagging in freely behaving mice, we further found that striatal SPNs in the direct and

indirect pathways show distinct activity profile, with D1-SPNs representing a strong bias

towards the preferred choice, while D2-SPNs encoding two choices equally.

Two prevailing models have been proposed to explain the functional distinction between

D1- and D2-SPNs. The canonical model of the basal ganglia suggests that the D1- and D2-SPNs

play antagonistic roles in controlling action as mediating "Go" and "No-go" signals, respectively

(Albin et al., 1989; DeLong, 1990; Kravitz et al., 2010). A more recent model, however, implies

that as D1-SPNs initiate an action, D2-SPNs co-activate with D1-SPNs to inhibit other

competing actions (Cui et al., 2013; Hikosaka et al., 2000; Isomura et al., 2013; Jin et al., 2014;

Mink, 1996). Essentially, these two models agree upon the functional role of D1-SPNs in

releasing or facilitating the desired action, but contradict on the function of D2-SPNs on which

targeted action of inhibiting. Here, our in vivo recording data indicate that both D1- and D2-

SPNs share similar neuronal dynamics during action selection, and the neural activity alone is

not sufficient to separate and determine whether "Go/No-go" or "Co-activation" model is

supported (Cui et al., 2013; Isomura et al., 2013; Jin et al., 2014). To resolve the functional

distinction of the direct vs. indirect pathway, we applied a series of cell-type-specific

manipulations on striatal D1- and D2-SPNs during action selection behavior. First, we generated

mutant mice in which NMDA receptors are deleted from either striatal D1- or D2-SPNs (Geddes

et al., 2018; Jin et al., 2014). Both the D1-NR1 KO and D2-NR1 KO mice showed learning

deficits and behavioral choice impairments when tested with probe trials, suggesting that both

D1- and D2-SPNs are necessary for learning and performing action selection. Notably, while the

D1-NR1 KO mice are mostly impaired in the choice associated with 8s, a less-preferred option

compared to 2s, the D2-NR1 KO mice are compromised in both 2s and 8s choice. Additional

experiments with cell-type specific ablation further confirmed these results, consistent the

distinct neuronal activity profile in these two pathways revealed during in vivo neuronal

recording. While both the "Go/No-go" and "Co-activation" models predict the suppression of

D1-SPNs activity leads to impaired action selection, supported by current KO and cell-ablation

data, the manipulation experiments on D2-SPNs favor the "Co-activation" but not the "Go/No-

go" model which the latter suggests D2-SPNs ablation would improve rather than impair action

selection.

Next, we directly introduced transient bidirectional manipulations to D1- and D2-SPNs

activity by optogenetics while mice performing the task. Our findings revealed that activation or

inhibition of D1-SPNs increased and decreased the correct rate of choice respectively, suggesting

a facilitating role of direct pathway in action selection, which again fits well with the "Go/No-go"

as well as the "Co-activation" model. In contrast, optogenetic activation of D2-SPNs decreased

the correct rate of choice, while inhibition of D2-SPNs promoted the correct choice. These

results were supportive to the "Go/No-go" model but contradicted to the prediction of "Co-

activation" theory, which the latter predicts that activation of D2-SPNs inhibits competing

actions to facilitate desired choice, whereas inhibition of D2-SPNs releases competing actions

and compromises the ongoing choice.

In summary, neither "Go/No-go" nor "Co-activation" models could fully explain the

experimental results we found, particularly for experiments on D2-SPNs in the indirect pathway.

Through computer simulation, we further demonstrated that a simple additive combination of

"Go/No-go" nor "Co-activation" models by linear addition cannot reproduce all the experimental

observations. To resolve these theoretical difficulties, we proposed a new center-surround-

context "Triple-control" model of basal ganglia pathways for action selection. Specifically, two

subpopulations of D2-SPNs in the indirect pathway function as "Co-activation" and "Go/No-go"

modules respectively, and an activity-dependent inhibition from "Co-activation" to "Go/No-go"

module mediates the dynamic switch between the dominant module depending on the inputs and

network state. Due to the dominant "Co-activation" module in the default state, excessive

inhibition of D2-SPNs or ablating the entire indirect pathway eliminates the promotive

contribution and impairs action selection in the "Triple-control" model, consistent with the

experimental observations. In contrast, transient increase of D2-SPNs firing activity during

optogenetic stimulation introduces shift toward the "Go/No-go" dominance from the "Co-

activation" module via firing-rate-dependent short-term depression of the inhibitory synapses

between them, which amplifies the "No-go" signal and compromises action selection as

experimentally found. In contrast, transient decrease of D2-SPNs firing activity during

optogenetic inhibition results in disinhibition of "Go/No-go" module from inhibitory control of

"Co-activation" module, with an attenuated "No-go" signal which leads to better performance in

choice. These results from our new "Triple-control" model thus suggest that the basal ganglia

circuitry could be much more dynamic than previously thought, and it could employ a complex

mechanism of functional module reconfiguration for context- or state-dependent flexible control.

In the "Triple-control" model, we posited the collateral connections among striatal D2-

SPNs and its short-term plasticity could serve as an operational mechanism for the dominant

module switching. However, besides these well-known striatal local connections as one of the

simplest possible mechanisms, other anatomical circuits within basal ganglia circuitry could

potentially fulfill this functional role alone or additionally as well. For example, striatal D2-SPNs

project to external global pallidus (GPe) through striatopallidal pathway, and meanwhile they

receive arkypallidal projections from GPe to both the striatal SPNs and interneurons (Abdi et al.,

2015; Fujiyama et al., 2016; Mallet et al., 2016). It is thus also possible that the dynamic

interaction between "Co-activation" and "Go/No-go" modules is mediated through di-synaptic or

tri-synaptic modulation with GPe and/or striatal interneurons involved. Furthermore, in theory

this dynamic interaction between "Co-activation" and "Go/No-go" modules can also occur

outside striatum in the downstream nuclei including GPe and SNr, given their specific neuronal

subpopulations receiving inputs from corresponding striatal D2-SPNs subgroups and proper

collateral connections within the nuclei (Atherton et al., 2013; Cazorla et al., 2014; Fujiyama et

al., 2011; Lee et al., 2020; Wu et al., 2000). Considering the crucial role of dopamine in basal

ganglia circuitry, the new "Triple-control" model can also reproduce our previous experiments

results on the effect of nigrostriatal dopamine on action selection (Howard et al., 2017).

Importantly, it unveils that there is an inverted-U-shaped relationship between dopamine

concentration change and action selection (Cools and D'Esposito, 2011). The model simulation

suggests while moderate dopamine increase improves decision making, too much dopamine

changes, either increase or decrease, dramatically impairs the choice behavior. These results

might be able to explain some of behavioral observations involving in obscure decision making

under the influence of addictive substances.

Our findings also have important implications in many neurological and psychiatric

diseases. It was known that the loss of dopamine leads to hyperactivity of D2-SPNs and

disruption of local D2-SPNs collaterals in Parkinson's disease (Taverna et al., 2008; Wei and

Wang, 2016). These alterations will not only break the balance of direct vs. indirect pathway, but

also disrupt the multiple dynamic controls from the indirect pathway. The action selection will

thus be largely problematic, even with L-DOPA treatment, which might restore the dopamine

partially but not necessarily the altered basal ganglia circuitry and its circuit dynamics (Bastide et

al., 2015). The current "Triple-control" model also provides some mechanistic insights into the

inhibitory control deficits observed Schizophrenia (Taverna et al., 2008; Wei and Wang, 2016).

For instance, an increase in the density and occupancy of the striatal D2 receptors (D2R) has

been frequently reported in schizophrenia patients (Abi-Dargham et al., 2000; Howes and Kapur,

2009; Laruelle et al., 1997; Wong et al., 1986). Many antipsychotic medications primarily aim to

block the D2R (la Fougere et al., 2005; Lally and MacCabe, 2015; Yokoi et al., 2002), but the

drug dose is the key to the treatment and severer adverse effects are associated with overdose of

D2R antagonism (Levine and Ruha, 2012). In addition, prolonged exposure to antipsychotics

often causes extrapyramidal symptoms, including Parkinsonian symptoms and tardive dyskinesia

(Jarskog et al., 2007; Seeman, 2002). The dose-dependent effects when modulating D2R were

also found in cognitive functions such as serial discrimination, in which relatively low and high

dose of D2R agonist in striatum impair the performance in the discrimination task, while the

intermediate dose of D2R agonist produces significant improvement (Cools and D'Esposito,

2011; Goldman-Rakic et al., 2000; Horst et al., 2019; Mattay et al., 2003). These observations

thus further underscore the dynamic interplays and complexity of basal ganglia pathways in

action control, as demonstrated in current study and the new triple-control functional model.

Acknowledgments: The authors would like to thank Tom Jessell and members of Jin lab for

discussion and comments on the manuscript. This research was supported by grants from the

NIH (R01NS083815), the Dystonia Medical Research Foundation and the McKnight Memory

and Cognitive Disorders Award to X.J.

Author Contributions: X.J. conceived the project. X.J. and H.L. designed the experiments. H.L.

performed the experiments and analyzed the data. X.J. supervises all aspects of the work. H.L.

and X.J. wrote the manuscript.

Conflicts of Interest: None of the authors declare any conflict of interest, financial or otherwise.

Figure Legends:

Figure 1. The neuronal dynamics in SNr during the 2-8 s action selection task. (A)

Schematic diagram for the design of 2-8 s task. (B) Correct rate for wild type mice across 14

days training (n = 10 mice, one-way repeated-measures ANOVA, significant effect of training

days, $F_{13,117} = 32.54$, $p < 0.0001$). (C) Movement trajectory of an example mouse in correct

(left panel) and incorrect (right panel) 8-s trials (gray line: trajectory of each trials; red/black line:

the average trajectory). (D) Diagram of electrode array implanted into substantia nigra pars

reticulata (SNr). (E) Firing Rate Index (FRI) of neuronal activity for all task-related SNr neurons

in correct 8-s trials. The magnitude of FRI is color coded and the SNr neurons are classified as

four different types based on the activity dynamics. (F-I) Averaged FRI for Type 1 (F), Type 2

(G), Type 3 (H), Type 4 (I) of SNr neurons in correct (red) and incorrect 8-s trials (gray). (J) The

proportion of four types of SNr neurons. Type 1 and Type 2 are major types and significantly

more than Type 3 and Type 4 (Z-test, $p < 0.05$). (K) Integrated SNr output defined as the

subtraction of averaged FRI between Type 1 and Type 2 SNr neurons. (L) Averaged

psychometric curve (n = 10 mice) of choice behavior. (M) The correlation between the Type 1

and Type 2 FRI subtraction and the behavioral choice (R = 0.98, $p < 0.0005$). Error bars denote

s.e.m., same for below unless stated otherwise.

Figure 2. SNr neuronal dynamics reflect action selection but not interval time or reward

value. (A) Task diagram of 2-8 s control task with 10% 16-s probe trials. (B) Percentage of

behavioral choice in 2-s, 8-s and 16-s trials (blue: left choice; red: right choice) (n = 9 mice,

paired t-test, $p < 0.05$). (C) Movement trajectory of an example mouse in 16-s trials (blue: left

choice; red: right choice). (D) Averaged SNr Type 1 FRI in 16-s trials (red: left choice; black:

right choice). Firing rates from 8s to 16s (highlighted area) are compared between left and right

choice (n = 26 neurons, two-way repeated-measures ANOVA, significant difference between left

and right choices, F1,25 = 6.646, p = 0.016). (E) Averaged SNr Type 2 FRI in 16-s trials (red:

left choice; black: right choice). Firing rates from 8s to 16s are compared between left and right

choice (n = 16 neurons, two-way repeated-measures ANOVA, significant difference between left

and right choices, F1,15 = 5.785, p = 0.029). (F) Subtraction of FRI for SNr Type 1 and Type 2

neurons in 16-s probe trials (red: left choice; black: right choice). (G) Task design of 2-8 s

standard task. (H) Percentage of behavioral choice in 2-s and 8-s trials (blue: left choice; red:

right choice) (n = 6 mice, paired t-test, p < 0.05). (I) Movement trajectory of an example mouse

in 8-s trials (blue: left choice; red: right choice). (J) Task design of reversed 2-8 s task. (K)

Percentage of behavioral choice in 2-s and 8-s trials in the reversed 2-8 s task (blue: left choice;

red: right choice) (n = 6 mice, paired t-test, p < 0.05). (L) Movement trajectory of the same

mouse as (I) in 8-s trials in the reversed 2-8 s task (blue: left choice; red: right choice). (M)

Averaged FRI of the SNr Type 1 neurons in correct 8-s trials (n = 14 neurons). (N) Averaged

FRI of the SNr Type 2 neurons in correct 8-s trials (n = 11 neurons). (O) Integrated SNr output

as the subtraction of FRI for SNr Type 1 (M) and Type 2 neurons (N) in the standard 2-8 s task.

(P) Averaged FRI of the same neurons as (M) in correct 8-s trials of the reversed 2-8 s task. (Q)

Averaged FRI of the same neurons as (N) in correct 8-s trials of the reversed 2-8 s task. (R)

Integrated SNr output as the subtraction of FRI for SNr Type 1 (P) and Type 2 neurons (Q) in the

reversed 2-8 s task.

Figure 3. Neuronal activity of striatal D1- and D2-SPNs during action selection. (A) FRI of

neuronal activity for all task-related SPNs in correct 8-s trials. SPNs were classified as Type 1 - 4.

(B-E) Averaged FRI for Type 1 (B), Type 2 (C), Type 3 (D), Type 4 (E) of SPNs in correct (red)

and incorrect 8-s trials (gray). (F) Diagram of simultaneous neuronal recording and optogenetic

identification of D1- vs. D2-SPNs in dorsal striatum. (G) Top panel: Raster plot for a

representative D1-SPN response to 100 ms optogenetic stimulation. Each row represents one

trial and each black dot represents a spike. Bottom panel: Peristimulus time histogram (PETH)

aligned to light onset at time zero. (H) PETH for the same neuron as shown in (G) with a finer

time scale. (I) Distribution of light response latencies for D1- and D2-SPNs. (J) Action potential

waveforms of the same neuron in (G) for spontaneous (black) and light-evoked (orange) spikes

($R = 0.998$, $P < 0.0001$, Pearson's correlation). (K) Principal component analysis (PCA) of

action potential waveforms showing the overlapped clusters of spontaneous (black) and light-

evoked (orange) spikes. (L) Proportion of D1-SPN subtypes. Type 1 neurons are significantly

more than other three types of neurons in D1-SPNs (Z-test, $p < 0.05$). (M) Averaged FRI for

Type 1 (blue) and Type 2 (red) D1-SPNs in correct 8-s trials. (N) Proportion of D2-SPN

subtypes. (O) Averaged FRI for Type 1 (blue) and Type 2 (red) D2-SPNs in correct 8-s trials.

**Figure 4. Selective genetic knockout and ablation of D1- or D2-SPNs distinctly alters action
selection.** (A) Correct rate of control (n = 11 mice) and D1-NR1 KO mice (n = 16) in 2-8 s task

during 14 days training (two-way repeated-measures ANOVA, significant difference between

control and KO mice, $F_{1,25} = 10.8$, $p = 0.003$). (B) Correct rate of control (n = 17) and D2-NR1

KO mice (n = 10) in 2-8 s task during 14 days training (two-way repeated-measures ANOVA,

significant difference between control and KO mice, $F_{1,25} = 8.728$, $p = 0.007$). (C) The

psychometric curve for control (n = 11) and D1-NR1 KO mice (n = 16) (two-way repeated-

measures ANOVA, significant difference between control and KO mice, $F_{1,25} = 12.27$, $p =$

0.002). (D) The psychometric curve for control (n = 17) and D2-NR1 KO mice (n = 10) (two-way repeated-measures ANOVA, significant difference between control and KO mice, $F_{1,25}$ = 9.64, p = 0.005). (E) Schematic of muscimol infusion into the dorsal striatum in trained mice. (F) Correct rate for control (black: pre-muscimol, gray: post-muscimol) and mice with muscimol infusion (magenta) in dorsal striatum (n = 9 mice, paired t-test, p < 0.01). (G) The psychometric curve for control (n = 9 mice, black: pre-muscimol, gray: post-muscimol control) and mice with muscimol infusion (n = 9 mice, magenta) in dorsal striatum (two-way repeated-measures ANOVA, significant difference between control and muscimol infusion, $F_{2,16}$ = 11.74, p = 0.0007). (H) Timeline for selective diphtheria toxin (DT) ablation experiments. (I) Schematic of diphtheria toxin receptor (DTR) virus (AAV-FLEX-DTR-GFP) injection in dorsal striatum of D1-Cre mice. (J) Correct rate for control (n = 9 mice) and mice with dorsal striatum D1-SPNs ablation (D1-DTR, n = 8 mice) (two-sample t-test, p = 0.0016). (K) The psychometric curve for control (n = 9 mice) and D1-SPNs ablation mice (n = 8 mice) (two-way repeated-measures ANOVA, main effect of ablation, $F_{1,15}$ = 1.84, p = 0.195; interaction between trial intervals and ablation, $F_{6,90}$ = 4.14, p = 0.001). (L) Movement trajectory of a control mouse in 8-s trials. (M) Movement trajectory of a D1-DTR mouse in 8-s trials. (N) Schematic of diphtheria toxin receptor (DTR) virus (AAV-FLEX-DTR-GFP) injection in dorsal striatum of A2a-Cre mice. (O) Correct rate for control (n = 8 mice) and mice with dorsal striatum D2-SPNs ablation (D2-DTR, n = 8 mice) (two-sample t-test, p = 0.005). (P) The psychometric curve for control (n = 9 mice) and D2-SPNs ablation mice (n = 8 mice) (two-way repeated-measures ANOVA, main effect of ablation, $F_{1,15}$ = 0.477, p = 0.5; interaction between trial intervals and ablation, $F_{6,90}$ = 12.6, p < 0.001). (Q) Movement trajectory of a control mouse in 8-s trials. (R) Movement trajectory of a D2-DTR mouse in 8-s trials. (S) Schematic of center-surround receptive field diagram for

Go/No-Go (left) and Co-activation (right) models. '+' indicates facilitating effect to selection. '-'

indicates inhibitory effect to selection.

Figure 5. Optogenetic manipulation of D1- vs. D2-SPNs differently regulates action

selection. (A) Schematic of optic fiber implantation for experimentally optogenetic excitation or

inhibition of D1- or D2-SPNs in the dorsal striatum. (B, C) Schematic for optogenetic excitation

(B) and inhibition (C) of D1-/D2-SPNs for 1 s right before lever extension in 2-8 s task. (D)

Change of correct rate for optogenetic excitation of D1-SPNs in 2-s and 8-s trials (n = 11 mice,

one-sample t-test, 2-s trials: p = 0.248; 8-s trials: p < 0.05). (E) Change of correct rate for

optogenetic inhibition of D1-SPNs in 2-s and 8-s trials (n = 6 mice, one-sample t-test, 2-s trials:

p = 0.557; 8-s trials: p < 0.05). (F) Change of correct rate for optogenetic excitation of D2-SPNs

in 2-s and 8-s trials (n = 8 mice, one-sample t-test, 2-s trials: p < 0.05; 8-s trials: p < 0.05). (G)

Change of correct rate for optogenetic inhibition of D2-SPNs in 2-s and 8-s trials (n = 5 mice,

one-sample t-test, 2-s trials: p < 0.05; 8-s trials: p < 0.05). (H) Schematic of center-surround

receptive field diagram for Go/No-Go (left) and Co-activation (right) models. '+' indicates

facilitating effect to selection. '-' indicates inhibitory effect to selection.

Figure 6. A triple-control computational model of basal ganglia direct and indirect

pathways for action selection. (A) Network structure of the cortico-basal ganglia model based

on realistic anatomy and synaptic connectivity. (B) Schematic of center-surround-context

receptive field diagram for 'Triple-control' model. '+' indicates facilitating effect to selection. '-'

indicates inhibitory effect to selection. (C) The psychometric curves of behavioral output in

control (black) and D1-SPNs ablation condition (blue) in 'Triple-control' model (n = 10, two-

way repeated-measures ANOVA, main effect of ablation, $F_{1,18} = 98.72$, $p < 0.0001$; interaction

between trial intervals and ablation, $F_{6,108} = 7.799$, $p < 0.0001$). (D) The psychometric curves

of behavioral output in control (black) and D2-SPNs ablation condition (red) in 'Triple-control'

model (n = 10, two-way repeated-measures ANOVA, main effect of ablation, $F_{1,18} = 99.54$, $p <$

0.0001; interaction between trial intervals and ablation, $F_{6,108} = 177.6$, $p < 0.0001$). (E) Change

of correct rate for optogenetic excitation of D1-SPNs in 2-s and 8-s trials (n = 10, one-sample t-

test, 2-s trials: $p = 0.407$; 8-s trials: $p < 0.05$). (F) Change of correct rate for optogenetic

excitation of D2-SPNs in 2-s and 8-s trials (n = 10, one-sample t-test, 2-s trials: $p < 0.05$; 8-s

trials: $p < 0.05$). (G) Change of correct rate for optogenetic inhibition of D1-SPNs in 2-s and 8-s

trials (n = 10, one-sample t-test, 2-s trials: $p = 0.28$; 8-s trials: $p < 0.05$). (H) Change of correct

rate for optogenetic inhibition of D2-SPNs in 2-s and 8-s trials (n = 10, one-sample t-test, 2-s

trials: $p < 0.05$; 8-s trials: $p < 0.05$).

Figure 7. Computational modeling reveals direct and indirect pathways regulating action

selection in a distinct manner. (A) Schematic for manipulation of D1-SPNs in 'Triple-control'

model. (B) Schematic of manipulation of D1-SPNs in the center-surround-context receptive field

diagram for 'Triple-control' model. '+' indicates facilitating effect to selection. '-' indicates

inhibitory effect to selection. (C) Correct rate change in 2s trials when manipulating D1-SPNs

with different manipulation strengths (n = 10, one-way repeated-measures ANOVA, effect of

manipulation strength, $F_{36,324} = 1.171$, $p = 0.238$). (D) Correct rate change in 8s trials when

manipulating D1-SPNs with different manipulation strengths (n = 10, one-way repeated-

measures ANOVA, effect of manipulation strength, $F_{36,324} = 13.71$, $p < 0.0001$). (E) Schematic

for optogenetic manipulation of D2-SPNs in 'Triple-control' model. (F) Schematic of

manipulation of D2-SPNs in the center-surround-context receptive field diagram for 'Triple-

control' model. '+' indicates facilitating effect to selection. '-' indicates inhibitory effect to

selection. (G) Correct rate change in 2s trials when manipulating D2-SPNs with different

manipulation strengths (n = 10, one-way repeated-measures ANOVA, effect of manipulation

strength, $F_{36,324} = 59.13$, $p < 0.0001$). (H) Correct rate change in 8s trials when manipulating

D2-SPNs with different manipulation strengths (n = 10, one-way repeated-measures ANOVA,

effect of manipulation strength, $F_{36,324} = 40.75$, $p < 0.0001$). (I) Diagram of linear modulation

of direct pathway. (J) Diagram of nonlinear modulation of indirect pathway.

STAR METHODS

Animals

All experiments were approved by the Salk Institute Animal Care, and done in accordance with NIH guidelines for the Care and Use of Laboratory Animals. Experiments were performed on both male and female mice, at least two months old, housed on a 12-hour light/dark cycle. C57BL/6J mice were purchased from the Jackson Laboratory at 8 weeks of age and used as wildtype mice. BAC transgenic mice expressing cre recombinase under the control of the dopamine D1 receptor (referred as D1-cre, GENSAT: EY217; minimal labeling in cortex; mostly dorsal labeling in striatum) or the A2a receptor (referred as A2a-cre, GENSAT: KG139) promoter were obtained from MMRRC, and either crossed to C57BL/6 or Ai32 (012569) mice obtained from Jackson Laboratory (Cui et al., 2013; Geddes et al., 2018; Jin et al., 2014; Madisen et al., 2012; Tecuapetla et al., 2016). Striatal neuron-type-specific NMDAR1-knockout (referred as NR1-KO) and littermate controls were generated by crossing D1-cre and A2a-cre mice with NMDAR1-loxP (also denoted as Grin1 flox/flox in the Jackson Laboratory database) mice. The behavioral experiments using NR1-KO mice were performed on 8 to 12 weeks old D1/A2a-cre + / NMDAR1-loxP homozygous mice and their littermate controls, including D1/A2a-cre +, D1/A2a-cre + / NMDAR1-loxP heterozygous and NMDAR1-loxP homozygous mice. There was no difference between the three control groups, so the data were combined.

Behavior task

Mice were trained on a temporal bisection task in the operant chamber (21.6 cm L × 17.8 cm W × 12.7 cm H), which was isolated within a sound attenuating box (Med-Associates, St. Albans, VT). The food magazine was located in the middle of one wall, and two retractable

levers were located to the left and right side of the magazine. A house light was (3 W, 24

V) mounted on the opposite wall of the magazine. Sucrose solution (10 %) was delivered

into a metal bowl in the magazine through a syringe pump. When a training session

started, the house light was turned on and two levers were extended. After a random time

interval (30 s on average), left and right levers were retracted and extended

simultaneously. Mice were able to make a choice by pressing either left or right lever.

Only the very first lever press after levers extension was registered as animals' choice. If

the interval between the levers retraction and extension was 2s, then only the left lever

press was active to trigger the sucrose reward; if the interval between the lever retraction

and extension was 8s, then only the right lever press was active to trigger the sucrose

reward (Howard et al., 2017). There was no punishment when mice made an unrewarded

choice. 2s-trial and 8s-trial were randomized and interleaved by random inter-trial

intervals (30 s on average). The mice were also trained in the reversed 2-8s task. In the

reversed 2-8s task, if the interval between the levers retraction and extension was 2s, then

only the right lever press was active to trigger the sucrose reward; if the interval between

the lever retraction and extension was 8s, then only the left lever press was active to

trigger the sucrose reward. Representative behavioral tracks were captured by EthoVision

(Noldus).

Behavior training

Mice were placed on food restriction throughout the training, and fed daily after

the training sessions with ~2.5 g of regular food to allow them to maintain a body weight

of around 85 % of their baseline weight. Training started with continuous reinforcement

(CRF), in which animals obtained a reinforcer after each lever press. The session began with the

illumination of the house light and extension of either left or right lever, and ended with the

retraction of the lever and the offset of the house light. On the first day of CRF, mice received 5

reinforcers on left and right lever. On the second day of CRF, mice received 10 reinforcers on

left and right lever. On the third day of CRF, mice received 15 reinforcers on left and right lever.

The order of left lever CRF and right lever CRF on each day was randomized across all the CRF

training days. After the training of CRF, animals started on the temporal bisection task (day 1).

Mice were trained in the temporal bisection task for 14 consecutive days. On each day, there

were 240 trials with 2s-trial and 8s-trial randomly intermixed at 50:50. After 14 days training,

mice received an interval discrimination test, in which 20% of 2s/8s trials were replaced by

probe trials. In probe trials, the levers retraction intervals were randomly selected from 2.3s, 3.2s,

4s, 5s and 6.3s. Neither choice in the probe trials was rewarded. Mice received 4 days of test,

interleaved by training days without probes. The animals were trained daily without interruption

and every day the training started approximately at same time (Howard et al., 2017). All

timestamps of lever presses, magazine entries and licks for each animal were recorded with 10

ms resolution. The training chambers and procedures for NR1-KO mice and their littermate

controls were exactly the same for C57BL/6J mice.

For the reversed task training, mice were trained in both the 2-8 s control task and

reversed version of 2-8s task on the same day for at least 14 days. During each day, mice were

trained in the 2-8s task first, and then mice were put back in the home cage for a 3-4 hours rest.

After the rest period, the same mice were trained in the reversed 2-8s task. The order of these

two tasks is fixed throughout the 14 days training.

Surgery

For in vivo electrophysiological data recording, each mouse was chronically

implanted with an electrode array which consists of an array of 2 rows × 8 columns

Platinum-coated tungsten microwire electrodes (35 μm diameter) with 150 μm spacing

between microwires in a row, and 250 μm spacing between 2 rows. The craniotomies

were made at the following coordinates: 0.5 mm rostral to bregma and 1.5 mm laterally

for dorsal striatum; 3.4 mm caudal to bregma and 1.0 mm laterally for SNr (Jin and Costa,

2010; Jin et al., 2014). During surgeries, the electrode arrays were gently lowered ~ 2.2

mm from the surface of the brain for dorsal striatum and ~ 4.3 mm for SNr, while

simultaneously monitoring neural activity. Final placement of the electrodes was

monitored online during the surgery based on the neural activity, and then confirmed

histologically at the end of the experiment after perfusion with 10 % formalin, brain

fixation in a solution of 30 % sucrose and 10 % formalin, followed by cryostat sectioning

(coronal slices of 40 - 60 μm).

For the cell type identification in striatum, the cre-inducible adeno-associated

virus (AAV) vector carrying the gene encoding the light-activated cation channel

channelrhodopsin-2 (ChR2) and a fluorescent reporter (DIO-ChR2-YFP/DIO-ChR2-

mCherry) was stereotactic injected into the dorsal striatum of D1-Cre or A2a-Cre mice,

enabling cell-type-specific expression of ChR2 in striatal D1-expressing or D2-

expressing projection neurons (at exactly the same coordinates of electrode array

implantation in striatum stated above). DIO-ChR2-YFP/DIO-ChR2-mCherry virus (1 μl,

one site, 10^{12} titer) was injected through a micro-injection Hamilton syringe, with the

whole injection taking ~10 min in total. The syringe needle was left in the position for 5–

10 min after the injection and then slowly moved out. Following viral injections or for

mice genetically expressing ChR2 under cre control (D1-Ai32, A2a-Ai32), electrode was

implanted as previously described (Geddes et al., 2018; Jin et al., 2014). The electrode array was

the same as used for dorsal striatum recording, but with a guiding cannula attached (Innovative

Neurophysiology) terminating ~300 μm above the electrode tips, and was implanted into the

same site after virus injection, allowing for simultaneous electrophysiological recording and light

stimulation. Following the implantation, a medal needle was inserted in the cannula and mice

were placed in the home cage for 2 weeks, allowing both viral expression and surgery recovery,

before further training and recording experiments.

For the optogenetic manipulation in striatum, we injected the AAV virus carrying the

gene for coding ChR2 (DIO-ChR2-YFP/DIO-ChR2-mCherry) or Halorhodopsin (DIO-

eNpHR3.0-eYFP). Virus was injected bilaterally at 0.5 mm rostral to bregma, 2 mm laterally and

~ 2.2 mm from the surface of the brain with 1 μl per site. 10 min after the virus injection, we

bilaterally implanted optical fiber units in dorsal striatum to the same site as virus injection. An

optical fiber unit was made by threading a 200-μm diameter, 0.37 NA optical fiber (Thor Labs)

with epoxy resin into a plastic ferrule (Geddes et al., 2018; Howard et al., 2017). Optical fiber

units were then cut and polished before the implantation.

For muscimol infusion in striatum, we bilaterally implanted cannulas (Plastics One, VA)

in wildtype mice to the site at 0.5 mm rostral to bregma, 2 mm laterally and ~ 2.2 mm from the

surface of the brain. After the implantation, cannulas were covered by dummy cannulas. Mice

were placed in the home cage for 2 weeks, allowing surgery recovery, before further training and

muscimol experiments.

For striatal neuron-type-specific ablation experiments, D1-cre and A2a-cre mice were

stereotaxically injected with a cre-inducible adeno-associated virus carrying the diphtheria toxin

receptor (Azim et al., 2014; Geddes et al., 2018) (AAV9-FLEX-DTR-GFP; Salk GT3

Core, CA). Virus was injected in eight different sites. We used two sets of AP/ML

coordinates for each hemisphere followed by two DV depths at each AP/ML site. The

coordinates were +0.9 mm AP, ±1.6 mm ML, -2.2 and -3.0 mm DV and 0.0 mm AP, ±2.1

mm ML, -2.2 and -3.0 mm DV. A Hamilton syringe was used to inject 1 uL at the four -

3.0 mm DV sites and another 0.5 uL at the four -2.2 mm DV sites for a total of 3 uL

injected per hemisphere. Following each injection, the needle was left in place for ~5

minutes and then raised over ~5 minutes. This same protocol was used for each injection

site.

Muscimol infusion

We daily trained wildtype mice with guide cannulas (Plastics One, VA) implanted

until they achieved at least 80% correct rate for 3 consecutive days, we started muscimol

infusion experiments. Muscimol was dissolved in saline before infusion (Sigma-Aldrich;

0.05 ug/ul). For the infusions, mice were briefly anesthetized with isoflurane and

injection cannulas (Plastics One, VA) were bilaterally inserted into the guide cannulas,

with the injection cannulas projecting 0.1 mm beyond the implanted guide cannulas. Each

injection cannula was attached to an infusion pump (BASi, IN) via polyethylene tubing.

Animals were bilaterally infused with 200 nL of liquid (saline or muscimol) followed by

a five-minute waiting period before removal of the infusion cannulas. Mice were returned

to their home cage and started in the behavioral task 30 minutes after infusion (Geddes et

al., 2018). To estimate the effects of muscimol on choice, we repeated saline controls and

muscimol infusions at least 3 times on a single mouse to gain enough probe trials for

psychometric curve fitting.

DTR-mediated cell ablation

For striatal neuron-type-specific ablation experiments, D1-cre and A2a-cre were injected

with AAV9-FLEX-DTR-GFP in striatum using the same coordinates described above. After

three-week recovery, mice were food-restricted and, following completion of CRF, underwent

training in the 2-8s task for two weeks. Immediately after day 14 of 2-8s task training, mice were

randomly divided into control and treatment groups. Treatment mice were administered mice 1

ug of diphtheria toxin (DT) dissolved in 300 uL of phosphate buffered saline (PBS) via

intraperitoneal (IP) injection on two consecutive days (Azim et al., 2014; Geddes et al., 2018),

whereas control mice received IP injections of PBS. To allow for neuronal ablation, animals

were stopped in behavioral training and placed back on food. Animals resumed 2-8s task training

with probe trials 14 days after the first DT or PBS injection.

Neural recordings during the task

The mice with electrode array implanted were trained with exactly the same procedure as

described above. When mice performed the 2-8s task with 80% correct rate for 3 consecutive

days, we connected the array with recording cable and continued training until mice adapted to

the mechanics of the recording cable and were able to maintain the correct rate greater than 80%

(Howard et al., 2017).

Neural activity was recorded using the MAP system (Plexon Inc., TX). The spike

activities were initially online sorted using a sorting algorithm (Plexon Inc., TX). Only spikes

with a clearly identified waveforms and relatively high signal-to-noise ratio were used for

further analysis. After the recording session, the spike activities were further sorted to

isolate single units by offline sorting software (Plexon Inc., TX). Single units displayed a

clear refractory period in the inter-spike interval histogram, with no spikes during the

refractory period (larger than 1.3 ms) (Geddes et al., 2018; Howard et al., 2017; Jin and

Costa, 2010; Jin et al., 2014). All the timestamps of animal's behavioral events were

recorded as TTL pulses which were generated by a Med-Associates interface board and

sent to the MAP recording system through an A/D board (Texas Instrument Inc., TX).

The animal's behavioral timestamps during the training session were synchronized and

recorded together with the neural activity.

1005

Neural dynamic analysis

The animal's behavior taking place during the lever retraction time period was

critical to the choice to be made, so we focused on the analysis of the neural activity from

levers retraction to levers extension. Neuronal firings aligned to lever retraction were

averaged across trials in 20-ms bins, and smoothed by a Gaussian filter (Gaussian filter

window size = 10, standard deviation = 5) to construct the peri-event histogram (PETH).

The neurons showing significant firing changes during the lever retraction period were

defined as task-related neurons (ANOVA); those showing no significant changes were

defined as non-task-related neurons, which were not included in the further dynamic

analysis.

During 2s trials, mice behaved exactly the same as they did during the 0s - 2s

period in the rewarded 8s trials, so we mainly analyzed firing activities in 8s trials. To

avoid confounding effect by the sensory responses triggered by the lever retraction, only neural

activity from 1s to 8s following lever retraction were included (Howard et al., 2017). Then we

calculated firing rate index (FRI) based on the PETH from 1s to 8s for each individual neuron as

follows:

$$\text{FRI} = \frac{\text{PETH} - \text{mean}(\text{PETH})}{\text{std}(\text{PETH})}$$

1022   We then used principal component analysis (PCA) and classification algorithm, a build-in

1023   toolbox in Matlab, to classify the task-related neurons based on types of dynamics. For striatum

1024   and SNr, we used the same algorithm to classify neurons, and we found the same types of

1025   dynamics in striatum and SNr: Type 1, monotonic decreasing; Type 2, monotonic increasing;

1026   Type 3, peak at around 4s; Type 4, trough at around 4s.

1027

Cell type classification

In dorsal striatum, we classified neurons as putative striatal projection neurons (SPNs) if

they showed waveform trough half-width between 100 μs and 250 μs and the baseline firing rate

less than 10 Hz. In substantia nigra pars reticulate (SNr), neurons with firing rate higher than 15

Hz were classified as putative SNr GABA neurons, which are most likely the SNr projection

neurons, because the percentage of GABAergic interneurons in the SNr is rather small (Deniau

JM, 2007; Jin and Costa, 2010).

To further identify the D1 and D2 SPNs in striatum, we utilized cre-loxp technique to

exclusively express ChR2 on D1-SPNs or D2-SPNs by injecting the AAV-DIO-ChR2-YFP/

AAV-DIO-ChR2-mCherry virus into dorsal striatum or genetically express ChR2 by D1-Ai32

and A2a-Ai32. Optical stimulation on ChR2-expressed cells is able to directly evoke spiking

activity with short latency (Geddes et al., 2018; Jin and Costa, 2010; Jin et al., 2014). Before the

training session, we connected the recording cable to the electrode array for neuronal

recording and inserted an optic fiber through the cannula attached to the array to conduct

light into striatum for light stimulation. For better monitoring of the same cells stably

during behavioral training and the later optogenetic identification, the optic fiber was

well fixed to the array. After each training session, we delivered blue light stimulation

through the optic fiber from a 473-nm laser (Laserglow Technologies) via a fiber-optic

patch cord, and simultaneously recorded the neuronal responses, to testify the molecular

identity of cells previously recorded during the behavioral training. The stimulation

pattern was 100-ms pulse width with 4s interval. The stimulation pattern was repeatedly

delivered for 100 trials. We very carefully regulated the laser power to a relatively low

level for each individual recording session which was strong enough to evoke reliable

spikes in a small population of neurons recorded from certain electrodes, since high laser

powers usually induced an electrical signal much larger and very different from the spike

waveforms previously recorded in the same electrode, presumably resulting from

synchronized activation of a large population of cells surrounding the electrode. For

neuron identification in different sessions in the same mouse, substantial effort was made

to optimize the position of optic fiber to identify those units recorded from different

electrodes and that were not being able to be identified in the previous session. The final

laser power used for reliable identification of D1/D2-SPNs was between 1.0 and 1.5 mW

measured at the tip of the optical fiber (slightly varying for different mice and different

sessions). Only those units showing very short (≤6 ms) response latency to light

stimulation and exhibiting exactly the same spike waveforms (R ≥ 0.95, Pearson's

correlation coefficient) during the behavioral performance and light response were

considered as direct light-activated and cre recombinase positive neurons thus D1-SPNs or D2-

SPNs (Geddes et al., 2018; Howard et al., 2017; Jin and Costa, 2010). Strict criteria were

employed to minimize the possibility of false positives (with the risk of increasing false

negatives, and hence having to perform more recordings/mice to achieve the same number of

neurons).

1068

Optical stimulation during the task

For optogenetic manipulation experiments, mice were injected with AAV virus carrying

were pre-trained in 2-8s task for two weeks and bilaterally implanted with optic fibers. After

achieving a correct rate of 80%, stimulation trials began. D1-SPNs and D2-SPNs neurons were

stimulated or inhibited bilaterally in 50% of trials using a single pulse of light (Laserglow, 473

nm, 5 mW, 1 s constant for ChR2 experiments; Laserglow, 532 nm 10 mW, 1 s constant for

Halorhodopsin experiments). Rewards were delivered only at correct responses during 2 and 8 s

trials. Within 50% of any type of trials, mice were optogenetically stimulated (or inhibited) for 1

s before lever extension (Howard et al., 2017). Mice only received stimulation (or inhibition)

once per trial. Sessions with correct rate below 75% for control trials were excluded from further

analysis.

1080

Computational model

We constructed a neuronal network model, including cortico-basal ganglia circuitry, to

simulate the behavioral effects of ablation and optogenetic manipulation on SPNs. Specifically,

cortical information corresponding to left or right choice is sent to D1- and D2-SPNs associated

with these two action options (Lo and Wang, 2006; Wang, 2002). One-way collateral inhibition

is added between D2 SPNs subgroups. Signals from D1- and D2-SPNs eventually

converge to two separate SNr populations through distinct pathways (Hikosaka et al.,

2000; Jin et al., 2014; Mink, 2003), and exert opposing effects on SNr activity (Smith et

al., 1998). Behavioral output is then determined by the dominant activity between the

mutually inhibiting left and right SNr populations (Mailly et al., 2003), which could

control the final motor output either through brainstem circuits or motor cortices (Aoki et

al., 2019; Hikosaka, 2007; Lo and Wang, 2006; Redgrave et al., 1999). Here for

simplicity, other basal ganglia nuclei such as globus pallidus and subthalamic nucleus are

not included in the model.

Cortical neurons firing activities are defined as:

$$f_{left}^{cortex}(t) = k_{left}^{cortex} e^{-t_m \cdot t} + I_{noise}^{left}(t)$$

$$f_{right}^{cortex}(t) = k_{right}^{cortex} e^{-t_m \cdot t} + I_{noise}^{right}(t)$$

1096 where $k_{left}^{cortex} = 2$, $k_{right}^{cortex} = -2$, $t_m = 0.4$ and $I_{noise}(t)$ is defined as Gaussian white noise

1097 (mean(I_{noise}^{left}) = 1, mean(I_{noise}^{right}) = 2, SD = 0.01).

1098 Dopamine neuron firing activities is defined as:

$$f_{DA}(t) = k_{DA} e^{-t_{DA} \cdot t} + I_{noise}^{DA}(t)$$

1099 where $k_{DA} = 3$, $t_{DA} = 0.4$ and $I_{noise}^{DA}(t)$ is defined as Gaussian white noise (mean (I_{noise}^{DA}) = 1,

1100 SD = 0.01).

1101   Neuronal activities of D1-SPNs are defined as:

$$\tau \frac{df_{left}^{D1}(t)}{dt} = w_0\left(E - f_{left}^{D1}(t)\right) + \widetilde{w}_{D1\_left} f_{left}^{cortex}(t) + w_{D1} f_{DA}(t) + I_{noise}(t)$$

$$\tau \frac{df_{right}^{D1}(t)}{dt} = w_0\left(E - f_{right}^{D1}(t)\right) + \widetilde{w}_{D1\_right} f_{right}^{cortex}(t) + w_{D1} f_{DA}(t) + I_{noise}(t)$$

1102   where $w_0 = 1$, $E = 20$, $\widetilde{w}_{D1\_left} = 3$, $\widetilde{w}_{D1\_right} = 6$, $w_{D1} = 2$, $\tau = 0.1$, $I_{noise}(t)$ is defined as

1103   Gaussian white noise (mean ($I_{noise}(t)$) = 0, SD = 0.5).

1104   Neuronal activities of D2-SPNs in "Co-activation" module (labelled as D2-SPN 1) are

1105   defined as:

$$\tau \frac{df_{left1}^{D2}(t)}{dt} = w_0\left(E - f_{left1}^{D2}(t)\right) + \widetilde{w}_{D2\_left}f_{left}^{cortex}(t) + w_{D2}f_{DA}(t) + I_{noise}(t)$$

$$\tau \frac{df_{right1}^{D2}(t)}{dt} = w_0\left(E - f_{right1}^{D2}(t)\right) + \widetilde{w}_{D2\_right}f_{right}^{cortex}(t) + w_{D2}f_{DA}(t) + I_{noise}(t)$$

where $w_0 = 1$, $E = 21$, $\widetilde{w}_{D2\_left} = 5$, $\widetilde{w}_{D2\_right} = 5$, $w_{D2} = -0.3$, $\tau = 0.1$, $I_{noise}(t)$ is defined

as Gaussian white noise (mean ($I_{noise}(t)$) = 0, SD = 0.5).

Neuronal activities of D2-SPNs in "Go/No-go" module (labelled as D2-SPN 2) are

defined as:

$\tau \frac{df_{left2}^{D2}(t)}{dt} = w_0\left(E - f_{left2}^{D2}(t)\right) + \widetilde{w}_{D2\_left}f_{left}^{cortex}(t) + w_{D2\_left}S_{left}(t)f_{right1}^{D2}(t) +$

$w_{D2}f_{DA}(t) + I_{noise}(t)$

$\tau \frac{df_{right2}^{D2}(t)}{dt} = w_0\left(E - f_{right2}^{D2}(t)\right) + \widetilde{w}_{D2\_right}f_{right}^{cortex}(t) + w_{D2\_right}S_{right}(t)f_{left1}^{D2}(t) +$

$w_{D2}f_{DA}(t) + I_{noise}(t)$

where $w_0 = 1$, $E = 21$, $\widetilde{w}_{D2\_left} = 5$, $\widetilde{w}_{D2\_right} = 5$, $w_{D2\_left} = -0.7$, $w_{D2\_right} = -0.5$,

$w_{D2} = -0.3$, $\tau = 0.1$, $I_{noise}(t)$ is defined as Gaussian white noise (mean ($I_{noise}(t)$) = 0, SD =

0.5). $S_{left}(t)$ and $S_{right}(t)$ are short-term depression functions:

$$S_{left}(t) = 3/(1 + e^{0.3(f_{right1}^{D2}(t)-15)})$$

$$S_{right}(t) = 3/(1 + e^{0.3(f_{left1}^{D2}(t)-15)})$$

SNr neurons receive striatal inputs as well as the local inhibitory inputs from other SNr

neurons. The SNr activities are defined as:

$\tau_{SNr}\dfrac{df_{left}^{SNr}(t)}{dt} = w_0\left(E - f_{left}^{SNr}(t)\right) + \widetilde{w1}_{SNr}f_{left}^{D1}(t) + \widetilde{w2}_{SNr}f_{right1}^{D2}(t) + \widetilde{w3}_{SNr}f_{left2}^{D2}(t) +$

$w_{left}^{SNr}f_{right}^{SNr}(t) + I_{noise}^{s}(t)$

$\tau_{SNr}\dfrac{df_{right}^{SNr}(t)}{dt} = w_0\left(E - f_{right}^{SNr}(t)\right) + \widetilde{w1}_{SNr-D1}f_{right}^{D1}(t) + \widetilde{w2}_{SNr}f_{left1}^{D2}(t) + \widetilde{w3}_{SNr}f_{right2}^{D2}(t) +$

$w_{right}^{SNr}f_{left}^{SNr}(t) + I_{noise}^{s}(t)$

where $w_0 = 1$, $E = 40$, $\widetilde{w1}_{SNr} = -0.1$, $\widetilde{w2}_{SNr} = 0.15$, $\widetilde{w3}_{SNr} = 0.07$, $w_{left}^{SNr} = -0.027$,

$w_{right}^{SNr} = -0.01$, $\tau_{SNr} = 0.2$. $I_{noise}^{s}(t)$ is defined as Gaussian white noise (mean (I_{noise}^{s})= 0, SD

$= 0.3$).

The time-dependent choice $C(t)$ is then determined by SNr outputs $f_{left}^{SNr}(t)$ and $f_{right}^{SNr}(t)$ as

follows:

$C(t) = \begin{cases} \text{left choice (short} - \text{duration choice)}, & f_{left}^{SNr}(t) - f_{right}^{SNr}(t) < 0 \\ \text{right choice (long} - \text{duration choice)}, & f_{left}^{SNr}(t) - f_{right}^{SNr}(t) \geq 0 \end{cases}$

For optogenetic manipulation of striatal neurons, the stimulation pattern is defined as:

$$F_{activation}(t) = \begin{cases} amp, & t_s \leq t \leq t_s + 1 \\ 0, & t < t_s \ or \ t > t_s + 1 \end{cases}$$

1130   and for inhibition, the pattern is defined as:

$$F_{inhibition}(t) = \begin{cases} -amp, & t_s \leq t \leq t_s + 1 \\ 0, & t < t_s \ or \ t > t_s + 1 \end{cases}$$

1131   where t_s is the onset of stimulation/inhibition, which lasts for 1 s. *amp* is defined as the strength

1132   of the optogenetic manipulation within the range of [1, 25].

1133   To add D1-D1 collateral connections to the 'Triple-control' model, the neuronal

1134   activities of D1-SPNs are defined as:

$$\tau\dfrac{df_{left}^{D1}(t)}{dt} = w_0\left(E - f_{left}^{D1}(t)\right) + \widetilde{w}_{D1_{left}}f_{left}^{cortex}(t) + w_{D1\_right}f_{right}^{D1}(t) + w_{D1}f_{DA}(t)$$

$$+ I_{noise}(t)$$

$$\tau \frac{df_{right}^{D1}(t)}{dt} = w_0\left(E - f_{right}^{D1}(t)\right) + \widetilde{w}_{D1_{right}}f_{right}^{cortex}(t) + w_{D1\_left}f_{left}^{D1}(t) + w_{D1}f_{DA}(t)$$

$$+ I_{noise}(t)$$

where $w_{D1\_left} = -0.3, w_{D1\_right} = -0.3$.

To add D1-D2 collateral connections to the 'Triple-control' model, the neuronal activities

of D2-SPNs in "Go/No-go" module (labelled as D2-SPN 2) are defined as:

$\tau \frac{df_{left2}^{D2}(t)}{dt} = w_0\left(E - f_{left2}^{D2}(t)\right) + \widetilde{w}_{D2\_left}f_{left}^{cortex}(t) + w_{D2_{left}}S_{left}(t)f_{right1}^{D2}(t) +$

$w_{D1_{left}}f_{left}^{D1}(t) + w_{D2}f_{DA}(t) + I_{noise}(t)$

$\tau \frac{df_{right2}^{D2}(t)}{dt} = w_0\left(E - f_{right2}^{D2}(t)\right) + \widetilde{w}_{D2\_right}f_{right}^{cortex}(t) + w_{D2\_right}S_{right}(t)f_{left1}^{D2}(t) +$

$w_{D1\_right}f_{right}^{D1}(t) + w_{D2}f_{DA}(t) + I_{noise}(t)$

where $w_{D1\_left} = -0.3, w_{D1\_right} = -0.3$.

To add D2-D1 collateral connections to the 'Triple-control' model, the neuronal activities

of D1-SPNs are defined as:

$$\tau \frac{df_{left}^{D1}(t)}{dt} = w_0\left(E - f_{left}^{D1}(t)\right) + \widetilde{w}_{D1_{left}}f_{left}^{cortex}(t) + w_{D2\_left}f_{left2}^{D2}(t) + w_{D1}f_{DA}(t)$$

$$+ I_{noise}(t)$$

$$\tau \frac{df_{right}^{D1}(t)}{dt} = w_0\left(E - f_{right}^{D1}(t)\right) + \widetilde{w}_{D1_{right}}f_{right}^{cortex}(t) + w_{D2\_right}f_{right2}^{D2}(t) + w_{D1}f_{DA}(t)$$

$$+ I_{noise}(t)$$

1145   where $w_{D2\_left} = -0.3, w_{D2\_right} = -0.3$.

1146   All the modeling programs were coded in Matlab.

1147

1148   **Psychometric curve fitting**

1149   Psychometric curves for behavioral data and for theoretical curves were fit using

1150   the following equation (Brunton et al., 2013; Howard et al., 2017):

$$y = a + \frac{b}{1 + e^{\frac{c-x}{d}}}$$

1151   where a is the percentage of long-lever selection during short duration trials, b is the difference

1152   between a and the percentage of long-lever selection during long duration trials, c is the x-

1153   intercept where long-duration selection equals 0.5, and d is the rate of increase or decrease in the

1154   curve (slope). These can be interpreted as change in overall choice, long-duration choice, time,

1155   and sensitivity, respectively (Brunton et al., 2013).

1156

Statistical procedures

Statistics for the wild-type and KO mice learning data were performed on the basis of

values for each mouse per day. One-way and two-way repeated measures ANOVA were used to

investigate general main effects; and paired or unpaired t-test were used in all planned and post-

hoc comparisons. Z-test was used for the comparison of neuron proportions (Sheskin, 2003).

Statistics for the optogenetic data were performed on the basis of control and stimulated values

for each mouse per stimulation condition. Statistical analyses were conducted in Matlab using

the statistics toolbox (The MathWorks Inc., MA) and GraphPad Prism 7 (GraphPad Software

Inc., CA). Results are presented as mean ± SEM for behavior readouts and the neuronal

recording data. $p < 0.05$ was considered significant. All statistical details are located within the

figure legends. The number of animals used in each experiment and the number of neurons are

specified in the text and figure legend.

1169

1170

References:

Abdi, A., Mallet, N., Mohamed, F.Y., Sharott, A., Dodson, P.D., Nakamura, K.C., Suri, S., Avery, S.V., Larvin, J.T., Garas, F.N.*, et al.* (2015). Prototypic and arkypallidal neurons in the dopamine-intact external globus pallidus. The Journal of neuroscience : the official journal of the Society for Neuroscience *35*, 6667-6688.

Abi-Dargham, A., Rodenhiser, J., Printz, D., Zea-Ponce, Y., Gil, R., Kegeles, L.S., Weiss, R., Cooper, T.B., Mann, J.J., Van Heertum, R.L.*, et al.* (2000). Increased baseline occupancy of D2 receptors by dopamine in schizophrenia. Proceedings of the National Academy of Sciences of the United States of America *97*, 8104-8109.

Albin, R.L., Young, A.B., and Penney, J.B. (1989). The functional anatomy of basal ganglia disorders. Trends in neurosciences *12*, 366-375.

Aoki, S., Smith, J.B., Li, H., Yan, X., Igarashi, M., Coulon, P., Wickens, J.R., Ruigrok, T.J., and Jin, X. (2019). An open cortico-basal ganglia loop allows limbic control over motor output via the nigrothalamic pathway. eLife *8*.

Atherton, J.F., Menard, A., Urbain, N., and Bevan, M.D. (2013). Short-term depression of external globus pallidus-subthalamic nucleus synaptic transmission and implications for patterning subthalamic activity. The Journal of neuroscience : the official journal of the Society for Neuroscience *33*, 7130-7144.

Azim, E., Jiang, J., Alstermark, B., and Jessell, T.M. (2014). Skilled reaching relies on a V2a propriospinal internal copy circuit. Nature *508*, 357-363.

Barbera, G., Liang, B., Zhang, L., Gerfen, C.R., Culurciello, E., Chen, R., Li, Y., and Lin, D.T. (2016). Spatially Compact Neural Clusters in the Dorsal Striatum Encode Locomotion Relevant Information. Neuron *92*, 202-213.

Bastide, M.F., Meissner, W.G., Picconi, B., Fasano, S., Fernagut, P.O., Feyder, M., Francardo, V., Alcacer, C., Ding, Y., Brambilla, R.*, et al.* (2015). Pathophysiology of L-dopa-induced motor and non-motor complications in Parkinson's disease. Progress in neurobiology *132*, 96-168.

Benecke, R., Rothwell, J.C., Dick, J.P., Day, B.L., and Marsden, C.D. (1987). Disturbance of sequential movements in patients with Parkinson's disease. Brain : a journal of neurology *110* *(Pt 2)*, 361-379.

Brunton, B.W., Botvinick, M.M., and Brody, C.D. (2013). Rats and humans can optimally accumulate evidence for decision-making. Science *340*, 95-98.

Calabresi, P., Picconi, B., Tozzi, A., Ghiglieri, V., and Di Filippo, M. (2014). Direct and indirect
pathways of basal ganglia: a critical reappraisal. Nature neuroscience *17*, 1022-1030.

Cazorla, M., de Carvalho, F.D., Chohan, M.O., Shegda, M., Chuhma, N., Rayport, S., Ahmari,
S.E., Moore, H., and Kellendonk, C. (2014). Dopamine D2 receptors regulate the anatomical and
functional balance of basal ganglia circuitry. Neuron *81*, 153-164.

Cools, R., and D'Esposito, M. (2011). Inverted-U-shaped dopamine actions on human working
memory and cognitive control. Biological psychiatry *69*, e113-125.

Cui, G., Jun, S.B., Jin, X., Pham, M.D., Vogel, S.S., Lovinger, D.M., and Costa, R.M. (2013).
Concurrent activation of striatal direct and indirect pathways during action initiation. Nature *494*,
238-242.

DeLong, M.R. (1990). Primate models of movement disorders of basal ganglia origin. Trends in
neurosciences *13*, 281-285.

Deniau JM, M.P., Maurice N, Charpier S. (2007). Gaba and the basal ganglia - from molecules
to systems, Vol 160 (Elsevier).

Ding, L., and Gold, J.I. (2012). Separate, causal roles of the caudate in saccadic choice and
execution in a perceptual decision task. Neuron *75*, 865-874.

Durieux, P.F., Schiffmann, S.N., and de Kerchove d'Exaerde, A. (2012). Differential regulation
of motor control and response to dopaminergic drugs by D1R and D2R neurons in distinct dorsal
striatum subregions. The EMBO journal *31*, 640-653.

Fujiyama, F., Nakano, T., Matsuda, W., Furuta, T., Udagawa, J., and Kaneko, T. (2016). A
single-neuron tracing study of arkypallidal and prototypic neurons in healthy rats. Brain structure
& function *221*, 4733-4740.

Fujiyama, F., Sohn, J., Nakano, T., Furuta, T., Nakamura, K.C., Matsuda, W., and Kaneko, T.
(2011). Exclusive and common targets of neostriatofugal projections of rat striosome neurons: a
single neuron-tracing study using a viral vector. Eur J Neurosci *33*, 668-677.

Gallistel, C.R. (1980). The Organization of Action: A New Synthesis.

Geddes, C.E., Li, H., and Jin, X. (2018). Optogenetic Editing Reveals the Hierarchical
Organization of Learned Action Sequences. Cell *174*, 32-43 e15.

Goldman-Rakic, P.S., Muly III, E.C., and Williams, G.V. (2000). D_1 receptors in prefrontal cells
and circuits. Brain Research Reviews.

Gong, S., Doughty, M., Harbaugh, C.R., Cummins, A., Hatten, M.E., Heintz, N., and Gerfen, C.R. (2007). Targeting Cre recombinase to specific neuron populations with bacterial artificial chromosome constructs. The Journal of neuroscience : the official journal of the Society for Neuroscience *27*, 9817-9823.

Gradinaru, V., Zhang, F., Ramakrishnan, C., Mattis, J., Prakash, R., Diester, I., Goshen, I., Thompson, K.R., and Deisseroth, K. (2010). Molecular and cellular approaches for diversifying and extending optogenetics. Cell *141*, 154-165.

Graybiel, A.M. (1998). The basal ganglia and chunking of action repertoires. Neurobiology of learning and memory *70*, 119-136.

Graybiel, A.M., and Rauch, S.L. (2000). Toward a neurobiology of obsessive-compulsive disorder. Neuron *28*, 343-347.

Gustafson, N., Gireesh-Dharmaraj, E., Czubayko, U., Blackwell, K.T., and Plenz, D. (2006). A comparative voltage and current-clamp analysis of feedback and feedforward synaptic transmission in the striatal microcircuit in vitro. Journal of neurophysiology *95*, 737-752.

Hikosaka, O. (2007). GABAergic output of the basal ganglia. Progress in brain research *160*, 209-226.

Hikosaka, O., Miyashita, K., Miyachi, S., Sakai, K., and Lu, X. (1998). Differential roles of the frontal cortex, basal ganglia, and cerebellum in visuomotor sequence learning. Neurobiology of learning and memory *70*, 137-149.

Hikosaka, O., Takikawa, Y., and Kawagoe, R. (2000). Role of the basal ganglia in the control of purposive saccadic eye movements. Physiol Rev *80*, 953-978.

Hikosaka, O., and Wurtz, R.H. (1983). Visual and oculomotor functions of monkey substantia nigra pars reticulata. IV. Relation of substantia nigra to superior colliculus. Journal of neurophysiology *49*, 1285-1301.

Horst, N.K., Jupp, B., Roberts, A.C., and Robbins, T.W. (2019). D2 receptors and cognitive flexibility in marmosets: tri-phasic dose–response effects of intra-striatal quinpirole on serial reversal performance. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology *44*, 564-571.

Howard, C.D., Li, H., Geddes, C.E., and Jin, X. (2017). Dynamic Nigrostriatal Dopamine Biases Action Selection. Neuron *93*, 1436-1450 e1438.

Howes, O.D., and Kapur, S. (2009). The Dopamine Hypothesis of Schizophrenia: Version III—The Final Common Pathway. Schizophrenia Bulletin *35*, 549-562.

Isomura, Y., Takekawa, T., Harukuni, R., Handa, T., Aizawa, H., Takada, M., and Fukai, T.
(2013). Reward-modulated motor information in identified striatum neurons. The Journal of
neuroscience : the official journal of the Society for Neuroscience *33*, 10209-10220.

Jarskog, L.F., Miyamoto, S., and Lieberman, J.A. (2007). Schizophrenia: new pathological
insights and therapies. Annual review of medicine *58*, 49-61.

Jiang, H., Stein, B.E., and McHaffie, J.G. (2003). Opposing basal ganglia processes shape
midbrain visuomotor activity bilaterally. Nature *423*, 982-986.

Jin, X., and Costa, R.M. (2010). Start/stop signals emerge in nigrostriatal circuits during
sequence learning. Nature *466*, 457-462.

Jin, X., and Costa, R.M. (2015). Shaping action sequences in basal ganglia circuits. Current
opinion in neurobiology *33*, 188-196.

Jin, X., Tecuapetla, F., and Costa, R.M. (2014). Basal ganglia subcircuits distinctively encode
the parsing and concatenation of action sequences. Nature neuroscience *17*, 423-430.

Klaus, A., and Plenz, D. (2016). A Low-Correlation Resting State of the Striatum during Cortical
Avalanches and Its Role in Movement Suppression. PLoS biology *14*, e1002582.

Kravitz, A.V., Freeze, B.S., Parker, P.R., Kay, K., Thwin, M.T., Deisseroth, K., and Kreitzer,
A.C. (2010). Regulation of parkinsonian motor behaviours by optogenetic control of basal
ganglia circuitry. Nature *466*, 622-626.

la Fougere, C., Meisenzahl, E., Schmitt, G., Stauss, J., Frodl, T., Tatsch, K., Hahn, K., Moller,
H.J., and Dresel, S. (2005). D2 receptor occupancy during high- and low-dose therapy with the
atypical antipsychotic amisulpride: a 123I-iodobenzamide SPECT study. Journal of nuclear
medicine : official publication, Society of Nuclear Medicine *46*, 1028-1033.

Lally, J., and MacCabe, J.H. (2015). Antipsychotic medication in schizophrenia: a review.
British medical bulletin *114*, 169-179.

Laruelle, M., D'Souza, C.D., Baldwin, R.M., Abi-Dargham, A., Kanes, S.J., Fingado, C.L.,
Seibyl, J.P., Zoghbi, S.S., Bowers, M.B., Jatlow, P.*, et al.* (1997). Imaging D2 receptor
occupancy by endogenous dopamine in humans. Neuropsychopharmacology : official
publication of the American College of Neuropsychopharmacology *17*, 162-174.

Lauwereyns, J., Watanabe, K., Coe, B., and Hikosaka, O. (2002). A neural correlate of response
bias in monkey caudate nucleus. Nature *418*, 413-417.

Lee, Y., Han, N.E., Kim, W., Kim, J.G., Lee, I.B., Choi, S.J., Chun, H., Seo, M., Lee, C.J., Koh,
H.Y.*, et al.* (2020). Dynamic Changes in the Bridging Collaterals of the Basal Ganglia Circuitry
Control Stress-Related Behaviors in Mice. Molecules and cells *43*, 360-372.

Levine, M., and Ruha, A.M. (2012). Overdose of atypical antipsychotics: clinical presentation,
mechanisms of toxicity and management. CNS drugs *26*, 601-611.

Lima, S.Q., Hromadka, T., Znamenskiy, P., and Zador, A.M. (2009). PINP: a new method of
tagging neuronal populations for identification during in vivo electrophysiological recording.
PloS one *4*, e6099.

Lo, C.C., and Wang, X.J. (2006). Cortico-basal ganglia circuit mechanism for a decision
threshold in reaction time tasks. Nature neuroscience *9*, 956-963.

Madisen, L., Mao, T., Koch, H., Zhuo, J.-m., Berenyi, A., Fujisawa, S., Hsu, Y.-W.A., Garcia,
A.J., Gu, X., and Zanella, S. (2012). A toolbox of Cre-dependent optogenetic transgenic mice for
light-induced activation and silencing. Nature neuroscience *15*, 793-802.

Mailly, P., Charpier, S., Menetrey, A., and Deniau, J.-M. (2003). Three-dimensional organization
of the recurrent axon collateral network of the substantia nigra pars reticulata neurons in the rat.
Journal of Neuroscience *23*, 5247-5257.

Mallet, N., Schmidt, R., Leventhal, D., Chen, F., Amer, N., Boraud, T., and Berke, J.D. (2016).
Arkypallidal Cells Send a Stop Signal to Striatum. Neuron *89*, 308-316.

Markowitz, J.E., Gillis, W.F., Beron, C.C., Neufeld, S.Q., Robertson, K., Bhagat, N.D., Peterson,
R.E., Peterson, E., Hyun, M., Linderman, S.W.*, et al.* (2018). The Striatum Organizes 3D
Behavior via Moment-to-Moment Action Selection. Cell *174*, 44-58 e17.

Mattay, V.S., Goldberg, T.E., Fera, F., Hariri, A.R., Tessitore, A., Egan, M.F., Kolachana, B.,
Callicott, J.H., and Weinberger, D.R. (2003). Catechol O-methyltransferase val158-met genotype
and individual variation in the brain response to amphetamine. Proceedings of the National
Academy of Sciences of the United States of America *100*, 6186-6191.

Mink, J.W. (1996). The basal ganglia: focused selection and inhibition of competing motor
programs. Progress in neurobiology *50*, 381-425.

Mink, J.W. (2003). The Basal Ganglia and involuntary movements: impaired inhibition of
competing motor patterns. Archives of neurology *60*, 1365-1368.

Nonomura, S., Nishizawa, K., Sakai, Y., Kawaguchi, Y., Kato, S., Uchigashima, M., Watanabe,
M., Yamanaka, K., Enomoto, K., Chiken, S.*, et al.* (2018). Monitoring and Updating of Action

Selection for Goal-Directed Behavior through the Striatal Direct and Indirect Pathways. Neuron *99*, 1302-1314 e1305.

Phillips, J.G., Chiu, E., Bradshaw, J.L., and Iansek, R. (1995). Impaired movement sequencing in patients with Huntington's disease: a kinematic analysis. Neuropsychologia *33*, 365-369.

Redgrave, P., Prescott, T.J., and Gurney, K. (1999). The basal ganglia: a vertebrate solution to the selection problem? Neuroscience *89*, 1009-1023.

Schmidt, R., and Berke, J.D. (2017). A Pause-then-Cancel model of stopping: evidence from basal ganglia neurophysiology. Philosophical transactions of the Royal Society of London Series B, Biological sciences *372*.

Seeman, P. (2002). Atypical antipsychotics: mechanism of action. Canadian journal of psychiatry Revue canadienne de psychiatrie *47*, 27-38.

Sheskin, D.J. (2003). Handbook of parametric and nonparametric statistical procedures (Chapman and Hall/CRC).

Smith, Y., Bevan, M., Shink, E., and Bolam, J.P. (1998). Microcircuitry of the direct and indirect pathways of the basal ganglia. Neuroscience *86*, 353-387.

Tai, L.H., Lee, A.M., Benavidez, N., Bonci, A., and Wilbrecht, L. (2012). Transient stimulation of distinct subpopulations of striatal neurons mimics changes in action value. Nature neuroscience *15*, 1281-1289.

Taverna, S., Ilijic, E., and Surmeier, D.J. (2008). Recurrent collateral connections of striatal medium spiny neurons are disrupted in models of Parkinson's disease. The Journal of neuroscience : the official journal of the Society for Neuroscience *28*, 5504-5512.

Tecuapetla, F., Carrillo-Reid, L., Bargas, J., and Galarraga, E. (2007). Dopaminergic modulation of short-term synaptic plasticity at striatal inhibitory synapses. Proceedings of the National Academy of Sciences *104*, 10258-10263.

Tecuapetla, F., Jin, X., Lima, S.Q., and Costa, R.M. (2016). Complementary Contributions of Striatal Projection Pathways to Action Initiation and Execution. Cell *166*, 703-715.

Wang, X.-J. (2002). Probabilistic decision making by slow reverberation in cortical circuits. Neuron *36*, 955-968.

Wei, W., and Wang, X.J. (2016). Inhibitory Control in the Cortico-Basal Ganglia-Thalamocortical Loop: Complex Regulation and Interplay with Memory and Decision Processes. Neuron *92*, 1093-1105.

Wong, D.F., Wagner, H.N., Jr., Tune, L.E., Dannals, R.F., Pearlson, G.D., Links, J.M.,
Tamminga, C.A., Broussolle, E.P., Ravert, H.T., Wilson, A.A*., et al.* (1986). Positron emission
tomography reveals elevated D2 dopamine receptors in drug-naive schizophrenics. Science *234*,
1558-1563.

Wu, Y., Richard, S., and Parent, A. (2000). The organization of the striatal output system: a
single-cell juxtacellular labeling study in the rat. Neurosci Res *38*, 49-62.

Wurtz, R.H., and Hikosaka, O. (1986). Role of the basal ganglia in the initiation of saccadic eye
movements. Progress in brain research *64*, 175-190.

Yokoi, F., Gründer, G., Biziere, K., Stephane, M., Dogan, A.S., Dannals, R.F., Ravert, H., Suri,
A., Bramer, S., and Wong, D.F. (2002). Dopamine D 2 and D 3 receptor occupancy in normal
humans treated with the antipsychotic drug aripiprazole (OPC 14597): a study using positron
emission tomography and [11 C] raclopride. Neuropsychopharmacology : official publication of
the American College of Neuropsychopharmacology *27*, 248-259.

1369

1370

Figure 1. The neuronal dynamics in SNr during the 2-8 s action selection task. (A) Schematic diagram for the design of 2-8 s task. (B) Correct rate for wild type mice across 14 days training (n = 10 mice, one-way repeated-measures ANOVA, significant effect of training days, $F_{13,117} = 32.54$, p < 0.0001). (C) Movement trajectory of an example mouse in correct (left panel) and incorrect (right panel) 8-s trials (gray line: trajectory of each trials; red/black line: the average trajectory). (D) Diagram of electrode array implanted into substantia nigra pars reticulata (SNr). (E) Firing Rate Index (FRI) of neuronal activity for all task-related SNr neurons in correct 8-s trials. The magnitude of FRI is color coded and the SNr neurons are classified as four different types based on the activity dynamics. (F-I) Averaged FRI for Type 1 (F), Type 2 (G), Type 3 (H), Type 4 (I) of SNr neurons in correct (red) and incorrect 8-s trials (gray). (J) The proportion of four types of SNr neurons. Type 1 and Type 2 are major types and significantly more than Type 3 and Type 4 (Z-test, p < 0.05). (K) Integrated SNr output defined as the subtraction of averaged FRI between Type 1 and Type 2 SNr neurons. (L) Averaged psychometric curve (n = 10 mice) of choice behavior. (M) The correlation between the Type 1 and Type 2 FRI subtraction and the behavioral choice (R = 0.98, p < 0.0005). Error bars denote s.e.m., same for below unless stated otherwise.

Figure 2. SNr neuronal dynamics reflect action selection but not interval time or reward value. (A) Task diagram of 2-8 s control task with 10% 16-s probe trials. (B) Percentage of behavioral choice in 2-s, 8-s and 16-s trials (blue: left choice; red: right choice) (n = 9 mice, paired t-test, p < 0.05). (C) Movement trajectory of an example mouse in 16-s trials (blue: left choice; red: right choice). (D) Averaged SNr Type 1 FRI in 16-s trials (red: left choice; black: right choice). Firing rates from 8s to 16s (highlighted area) are compared between left and right choice (n = 26 neurons, two-way repeated-measures ANOVA, significant difference between left and right choices, $F_{1,25}$ = 6.646, p = 0.016). (E) Averaged SNr Type 2 FRI in 16-s trials (red: left choice; black: right choice). Firing rates from 8s to 16s are compared between left and right choice (n = 16 neurons, two-way repeated-measures ANOVA, significant difference between left and right choices, $F_{1,15}$ = 5.785, p = 0.029). (F) Subtraction of FRI for SNr Type 1 and Type 2 neurons in 16-s probe trials (red: left choice; black: right choice). (G) Task design of 2-8 s standard task. (H) Percentage of behavioral choice in 2-s and 8-s trials (blue: left choice; red: right choice) (n = 6 mice, paired t-test, p < 0.05). (I) Movement trajectory of an example mouse in 8-s trials (blue: left choice; red: right choice). (J) Task design of reversed 2-8 s task. (K) Percentage of behavioral choice in 2-s and 8-s trials in the reversed 2-8 s task (blue: left choice; red: right choice) (n = 6 mice, paired t-test, p < 0.05). (L) Movement trajectory of the same mouse as (I) in 8-s trials in the reversed 2-8 s task (blue: left choice; red: right choice). (M) Averaged FRI of the SNr Type 1 neurons in correct 8-s trials (n = 14 neurons). (N) Averaged FRI of the SNr Type 2 neurons in correct 8-s trials (n = 11 neurons). (O) Integrated SNr output as the subtraction of FRI for SNr Type 1 (M) and Type 2 neurons (N) in the standard 2-8 s task. (P) Averaged FRI of the same neurons as (M) in correct 8-s trials of the reversed 2-8 s task. (Q) Averaged FRI of the same neurons as (N) in correct 8-s trials of the reversed 2-8 s task. (R) Integrated SNr output as the subtraction of FRI for SNr Type 1 (P) and Type 2 neurons (Q) in the reversed 2-8 s task.

Figure 3. Neuronal activity of striatal D1- and D2-SPNs during action selection. (A) FRI of neuronal activity for all task-related SPNs in correct 8-s trials. SPNs were classified as Type 1 - 4. (B-E) Averaged FRI for Type 1 (B), Type 2 (C), Type 3 (D), Type 4 (E) of SPNs in correct (red) and incorrect 8-s trials (gray). (F) Diagram of simultaneous neuronal recording and optogenetic identification of D1- vs. D2-SPNs in dorsal striatum. (G) Top panel: Raster plot for a representative D1-SPN response to 100 ms optogenetic stimulation. Each row represents one trial and each black dot represents a spike. Bottom panel: Peristimulus time histogram (PETH) aligned to light onset at time zero. (H) PETH for the same neuron as shown in (G) with a finer time scale. (I) Distribution of light response latencies for D1- and D2-SPNs. (J) Action potential waveforms of the same neuron in (G) for spontaneous (black) and light-evoked (orange) spikes (R = 0.998, P < 0.0001, Pearson's correlation). (K) Principal component analysis (PCA) of action potential waveforms showing the overlapped clusters of spontaneous (black) and light-evoked (orange) spikes. (L) Proportion of D1-SPN subtypes. Type 1 neurons are significantly more than other three types of neurons in D1-SPNs (Z-test, p < 0.05). (M) Averaged FRI for Type 1 (blue) and Type 2 (red) D1-SPNs in correct 8-s trials. (N) Proportion of D2-SPN subtypes. (O) Averaged FRI for Type 1 (blue) and Type 2 (red) D2-SPNs in correct 8-s trials.

Figure 4. Selective genetic knockout and ablation of D1- or D2-SPNs distinctly alters action selection. (A) Correct rate of control (n = 11 mice) and D1-NR1 KO mice (n = 16) in 2-8 s task during 14 days training (two-way repeated-measures ANOVA, significant difference between control and KO mice, $F_{1,25}$ = 10.8, p = 0.003). (B) Correct rate of control (n = 17) and D2-NR1 KO mice (n = 10) in 2-8 s task during 14 days training (two-way repeated-measures ANOVA, significant difference between control and KO mice, $F_{1,25}$ = 8.728, p = 0.007). (C) The psychometric curve for control (n = 11) and D1-NR1 KO mice (n = 16) (two-way repeated-measures ANOVA, significant difference between control and KO mice, $F_{1,25}$ = 12.27, p = 0.002). (D) The psychometric curve for control (n = 17) and D2-NR1 KO mice (n = 10) (two-way repeated-measures ANOVA, significant difference between control and KO mice, $F_{1,25}$ = 9.64, p = 0.005). (E) Schematic of muscimol infusion into the dorsal striatum in trained mice. (F) Correct rate for control (black: pre-muscimol, gray: post-muscimol) and mice with muscimol infusion (magenta) in dorsal striatum (n = 9 mice, paired t-test, p < 0.01). (G) The psychometric curve for control (n = 9 mice, black: pre-muscimol, gray: post-muscimol control) and mice with muscimol infusion (n = 9 mice, magenta) in dorsal striatum (two-way repeated-measures ANOVA, significant difference between control and muscimol infusion, $F_{2,16}$ = 11.74, p = 0.0007). (H) Timeline for selective diphtheria toxin (DT) ablation experiments. (I) Schematic of diphtheria toxin receptor (DTR) virus (AAV-FLEX-DTR-GFP) injection in dorsal striatum of D1-Cre mice. (J) Correct rate for control (n = 9 mice) and mice with dorsal striatum D1-SPNs ablation (D1-DTR, n = 8 mice) (two-sample t-test, p = 0.0016). (K) The psychometric curve for control (n = 9 mice) and D1-SPNs ablation mice (n = 8 mice) (two-way repeated-measures ANOVA, main effect of ablation, $F_{1,15}$ = 1.84, p = 0.195; interaction between trial intervals and ablation, F6,90 = 4.14, p = 0.001). (L) Movement trajectory of a control mouse in 8-s trials. (M) Movement trajectory of a D1-DTR mouse in 8-s trials. (N) Schematic of diphtheria toxin receptor (DTR) virus (AAV-FLEX-DTR-GFP) injection in dorsal striatum of A2a-Cre mice. (O) Correct rate for control (n = 8 mice) and mice with dorsal striatum D2-SPNs ablation (D2-DTR, n = 8 mice) (two-sample t-test, p = 0.005). (P) The psychometric curve for control (n = 9 mice) and D2-SPNs ablation mice (n = 8 mice) (two-way repeated-measures ANOVA, main effect of ablation, $F_{1,15}$ = 0.477, p = 0.5; interaction between trial intervals and ablation, $F_{6,90}$ = 12.6, p < 0.001). (Q) Movement trajectory of a control mouse in 8-s trials. (R) Movement trajectory of a D2-DTR mouse in 8-s trials. (S) Schematic of center-surround receptive field diagram for Go/No-Go (left) and Co-activation (right) models. '+' indicates facilitating effect to selection. '-' indicates inhibitory effect to selection.

Figure 5. Optogenetic manipulation of D1- vs. D2-SPNs differently regulates action selection. (A) Schematic of optic fiber implantation for experimentally optogenetic excitation or inhibition of D1- or D2-SPNs in the dorsal striatum. (B, C) Schematic for optogenetic excitation (B) and inhibition (C) of D1-/D2-SPNs for 1 s right before lever extension in 2-8 s task. (D) Change of correct rate for optogenetic excitation of D1-SPNs in 2-s and 8-s trials (n = 11 mice, one-sample t-test, 2-s trials: p = 0.248; 8-s trials: p < 0.05). (E) Change of correct rate for optogenetic inhibition of D1-SPNs in 2-s and 8-s trials (n = 6 mice, one-sample t-test, 2-s trials: p = 0.557; 8-s trials: p < 0.05). (F) Change of correct rate for optogenetic excitation of D2-SPNs in 2-s and 8-s trials (n = 8 mice, one-sample t-test, 2-s trials: p < 0.05; 8-s trials: p < 0.05). (G) Change of correct rate for optogenetic inhibition of D2-SPNs in 2-s and 8-s trials (n = 5 mice, one-sample t-test, 2-s trials: p < 0.05; 8-s trials: p < 0.05). (H) Schematic of center-surround receptive field diagram for Go/No-Go (left) and Co-activation (right) models. '+' indicates facilitating effect to selection. '-' indicates inhibitory effect to selection.

Figure 6. A triple-control computational model of basal ganglia direct and indirect pathways for action selection. (A) Network structure of the cortico-basal ganglia model based on realistic anatomy and synaptic connectivity. (B) Schematic of center-surround-context receptive field diagram for 'Triple-control' model. '+' indicates facilitating effect to selection. '-' indicates inhibitory effect to selection. (C) The psychometric curves of behavioral output in control (black) and D1-SPNs ablation condition (blue) in 'Triple-control' model (n = 10, two-way repeated-measures ANOVA, main effect of ablation, $F_{1,18} = 98.72$, p < 0.0001; interaction between trial intervals and ablation, $F_{6,108} = 7.799$, p < 0.0001). (D) The psychometric curves of behavioral output in control (black) and D2-SPNs ablation condition (red) in 'Triple-control' model (n = 10, two-way repeated-measures ANOVA, main effect of ablation, $F_{1,18} = 99.54$, p < 0.0001; interaction between trial intervals and ablation, $F_{6,108} = 177.6$, p < 0.0001). (E) Change of correct rate for optogenetic excitation of D1-SPNs in 2-s and 8-s trials (n = 10, one-sample t-test, 2-s trials: p = 0.407; 8-s trials: p < 0.05). (F) Change of correct rate for optogenetic excitation of D2-SPNs in 2-s and 8-s trials (n = 10, one-sample t-test, 2-s trials: p < 0.05; 8-s trials: p < 0.05). (G) Change of correct rate for optogenetic inhibition of D1-SPNs in 2-s and 8-s trials (n = 10, one-sample t-test, 2-s trials: p = 0.28; 8-s trials: p < 0.05). (H) Change of correct rate for optogenetic inhibition of D2-SPNs in 2-s and 8-s trials (n = 10, one-sample t-test, 2-s trials: p < 0.05; 8-s trials: p < 0.05).

Figure 7. Computational modeling reveals direct and indirect pathways regulating action selection in a distinct manner. (A) Schematic for manipulation of D1-SPNs in 'Triple-control' model. (B) Schematic of manipulation of D1-SPNs in the center-surround-context receptive field diagram for 'Triple-control' model. '+' indicates facilitating effect to selection. '-' indicates inhibitory effect to selection. (C) Correct rate change in 2s trials when manipulating D1-SPNs with different manipulation strengths (n = 10, one-way repeated-measures ANOVA, effect of manipulation strength, $F_{36,324} = 1.171$, p = 0.238). (D) Correct rate change in 8s trials when manipulating D1-SPNs with different manipulation strengths (n = 10, one-way repeated-measures ANOVA, effect of manipulation strength, $F_{36,324} = 13.71$, p < 0.0001). (E) Schematic for optogenetic manipulation of D2-SPNs in 'Triple-control' model. (F) Schematic of manipulation of D2-SPNs in the center-surround-context receptive field diagram for 'Triple-control' model. '+' indicates facilitating effect to selection. '-' indicates inhibitory effect to selection. (G) Correct rate change in 2s trials when manipulating D2-SPNs with different manipulation strengths (n = 10, one-way repeated-measures ANOVA, effect of manipulation strength, $F_{36,324} = 59.13$, p < 0.0001). (H) Correct rate change in 8s trials when manipulating D2-SPNs with different manipulation strengths (n = 10, one-way repeated-measures ANOVA, effect of manipulation strength, $F_{36,324} = 40.75$, p < 0.0001). (I) Diagram of linear relationship between the modulation of direct pathway and correct rate of action selection. (J) Diagram of nonlinear relationship between the modulation of indirect pathway and correct rate of action selection.

Figure S1. Behavioral performance across 14 days of training and the SNr neuronal recording on day 1. (A) Lever press latency after lever extension for wild type mice across 14 days training (n = 10 mice, one-way repeated-measures ANOVA, effect of training days, $F_{13,117} = 21.32$, $p < 0.0001$). (B) Lever press ratio for wild type mice across 14 days training (n = 10 mice, one-way repeated-measures ANOVA, effect of training days, $F_{13,117} = 6.472$, $p < 0.0001$). (C) Example of recording array placement in SNr (left) and validation of array placement in a cohort of wildtype mice (right). Inset better demonstrates small tracts formed by the array implant. (D, E) Firing activities of an example SNr neuron in correct 2-s (D) and 8-s trials (E) after 14 days training. Top panels: raster plot of spikes across trials aligned to lever retraction at time 0. Bottom panel: PETH plot. Red shaded areas highlight the initial 2-s period after the lever retraction in 2-s (D) and 8-s trials (E). (F) Averaged FRI for Type 1, Type 2, Type 3 and Type 4 of SNr neurons in correct (red) and incorrect 2-s trials (gray). (G) Firing Rate Index (FRI) of neuronal activity for all task-related SNr neurons in correct 8-s trials on day 1 of training. The magnitude of FRI is color coded and the SNr neurons are categorized as four subgroups based on the activity dynamics. (H-K) Averaged FRI of SNr neurons in correct (red) and incorrect 8-s trials (gray) on day 1 of training.

Figure S2. Behavioral statistics and neuronal dynamics of SNr neurons in the standard and reversed 2-8s tasks . (A) Correct rates of the same group of mice both in the standard and reversed 2-8s tasks (n = 6 mice, paired t-test, $p = 0.33$). (B) Lever press ratios of the same group of mice both in the standard and reversed 2-8s tasks (n = 6 mice, paired t-test, $p < 0.05$). (C) Averaged FRI of the SNr Type 3 neurons in correct 8-s trials of the standard 2-8s task. (D) Averaged FRI of the SNr Type 3 neurons in correct 8-s trials of the reversed 2-8s task. (E) Averaged FRI of the SNr Type 4 neurons in correct 8-s trials of the standard 2-8s task. (F) Averaged FRI of the SNr Type 4 neurons in correct 8-s trials of the reversed 2-8s task.

Figure S3. Striatum neuronal recording on day 1 of training, recording array and optic fiber placement validation. (A) Averaged FRI for Type 1, Type 2, Type 3 and Type 4 of SPNs in correct (red) and incorrect 2-s trials (gray). (B) Firing Rate Index (FRI) of neuronal activity for all task-related SPNs in correct 8-s trials on day 1 of training. The magnitude of FRI is color coded and the SPNs are categorized as four subgroups based on the activity dynamics. (C-F) Averaged FRI of SPNs in correct (red) and incorrect 8-s trials (gray). (G) Recording array affixed with a cannula implanted in D1-Ai32 or A2a-Ai32 mice. Light emitted by optic fiber placed through the attached cannula is in close proximity to the tips of the recording array. (H) Top-down view of the array implantation. (I) Example of array placement in dorsal striatum of a D1-Ai32 mouse (left) and validation of fiber placement in a cohort of D1-Ai32 mice (right). Inset better demonstrates the tract formed by the array implant. (J) Example of array placement in dorsal striatum of a A2a-Ai32 mouse (left) and validation of fiber placement in a cohort of A2a-Ai32 mice (right). Inset better demonstrates the tract formed by the array implant. (K) Example of fiber placement in dorsal striatum of a D1-cre mouse with AAV5-DIO-ChR2-mcherry injected (left) and validation of fiber placement in a cohort of D1-cre mice (right). Inset better demonstrates the tract formed by the fiber implant. (L) Example of fiber placement in dorsal striatum of a A2a-cre mouse with AAV5-DIO-ChR2-mcherry injected (left) and validation of fiber placement in a cohort of A2a-cre mice (right). Inset better demonstrates the tract formed by the fiber implant.

Figure S4. Simulation of lesion experiments in Go/No-Go, Co-activation and combination models. (A) Diagram of Go/No-Go model. (B) The psychometric curves of behavior outputs simulated by Go/No-Go model in control (black) and D1-SPNs ablation condition (blue). (C) The psychometric curves of behavior outputs simulated by Go/No-Go model in control (black) and D2-SPNs ablation condition (red). (D) Diagram of Co-activation model. (E) The psychometric curves of behavior outputs simulated by Co-activation model in control (black) and D1-SPNs ablation condition (blue). (F) The psychometric curves of behavior outputs simulated by Co-activation model in control (black) and D2-SPNs ablation condition (red). (G) Diagram of combination of Go/No-Go and Co-activation model. (H) The psychometric curves of behavior outputs simulated by combined model in control (black) and D1-SPNs ablation condition (blue). (I) The psychometric curves of behavior outputs simulated by combined model in control (black) and D2-SPNs ablation condition (red).

Figure S5. Simulation of optogenetic manipulation in Go/No-Go and Co-activation models. (A, B) Simulating optogenetic activation (A) and inhibition (B) of D1-SPNs at 2s. Blue bar above indicates optogenetic activation. Yellow bar above indicates optogenetic inhibition. (C, D) Simulating optogenetic activation (C) and inhibition (D) of D2-SPNs at 2s. (E, F) Simulating optogenetic activation (E) and inhibition (F) of D1-SPNs at 8s. (G, H) Simulating optogenetic activation (G) and inhibition (H) of D2-SPNs at 8s. (I, J) Diagram of D1-SPN (I) and D2-SPN (J) manipulation in Go/No-Go model. (K, L) Change of correct rate in 2-s and 8-s trials when activating (K) and inhibiting (L) D1-SPNs in Go/No-Go model. (M, N) Change of correct rate in 2-s and 8-s trials when activating (M) and inhibiting (N) D2-SPNs in Go/No-Go model. (O, P) Diagram of D1-SPN (O) and D2-SPN (P) manipulation in Co-activation model. (Q, R) Change of correct rate in 2-s and 8-s trials when activating (Q) and inhibiting (R) D1-SPNs in Co-activation model. (S, T) Change of correct rate in 2-s and 8-s trials when activating (S) and inhibiting (T) D2-SPNs in Co-activation model. One-sample test for all the change of correct rate. $*p < 0.05$.

Figure S6. The neuronal activities in the "Triple-control" model and simulation of lesion experiments. (A) The simulated neuronal dynamics quantified as FRI for the cortical neurons in 8s trials. (B) The simulated neuronal dynamics quantified as FRI for the D1-SPN in 8s trials. (C) The simulated neuronal dynamics quantified as FRI for the D2-SPN 1 in 8s trials. (D) The simulated neuronal dynamics quantified as FRI for the D2-SPN 2 in 8s trials. (E) Schematic of selective ablation of D1-SPNs in the "Triple-control" model. (F) The model's Type 1 and Type 2 SNr FRI in control condition (black) and under D1-SPNs ablation (blue). (G) The subtraction of FRI between Type 1 and Type 2 SNr neurons in control (black) and D1-SPNs ablation condition (blue). (H) Schematic of selective ablation of D2-SPNs in the "Triple-control" model. (I) The model's Type 1 and Type 2 SNr FRI in control condition (black) and under D2-SPNs ablation (red). (J) The subtraction of FRI between Type 1 and Type 2 SNr neurons in control (black) and D2-SPNs ablation condition (red).

Figure S7. Optogenetic activation of D1- vs. D2-SPNs differently regulates SNr activities in model and experiments. (A) A computational motif of indirect pathway with collateral inhibitory synapse D2-SPN 1→ D2-SPN 2. The collateral synapse between D2-SPNs exhibits short-term depression. (B) Relationship between synaptic strength of D2-SPN 1→ D2-SPN 2 and the firing rate of D2-SPN 1. (C) The inhibition effect of the collateral synapse between D2-SPNs. (D) Activation of presynaptic D2-SPN 1 at 1s. (E) Synaptic inhibition effect D2-SPN 1→ D2-SPN 2 synapse when activating D2-SPN 1 at 1s. (F) Activation of D2-SPN 2 at 1s. (G) Activation of presynaptic D2-SPN 1 at 7s. (H) Synaptic inhibition effect D2-SPN 1→ D2-SPN 2 synapse when activating D2-SPN 1 at 7s. (I) Activation of D2-SPN 2 at 7s. (J) SNr neuron activities responding to activation of D2-SPNs at 1s (blue) and 7s (purple). (K) Comparison of FRI changes in SNr caused by activation of D2-SPNs at 1s and 7s. (L) Schematic of simultaneous optogenetic excitation of D1- or D2-SPNs in the dorsal striatum and recording in SNr during action selection. (M) Averaged neuronal activities of an example SNr Type 1 neuron responding to optogenetic activation of D1-SPNs at 1s during 8-s trials. (N) The percentage of SNr Type 1 (left) and Type 2 (right) neurons showing excitation (blue) and inhibition (red) when stimulating D1-SPNs. (O) The percentage of SNr Type 1 (left) and Type 2 (right) neurons showing excitation (blue) and inhibition (red) when stimulating D2-SPNs. (P) Averaged neuronal activities of an example SNr Type 2 neuron responding to optogenetic activation of D1-SPNs at 1s (blue) and 8s (purple) during 8-s trials. (Q, R) Averaged neuronal activities of SNr Type 1 (Q) and Type 2 (R) neuron responding to optogenetic activation of D2-SPNs at 1s (blue) and 7s (purple) during 8-s trials. (S) Comparison of FRI changes in SNr Type 1 (left) and Type 2 (right) neurons caused by optogenetic activation of D1-SPNs at 1s and 7s. (T) Comparison of FRI changes in SNr Type 1 (left) and Type 2 (right) neurons caused by optogenetic activation of D2-SPNs at 1s and 7s (paired t-test, $p < 0.05$).

Figure S8. Computational modeling of optogenetic manipulation reveals that D1- vs. D2-SPNs differently regulates SNr outputs in the "Triple-control" model. (A, B) Schematic for optogenetic manipulation of D1-SPNs (A) and D2-SPNs (B) in the 'Triple-control' model. (C) Modeling of neuronal dynamics of SNr Type 1/Type 2 (left panel) and integrated output (right panel) under control (black) and activation (blue) of D1-SPNs at 2s. (D) Modeling of neuronal dynamics of SNr Type 1/Type 2 (left panel) and integrated output (right panel) under control (black) and activation (red) of D2-SPNs at 2s. (E) Modeling of neuronal dynamics of SNr Type 1/Type 2 (left panel) and integrated output (right panel) under control (black) and inhibition (blue) of D1-SPNs at 2s. (F) Modeling of neuronal dynamics of SNr Type 1/Type 2 (left panel) and integrated output (right panel) under control (black) and inhibition (red) of D2-SPNs at 2s. (G) Modeling of neuronal dynamics of SNr Type 1/Type 2 (left panel) and integrated output (right panel) under control (black) and activation (blue) of D1-SPNs at 8s. (H) Modeling of neuronal dynamics of SNr Type 1/Type 2 (left panel) and integrated output (right panel) under control (black) and activation (red) of D2-SPNs at 8s. (I) Modeling of neuronal dynamics of SNr Type 1/Type 2 (left panel) and integrated output (right panel) under control (black) and inhibition (blue) of D1-SPNs at 8s. (J) Modeling of neuronal dynamics of SNr Type 1/Type 2 (left panel) and integrated output (right panel) under control (black) and inhibition (red) of D2-SPNs at 8s.

Figure S9. Computational modeling of manipulation reveals that Go/No-Go and Co-activation model differently predicts the behavioral outcomes. (A) Diagram of Go/No-Go model. (B) Diagram of Co-activation model. (C) Correct rate change in 2s (left panel) and 8s trials (right panel) when manipulating D1-SPNs in Go/No-Go model with different manipulation strengths. (D) Correct rate change in 2s (left panel) and 8s trials (right panel) trials when manipulating D1-SPNs in Co-activation model with different manipulation strengths. (E) Correct rate change in 2s (left panel) and 8s trials (right panel) when manipulating D2-SPNs in Go/No-Go model with different manipulation strengths. (F) Correct rate change in 2s (left panel) and 8s trials (right panel) trials when manipulating D2-SPNs in Co-activation model with different manipulation strengths.

Figure S10. Computational modeling reveals that the linear and nonlinear modulation of action selection by direct versus indirect pathway qualitatively hold with additional striatal collateral connections. (A) Schematic for 'Triple-control' model with D1-D1 collateral connections. (B) Correct rate change in 2s trials (upper panel) and 8s trials (bottom panel) when manipulating D1-SPNs with different manipulation strengths (n = 10, one-way repeated-measures ANOVA, effect of manipulation strength, 2s trials: $F_{40,369} = 1.328$, $p = 0.0945$; 8s trials: $F_{40,369} = 7.595$, $p < 0.0001$). For comparison, the same simulation results as in Figure 7(C, G) are shown in gray. (C) Correct rate change in 2s trials (upper panel) and 8s trials (bottom panel) when manipulating D2-SPNs with different manipulation strengths (n = 10, one-way repeated-measures ANOVA, effect of manipulation strength, 2s trials: $F_{40,369} = 38.22$, $p < 0.0001$; 8s trials: $F_{40,369} = 34.29$, $p < 0.0001$). For comparison, the same simulation results as in Figure 7(D, H) are shown in gray. (D) Schematic for 'Triple-control' model with D1-D2 collateral connections. (E) Correct rate change in 2s trials (upper panel) and 8s trials (bottom panel) when manipulating D1-SPNs with different manipulation strengths (n = 10, one-way repeated-measures ANOVA, effect of manipulation strength, 2s trials: $F_{40,369} = 0.9335$, $p = 0.5893$; 8s trials: $F_{40,369} = 8.778$, $p < 0.0001$). For comparison, the same simulation results as in Figure 7(C, G) are shown in gray. (F) Correct rate change in 2s trials (upper panel) and 8s trials (bottom panel) when manipulating D2-SPNs with different manipulation strengths (n = 10, one-way repeated-measures ANOVA, effect of manipulation strength, 2s trials: $F_{40,369} = 40.94$, $p < 0.0001$; 8s trials: $F_{40,369} = 26.61$, $p < 0.0001$). For comparison, the same simulation results as in Figure 7(D, H) are shown in gray. (G) Schematic for 'Triple-control' model with D2-D1 collateral connections. (H) Correct rate change in 2s trials (upper panel) and 8s trials (bottom panel) when manipulating D1-SPNs with different manipulation strengths (n = 10, one-way repeated-measures ANOVA, effect of manipulation strength, 2s trials: $F_{40,369} = 0.6827$, $p = 0.9299$; 8s trials: $F_{40,369} = 10.06$, $p < 0.0001$). For comparison, the same simulation results as in Figure 7(C, G) are shown in gray. (I) Correct rate change in 2s trials (upper panel) and 8s trials (bottom panel) when manipulating D2-SPNs with different manipulation strengths (n = 10, one-way repeated-measures ANOVA, effect of manipulation strength, 2s trials: $F_{40,369} = 153.3$, $p < 0.0001$; 8s trials: $F_{40,369} = 38.38$, $p < 0.0001$). For comparison, the same simulation results as in Figure 7(D, H) are shown in gray.

Figure S11. Computational modeling of dopaminergic modulation in the "Triple-control" model. (A) Diagram of Triple-control model with dopaminergic modulation on SPNs. (B) Schematic of center-surround-context receptive field diagram with dopaminergic modulation added for 'Triple-control' model. '+' indicates facilitating effect to selection. '-' indicates inhibitory effect to selection. (C) Simulation of two types of dopamine dynamics (black: decreasing dopamine; blue: constant dopamine with no change) in 8s trials. (D) Psychometric curves corresponding to each dopamine dynamics (n = 10, two-way repeated-measures ANOVA, main effect of ablation, $F_{1,18} = 0.8743$, $p = 0.362$; interaction between trial intervals and ablation, $F_{6,108} = 8.261$, $p < 0.0001$). (E, F) Correct rate change in 2s (E) and 8s trials (F) trials when manipulating dopamine in 'Triple-control' model with different manipulation strengths (n = 10, one-way repeated-measures ANOVA, effect of manipulation strength, 2-s trials: $F_{36,324} = 3.868$, $p < 0.0001$; 8-s trials: $F_{36,324} = 39.98$, $p < 0.0001$).